SARS-CoV-2 RdRp is a versatile enzyme with proofreading activity and ability

to incorporate NHC into RNA by using diphosphate form molnupiravir as a

substrate

Maofeng Wang, Cancan Wu, Nan Liu, Fengyu Zhang, Hongjie Dong, Shuai Wang,

Min Chen, Xiaoqiong Jiang ＆ Lichuan Gu*

State Key Laboratory of Microbial Technology, Shandong University, 72 Binhai Road,

Qingdao 266237, P.R. China

*To whom correspondence should be addressed: L.G. (e-mail: lcgu@sdu.edu.cn)

Abstract

The coronavirus disease 2019 (COVID-19) has been ravaging throughout the world for almost two years and has severely impaired both human health and the economy. The causative agent, severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) employs the viral RNA-dependent RNA polymerase (RdRp) complex for genome replication and transcription, making RdRp an appealing target for antiviral drug development. Although the structure of the RdRp complex has been determined, the function of RdRp has not been fully characterized. Here we reveal that in addition to RNA dependent RNA polymerase activity, RdRp also shows exoribonuclease activity and consequently proofreading activity. We observed that RdRp and nsp14-ExoN, when combined, exhibit higher proofreading activity compared to RdRp alone. Moreover, RdRp can recognize and utilize nucleoside diphosphate (NDP) as substrate to synthesize RNA and can also incorporate β-d-N4-hydroxycytidine (NHC) into RNA while using diphosphate form molnupiravir as substrate.

Introduction

The pandemic of coronavirus disease 2019 (COVID-19) has severely impacted human health and global economy since the outbreak in late 2019[1,2]. The causative agent，severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) is highly homologous to 2002 SARS-CoV with a similarity of 79.5%, and considered as new human infection along with β-coronavirus[3,4]. As of the date of writing, more than 254.02 million infections and more than 5.11 million deaths have been confirmed (https://www.worldometers.info/coronavirus/?utm_campaign). Although many kinds of vaccines have been developed to contain the pandemic, the rapidly spreading mutant strains arise serious concern that the vaccines would not yield the expected outcome[5-8]. As result, there is an urgent need to find effective medication or treatment to counter the virus[9-11].

SARS-CoV-2 is a positive-sense single-stranded RNA virus with a large genome of approximately 30 kb organized in 14 open reading frames[9,12]. Replication of the genome and transcription of genes are all dependent on a protein complex known as RNA-dependent RNA polymerase (RdRp). The RdRp of SARS-CoV-2 contains a catalytic subunit (non-structural protein 12, nsp12) and two accessory subunits (nsp7 and nsp8)[13,14]. Due to the significant role of RdRp and the lack of homologues in humans, RdRp has become the most appealing target for anti-coronavirus drug development[15,16].

The structure of the nsp12-nsp7-nsp8 complex has been determined in various forms, which includes active and inhibited forms[17-21]. The structure of SARS-CoV-2 RdRp

complex indicate that nsp12 contains an N-terminal β hairpin (residues 31 to 50) and

a nucleotide transferase domain (NiRAN; residues 51 to 250)[20-22]. After the NiRAN

domain is an interface domain (residues 251 to 365), connected to the RdRp catalytic

domain (residues 366 to 920). The catalytic domain of RdRp is described as a

cup-shaped right-hand conformation, composed of fingers, palm and thumb

subdomains, forming a closed circle[23]. The combination of nsp7 and nsp8 stabilizes

the closed conformation. The active site of the catalytic domain of SARS-CoV-2

RdRp is formed by seven conserved catalytic motifs (motifs A to G) which are

necessary for catalysis[24-28].

These structures are crucial to understand the mechanism of RdRp activity and

inhibition by remdesivir or other inhibitors[23]. The understanding of these structures

can play an influential role in the development of more effective anti-coronavirus

drugs. However, an analysis of the structure does not mean we have complete

knowledge of RdRp. During an experiment to screen an array of FDA-approved drugs

for RdRp inhibitors, we unexpectedly discovered that RdRp exhibits exoribonuclease

activity. Additional research has indicated that RdRp displays proofreading activity as

well. Further observation suggests that RdRp and nsp14-ExoN, when combined,

exhibit higher proofreading activity compared to RdRp alone. Moreover, RdRp can

also use nucleoside diphosphate (NDP) as substrate to synthesize RNA, which may

give SARS-CoV-2 a great advantage for survival when ATP level inside host cells is

low. In addition, we proved that molnupiravir is fully functionable in its diphosphate

form in addition to triphosphate form. This can imply a new strategy for designing

COVID-19 treatment nucleoside analogue drugs with better efficacy and fewer side

effects.

Results

Exoribonuclease activity was observed during characterization of RdRp

polymerase activity

Since RdRp is by far the most attractive target for anti-coronavirus drug development,

thorough characterization of RdRp polymerase activity would greatly without doubt

contribute to discovering the effective strategy for drug screening[15,16]. Based on

previous research, SARS-CoV-2 RdRp polymerase activity was measured by

conducting a conventional RNA elongation assay[23]. When nucleoside triphosphate

(NTP) substrate is added, elongation catalyzed by RdRp complex occurred on the

RNA duplex, resulting in an intact double-stranded RNA product. Nsp12 alone could

not catalyze the extension reaction (Fig.1a). Surprisingly, when NTP was absent from

the system for control, the RNA duplex was markedly degraded. Degradation of RNA

duplex was inhibited when EDTA was added (Fig.1b). Our results indicated that in

addition to polymerase activity, RdRp also displays exoribonuclease activity despite

that neither sequence nor structure gives any clue of the existence of a functional

domain.

SARS-CoV-2 RdRp also has proofreading activity

Our finding that SARS-CoV-2 RdRp exhibits exoribonuclease activity prompted us to

speculate that RdRp may also possess proofreading activity. To test our speculation,

an RNA duplex (M1 RNA) consisting of a 13 nt and a 23 nt RNA with a U: G

mismatch at the 3'-end of the extending strand of RNA was prepared (Fig.2b). An

extension assay was then performed by using M1 RNA as the template. Result

showed that RdRp successfully extended the shorter RNA strand, just as it did with

the RNA duplex without mismatch, although the productivity was slightly lower

(Fig.3a). We hypothesized two possibilities: first, RdRp simply ignored the mismatch

and processed to 3' end leaving the U: G mismatch unchanged; second, RdRp

corrected the mismatch and then finished the extension. The nucleotide mismatch

always slowed down the extension reaction, no matter the case.

To figure out this process, nascent RNA strands were reverse transcribed into cDNAs.

The cDNAs were then amplified by 10 cycles of PCR and sequenced by

high-throughput sequencing. For each product, at least 20,000 sequences were

obtained (SI). The data showed three kinds of nascent RNAs were produced: (1)

mutation repaired GCUAUGUGAGAU**G**AAAGUUAACU, 39.29%; (2) mutation

ignored GCUAUGUGAGAUUAAAGUUAACU, 13.82%; (3) mutation leading to

frame shift GCUAUGUGAGAU**UG**AAAGUUAACU, 36.89%.

We also challenged RdRp with the RNA template (M2 RNA) consisting of a 13 nt and

a 23 nt RNA with a UU: GC double mismatch at the 3'-end of the extending strand of

RNA (Fig.2c). In this experiment, RNA extension was observed as well, but with a

much lower productivity (Fig.3a). The products were then processed and sequenced

as mentioned above. The results included four kinds of RNAs: (1) mutations half

repaired GCUAUGUGAGA**UG**AAAGUUAACU, 54.88%; (2) mutations completely

repaired GCUAUGUGAGA**CG**AAAGUUAACU, 20.69%; (3) mutation ignored

GCUAUGUGAGA**UU**AAAGUUAACU,   3.01%;   (4)   frame   shift,

GCUAUGUGAGA**UUG**AAAGUUAACU, 11.42%.

The data strongly suggested that SARS-CoV-2 RdRp itself possesses proofreading

activity.

Combination of RdRp and nsp14-ExoN exhibited higher proofreading activity

Previous studies indicated that nsp14 is highly conserved within the Coronaviridae

family and critical for the SARS-CoV replication fidelity[7,29,30]. Since we discovered

that SARS-CoV-2 RdRp itself has proofreading activity, we supposed that the

combination of the two proteins may exhibit higher proofreading activity. To confirm

this hypothesis, we first tested the exoribonuclease activity of nsp14-nsp10 complex

in vitro and chose 300 nM as the proper concentration of nsp14 for the next

experiments[31] (Fig.3b).

We then conducted a proofreading activity assay for RdRp in the presence of 300 nM

of nsp14 with a RNA duplex containing a single mismatch (Fig.3c). High-throughput

sequencing indicated three kinds of nascent RNAs were produced: (1) mutation

repaired GCUAUGUGAGAU**G**AAAGUUAACU, 56.91%; (2) mutation ignored

GCUAUGUGAGA**UU**AAAGUUAACU, 22.85%; (3) mutation leading to frame shift

GCUAUGUGAGA**UUG**AAAGUUAACU, 10.13%. In the case of double mismatch,

nsp14 did not show significant role in improving the proofreading activity of RdRp:

(1) mutations half repaired 61.17%; (2) mutations completely repaired 17.21%; (3)

mutation ignored 5.72%; (4) frame shift 5.90% (Fig.3c). These data unambiguously

indicated that combination of RdRp and nsp14 is conducive to maintaining higher

replication fidelity especially for single mutation correction.

RdRp synthesizes RNA by using NDP as substrate

It is common sense that RNA polymerases synthesize RNA by using four types of

nucleoside triphosphate (ATP, UTP, CTP and GTP) as substrate. However, the

unexpected discovery of the exoribonuclease activity of RdRp gave us the unusual

idea that RdRp may also be able to use nucleoside diphosphate (NDP) as substrate. To

verify our hypothesis, we performed a RNA extension assay by using NTP and NDP

as substrate respectively. The result was fascinating, for RdRp synthesized the same

amount of RNA regardless NTP or NDP as substrate (Fig.4a). This result also raised a

question of whether the RNA made from NDP has the right sequence. To answer this

question, cDNA was synthesized through reverse transcription of the RNA product,

and then amplified by 10 cycles of PCR with primers containing the restriction sites

of Xho1 and BamH1. PCR products were then ligated into the vector pET-15b and

used to transform *E. coli* DH5α strain. Plasmids extracted from the positive clones

were sequenced by Sanger sequencing. The results confirmed that RNA synthesized

from NDP has the right sequence (Fig.4b). This means that unlike most other RNA

polymerase RdRp uses both NTP and NDP as substrate.

SARS-CoV-2 RdRp incorporates NHC into RNA by using diphosphate form

molnupiravir as a substrate

Since RdRp has become the most important target for the development of antiviral

drugs, many nucleoside analogue drugs have been constructed to target RdRp for

COVID-19 treatment[32-35]. Among all these chemicals reported by far, molnupiravir

has been viewed as a promising drug candidate. The molecular mechanism of

molnupiravir has been determined recently (Fig.5a). Once ingested by patients,

molnupiravir undergoes stepwise phosphorylation to yield the active nucleoside

triphosphate analogue (β-d-N4-hydroxycytidine (NHC) triphosphate, or called MTP).

The RdRp then uses MTP as substrate and incorporates NHC monophosphate into

RNA at the position where the nucleotide in the template is A or G, resulting in "error

catastrophe" and virus death[36-38].

Although the mechanism of molnupiravir has been well established, our finding that

RdRp can use NDP as a substrate still raises a question: is NHC diphosphate (MDP)

also an active form and recognized by RdRp as a substrate? To address this question,

the 13+23 nt RNA duplex was used to perform the RNA extension assay with NDPs

and MDP as the substrates in comparison with the extension using NTPs and MTP as

the substrates. 13+23 nt RNA duplex allows ten nucleotides (nt) of extension (Fig.5b).

In this case, when uridine diphosphate (UDP) was replaced by MDP, the incorporation

of M did not obstruct the incorporation of next subsequent nucleotides, RdRp still

finished the extension reaction. Similar results were obtained, in the case of using

MTP as a substrate (Fig.5c). Thus, it is reasonable to speculate that both MTP and

MDP are the active form of monulpiravir inside the human body.

Discussion

Since its outbreak in late 2019, the COVID-19 pandemic has been plaguing people for

two years with no sign of abating[3,12]. Although many kinds of vaccines have been

developed to suppress the pandemic, they had not produced the anticipated result to

withhold the spread of SARS-CoV-2. Development of potent antivirals against

SARS-CoV-2 is still an urgent need[10,11].

Since RNA-dependent RNA polymerase (RdRp) is critical for viral genome

replication and transcription and conserved in RNA virus species, RdRp has become

the most appealing target for antivirals development[11,15,23,26]. Thoroughly

characterization of RdRp both structurally and biochemically would no doubt be

beneficial to drug development. The discovery that RdRp has exoribonuclease activity

and proofreading activity is a complete surprise. Although it is general knowledge that

SARS-CoV-2 has proofreading activity, this activity was thought to be attributed to

nsp14-ExoN[29-31,39]. Sequence analysis of RdRp does not give any sign of the existence

of an exoribonuclease domain or a domain of unknown function[21,23]. We speculate the

exoribonuclease activity to be the "reverse reaction" of the polymerization. More

research is needed to confirm and explain the mechanism.

The substrate usage of RdRp surprised us greatly. As far as we know, all RNA

polymerases, including DNA dependent and RNA dependent RNA polymerases, use

nucleoside triphosphate (NTP) as substrate to synthesize RNA under the guidance of

the template chain[40]. Our finding that SARS-CoV-2 RdRp uses both NTP and NDP as

substrate to synthesize RNA with equal efficiency makes it an extremely unique RNA

polymerase. By far, we don't know why SARS-CoV-2 RdRp is so distinctive or if

there are any other enzyme–possessing similar features. We speculate that the ability

to use NDP as substrate could be a great advantage for SARS-CoV-2. During the late

stage of infection, the host cell may fail to produce enough ATP and thus other NTPs.

By using NDP as substrate, SARS-CoV-2 can continue the process of genome

replication and assembly of progeny virus while most of the metabolic activities stall.

The newly discovered feature of RdRp may profoundly change the strategy of

antiviral drug design. Many antiviral drugs are nucleoside analogues that target viral

polymerases as inhibitors to terminate RNA chain elongation[41,42]. However, such

chain-terminating antivirals often do not work as effectively as expected in the

treatment of COVID-19 because SARS-CoV-2 has exonucleolytic proofreading

activity that can remove the incorporated inhibitors[23,32,33,42-44]. It was thought that the

proofreading activity was carried out by nsp14-ExoN and incapacitating nsp14-ExoN

may be a way to deal with this issue[29,31,43]. The proofreading activity of RdRp is

unfavorable to this kind of RNA chain terminators. This may also explain why

molnupiravir, a mutagenizing nucleoside analog, is much more effective against

SARS-CoV-2 than remdesivir since molnupiravir can evade the proofreading

mechanism by both RdRp and nsp14-ExoN[36,37].

It has long been general knowledge that to become active against virus nucleoside

analogue drugs must undergo stepwise addition of phosphate groups to become the

triphosphate form. For this outcome, this type of antivirals must be recognized by

three kinds of kinases[45]. Which raises a concern that the triphosphate form of

antivirals may be incorporated into host mRNA[38]. Even worse, mutagenic

ribonucleoside analogs could be reduced into the 2'-deoxyribonucleotide form by host

ribonucleotide reductase and then incorporated into DNA[36,38]. Our finding that RdRp

also uses MDP as a substrate could largely resolve these concerns. There'd be no

issue about whether the prodrugs can become triphosphate form inside host cells since

the diphosphate form is also active. Furthermore, nucleoside diphosphate analogues

which can evade nucleoside diphosphate kinase would cease to have possibility of

being incorporated into mRNA or DNA. The best strategy is to use the

membrane-permeable nucleotide diphosphate analog for COVID-19 treatment[45,46].

The advantage of this strategy is that the drug is not only able to bypass all the

phosphorylation steps but also have no risk of incorporation into host mRNA or DNA

(Fig.6).

Materials and Methods

Protein expression and purification

Preparation of SARS-CoV-2 RdRp, composed of nsp12, nsp7 and two copied of the

nsp8 subunits, was carried out as described in ref.[23]. The SARS-CoV-2 *nsp12*, *nsp10*

and *nsp7* gene were cloned into a modified pET-21b vector with the C-terminus

possessing a 6×His-tag. The *nsp8* gene was cloned into the modified pET-32a vector

with the N-terminus possessing a trx-His$_6$-tag and PreScission Protease site. The

plasmids were transformed into *E. coli* BL21 (DE3) and the transformed cells were

cultured at 37 °C in LB with a final concentration of 100 μg/ml ampicillin. The

nsp7-pET-21b plasmid was transformed into *E. coli* Rosetta-gami2 (DE3) and the

transformed cells were cultured at 37 °C in LB with a final concentration of 100

μg/ml ampicillin and 25 μg/ml chloramphenicol. The *nsp14* gene (nsp14-ExoN

(residues 1–289)) was cloned into the modified pET-28b vector with the N-terminus

possessing a MBP-tag and PreScission Protease site. The plasmid was then

transformed into *E. coli* BL21 (DE3) and the transformed cells were cultured at 37 °C

in LB with a final concentration of 100 μg/ml kanamycin.

Bacterial cultures were incubated by shaking at 200 rpm, 37 °C to an OD600 of 0.8,

then the temperature was lowered to 16 °C and a final concentration of 0.2 mM of

isopropyl β-D-1-thiogalactopyranoside (IPTG) was added to induce protein

expression for 20 h. Subsequently, the cells were harvested by centrifugation at 5000

$\times g$ for 15 min at 4 °C. The pellet was resuspended in lysis buffer containing 25 mM

TrisHCl pH 8.0, 250 mM NaCl, 4 mM $MgCl_2$, 10% Glycerol and homogenized with a

high-pressure cell disrupter at 4 °C. The lysate was centrifuged at 25,200 $\times g$ for 50

min at 4 °C, and the supernatant was then loaded onto Ni-NTA affinity

chromatography column for purification. Nsp12, nsp10, nsp7 were eluted with elution

buffer (25 mM TrisHCl pH 8.0, 200 mM NaCl, 4 mM $MgCl_2$, 250 mM Imidazole).

Nsp8 was subject to on-column tag cleavage by PPase (PreScission Protease) and then

eluted by the lysis buffer. The supernatant of nsp14-ExoN was loaded onto a

Dextrin-Sepharose gravity column (GE Healthcare), washed and then eluted with

elution buffer containing 25 mM TrisHCl pH 8.0, 100 mM NaCl and 20 mM maltose.

The MBP tag was then removed by PPase digestion. Next, all proteins were purified

by ion exchange chromatography (Source 15Q HR 16/10, GE Healthcare) and size

exclusion chromatography (Superdex 200 10/300GL, GE Healthcare) with 25 mM

TrisHCl pH 8.0, 200 mM NaCl, 4 mM $MgCl_2.$

For nsp12-nsp7-nsp8 complex assembly, protein samples were mixed with the molar

ratio of nsp7:nsp8:nsp12=2:2:1 at 4 °C overnight. The incubated RdRp complex was

then concentrated with a 100 kDa molecular weight cut-off centrifugal filter unit

(Millipore Corporation) and then further purified by size exclusion chromatography

using a SuperdexS200 10/300 GL column in 25 mM TrisHCl pH 8.0, 200 mM NaCl,

4 mM $MgCl_2$.

Preparation of fluorescently labeled RNA for polymerase activity and

proofreading assays

RNA template-product duplex was designed according to the published SARS-CoV-2

RNA extension assays[23]. FAM labeled 13 nt oligonucleotide with the sequence of

FAM - GCUAUGUGAGAUU and the 23 nt complementary oligonucleotide with the

sequence of CGAUACACUCUAAUUUCAAUUGA were synthesized for

polymerase activity assay. Two 23 nt oligonucleotides with U: G mutation (M1 RNA,

CGAUACACUCUACUUUCAAUUGA) or UU: CG mutation (M2 RNA,

CGAUACACUCUGCUUUCAAUUGA) were synthesized for proofreading assays.

All unmodified and 5' FAM-labeled RNA oligonucleotides were purchased from

Tsingke Biotechnology Co.,Ltd.. The oligonucleotides were mixed in equal molar

ratio in DEPC water, annealed by heating it to 95 °C for 10 min and gradually cooling

to room temperature to make the RNA duplex (Fig.2a, 2b, 2c).

RdRp polymerase activity assays

DEPC-treated water was used for all buffer preparation. RdRp at final concentration

of 2 μM was incubated with 200 nM 13+23 nt RNA duplex and 25 μM NTP (or

NDP/NMP) in a 20 μl reaction buffer containing 20 mM TrisHCl pH8.0, 10 mM KCl,

10 mM MgCl$_2$, 0.01% Triton-X100, 1 mM DTT for 40 min at 37 °C[23]. Eventually,

glycerol was added to a concentration of 6.5%. 10 μl RNA product for each reaction

was resolved on 20% denaturing polyacrylamide-urea gels and imaged with a

Tanon-5200 Multi Fluorescence Imager.

RdRp exoribonuclease activity assay

RdRp at final concentration of 2 μM was incubated with 200 nM 13+23nt RNA

duplex in 20 μl reaction buffer for 40 min at 37 °C. Reaction systems containing 50

mM EDTA with or without RdRp were prepared as the control group. Eventually,

glycerol was added to a concentration of 6.5%. 10 μl RNA product for each reaction

was resolved on 20% denaturing polyacrylamide-urea gels and imaged with a Clinx

Science Instruments Co., Ltd. (CSI).

Nsp14-ExoN exoribonuclease activity assay

Nsp10 and nsp14 were incubated at a molar ratio of 3:1 for nsp14-nsp10 complex

assembly[31]. Reaction buffer containing 300 nM of nsp14-nsp10 complex, 20 mM

TrisHCl pH 8.0, 10 mM KCl, 10 mM MgCl$_2$, 0.01% Triton-X100, 1 mM DTT and

250 nM M1 RNA was incubated at 37 °C for 1 h. Reaction system without

nsp14-nsp10 complex was used as a negative control. After incubation, glycerol was

added to a concentration of 6.5%. The product was resolved on 20% denaturing

polyacrylamide-urea gels, and imaged with a Tanon-5200 Multi Fluorescence Imager.

The proofreading activities assays for RdRp and nsp14

Double stranded RNAs were prepared by using the extension reaction catalyzed by 2 μM RdRp containing 250 nM M1 RNA / M2 RNA, 25 μM NTP, with or without 300 nM nsp14-nsp10 in a 20 μl reaction system at 37 °C for 1 h. 10 μl product for each reaction was resolved on 20% denaturing polyacrylamide-urea gels to confirm the dsRNAs production. The remaining dsRNA products were mixed with primer1 (acgatgcaaagtctcgacaaatggtcgataccaattcaCCGCTCGAGCGGAGTTAACTTT) and dNTPs at 65 °C for 10 min and then cooled down to 24 °C for primer annealing. 10 × MMulV buffer, RNase inhibitor, and MMulV reverse transcriptase were added to the above system at 24 °C for 2 h to obtain cDNA, and then the enzymes were inactivated at 70 °C for 15 min. Subsequently, primer2 (cagataaactataattcctaatcgcgaggtggcactgcaaCGCGGATCCGCGGCTATGTGAGA(T)) was added, and the high-fidelity DNA polymerase pfu-Phusion was used to perform PCR amplification. Individual PCR products were then purified by agarose gel electrophoresis for high-throughput sequencing (Sangon Biotech).

RNA extension assays with NDP as substrate.

RdRp at final concentration of 2 μM was incubated with 200 nM 13+23 nt RNA duplex and 25 μM NTP (or NDP/NMP) in a 20 μl reaction buffer containing 20 mM TrisHCl pH8.0, 10 mM KCl, 10 mM $MgCl_2$, 0.01% Triton-X100, 1 mM DTT for 40 min at 37 °C.

RNA extension was performed by incubating 2 μM RdRp, 250 nM 13+23 nt RNA duplex, 25 μM NDP in a 20μl reaction system at 37 °C for 1 h. 10 μl RNA product was resolved on 20% denaturing polyacrylamide-urea gels and imaged with a

Tanon-5200 Multi Fluorescence Imager to confirm the extension. The remaining RNA product, primer1 (CCGCTCGAGCGGAGTTAACTTT) and dNTPs were gently mixed at 65 °C for 10 min. The mixture was cooled down to 24 °C then 10× MMulV buffer and MMulV reverse transcriptase were added and incubated for 2 h to obtain cDNA. Thereafter, the reverse transcriptase was inactivated at 70 °C for 15 min. Subsequently, primer2 (CGCGGATCCGCG GCTATGTGAGAT) was added, and the product was amplified by PCR with catalyzed by the high-fidelity DNA polymerase pfu-Phusion. The PCR product was digested with Xho1 and BamH1 and ligated into pET-15b vector, and transformed into *E-coli* DH5α to obtain clones. Plasmids from positive clones were then sequenced with T7 universal primers.

RNA extension assays with MDP as substrate.

A RNA extension assay was performed as mentioned previously[37]. In this assay the 13+23 nt RNA duplex was used (Fig.5e). The 5'end of the RNA product strand contains a 6-carboxyfluorescein (FAM) fluorescent label, which allows us to monitor RNA elongation. RdRp can incorporate NHC monophosphate into RNA at the position where the nucleotide in the template is A. MDP and MTP were purchased from MedChemExpress.

For RNA extension RdRp at final concentration of 2 μM was incubated with 200 nM 13+23 nt RNA duplex and 50 μM NTP/NDP with or without MTP/MDP in 20 μl reaction buffer for 40 min at 37 °C. 10 μl RNA product for each reaction was resolved on 20% denaturing polyacrylamide-urea gels and imaged with a Tanon-5200 Multi Fluorescence Imager to confirm the extension.

Data availability

All data generated or analyzed during this study are included in this published article

(and its Supplementary Information files).

Acknowledgements

The authors thank Yan Zhang at Zhejiang University for offering SARS-CoV-2 RdRp

gene, Carina Muyao Gu at CBe-Learn School for helping with the English expression

of the manuscript. This work was supported by Shandong Provincial Key Research

and Development Program (2020CXGC011305).

Author contributions

M.W. and L.G. conceived, designed the experiments and analyzed the data. M.W.

performed functional assays of RdRp. M.W., C.W., and X.J. participated in expression,

purification of the SARS-CoV-2 RdRp complex and other proteins. N.L., F.Z., H.D.,

S.W. and M.C. helped with the experiments and participated in figure and manuscript

preparation. M.W. and L.G. wrote the manuscript.

Competing interests

The authors declare no competing interests.

References

1   Chen, Y., Liu, Q. & Guo, D. Emerging coronaviruses: Genome structure,
     replication, and pathogenesis. *J Med Virol* **92**, 418-423,
     doi:10.1002/jmv.25681 (2020).

2   Zhu, N. *et al.* A Novel Coronavirus from Patients with Pneumonia in China,
     2019. *N Engl J Med* **382**, 727-733, doi:10.1056/NEJMoa2001017 (2020).

3   Lu, R. *et al.* Genomic characterisation and epidemiology of 2019 novel

coronavirus: implications for virus origins and receptor binding. *The Lancet*

395, 565-574, doi:10.1016/s0140-6736(20)30251-8 (2020).

4   Chan, J. F. *et al.* Genomic characterization of the 2019 novel

human-pathogenic coronavirus isolated from a patient with atypical

pneumonia after visiting Wuhan. *Emerg Microbes Infect* **9**, 221-236,

doi:10.1080/22221751.2020.1719902 (2020).

5   Hoffmann, M. *et al.* SARS-CoV-2 variant B.1.617 is resistant to

bamlanivimab and evades antibodies induced by infection and vaccination.

Cell Rep **36**, 109415, doi:10.1016/j.celrep.2021.109415 (2021).

6   Liu, C. *et al.* Reduced neutralization of SARS-CoV-2 B.1.617 by vaccine and

convalescent serum. *Cell* **184**, 4220-4236 e4213,

doi:10.1016/j.cell.2021.06.020 (2021).

7   Mlcochova, P. *et al.*, doi:10.1101/2021.05.08.443253 (2021).

8   Planas, D. *et al.* Reduced sensitivity of SARS-CoV-2 variant Delta to antibody

neutralization. *Nature* **596**, 276-280, doi:10.1038/s41586-021-03777-9 (2021).

9   Rudd, C. E. GSK-3 Inhibition as a Therapeutic Approach Against SARs CoV2:

Dual Benefit of Inhibiting Viral Replication While Potentiating the Immune

Response. *Front Immunol* **11**, 1638, doi:10.3389/fimmu.2020.01638 (2020).

10   Liu, C. *et al.* Research and Development on Therapeutic Agents and Vaccines

for COVID-19 and Related Human Coronavirus Diseases. *ACS Cent Sci* **6**,

315-331, doi:10.1021/acscentsci.0c00272 (2020).

11 Quimque, M. T. J. *et al.* Virtual screening-driven drug discovery of SARS-CoV2 enzyme inhibitors targeting viral attachment, replication, post-translational modification and host immunity evasion infection mechanisms. *J Biomol Struct Dyn* **39**, 4316-4333, doi:10.1080/07391102.2020.1776639 (2021).

12 Wu, F. *et al.* A new coronavirus associated with human respiratory disease in China. *Nature* **579**, 265-269, doi:10.1038/s41586-020-2008-3 (2020).

13 Wu, C. *et al.* Analysis of therapeutic targets for SARS-CoV-2 and discovery of potential drugs by computational methods. *Acta Pharm Sin B* **10**, 766-788, doi:10.1016/j.apsb.2020.02.008 (2020).

14 Subissi, L. *et al.* One severe acute respiratory syndrome coronavirus protein complex integrates processive RNA polymerase and exonuclease activities. *Proc Natl Acad Sci U S A* **111**, E3900-3909, doi:10.1073/pnas.1323705111 (2014).

15 McKee, D. L., Sternberg, A., Stange, U., Laufer, S. & Naujokat, C. Candidate drugs against SARS-CoV-2 and COVID-19. *Pharmacol Res* **157**, 104859, doi:10.1016/j.phrs.2020.104859 (2020).

16 Chu, C. K. *et al.* Antiviral Activity of Nucleoside Analogues against SARS-coronavirus (SARS-CoV). **17**, 285-289, doi:10.1177/095632020601700506 (2006).

17 Zhai, Y. *et al.* Insights into SARS-CoV transcription and replication from the structure of the nsp7-nsp8 hexadecamer. *Nat Struct Mol Biol* **12**, 980-986,

doi:10.1038/nsmb999 (2005).

18   Johnson, M. A., Jaudzems, K. & Wuthrich, K. NMR Structure of the

SARS-CoV Nonstructural Protein 7 in Solution at pH 6.5. *J Mol Biol* **402**,

619-628, doi:10.1016/j.jmb.2010.07.043 (2010).

19   Peti, W. *et al.* Structural genomics of the severe acute respiratory syndrome

coronavirus: nuclear magnetic resonance structure of the protein nsP7. *J Virol*

79, 12905-12913, doi:10.1128/JVI.79.20.12905-12913.2005 (2005).

20   Gao, Y. *et al.* Structure of the RNA-dependent RNA polymerase from

COVID-19 virus. *Science* **368**, 779-782, doi:10.1126/science.abb7498 (2020).

21   Kirchdoerfer, R. N. & Ward, A. B. Structure of the SARS-CoV nsp12

polymerase bound to nsp7 and nsp8 co-factors. *Nat Commun* **10**, 2342,

doi:10.1038/s41467-019-10280-3 (2019).

22   Lehmann, K. C. *et al.* Discovery of an essential nucleotidylating activity

associated with a newly delineated conserved domain in the RNA

polymerase-containing protein of all nidoviruses. *Nucleic Acids Res* **43**,

8416-8434, doi:10.1093/nar/gkv838 (2015).

23   Yin, W. *et al.* Structural basis for inhibition of the RNA-dependent RNA

polymerase from SARS-CoV-2 by remdesivir. *Science* **368**, 1499-1504,

doi:10.1126/science.abc1560 (2020).

24   Vázquez, A. L., Alonso, J. M. & Parra, F. Mutation analysis of the GDD

sequence motif of a calicivirus RNA-dependent RNA polymerase. *Journal of*

virology **74**, 3888-3891, doi:10.1128/jvi.74.8.3888-3891.2000 (2000).

25   Wang, Y. *et al.* Mutational analysis of the GDD sequence motif of classical

swine fever virus RNA-dependent RNA polymerases. *Virus Genes* **34**, 63-65,

doi:10.1007/s11262-006-0001-z (2007).

26   Jiang, Y., Yin, W. & Xu, H. E. RNA-dependent RNA polymerase: Structure,

mechanism, and drug discovery for COVID-19. *Biochem Biophys Res*

Commun **538**, 47-53, doi:10.1016/j.bbrc.2020.08.116 (2021).

27   Garriga, D., Ferrer-Orta, C., Querol-Audi, J., Oliva, B. & Verdaguer, N. Role

of motif B loop in allosteric regulation of RNA-dependent RNA

polymerization activity. *J Mol Biol* **425**, 2279-2287,

doi:10.1016/j.jmb.2013.03.034 (2013).

28   Gong, P. & Peersen, O. B. Structural basis for active site closure by the

poliovirus RNA-dependent RNA polymerase. *Proc Natl Acad Sci U S A* **107**,

22505-22510, doi:10.1073/pnas.1007626107 (2010).

29   Becares, M. *et al.* Mutagenesis of Coronavirus nsp14 Reveals Its Potential

Role in Modulation of the Innate Immune Response. *J Virol* **90**, 5399-5414,

doi:10.1128/JVI.03259-15 (2016).

30   Ma, Y. *et al.* Structural basis and functional analysis of the SARS coronavirus

nsp14-nsp10 complex. *Proc Natl Acad Sci U S A* **112**, 9436-9441,

doi:10.1073/pnas.1508686112 (2015).

31   Bouvet, M. *et al.* RNA 3'-end mismatch excision by the severe acute

respiratory syndrome coronavirus nonstructural protein nsp10/nsp14

exoribonuclease complex. *Proc Natl Acad Sci U S A* **109**, 9372-9377,

doi:10.1073/pnas.1201130109 (2012).

32   Choy, K. T. *et al.* Remdesivir, lopinavir, emetine, and homoharringtonine

inhibit SARS-CoV-2 replication in vitro. *Antiviral Res* **178**, 104786,

doi:10.1016/j.antiviral.2020.104786 (2020).

33   Wang, M. *et al.* Remdesivir and chloroquine effectively inhibit the recently

emerged novel coronavirus (2019-nCoV) in vitro. *Cell Res* **30**, 269-271,

doi:10.1038/s41422-020-0282-0 (2020).

34   Williamson, B. N. *et al.* Clinical benefit of remdesivir in rhesus macaques

infected with SARS-CoV-2. *Nature* **585**, 273-276,

doi:10.1038/s41586-020-2423-5 (2020).

35   Xie, X. *et al.* A nanoluciferase SARS-CoV-2 for rapid neutralization testing

and screening of anti-infective drugs for COVID-19. *Nat Commun* **11**, 5214,

doi:10.1038/s41467-020-19055-7 (2020).

36   Zhou, S. *et al.* beta-d-N4-hydroxycytidine Inhibits SARS-CoV-2 Through

Lethal Mutagenesis But Is Also Mutagenic To Mammalian Cells. *J Infect Dis*

224, 415-419, doi:10.1093/infdis/jiab247 (2021).

37   Kabinger, F. *et al.* Mechanism of molnupiravir-induced SARS-CoV-2

mutagenesis. *Nat Struct Mol Biol* **28**, 740-746,

doi:10.1038/s41594-021-00651-0 (2021).

38   Malone, B. & Campbell, E. A. Molnupiravir: coding for catastrophe. *Nat*

Struct Mol Biol **28**, 706-708, doi:10.1038/s41594-021-00657-8 (2021).

39   Eckerle, L. D., Lu, X., Sperry, S. M., Choi, L. & Denison, M. R. High fidelity

of murine hepatitis virus replication is decreased in nsp14 exoribonuclease mutants. *J Virol* **81**, 12135-12144, doi:10.1128/JVI.01296-07 (2007).

Malone, B. *et al.* Structural basis for backtracking by the SARS-CoV-2 replication-transcription complex. *bioRxiv*, doi:10.1101/2021.03.13.435256 (2021).

Sheahan, T. P. *et al.* An orally bioavailable broad-spectrum antiviral inhibits SARS-CoV-2 in human airway epithelial cell cultures and multiple coronaviruses in mice. *Sci Transl Med* **12**, eabb5883, doi:10.1126/scitranslmed.abb5883 (2020).

Zandi, K. *et al.* Repurposing Nucleoside Analogs for Human Coronaviruses. *Antimicrob Agents Chemother* **65**, e01652-01620, doi:10.1128/AAC.01652-20 (2020).

Robson, F. *et al.* Coronavirus RNA Proofreading: Molecular Basis and Therapeutic Targeting. *Mol Cell* **79**, 710-727, doi:10.1016/j.molcel.2020.07.027 (2020).

Liu, C. *et al.* Structural basis of mismatch recognition by a SARS-CoV-2 proofreading enzyme. *Science* **373**, 1142-1146, doi:10.1126/science.abi9310 (2021).

Meier, C. Nucleoside diphosphate and triphosphate prodrugs - An unsolvable task? *Antivir Chem Chemother* **25**, 69-82, doi:10.1177/2040206617738656 (2017).

Gollnest, T. *et al.* Membrane-permeable Triphosphate Prodrugs of Nucleoside

526   Analogues. *Angew   Chem   Int   Ed   Engl* **55**,   5255-5258,

527   doi:10.1002/anie.201511808 (2016).

 Figure legends

 Fig.1 RdRp both synthesizes and degrades RNA. **a** RdRp complex shows polymerase

activity in vitro. The primer strand is labeled with fluorescence at the 5′ end. The

positions of the primer and the full-length extension product are indicated. **b** RNA

degradation catalyzed by RdRp (RNA substrate is corresponding to 13+23nt RNA

duplex without fluorescence) when NTP is absent. EDTA prevents RNA degradation

by RdRp.

Fig.2 The RNA duplexes for primer extension and RdRp proofreading. **a** The 13+23

nt RNA template-product duplex. The direction of RNA extension is shown. The color

of the depicted circles indicates the experimental design: blue, RNA template strand;

red, RNA product strand. The 5'end of the RNA product contains a FAM fluorescent

label. **b** The primer/template bearing a U: C mismatch (M1 RNA). **c** The

primer/template bearing a UU: GC double mismatches (M2 RNA).

Fig.3 Proofreading activity of RdRp and its combination with nsp14-ExoN. **a** The

13+23 nt RNA, M1 RNA, M2 RNA were used as template respectively. The positions

of the template RNA and the full-length extension product are indicated. **b**

Fluorescence display of M1 RNA cleavage products. **c** The 13+23 nt RNA, M1 RNA,

M2 RNA were used as template respectively. Primer strand extension by RdRp alone

or with nsp10 + nsp14. The positions of the template RNA and the full-length

extension product and degraded product are indicated.

Fig.4 RdRp uses both NTP and NDP as substrate. **a** RdRp was incubated with the

13+23 nt primer/template in the presence of NMP, NDP or NTP respectively. The

positions of the template RNA and the full-length extension product are indicated. **b**

Part of the Sanger sequencing chromatogram of the RT-PCR products corresponding

to the extended 10 nt RNA on the 13+23 nt RNA duplex with NDP as substrate. The

dash line circle represents the 10 nt extension.

Fig.5 RdRp incorporates NHC into RNA with MDP as substrate. **a** Metabolism of

nucleoside analogues molnupiravir in host cell. **b** The 13+23 nt RNA allows for RNA

extension by 10 nucleotides. NHC monophosphate can be incorporated into growing

RNA instead of U or C. **c** RNA elongation stalls at the expected positions when the

cognate NDP is withheld from the reaction. Extension to the end of the template is

possible when MDP replaces UDP in the presence of other substrate NDPs, showing

that using MDP as a substrate RdRp incorporates M into RNA and incorporation of M

does not prevent RNA extension.

Fig.6 Nucleotide diphosphate analog prodrugs designed by our strategy would have

less potential to cause side effects. The terminal phosphate group (β-phosphate) is

modified by lipophilic masks to give the prodrug more membrane permeability. Inside

cell the masks are removed by enzymatic or chemical reactions. The ribose moiety

and the base group should be designed in such a way that the diphosphate form drug

can evade NDPK thus eliminating the risk of being incorporated into host mRNA or

DNA.

a

a

Molnupiravir → NHC → NHC-monophosphate=MMP

NMP-K ↓

NHC-diphosphate=MDP

NDP-K ↓

NHC-triphosphate=MTP

b

FAM
5' G C U A U G U G A G A U U → polymerization
3' C G A U A C A C U C U A A U U U C A A U U G A 5'

c

RNA+RdRp

RNA only | All NTPs | no UTP | +MTP | All NDPs | no UDP | +MDP

← Full-length product
← Stalled product
← Input RNA