**RNA-guided *As*Cas12a- and *Sp*Cas9-catalyzed knockout and homology
directed repair of the *omega-1* locus of the human blood fluke, *Schistosoma*
*mansoni***

**Wannaporn Ittiprasert[1,*,#], Chawalit Chatupheeraphat[1, 2,*], Victoria H. Mann[1], Wenhui
Li[1,3], André Miller[4], Taiwo Ogunbayo[4], Kenny Tran[4], Yousef Alrefaei[1,5], Margaret
Mentink-Kane[4], Paul J. Brindley[1,#]**

[1] Department of Microbiology, Immunology & Tropical Medicine, & Research Center for
Neglected Diseases of Poverty, School of Medicine & Health Sciences, George Washington
University, Washington, D.C. 20037, USA.

[2] Center for Research and Innovation, Faculty of Medical Technology, Mahidol University,
Salaya, Nakhon Pathom, 73170 Thailand

[3] Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou,
730050, Gansu Province, China

[4] Schistosomiasis Resource Center, Biomedical Research Institute, Rockville, MD 20850

[5] College of Health Sciences, PAEET, Adailiya, Kuwait City, 73101, Kuwait

*, equal contribution

#, correspondence: Wannaporn Ittiprasert, wannaporni@gwu.edu, Paul J. Brindley,
pbrindley@gwu.edu

Keywords: *As*Cas12a; *Sp*Cas9; ribonucleoprotein complex, CRISPR; omega-1; genome editing;
homology directed repair; non-homologous end-joining; schistosome egg

Abstract

We compared the efficiency of gene knockout (KO) and precision of insertion (knock-in, KI) of
the RNA-guided *As*Cas12a nuclease of *Acidaminococcus* sp. with that of *Sp*Cas9 from
Streptococcus pyogenes, aiming to enhance the functional genomics toolkit for *Schistosoma*
mansoni. Programmed DNA cleavages catalyzed by Cas12a and Cas9 result in staggered and
blunt ended strand breaks, respectively. TTTV, the optimal protospacer adjacent motif for
*As*Cas12a would occur frequently within the AT-rich genome of this platyhelminth. We
deployed optimized conditions for the ratio of guide RNAs to the nuclease, donor templates, and
electroporation parameters, to target a key enzyme termed omega-1 that is secreted by the
schistosome egg. *As*Cas12a was more efficient than *Sp*Cas9 for gene knockout of *omega-1* as
determined by tracking of indels by decomposition ($P < 0.001$). Resulting from CRISPREsso2
analysis, most mutations were deletions; *Sp*Cas9 induced short deletions of 3 nt in length
whereas *As*Cas12a induced deletions of 2 to 26 nt. Knockout efficiency of both nucleases
markedly increased in the presence of short, single stranded oligodeoxynucleotide (ssODN)
donor templates. With *As*Cas12a, ssODNs representative of both the non-CRISPR target (NT)
and target (T) strands of the targeted gene were tested, resulting in KO efficiencies of 15.67,
28.71 and 21.43% in the *Sp*Cas9 plus donor ssODN, *As*Cas12a plus NT-ssODN, and *As*Cas12a
plus T-ssODN groups, respectively. *Trans* cleavage activity against the ssODNs by activated
*As*Cas12a was not apparent *in vitro*. Programmed *Sp*Cas9 editing led to more precise transgene
insertion than *As*Cas12a, with KI efficiencies of 17.07% for the KI_SpCas9 group, 14.58% for
KI_*As*Cas12a-NT-ssODN and 12.37% for KI_*As*Cas12a-T-ssODN. Although *As*Cas12a induced
fewer mutations per genome than *Sp*Cas9, the phenotypic impact on transcription and expression
of omega-1 was similar for both nucleases. These findings revealed that *As*Cas12a and *Sp*Cas9
both provide tractable routes for RNA-guided programmed mutation of the genome of the
schistosome egg.
47

Introduction

Interest is increasing in functional genomics to investigate helminth parasites, specifically genome editing to investigate pathophysiology, carcinogenesis, and biology. Recent progress showcased the deployment of CRISPR/Cas programmed genome editing in parasitic flatworms, in particular with the human blood fluke *Schistosoma mansoni* (Ittiprasert *et al*, 2019) and the carcinogenic liver fluke, *Opisthorchis viverrini* (Arunsan *et al*, 2019) as exemplars. However, given that these helminth parasites are non-model organisms, that their genome sequences are often not available or well curated, and that they are frequently refractory to maintenance in the laboratory, approaches to advance functional genomics using CRISPR/Cas are not yet established or available for wide use in the field of tropical medical and neglected tropical diseases. Nonetheless, our capacity to decipher the molecular pathogenesis of these diseases will be aided by advances in gene editing methodologies (Hoffmann *et al*, 2014).

RNA-guided nucleases evolved in archaea and bacteria as adaptive immunity mechanism to destroy phage viruses and other nucleic acid invaders (Jinek *et al*, 2012; Koonin *et al*, 2017). The catalogue of these CRISPR/Cas nucleases is expanding. The *Sp*Cas9 of *Streptococcus pyogenes,* a Class 2, type II Cas enzyme in the phylogeny of Koonin and coworkers (Koonin *et al.*, 2017), dominates diverse gene-editing applications in biomedicine and beyond (Charpentier, 2015; Huang *et al*, 2018; Mustafa & Makhawi, 2020), given its programmable, precise double-stranded DNA (dsDNA) cleavage, accomplished by two catalytic domains, RuvC and HNH, of the nuclease at the target site complementary to the guide RNA (Nishimasu *et al*, 2014). A second Class 2 family, the type V nucleases as represented by *As*Cas12a (formerly known as Cpf1) uses a single RuvC catalytic domain for guide RNA-directed dsDNA cleavage (Jeon *et al*, 2018; Zetsche *et al*, 2015). Unlike *Sp*Cas9, *As*Cas12a orthologues recognize a T nucleotide–rich protospacer-adjacent motif (PAM), catalyze the maturation of their own guide CRISPR RNA (crRNA) and a dsDNA break, distal to the PAM, bearing staggered 5′- and 3′-ends (Jeon *et al.*, 2018). These Cas12a features are attractive for genome editing (Malzahn *et al*, 2019). For their maturation, the crRNAs of type II CRISPR-Cas systems require processing by a cognate trans-activating CRISPR RNA (Charpentier *et al*, 2015). Moreover, the *As*Cas12a nuclease is smaller in size than *Sp*Cas9 and uses a shorter CRISPR RNA (crRNA) for activity (Liu *et al*, 2017). Notably, target site activated *As*Cas12a exhibits non-specific *trans* activity against bystander nucleic acids (Chen *et al*, 2018; Smith *et al*, 2020), an attribute that has enabled its application in a burgeoning range of biosensors (Kellner *et al*, 2019; Li *et al*, 2018a).

Delivery of CRISPR reagents into the cells of the multicellular parasite *S. mansoni* in the form of ribonucleoprotein (RNP) complexes may enable immediate and permanent programmed gene editing and obviate concerns with CRISPR reagent longevity or integration in the host cell genome. Programmed CRISPR/Cas editing using RNP-based delivery can precede analysis of mutated target regions that rely on cloning and vector construction. Previously, we deployed both RNP and lentiviral-delivered CRISPR/*Sp*Cas9 approaches to knock-out (KO) and knock-in (KI) into a blunt end at *Sp*Cas9 cleavage site at the multiple copies of the omega-1 (*ω1*) gene in parasite eggs. To advance functional genomics for schistosomes and aiming to improve the mutagenesis and transgene knock-in of CRISPR efficiency in schistosomes, which exhibit AT-rich genomes, we investigated the nuclease activity of *As*Cas12a in comparison with *Sp*Cas9. These CRISPR systems differ in key ways: first, *Sp*Cas9 recognizes NGG as the protospacer adjacent motif (PAM) whereas *As*Cas12a recognizes the T-rich, TTTV, and second, catalysis by

*Sp*Cas9 results in blunt double stranded DNA at three nucleotides 5' to the PAM whereas
*As*Cas12a results in a staggered cleavage at 23 nt (target strand; T) and 18 nt (non-target strand;
NT) 3' to the PAM (Banakar *et al*, 2020). Comparisons of these two RNA-programmed
nucleases in the context of the *ω1* gene of *S. mansoni* may inform decisions for functional
genomics manipulations aiming to enhance efficiency of homology directed repair (HDR)
targeting genes expressed by the developing schistosome egg, and indeed other developmental
stages (Ittiprasert *et al.*, 2019; Sankaranarayanan *et al*, 2021; Takaki *et al*, 2021a; Takaki *et al*,
2021b).

Results

The omega-1 multicopy gene, guide RNAs, and single stranded DNA donor templates

The gene editing efficiency of *Sp*Cas9 and *As*Cas12a was compared in assays targeting the *ω1*
gene which encoded a T2 ribonuclease that is secreted by the mature egg of *S. mansoni* into the
surrounding host tissues and circumoval granuloma (Schwartz & Fallon, 2018). Specifically, the
four copies of *ω1*, well annotated in WormBase Parasite (Bolt *et al*, 2018) (update of 16
September 2021) as hepatotoxic ribonuclease – Smp_334170.1, Smp_334170.2, Smp_334240,
and Smp_333930, were targeted for design of guide RNAs (gRNA) and donor repair templates.
These four copies reside on chromosome 1 of *S. mansoni*, with reverse strand positions at
3,980,364 to 3,984675, 3,992,964 to 3,995,246 and 3,908,953 to 3,911,250 (PRJEA36577),
respectively. These copies share >99% nucleotide sequence identity, are ~2.3 kb in length, are
highly AT rich (65%), and encode an enzyme of 127 (Smp_334170.2, Smp_334240 and
Smp_333930) or 115 (Smp_334170.1) amino acid residues in length. Smp_333220, also located
on chromosome 1 at 3,925,420-3,927,716 but lacking annotation on the *S. mansoni* annotated
genome version 7, Year 2021, was also included. Smp_333220 shares >99% chromosomal and
mRNA sequence identity with the three *ω1* copies (Fig S1B). Guide RNAs for *Sp*Cas9 and
*As*Cas12a designed with the assistance of CHOPCHOP webtool both target the same region of
exon 1 of *ω1*, disrupting its catalytic active site histidine (Luhtala & Parker, 2010), with the
programmed cleavage sites for Cas in close proximity, separated by three nucleotides (Fig 1A
and B).

We designed three single stand oligodeoxynucleotide (ssODN) donors for homology directed
repair (HDR) of *ω1* following programmed mutation. Given the differing PAM requirements of
*Sp*Cas9 and *As*Cas12a as well as the divergent formats of the DSB, blunted ended by *Sp*Cas9 and
sticky ended by *As*Cas12a, the designs differed for the ssODNs for use with each nuclease.
Based on earlier findings (Ittiprasert *et al.*, 2019), the design of the ssODN donors included 3'
and 5' homology arms (HA) of each of 50 nt in length, complementary to the target strand for
*Sp*Cas9 and for *As*Cas12a, for either the target strand, termed *Sp*Cas9-T-ssODN, or for the non-
target strand donor, termed *As*Cas12a-NT-ssODN. Note that sequences identical to the PAM for
*As*Cas12a and ~18 nt of the protospacer region were retained within the 5'-HA of *As*Cas12a-T-
ssODN (Fig 1B and C).

Absence of *trans* activity of activated *As*Cas12a against the ssODN donors

Previous reports dealing with *Lachnospiraceae bacterium* ND2006 and *Francisella novicida* *As*Cas12a describe indiscriminate *trans*-activity unleashed on ssDNA after gRNA activation and target binding of *As*Cas12a (Chen *et al.*, 2018; Gier *et al*, 2020). Given that our goal here was to compare *Sp*Cas9 and *As*Cas12a in schistosomes, and given that we used an orthologous *As*Cas12a from *Acidaminococcus* sp. BV3L6 (IDT), in both KO and KI assays, we sought to confirm whether the omega-1 target gene activated *As*Cas12a might exhibit also *trans*-activity against the donor single stranded templates that we planned to use in KI assays, *As*Cas12a-T-ssODN and *As*Cas12a-NT-ssODN, in addition to RNA-guided double stranded DNA cleavage activity.

Accordingly, we also carried out *in vitro As*Cas12a nuclease activity assays using RNPs targeting a 426 bp amplicon of the exon 1 of *ω1* (Fig A, 2B and 2C). These assays confirmed that the crRNA was highly active and following incubation at 37°C for 30 min had cleaved the double stranded amplicon at the programmed cleavage site into two fragments of 265 and 161 bp, as expected (Fig 2B; Agilent Bioanalyzer electropherogram). In contrast, the activated *As*Cas12a-RNP complex failed to digest either of the single stranded templates, ssODNs *As*Cas12a-T-ssODN or *As*Cas12a-NT-ssODN after 120 min at 37°C (Fig 2C). As seen in the electropherogram, the ssODNs remained intact at 124 nt. Single stranded DNA standards ranging in size from <40 to 200 nt were included in the assay (Fig 2C). In summary, the findings suggested that *omega-1* target gene activated *As*Cas12a would not exhibit indiscriminate *trans*-nuclease activity against the ssODNs within schistosome cells, although an *in vitro* surrogate assay cannot faithfully mimic conditions within a viable schistosome egg.

Quantification of transgene copy number by external standard curve

The external standard curve design that we employed to estimate transgene copy number in genomics DNA of the schistosome eggs (Fig 2E and F). The standard curve on simple linear regression (y = -4.0964ln[10] + 44.821, R² = 0.9923) was established with logarithm-10 transformed initial DNA input, plasmid pCR4, which encodes the six stop codon cassette (Ittiprasert *et al.*, 2019), as the dependent and the Ct value from the qPCRs as the independent variable (not shown). We estimated the transgene copy number on the 5' and 3' of KI site in *ω1* by converting the measured Ct values to log copy numbers using the equations obtained from the standard curves. Subsequently, dividing the means by the antilog of each result, we arrived at an estimate of the transgene copy number. A 790 bp product amplified for the control wildtype sequence and amplicons of 235 bp and 182 bp for the 5' and 3' flanking regions, respectively, of the KI genomic DNA containing the six stop codon transgene were analyzed, which revealed 3'-side transgene integration values of 3.61±0.25, 4.44±0.84 and 4.19±1.22 from the KI_*Sp*Cas9, KI_*As*Cas12a-NT-ssODN and KI_*As*Cas12a-T-ssODN treatment groups, respectively (P = 0.4057, = 0.6264 and = 0.9155; NS; one-way ANOVA, multiple comparisons, 95% CI for differences of the means) (Fig 2E). However, the presence of asymmetrical HDR was revealed by significant differences among the copy number values for the 5' side transgene integration estimates: 2.72±0.42, 5.06±0.54, and 1.64±0.39 for the KI_*Sp*Cas9, KI_*As*Cas12a-NT-ssODN and KI_*As*Cas12a-T-ssODN groups, respectively (Fig 2F; P = 0.0214, P=0.0001 and $P \leq 0.0001$, *,***, ****, respectively; one-way ANOVA, multiple comparisons).

Estimation of CRISPR efficiency of *Sp*Cas9 and *As*Cas12a

RNP complexes of *Sp*Cas9-sgRNA and *As*Cas12a-CRISPR RNA (crRNA) targeting *ω1* were delivered to the LE by electroporation (Ittiprasert *et al.*, 2019). The efficiency of programmed mutation was assessed by TIDE analysis of Sanger sequence chromatograms (Gier *et al.*, 2020; Jacobsen *et al*, 2020), followed by targeted amplicon NGS and analysis of the reads using CRISPResso2 (Clement *et al*, 2019; Pinello *et al*, 2016). Twelve independent biological replicates were carried out by TIDE analysis. *As*Cas12a programmed KO led to significantly more indels than were induced with *Sp*Cas9 ($P < 0.001$, paired, two-tailed $t = 7.450$, df = 11) (Fig 2G). The mean CRISPR efficiency for KO with *Sp*Cas9 was 0.89%±0.43 (range, 0.30-1.31%) and 1.83%±0.29 (range, 1.39-2.54%) with *As*Cas12a, when compared with the control (mock) groups. Programmed mutation by *Sp*Cas9 was more efficient in the presence of donor ssODN, i.e., KI *versus* KO groups, 1.43±0.4, range 0.51-2.31% (Fig 2G and H). By contrast, differences in mutation efficiency were not apparent between the KO and KI groups using *As*Cas12a; 1.75%±0.63 (range 1.11-2.69%) and 1.54%±0.49 (range 1.82-2.20%) for *As*Cas12a-NT ssODN and *As*Cas12a-T ssODN, respectively. For deeper analysis, both on target KO and KI efficiency also was investigated by CRISPResso2 from pooled samples by target amplicon reads by MiSeq 2×250 bp configuration (Illumina).

Estimates of indels from NHEJ and/or HDR using NGS reads from targeted amplicons

Pooled gDNAs (above) also were used as templates for targeted amplicons for NGS and CRISPResso2 employed to assess the efficiency of transgene KI from HDR in the NGS reads. Comparison of the mutations among all four copies of the *ω1* gene also were undertaken. From whereas TIDE analysis using Sanger direct sequencing chromatograms was used above to estimate % efficiency of CRISPR manipulations in individual biological replicates, here targeted amplicons from pools of gDNAs from the biological replicates was used to acquire libraries of reads (Amplicon EZ, Genewiz). More than 150,000 reads from each NGS library were obtained. Analysis of the reads using CRISPResso2 provided increased sensitivity of detection of programmed gene editing than TIDE, as expected (Sentmanat *et al*, 2018). Specifically, there were 12.26%, 1.66%, 0.12% modified sequences (NHEJ) for KO-*Sp*Cas9 and 9.27%, 1.33% and 0.28% of NHEJ for KO-*As*Cas12a, when compared with the WormBase Parasite reference sequences (identical amplicon from Smp_334170.1, Smp_334170.2, Smp_334240), Smp_333930 and Smp_333870, respectively (Fig 3A and B). A small number of reads were assessed to be ambiguous by CRISPResso2, 0.49% and 0.93% for KO-*Sp*Cas9 and KO-*As*Cas12a, respectively (Fig 3A and B). Deletions were the only type of indel detected, for both *Sp*Cas9 and *As*Cas12a. In like fashion to other studies (An *et al*, 2020; Dumeau *et al*, 2019; Kissling *et al*, 2019), deletion of large sizes resulted from programmed *As*Cas12a editing (from -2 to 26 bp deletion size) at the programmed DSB, compared to -3 bp at the DSB programmed from *Sp*Cas9 (Fig 3C, D provide representative allele sequences). Both enzymes showed a preference to mutate the reference Smp_334170 copy, and also Smp_334240, rather than the Smp_333930 and/or Smp_333870 copies of omega-1. The numbers of indels arising from NHEJ with *Sp*Cas9 and with *As*Cas12a both increased in the presence of a ssODN donor repair template. The total NHEJ were 15.67%, 28.71% and 21.43% from KI-*Sp*Cas9-NT-ssODN, KI-*As*Cas12a-NT-ssODN and KI-*As*Cas12a-T-ssODN, respectively (Fig 4A, B and C). Deletion size as determined by CRISPResso2 for the KI-*Sp*Cas9 group was larger than for KO-*Sp*Cas9,

increasing up to 13 bp length deletions. Similarly, base deletion size from KI-*As*Cas12a was
larger than KO-*As*Cas12a with deletion of 42 bp detected (Fig 4D, E and F). Levels of
ambiguous results were estimated at 1.66%, 1.03% and 0.73% from KI-*Sp*Cas9, KI-*As*Cas12a-
NT-ssODN and KI-*As*Cas12a-T-ssODN, respectively.

To estimate the HDR events, we included the donor sequence as a parameter in the
CRISPResso2 analysis. More HDR events mediated by CRISPR/*Sp*Cas9 were detected than with
CRISPR/*As*Cas12a; 17.07% *vs* 14.56% and 12.37%, in like fashion to our earlier findings
(Santos *et al*, 2021). However, in *Sp*Cas9 sample, more imperfect HDR was seen (0.94%) than
in the *As*Cas12a groups (0.41% and 0.68%) (Fig 4A, B, C, F, G and H). Frequently, imperfect
HDR event alleles from both Cas nucleases were missing several residues (2-5 nt) on the 5' side
of the programmed DSB. We note that this mutation repair outcome may have impacted
quantification of the 5' KI transgene copy number, especially in KI-*As*Cas12a-T-ssODN samples
which contained 5 nt imperfect HDR repair on the six stop codon cassette (primer probing
sequence). Imperfect HDR at the 5' side might result from synthesis-dependent strand annealing
(Straume *et al*, 2020).

Transcription of ω1 interrupted by programmed mutation by *Sp*Cas9 and *As*Cas12a

Primers specific for the four annotated copies of *ω1* were employed to investigate transcription
by real time quantitative PCR (Fig S1C). Relative expression levels of *ω1* mRNA in negative
controls and experimental KO or KI eggs were compared with wild type. Significantly reduced
levels of *ω1* were seen in eggs following the experimental the CRISPR manipulations for gene
KO and KI. Transcription of *ω1* was reduced, as follows: 63.76%±5.57, 76.88%±11.33,
71.81%±10.91, 98.69%±0.44 and 80.04%±0.09 in the KO_*Sp*Cas9, KI_*Sp*Cas9, KO_*As*Cas12a,
KI_*As*Cas12a-NT-ssODN and *As*Cas12a-T-ssODN groups, respectively. Notably, LE subjected
to KI_*As*Cas12a_NT-ssODN retained <2% ω1 transcript levels of the mock control (Fig 5A),
and the level in the KI_*As*Cas12a-NT-ssODN group was significantly lower than the
KI_*As*Cas12a-T-ssODN group, KI_*Sp*Cas9-ssODN. Significant differences were not apparent
between the KO_*Sp*Cas9 and KO_*As*Cas12a groups, generally. There were no statistic
differences in levels of transcription among the negative control groups (Fig 5A). Differences
among treatment groups were assessed using ANOVA with multiple comparisons ($P \leq 0.01$ to $\leq$
0.0001; n= 4; F = 11.77).

Protein levels ascertained by western blotting

An anti-rω1 rabbit IgG was used to investigate levels of omega-1 in lysates (soluble egg antigen;
SEA) of LE. SEA (200 ng) from the three controls, i.e., the mock, enzyme only, and mixed
sgRNAs only groups, and from the KO and KI groups with *Sp*Cas9 or *As*Cas12a was size
fractionated by SDS-PAGE and transferred to PVDF and probed with the rabbit antibody (Fig
5B). Following densitometric analysis of the resulting signals, the levels of omega-1 remaining
in the SEA were estimated at 64%±11.65, 43.88%±6.02, 49.27%±4.19, 37.9%±2.28 and
52.16%±10 in the KO_*Sp*Cas9, KI_*Sp*Cas9, KO_*As*Cas12a, KI_*As*Cas12a-NT-ssODN and
KI_*As*Cas12a-T-ssODN groups, respectively. The differential intensity of ω1expression was not
diminished among the sgRNA only groups (Fig 5B and C). By contrast, ω1 protein levels were
markedly diminished among all the experimental treatment groups when compared with the

controls ($P \leq 0.0001$) (ANOVA, F = 27.05 , $n = 4$). Significant differences among for levels of
ω1 among the KO and KI SEA samples were not apparent (Fig 5C).

Discussion

Given the central role of the egg in disease and in the transmission of schistosomiasis (Costain *et*
al, 2018), and given the potential of CRISPR/Cas gene editing to advance understanding of this
and other neglected tropical disease pathogens, access to omega-1-negative eggs will provide
information to confirm the role of this ribonuclease in disease transmission. Functional knockout
(KO) of the *omega-1* gene and the resulting immunologically impaired phenotype have
showcased the novel application of CRISPR/*Sp*Cas9 and its utility for functional genomics in
schistosomes (Hsu *et al*, 2014; Jinek *et al.*, 2012; McVeigh & Maule, 2019; Sankaranarayanan *et*
al., 2021). In view of this progress, and the demonstrated tractability of *Sp*Cas9-catalyzed KO of
the *omega-1* gene, here we compared the performance of a second RNA-guided Cas nuclease,
the *As*Cas12a nuclease of *Acidaminococcus* sp., in terms of efficiency and precision with those
for *Sp*Cas9. We now report that *As*Cas12a and *Sp*Cas9 both provide tractable routes for RNA-
guided programmed mutation of the genome of the schistosome egg.

We targeted the well-known omega-1 extracellular T2 ribonuclease secreted by the mature egg
of *S. mansoni*, which is a key modulator of pathogenesis and transmission (Costain *et al.*, 2018;
Everts *et al*, 2009; Schwartz & Fallon, 2018; Steinfelder *et al*, 2009). Guide RNAs for both
*Sp*Cas9 and *As*Cas12a were designed to anneal to protospacer regions that are essentially at the
same target site; the programmed DSB were separated by only three nucleotide residues, on the
non-target strand of the target gene. The optimal protospacer adjacent motif for *As*Cas12a, TTTV
(Swarts & Jinek, 2019; Zetsche *et al*, 2015), is expected to occur abundantly in the AT-rich
genome of *S. mansoni* (Protasio et al, 2012). Programmed DSB catalyzed by *As*Cas12a results in
a staggered strand break of the target gene sequence, which can enhance efficiency of homology
directed repair (HDR) (Swarts & Jinek, 2019). For RNPs assembled from the nuclease and
cognate crRNA, *As*Cas12a was significantly more efficient for gene knockout than *Sp*Cas9, as
assessed by TIDE, presumably reflecting non-homology end joining (NHEJ)-catalyzed repair
and gene silencing.

We also investigated HDR-mediated gene editing with *Sp*Cas9 and *As*Cas12a delivered by
electroporation in the presence of the RNPs and donor repair templates. Each donor ssODN
template for *Sp*Cas9- and *As*Cas12a-programmed KI included a transgene cassette of 24 nt in
length encoding six stop codons (Ittiprasert *et al.*, 2019) but in which the homology arms
differed slightly on the non-target strand due to the discrete positions of the predicted DSB for
each nuclease. For programmed KI catalyzed by CRISPR/*As*Cas12a, we also included as the
donor template, the reverse complement sequence of the six stop codon cassette flanked by the
target stand sequence as the homology arms. This additional donor was included in our analysis,
given concern with respect to the known indiscriminate t*rans*-activity by activated *As*Cas12a
against single stranded DNA(Chen *et al.*, 2018), and especially for the ssODN with NT
homology arms which retains the PAM and partial (~18 nt) protospacer sequence (Swarts, 2019;
Swarts & Jinek, 2018, 2019)

RNA-guided *Sp*Cas9 cleavage led to higher levels (~4% higher) of precise knock-out compared
to that obtained with *As*Cas12a but markedly less precise knock-in (~6-13%). Deletions were

the frequent mutation type seen with both *Sp*Cas9 (deletions of 3 nt in length) and *As*Cas12a
(deletions of 2 to 26 nt in length) due to NHEJ repair. Using conditions optimized previously for
*Sp*Cas9 KI (Ittiprasert *et al.*, 2019), CRISPR/*Sp*Cas9 induced higher (~3%) HDR levels than
CRISPR/*As*Cas12a to insert the transgene for both *As*Cas12a-NT-ssODN and *As*Cas12a-T-
ssODN donors. Both *Sp*Cas9-and *As*Cas12a-edited eggs exhibited reduced levels of transcription
and expression of omega-1. A spectrum of chromosomal alleles was apparent following
programmed knock-in of the ssODN repair templates. Both target strand and non-target strand
donors were provided. Whereas accurate HDR was catalyzed by both *Sp*Cas9 and *As*Cas12a,
precise repair accounted for only a minority of the payload integrations. In some cases, the two
sides of the DSB programmed by *As*Cas12a were repaired by different pathways since mutant
alleles that included only the left or only the right side of the transgene were observed (Canaj *et
al*, 2019). Moreover, accurate HDR was significantly more frequent in the presence of the non-
target strand compared with target strand donor.

These findings may have been expected based on reports with other cells and species that
indicate that integration of a repair template is not always precise (Paix *et al*, 2017; Ranawakage
et al, 2020). In addition, *Sp*Cas9 may lead to higher levels of HDR correction than *As*Cas12a in
some contexts including in HEK293T and pluripotent stem cells where mis-integrations patterns
indicate that two sides of a DSB are repaired by separate cellular repair pathways including the
involvement of microhomology as a driver of the mis-integration (Santos *et al.*, 2021).

In addition to its guide RNA programmed gene editing and DSB activity, *As*Cas12a and related
Type 2 subtypes V and VI nucleases display target substrate bound, non-specific *trans*-activity
against ssDNA, dsDNA and/or RNA (Fuchs *et al*, 2019; Nguyen *et al*, 2020). Accordingly, we
speculated that activated *As*Cas12a associated with the target exon 1 site in *omega-1* may digest
the donor ssODN provided as the repair template. If so, investigation of comparative efficiency
of *As*Cas12a *versus* *Sp*Cas9 catalyzed knock-in would have been compromised. To investigate
this possibility, we performed a series of assays *in vitro* with both non-activated and activated
*As*Cas12a. Under the conditions and time frame of the reactions - commercially sourced
*As*Cas12a, 37°C, 100 mM NaCl, and 120 min reaction duration, we failed to detect degradation
of the ssODNs. Although our assays obviously cannot exactly reproduce the physiological
environment of the transfected schistosome eggs *in vivo*, we failed to detect indiscriminate *trans*-
activity against the ssODNs. However, lack of apparent *As*Cas12a trans-activity *in vitro* does not
preclude that indiscriminate *trans*-activity does not take place within the nucleus of the cells
within the egg. The RNP may not have been retained at the programmed target site, perhaps
degraded by cellular enzymes and/or displaced by the cellular repair machinery following
cleavage before *trans*-activity damaged the donor ssODN or other phenomena including
protection of cellular DNA by histones. Nonetheless, both *Sp*Cas9 and *As*Cas12a delivered
similar knock-in performance in these schistosome eggs.

Whereas enhanced mutation efficacy of *As*Cas12a over *Sp*Cas9 has been reported, the reverse
outcome also has been seen (Hur *et al*, 2016; Kim *et al*, 2016). Comparisons among studies are
not straightforward because of differences in species, target genes, delivery method, and
temperature, among other confounders. In addition, there are three widely used orthologues of
*As*Cas12a, *Fn*Cas12a from *Francisella novicida U112*, *Lb*Cas12a from *Lachnospiraceae
bacterium ND2006*, and *As*Cas12a from *Acidaminococcus* sp. *BV3L6* and these have been tested
in diverse eukaryotes (Mustafa & Makhawi, 2020). Other orthologues have also been studied

(Tóth *et al*, 2018; Zhang *et al*, 2021). *As*Cas12a in active at 37°C whereas other orthologues are
catalytically active at lower temperatures, which has enabled studies with zebrafish (Liu *et al*,
2019), insects, and maize (Zhang *et al.*, 2021), and environmental and diagnostic biosensing
(Chen *et al.*, 2018; Ma *et al*, 2020; Nguyen *et al*, 2021).

*As*Cas12a provided enhanced programmed knockout activity compared to *Sp*Cas9 at a specific
target site in the *omega-1* gene locus but both exhibited similar performance for HDR based
transgene insertion. Although these findings do not enable extrapolation to performance of
*As*Cas12a for CRISPR/Cas-based gene editing of other schistosome genes, they demonstrate that
Cas12a is active in schistosomes and expand the toolkit for functional genomics in helminth
parasites (Sankaranarayanan *et al.*, 2021; Zamanian & Andersen, 2016). Optimization of the
*As*Cas12a-based approach with regard to ratios of guide RNA:*As*Cas12a and/or ssODN, format
of donor(s), and/or temperature will be informative (Chen *et al.*, 2018; Hendel *et al*, 2015;
Swarts & Jinek, 2018). We anticipate that *As*Cas12a will find increasing use in functional
genomes for AT-rich genomes, including schistosomes and other platyhelminths (International
Helminth Genomes, 2019). Since transgene insertion into the zygote within the newly released
egg has already been described and shown to enable establishment of lines of transgenic
schistosomes, advances with gene editing approaches to this key developmental stage of the
schistosome life cycle portend progress in the near future with informative gain-of-function lines
of transgenic schistosomes.

Materials and Methods

Reagent/Resource	Reference or Source	Identifier or Catalog Number
Experimental Models		
Schistosoma mansoni eggs	Schistosomiasis Resource Center, Biomedical Research Institute, Maryland, USA	
Recombinant DNA		
recombinant omega-1 protein	Dr. Ron Hokke	Gifted from Dr. Ron Hokke
Antibodies		
Polyclonal antibody against recombinant omega-1 protein	Research Resource Identifier (RRID)	AB_2889886
goat anti-rabbit IgG (H+L) horse radish peroxidase conjugate	Life Technologies, Thermo Fisher Sciences	Cat # 31460
Oligonucleotides and sequence-based reagents PCR primers	This study	Supplement Table S2. They all synthesized from IDT
ssODN	Ittiprasert *et al*., 2019 and this study	Ultramer DNA Oligos, Synthesized from IDT
synthetic guide RNA	Ittiprasert *et al*., 2019 and this study	Synthesized from IDT
Chemicals, enzymes and other reagents		
Alt-R® CRISPR-Sp HiFi Cas9 enzyme	Integrated DNA Technologies	1081060
Alt-R® CRISPR-AsCas12a enzyme	Integrated DNA Technologies	1081068
DNAzol® ES reagent	Molecular Research Center	DN127
RNAzol® RT Reagent	Molecular Research Center	RN190
Opti-MEM	Fisher Scientific	Gibco™ 31985062
DMEM	Fisher Scientific	88364
fetal bovine serum	Fisher Scientific	Gibco™ 26140079
100× Penicillin-Streptomycin-Fungizone	Fisher Scientific	SV30079.01
NucleoSpin® Gel and PCR clean up	Takara	740609
SsoAdvanced Universal SYBR Green Supermix	Bio-Rad	1725271
RIPA buffer	Millipore-Sigma	R0278-500ML
Bradford reagent	Bio-Rad	5000201
SDS-PAGE polyacrylamide gradient gels, 4-12% gel, w/v	Thermo Fisher Scientific	NP0322PK2
Maxima First Strand cDNA Synthesis Kit for RT-qPCR, with dsDNase	Thermo Fisher Scientific	K1672
Peroxidase substate; Clarity™ Western ECL Substrate	Bio-Rad	1705062S
GoTaq® Green Master Mix	Promega	M712
NEB buffer 2.1	NEB	B720S
Proteinase K	Thermo Fisher Scientific	EO0491
Agilent DNA 1000 Kit	Agilent Technologies	5067-1505
Agilent small RNA kit	Agilent Technologies	5067-1548

Software		
CHOPCHOP	http://chopchop.cbu.uib.no/	
TIDE	https://tide.deskgen.com/	
CRISPREsso2	https://crispresso.pinellolab.partners.org/submission	
Snap Gene	https://www.snapgene.com/	Academic Subscription Plan
GraphPad Prism	Prism software	Monthly Subscription Plan
Other		
CFX96 Touch Real-Time PCR	Bio-Rad	
ElectroSquarePorator	BTX	ECM830
Chemi-Doc MP System	Bio-Rad	
BioAnalyzer	Agilent Technology	

Ethics statement

Mice experimentally infected with *S. mansoni*, obtained from Schistosomiasis Resource Center
(SRC) at the Biomedical Research Institute (BRI), MD were housed at the Animal Research
Facility of the George Washington University (GWU), which is accredited by the American
Association for Accreditation of Laboratory Animal Care (AAALAC no. 000347) and has an
Animal Welfare Assurance on file with the National Institutes of Health, Office of Laboratory
Animal Welfare, OLAW assurance number A3205-01. All procedures employed were consistent
with the Guide for the Care and Use of Laboratory Animals. The Institutional Animal Care and
Use Committee (IACUC) at GWU approved the protocol used for maintenance of mice and
recovery of schistosomes. Swiss albino mice were euthanized seven weeks after infection with *S.
mansoni*, livers were removed at necropsy, and schistosome eggs recovered from the livers
(Dalton *et al*, 1997). The 'liver eggs, termed LE' population is expected to include eggs of a
spectrum of age ranging from newly released by the female schistosome through to mature eggs
containing a fully developed miracidium (Jurberg *et al*, 2009). LE were maintained in DMEM
medium supplemented with 20% heat-inactivated fetal bovine serum (FBS), 2%
streptomycin/penicillin at 37°C under 5% CO_2 in air for 18-24 hours before use in gene editing
assays (below) (Dalton *et al.*, 1997; Ittiprasert *et al.*, 2019; Lu *et al*, 2021; Mann *et al*, 2010;
Mann *et al*, 2014).

CRISPR/Cas target design and single stranded DNA donors

The *ω1*-gene specific gRNA for *Sp*Cas9 and *As*Cas12a were designed to target all four copies of
the hepatotoxic ribonuclease *omega-1* gene. All four are located on *S. mansoni* genome update
annotation of version 7 (updated 16-September-2021 on WormBase ParaSite,
https://parasite.wormbase.org/Multi/Search/Results?species=all;idx=;q=Hepatotoxic%20ribonucl
ease%20omega-1;site=ensemblunit&filter_species=*Schistosoma_mansoni*_prjea36577);
Smp_334170.1, Smp_334170.2, Smp_334240 and Smp_333930 by an online CRISPR design
tool, CHOCHOP (Labun *et al*, 2016; Labun *et al*, 2019; Montague *et al*, 2014). There are 99-
100% sequence and coding sequence identity among those copies (Fig S1A), which encoding an
enzyme of 115amino acids in length, respectively. These *ω1* copies are located on chromosome
1 at nucleotide (nt) positions 3980364 to 3984675, 3992964 to 3995246, and 3908953 to
3911250, respectively. The guide RNA (gRNA), which targets exon 1 of *ω1* complimentary to
the catalytic active site histidine [within FRKHEFEKHGLCAVEDPQV] codons, for the gRNAs

programming both *Sp*Cas9 and *As*Cas12a. In addition, the gRNAs exhibited similarly high
predicted CRISPR efficiency scores along with an absence of predicted off-target cleavages, as
predicted using the CHOPCHOP algorithm using Smp_334170.2 as template (Fig 1A and B).
Specifically, the sequence of the gRNA used with *Sp*Cas9 was 5'-
GCATGGTTTGTGTGCAGTTG-3', 20 nt in length and that for *As*Cas12a was 5'-
AAAAGCATGGTTTGTGTGCA -3', 20 nt on exon 1 of Smp_334170.2, Smp_334240 and
Smp_333930 or exon 4 of Smp_334170.1 (Fig S1A) The expected cut sites of two nucleases are
discrete, being 3 and 2 nt apart from each other on the non-target (NT) and target (T) strands,
respectively (Fig 1A).

Single stranded oligonucleotide donors (ssODN) were designed to insert the 24 bp encoding six-
stop codons into the programmed cleavage target. The homology arms used in the ssODN donor
with *Sp*Cas9 was the non-target strand flanking both the 3' and 5' sides of the blunt ended DSB,
termed here *Sp*Cas9-ssODN. The homology arms of the two donor ssODNs used with *As*Cas12a
(cut at 18 nt and 23 nt on NT and T strands, leading to a sticky ended DSB) were either the target
stand or the non-target strand sequences (Fig 1B and C), here termed *As*Cas12a-NT-ssODN and
*As*Cas12a-T-ssODN, respectively. The complementary reverse of the 24 nt stop codon cassette
was used in the *As*Cas12a-T-ssODN donor, aiming for HDR insertion of the stop codons in the
translated forward strand of the gene (Fig 1C).

CRISPR reagents

Alt-R® CRISPR-*Sp*Cas9 and Alt-R® CRISPR-*As*Cas12a reagents (Alt-R® S.p. HiFi Cas9
nuclease V3, synthetic gRNA (sgRNA) (Labun *et al.*; Labun *et al.*; Nishimasu *et al.*), Alt-R® As
*As*Cas12a V3 and crRNA), and ssODN HDR donor templates (Ultramer DNA oligos) and Alt-
R® *Sp*Cas9 or *As*Cas12a Electroporation enhancer buffers were from Integrated DNA
Technologies (IDT) (Coralville, IA). Figure 1C provides the nucleotide sequences of the gRNAs
and donor ssODNs.

Ribonucleoprotein (RNP) assembly and delivery by electroporation

Stock solutions of the gRNAs for use with *Sp*Cas9 and *As*Cas12a and of the ssODNs were
prepared at 1.0 µg/µl in Opti-MEM medium (Sigma, St. Louis, MO). Each gRNA was
combined with the respective nuclease at 1:1 ratio to form an RNP complex as described
(Ittiprasert *et al.*, 2019). The mixture was incubated at room temperature for 15 min, after which
3 µl of Alt-R® *Sp*Cas9 or *As*Cas12a electroporation enhancer was added. The mixture was
transferred to a chilled 2 mm electroporation cuvette (BTX), containing ~ 5,000 LE in 100 µl
Opti-MEM. LE in the presence of RNPs and the other reagents were subjected to square wave
electroporation with a single 125 volts, 1 pulse for 20 ms (ElectroSquarePorator, ECM830, BTX
San Diego, CA). In KI assays, 6 µg of ssODN was included in the cuvette before
electroporation. Following delivery of the RNPs, LE were maintained for 10 days at 37°C, 5%
CO_2 in air. In addition to experimental gene-edited groups, control groups included LE
electroporated in Opti-MEM (mock), mixed-Cas enzymes only, mixed-sgRNAs only, and
mixed-ssODNs only. Experimental and control groups were otherwise treated similarly.

PCR to detect a multiple stop codon containing transgene

Genomic DNAs were extracted from LE using the DNAzol® RT reagent (Molecular Research
Center Inc, Cincinnati, OH), and the concentration and purity determined by spectrophotometry
(Nanodrop 1000, Thermo Fisher Scientific Inc., Waltham, MA). The HDR target site(s) on
either the 5' or 3' KI transgene was amplified independently with two pairs of primers (Fig 2A):
1) 5'KI: 5'int-F and 24nt-6stp-R primers and 2) 3'KI: 24nt-6stp-F and 3'int-R using the GoTaq
DNA polymerase mix (Promega, Madison, WI) and 200 nM of each primer. The thermal cycling
included denaturation at 95°C, 3 min followed by 30 cycles of 94°C, 30s, 56°C, 30s and 72°C,
30s and a final extension for 5 min at 72°C. Thereafter, the amplicons (10 μl) were size separated
by agarose gel electrophoresis, stained with ethidium bromide, and imaged using the ChemiDoc
Imaging System (Bio-Rad, Hercules, CA). The expected sizes for the 5'KI and 3'KI amplicons
were 235 and 182 bp, respectively (Fig 2D). A positive control PCR to amplify ~790 bp flanking
the DSB was included.

Real time qPCR to estimate transgene copy numbers

A SYBR green quantitative PCR-based approach was used to estimate transgene copy numbers
using two pairs of primers; 5'KI and 3'KI as detailed (Fig 2A and D) with the SsoAdvanced
Universal SYBR Green Supermix (Bio-Rad). Briefly, one ng of genomic DNA (gDNA) was
amplified with 10 µl of 2× SsoAdvanced Universal SYBR Green Supermix and 200 nM of each
primer, using the CFX Connect Real-Time PCR Detection System (Bio-Rad). PCR conditions
included denaturation at 95°C, 3 min followed by 30 cycles of 94°C, 30 s, 56°C, 30 s and 72°C,
30 s and a final extension at 72°C for 5 min, with the signals collected at the annealing step in
each cycle. Analysis of transgene copy number in the pooled gDNAs from LE involved
comparison of the qPCR signals obtained using a 10-fold serial dilution series of known
quantities of the 24 nt six-stop codon transgene in 235 bp (5' integration PCR fragment) ligated-
pCR4-TOPO plasmid DNA (size 4,191 bp) to establish a standard curve (Shirvani *et al*, 2020).
The standard curve was based on a linear regression generated using logarithm-10 transformed
initial DNA input as the dependent variable and the Ct number of the qPCR signal as the
independent variable (Ct values in each dilution were measured in triplicate). Ct numbers were
used to fit into the linear regression to derive the estimation of transformed DNA amount for the
transgene (Lee *et al*, 2006; Yuan *et al*, 2007). The concentration of the 5'KI-transgene ligated-
pCR4-TOPO plasmid was measured (NanoDrop 1000) and the corresponding copy number was
calculated as follows:

$$\text{DNA copy number} = \frac{6.02 \times 10^{23} \,(\text{copy/mol}) \times \text{DNA quantity (g)}}{\text{DNA length (bp)} \times 660 \,(\text{g/mol/bp})}$$

466 Data were collected from four biological replicates. 5'KI and 3'KI transgene copy numbers
467 from the *Sp*Cas9 and *As*Cas12a treatments were plotted using GraphPad Prism v9 (Fig 2E and
468 F).

Tracking of INDELs by decomposition (TIDE) analysis

Genomic DNAs from LE exposed to the RNPs were isolated as above. The target *omega-1* locus
in the replicates of gDNAs was amplified using a control primer pair (Fig 2A). The amplicon of
790 bp in size from control and experimental groups, both KO and KI focused manipulations,
was purified by NucleoSpin® Gel and PCR clean up (Macherey-Nagel, Bethlehem, PA) after

which the nucleotide sequence determined by the Sanger method (Genewiz, South Plainland,
NJ). Sanger sequence traces from the 12 independent replicates of the experimental groups for
both *Sp*Cas9 or *As*Cas12a- (Jacobsen *et al.*, 2020) and *Sp*Cas9-catalyzed gene editing
manipulations were analyzed and compared with reads from the control (mock) group using the
Tracking of Indels by Decomposition (TIDE) algorithm (Brinkman *et al*, 2014; Brinkman & van
Steensel, 2019; Meshalkina *et al*, 2020). TIDE provides a rapid and informative assay and also
inform decisions on whether proceed to more detailed high throughput, next generation
sequencing (NGS) of targeting amplicons (Germini *et al*, 2018).

On-target amplicon next-generation sequencing

Subsequently, the Amplicon-EZ next generation sequencing approach (Genewiz) was used for
deeper coverage of the $\omega1$ coding sequence mutations and provided > 50,000 reads per sample.
Amplicons of 426 bp obtained with the primers bearing Illumina partial adapter sequences at the
5' end - NGS-F 5'- TATTGTCAACGGCGTACAGG-3' and NGS-R 5'-
CAATCGGCACTGAGACGCA-3' - flanking the programmed DSBs for CRISPR/*Sp*Cas9 and
CRISPR/*As*Cas12a were sequenced using Illumina chemistry. Partial Illumina adapters were
added to 5' end of the forward and reverse primers forward sequencing read: 5'-
ACACTCTTTCCCTACACGACGCTCTTCCGATCT-3', reverse sequencing read: 5'-
GACTGGAGTTCAGACGTGTGCTCTTCCGATCT-3'. The amplicons were purified using the
NucleoSpin® Gel and PCR clean up kits from Macherey-Nagel (Bethlehem, PA), after which
quantification and qualification of purified band(s) were undertaken using the Agilent DNA 1000
kit and BioAnalyzer (Agilent). Illumina NGS was performed with 2×250 bp sequencing without
fragmenting the amplicons. Raw reads in FASTQ format were analyzed for mutations resulting
from NHEJ and HDR by CRISPREsso2 (Clement *et al.*, 2019) using the reference $\omega1$ sequences,
Smp_334170 and Smp_333930 including Smp_333870 (this latter is annotated as an
uncharacterized protein but shares >99% identity) (Fig S1B). The default parameters were used
to estimate indel and HDR events. The sequence reads from the Amplicon-EZ runs are available
at GenBank Bioproject PRJNA415471, BioSample SAMN07823308, SRA study SRP126685,
accessions SRR: 13498338-13498342, 15971309-15971320, and 6374209-6374210.

Quantitative real time PCR

Total RNA of LE was isolated using the RNAzol® RT reagent (Molecular Research Center)
according to the manufacturer's instruction. RNA concentration and purity were determined
(Nanodrop 1000 spectrophotometer, Thermo Fisher). Ten nanograms of RNA was treated with
DNase I to remove residual DNA then converted to cDNA by Maxima First Strand cDNA
Synthesis Kit (Thermo Fisher Scientific). The $\omega1$-specific product was amplified using
SsoAdvanced Universal SYBR Green Supermix (Bio-Rad) on CFX96 real time PCR (Bio-Rad)
using the primers as described (Ittiprasert *et al.*, 2019), $\omega1$-RT-F: 5'-
GGTTACTTTAAATTTGGTA-3' and $\omega1$-RT-R; 5'-AGTTCCAAGGAACGGGCAG-3'.
These primers amplify all annotated $\omega1$ transcript copies; Smp_334170.1, Smp_334170.2,
Smp_334240 and Smp_333930 (Fig S1C). The PCR reaction was denatured at 95°C for 30 s, 40
amplification cycles each consisting of denaturation at 95°C for 15 s and annealing/extension at
60°C for 30 s. The output was analyzed in CFX manager software (Bio-Rad). Relative
expression of $\omega1$ was calculated using the $2^{-\Delta\Delta Ct}$ method and normalized to schistosome GAPDH
(Smp_0569701) expression using specific primers; forward 5'-

ATGGGACATTTCCAGGCGAG-3', reverse 5'-CCAACAACGAACATGGGTGC-3' (Livak &
Schmittgen, 2001). Data four biological replicates were compared to wild type LE and fold
change reported as mean ± SE ($n = 4$).

Western blot analysis

Soluble egg antigen (SEA) of LE was extracted by mechanical lysis (Funakoshi motorized
pestle; Diagnocine, Hackensack, NJ) in RIPA buffer (Millipore Sigma) supplemented with
protease inhibitors (Protease Inhibitor Cocktail I, Millipore Sigma), as described (Ittiprasert *et*
al., 2019). The lysate was clarified by centrifugation at 13,000 rpm, 15 min, 4°C after which
protein concentration was determined by the Bradford method (Kielkopf *et al*, 2020). Ten
micrograms of SEA were size-separated through SDS-PAGE polyacrylamide gradient gels, 4-
12% gel, w/v (Bolt Bis-Tris Plus, Invitrogen, Thermo Fisher Scientific) before transfer to PVDP
membranes (Transblot Turbo Transfer System, Bio-Rad). Polyclonal anti-ω1 IgG antibodies
were purified from the rabbit anti-ω1 sera, raised as follows. Two NZ white rabbits, E12983 and
E12984, were immunized with 30 μg of recombinant *omega-1* RNase T2 of *Schistosoma*
mansoni (rω1) (produced *in planta* by agroinfiltration of leaves of *Nicotiana benthamiana*
(Wilbers *et al*, 2017)) in Complete Freund's Adjuvant and boosted four times at two-week
intervals using 15 μg rω1 in Incomplete Freund's Adjuvant each time. Blood was collected 66
days after the first immunization, IgGs was isolated from sera using an immobilized protein A
ligand. The concentrations of the purified antibodies were 1.31 mg/ml (E12983) and 1.51 mg/ml
(E12984). These antibodies have been assigned Research Resource Identifier (RRID)
AB_2889886 (Bandrowski *et al*, 2016). PVDF membranes were probed with the E12983 IgG
antibody, diluted 1:1,000, for 18 h at 4°C in PBS-Tween (0.5%), 5% non-fat powdered milk,
washed 3 times in PBS-Tween (wash buffer), and probed with goat anti-rabbit IgG (H+L) horse
radish peroxidase conjugate (Life Technologies, Thermo Fisher Sciences), diluted 1:5,000, for
60 min at 23°C with gentle shaking. Following washing (3 times in wash buffer), the membrane
was exposed to a peroxidase substrate (Clarity Western ECL Substrate, Bio-Rad) for 5 min at
23°C, after which chemiluminescence was quantified and imaged (Chemi-Doc MP System, Bio-
Rad). The Image Lab software (Bio-Rad) was used to quantify the ω1 signals. The percentage of
ω1 expression from the experimental and control groups was established by comparison with
that of wild type SEA, assigned as 100%.

**Single stranded DNA donors as a potential substrate for indiscriminate *trans* activity of
activated *As*Cas12a**

Previous reports reveal the *trans*-activity of wild type *As*Cas12a prefers ssDNAs over dsDNAs,
and was inactive toward dsDNA in the presence of magnesium (II) ions (Li *et al*, 2020). The
*As*Cas12a exhibited not only endonuclease activity on both dsDNA, *cis*- and *trans*-activity (Li *et*
al, 2018b). Accordingly, to investigate whether activated *As*Cas12a could degrade the ssODN
donors used in this study by indiscriminate *trans*-activity, several *in vitro As*Cas12a-RNP
digestion assays were carried out. These included the dsDNAs containing the *As*Cas12a target
sequence used in this study and 5'PAM (TTTG) as positive enzymatic activity. A 426 bp
dsDNA template was amplified using the NGS study (above) primers. Activity of *As*Cas12a-
RNP against the dsDNA template released fragments of 265 and 161 bp (Fig 2B), at 37°C for
≥30 min as predicted in 1× NEB Buffer 2.1 (500 mM NaCl, 20 mM sodium acetate, 0.1 mM
EDTA, 0.1 mM tris(2-carboxyethyl) phosphine (TCEP), 50% glycerol). *As*Cas12a activity was

inactivated by the addition of Proteinase K (1 µl at 20 mg/ml) at 56°C for 10 min.
Electropherograms of the digestion products were obtained using the BioAnalyzer (Agilent
DNA1000) to visualize the size of templates and fragments.

To investigate indiscriminate *trans*-cleavage of ssODNs, similar *in vitro* digestion reaction was
carried out, as above, but with the addition of 50 ng of either *As*Cas12a-NT-ssODN or
*As*Cas12a-T-ssODN. A mixture of *As*Cas12a-RNP complex in 1× NEB Buffer 2.1 and ssODN
was incubated at 37°C for 30, 60 and 120 min before terminating the reaction by addition of
proteinase K at 56°C for 10 min, and visualized as above. The ssDNA ladder (SimPlex Sciences,
NJ) ranging from 10-200 bp was included as the single stranded DNA marker. *Trans*-activity of
Alt-R *As*Cas12a V3-RNP complex was not apparent against the ssODNs under these conditions
in vitro (Fig. 2C).

Statistical analysis

Statistical analysis was performed using GraphPad Prism version 9.1 (GraphPad Prism, La Jolla,
CA). Graphs were plotted as the mean with bars to show the standard error of the mean. The
differences among groups were assessed by one-way ANOVA. *P* values of ≤ 0.05 were the
statistically significant.

GenBank/EMBL/DDBJ

Database accessions

Nucleotide sequences reads of the amplicons are available at Bioproject
PRJNA415471, https://www.ncbi.nlm.nih.gov/bioproject/PRJNA415471 and GenBank
accession PRJNA415471, BioSample, SAMN07823308, SRA, SRP126685.

Acknowledgements

We thank Professor Cornelis Hokke, Leiden University Medical Center, The Netherlands for the
gift of recombinant *Schistosoma mansoni* omega-1. We thank the Thailand Research Fund, the
Royal Golden Jubilee Ph.D. program, and Dr. Dalina Tanyong, Mahidol University, Thailand,
for support for CC to perform the work at GWU. This research was funded in whole, or in part,
by the Wellcome Trust (award number 107475/Z/15/Z, titled 'Functional Genomics Flatworms
Initiative', K. H. Hoffmann, principal investigator). For the purpose of Open Access, the authors
have applied a CC BY public copyright license to any Author Accepted Manuscript version
arising from this submission. Schistosome-infected mice and snails were provided by the NIAID
Schistosomiasis Resource Center of the Biomedical Research Institute, Rockville, Maryland
through NIH-NIAID Contract HHSN272201000005I for distribution through BEI Resources.

Author contributions

WI, CC, and PJB designed the experiments, analyzed the data, and drafted the manuscript with
input from all the co-authors; CC and WI performed the gene editing focused experiments;
VHM, WL, AM, TO, YNA and MMK contributed the parasite materials to the experiments, to
data collection, and reviewed and edited the manuscript; PJB and WI supervised the project.
PJB, WI and MMK arranged the funding.

Conflict of interest

The authors declare that do not have competing interests.

Figure legends

Figure 1. Schematic of the *Schistosoma mansoni* hepatotoxic ribonuclease omega 1 (ω1)
gene structure, target sites for the *Sp*Cas9 and *As*Cas12a, ssODN donors and partial amino
acid sequence with histidine active site.

A   Diagram of the omega-1 (*ω1*) gene (Gene ID; Smp_334170.2 of *S. mansoni* annotated
genome version 7_2021) and CRISPR/Cas target sites on coding sequence of the first exon
coding for histidine active site, FRKHEFEKHGLCAVEDPQV. The relative positions of
crRNA for CRISPR-*Sp*Cas9 (red arrow) and -*As*Cas12a (blue arrow) are indicated along
with their 3'PAM (yellow arrow) and 5'PAM (yellow arrow), respectively.

B   Programmed cleavage site for *Sp*Cas9/sgRNA ~3nt upstream of PAM 3'TGG (red arrow);
and cleavage site for *As*Cas12a/crRNA at 18-24 nt downstream of a prospective 5-'TTTV
PAM (blue arrow). The programmed cleavage sites for both enzymes, *Sp*Cas9 and
*As*Cas12a, were three to six nt distant from each other. Fifty residues upstream and
downstream of the double strand break (DSB) were included as the homology arms (HA) of
the six stop codon cassette transgene (payload/cargo). Because cleavage by *Sp*Cas9
liberates blunt ended DNA, only a non-target strand (NT) repair template was provided for
HDR (red bar). Sticky ended DNA results from cleavage by *As*Cas12a and hence both non
target and target strand (T) repair templates for HDR were provided (light and dark blue
bars).

C   The DNA donor sequences of CRISPR/*Sp*Cas9 and CRISPR/*As*Cas12a with non-target
(NT) or target (T) sequence HA where the *As*Cas12a-NT-ssODN donor repairs the anti-
sense DNA and *As*Cas12a-T-ssODN donor repairs the sense stand.

Figure 2. Knock-in by homology directed repair.

A   The location of primers used for 5'- and 3'- transgene (purple lines) knock-in PCR (5'int-F
with 24nt-6stp-R and 24nt-6stp-F with 3'int-R) and internal positive control (control-F and
-R). The primers for next generation sequencing indicate as NGS-F and NGS-R. All
primers are located outside of the HA (purple boxes) sequences to avoid also positive
amplification from DNA donors. The 24 nt-6stp-F and -R primers are highly specific for
the stop codon transgene, and should not anneal to the wild type schistosome gene.

B   The *As*Cas12a target recognition activates specific 426 bp dsDNA amplicon template (lane
1) cleavage into 265bp and 161 bp (lane 3) products. The crRNA/*As*Cas12a RNP complex
of ~40 nt is seen in lanes 2 and 3.

C   No specific ssDNA cleavage was seen on the Cas12a-NT-ssODN or Cas12a-T-ssODN
donors at 124 nt as in lanes 1 and 2, respectively. The crRNA/*As*Cas12a RNP (~40 nt in
lanes 3-9) containing activated *As*Cas12a did not show indiscriminate ssDNA digestion at
30 min, 60 and 120 min in lanes 4-5, 6-7 and 8-9, respectively. Lanes 4, 6 and 8 show the

uncut 124 nt of Cas12a-NT-ssODN (top band) with Cas12a-crRNA (lower band), and lanes
5, 7 and 9 shows 124 nt of Cas12a-T-ssODN (top band) with Cas12a-crRNA (lower band).

D   The PCR detection of the integration sites in the transgenic eggs and control. The
amplified control product (lanes C) for the wildtype sequence was 790 bp in length while
those for the 5'(lanes 5') and 3' (lanes 3') flanking regions of the transgenic sequence were
235 bp and 182 bp due to KI of Cas-9 ssODN, Cas12a_NT-ssODN and Cas12a_T-ssODN
donor templates, respectively. Lane M is a 1 kb Plus DNA ladder; mock (only buffer
electroporation) served as the negative control for transgene integration.

E, F The estimation of transgene copy number in egg DNA were calculated by qPCR comparing
with known quantity of the 24 nt transgene in 235 bp (5' integration PCR fragment)
ligated-pCR4-TOPO plasmid DNA (size 4,191 bp) standard curve; $y = -4.0964\ln(10) +$
44.821, $R^2 = 0.9923$ (data not shown). There was the confirmation of 3' integration of
3.61 ± 0.25, 4.44 ± 0.84 and 4.19 ± 1.22 of transgene copy from KI_Cas9, KI_Cas12a_NT-
ssODN and KI_Cas12a_T-ssODN, respectively (one-way ANOVA, multiple comparisons,
95% CI of diff, P-value at 0.4057, 0.6264 and 0.9155, not significant). While asymmetric
HDR was revealed by 5' transgene integration of 2.72 ± 0.42, 5.06 ± 0.54 and 1.64 ± 0.39 copy
number from KI_Cas9, KI_Cas12a_NT-ssODN and KI_Cas12a_T-ssODN, respectively.
The asterisk symbols *,*** and **** indicate statistical significance at $P = 0.0214$,
$P=0.0001$ and $P < 0.0001$, respectively (one-way ANOVA, multiple comparison). The
control primers used for amplification from both the control and experimental groups,
followed by Sanger sequencing results, were monitored for efficiency of programmed
CRISPR editing firstly by TIDE followed by NGS and CRISPresso2 of the NGS reads.

G,H 5'int-F and 3'int-R are genome-specific primers, and 24nt-6stp-F and 24nt-6stp-R are
transgene-specific primers. From the findings, we confirmed that programmed
CRISPR/*As*Cas12a gene editing was active in schistosome eggs, and higher than for
*Sp*Cas9 by TIDE analysis (ANOVA, ****, P 0.0001, n = 12). The estimation of CRISPR
efficiency from the 12 biological replicates (each black dot in panels G and H) KI_Cas9
(wide range of CRISPR efficiency from 0.46-2.23 among 12 biological replicates) was
higher than for KO based treatments. The estimated CRISPR efficiency in schistosome
eggs (LE) from CRISPR/*As*Cas12a (green dots in panels G and H, $n = 8$) was at least 1.5%
in most samples. There were no statistically significant differences in CRISPR efficiency
among the KI_Cas9 vs. KI_Cas12a (panel H) from Sanger sequencing along with TIDE
analysis.

Figure 3. NHEJ assessment from *S. mansoni* eggs at the target *ω1* gene following delivery
of *Sp*Cas9 and *As*Cas12a RNP complexes by electroporation. CRISPResso2 analysis of the
Illumina sequence reads revealed non-homologous end joining (NHEJ) mutations at targeted loci
in schistosome eggs. Allele-specific editing outcomes of CRISPR-*Sp*Cas9 and -*As*Cas12a gene
editing in schistosome multiple copies of *ω1* Smp_334170.2, Smp_334170.1, Smp_334240
(100% gene sequence identical; Fig S1B). Mutagenesis frequencies for all loci including
Smp_333930 and Smp_333870 were included in CRISPResso2 analysis (Fig S1B). The results
were from NGS libraries of pooled genomic DNA from seven biological replicates of KO_Cas9
(Panel A and C) and KO_Cas12a (Panel B and D). The sequence reads assigned to reference
genes, Smp_333930 and Smp_333870 loci by the CRISPResso2 to achieve accurate

quantification of programmed mutation of the multiple copies of *ω1* from *Sp*Cas9 and *As*Cas12a.
The bar chart shows the assignment of each read to the wild type (unmodified) and NHEJ with
12.26%, 1.66% and 0.12% (total 14.04%) from KO_Cas9 on references and other two loci,
respectively (panel A). Panel B shows the assigned NHEJ with 9.27%, 1.33% and 0.28% (total
10.88%) from KO_Cas12a on references and two highly identical copies of *ω1* (panel B). The
majority of NHEJ indels from both nucleases were nucleotide substitutions followed by gene
deletions (Panel C and D). Mutations were induced in all copies of *ω1*; however, most editing
was seen on the reference target site. Only deletions were revealed in this experiment, 3 nt
deletion downstream from expected cut side (red arrow) was the majority of NHEJ mediated
from *Sp*Cas9 (Panel C). The larger deletion from 2 to 26 nt at downstream of expected sticky
end cut sites (red arrows) were revealed from NHEJ. CRISPResso2 interpreted ≤ 1.0% of the
reads as ambiguous. Ambiguous alignments: 0.49% and 0.93% that could not be attributed
uniquely to one of these loci are shown in the right-hand side bar of each graph.

Figure. 4. HDR and induction of NHEJ at the *ω1* target in the presence of donor ssODN
templates. Panels A to C show the percentage of NHEJ from schistosome eggs that introduced
RNP along with ssODN template were higher in the presence of ssODN donor templates than in
their absence. There were totals of 15.67% (13.22+1.16+0.52), 28.71% (22.31+6.17+0.23) and
21.43% (9.62+11.56+0.25) editing efficiency in the +KI_Cas9, KI_Cas12a_NT-ssODN and KI-
Cas12a_T-ssODN groups, respectively (blue or green bars in each graph). Most mutations were
substitutions and deletions located downstream of the programmed double stranded breaks
(DSB), with the DSBs indicated by red arrows (Panel D, E and F). A variety of sizes of base
deletions from 1 to 15 bp in length were seen in the KI_Cas9, KI_Cas12a_NT-ssODN groups
while large deletions, up to 40 bp, were evident as major deletions detected in the KI_Cas12a_T-
ssODN group. The percentage of perfect HDR in the three experimental groups, KI_Cas9,
KI_Cas12a_NT-ssODN and KI_Cas12a_T-ssODN were 17.07%, 14.58% and 12.37%,
respectively (red bars, Panel G, H and I). In addition, imperfect HDRs occurred in all KI
treatment groups, likely as the result of interruption of 5' end HDR gene repair by competing
NHEJ activity with the result that deletion of 2-5 nt were seen along with donor transgene
insertion (black bars in Panel G, H and I); the frequency of this event was 0.94, 0.41 and 0.68%
among the KI_Cas9, KI_Cas12a_NT-ssODN and KI_Cas12a_T-ssODN groups.

Figure 5. Reduction of the expression of ω1 as assessed by quantitative real time PCR and
immunoblot analysis.

A   Reduction of ω1 transcript levels from KO_Cas9, KI_Cas9, KO_Cas12a, KI_Cas12a_NT-
ssODN and KI_Cas12a_T-ssODN eggs compared with control groups after normalization
with wild type egg RNA (100% as discontinued line). The means represent four
independent biological replicates.

B, C Soluble egg antigen; SEA (in duplicates) was sized in SDS-PAGE gels, transferred to
PVDF and probed with anti-recombinant omega-1 (rω1) rabbit IgG. Levels of ω1 protein,
migrating at ~30 kDa, were reduced in all treatment groups. The percentage of ω1
expression in both KO and KI experimental groups were compared to ω1 levels in wild
type (WT) SEA (100%) (panel C) (one-way ANOVA, multiple comparison, *, $P \leq 0.05$,
***, $P \leq 0.001$, ****, $P \leq 0.0001$).

Supporting information

Figure S1. **Gene structure encoding the hepatotoxic ribonuclease, omega 1 *(ω1)*. At least**
five copies of the gene are located on chromosome 1, as apparent in the draft sequence of *S.*
***mansoni*, version 7, scaffold SM_V7_1:3,885,722-4,002,365.** The positions of the several *ω1*
genes of the schistosome genome (WormBase Parasite, September 2021), including two loci of
Smp_334170, Smp_334240, Smp_334070, Smp_3332220, Smp_333930, Smp_333870 and
Smp_179960 are shown to for their physical relationship of the copies on the chromosome.

A　　Gene structures and sizes of Smp_334170.1, Smp_334170.2 and Smp_334240 and the
CRISPR/Cas programmed target sites for *Sp*Cas9 (blue arrow) and *As*Cas12a (green).

B　　All gene copies of *omega-1* share >99% DNA sequence identity; DNA sequence alignment
of the *omega-1* orthologues and target amplicon primer indicated as NGS-F and NGS-R.
Four copies of the gene, Smp_334240, Smp_334170 and Smp_333930, are reported in
Uniprot.

C　　Alignment of mRNA sequences and primers used for gene transcript analysis (purple).

Table S1. The CRISPREsso2 analysis findings from four negative controls: mixed sgRNAs
only, *Sp*Cas9 only, *As*Cas12a only, and mixed donors only. There were over 180,000 reads from
each library. There was < 1% of background modified reads from all negative control samples.

Table S2. Nucleotide sequences of oligonucleotide primers used in the investigation.

References

An Y, Geng Y, Yao J, Fu C, Lu M, Wang C, Du J (2020) Efficient Genome Editing in Populus
Using CRISPR/Cas12a. *Front Plant Sci* 11: 593938
Arunsan P, Ittiprasert W, Smout MJ, Cochran CJ, Mann VH, Chaiyadet S, Karinshak SE, Sripa
B, Young ND, Sotillo J *et al* (2019) Programmed knockout mutation of liver fluke granulin
attenuates virulence of infection-induced hepatobiliary morbidity. *Elife* 8
Banakar R, Schubert M, Collingwood M, Vakulskas C, Eggenberger AL, Wang K (2020)
Comparison of CRISPR-Cas9/Cas12a Ribonucleoprotein Complexes for Genome Editing
Efficiency in the Rice Phytoene Desaturase (OsPDS) Gene. *Rice* 13: 4
Bandrowski A, Brush M, Grethe JS, Haendel MA, Kennedy DN, Hill S, Hof PR, Martone ME,
Pols M, Tan SC *et al* (2016) The Resource Identification Initiative: a cultural shift in publishing.
Brain Behav 6: e00417
Bolt BJ, Rodgers FH, Shafie M, Kersey PJ, Berriman M, Howe KL (2018) Using WormBase
ParaSite: An Integrated Platform for Exploring Helminth Genomic Data. *Methods Mol Biol*
1757: 471-491
Brinkman EK, Chen T, Amendola M, van Steensel B (2014) Easy quantitative assessment of
genome editing by sequence trace decomposition. *Nucleic Acids Res* 42: e168
Brinkman EK, van Steensel B (2019) Rapid Quantitative Evaluation of CRISPR Genome Editing
by TIDE and TIDER. *Methods Mol Biol* 1961: 29-44
Canaj H, Hussmann JA, Li H, Beckman KA, Goodrich L, Cho NH, Li YJ, Santos DA,
McGeever A, Stewart EM *et al* (2019) Deep profiling reveals substantial heterogeneity of
integration outcomes in CRISPR knock-in experiments. *bioRxiv*: 841098
Charpentier E (2015) CRISPR-Cas9: how research on a bacterial RNA-guided mechanism
opened new perspectives in biotechnology and biomedicine. *EMBO Mol Med* 7: 363-365
Charpentier E, Richter H, van der Oost J, White MF (2015) Biogenesis pathways of RNA guides
in archaeal and bacterial CRISPR-Cas adaptive immunity. *FEMS Microbiol Rev* 39: 428-441
Chen JS, Ma E, Harrington LB, Da Costa M, Tian X, Palefsky JM, Doudna JA (2018) CRISPR-
Cas12a target binding unleashes indiscriminate single-stranded DNase activity. *Science* 360:
436-439
Clement K, Rees H, Canver MC, Gehrke JM, Farouni R, Hsu JY, Cole MA, Liu DR, Joung JK,
Bauer DE *et al* (2019) CRISPResso2 provides accurate and rapid genome editing sequence
analysis. *Nat Biotechnol* 37: 224-226
Costain AH, MacDonald AS, Smits HH (2018) Schistosome Egg Migration: Mechanisms,
Pathogenesis and Host Immune Responses. *Front Immunol* 9: 3042
Dalton JP, Day SR, Drew AC, Brindley PJ (1997) A method for the isolation of schistosome
eggs and miracidia free of contaminating host tissues. *Parasitology* 115 (Pt 1): 29-32
Dumeau CE, Monfort A, Kissling L, Swarts DC, Jinek M, Wutz A (2019) Introducing gene
deletions by mouse zygote electroporation of Cas12a/Cpf1. *Transgenic Res* 28: 525-535
Everts B, Perona-Wright G, Smits HH, Hokke CH, van der Ham AJ, Fitzsimmons CM, Doenhoff
MJ, van der Bosch J, Mohrs K, Haas H *et al* (2009) Omega-1, a glycoprotein secreted by
Schistosoma mansoni eggs, drives Th2 responses. *J Exp Med* 206: 1673-1680
Fuchs RT, Curcuru J, Mabuchi M, Yourik P, Robb GB (2019) Cas12a trans-cleavage can be
modulated in vitro and is active on ssDNA, dsDNA, and RNA. *bioRxiv*: 600890
Germini D, Tsfasman T, Zakharova VV, Sjakste N, Lipinski M, Vassetzky Y (2018) A
Comparison of Techniques to Evaluate the Effectiveness of Genome Editing. *Trends Biotechnol*
36: 147-159

Gier RA, Budinich KA, Evitt NH, Cao Z, Freilich ES, Chen Q, Qi J, Lan Y, Kohli RM, Shi J
(2020) High-performance CRISPR-Cas12a genome editing for combinatorial genetic screening.
Nat Commun 11: 3455
Hendel A, Bak RO, Clark JT, Kennedy AB, Ryan DE, Roy S, Steinfeld I, Lunstad BD, Kaiser
RJ, Wilkens AB *et al* (2015) Chemically modified guide RNAs enhance CRISPR-Cas genome
editing in human primary cells. *Nat Biotechnol* 33: 985-989
Hoffmann KF, Brindley PJ, Berriman M (2014) Medicine. Halting harmful helminths. *Science*
346: 168-169
Hsu PD, Lander ES, Zhang F (2014) Development and applications of CRISPR-Cas9 for
genome engineering. *Cell* 157: 1262-1278
Huang CH, Lee KC, Doudna JA (2018) Applications of CRISPR-Cas Enzymes in Cancer
Therapeutics and Detection. *Trends Cancer* 4: 499-512
Hur JK, Kim K, Been KW, Baek G, Ye S, Hur JW, Ryu SM, Lee YS, Kim JS (2016) Targeted
mutagenesis in mice by electroporation of Cpf1 ribonucleoproteins. *Nat Biotechnol* 34: 807-808
International Helminth Genomes C (2019) Comparative genomics of the major parasitic worms.
Nat Genet 51: 163-174
Ittiprasert W, Mann VH, Karinshak SE, Coghlan A, Rinaldi G, Sankaranarayanan G, Chaidee A,
Tanno T, Kumkhaek C, Prangtaworn P *et al* (2019) Programmed genome editing of the omega-1
ribonuclease of the blood fluke, Schistosoma mansoni. *Elife* 8
Jacobsen T, Ttofali F, Liao C, Manchalu S, Gray BN, Beisel CL (2020) Characterization of
Cas12a nucleases reveals diverse PAM profiles between closely-related orthologs. *Nucleic Acids*
Res 48: 5624-5638
Jeon Y, Choi YH, Jang Y, Yu J, Goo J, Lee G, Jeong YK, Lee SH, Kim IS, Kim JS *et al* (2018)
Direct observation of DNA target searching and cleavage by CRISPR-Cas12a. *Nat Commun* 9:
2777
Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E (2012) A programmable
dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 337: 816-821
Jurberg AD, Goncalves T, Costa TA, de Mattos AC, Pascarelli BM, de Manso PP, Ribeiro-Alves
M, Pelajo-Machado M, Peralta JM, Coelho PM *et al* (2009) The embryonic development of
Schistosoma mansoni eggs: proposal for a new staging system. *Dev Genes Evol* 219: 219-234
Kellner MJ, Koob JG, Gootenberg JS, Abudayyeh OO, Zhang F (2019) SHERLOCK: nucleic
acid detection with CRISPR nucleases. *Nat Protoc* 14: 2986-3012
Kielkopf CL, Bauer W, Urbatsch IL (2020) Bradford Assay for Determining Protein
Concentration. *Cold Spring Harb Protoc* 2020: 102269
Kim D, Kim J, Hur JK, Been KW, Yoon SH, Kim JS (2016) Genome-wide analysis reveals
specificities of Cpf1 endonucleases in human cells. *Nat Biotechnol* 34: 863-868
Kissling L, Monfort A, Swarts DC, Wutz A, Jinek M (2019) Preparation and electroporation of
Cas12a/Cpf1-guide RNA complexes for introducing large gene deletions in mouse embryonic
stem cells. *Methods Enzymol* 616: 241-263
Koonin EV, Makarova KS, Wolf YI (2017) Evolutionary Genomics of Defense Systems in
Archaea and Bacteria. *Annu Rev Microbiol* 71: 233-261
Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E (2016) CHOPCHOP v2: a web tool
for the next generation of CRISPR genome engineering. *Nucleic Acids Res* 44: W272-276
Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E (2019)
CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. *Nucleic Acids*
Res 47: W171-W174

Lee C, Kim J, Shin SG, Hwang S (2006) Absolute and relative QPCR quantification of plasmid
copy number in Escherichia coli. *J Biotechnol* 123: 273-280
Li B, Yan J, Zhang Y, Li W, Zeng C, Zhao W, Hou X, Zhang C, Dong Y (2020) CRISPR-
Cas12a Possesses Unconventional DNase Activity that Can Be Inactivated by Synthetic
Oligonucleotides. *Mol Ther Nucleic Acids* 19: 1043-1052
Li S-Y, Cheng Q-X, Wang J-M, Li X-Y, Zhang Z-L, Gao S, Cao R-B, Zhao G-P, Wang J
(2018a) CRISPR-Cas12a-assisted nucleic acid detection. *Cell Discovery* 4: 20
Li SY, Cheng QX, Liu JK, Nie XQ, Zhao GP, Wang J (2018b) CRISPR-Cas12a has both cis-
and trans-cleavage activities on single-stranded DNA. *Cell Res* 28: 491-493
Liu P, Luk K, Shin M, Idrizi F, Kwok S, Roscoe B, Mintzer E, Suresh S, Morrison K, Frazão JB
et al (2019) Enhanced Cas12a editing in mammalian cells and zebrafish. *Nucleic Acids Res* 47:
4169-4180
Liu Y, Han J, Chen Z, Wu H, Dong H, Nie G (2017) Engineering cell signaling using tunable
CRISPR-Cpf1-based transcription factors. *Nat Commun* 8: 2095
Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25: 402-408
Lu Z, Sankaranarayanan G, Rawlinson KA, Offord V, Brindley PJ, Berriman M, Rinaldi G
(2021) The Transcriptome of Schistosoma mansoni Developing Eggs Reveals Key Mediators in
Pathogenesis and Life Cycle Propagation. *Frontiers in Tropical Diseases* 2
Luhtala N, Parker R (2010) T2 Family ribonucleases: ancient enzymes with diverse roles. *Trends*
Biochem Sci 35: 253-259
Ma P, Meng Q, Sun B, Zhao B, Dang L, Zhong M, Liu S, Xu H, Mei H, Liu J *et al* (2020)
MeCas12a, a Highly Sensitive and Specific System for COVID-19 Detection. *Advanced Science*
7: 2001300
Malzahn AA, Tang X, Lee K, Ren Q, Sretenovic S, Zhang Y, Chen H, Kang M, Bao Y, Zheng X
et al (2019) Application of CRISPR-Cas12a temperature sensitivity for improved genome
editing in rice, maize, and Arabidopsis. *BMC Biol* 17: 9
Mann VH, Morales ME, Rinaldi G, Brindley PJ (2010) Culture for genetic manipulation of
developmental stages of Schistosoma mansoni. *Parasitology* 137: 451-462
Mann VH, Suttiprapa S, Skinner DE, Brindley PJ, Rinaldi G (2014) Pseudotyped murine
leukemia virus for schistosome transgenesis: approaches, methods and perspectives. *Transgenic*
Res 23: 539-556
McVeigh P, Maule AG (2019) Can CRISPR help in the fight against parasitic worms? *Elife* 8
Meshalkina DA, Glushchenko AS, Kysil EV, Mizgirev IV, Frolov A (2020) SpCas9- and
LbCas12a-Mediated DNA Editing Produce Different Gene Knockout Outcomes in Zebrafish
Embryos. *Genes (Basel)* 11
Montague TG, Cruz JM, Gagnon JA, Church GM, Valen E (2014) CHOPCHOP: a
CRISPR/Cas9 and TALEN web tool for genome editing. *Nucleic Acids Res* 42: W401-407
Mustafa MI, Makhawi AM (2020) SHERLOCK and DETECTR: CRISPR-Cas Systems as
Potential Rapid Diagnostic Tools for Emerging Infectious Diseases. *J Clin Microbiol*
Nguyen LT, Smith BM, Jain PK (2020) Enhancement of trans-cleavage activity of Cas12a with
engineered crRNA enables amplified nucleic acid detection. *Nature Communications* 11: 4906
Nguyen PQ, Soenksen LR, Donghia NM, Angenent-Mari NM, de Puig H, Huang A, Lee R,
Slomovic S, Galbersanini T, Lansberry G *et al* (2021) Wearable materials with embedded
synthetic biology sensors for biomolecule detection. *Nat Biotechnol*

Nishimasu H, Ran FA, Hsu PD, Konermann S, Shehata SI, Dohmae N, Ishitani R, Zhang F,
Nureki O (2014) Crystal structure of Cas9 in complex with guide RNA and target DNA. *Cell*
156: 935-949
Paix A, Folkmann A, Goldman DH, Kulaga H, Grzelak MJ, Rasoloson D, Paidemarry S, Green
R, Reed RR, Seydoux G (2017) Precision genome editing using synthesis-dependent repair of
Cas9-induced DNA breaks. *Proc Natl Acad Sci U S A* 114: E10745-e10754
Pinello L, Canver MC, Hoban MD, Orkin SH, Kohn DB, Bauer DE, Yuan GC (2016) Analyzing
CRISPR genome-editing experiments with CRISPResso. *Nat Biotechnol* 34: 695-697
Protasio AV, Tsai IJ, Babbage A, Nichol S, Hunt M, Aslett MA, De Silva N, Velarde GS,
Anderson TJ, Clark RC *et al* (2012) A systematically improved high quality genome and
transcriptome of the human blood fluke Schistosoma mansoni. *PLoS Negl Trop Dis* 6: e1455
Ranawakage DC, Okada K, Sugio K, Kawaguchi Y, Kuninobu-Bonkohara Y, Takada T,
Kamachi Y (2020) Efficient CRISPR-Cas9-Mediated Knock-In of Composite Tags in Zebrafish
Using Long ssDNA as a Donor. *Front Cell Dev Biol* 8: 598634
Sankaranarayanan G, Berriman M, Rinaldi G (2021) An uneven race: genome editing for
parasitic worms. *Nat Rev Microbiol* 19: 621
Santos L, Mention K, Cavusoglu-Doran K, Sanz DJ, Bacalhau M, Lopes-Pacheco M, Harrison
PT, Farinha CM (2021) Comparison of Cas9 and Cas12a CRISPR editing methods to correct the
W1282X-CFTR mutation. *J Cyst Fibros*
Schwartz C, Fallon PG (2018) Schistosoma "Eggs-Iting" the Host: Granuloma Formation and
Egg Excretion. *Front Immunol* 9: 2492
Sentmanat MF, Peters ST, Florian CP, Connelly JP, Pruett-Miller SM (2018) A Survey of
Validation Strategies for CRISPR-Cas9 Editing. *Sci Rep* 8: 888
Shirvani R, Barshan-Tashnizi M, Shahali M (2020) An investigation into gene copy number
determination in transgenic yeast; The importance of selecting a reliable real-time PCR standard.
Biologicals 65: 10-17
Smith CW, Nandu N, Kachwala MJ, Chen YS, Uyar TB, Yigit MV (2020) Probing CRISPR-
Cas12a Nuclease Activity Using Double-Stranded DNA-Templated Fluorescent Substrates.
Biochemistry 59: 1474-1481
Steinfelder S, Andersen JF, Cannons JL, Feng CG, Joshi M, Dwyer D, Caspar P, Schwartzberg
PL, Sher A, Jankovic D (2009) The major component in schistosome eggs responsible for
conditioning dendritic cells for Th2 polarization is a T2 ribonuclease (omega-1). *J Exp Med* 206:
1681-1690
Straume AH, Kjaerner-Semb E, Ove Skaftnesmo K, Guralp H, Kleppe L, Wargelius A,
Edvardsen RB (2020) Indel locations are determined by template polarity in highly efficient in
vivo CRISPR/Cas9-mediated HDR in Atlantic salmon. *Sci Rep* 10: 409
Swarts DC (2019) Making the cut(s): how Cas12a cleaves target and non-target DNA. *Biochem*
Soc Trans 47: 1499-1510
Swarts DC, Jinek M (2018) Cas9 versus Cas12a/Cpf1: Structure-function comparisons and
implications for genome editing. *Wiley Interdiscip Rev RNA*: e1481
Swarts DC, Jinek M (2019) Mechanistic Insights into the cis- and trans-Acting DNase Activities
of Cas12a. *Mol Cell* 73: 589-600 e584
Takaki KK, Rinaldi G, Berriman M, Pagan AJ, Ramakrishnan L (2021a) Schistosoma mansoni
Eggs Modulate the Timing of Granuloma Formation to Promote Transmission. *Cell Host*
Microbe 29: 58-67 e55

Takaki KK, Roca FJ, Schramm G, Wilbers RHP, Ittiprasert W, Brindley PJ, Rinaldi G, Berriman
M, Ramakrishnan L, Pagan AJ (2021b) Tumor Necrosis Factor and Schistosoma mansoni egg
antigen omega-1 shape distinct aspects of the early egg-induced granulomatous response. *PLoS*
Negl Trop Dis 15: e0008814
Tóth E, Czene BC, Kulcsár PI, Krausz SL, Tálas A, Nyeste A, Varga É, Huszár K, Weinhardt N,
Ligeti Z *et al* (2018) Mb- and FnCpf1 nucleases are active in mammalian cells: activities and
PAM preferences of four wild-type Cpf1 nucleases and of their altered PAM specificity variants.
Nucleic Acids Res 46: 10272-10285
Wilbers RH, Westerhof LB, van Noort K, Obieglo K, Driessen NN, Everts B, Gringhuis SI,
Schramm G, Goverse A, Smant G *et al* (2017) Production and glyco-engineering of
immunomodulatory helminth glycoproteins in plants. *Sci Rep* 7: 45910
Yuan JS, Burris J, Stewart NR, Mentewab A, Stewart CN, Jr. (2007) Statistical tools for
transgene copy number estimation based on real-time PCR. *BMC Bioinformatics* 8 Suppl 7: S6
Zamanian M, Andersen EC (2016) Prospects and challenges of CRISPR/Cas genome editing for
the study and control of neglected vector-borne nematode diseases. *Febs j* 283: 3204-3221
Zetsche B, Gootenberg JS, Abudayyeh OO, Slaymaker IM, Makarova KS, Essletzbichler P, Volz
SE, Joung J, van der Oost J, Regev A *et al* (2015) Cpf1 is a single RNA-guided endonuclease of
a class 2 CRISPR-Cas system. *Cell* 163: 759-771
Zhang Y, Ren Q, Tang X, Liu S, Malzahn AA, Zhou J, Wang J, Yin D, Pan C, Yuan M *et al*
(2021) Expanding the scope of plant genome engineering with Cas12a orthologs and highly
multiplexable editing systems. *Nat Commun* 12: 1944
942

C

Cas9-ssODN:
cttgtccacttttaggaaacatgaatttgaaaagcatggtttgtgtgcag<u>TAAGTGACTAGGTAACTGAGTAGC</u>ttgaggatcctcaagtgtttaatcaatacggttactttaaatttggtatt

Cas12a- NT-ssODN:
tgtccacttttaggaaacatgaatttgaaaagcatggtttgtgtgcagtt<u>TAAGTGACTAGGTAACTGAGTAGC</u>gaggatcctcaagtgtttaatcaatacggttactttaaatttggtattaa

Cas12a- T-ssODN:
attggaacaggtgaaaatcctttgtacttaaacttttcgtaccaaacacac<u>ATTCACTGATCCATTGACTCATCG</u>gtcaactcctaggagttcacaaattagttatgccaatgaaatttaaacca

figure 2

A

B

C

D