RNA-guided *As*Cas12a- and *Sp*Cas9-catalyzed knockout and homology
directed repair of the *omega-1* locus of the human blood fluke, *Schistosoma*
mansoni

Wannaporn Ittiprasert[1,*,#], Chawalit Chatupheeraphat[1, 2,*], Victoria H. Mann[1], Wenhui
Li[1,3], André Miller[4], Taiwo Ogunbayo[4], Kenny Tran[4], Yousef Alrefaei[1,5], Margaret
Mentink-Kane[4], Paul J. Brindley[1,#]

[1] Department of Microbiology, Immunology & Tropical Medicine, & Research Center for
Neglected Diseases of Poverty, School of Medicine & Health Sciences, George Washington
University, Washington, DC 20037, USA.

[2] Center for Research and Innovation, Faculty of Medical Technology, Mahidol University,
Salaya, Nakhon Pathom, 73170 Thailand

[3] Lanzhou Veterinary Research Institute, Chinese Academy of Agricultural Sciences, Lanzhou,
730050, Gansu Province, China

[4] Schistosomiasis Resource Center, Biomedical Research Institute, Rockville, MD 20850

[5] College of Health Sciences, PAEET, Adailiya, Kuwait City, 73101, Kuwait

*, equal contribution

#, correspondence: Wannaporn Ittiprasert, wannaporni@gwu.edu, Paul J. Brindley,
pbrindley@gwu.edu

Keywords: *As*Cas12a; *Sp*Cas9; ribonucleoprotein complex, CRISPR; omega-1; genome editing;
homology directed repair; non-homologous end-joining; schistosome egg

Abstract

We compared the efficiency and precision of the RNA-guided *As*Cas12a nuclease of *Acidaminococcus* sp. with *Sp*Cas9 of *Streptococcus pyogenes* aiming to advance functional genomics tools for *Schistosoma mansoni*. Programmed double stranded cleavage catalyzed by *As*Cas12a results in a staggered strand break whereas *Sp*Cas9 produces a blunt ended chromosomal break. The TTTV, the optimal protospacer adjacent motif for *As*Cas12a is expected frequently within the AT-rich genome of this platyhelminth. We deployed optimized conditions (gRNA:*Sp*Cas9:DNA donor ratio and electroporation condition) to edit the *ω1* gene. *Sp*Cas9 delivered higher efficiency to mutate *ω1* target compared to *As*Cas12a for non-homology end joining (NHEJ)-catalyzed repair (14.04% *vs.* 10.88%, n = 7 replicates). Most mutations were deletions; *Sp*Cas9 induced -3 nt size deletions whereas *As*Cas12a induced deletions ranging in size from -25 to -2 nt. Although these were less absolute percentage *As*Cas12a than *Sp*Cas9 programmed mutations, the phenotypic outcomes on levels of *ω1* transcripts and expressed omega-1 protein were similar. Gene editing efficiency of *Sp*Cas9 and *As*Cas12a markedly increased in the presence of short single stranded donor templates bearing symmetrical homolog arms of 50 nt in length. With *As*Cas12a, both non-CRISPR target (NT) and target (T) strands of the *ω1* gene were tested as homology direct repair (HDR) donor templates. There were 15.67%, 28.71% and 21.43% of NHEJ from 7 pooled genomic DNA from KI_*Sp*Cas9, KI_*As*Cas12a-NT-ssODN and KI_*As*Cas12a-T-ssODN experiments, respectively. Programmed *Sp*Cas9 cleavage led to higher levels than *As*Cas12a of precise HDR mediated; 17.07%, KI_*Sp*Cas9 *vs.* 14.58%, KI_*As*Cas12a-NT-ssODN and 12.35%, KI_*As*Cas12a-T-ssODN ($P < 0.0.5$), although no significant differences in reduction in ω1 transcripts or of protein levels were apparent. These findings revealed that both *As*Cas12a and *Sp*Cas9 can provide programmed knockout and transgene insertion into genes expressed in the schistosome egg.

Introduction

Interest has increased in the use of functional genomics to investigate helminth parasites, specifically genome editing to investigate pathophysiology, carcinogenesis and biology. Recent progress has ushered in and showcased the deployment of CRISPR/Cas programmed genome editing in parasitic flatworms, in particular with the human blood fluke *Schistosoma mansoni* (Ittiprasert *et al*, 2019) and the carcinogenic liver fluke, *Opisthorchis viverrini* (Arunsan *et al*, 2019) as exemplars. However, given that these helminth parasites are non-model organisms, that the genome sequences are often not well established or curated, and that they are frequently refractory or even intractable to maintain in the laboratory or laboratory rodents, approaches to advance functional genomics investigation using the CRISPR/Cas technologies are not yet established or available for use in the field of tropical medical and neglected tropical diseases research. These fields, nonetheless, and our capacity to manage these diseases will surely be aided by advanced in gene editing technologies (Hoffmann *et al*, 2014).

RNA-guided nucleases evolved in archaea and bacteria as an for adaptive immunity mechanism to deal with and destroy phage viruses and other nucleic acid invaders (Jinek *et al*, 2012; Koonin *et al*, 2017). The catalogue of these CRISPR-Cas nucleases is expanding. The *Sp*Cas9 of *Streptococcus pyogenes,* a Class 2, Type II Cas enzyme in the phylogeny of Koonin and coworkers (Koonin *et al.*, 2017), dominates diverse gene-editing applications in biomedicine and beyond (Charpentier, 2015; Huang *et al*, 2018; Mustafa & Makhawi, 2020), given its precise double-stranded DNA (dsDNA) cleavage induced by two catalytic domains, RuvC and HNH, at sequences complementary to the guide RNA (Nishimasu *et al*, 2014). A second family of Class 2 enzymes, type V nucleases as represented by *As*Cas12a (formerly known as Cpf1), uses a single RuvC catalytic domain for guide RNA-directed dsDNA cleavage (Jeon *et al*, 2018; Zetsche *et al*, 2015). Distinct from *Sp*Cas9, *As*Cas12a orthologues recognize a T nucleotide–rich protospacer-adjacent motif (PAM), catalyze the maturation of their own guide CRISPR RNA (crRNA) and a dsDNA break distal to the PAM bearing staggered 5′- and 3′-ends (Jeon *et al.*, 2018), features that are attractive for genome editing (Malzahn *et al*, 2019). For their maturation, the crRNAs of type II CRISPR-Cas systems require processing by a cognate trans-activating CRISPR RNA (Charpentier *et al*, 2015). Moreover, the *As*Cas12a endonuclease is smaller than *Sp*Cas9 in size and uses a shorter CRISPR RNA (crRNA) for activity (Liu *et al*, 2017). Notably, target site activated *As*Cas12a exhibits non-specific *trans* activity against bystander nucleic acids (Chen *et al*, 2018; Smith *et al*, 2020), an attribute that has enabled its application in a burgeoning range of biosensors (Kellner *et al*, 2019; Li *et al*, 2018a).

Delivery of CRISPR reagents into the cells of the multicellular parasite *S. mansoni* in the form of ribonucleoprotein (RNP) complexes can enable immediate and permanent programmed gene editing and obviate concerns with CRISPR reagent longevity or integration in the host cell genome. Programmed CRISPR-editing using RNP-based delivery can precede analysis of mutated target regions that rely on cloning and vector construction steps. Previously, we deployed both RNPs and lentiviral-delivered CRISPR/*Sp*Cas9 approaches to knock-out (KO) and knock-in (KI) into a blunt end at *Sp*Cas9 cleavage site at the multiple copies of the omega-1 (*ω1*) gene in parasite eggs. To advance functional genomics for schistosomes and aiming to improve the mutagenesis and transgene knock-in of CRISPR efficiency in schistosomes, which exhibit AT-rich genomes, we investigated the nuclease activity of Cas12a of *Acidaminococcus* sp. (*As*Cas12a) in contrast with Cas9 of *S. pyogenes* (*Sp*Cas9). These CRISPR systems differ in two

main aspects: 1) *Sp*Cas9 recognizes NGG as the protospacer adjacent motif (PAM) whereas
*As*Cas12a recognizes the T-rich, TTTV; 2) Catalysis by *Sp*Cas9 results in blunt double stranded
DNA at three nucleotides 5' to the PAM whereas *As*Cas12a results in a staggered cleavage at 23
nt (target strand; T) and 18 nt (non-target strand; NT) 3' to the PAM (Banakar *et al*, 2020).
Comparisons of these two RNA-programmed nucleases in the context of the *ω1* gene of *S.*
mansoni may inform decisions for functional genomics manipulations aiming to enhance
efficiency of homology directed repair (HDR) targeting genes expressed by the developing
schistosome egg and other developmental stages (Ittiprasert *et al.*, 2019; Sankaranarayanan *et al*,
2021; Takaki *et al*, 2021a; Takaki *et al*, 2021b).

Results

The *omega-1* multicopy gene, guide RNAs, and single stranded DNA donor templates

The gene editing efficiency of *Sp*Cas9 and *As*Cas12a was compared in assays targeting the
omega-1 gene which encoded a T2 ribonuclease that is secreted by the mature egg of *S. mansoni*
into the surrounding host tissues and circumoval granuloma (Schwartz & Fallon, 2018).
Specifically, the four copies of *omega-1*, well annotated in WormBase Parasite (Bolt *et al*, 2018)
(update of 16 September 2021) as hepatotoxic ribonuclease – Smp_334170.1, Smp_334170.2,
Smp_334240, and Smp_333930, were targeted for design of gRNAs and donor repair templates.
These four copies reside on chromosome 1 of *S. mansoni*, with reverse strand positions at
3,980,364 to 3,984675, 3,992,964 to 3,995,246 and 3,908,953 to 3,911,250 (PRJEA36577),
respectively. These copies share >99% nucleotide sequence identity, are ~2.3 kb in length, are
highly AT rich (65%), and encode an enzyme of 127 (Smp_334170.2, Smp_334240 and
Smp_333930) or 115 (Smp_334170.1) amino acid residues in length. Smp_333220, also located
on chromosome 1 at 3,925,420-3,927,716 but lacking annotation on the *S. mansoni* annotated
genome version 7_2021, was also included. Smp_333220 shares >99% chromosomal and
mRNA sequence identity with the three *ω1* copies (Fig S1B). Guide RNAs for *Sp*Cas9 and
*As*Cas12a designed with the assistance of CHOPCHOP both target the same region of exon 1 of
omega-1, disrupting its catalytic active site histidine (Luhtala & Parker, 2010), with the
programmed cleavage sites for Cas in close proximity, separated by three nucleotides (Fig 1A
and B).

We designed three single stand oligodeoxynucleotide (ssODN) donors for homology directed
repair (HDR) of *ω1* following programmed mutation. Given the differing PAM requirements of
*Sp*Cas9 and *As*Cas12a as well as the divergent formats of the DSB, blunted ended by *Sp*Cas9 and
sticky ended by *As*Cas12a, the designs differed for the ssODNs for use with each nuclease.
Based on earlier findings (Ittiprasert *et al.*, 2019), the design of the ssODN donors included 3'
and 5' homology arms (HA) of each of 50 nt in length, complementary to the target strand for
*Sp*Cas9 and for *As*Cas12a, for either the target strand, termed *Sp*Cas9-T-ssODN, or for the non-
target strand donor, termed *As*Cas12a-NT-ssODN. Note that sequences identical to the PAM for
*As*Cas12a and ~18 nt of the protospacer region were retained within the 5'-HA of *As*Cas12a-T-
ssODN (Fig 1B and C).

Absence of *trans* activity of activated *As*Cas12a against the ssODN donors

Previous reports dealing with *Lachnospiraceae bacterium* ND2006 and *Francisella novicida* *As*Cas12a describe indiscriminate *trans*-activity unleashed on ssDNA after gRNA activation and target binding of *As*Cas12a (Chen *et al.*, 2018; Gier *et al*, 2020). Given that our goal here was to compare *Sp*Cas9 and *As*Cas12a in schistosomes, and given that we used an orthologous *As*Cas12a from *Acidaminococcus* sp*.* BV3L6 (IDT), in both KO and KI assays, we sought to confirm whether the *omega-1* target gene activated *As*Cas12a might exhibit also *trans*-activity against the donor single stranded templates that we planned to use in KI assays, *As*Cas12a-T-ssODN and *As*Cas12a-NT-ssODN, in addition to RNA-guided double stranded DNA cleavage activity.

Accordingly, we also carried out *in vitro As*Cas12a nuclease activity assays using RNPs targeting a 426 bp amplicon of the exon 1 of *omega-1* (

A, 2B and 2C). These assays confirmed that the crRNA was highly active and following incubation at 37°C for 30 min had cleaved the double stranded amplicon at the programmed cleavage site into two fragments of 265 and 161 bp, as expected (Fig 2B; Agilent Bioanalyzer electropherogram). In contrast, the activated *As*Cas12a-RNP complex failed to digest either of the single stranded templates, ssODNs *As*Cas12a-T-ssODN or *As*Cas12a-NT-ssODN after 120 min at 37°C (Fig 2C). As seen in the electropherogram, the ssODNs remained intact at 124 nt. Single stranded DNA standards ranging in size from 200 to < 40 nt were included in the assay (Fig 2C). In summary, the findings suggested that *omega-1* target gene activated As*As*Cas12a would not exhibit indiscriminate *trans*-nuclease activity against the ssODNs within schistosome cells, although an *in vitro* surrogate assay cannot faithfully mimic conditions within a viable schistosome egg.

Estimation of CRISPR efficiency of *Sp*Cas9 and *As*Cas12a

RNP complexes of *Sp*Cas9-sgRNA and *As*Cas12a-CRISPR RNA (crRNA) targeting *ω1* were delivered to the LE by electroporation (Ittiprasert *et al.*, 2019). The efficiency of programmed mutation was assessed by TIDE analysis of Sanger sequence chromatograms (Gier *et al.*, 2020; Jacobsen *et al*, 2020), followed by targeted amplicon NGS and analysis of the reads using CRISPResso2 (Pinello *et al*, 2016). Twelve independent biological replicates were carried out. *As*Cas12a programmed KO led to significantly more indels than were induced with *Sp*Cas9 *(P <* 0.001, paired, two-tailed *t* = 7.450, df = 11) (Fig 2G and H). The mean CRISPR efficiency for KO with *Sp*Cas9 was 0.89%±0.43 (range, 0.30-1.31%) and 1.83%±0.29 (range, 1.39-2.54%) with *As*Cas12a, when compared with the control (mock) groups. KI efficiency also was investigated for individual replicates by TIDE and for pooled samples by analysis amplicon NGS (Illumina) reads. Programmed mutation by *Sp*Cas9 was more efficient in the presence or absence of donor ssODN, i.e., KI *versus* KO groups, 1.43±0.4, range 0.51-2.31% (Fig 2G and H). By contrast, differences in mutation efficiency were not apparent between the KO and KI groups using *As*Cas12a; 1.75%±0.63 (range 1.11-2.69%) and 1.54%±0.49 (range 1.82-2.20%) for *As*Cas12a-NT ssODN and *As*Cas12a-T ssODN, respectively.

Quantification of transgene copy number by external standard curve

The external standard curve design that we employed to estimate transgene copy number in
genomics DNA of the schistosome eggs estimates transgene copy number by comparison with an
external standard curve (Fig 2E and F). The standard curve on simple linear regression (y = -
$4.0964\ln[10] + 44.821$, $R^2 = 0.9923$) was established with logarithm-10 transformed initial DNA
input, plasmid pCR4, which encodes the six stop codon cassette (Ittiprasert *et al.*, 2019), as the
dependent and the Ct value from the qPCRs as the independent variable (not shown). We
estimated the transgene copy number on the 5' and 3' of KI site in *ω1* by converting the
measured Ct values to log copy numbers using the equations obtained from the standard curves.
Subsequently, dividing the means by the antilog of each result, we arrived at an estimate of the
transgene copy number. A 790 bp product amplified for the control wildtype sequence and
amplicons of 235 bp and 182 bp for the 5' and 3' flanking regions, respectively, of the KI
genomic DNA containing the six stop codon transgene were analyzed, which revealed 3'-side
transgene integration values of 3.61±0.25, 4.44±0.84 and 4.19±1.22 from the KI_*Sp*Cas9,
KI_*As*Cas12a-NT-ssODN and KI_*As*Cas12a-T-ssODN treatment groups, respectively (*P* =
0.4057, = 0.6264 and = 0.9155; NS; one-way ANOVA, multiple comparisons, 95% CI for
differences of the means) (Fig 2E). However, the presence of asymmetrical HDR was revealed
by significant differences among the copy number values for the 5' side transgene integration
estimates: 2.72±0.42, 5.06±0.54, and 1.64±0.39 for the KI_*Sp*Cas9, KI_*As*Cas12a-NT-ssODN
and KI_*As*Cas12a-T-ssODN groups, respectively (Fig 2F; *P* = 0.0214, *P*=0.0001 and *P* < 0.0001,
*,***, ****, respectively; one-way ANOVA, multiple comparisons).

Estimates of indels from NHEJ and/or HDR using NGS reads from targeted amplicons

Pooled gDNAs (above) also were used as templates for targeted amplicons for NGS and
CRISPResso2 employed to assess the efficiency of transgene KI from HDR in the NGS reads.
Comparison of the mutations among all four copies of the *ω1* gene also were undertaken. From
Whereas TIDE analysis using Sanger direct sequencing chromatograms was used above to
estimate % efficiency of CRISPR manipulations in individual biological replicates, here targeted
amplicons from pools of gDNAs from the biological replicates was used to acquire libraries of
reads (Amplicon EZ, Genewiz). More than 150,000 reads from each NGS library were obtained.
Analysis of the reads using CRISPResso2 provided increased sensitivity of detection of
programmed gene editing than TIDE, as expected (Sentmanat *et al*, 2018). Specifically, there
were 12.26%, 1.66%, 0.12% modified sequence (NHEJ) for KO-*Sp*Cas9 and 9.27%, 1.33% and
0.28% of NHEJ for KO-*As*Cas12a, when compared with the WormBase Parasite reference
sequence (identical amplicon from Smp_334170.1, Smp_334170.2, Smp_334240), Smp_333930
and Smp_333870, respectively (Fig 3A and B). A small number of reads were assessed to be
ambiguous by CRISPResso2, 0.49% and 0.93% for KO-*Sp*Cas9 and KO-*As*Cas12a, respectively
(Fig 3A and B). Deletions were the only type of indel detected, for both *Sp*Cas9 and *As*Cas12a.
In like fashion to other studies (An *et al*, 2020; Dumeau *et al*, 2019; Kissling *et al*, 2019),
deletion of large sizes resulted from programmed *As*Cas12a editing (from -2 bp to -26 bp
deletion size) a the programmed DSB, compared to -3 bp at the DSB programmed from *Sp*Cas9
(Fig 3C and D provide representative allele sequences). Both enzymes showed a preference to
mutate the reference Smp_334170 copy, and also Smp_334240, rather than the Smp_333930
and/or Smp_333870 copies of *omega-1*. The numbers of indels arising from NHEJ with *Sp*Cas9
and with *As*Cas12a both increased in the presence of a ssODN donor repair template. The total
NHEJ were 15.67%, 28.71% and 21.43% from KI-*Sp*Cas9-NT-ssODN, KI-*As*Cas12a-NT-
ssODN and KI-*As*Cas12a-T-ssODN, respectively (Fig 4A, B and C). Deletion size as

determined by CRISPResso2 for the KI-*Sp*Cas9 group was larger than for KO-*Sp*Cas9,
increasing up to -13 bp length deletions. Similarly, base deletion size from KI-*As*Cas12a was
larger than KO-*As*Cas12a with deletion of -42 bp detected (Fig 4D, E and F). Levels of
ambiguous results were estimated at 1.66%, 1.03% and 0.73% from KI-*Sp*Cas9, KI-*As*Cas12a-
NT-ssODN and KI-*As*Cas12a-T-ssODN, respectively.

To estimate the HDR events, we included the donor sequence as a parameter in the
CRISPResso2 analysis. More HDR events mediated by CRISPR/*Sp*Cas9 were detected than with
CRISPR/*As*Cas12a; 17.07% *vs* 14.56% and 12.37%, in like fashion to our earlier findings
(Santos *et al*, 2021). However, in *Sp*Cas9 sample, more imperfect HDR was seen (0.94%) than
in the *As*Cas12a groups (0.41% and 0.68%) (Fig 4A, B, C, G, H and F). Frequently, imperfect
HDR event alleles from both Cas nucleases were missing several residues (2-5 nt) on the 5' side
of the programmed DSB. We note that this mutation repair outcome may have impacted
quantification of the 5' KI transgene copy number, especially in KI-*As*Cas12a-T-ssODN samples
which contained 5 nt imperfect HDR repair on the six stop codon cassette (primer probing
sequence). Imperfect HDR at the 5' side might result from synthesis-dependent strand annealing
(Straume *et al*, 2020).

Transcription of *ω1* interrupted by programmed mutation by *Sp*Cas9 and *As*Cas12a

Primers specific for the four annotated copies of *ω1* were employed to investigate transcription
by real time quantitative PCR (Fig S1C). Relative expression levels of *ω1* mRNA in negative
controls and experimental KO or KI eggs were compared with wild type. Significantly reduced
levels of *ω1* were seen in eggs following the experimental the CRISPR manipulations for gene
KO and KI. Transcription of *ω1* was reduced, as follows: 63.76%±5.57, 76.88%±11.33,
71.81%±10.91, 98.69%±0.44 and 80.04%±0.09 in the KO_*Sp*Cas9, KI_*Sp*Cas9, KO_*As*Cas12a,
KI_*As*Cas12a-NT-ssODN and *As*Cas12a-T-ssODN groups, respectively. Notably, LE subjected
to KI_*As*Cas12a_NT-ssODN retained <2% ω1 transcript levels of the mock control (Fig 5A),
and the level in the KI_*As*Cas12a-NT-ssODN group was significantly lower than the
KI_*As*Cas12a-T-ssODN group, KI_*Sp*Cas9-ssODN. Significant differences were not apparent
between the KO_*Sp*Cas9 and KO_*As*Cas12a groups, generally. There were minor differences in
levels of transcription among the negative control groups (Fig 5A). Differences among treatment
groups were assessed using ANOVA with multiple comparisons (*P* values < 0.01 to < 0.0001; n=
4; F = 11.77).

Protein levels ascertained by western blotting

An anti-rω1 rabbit IgG was used to investigate levels of omega-1 in lysates (SEA) of LE. SEA
(200 ng) from the three controls, i.e., the mock, enzyme only, and mixed sgRNAs only groups,
and from the KO and KI groups with *Sp*Cas9 or *As*Cas12a was size fractionated by SDS-PAGE
and transferred to PVDF and probed with the rabbit antibody (Fig 5B). Following densitometric
analysis of the resulting signals, the levels of omega-1 remaining in the SEA were estimated at
64%±11.65, 43.88%±6.02, 49.27%±4.19, 37.9%±2.28 and 52.16%±10 in the KO_*Sp*Cas9,
KI_*Sp*Cas9, KO_*As*Cas12a, KI_*As*Cas12a-NT-ssODN and KI_*As*Cas12a-T-ssODN groups,
respectively. The differential intensity of ω1expression was not diminished among the sgRNA
only groups (Fig 5B and C). By contrast, ω1 protein levels were markedly diminished among all
the experimental treatment groups when compared with the controls (*P* <0.0001) (ANOVA, F =

27.05 , n = 4). Significant differences among for levels of ω1 among the KO and KI SEA
samples were not apparent (Fig 5C).

Discussion

Given the central role of the egg and its secretions in disease and in the transmission of
schistosomiasis (Costain *et al*, 2018), and given the potential of CRISPR/Cas gene editing to
advance understanding of this and other neglected tropical disease pathogens, we posit that
access to omega-1-negative eggs will provide an informative tool to confirm the role of this
ribonuclease in disease transmission. Functional knockout (KO) of the *omega-1* gene and the
resulting immunologically impaired phenotype have showcased the novel application of
CRISPR/*Sp*Cas9 and its utility for functional genomics in schistosomes (Hsu *et al*, 2014; Jinek *et*
al., 2012). In view of this progress, and the demonstrated tractability of *Sp*Cas9-catalyzed KO of
the *omega-1* gene, here compared the performance of a second RNA-guided Cas nuclease, in
terms of efficiency and precision, with *Sp*Cas9 and the *As*Cas12a nuclease of *Streptococcus*
pyogenes and *Acidaminococcus* sp. We now report that the efficiency of RNA guided
programmed knock-out of ω1 by *As*Cas12a was superior to *Sp*Cas9.

In this investigation, the RNA-guided target region for programmed mutation included PAMs for
both *Sp*Cas9 and *As*Cas12a within three nucleotides of each other within exon 1 of the coding
region of the omega-1 gene. The optimal protospacer adjacent motif for *As*Cas12a, TTTV
(Swarts & Jinek, 2019; Zetsche *et al.*, 2015), is expected to occur abundantly in the AT rich
genome of *S. mansoni* (Protasio *et al*, 2012; Zetsche *et al.*, 2015). Programmed double stranded
cleavage catalyzed by *As*Cas12a results in a staggered strand break of the target gene sequence
that may enhance efficiency of homology directed repair (HDR) (Swarts & Jinek, 2019). When
RNPs assembled from the nuclease and cognate crRNA, *As*Cas12a was significantly more
efficient than *Sp*Cas9 for KO, presumably reflected non-homology end joining (NHEJ)-catalyzed
repair and gene silencing. Specifically, the KO efficacy of *As*Cas12a was significantly higher
than that to *Sp*Cas9; 1.8% versus 0.9%, respectively. Greater mutation efficacy for versions of
*As*Cas12a over *Sp*Cas9 have been reported but, on the other hand, the reverse outcome has been
described (Hur *et al*, 2016; Kim *et al*, 2016). Comparisons among studies are not
straightforward because of differences in species, target genes, delivery method, and
temperature, among other confounders. In addition, there are three widely used orthologues of
*As*Cas12a, *Fn*Cas12a from *Francisella novicida U112*, *Lb*Cas12a from *Lachnospiraceae*
bacterium ND2006, and *As*Cas12a from *Acidaminococcus* sp. *BV3L6* and these have been tested
in diverse target eukaryotic species (Mustafa & Makhawi, 2020). (Other orthologues have also
been studied (Tóth *et al*, 2018; Zhang *et al*, 2021).) *As*Cas12a in active at 37°C whereas other
orthologues are catalytically active at lower temperatures, which has enabled their deployment in
research targeting zebrafish (Liu *et al*, 2019), insects, and maize (Zhang *et al.*, 2021), and
environmental and diagnostic biosensing (Chen *et al.*, 2018; Ma *et al*, 2020; Nguyen *et al*, 2021).

We compared these two CRISPR methods for programmed gene editing of the well-known
omega-1 extracellular T2 ribonuclease secreted by the mature egg of *S. mansoni*, which is a key
modulator of pathogenesis and transmission (Costain *et al.*, 2018; Everts *et al*, 2009; Schwartz &
Fallon, 2018; Steinfelder *et al*, 2009).   We investigated HDR-mediated gene editing with HiFi-
*Sp*Cas9 or *As*Cas12a delivered by electroporation in the presence of the RNPs and donor repair
templates donors. We designed the guide RNAs to anneal to protospacer regions that we

essentially at the same site of the gene. The programmed DSB were separated by only three
nucleotide residues, on the non-target strand of the target gene. Each donor ssODN template for
*Sp*Cas9- and *As*Cas12a-programmed KI included a transgene cassette of 24 nt in length with six
stop codons (Ittiprasert *et al.*, 2019) but in which the homology arms differed slightly on the non-
target strand due to the position of the predicted DSB. For programmed KI with
CRISPR/*As*Cas12a, we also included as the transgene, the reverse complement sequence of the
six stop codon cassette flanked by the target stand sequence as the homology arms, given
concern with respect to indiscriminate trans-activity by activated *As*Cas12a against random
single stranded DNA, and especially for the ssODN with NT homology arms which retains the
PAM and partial (~18 nt) protospacer sequence (Swarts, 2019; Swarts & Jinek, 2018, 2019).
RNA-guided *Sp*Cas9 cleavage led to higher level (~4% higher) of precise KO compared to that
obtained with *As*Cas12a but markedly less, ~6-13%, precise KI. Deletions were the frequent
mutation type seen with both *Sp*Cas9 (deletions of -3 nt in length) and *As*Cas12a (deletions of -
26 nt to -2 nt in length) due to NHEJ repair. Using conditions optimizing previously for *Sp*Cas9
KI (Ittiprasert *et al.*, 2019), here CRISPR/*Sp*Cas9 induced higher (~3%) HDR levels than
CRISPR/*As*Cas12a to insert the transgene into the *ω1* target for both *As*Cas12a-NT-ssODN and
*As*Cas12a-T-ssODN donors. Characterization of both *Sp*Cas9-and *As*Cas12a-edited eggs
revealed reductions in levels of both of *ω1* mRNAs and of expressed ω1 protein. Together, these
outcomes indicated that both of the CRISPR/Cas systems mutated this key schistosome egg
gene. Further optimization of the *As*Cas12a-based approach with regard to gRNA:*As*Cas12a
and/or gRNA:*As*Cas12a:ssODN ratios, format of donor(s), and/or temperature manipulation
likely will be informative.

A spectrum of chromosomal alleles was apparent following programmed KI of single stranded
DNA (ssDNA) repair templates of 124 nt in length with 5'- and 3'-homology arms specific for
the predicted cleavage sites of each nuclease. Both target strand (T) and non-target strand (NT)
ssDNA donors were tested. Accurate HDR was catalyzed by both *Sp*Cas9 and *As*Cas12a, and
these accounted for a minority of the payload integration events. The two sides of the
programmed *As*Cas12a double-strand break were repaired by apparently separate pathways since
mutant alleles that included only the left or only the right side of the donor transgene were
frequent although accurate HDR was significantly more likely in the presence of the NT
compared with T donor. Overall, *As*Cas12a was more efficient than *Sp*Cas9 for RNA-guided
knockout whereas both *Sp*Cas9 and *As*Cas12a catalyzed similar levels of accurate transgene
insertion at this model *ω1* gene target. These findings may be expected based on reports with
other cells and species that indicate that integration of repair template is not always precise (Paix
et al, 2017; Ranawakage *et al*, 2020) and *Sp*Cas9 shown even higher levels of HDR correction
than *As*Cas12a (Santos *et al.*, 2021), including in HEK293T and pluripotent stem cells where
mis-integrations patterns indicate that two sides of a DSB are repaired by separate cellular repair
pathways including the involvement of microhomology as a driver of the mis-integration.

In addition to its guide RNA programmed gene editing and double stranded DNA cleavage
activity, *As*Cas12a and related Type 2 subtypes V and VI nucleases display target substrate
bound non-specific trans-activity against ssDNA, dsDNA and/or RNA (Fuchs *et al*, 2019;
Nguyen *et al*, 2020). Accordingly, we speculated that activated *As*Cas12a associated with the
target exon 1 site in omega-1 may digest the donor ssODN provided as the repair template. If so,
investigation of comparative efficiency of *As*Cas12a *versus Sp*Cas9 catalyzed knock-in would
have been compromised. To investigate this possibility, we performed a series of in vitro assays

using non-activated and activated *As*Cas12a. Under the conditions and time frame of the
reactions (commercial *As*Cas12a, 37°C, 100 mM NaCl, 120 min reaction duration), we failed to
detect any marked degradation of the ssODN. Although this assay did not reflect the condition *in*
vivo in the transfected schistosome eggs, lack of apparent *As*Cas12a trans-activity *in vitro* may
not accurately reflect the situation within the nucleus of the cells within the egg: the RNP may
not have been retained at the programmed target site - degraded by cellular enzymes and/or
displaced by the cellular repair machinery-following cleavage before trans activity damaged the
donor ssODN and/pr other phenomena including protection of cellular DNA by histones.
Nonetheless, both *Sp*Cas9 and *As*Cas12a delivered similar knock-in performance in the eggs.

*As*Cas12a provided enhanced programmed knockout activity compared to *Sp*Cas9 at a specific
target site in the *omega-1* gene locus but both exhibited similar performance for HDR based
transgene insertion. Whereas these findings do not enable extrapolation to similarly superior
performance of *As*Cas12a for CRSISR/Cas-based gene editing of other schistosome genes and
for insertion of transgenes, and yet given the central role of the schistosome egg in
schistosomiasis, the approach demonstrated here does expand the toolkit for functional genomics
in helminth parasites (Sankaranarayanan *et al.*, 2021; Zamanian & Andersen, 2016). *As*Cas12a
will find use or gene editing AT rich genomes, such as in schistosomes and other parasitic
platyhelminths (International Helminth Genomes, 2019) although optimizing editing protocols
including time, temperature, metal ion concentration of editing reactions, donor format, donor
length, and guide RNA modifications, and other parameters will influence performance (Chen *et*
al., 2018; Hendel *et al*, 2015; Swarts & Jinek, 2018). Lastly, since transgene insertion into the
zygote within the newly released egg has already been described and shown to enable
establishment of lines of transgenic schistosomes, advances with gene editing approaches to this
key developmental stage of the schistosome life cycle portend progress in the near future with
informative gain-of-function lines of transgenic schistosomes.

Materials and Methods

Reagent/Resource	Reference or Source	Identifier or Catalog Number
Experimental Models		
Schistosoma mansoni eggs	Schistosomiasis Resource Center, Biomedical Research Institute, Maryland, USA	
Swiss Inbred mice (*M. musculus*)	The Jackson Laboratory	SWR/J
Recombinant DNA		
rocombinant omega-1 protein	Dr. Ron Hokke	Gifted from Dr. Ron Hokke
Antibodies		
Polyclonal antibody against recombinant omega-1 protein	Research Resource Identifier (RRID)	AB_2889886
goat anti-rabbit IgG (H+L) horse radish peroxidase conjugate	Life Technologies, Thermofisher Sciences	Cat # 31460
Oligonucleotides and sequence-based reagents		
PCR primers	This study	Synthesized from IDT
ssODN	Ittiprasert *et al*., 2019 and this study	Ultramer DNA Oligos, Synthesized from IDT
synthetic guide RNA	Ittiprasert *et al*., 2019 and this study	Synthesized from IDT
Chemicals, enzymes and other reagents		
Alt-R® CRISPR-Sp HiFi Cas9 enzyme V3	Integrated DNA Technologies	1081060
Alt-R® CRISPR-AsCas12a enzyme V3	Integrated DNA Technologies	1081068
DNAzol® ES reagent	Molecular Research Center	DN127
RNAzol® RT Reagent	Molecular Research Center	RN190
Opti-MEM	FisherScientific	Gibco™ 31985062
DMEM	FisherScientific	88364
fetal bovine serum	FisherScientific	Gibco™ 26140079
100x Penicillin-Streptomycin-Fungizone	FisherScientific	SV30079.01
NucleoSpin® Gel and PCR clean up	Takara	740609

SsoAcvanced Universal SYBR Green Supermix	BioRad	1725271
RIPA buffer	Millipore-Sigma	R0278-500ML
Bradford reagent	BioRad	5000201
SDS-PAGE polyacrylamide gradient gels, 4-12% gel, w/v	ThermoFisher Scientific	NP0322PK2
Maxima First Strand cDNA Synthesis Kit for RT-qPCR, with dsDNase	ThermoFisher Scientific	K1672
Peroxidate substate; Clarity^{TM} Western ECL Substrate	BioRad	1705062S
GoTaq® Green Master Mix	Promega	M712
NEB buffer 2.1	NEB	B720S
Proteinase K	ThermoFisher Scientific	EO0491
Agilent DNA 1000 Kit	Agilent Technologies	5067-1505
Agilent small RNA kit	Agilent Technologies	5067-1548
Software		
CHOPCHOP	http://chopchop.cbu.uib.no/	
TIDE	https://tide.deskgen.com/	
CRISPREsso2	https://crispresso.pinellolab.partners.org/submission	
SnapGene	https://www.snapgene.com/	Academic Subscription Plan
GraphPad Prism	Prism software	Monthly Subscription Plan
Other		
CFX96 Touch Real-Time PCR Detection System	BioRad	
ElectroSquarePorator	BTX	ECM830
Chemi-Doc MP System	BioRad	
BioAnalyzer	Agilent Technology	

372
373

Ethics statement

Mice experimentally infected with *S. mansoni*, obtained from Schistosomiasis Resource Center (SRC) at the Biomedical Research Institute (BRI), MD were housed at the Animal Research Facility of the George Washington University (GWU), which is accredited by the American Association for Accreditation of Laboratory Animal Care (AAALAC no. 000347) and has an Animal Welfare Assurance on file with the National Institutes of Health, Office of Laboratory Animal Welfare, OLAW assurance number A3205-01. All procedures employed were consistent with the Guide for the Care and Use of Laboratory Animals. The Institutional Animal Care and Use Committee (IACUC) at GWU approved the protocol used for maintenance of mice and recovery of schistosomes. Swiss albino mice were euthanized seven weeks after infection with *S. mansoni*, livers were removed at necropsy, and schistosome eggs recovered from the livers (Dalton *et al*, 1997). The 'liver eggs, termed LE' population is expected to include eggs of a spectrum of age ranging from newly released by the female schistosome through to mature eggs containing a fully developed miracidium (Jurberg *et al*, 2009). LE were maintained in DMEM medium supplemented with 10% heat-inactivated fetal bovine serum (FBS), 2% streptomycin/penicillin at 37°C under 5% CO_2 in air for 18-24 hours before use in gene editing assays (below) (Dalton *et al.*, 1997; Ittiprasert *et al.*, 2019; Lu *et al*, 2021; Mann *et al*, 2010; Mann *et al*, 2014).

CRISPR/Cas target design and single stranded DNA donors

The *ω1*-gene specific gRNA for *Sp*Cas9 and *As*Cas12a were designed to target all four copies of the hepatotoxic ribonuclease *omega-1* gene. All four are located on *S. mansoni* genome update annotation of version 7 (updated 16-September-2021 on WormBase ParaSite, https://parasite.wormbase.org/Multi/Search/Results?species=all;idx=;q=Hepatotoxic%20ribonuclease%20omega-1;site=ensemblunit&filter_species=*Schistosoma_mansoni*_prjea36577); Smp_334170.1, Smp_334170.2, Smp_334240 and Smp_333930 by an online CRISPR design tool, CHOCHOP (Labun *et al*, 2016; Labun *et al*, 2019; Montague *et al*, 2014). There are 99-100% sequence and coding sequence identity among those copies (Fig S1A), which encoding an enzyme of 115amino acids in length, respectively. These *ω1* copies are located on chromosome 1 at nucleotide (nt) position 3980364 to 3984675, 3992964 to 3995246, and 3908953 to 3911250, respectively. The guide RNA (gRNA), which targets exon 1 of *ω1* complimentary to the catalytic active site histidine [within FRKHEFEKHGLCAVEDPQV] codons, for the gRNAs programming both *Sp*Cas9 and *As*Cas12a. In addition, the gRNAs exhibited similarly high predicted CRISPR efficiency scores along with an absence of predicted off-target cleavages, as predicted using the CHOPCHOP algorithm using Smp_334170.2 as template (Fig 1A and B). Specifically, the sequence of the gRNA used with *Sp*Cas9 was 5'-GCATGGTTTGTGTGCAGTTG-3', 20 nt in length and that for *As*Cas12a was 5'-AAAAGCATGGTTTGTGTGCA -3', 20 nt on exon 1 of Smp_334170.2, Smp_334240 and Smp_333930 or exon 4 of Smp_334170.1 (Fig S1A) The expected cut sites of two nucleases are discrete, being 3 and 2 nt apart from each other on the non-target (NT) and target (T) strands, respectively (Fig 1A).

Single stranded oligonucleotide donors (ssODN) were designed to insert the 24 bp encoding six-stop codons into the programmed cleavage target. The homology arms used in the ssODN donor with *Sp*Cas9 was the non-target strand flanking both the 3' and 5' sides of the blunt ended DSB,

termed here *Sp*Cas9-ssODN. The homology arms of the two donor ssODNs used with *As*Cas12a
(cut at 18 nt and 23 nt on NT and T strands, leading to a sticky ended DSB) were either the target
stand or the non-target strand sequences (Fig 1B and C), here termed *As*Cas12a-NT-ssODN and
*As*Cas12a-T-ssODN, respectively. The complementary reverse of the 24 nt stop codon cassette
was used in the *As*Cas12a-T-ssODN donor, aiming for HDR insertion of the stop codons in the
translated forward strand of the gene (Fig 1C).

CRISPR reagents

Alt-R® CRISPR-*Sp*Cas9 and Alt-R® CRISPR-*As*Cas12a reagents (Alt-R® S.p. HiFi Cas9
nuclease V3, synthetic gRNA (sgRNA) (Labun *et al.*; Labun *et al.*; Nishimasu *et al.*), Alt-R® As
*As*Cas12a V3 and crRNA), and ssODN HDR donor templates (Ultramer DNA oligos) and Alt-
R® *Sp*Cas9 or *As*Cas12a Electroporation enhancer buffers were from Integrated DNA
Technologies (IDT) (Coralville, IA). Figure 1C provides the nucleotide sequences of the gRNAs
and donor ssODNs.

Ribonucleoprotein (RNP) assembly and delivery by electroporation

Stock solutions of the gRNAs for use with *Sp*Cas9 and *As*Cas12a and of the ssODNs were
prepared at 1.0 µg/µl in Opti-MEM medium (Sigma, St. Louis, MO). Each gRNA was
combined with the respective nuclease at 1:1 ratio to form an RNP complex (Ittiprasert *et al.*,
2019). The mixture was incubated at 23°C for 15 min, after which 3 µl of Alt-R® *Sp*Cas9 or
*As*Cas12a electroporation enhancer was added. The mixture was transferred to a chilled 4 mm
electroporation cuvette (BTX), containing ~ 5,000 LE in 100 µl Opti-MEM. LE in the presence
of RNPs and the other reagents were subjected to square wave electroporation with a single 125
volts, 1 pulse for 20 ms (ElectroSquarePorator, ECM830, BTX San Diego, CA). In KI assays, 6
µg of ssODN was included in the cuvette before electroporation. Following delivery of the
RNPs, LE were maintained for 10 days at 37°C, 5% CO_2 in air. In addition to experimental gene-
edited groups, control groups included LE electroporated in Opti-MEM (mock), mixed-Cas
enzymes only, mixed-sgRNAs only, and mixed-ssODNs only. Experimental and control groups
were otherwise treated similarly.

PCR to detect a multiple stop codon containing transgene

Genomic DNAs were extracted from LE using the DNAzol RT reagent (Molecular Research
Center Inc, Cincinnati, OH), and the concentration and purity determined by spectrophotometry
(Nanodrop 1000, Thermo Fisher Scientific Inc., Waltham, MA). The HDR target site(s) on
either the 5' or 3' KI transgene was amplified independently with two pairs of primers (Fig 2A):
1) 5'KI: 5'int-F and 24nt-6stp-R primers and 2) 3'KI: 24nt-6stp-F and 3'int-R using the GoTaq
DNA polymerase mix (Promega, Madison, WI) and 200 nM of each primer. The thermal cycling
included denaturation at 95°C, 3 min followed by 30 cycles of 94°C, 30s, 56°C, 30s and 72°C,
30s and a final extension for 5 min at 72°C. Thereafter, the amplicons (10 µl) were size separated
by agarose gel electrophoresis, stained with ethidium bromide, and imaged using the ChemiDoc
Imaging System (Bio-Rad, Hercules, CA). The expected sizes for the 5'KI and 3'KI amplicons
were 235 and 182 bp, respectively (Fig 2D). A control PCR to amplify ~790 bp flanking the
DSB was included.

Real time qPCR to estimate transgene copy numbers

A SYBR green quantitative PCR-based approach was used to estimate transgene copy numbers
using two pairs of primers; 5'KI and 3'KI as detailed (Fig 2A and D) with the SsoAdvanced
Universal SYBR Green Supermix (Bio-Rad). Briefly, one ng of gDNA was amplified with 10 µl
of 2× SsoAdvanced Universal SYBR Green Supermix and 200 nM of each primer, using the
CFX Connect Real-Time PCR Detection System (Bio-Rad). PCR conditions included
denaturation at 95°C, 3 min followed by 30 cycles of 94°C, 30 s, 56°C, 30 s and 72°C, 30 s and a
final extension at 72°C for 5 min, with the signals collected at the annealing step in each cycle.
Analysis of transgene copy number in the pooled gDNAs from LE involved comparison of the
qPCR signals obtained using a 10-fold serial dilution series of known quantities of the 24 nt six-
stop codon transgene in 235 bp (5' integration PCR fragment) ligated-pCR4-TOPO plasmid
DNA (size 4,191 bp) to establish a standard curve (Shirvani *et al*, 2020). The standard curve was
based on a linear regression generated using logarithm-10 transformed initial DNA input as the
dependent variable and the Ct number of the qPCR signal as the independent variable (Ct values
in each dilution were measured in triplicate). Ct numbers were used to fit into the linear
regression to derive the estimation of transformed DNA amount for the transgene (Lee *et al*,
2006; Yuan *et al*, 2007). The concentration of the 5'KI-transgene ligated-pCR4-TOPO plasmid
was measured (NanoDrop 1000) and the corresponding copy number was calculated as follows:

$$\text{DNA copy number} = \frac{6.02 \times 10^{23} \text{ (copy/mol)} \times \text{DNA quantity (g)}}{\text{DNA length (bp)} \times 660 \text{ (g/mol/bp)}}$$

Data were collected from four biological replicates. 5'KI and 3'KI transgene copy numbers
from the *Sp*Cas9 and *As*Cas12a treatments were plotted using GraphPad Prism v9 (Fig 2E and
F).

Tracking of INDELs by decomposition (TIDE) analysis

Genomic DNAs from LE exposed to the RNPs were isolated as above. The target *omega-1* locus
in the replicates of gDNAs was amplified using a control primer pair (Fig 2A). The amplicon of
790 bp in size from control and experimental groups, both KO and KI focused manipulations,
was purified by NucleoSpin® Gel and PCR clean up (Macherey-Nagel, Bethlehem, PA) after
which the nucleotide sequence determined by the Sanger method (Genewiz, South Plainland,
NJ). Sanger sequence traces from the 12 independent replicates of the experimental groups for
both *Sp*Cas9 or *As*Cas12a- (Jacobsen *et al.*, 2020) and *Sp*Cas9-catalyzed gene editing
manipulations were analyzed and compared with reads from the control (mock) group using the
Tracking of Indels by Decomposition (TIDE) algorithm (Brinkman *et al*, 2014; Brinkman & van
Steensel, 2019; Meshalkina *et al*, 2020). TIDE provides a rapid and informative assay and also
inform decisions on whether proceed to more detailed high throughput, next generation
sequencing (NGS) of targeting amplicons (Germini *et al*, 2018).

On-target amplicon next-generation sequencing

Subsequently, the Amplicon-EZ next generation sequencing approach (Genewiz) was used for
deeper coverage of the *ω1* coding sequence mutations and provided > 50,000 reads per sample.

Amplicons of 426 bp obtained with the primers bearing Illumina partial adapter sequences at the
5' end - NGS-F 5'- TATTGTCAACGGCGTACAGG-3' and NGS-R 5'-
CAATCGGCACTGAGACGCA-3' - flanking the programmed DSBs for CRISPR/*Sp*Cas9 and
CRISPR/*As*Cas12a were sequenced using Illumina chemistry. Partial Illumina adapters were
added to 5' end of the forward and reverse primers forward sequencing read: 5'-
ACACTCTTTCCCTACACGACGCTCTTCCGATCT-3', reverse sequencing read: 5'-
GACTGGAGTTCAGACGTGTGCTCTTCCGATCT-3'. The amplicons were purified using the
NucleoSpin® Gel and PCR clean up kits from Macherey-Nagel (Bethlehem, PA), after which
quantification and qualification of purified band(s) were undertaken using the Agilent DNA 1000
kit and BioAnalyzer (Agilent). Illumina NGS was performed with 2×250 bp sequencing without
fragmenting the amplicons. Raw reads in FASTQ format were analyzed for mutations resulting
from NHEJ and HDR by CRISPREsso2 (Clement *et al*, 2019) using the reference *ω1* sequences,
Smp_334170 and Smp_333930 including Smp_333870 (this latter is annotated as an
uncharacterized protein but shares >99% identity) (Fig S1B). The default parameters were used
to estimate indel and HDR events. The sequence reads from the Amplicon-EZ runs are available
at GenBank Bioproject PRJNA415471, BioSample SAMN07823308, SRA study SRP126685,
accessions SRR: 13498338-13498342, 15971309-15971320, and 6374209-6374210.

Quantitative real time PCR

Total RNA of LE was isolated using the RNAzol RT reagent (Molecular Research Center)
according to the manufacturer's instruction. RNA concentration and purity were determined
(Nanodrop 1000 spectrophotometer, Thermo Fisher). Ten nanograms of RNA was treated with
DNase I to remove residual DNA then converted to cDNA by Maxima First Strand cDNA
Synthesis Kit (Thermo Fisher Scientific). The *ω1*-specific product was amplified using
SsoAdvanced Universal SYBR Green Supermix (Bio-Rad) on CFX96 real time PCR (Bio-Rad)
using the primers as described (Ittiprasert *et al.*, 2019), ω1-RT-F: 5'-
GGTTACTTTAAATTTGGTA-3' and ω1-RT-R; 5'-AGTTCCAAGGAACGGGCAG-3'.
These primers amplify all annotated *ω1* transcript copies; Smp_334170.1, Smp_334170.2,
Smp_334240 and Smp_333930 (Fig S1C). The PCR reaction was denatured at 95°C for 30 s, 40
amplification cycles each consisting of denaturation at 95°C for 15 s and annealing/extension at
60°C for 30 s. The output was analyzed in CFX manager software (Bio-Rad). Relative
expression of ω1 was calculated using the $2^{-\Delta\Delta Ct}$ method and normalized to schistosome GAPDH
(Smp_0569701) expression using specific primers; forward 5'-
ATGGGACATTTCCAGGCGAG-3', reverse 5'-CCAACAACGAACATGGGTGC-3' (Livak &
Schmittgen, 2001). Data four biological replicates were compared to wild type LE and fold
change reported as mean ± SE (n = 4).

Western blot analysis

Soluble egg antigen (SEA) of LE was extracted by mechanical lysis (Funakoshi motorized
pestle; Diagnocine, Hackensack, NJ) in RIPA buffer (Millipore Sigma) supplemented with
protease inhibitors (Protease Inhibitor Cocktail I, Millipore Sigma), as described (Ittiprasert *et*
al., 2019). The lysate was clarified by centrifugation at 13,000 rpm, 15 min, 4°C after which
protein concentration was determined by the Bradford method (Kielkopf *et al*, 2020). Ten
micrograms of SEA were size-separated through SDS-PAGE polyacrylamide gradient gels, 4-
12% gel, w/v (Bolt Bis-Tris Plus, Invitrogen, Thermo Fisher Scientific) before transfer to PVDP

membranes (Transblot Turbo Transfer System, Bio-Rad). Polyclonal anti-ω1 IgG antibodies were purified from the rabbit anti-ω1 sera, raised as follows. Two NZ white rabbits, E12983 and E12984, were immunized with 30 μg of recombinant *omega-1* RNase T2 of *Schistosoma mansoni* (rω1) (produced *in planta* by agroinfiltration of leaves of *Nicotiana benthamiana* (Wilbers *et al*, 2017)) in Complete Freund's Adjuvant and boosted four times at two-week intervals using 15 μg rω1 in Incomplete Freund's Adjuvant each time. Blood was collected 66 days after the first immunization, IgGs was isolated from sera using an immobilized protein A ligand. The concentrations of the purified antibodies were 1.31 mg/ml (E12983) and 1.51 mg/ml (E12984). These antibodies have been assigned Research Resource Identifier (RRID) AB_2889886 (Bandrowski *et al*, 2016).   PVDF membranes were probed with the E12983 IgG antibody, diluted 1:1,000, for 18 h at 4°C in PBS-Tween (0.5%), 5% non-fat powdered milk, washed 3 times in PBS-Tween (wash buffer), and probed with goat anti-rabbit IgG (H+L) horse radish peroxidase conjugate (Life Technologies, Thermo Fisher Sciences), diluted 1:5,000, for 60 min at 23°C with gentle shaking. Subsequently, following washing (3 times in wash buffer), the membrane was exposed to a peroxidase substrate (Clarity Western ECL Substrate, Bio-Rad) for 5 min at 23°C, after which chemiluminescence was quantified and imaged (Chemi-Doc MP System, Bio-Rad). The Image Lab software (Bio-Rad) was used to quantify the ω1 signals. The percentage of ω1 expression from the experimental and control groups was established by comparison with that of wild type SEA, assigned as 100%.

Single stranded DNA donors as a potential substrate for indiscriminate *trans* activity of activated *As*Cas12a

Previous reports reveal the *trans*-activity of wild type *As*Cas12a prefers ssDNAs over dsDNAs, and was inactive toward dsDNA in the presence of magnesium (II) ions (Li *et al*, 2020). The *As*Cas12a exhibited not only endonuclease activity on both dsDNA, *cis*- and *trans*-activity (Li *et al*, 2018b). Accordingly, to investigate whether activated *As*Cas12a could degrade the ssODN donors used in this study by indiscriminate *trans*-activity, several *in vitro As*Cas12a-RNP digestion assays were carried out. These included the dsDNAs containing the *As*Cas12a target sequence used in this study and 5'PAM (TTTG) as positive enzymatic activity. A 426 bp dsDNA template was amplified using the NGS study (above) primers. Activity of *As*Cas12a-RNP against the dsDNA template released fragments of 265 and 161 bp (Fig 2B), at 37°C for ≥30 min as predicted in 1× NEB Buffer 2.1 (500 mM NaCl, 20 mM sodium acetate, 0.1 mM EDTA, 0.1 mM tris(2-carboxyethyl) phosphine (TCEP), 50% glycerol). *As*Cas12a activity was inactivated by the addition of Proteinase K (1 ul at 20 mg/ml) at 56°C for 10 min. Electropherograms of the digestion products were undertaken using the BioAnalyzer (Agilent DNA1000) to visualize the size templates and fragments.

To investigate indiscriminate *trans*-cleavage of ssODNs, similar *in vitro* digestion reaction was carried out, as above, but with the addition of |50 ng of either *As*Cas12a-NT-ssODN or *As*Cas12a-T-ssODN. The *As*Cas12a-RNP complex in 1× NEB Buffer 2.1 and ssODN were incubated at 37°C for 30, 60 and 120 min before stopping the reaction by addition of proteinase K at 56°C for 10 min, as visualized as above. The ssDNA ladder (SimPlex Sciences, NJ) ranging from 10-200 bp was included as the single stranded DNA marker. *Trans*-activity of Alt-R *As*Cas12a V3-RNP complex was not apparent against the ssODNs under these conditions *in vitro* (Fig 2C).

Statistical analysis

Statistical analysis was performed using GraphPad Prism version 9.1 (GraphPad Prism, La Jolla, CA). Graphs were plotted as the mean with bars to show the standard error of the mean. The differences among groups were assessed by one-way ANOVA. P-values ≤ 0.05 considered to be the statistically significant.

GenBank/EMBL/DDBJ

Database accessions

cknowledgements

We thank Professor Cornelis Hokke, Leiden University Medical Center, The Netherlands for the gift of recombinant *Schistosoma mansoni* omega-1. We thank the Thailand Research Fund, the Royal Golden Jubilee Ph.D. program, and Dr. Dalina Tanyong, Mahidol University, Thailand, for support for CC to perform the work at GWU. This research was funded in whole, or in part, by the Wellcome Trust (award number 107475/Z/15/Z, titled 'Functional Genomics Flatworms Initiative', K. H. Hoffmann, principal investigator). For the purpose of Open Access, the authors have applied a CC BY public copyright license to any Author Accepted Manuscript version arising from this submission. Schistosome-infected mice and snails were provided by the NIAID Schistosomiasis Resource Center of the Biomedical Research Institute, Rockville, Maryland through NIH-NIAID Contract HHSN272201000005I for distribution through BEI Resources.

Author contributions.

WI, CC, and PJB designed the experiments, analyzed the data, and drafted the manuscript with input from all the co-authors; CC and WI performed the gene editing focused experiments; VHM, WL, AM, TO, YNA and MMK contributed the parasite materials to the experiments, to data collection, and reviewed and edited the manuscript; PJB and WI supervised the project. PJB, WI and MMK arranged the funding.

Conflict of interest

The authors declare that do not have competing interests.

Figure legends

Figure 1. Schematic of the *S. mansoni* hepatotoxic ribonuclease omega 1 (ω1) gene structure, target sites for the *Sp*Cas9 and *As*Cas12a, ssODN donors and partial amino acid sequence with histidine active site.

A   Diagram of the ω1 gene (Gene ID; Smp_334170.2 of *S. mansoni* annotated genome version 7_2021) and CRISPR/Cas target sites on coding sequencing of the first exon coding for

histidine active site, FRKHEFEKHGLCAVEDPQV. The relative positions of crRNA for
CRISPR-*Sp*Cas9 (red arrow) and -*As*Cas12a (blue arrow) are indicated along with their
3'PAM (yellow arrow) and 5'PAM (yellow arrow), respectively.

B   Programmed cleavage site for *Sp*Cas9/sgRNA ~3nt upstream of PAM 3'TGG (red arrow);
and cleavage site for *As*Cas12a/crRNA at 18-24 nt downstream of a prospective 5-'TTTV)
(blue arrow). The programmed cleavage sites for both enzymes, *Sp*Cas9 and *As*Cas12a,
were three to six nt distant from each other. Fifty residues upstream and downstream of the
double strand break (DSB) were included as the homology arms (HA) of the six stop codon
cassette transgene (payload/cargo). Because cleavage by *Sp*Cas9 liberates blunt ended
DNA, only a non-target strand (NT) repair template was provided for HDR (red bar). Sticky
ended DNA results from cleavage by *As*Cas12a and hence both non target and target strand
(T) repair templates for HDR were provided (light and dark blue bars).

C   The DNA donor sequences of CRISPR/*Sp*Cas9 and CRISPR/*As*Cas12a with non-target
(NT) or target (T) sequence HA where the *As*Cas12a-NT-ssODN donor repairs the anti-
sense DNA and *As*Cas12a-T-ssODN donor repairs the sense stand.

Figure 2. Knock-in by homology directed repair.

A   The location of primers used for 5'- and 3' transgene (purple lines) knock-in PCR (5'int-F
with 24nt-6stp-R and 24nt-6stp-F with 3'int-R) and internal positive control (control-F and
-R). The primers for next generation sequencing indicate as NGS-F and NGS-R. All
primers locates further from HA (purple boxes) to protect the false positive PCR from
DNA donor and 24 nt-6stp-F and -R primers are highly specific to stop codon transgene,
not schistosome genome.

B   The *As*Cas12a target recognition activates specific 426 bp dsDNA amplicon template (lane
1) cleavage into 265bp and 161 bp (lane 3). The crRNA/*As*Cas12a RNP complex is ~40 nt
are in lanes 2 and 3.

C   There were no specific ssDNA cleavage on either *As*Cas12a-NT-ssODN or *As*Cas12a-T-
ssODN donors at 124 nt as in lanes 1 and 2, respectively. The crRNA/*As*Cas12a RNP (~40
nt in lanes 3-9) containing activated *As*Cas12a enzyme shown no random ssDNA digestion
at 30 min, 60 and 120 min in lanes 4-5, 6-7 and 8-9, respectively. Lanes 4, 6 and 8 shown
the uncut 124 nt of *As*Cas12a-NT-ssODN (top band) with *As*Cas12a-crRNA (lower band),
and lanes 5, 7 and 9 shows 124 nt of *As*Cas12a-T-ssODN (top band) with *As*Cas12a-
crRNA (lower band).

D   The PCR detection of the integration sites in the transgenic eggs and control. The
amplified control product (lanes C) for the wildtype sequence was 790 bp, while those for
the 5'(lanes 5') and 3' (lanes 3') flanking regions of the transgenic sequence were 235 bp
and 182 bp from KI introducing CAS-9 ssODN, *As*Cas12a-NT-ssODN and *As*Cas12a-T-
ssODN, respectively. Lane M is a 1 kb Plus DNA ladder; Mock (only buffer
electroporation) used as the negative transgene integration control.

E and F   The estimation of transgene copy number analysis in egg DNA were calculated by
qPCR comparing with the known amount of 24 nt transgene in 235 bp (5' integration PCR

fragment) ligated-pCR4-TOPO plasmid DNA (size 4,191 bp) standard curve; y = -
4.0964ln(10) + 44.821, R² = 0.9923 (data not shown). There were the confirmation of 3'
integration of 3.61±0.25, 4.44±0.84 and 4.19±1.22 of transgene copy from KI_*Sp*Cas9,
KI_*As*Cas12a-NT-ssODN and KI_*As*Cas12a-T-ssODN, respectively with no statistic
significant (one-way ANOVA, multiple comparisons, 95% CI of diff, P Value at 0.4057,
0.6264 and 0.9155). While there was asymmetric HDR revealed by 5' transgene integration
of 2.72±0.42, 5.06±0.54 and 1.64±0.39 copy number from KI_*Sp*Cas9, KI_*As*Cas12a-NT-
ssODN and KI_*As*Cas12a-T-ssODN, respectively. The asterisk *,*** and **** indicates
statistic significant difference with $P = 0.0214$, $P=0.0001$ and $P < 0.0001$, respectively by
one-way ANOVA, multiple comparison. The control primers used for amplicon
preparation from both control and experimental groups, then their Sanger Direct
Sequencing results were pre-investigated for CRISPR efficiency by TIDE analysis prior to
proceed for NGS.

G and H   5'int-F and 3'int-R are genome-specific primers, and 24nt-6stp-F and 24nt-6stp-R
are transgene-specific primers. From these results, we confirmed that programmed
CRISPR/*As*Cas12a gene editing was active in schistosome eggs, and higher than *Sp*Cas9 by
TIDE analysis with statistic significant (ANOVA, ****, P 0.0001, n = 12). The estimation
of CRISPR efficiency from the 12 biological replicates (each black dot in panels G and H)
KI_*Sp*Cas9 experiment (wide range of CRISPR efficiency from 0.46-2.23 among 12
biological replicates) was higher than KO experiment. The estimated CRISPR efficiency
in LE from CRISPR/*As*Cas12a (green dots in panels G and H, $n = 8$) shown more than
1.5% in most samples. There was no statistically significant differences in CRISPR
efficiency among KI_*Sp*Cas9 vs. KI_*As*Cas12a (panel H) from Sanger sequencing along
with TIDE analysis. Samples of the genomic DNAs were pooled, targeted amplicons
prepared, and Illumina NGS performed.

Figure 3. NHEJ assessment from *S. mansoni* eggs on target ω1 gene after *Sp*Cas9 and
***As*Cas12a RNP complex electroporation.** CRISPResso2 analysis of the Illumina sequence
reads revealed non-homologous end joining mutations at targeted loci in schistosome eggs.
Allele-specific editing outcomes of CRISPR-*Sp*Cas9 and -*As*Cas12a gene editing in schistosome
multiple copies of *ω1* Smp_334170.2, Smp_334170.1, Smp_334240 (100% gene sequence
identical; Fig. S1b). Mutagenesis frequencies for all loci including Smp_333930 and
Smp_333870 were included in CRISPResso2 analysis (Fig S1B). The results were from NGS
libraries of pooled genomic DNA from 7 biological replicates of ω1 gene editing experiments of
KO-*Sp*Cas9 (Panel A and C) and KO-*As*Cas12a (Panel B and D). The sequence reads assigned
to references, Smp_333930 and Smp_333870 loci by the software to achieve accurate
quantification of genome editing at genomic loci with multiple copies from *Sp*Cas9 and
*As*Cas12a. The bar chart shows the assignment of each read to the wild type (unmodified) and
NHEJ with 12.26%, 1.66% and 0.12% (total 14.04%) from Δω1KO-*Sp*Cas9 on references and
other two loci, respectively (panel a). Panel b shows the assigned NHEJ with 9.27%, 1.33% and
0.28% (total 10.88%) from Δω1-*As*Cas12a on references and two high identity of *ω1* gene loci
(panel b). Ambiguous aligments; 0.49% and 0.93% that could not be attributed uniquely to one
of these loci are shown in the last bar of each graph. The majority of NHEJ indels from both
nucleases were nucleotide substitutions followed by gene deletions (Panel C and D) on the
references. The mutations occurred in all ω1 gene copies with the most editing on the reference
target site. CRISPResso2 software interpreted ≤ 1.0% of the reads as ambiguous. Only deletions

were revealed in these experiment, -3 nt deletion downstream from expected cut side (red arrow)
was the majority of NHEJ mediated from *Sp*Cas9 (Panel C). The larger deletion from -2 to -26
nt at downstream of expected sticky end cut sites (red arrows) were revealed from NHEJ.

Figure. 4. HDR and induction of NHEJ at the ω1 target sites using ssODN template. Panel
A to C shown the percentage of NHEJ from egg samples that introduced RNP along with ssODN
template were higher than without ssODN. There were total 15.67% (13.22+1.16+0.52), 28.71%
(22.31+6.17+0.23) and 21.43% (9.62+11.56+0.25) from Δω1KI-*Sp*Cas9, Δω1KI-*As*Cas12a-NT-
ssODN and Δω1KI-*As*Cas12a-T-ssODN, respectively (blue or green bars in each graph).
Majority of mutations are substitutions and deletions toward to downstream of the expected cut
sites indicating by red arrows (Panel D, E and F). The various base deletions from -1 to -15 bp
found in Δω1KI-*Sp*Cas9, Δω1KI-*As*Cas12a-NT-ssODN egg DNAs, while large deletion as -40
bp found as a major deletion in Δω1KI-*As*Cas12a-T-ssODN. The percentage of perfect HDR in
3 experimental groups; Δω1KI-*Sp*Cas9, Δω1KI-*As*Cas12a-NT-ssODN and Δω1KI-*As*Cas12a-T-
ssODN were 17.07%, 14.58% and 12.37%, respectively (red bars, Panel G, H and I). The
imperfect HDRs occurred in all KI samples resulting from interrupting 5' end HDR gene repair
with 2-5 nucleotide base missing (black bars in Panel G, H and I) with small percentage at 0.94,
0.41 and 0.68.

Figure 5. Reduction of the expression of ω1 by quantitative real time PCR and immunoblot
analysis.

A   Reduction of ω1 transcript from Δω1KO-*Sp*Cas9, Δω1KI-*Sp*Cas9, Δω1KO-*As*Cas12a,
Δω1KI-*As*Cas12a NTS and Δω1KI-*As*Cas12a TS eggs comparing with control groups after
normalization with wild type egg RNA (100% as discontinued line). The means of four
independent biological replicates.

B and C   Soluble egg antigen; SEA (in duplicates) was solubilized, electrophoresed in
SDS-PAGE gels, transferred to nitrocellulose membrane, and probed with anti-recombinant
plant omega-1 (rω1) rabbit IgG. There was ω1 protein reduction in all SEA samples (~30
kDa as arrow, Panel B). The percentage of ω1 expression in both KO and KI experimental
samples were comparing ω1 amount of wild-type SEA (100%) (Panel C); (one-way
ANOVA, multiple comparison, *, $P< 0.05$, ***, $P< 0.001$, ****, $P<0.0001$).

Supporting information

Figure S1. **Gene structure encoding the hepatotoxic ribonuclease, omega 1 (ω1). At least**
five copies of the gene are located on chromosome 1, as apparent in the draft sequence of *S.*
***mansoni*, version 7, scaffold SM_V7_1:3,885,722-4,002,365.** The positions of several ω1
genes coding on updated annotation (year 2021) on WormBase Parasite, including two loci of
Smp_334170, Smp_334240, Smp_334070, Smp_3332220, Smp_333930, Smp_333870 and
Smp_179960 are shown for their physical relationship of the copies on the chromosome.

A   Map of gene structures and sizes of Smp_334170.1, Smp_334170.2 and Smp_334240 and
the CRISPR/Cas programmed target sites for *Sp*Cas9 (blue arrow) and *As*Cas12a (green
arrow).

B   All gene copies shown DNA sequence identity >99% as DNA sequence alignment and
target amplicon primer indicated as NGS-F and NGS-R. Only four of these gene copies;
Smp_334240, Smp_334170 and Smp_333930 were reported for Uniprot.

C   Alignment of mRNA sequences and primers used for gene transcript study (purple).

Table S1. The CRISPREsso2 analysis results from four negative controls; mixed sgRNAs only,
*Sp*Cas9 only, *As*Cas12a only and mixed donors only. There were over 180,000 reads from each
library. There was < 1% of background modified reads from all negative control samples.

Table S2. Nucleotide sequences of oligonucleotide primers used in the investigation.

References

An Y, Geng Y, Yao J, Fu C, Lu M, Wang C, Du J (2020) Efficient Genome Editing in Populus
Using CRISPR/Cas12a. *Front Plant Sci* 11: 593938

Arunsan P, Ittiprasert W, Smout MJ, Cochran CJ, Mann VH, Chaiyadet S, Karinshak SE, Sripa
B, Young ND, Sotillo J *et al* (2019) Programmed knockout mutation of liver fluke granulin
attenuates virulence of infection-induced hepatobiliary morbidity. *Elife* 8

Banakar R, Schubert M, Collingwood M, Vakulskas C, Eggenberger AL, Wang K (2020)
Comparison of CRISPR-Cas9/Cas12a Ribonucleoprotein Complexes for Genome Editing
Efficiency in the Rice Phytoene Desaturase (OsPDS) Gene. *Rice* 13: 4

Bandrowski A, Brush M, Grethe JS, Haendel MA, Kennedy DN, Hill S, Hof PR, Martone ME,
Pols M, Tan SC *et al* (2016) The Resource Identification Initiative: a cultural shift in publishing.
Brain Behav 6: e00417

Bolt BJ, Rodgers FH, Shafie M, Kersey PJ, Berriman M, Howe KL (2018) Using WormBase
ParaSite: An Integrated Platform for Exploring Helminth Genomic Data. *Methods Mol Biol*
1757: 471-491

Brinkman EK, Chen T, Amendola M, van Steensel B (2014) Easy quantitative assessment of
genome editing by sequence trace decomposition. *Nucleic Acids Res* 42: e168

Brinkman EK, van Steensel B (2019) Rapid Quantitative Evaluation of CRISPR Genome Editing
by TIDE and TIDER. *Methods Mol Biol* 1961: 29-44

Charpentier E (2015) CRISPR-Cas9: how research on a bacterial RNA-guided mechanism
opened new perspectives in biotechnology and biomedicine. *EMBO Mol Med* 7: 363-365

Charpentier E, Richter H, van der Oost J, White MF (2015) Biogenesis pathways of RNA guides
in archaeal and bacterial CRISPR-Cas adaptive immunity. *FEMS Microbiol Rev* 39: 428-441

Chen JS, Ma E, Harrington LB, Da Costa M, Tian X, Palefsky JM, Doudna JA (2018) CRISPR-
Cas12a target binding unleashes indiscriminate single-stranded DNase activity. *Science* 360:
436-439

Clement K, Rees H, Canver MC, Gehrke JM, Farouni R, Hsu JY, Cole MA, Liu DR, Joung JK,
Bauer DE *et al* (2019) CRISPResso2 provides accurate and rapid genome editing sequence
analysis. *Nat Biotechnol* 37: 224-226

Costain AH, MacDonald AS, Smits HH (2018) Schistosome Egg Migration: Mechanisms,
Pathogenesis and Host Immune Responses. *Front Immunol* 9: 3042

Dalton JP, Day SR, Drew AC, Brindley PJ (1997) A method for the isolation of schistosome
eggs and miracidia free of contaminating host tissues. *Parasitology* 115 (Pt 1): 29-32

Dumeau CE, Monfort A, Kissling L, Swarts DC, Jinek M, Wutz A (2019) Introducing gene
deletions by mouse zygote electroporation of Cas12a/Cpf1. *Transgenic Res* 28: 525-535

Everts B, Perona-Wright G, Smits HH, Hokke CH, van der Ham AJ, Fitzsimmons CM, Doenhoff
MJ, van der Bosch J, Mohrs K, Haas H *et al* (2009) Omega-1, a glycoprotein secreted by
Schistosoma mansoni eggs, drives Th2 responses. *J Exp Med* 206: 1673-1680

Fuchs RT, Curcuru J, Mabuchi M, Yourik P, Robb GB (2019) Cas12a trans-cleavage can be
modulated in vitro and is active on ssDNA, dsDNA, and RNA. *bioRxiv*: 600890

Germini D, Tsfasman T, Zakharova VV, Sjakste N, Lipinski M, Vassetzky Y (2018) A
Comparison of Techniques to Evaluate the Effectiveness of Genome Editing. *Trends Biotechnol*
36: 147-159

Gier RA, Budinich KA, Evitt NH, Cao Z, Freilich ES, Chen Q, Qi J, Lan Y, Kohli RM, Shi J
(2020) High-performance CRISPR-Cas12a genome editing for combinatorial genetic screening.
Nat Commun 11: 3455

Hendel A, Bak RO, Clark JT, Kennedy AB, Ryan DE, Roy S, Steinfeld I, Lunstad BD, Kaiser
RJ, Wilkens AB *et al* (2015) Chemically modified guide RNAs enhance CRISPR-Cas genome
editing in human primary cells. *Nat Biotechnol* 33: 985-989

Hoffmann KF, Brindley PJ, Berriman M (2014) Medicine. Halting harmful helminths. *Science*
346: 168-169

Hsu PD, Lander ES, Zhang F (2014) Development and applications of CRISPR-Cas9 for
genome engineering. *Cell* 157: 1262-1278

Huang CH, Lee KC, Doudna JA (2018) Applications of CRISPR-Cas Enzymes in Cancer
Therapeutics and Detection. *Trends Cancer* 4: 499-512

Hur JK, Kim K, Been KW, Baek G, Ye S, Hur JW, Ryu SM, Lee YS, Kim JS (2016) Targeted
mutagenesis in mice by electroporation of Cpf1 ribonucleoproteins. *Nat Biotechnol* 34: 807-808

International Helminth Genomes C (2019) Comparative genomics of the major parasitic worms.
Nat Genet 51: 163-174

Ittiprasert W, Mann VH, Karinshak SE, Coghlan A, Rinaldi G, Sankaranarayanan G, Chaidee A,
Tanno T, Kumkhaek C, Prangtaworn P *et al* (2019) Programmed genome editing of the omega-1
ribonuclease of the blood fluke, Schistosoma mansoni. *Elife* 8

Jacobsen T, Ttofali F, Liao C, Manchalu S, Gray BN, Beisel CL (2020) Characterization of
Cas12a nucleases reveals diverse PAM profiles between closely-related orthologs. *Nucleic Acids*
Res 48: 5624-5638

Jeon Y, Choi YH, Jang Y, Yu J, Goo J, Lee G, Jeong YK, Lee SH, Kim IS, Kim JS *et al* (2018)
Direct observation of DNA target searching and cleavage by CRISPR-Cas12a. *Nat Commun* 9:
2777

Jinek M, Chylinski K, Fonfara I, Hauer M, Doudna JA, Charpentier E (2012) A programmable
dual-RNA-guided DNA endonuclease in adaptive bacterial immunity. *Science* 337: 816-821

Jurberg AD, Goncalves T, Costa TA, de Mattos AC, Pascarelli BM, de Manso PP, Ribeiro-Alves
M, Pelajo-Machado M, Peralta JM, Coelho PM *et al* (2009) The embryonic development of
Schistosoma mansoni eggs: proposal for a new staging system. *Dev Genes Evol* 219: 219-234

Kellner MJ, Koob JG, Gootenberg JS, Abudayyeh OO, Zhang F (2019) SHERLOCK: nucleic
acid detection with CRISPR nucleases. *Nat Protoc* 14: 2986-3012

Kielkopf CL, Bauer W, Urbatsch IL (2020) Bradford Assay for Determining Protein
Concentration. *Cold Spring Harb Protoc* 2020: 102269

Kim D, Kim J, Hur JK, Been KW, Yoon SH, Kim JS (2016) Genome-wide analysis reveals
specificities of Cpf1 endonucleases in human cells. *Nat Biotechnol* 34: 863-868

Kissling L, Monfort A, Swarts DC, Wutz A, Jinek M (2019) Preparation and electroporation of
Cas12a/Cpf1-guide RNA complexes for introducing large gene deletions in mouse embryonic
stem cells. *Methods Enzymol* 616: 241-263

Koonin EV, Makarova KS, Wolf YI (2017) Evolutionary Genomics of Defense Systems in
Archaea and Bacteria. *Annu Rev Microbiol* 71: 233-261

Labun K, Montague TG, Gagnon JA, Thyme SB, Valen E (2016) CHOPCHOP v2: a web tool
for the next generation of CRISPR genome engineering. *Nucleic Acids Res* 44: W272-276

Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E (2019)
CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. *Nucleic Acids*
Res 47: W171-W174

Lee C, Kim J, Shin SG, Hwang S (2006) Absolute and relative QPCR quantification of plasmid
copy number in Escherichia coli. *J Biotechnol* 123: 273-280

Li B, Yan J, Zhang Y, Li W, Zeng C, Zhao W, Hou X, Zhang C, Dong Y (2020) CRISPR-
Cas12a Possesses Unconventional DNase Activity that Can Be Inactivated by Synthetic
Oligonucleotides. *Mol Ther Nucleic Acids* 19: 1043-1052

Li S-Y, Cheng Q-X, Wang J-M, Li X-Y, Zhang Z-L, Gao S, Cao R-B, Zhao G-P, Wang J
(2018a) CRISPR-Cas12a-assisted nucleic acid detection. *Cell Discovery* 4: 20

Li SY, Cheng QX, Liu JK, Nie XQ, Zhao GP, Wang J (2018b) CRISPR-Cas12a has both cis-
and trans-cleavage activities on single-stranded DNA. *Cell Res* 28: 491-493

Liu P, Luk K, Shin M, Idrizi F, Kwok S, Roscoe B, Mintzer E, Suresh S, Morrison K, Frazão JB
et al (2019) Enhanced Cas12a editing in mammalian cells and zebrafish. *Nucleic Acids Res* 47:
4169-4180

Liu Y, Han J, Chen Z, Wu H, Dong H, Nie G (2017) Engineering cell signaling using tunable
CRISPR-Cpf1-based transcription factors. *Nat Commun* 8: 2095

Livak KJ, Schmittgen TD (2001) Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. *Methods* 25: 402-408

Lu Z, Sankaranarayanan G, Rawlinson KA, Offord V, Brindley PJ, Berriman M, Rinaldi G
(2021) The Transcriptome of Schistosoma mansoni Developing Eggs Reveals Key Mediators in
Pathogenesis and Life Cycle Propagation. *Frontiers in Tropical Diseases* 2

Luhtala N, Parker R (2010) T2 Family ribonucleases: ancient enzymes with diverse roles. *Trends*
Biochem Sci 35: 253-259

Ma P, Meng Q, Sun B, Zhao B, Dang L, Zhong M, Liu S, Xu H, Mei H, Liu J *et al* (2020)
MeCas12a, a Highly Sensitive and Specific System for COVID-19 Detection. *Advanced Science*
7: 2001300

Malzahn AA, Tang X, Lee K, Ren Q, Sretenovic S, Zhang Y, Chen H, Kang M, Bao Y, Zheng X
et al (2019) Application of CRISPR-Cas12a temperature sensitivity for improved genome
editing in rice, maize, and Arabidopsis. *BMC Biol* 17: 9

Mann VH, Morales ME, Rinaldi G, Brindley PJ (2010) Culture for genetic manipulation of
developmental stages of Schistosoma mansoni. *Parasitology* 137: 451-462

Mann VH, Suttiprapa S, Skinner DE, Brindley PJ, Rinaldi G (2014) Pseudotyped murine
leukemia virus for schistosome transgenesis: approaches, methods and perspectives. *Transgenic*
Res 23: 539-556

Meshalkina DA, Glushchenko AS, Kysil EV, Mizgirev IV, Frolov A (2020) SpCas9- and
LbCas12a-Mediated DNA Editing Produce Different Gene Knockout Outcomes in Zebrafish
Embryos. *Genes (Basel)* 11

Montague TG, Cruz JM, Gagnon JA, Church GM, Valen E (2014) CHOPCHOP: a
CRISPR/Cas9 and TALEN web tool for genome editing. *Nucleic Acids Res* 42: W401-407

Mustafa MI, Makhawi AM (2020) SHERLOCK and DETECTR: CRISPR-Cas Systems as
Potential Rapid Diagnostic Tools for Emerging Infectious Diseases. *J Clin Microbiol*

Nguyen LT, Smith BM, Jain PK (2020) Enhancement of trans-cleavage activity of Cas12a with
engineered crRNA enables amplified nucleic acid detection. *Nature Communications* 11: 4906

Nguyen PQ, Soenksen LR, Donghia NM, Angenent-Mari NM, de Puig H, Huang A, Lee R,
Slomovic S, Galbersanini T, Lansberry G *et al* (2021) Wearable materials with embedded
synthetic biology sensors for biomolecule detection. *Nat Biotechnol*

Nishimasu H, Ran FA, Hsu PD, Konermann S, Shehata SI, Dohmae N, Ishitani R, Zhang F,
Nureki O (2014) Crystal structure of Cas9 in complex with guide RNA and target DNA. *Cell*
156: 935-949

Paix A, Folkmann A, Goldman DH, Kulaga H, Grzelak MJ, Rasoloson D, Paidemarry S, Green
R, Reed RR, Seydoux G (2017) Precision genome editing using synthesis-dependent repair of
Cas9-induced DNA breaks. *Proc Natl Acad Sci U S A* 114: E10745-e10754

Pinello L, Canver MC, Hoban MD, Orkin SH, Kohn DB, Bauer DE, Yuan GC (2016) Analyzing
CRISPR genome-editing experiments with CRISPResso. *Nat Biotechnol* 34: 695-697

Protasio AV, Tsai IJ, Babbage A, Nichol S, Hunt M, Aslett MA, De Silva N, Velarde GS,
Anderson TJ, Clark RC *et al* (2012) A systematically improved high quality genome and
transcriptome of the human blood fluke Schistosoma mansoni. *PLoS Negl Trop Dis* 6: e1455

Ranawakage DC, Okada K, Sugio K, Kawaguchi Y, Kuninobu-Bonkohara Y, Takada T,
Kamachi Y (2020) Efficient CRISPR-Cas9-Mediated Knock-In of Composite Tags in Zebrafish
Using Long ssDNA as a Donor. *Front Cell Dev Biol* 8: 598634

Sankaranarayanan G, Berriman M, Rinaldi G (2021) An uneven race: genome editing for
parasitic worms. *Nat Rev Microbiol* 19: 621

Santos L, Mention K, Cavusoglu-Doran K, Sanz DJ, Bacalhau M, Lopes-Pacheco M, Harrison
PT, Farinha CM (2021) Comparison of Cas9 and Cas12a CRISPR editing methods to correct the
W1282X-CFTR mutation. *J Cyst Fibros*

Schwartz C, Fallon PG (2018) Schistosoma "Eggs-Iting" the Host: Granuloma Formation and
Egg Excretion. *Front Immunol* 9: 2492

Sentmanat MF, Peters ST, Florian CP, Connelly JP, Pruett-Miller SM (2018) A Survey of
Validation Strategies for CRISPR-Cas9 Editing. *Sci Rep* 8: 888

Shirvani R, Barshan-Tashnizi M, Shahali M (2020) An investigation into gene copy number
determination in transgenic yeast; The importance of selecting a reliable real-time PCR standard.
Biologicals 65: 10-17

Smith CW, Nandu N, Kachwala MJ, Chen YS, Uyar TB, Yigit MV (2020) Probing CRISPR-
Cas12a Nuclease Activity Using Double-Stranded DNA-Templated Fluorescent Substrates.
Biochemistry 59: 1474-1481

Steinfelder S, Andersen JF, Cannons JL, Feng CG, Joshi M, Dwyer D, Caspar P, Schwartzberg
PL, Sher A, Jankovic D (2009) The major component in schistosome eggs responsible for
conditioning dendritic cells for Th2 polarization is a T2 ribonuclease (omega-1). *J Exp Med* 206:
1681-1690

Straume AH, Kjaerner-Semb E, Ove Skaftnesmo K, Guralp H, Kleppe L, Wargelius A,
Edvardsen RB (2020) Indel locations are determined by template polarity in highly efficient in
vivo CRISPR/Cas9-mediated HDR in Atlantic salmon. *Sci Rep* 10: 409

Swarts DC (2019) Making the cut(s): how Cas12a cleaves target and non-target DNA. *Biochem*
Soc Trans 47: 1499-1510

Swarts DC, Jinek M (2018) Cas9 versus Cas12a/Cpf1: Structure-function comparisons and
implications for genome editing. *Wiley Interdiscip Rev RNA*: e1481

Swarts DC, Jinek M (2019) Mechanistic Insights into the cis- and trans-Acting DNase Activities
of Cas12a. *Mol Cell* 73: 589-600 e584

Takaki KK, Rinaldi G, Berriman M, Pagan AJ, Ramakrishnan L (2021a) Schistosoma mansoni
Eggs Modulate the Timing of Granuloma Formation to Promote Transmission. *Cell Host*
Microbe 29: 58-67 e55

Takaki KK, Roca FJ, Schramm G, Wilbers RHP, Ittiprasert W, Brindley PJ, Rinaldi G, Berriman
M, Ramakrishnan L, Pagan AJ (2021b) Tumor Necrosis Factor and Schistosoma mansoni egg
antigen omega-1 shape distinct aspects of the early egg-induced granulomatous response. *PLoS*
Negl Trop Dis 15: e0008814

Tóth E, Czene BC, Kulcsár PI, Krausz SL, Tálas A, Nyeste A, Varga É, Huszár K, Weinhardt N,
Ligeti Z *et al* (2018) Mb- and FnCpf1 nucleases are active in mammalian cells: activities and
PAM preferences of four wild-type Cpf1 nucleases and of their altered PAM specificity variants.
Nucleic Acids Res 46: 10272-10285

Wilbers RH, Westerhof LB, van Noort K, Obieglo K, Driessen NN, Everts B, Gringhuis SI,
Schramm G, Goverse A, Smant G *et al* (2017) Production and glyco-engineering of
immunomodulatory helminth glycoproteins in plants. *Sci Rep* 7: 45910

Yuan JS, Burris J, Stewart NR, Mentewab A, Stewart CN, Jr. (2007) Statistical tools for
transgene copy number estimation based on real-time PCR. *BMC Bioinformatics* 8 Suppl 7: S6

Zamanian M, Andersen EC (2016) Prospects and challenges of CRISPR/Cas genome editing for
the study and control of neglected vector-borne nematode diseases. *Febs j* 283: 3204-3221

Zetsche B, Gootenberg JS, Abudayyeh OO, Slaymaker IM, Makarova KS, Essletzbichler P, Volz
SE, Joung J, van der Oost J, Regev A *et al* (2015) Cpf1 is a single RNA-guided endonuclease of
a class 2 CRISPR-Cas system. *Cell* 163: 759-771

Zhang Y, Ren Q, Tang X, Liu S, Malzahn AA, Zhou J, Wang J, Yin D, Pan C, Yuan M *et al*
(2021) Expanding the scope of plant genome engineering with Cas12a orthologs and highly
multiplexable editing systems. *Nat Commun* 12: 1944

figure 1

C   **Cas9-ssODN:**
cttgtccactttaggaaacatgaatttgaaaagcatggtttgtgtgcag<u>TAAGTGACTAGGTAACTGAGTAGC</u>ttgaggatcctcaagtgtttaatcaatacggttactttaaatttggtatt

Cas12a- NT-ssODN:
tgtccactttaggaaacatgaatttgaaaagcatggtttgtgtgcagtt<u>TAAGTGACTAGGTAACTGAGTAGC</u>gaggatcctcaagtgtttaatcaatacggttactttaaatttggtattaa

Cas12a- T-ssODN:
attggaacaggtgaaaatcctttgtacttaaactttttcgtaccaaacacac<u>ATTCACTGATCCATTGACTCATCG</u>gtcaactcctaggagttcacaaattagttatgccaatgaaatttaaacca

A

Δω1KO-Cas9

B

Δω1KO-Cas12a

C

D