Metabolic flux and flux balance analyses indicate the relevance of metabolic thermogenesis and aerobic glycolysis in cancer cells

Authors: Nobuyuki Okahashi[1], Tomoki Shima[1], Yuya Kondo[1], Chie Araki[1], Shuma Tsuji[1], Akane Sawai[1], Hikaru Uehara[1], Susumu Kohno[2], Hiroshi Shimizu[1], Chiaki Takahashi[2], and Fumio Matsuda[1,*]

Affiliations:

[1]Department of Bioinformatic Engineering, Graduate School of Information Science and Technology, Osaka University, 1-5 Yamadaoka, Suita, Osaka 565-0871, Japan

[2]Cancer Research Institute, Kanazawa University, Kanazawa, Ishikawa 920-1192, Japan.

*Corresponding author.

Email: *fmatsuda@ist.osaka-u.ac.jp*

Abstract: Adenosine triphosphate (ATP) regeneration by substrate-level phosphorylation is a general feature of cancer metabolism, even under normoxic conditions (aerobic glycolysis). However, it is unclear why cancer cells prefer inefficient aerobic glycolysis over the highly efficient process of oxidative phosphorylation for ATP regeneration. To investigate the metabolic principles underlying aerobic glycolysis, we performed [13]C-metabolic flux analysis of 12 cultured cancer cell lines and explored the metabolic constraints required to reproduce the results using *in silico* metabolic simulations. We found that the measured flux distribution can be reproduced by maximizing the ATP consumption in the flux balance analysis considering a limitation of metabolic heat dissipation (enthalpy change). It suggests that aerobic glycolysis may be preferable because metabolic heat production during one mol of ATP regeneration by aerobic glycolysis was less than that produced by oxidative phosphorylation (OXPHOS). Consistent with the simulation, OXPHOS inhibition induced metabolic redirection to aerobic glycolysis while maintaining the intracellular temperature. Furthermore, the dependency on aerobic glycolysis was partly alleviated upon culturing at low temperatures. Our data suggest that metabolic thermogenesis is an important factor in understanding aerobic glycolysis in cancer cells and that an advantage of aerobic glycolysis is the reduction in metabolic heat generation during ATP regeneration.

Introduction

Adenosine triphosphate (ATP) regeneration is one of the most important metabolic processes and is required to maintain various cellular functions. Generally, normal human cells efficiently regenerate ATP by oxidative phosphorylation (OXPHOS) through the electron transport chain (ETC) (32 ATP/glucose, denoted as the glucose → TCA cycle). In contrast, cancer cells depend on substrate-level phosphorylation for ATP regeneration, even under conditions of oxygen availability (2 ATP/glucose, aerobic glycolysis) [1]. This altered metabolic state is ascribed to the reduced ETC activity that triggers the metabolic rewiring toward aerobic glycolysis to meet the large ATP demand for active cell proliferation in cancer cells [2]. However, ETC deficiency depends on the type of cancer cell line as certain cell types activate ETC and more than 50% of ATP regeneration is attributed to OXPHOS [3]. Moreover, cancer cells commonly catabolize

glutamine as a carbon source to regenerate ATP by OXPHOS (glutaminolysis) [4,5]. Thus, the rationale for employing aerobic glycolysis for ATP regeneration is far from being understood, although several possible roles have been proposed [6].

Results

Total ATP regeneration flux is not correlated with cell proliferation rates

In this study, we performed ^{13}C-metabolic flux analysis (^{13}C-MFA) of 12 cancer cell lines and *in silico* metabolic simulations. These cell lines were selected based on the diversity of their expression patterns for genes associated with central carbon metabolism to obtain a global view of energy metabolism [7] (**Fig. S1**). Their ability to grow in Dulbecco's modified Eagle's medium (DMEM) was considered to reduce the bias derived from the culture conditions. The specific cell proliferation rate, cell adhesion area, and diameter of trypsinized cells were measured as visible phenotypes of the cell cultures (**Table S1**).

Metabolic flux distributions allowed quantitative comparison of flux levels among pathways [8,9] (**Figs. 1A and S2, Table S2, Data S1**). Details of the procedure for ^{13}C-MFA are provided in the supplementary Text (**Text S1**). ATP was mainly regenerated by three metabolic pathways in the 12 cancer cell lines: aerobic glycolysis (red arrow in **Fig. 1A**), glutaminolysis (blue arrow), and glucose → TCA cycle (gray arrow). In the case of MCF-7 cells, the metabolic flux level in glycolysis, represented by the glyceraldehyde-3-phosphate dehydrogenase (GAPDH) reaction, was 1104 nmol (10^6 cells)$^{-1}$ h^{-1}. The flux levels of glutamate dehydrogenase (GLUDH, representing the glutaminolysis flux) and the isocitrate dehydrogenase (IDH) reaction (representing the glucose → TCA cycle flux) were less than 10% of the glycolytic flux, that were 19 and 25 nmol (10^6 cells)$^{-1}$ h^{-1}, respectively (**Fig. 1A**). All 12 cell lines demonstrated similar trends, that is, the flux through aerobic glycolysis was generally higher than that through glutaminolysis and the glucose → TCA cycle (**Fig. S3**). However, significant variations were observed in the fluxes of GAPDH, GLUDH, and IDH reactions among the cell lines (**Fig. S3**). Among these, the glucose → TCA cycle flux was not universal because the metabolic flux levels of the IDH reaction were close to zero in HepG2 and MIA PaCa-2 cells (**Fig. S3**). Principal component analysis revealed no definite correlation between the metabolic flux distributions and cell origins (**Fig. 1B**), suggesting that the variations may be due to other biological contexts in each cell line.

Net ATP regeneration rates within the central carbon metabolic network were calculated from the metabolic flux distributions (**Fig. 1C**). For instance, the net ATP regeneration rate in MCF-7 cells was 1815 nmol (10^6 cells)$^{-1}$ h^{-1}, which varied 2.8 times among the 12 cell lines. An approximate estimate indicated that the net ATP regeneration rates were higher than those of most non-cancerous cells (See Supplementary **Text S2** for details). Moreover, the ATP regeneration rate in MCF-7 cells showed that 39% of ATP was regenerated by OXPHOS through the ETC (**Fig. 1C**). OXPHOS commonly occurred in all 12 cancer cell lines, suggesting that this mechanism is still mandatory for ATP regeneration in cancer cells. Consistent with this observation, the cell lines maintained their mitochondrial membrane potential (**Fig. S4**). However, a large variation was observed in the OXPHOS-dependent ATP regeneration rate, ranging from 24% (HepG2) to 68% (LS174T) of the net ATP regeneration rate (**Fig. 1C**) [10]. This variation is likely to be derived from the distinct ETC activities of each cell line depending on the biological context. Thus, it is reasonable to assume that the ETC must work within an optimal capacity as a loss of mitochondrial membrane potential by low ETC activity triggers cell

apoptosis [11,12] and an ETC overload excessively generates reactive oxygen species, which also induces cell apoptosis [13].

90　　Given that cells maintain a rapid turnover of ATP, it can be assumed that an equal amount of regenerated ATP must be consumed for some cellular functions, such as proliferation. However, no correlation was observed between ATP regeneration rates and specific cell proliferation rates ($r = 0.06$, **Figs. 1D and S5**), suggesting the presence of other major ATP demands in cultured cancer cells. The correlation analysis indicated that the cell adhesion area on the culture plates

95　　was positively correlated with the net ATP regeneration rates ($r = 0.76$, **Figs. 1D and S5**). This trend was also observed when an outlier (Caki-1) was removed from the dataset ($r = 0.68$; **Fig. S6**). Moreover, no correlation was observed between the cell adhesion area and diameter of the trypsinized floating cells ($r = 0.07$, **Fig. 1D**). Thus, the cell adhesion area is a phenotype of cancer cells that is distinct from the cell volume. Notably, the adhesion between the cell and the

100　　extracellular matrix mediated by cadherin promotes cancer cell invasion in conjunction with metabolic changes to produce the required amount of ATP [14]. Actin and myosin utilize ATP to regulate tumor cell adhesion and migration [15,16]. Taken together, these results indicate that cancer cell metabolism is not shaped by a proliferation-oriented mechanism.

Figure 1. Variation in the metabolic flux distribution among 12 cancer cell lines and their correlation with the phenotypes.
(**A**) Metabolic flux distribution of MCF-7 as determined by [13]C-MFA. (**B**) The score plot of the principal component analysis using the metabolic flux distributions of 12 cell lines. The color

110　　legend is identical to that of the next panel. (**C**) Variations in ATP regeneration rates by substrate-level phosphorylation and OXPHOS. Error bars represent 95% confidence intervals. (**D**) Correlation between ATP regeneration rates and visible cell phenotypes. Spearman's rank order correlation coefficients are depicted in the heat map ($n=12$).

115　　***In silico* simulation indicates that aerobic glycolysis is required to reduce metabolic heat**

To elucidate the principle underlying the coordination of cancer metabolism, we explored the metabolic constraints required to reproduce the measured flux distributions using *in silico*

metabolic simulations. We performed flux balance analysis (FBA) using a genome-scale model of human metabolism (RECON2) [17] (**Data S2**). FBA requires an objective function to simulate the intracellular metabolic flux distribution using a linear programming method [18]. Although maximal biomass production is commonly used as an objective function to simulate bacterial metabolism, a suitable objective function for simulating cancer cell metabolism remains under investigation [19]. In this study, the [13]C-MFA results suggested that the cultured cancer cells consumed a large amount of ATP to maintain certain cellular functions other than proliferation (**Fig. 1D**). Thus, the maximal ATP consumption was employed as the objective function in the *in silico* analysis.

The FBA also requires constraints on the solution space (upper and lower boundaries of the metabolic flux vector) [20]. Here, we performed FBA to determine additional constraints that drive cell metabolism toward aerobic glycolysis and glutaminolysis for ATP regeneration, as revealed in the [13]C-MFA (**Fig. 1**). The flux distribution of MCF-7 cells was chosen as an example for reproduction by FBA (**Fig. 2A**). The relatively minor metabolic fluxes, such as those related to biomass synthesis as well as the uptake and production of amino acids other than glutamine, were constrained at the measured values of MCF-7 (all constraints for the influx and efflux are presented in **Table S3**). In contrast, no specific constraints were imposed on the uptake rates of glucose, glutamine, and oxygen and the excretion rates of lactate, carbon dioxide, and water.

The FBA results without additional constraints showed that considerable amounts of glucose and glutamine were consumed until the ATP regeneration rate reached the upper limit (**Fig. 2B, Data S3**). The predicted metabolic flux distribution mainly exhibited OXPHOS-dependent metabolism through the glucose → TCA cycle and a low correlation coefficient with that observed in MCF-7 cells ($r = 0.62$). This indicated that additional constraints are required to reproduce the measured value for MCF-7 cells.

In this study, we considered various thermodynamic parameters that govern cell metabolism [21]. For example, the entire stoichiometry of aerobic glycolysis (glucose → 2 lactate) indicates that the enthalpy change ($\Delta_r H'^\circ$), or metabolic heat generation, per 1 mole of ATP regeneration by aerobic glycolysis is -55.0 kJ (mol ATP)$^{-1}$ (**Table S4**) [22]. Similarly, the enthalpy change in the glucose → TCA cycle reaction was -88.2 kJ (mol ATP)$^{-1}$ (**Table S4**). Thus, it is expected that if the enthalpy change is limited, aerobic glycolysis would be preferentially employed for ATP regeneration in cells. Thermogenesis has long been observed in patients with breast cancer [23]. The preference for aerobic glycolysis was not observed for other parameters, such as entropy and Gibbs free energy change (**Table S4**). Thus, the genome-scale metabolic model was modified to determine the levels of cellular dissipation of enthalpy change from the FBA results (**Table S5 and Data S3**) [22].

The FBA was performed by fixing the $\Delta_r H'^\circ$ level at an arbitrary value as an additional constraint. We found that when the $\Delta_r H'^\circ$ level was set at -105 mJ (10^6 cells)$^{-1}$ h^{-1}, the predicted metabolic flux distribution was more similar to the measured MCF-7 data (**Fig. 2C**, $r = 0.89$) because aerobic glycolysis was activated for ATP regeneration, as expected.

In addition to metabolic heat, the ETC activity was applied as an additional constraint because the level of ETC flux in the above simulation was considerably low (**Fig. 2C**), which was inconsistent with the measured data (**Figs. 2A and S4**). Thus, FBA was performed by fixing the metabolic flux level of complex IV of the ETC at the measured value of MCF-7 cells (282 nmol (10^6 cells)$^{-1}$ h^{-1}). The predicted metabolic flux distribution successfully reproduced ATP regeneration by both aerobic glycolysis and glutaminolysis, and its correlation with the measured

flux distributions was $r = 0.98$ (**Fig. 2D, Fig. S7, Data S3**). However, ATP regeneration by the glucose → TCA cycle did not occur in the predicted distribution (**Fig. 2D**), which was inconsistent with the [13]C-MFA result (**Fig. 2A**). This is because the metabolic heat dissipation per 1 mole of ATP regeneration by glutaminolysis is lower than that of the glucose → TCA cycle reaction, although the detailed $-\Delta_r H'^\circ$ level cannot be calculated by simple material balance equations. Notably, the glucose → TCA cycle pathway is not universal, as mentioned above (**Fig. S3**). The FBA showed that similar ATP regeneration occurred by both aerobic glycolysis and glutaminolysis in all 12 cell lines (**Fig. S7 and Data S3**). The simulation of cancer cell metabolism suggested that the ETC activity and metabolic heat generation are possible constraints of cancer cell metabolism.

	Measured (MCF-7)	Simulated case 1	Simulated case 2	Simulated case 3
Objective function		ATP consumption max	ATP consumption max	ATP consumption max
Metabolic heat dissipation		Predicted	Fixed at -105	Fixed at -105
ETC level (complex IV)		Predicted	Predicted	Fixed at 282
Correlation to MCF-7 data		$r = 0.62$	$r = 0.89$	$r = 0.98$

Figure 2. Metabolic heat dissipation and ETC activity govern cancer cell metabolism.
In silico simulation of metabolism was performed using the FBA with a genome-scale model of human metabolism (RECON2). Metabolic flux levels of the key reactions of aerobic glycolysis (red), glucose → TCA cycle (gray), and glutaminolysis (blue) are displayed in the boxes. The maximum ATP consumption was employed as an objective function. Additional constraints are represented by yellow and green numbers. (**A**) Measured metabolic flux distribution in MCF-7 cells. (**B-D**) Results of FBA (**B**) without additional constraints, (**C**) with fixed metabolic heat dissipation, and (**D**) with fixed metabolic flux dissipation and ETC flux.

ETC inhibitor treatment induces metabolic rewiring toward aerobic glycolysis while maintaining the intracellular temperature

The metabolic simulation indicates that the downregulation of ETC activity will be compensated by the elevation of aerobic glycolysis while maintaining the heat dissipation level, which corresponds to a metabolic shift from the level in **Fig. 2D** to that in **Fig. 2C**. To test this hypothesis, the MCF-7 cells were treated with various ETC inhibitors. Rotenone treatment (100 nM) partially inhibited cell proliferation by 25% after 24 h (data not shown). The time-course data showed that cell proliferation was temporarily halted by 6 h and then recovered between 6

and 24 h (**Fig. 3A**). The recovery in growth was accompanied by the activation of aerobic glycolysis as the specific rates for glucose consumption and lactate production between 12 and 24 h increased by 1.2-fold with rotenone treatment (**Fig. 3B**).

To investigate the switching point of the metabolic pathway, ^{13}C-tracer and metabolome analysis was performed. Tracer analysis using [U-^{13}C]glutamine revealed that the relative abundance of the m+4 isotopomer of citrate, which is produced from acetyl-CoA and oxaloacetate, was reduced after rotenone treatment, whereas that of m+5 citrate produced by reductive glutamine metabolism increased (**Fig. 3C**). The labeling pattern indicated a reduction in the flux of the glucose → TCA cycle in the rotenone-treated MCF-7 cells. The metabolic shift is reasonable because MCF-7 cells used the glucose → TCA pathway (**Fig. 1A**) although the use of this pathway could not be reproduced by metabolic simulation (**Fig. 2D**). Metabolome analysis showed that rotenone treatment induced an accumulation of malate and lactate, while also decreasing the citrate and isocitrate levels (**Fig. 3D**). A similar metabolic profile was commonly induced by other ETC inhibitors, antimycin and oligomycin, but not by the mitochondrial uncoupler, carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone (FCCP) (**Fig. 3D**). These results suggest that the downregulation of the entry point of the TCA cycle also contributes to a metabolic shift from the glucose → TCA cycle to aerobic glycolysis.

To evaluate the effect of changes in heat generation by metabolic redirection, the intracellular temperature of cells treated with ETC inhibitors was measured using a celluler thermoprobe. The temperature increase caused by the FCCP-induced decoupling was accurately captured (**Fig. 3E**). In contrast, treatment with rotenone and other ETC inhibitors, including antimycin and oligomycin, did not significantly change the intracellular temperature, demonstrating that aerobic glycolysis was elevated while maintaining the heat dissipation level (**Fig. 3E**). The metabolic simulation results indicate that a reduction in ATP regeneration by the restriction of ETC activity was circumvented by the activation of aerobic glycolysis within the heat dissipation capacity, supporting the validity of the metabolic model considering enthalpy change.

Figure 3. ETC inhibitor treatment induces metabolic rewiring toward aerobic glycolysis while maintaining the intracellular temperature.
(**A**) Proliferation of MCF-7 cells after 100 nM rotenone treatment. (**B**) Specific rates for glucose and glutamine uptake and lactate excretion between 12 and 24 h of culturing. Asterisks indicate the statistical significance relative to the control, as determined by a two-sided Student's *t*-test ($p < 0.05$). (**C**) Mass isotopomer distribution of citrate in MCF-7 cells in [U-^{13}C]glutamine tracing. (**D**) Metabolites of MCF-7 cells treated with OXPHOS inhibitors. Relative abundances of metabolites (Log$_2$(fold change)) are shown in the heatmap. All data are represented as mean ± SD (*n*=3). (**E**) Intracellular temperature of MCF-7 treated by ETC-inhibitors for 24 h. Intracellular temperatures of MCF-7 cells treated with DMSO, 100 nM rotenone, 100 nM antimycin, 100 nM oligomycin, and 5 mM FCCP for 24 h were measured using a cellular thermoprobe. The color scale represents the fluorescence intensity ratio (570-625/510-550 nm), which corresponds to the intracellular temperature.

Metabolic rewiring toward aerobic glycolysis is dependent on temperature

The above observations suggest that cancer cells rewire their metabolism to control the heat dissipation level. The glucose → TCA cycle pathway is preferable under lower and higher temperature conditions to maintain a constant intracellular temperature. To test this assumption, cells were cultured at control (37 °C), low (36 °C), and high (38 °C) temperatures. A comparison between the glucose and lactate consumption for 18 h showed that there was a limited effect on the glucose consumption (**Fig. 4A**), whereas lactate production tended to decrease and increase at 36 and 38 °C, respectively (**Fig. 4B**). A significant increase in lactate production was observed in both cell lines at 38 °C (*t*-test, *n*=3, α=0.05). In contrast, a significant decrease in lactate production was observed in five cell lines at 36 °C. The results indicated that the metabolic shift toward the glucose → TCA cycle pathway was temperature-sensitive.

Gene expression data were obtained from MCF-7 and HCT116 cells, which indicated a temperature-dependent increase in lactate production, as representative cell lines (*n*=1, **Table S6, Data S4**). We selected a total of 3154 and 4533 open reading frames (ORFs), whose expression levels were altered in MCF-7 and HCT116 cells in the following order: [36 °C]>[37 °C]>[38 °C]. A total of 683 ORFs were common in the MCF-7 and HCT116 datasets (**Fig. 4C, Data S5**), which was significantly larger than the expected number of overlapping ORFs (545 ORFs, χ^2 test, *p*-value =2.0×10^{-9}). An over-representation analysis was performed for the 683 ORFs using the Kyoto Encyclopedia of Genes and Genomes (KEGG) 2019 human dataset [24]. The results revealed that ORFs in the "TCA cycle" category (*PDHA1, MDH2, IDH3G, IDH2, IDH3B, OGDHL, DLAT,* and *SDHD*) were enriched in 683 ORFs (**Fig. 4D, Table S7**). Notably, *PDHA1* and *DLAT* encode subunits of the pyruvate dehydrogenase complex, which is located at the entry point of the TCA cycle; this was also indicated in the aforementioned metabolome analysis (**Fig. 3D**). No functional category was found in the ORF list that exhibited a reversed expression pattern in the following order: [36 °C]<[37 °C]<[38 °C] (data not shown). These results suggest that metabolic rewiring from the glucose → TCA cycle to aerobic glycolysis is dependent on temperature-induced changes in the metabolic thermogenesis.

Figure 4. Culture temperature induces metabolic rewiring by altering gene expression.
Cancer cells were cultured under control (37 °C), low (36 °C), and high (38 °C) temperatures.
Concentrations of **(A)** glucose and **(B)** lactate in the culture medium at 18 h. Data are represented
as mean ± SD (n=3). Asterisks indicate the statistical significance of concentrations under test
conditions as determined by one way analysis of variance (ANOVA) using the Tukey-Kramer
method (*$p < 0.05$). **(C)** Venn diagram of temperature-responsive genes. RNA-Seq of MCF-7
and HCT116 cells at 24 h identified 3154 and 4533 genes expressed with fold changes in the
following order: [36 °C]>[37 °C]>[38 °C]. **(D)** An over-represented analysis of gene categories
of 683 common genes by Enricher using the KEGG 2019 Human dataset [24].

Discussion

This study reaffirms that thermogenesis is an important aspect of cellular metabolism,
particularly when cancer cells require large amounts of ATP. The effects of thermogenesis on
cancer cell metabolism are shown in **Figure 5**. In non-cancerous cells, ATP is supplied by the
glucose → TCA cycle pathway through OXPHOS, which efficiently regenerates ATP (**Fig. 5A**).
Metabolic heat dissipation is maintained within the heat homeostasis. The ATP demand is
elevated during cancer development as ATP is required for cancer-specific features, such as
rapid cell proliferation, tumor cell adhesion, and migration. The ^{13}C-MFA results suggested that
a larger amount of ATP was utilized for cell adhesion rather than cell proliferation (**Figs. 1C and
D**).

To meet the elevated ATP demand, the active use of the OXPHOS pathway is infeasible because
of the production of excessive metabolic heat (**Fig. 5B**). Notably, intracellular heat homeostasis
is maintained by a balance between the generation and dissipation of a large amount of heat. ^{13}C-
MFA and metabolic simulation results indicated that the amount of heat dissipated from MCF-7
cells was estimated to be on the order of 100 mJ $(10^6$ cells$)^{-1}$ h^{-1} (**Fig. 2D**). This level is
substantially higher than that required to increase the temperature of cultured cells by 1 °C [7.4
micro J $(10^6$ cells$)^{-1}$], assuming that the cells contain 1.76 pico L of water.

In contrast to the activation of the OXPHOS pathway, the additional use of aerobic glycolysis
can compensate for the lack of ATP, which limits cellular thermogenesis in the glucose → TCA

cycle (**Fig. 5C**). This is the only feasible scenario for ATP production in cancer cells as all 12 cancer cell lines investigated in this study employed aerobic glycolysis in combination with OXPHOS. The relationship between heat and metabolism was elucidated by *in silico* simulation (**Fig. 2**) and validated by culture experiments at low and high temperatures (**Fig 4**). The inhibition of ETC activity was compensated by metabolic rewiring for increased aerobic glycolysis while maintaining the heat dissipation level (**Fig. 3**).

Thus, the cell metabolism operates within an appropriate range of metabolic heat generation. This indicates that external heat radiation regulates respiration, as demonstrated by hyperthermia [23]. These results provide new insights into the relevance of cancer metabolism and malignancy, cancer microenvironment interactions, and drug resistance [6]. Heat generation may also be involved in activating immune cells and controlling the cancer microenvironment [25,26]. Furthermore, the relationship between metabolism and heat is universal. Notably, aerobic glycolysis has been observed in normal cells, such as astrocytes, embryos, and stem cells [27–29]. These tissues and cells commonly require large amounts of ATP for biochemical processes; however, an embryo without blood vessels has a limited capacity to release heat. The role of aerobic glycolysis in cancer and healthy cells while maintaining heat homeostasis should be analyzed.

Figure 5. Cancer cell metabolism provides a large supply of ATP with less heat generation. (**A**) Normal cells can be assumed to regenerate ATP through the glucose → TCA cycle pathway and maintain metabolic heat dissipation and ETC activity within adequate levels. (**B**) Cancer cells have a high demand for ATP but the additional activation of the glucose → TCA cycle

pathway is limited as the ETC activity and metabolic heat dissipation would otherwise exceed the permissible range. **(C)** To avoid this scenario, cancer cells rewire their metabolism to favor aerobic glycolysis and glutaminolysis to increase their ATP supply and fulfill ETC activities, respectively, with less metabolic thermogenesis.

Materials and Methods

Cell lines and culture conditions

Twelve human cancer cell lines derived from colorectal cancer (HCT116, DLD-1, LS174T, and WiDr), pancreatic cancer (MIA Paca-2 and PANC-1), hepatoblastoma (HepG2), hepatocellular carcinoma (huH-1), breast cancer (MCF-7), renal cancer (Caki-1), glioblastoma (YKG-1), and sarcoma (HT1080) were obtained from the RIKEN Bioresource Research Center and Japanese Collection of Research Bioresources (JCRB) Cell Bank. Culture experiments were conducted as described previously [30]. In particular, 1.0×10^6 cells were seeded in 10 mL of Dulbecco's modified Eagle's medium (DMEM) supplemented with 10% fetal bovine serum (FBS) and 1% penicillin/streptomycin (Wako) in 100-mm (diameter) plates and cultured for 15 h at 37 °C under 5% CO_2. ^{13}C-labeling was performed in 10 mL of DMEM containing 20 mM [1-^{13}C] of glucose (Cambridge Isotope Laboratories, Andover, MA, USA, over 99% purity) and 10% dialyzed FBS (Life Technologies, Gaithersburg, MD, USA). The cells were counted noninvasively at 0, 12, 16, 20, and 24 h in triplicate cultures using a CKX53 inverted microscope (Olympus, Tokyo, Japan) equipped with a DP22 digital camera (Olympus) and CKX-CCSW software (Olympus). Cell images were obtained at 10 different locations in the culture dish, and the average cell numbers were used to generate the growth curve. Culture media (200 μL) were sampled at each time point. Trypsin was added to the plates and activated at 37 °C for 1 min to measure the diameter of the floating cells. After cell collection, the diameter of live cells was counted using trypan blue staining and a TC20 automated cell counter (Bio-Rad, Hercules, CA, USA). Cell sizes were analyzed using the ImageJ software (National Institutes of Health, Bethesda, MD, USA).

Extracellular metabolite measurements

The collected culture medium was mixed with an equal volume of 20 mM pimelate solution (internal standard) and filtered through a filter cartridge (pore size of 0.45 μm). Glucose, lactate, and acetate concentrations were determined by high-performance liquid chromatography (HPLC) equipped with a refractive index detector (Prominence, Shimadzu, Kyoto, Japan) and an Aminex HPX-87H column (Bio-Rad), as previously described [30]. Amino acid concentrations in the culture media were measured by HPLC using the AccQ•Tag method (Armenta et al., 2010). HPLC Prominence (Shimadzu) system equipped with a Luna C18 (2) column (250 mm, 4.6 mm, and 5.0 μm, SHIMADZU GLC, Kyoto, Japan) and a photodiode array detector (260 nm) was used. Derivatized amino acids were eluted with a 20 mM sodium acetate solution containing 0.04% (v/v) of trimethylamine and phosphate adjusted to a pH of 6.8 (A) and acetonitrile (B) under the following gradient conditions: 0 min, 0% (B); 0.5 min, 8% (B), 17.5 min, 12% (B), 19 min, 15% (B), 20 min, 20% (B), 30.6 min, 100% (B), and 33.1 min, 0% (B) at a flow rate of 1.0 mL/min. The column oven temperature was maintained at 40 °C.

Extraction and derivatization of intracellular metabolites

Intracellular metabolites were extracted using the methanol/water/chloroform method [30]. Cellular metabolites were rapidly quenched by adding 800 µL of precooled methanol after rapid medium removal and rinsing with phosphate-buffered saline (1 mL). This procedure was performed within 15 s. Cells and solutions were collected by scraping. Cell lysates were transferred into fresh sample tubes, followed by the addition of 800 µL of cold chloroform and 320 µL of cold water. After vortexing and centrifugation, 250 µL of the upper aqueous layer was collected and dried in a vacuum concentrating centrifuge (CVE-3110; Eyela, Tokyo, Japan) at room temperature. The dried metabolites were methoxyaminated and *tert*-butyldimethylsilyated for gas chromatography-mass spectrometry (GC-MS) analysis, as described previously [31].

Gas chromatography/mass spectrometry analysis

Gas chromatography/mass spectrometry (GC/MS) analysis was performed using a GCMS-QP2020 instrument (Shimadzu) equipped with a DB-5MS capillary column (Agilent Technologies). GC/MS was operated under electron impact (EI) ionization at 70 eV. One microliter of the sample was injected at 250 °C using helium as the carrier gas at a flow rate of 1 ml/min. To analyze central metabolite derivatives, the GC oven temperature was maintained at 60 °C, and then increased to 325 °C at 10 °C /min for a total run time of approximately 30 min. The MS source and quadrupole were maintained at 230 and 150 °C, respectively. The effects of naturally occurring isotopes were corrected [32].

^{13}C-metabolic flux and mass balance analyses

The specific uptake and secretion rates of extracellular metabolites were calculated as previously described [33]. The metabolic flux distribution was estimated by minimizing the variance-weighted residual sum of squares of measured and estimated mass isotopomer distributions of intracellular metabolites and effluxes using mfapy [34]. The net ATP regeneration flux was calculated using flux distributions and the following assumptions: the reactions catalyzed by hexokinase, phosphofructokinase, pyruvate carboxylase, acetyl-coenzyme A citrate lyase/phosphoglycerate kinase, pyruvate kinase, and succinyl-coenzyme A lyase were assumed to be ATP consuming/regenerating reactions, respectively. NADPH is also assumed to be regenerated in reactions involving glucose 6-phosphate dehydrogenase, 6-phosphogluconate dehydrogenase, isocitrate dehydrogenase, and malate NADP$^+$ oxidoreductase, and NADPH is consumed in the fatty acid and proline biosynthesis pathways. We assumed that excess NADPH was converted to NADH through a transhydrogenase reaction. NADH is regenerated in reactions involving glyceraldehyde-3 phosphate dehydrogenase, pyruvate dehydrogenase, alpha-ketoglutarate dehydrogenase, malate dehydrogenase, and transhydrogenase reactions and is consumed by lactate dehydrogenase. We assumed that excess NADH was converted to ATP through OXPHOS. The P/O ratio is assumed to be 2.5. Although FADH$_2$ was also converted to ATP during OXPHOS, the P/O ratio was assumed to be 1.5. The ATP required for cell biomass synthesis was assumed to be 35 mmol/g of the dry cell weight [17].

Clustering analysis

The gene expression dataset (E-MTAB-2706) of cancer cell lines was obtained from ArrayExpress (https://www.ebi.ac.uk/arrayexpress/) [7]. Genes related to central metabolism were obtained from the KEGG pathway database (glycolysis/gluconeogenesis, pentose phosphate pathway, citrate cycle, D-glutamine and D-glutamate metabolism, and related transporters). The 62 genes were selected because the dataset did not contain missing data. The transcript per million (TPM)-normalized, log$_2$-transformed, and Z-scored gene expression datasets of 622

cancer cell lines were used for hierarchical clustering using the Seaborn clustermap of Python 3.8 (Ward method combined with Euclidean distance).

Flux balance analysis

410   A human genome-scale model (RECON2) [17] was used with the following modifications: i) 41 reactions responsible for the degradation of essential amino acids were removed from the model; ii) terms of standard enthalpy of formation (MegaJ mol^{-1}) were added to intra/extracellular transport reactions (**Tables S7 and S8**); and iii) the reaction (R_ent) was added to the sum of the standard enthalpy of formation. The metabolic heat dissipation (h_{out}) was
415 calculated using the following equation:

$$h_{out} = -\left(\sum_i \Delta_f H'^0_i J_i - \sum_j \Delta_f H'^0_j J_j \right)$$

Here, Ji and Jj indicate the uptake and excretion flux levels of the ith and jth metabolite, respectively. Moreover, $\Delta_f H'^0$ is the standard enthalpy of formation of each metabolite; $\Delta_f H'^0$ values at a pH of 7.0 and ion strength of 0.1 M were obtained from published literature [22]. The
420 $\Delta_f H'^0$ level of biomass was approximated based on the data from yeast [35]. The approximation did not affect the FBA in this study because the metabolic flux levels for biomass synthesis were fixed at the measured values. FBA was performed using an in-house Python script with the GNU Linear Programming Kit (GLPK) in the pulp module as a linear programming solver.

425 RNA-sequencing

     The total RNA was extracted using the QIAzol lysis reagent (QIAGEN). Library preparation was performed using the TruSeq stranded mRNA sample prep kit (Illumina, San Diego, CA, USA) according to the manufacturer's instructions. Sequencing was performed on an Illumina HiSeq 3000 platform in the 101 bp single-end mode. Illumina Casava1.8.2 software was
430 used for base calling. Sequenced reads were mapped to the human reference genome sequences (hg19) using TopHat v2.0.13 in combination with Bowtie2 ver. 2.2.3 and SAMtools ver. 0.1.19. Fragments per kilobase of exon per million mapped fragments (FPKMs) were calculated using Cufflinks version 2.2.1. Gene enrichment analysis was conducted using the Enrichr web tool [24].

435 Fluorescence microscopy

     Cells were seeded on glass bottom plates (35 mm diameter, Matsunami Glass, Osaka, Japan) and stained according to the manufacturer's protocol. The mitochondria were stained using MitoBright LT Red (Dojindo, Kumamoto, Japan). The mitochondrial membrane potential was detected using the JC-1 MitoMP Detection Kit (Dojindo). ECLIPSE TE2000-E inverted
440 microscope (Nikon, Tokyo, Japan) equipped with an oil immersion objective lens (Plan Apo 60×/1.4 Oil Ph3 DM, Nicon), filter sets (excitation filter, 465-496 nm; emission filter, 515-555; dichroic mirror, 505, Nicon), and an iXon EMCCD camera (Andor Technology Ltd, Belfast, UK) were used. The intracellular temperature was measured using a cellular thermoprobe to determine the fluorescence ratio (Funakoshi, Tokyo, Japan). IX71 inverted microscope
445 (Olympus, Tokyo, Japan) equipped with an objective lens (LCPlanFl 20×/0.40 Ph1, Olympus) and ORCA-Spark digital CMOS camera (Hamamatsu) were used. Fluorescence images were taken using U-MWIBA3 (excitation filter, 460-495 nm; emission filter, 510-550 nm; dichroic mirror, 505 nm, Olympus) and custom-made (excitation filter, 460-495 nm; emission filter, 570-625 nm; dichroic mirror, 505 nm, Olympus) filter cubes. The temperature and CO_2 content of

450   cells were maintained at 37 °C and 5%, respectively, using a microscope incubator (BLAST, Kawasaki, Japan). Images were analyzed using the ImageJ software.

References

455   1.   De Berardinis, R. J. & Chandel, N. S. Fundamentals of cancer metabolism. *Sci. Adv.* **2**, (2016).

2.   Koppenol, W. H., Bounds, P. L. & Dang, C. V. Otto Warburg's contributions to current concepts of cancer metabolism. *Nat. Rev. Cancer* **11**, 325–337 (2011).

3.   Zu, X. L. & Guppy, M. Cancer metabolism: Facts, fantasy, and fiction. *Biochem. Biophys. Res. Commun.* **313**, 459–465 (2004).

460   4.   Fan, J. *et al.* Glutamine-driven oxidative phosphorylation is a major ATP source in transformed mammalian cells in both normoxia and hypoxia. *Mol. Syst. Biol.* **9**, 712 (2013).

5.   Yang, L., Venneti, S. & Nagrath, D. Glutaminolysis: A Hallmark of Cancer Metabolism. *Annu. Rev. Biomed. Eng.* **19**, 163–194 (2017).

465   6.   Liberti, M. V. & Locasale, J. W. The Warburg Effect: How Does it Benefit Cancer Cells? *Trends Biochem. Sci.* **41**, 211–218 (2016).

7.   Klijn, C. *et al.* A comprehensive transcriptional portrait of human cancer cell lines. *Nat. Biotechnol.* **33**, 306–312 (2015).

8.   Duckwall, C. S., Murphy, T. A. & Young, J. D. Mapping cancer cell metabolism with(13)C flux analysis: Recent progress and future challenges. *J. Carcinog.* **12**, 13 (2013).

9.   Antoniewicz, M. R. Methods and advances in metabolic flux analysis: a mini-review. *J. Ind. Microbiol. Biotechnol.* **42**, 317–325 (2015).

10.   Zhang, J., Pavlova, N. N. & Thompson, C. B. Cancer cell metabolism: the essential role of the nonessential amino acid, glutamine. *EMBO J.* **36**, 1302–1315 (2017).

11.   Ly, J. D., Grubb, D. R. & Lawen, A. The mitochondrial membrane potential ($\delta\psi$m) in apoptosis; an update. *Apoptosis* **8**, 115–128 (2003).

12.   Li, N. *et al.* Mitochondrial complex I inhibitor rotenone induces apoptosis through enhancing mitochondrial reactive oxygen species production. *J. Biol. Chem.* **278**, 8516–8525 (2003).

13.   Zhao, R. Z., Jiang, S., Zhang, L. & Yu, Z. Bin. Mitochondrial electron transport chain, ROS generation and uncoupling (Review). *International Journal of Molecular Medicine* **44**, 3–15 (2019).

14.   Sousa, B., Pereira, J. & Paredes, J. The crosstalk between cell adhesion and cancer metabolism. *International Journal of Molecular Sciences* **20**, (2019).

15.   Ouderkirk, J. L. & Krendel, M. Non-muscle myosins in tumor progression, cancer cell invasion, and metastasis. *Cytoskeleton* **71**, 447–463 (2014).

16.   Zanotelli, M. R. *et al.* Regulation of ATP utilization during metastatic cell migration by collagen architecture. *Mol. Biol. Cell* **29**, 1–9 (2018).

490   17.   Thiele, I. *et al.* A community-driven global reconstruction of human metabolism. *Nat. Biotechnol.* **31**, 419–425 (2013).

        18.   Feist, A. M. & Palsson, B. O. The biomass objective function. *Curr. Opin. Microbiol.* **13**, 344–349 (2010).

        19.   Zielinski, D. C. *et al.* Systems biology analysis of drivers underlying hallmarks of cancer
495   cell metabolism. *Sci. Reports 2017 71* **7**, 1–14 (2017).

        20.   Orth, J. D., Thiele, I. & Palsson, B. Ø. What is flux balance analysis? *Nat. Biotechnol. 2010 283* **28**, 245–248 (2010).

        21.   Niebel, B., Leupold, S. & Heinemann, M. An upper limit on Gibbs energy dissipation governs cellular metabolism. *Nat. Metab.* **1**, 125–132 (2019).

500   22.   Alberty, R. A. Calculation of standard transformed Gibbs energies and standard transformed enthalpies of biochemical reactants. *Arch. Biochem. Biophys.* **353**, 116–130 (1998).

        23.   Cavaliere, R. *et al.* Selective heat sensitivity of cancer cells. Biochemical and clinical studies. *Cancer* **20**, 1351–1381 (1967).

505   24.   Xie, Z. *et al.* Gene Set Knowledge Discovery with Enrichr. *Curr Protoc* **1**, e90 (2021).

        25.   Kokolus, K. M. *et al.* Baseline tumor growth and immune control in laboratory mice are significantly influenced by subthermoneutral housing temperature. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 20176–20181 (2013).

        26.   Repasky, E. A., Evans, S. S. & Dewhirst, M. W. Temperature matters! And why it should
510   matter to tumor immunologists. *Cancer immunology research* **1**, 210–216 (2013).

        27.   Abe, T., Takahashi, S. & Suzuki, N. Oxidative metabolism in cultured rat astroglia: Effects of reducing the glucose concentration in the culture medium and of D-aspartate or potassium stimulation. *J. Cereb. Blood Flow Metab.* **26**, 153–160 (2006).

        28.   Miyazawa, H. *et al.* Rewiring of embryonic glucose metabolism via suppression of PFK-1
515   and aldolase during mouse chorioallantoic branching. *Dev.* **144**, 63–73 (2017).

        29.   Ito, K. & Suda, T. Metabolic requirements for the maintenance of self-renewing stem cells. *Nature Reviews Molecular Cell Biology* **15**, 243–256 (2014).

        30.   Okahashi, N. *et al.* Metabolic characterization of cultured mammalian cells by mass balance analysis, tracer labeling experiments and computer-aided simulations. *J. Biosci.*
520   *Bioeng.* **120**, 725–731 (2015).

        31.   Araki, C., Okahashi, N., Maeda, K., Shimizu, H. & Matsuda, F. Mass spectrometry-based method to study inhibitor-induced metabolic redirection in the central metabolism of cancer cells. *Mass Spectrom.* **7**, 1–9 (2018).

        32.   Okahashi, N., Yamada, Y., Iida, J. & Matsuda, F. Isotope Calculation Gadgets : A Series
525   of Software for Isotope-Tracing Experiments in Garuda Platform. *Metabolites* **12**, (2022).

        33.   Matsuda, F., Toya, Y. & Shimizu, H. Learning from quantitative data to understand central carbon metabolism. *Biotechnol. Adv.* **35**, 971–980 (2017).

        34.   Matsuda, F. *et al.* mfapy : An open-source Python package for 13C-based metabolic flux analysis. *Metab. Eng. Commun.* **13**, e00177 (2021).

530   35.   Popovic, M. Thermodynamic properties of microorganisms: determination and analysis of enthalpy, entropy, and Gibbs free energy of biomass, cells and colonies of 32 microorganism species. *Heliyon* **5**, e01950 (2019).

Acknowledgments: We acknowledge the NGS core facility of the Genome Information
535   Research Center at the Research Institute for Microbial Diseases, Osaka University for their support in RNA sequencing and data analysis. We thank Prof. Shinya Kuroda, Dr. Satoshi Ohno (University of Tokyo), and Shimpei Kawaoka (Kyoto University) for helpful comments on this manuscript. We also thank Keiko Fukamoto for her skillful technical support. Grant-in-Aid for Scientific Research on Innovative Areas (Grant no. 17H06303). Extramural Collaborative
540   Research Grant of Cancer Research Institute, Kanazawa University (Grant no.04-13).

Author contributions:

N.O., S.K., H.S., C.T. and F.M. designed the study. N.O., T.S., Y.K., C.A., and H.U. performed ^{13}C-tracer experiments. F.M. performed flux balance analysis. N.O., S.T. and A.S. performed
545   metabolome analysis. N.O. performed microscopic analysis and RNA-sequencing. N.O., T.S., Y.K., C.A., S.T., A.S., H.U., S.K., H.S., C.T. and F.M. analyzed and interpreted data. N.O. and F.M. wrote original draft. N.O., H.S., S.K., C.T. and F.M. reviewed and edited manuscript with help from all authors.

550   **Competing interests:** Authors declare that they have no competing interests.

Data availability: RNA-Seq data is available at DDBJ of the National Institute of Genetics (https:// www.ddbj.nig.ac.jp/index-e.html) via the index of DRA012975.

Supplementary Materials

Metabolic flux and flux balance analyses indicate the relevance of metabolic thermogenesis and aerobic glycolysis in cancer cells

Nobuyuki Okahashi[1], Tomoki Shima[1], Yuya Kondo[1], Chie Araki[1], Shuma Tsuji[1], Akane Sawai[1], Hikaru Uehara[1], Susumu Kohno[2], Hiroshi Shimizu[1], Chiaki Takahashi[2], and Fumio Matsuda[1,*]

Correspondence to: *fmatsuda@ist.osaka-u.ac.jp*

Supplementary Text

570 **Text S1. Detailed explanation of the ^{13}C-metabolic flux analysis procedure.**
The metabolic model consisted of the major pathways of central carbon metabolism, including glycolysis, the pentose phosphate pathway, the TCA cycle, anaplerotic reactions, and biomass synthesis (**Table S8**). Folate metabolism was ignored, as MTHFD1 and MTHFD2 were not highly expressed (**Fig. S1**). The catabolism of the essential amino acids was ignored.
575 Biomass effluxes of amino acids for proteins, glucose-6-phosphate for glycogen, ribose-5-phosphate for ribonucleic acid, and cytosolic acetyl-CoA for fatty acids were calculated from the specific growth rates and biomass compositions of mammalian cells [36]. The constant for the conversion from cell number to dry cell weight was assumed to be 0.392 mg-dry cell weight (10^6 cells)$^{-1}$ for all cell lines [36]. This assumption was based on the observation that the detached cell
580 sizes from culture dishes were almost identical to all cell lines except for Caki-1 (**Table S1**). Additionally, the effect of this assumption on the total metabolic flux was considered to be minor because the effluxes for biomass synthesis accounted for only a minor percentage of carbon uptake flux. The goodness of fit of ^{13}C-MFA was evaluated by χ^2 statistics test, where the standard deviations of ^{13}C-labeling were assumed to be 0.01 [31]. The ^{13}C-MFA of all cultured
585 cancer cell lines successfully passed the χ^2-test (**Table S2**). All 95% confidence intervals were determined using the grid search method and are shown in **Table S2** [37].

Text S2. Comparison of the net ATP regeneration rate in cultured cancer cells with that of normal cells.
590 In this study, the net ATP regeneration rate of MCF-7 cells was determined as 1815 nmol (10^6 cells)$^{-1}$ h^{-1} based on the metabolic flux distribution (**Fig. 1a**). The net ATP regeneration rates ranged from 1460 (LS174T) to 4086 (Caki-1) nmol (10^6 cells)$^{-1}$ h^{-1} among the 12 cell lines. The total human cell number is 3.72×10^{13}, composed of 15% non-erythrocyte cells [38]. Moreover, ATP was presumed to be regenerated by OXPHOS using glucose (32 ATP/glucose). If all non-
595 erythrocyte cells reproduce ATP at the same level of LS174T, the required amount of glucose (g day^{-1}) is 1460×10^{-9} nmol (10^{-6} cells)$^{-1}$ h$^{-1} \times 24$ h$\times 3.72 \times 10^{13}$ cells$\times 0.15 \times 1/32$ (glucose/ATP) $\times 180$ g mol^{-1} = 1100 g day^{-1}, equal to 3686 kcal day^{-1}. The recommended daily calorie intake is 2,000 and 2,500 kcal day^{-1} for women and men, respectively, suggesting that the ATP regeneration level of LS174T cells was higher than that of normal cells.
600 Moreover, the metabolic flux analysis conducted in this study revealed that the ATP regeneration rates by OXPHOS in 12 cancer cells were 447–1727 nmol (10^6 cells)$^{-1}$ h^{-1}. If the

glucose equivalent of 2,500 kcal day^{-1} were catabolized to produce ATP by OXPHOS, the net ATP regeneration rate would be 994 nmol (10^6 cells)$^{-1}$ h^{-1}. This level is similar to reported data such as that of rat heart H9c2 (normal) cells (1282.5 nmol (10^6 cells)$^{-1}$ h^{-1}) [39].

Supplementary figures

610 **Figure S1. Hierarchical clustering of 622 cancer cell line using publicly available expression data of 62 central metabolism-related genes.**
62 central carbon metabolism-related genes were obtained from the KEGG pathway database and selected due to there being no missing data in the dataset. For hierarchical clustering, the Ward method combined with the Euclidean distance was used. From each cluster, colorectal
615 cancer (HCT116, DLD-1, LS174T, and WiDr), pancreatic cancer (MIA Paca-2, and PANC-1), hepatoblastoma (HepG2), hepatocellular carcinoma (huH-1), breast cancer (MCF-7), renal cancer (Caki-1), and sarcoma (HT1080) were selected. Moreover, glioblastoma (YKG-1) was arbitrarily selected for the analysis.

Figure S2. Metabolic flux distribution of 12 cancer cell lines as determined by ^{13}C-MFA.
All the values represent the metabolic flux level (nmol (10^6 cells)$^{-1}$ h^{-1}). Measured specific cell proliferation rate (μ[h^{-1}]) and cell adhesion area (cell area [μm^2]), as well as the ATP regeneration rates by OXPHOS, substrate-level phosphorylation, and the net ATP regeneration rate are shown in the figure.

 Figure S2. (continued) Metabolic flux distribution of 12 cancer cell lines as determined by [13]C-MFA. All values represent metabolic flux level (nmol (10^6 cells)$^{-1}$ h^{-1}).

635 **Figure S3. Variations in flux levels in representative metabolic pathways.**
Flux levels of (a) GAPDH and (b) GLUDH and IDH reactions are presented. Error bars represent 95% confidence intervals.

Figure S4. Mitochondrial membrane potential is retained in cultured cancer cells.
The potential-dependent dye, JC-1, was used to monitor the mitochondrial membrane potential.

Figure S5. Scatter plots among the ATP regeneration rates and the visible phenotype data.
Spearman's rank order correlation coefficients were also shown.

Figure S6. Correlation among ATP regeneration rates and the visible phenotypes in the cell upon removal of the outlier cell line (Caki-1) from the dataset.
Spearman's rank order correlation coefficients are shown in the heat map (*n*=11).

Figure S7. **Comparison of measured metabolic flux data and predicted metabolic flux distributions by FBA for 12 cell lines.**

In silico simulation of metabolism was performed by FBA using a genome scale model of human metabolism (RECON2).

Supplementary tables

Table S1. Culture profiles of 12 cell lines.

Measured	Unit	Caki-1	DLD-1	HCT116	HepG2	HT-1080	Huh-1
Specific cell growth rate	h^{-1}	0.0228	0.0437	0.031	0.0316	0.0249	0.0175
cell adhesion area	μm^2 ($n = 30$)	3486 ± 792	1623 ± 468	645 ± 159	891 ± 300	1284 ± 533	1015 ± 322
Diameter of floating cells	μm ($n = 56\text{-}595$)	18.3 ± 3.7	12.4 ± 2.2	12.8 ± 1.5	14.1 ± 1.5	12.6 ± 1.9	15.0 ± 2.0
Glucose uptake	nmol (10^6 cells)$^{-1}$ h^{-1}	606 ± 232	742 ± 32.0	361 ± 17.0	755 ± 250	604 ± 75.8	397 ± 45.0
Glutamine uptake	nmol (10^6 cells)$^{-1}$ h^{-1}	327 ± 30	84 ± 4.8	98 ± 5.3	295 ± 46	278 ± 7.07	168 ± 9.3
Arginine uptake	nmol (10^6 cells)$^{-1}$ h^{-1}	22 ± 2	25 ± 1.2	25 ± 1.2	84 ± 6	56.9 ± 2.78	22 ± 2.2
Valine uptake	nmol (10^6 cells)$^{-1}$ h^{-1}	-32.9 ± 21.3	7 ± 0.7	5 ± 1.6	66 ± 26	8.5 ± 4.8	13.6 ± 2.8
Leucine uptake	nmol (10^6 cells)$^{-1}$ h^{-1}	4.2 ± 17.2	16 ± 0.5	12 ± 1.9	76 ± 22	22.2 ± 4.9	17.5 ± 2.0
Isoleucine uptake	nmol (10^6 cells)$^{-1}$ h^{-1}	-24.4 ± 22.5	9.4 ± 0.4	13.9 ± 2.7	76 ± 22	17.5 ± 5.2	32.9 ± 5.3
Lactate excretion	nmol (10^6 cells)$^{-1}$ h^{-1}	2158 ± 27	1311 ± 40	776 ± 3	1282 ± 29	1514 ± 13.0	708 ± 3
Glutamate excretion	nmol (10^6 cells)$^{-1}$ h^{-1}	143 ± 5	1 ± 1	22 ± 0	132 ± 14	133 ± 2.96	43 ± 2
Alanine excretion	nmol (10^6 cells)$^{-1}$ h^{-1}	0 ± 0	23.2 ± 1.3	8.8 ± 0.2	80 ± 7	7.14 ± 0.317	0.0 ± 0.0
Proline excretion	nmol (10^6 cells)$^{-1}$ h^{-1}	34.6 ± 0.4	9.2 ± 0.5	1.7 ± 0.1	21 ± 2	12.4 ± 0.472	5.4 ± 0.3
Ornithine excretion	nmol (10^6 cells)$^{-1}$ h^{-1}	18.2 ± 3.03	21.9 ± 1.2	23.9 ± 0.7	112 ± 19	62.9 ± 2.6	6.4 ± 1

670 **(Continued) Table S1. Culture profiles of 12 cell lines.**

Measured	Unit	LS174T	MCF7	MIAPACA2	Panc-1	WiDr	YKG-1
Specific cell growth rate	h^{-1}	0.0349	0.039	0.0406	0.0139	0.0279	0.0245
cell adhesion area	μm^2 ($n = 30$)	296 ± 93	1152 ± 372	838 ± 174	1263 ± 405	603 ± 196	1611 ± 588

Diameter of floating cells	μm (*n* = 56-595)	14.0 ± 2.0	12.2 ± 1.8	13.6 ± 1.8	15.6 ± 2.1	12.9 ± 1.6	13.8 ± 1.4
Glucose uptake	nmol (10^6 cells)$^{-1}$ h^{-1}	222 ± 21.0	507 ± 9.9	1031 ± 81.7	538 ± 133.5	481 ± 37.0	400 ± 43.98
Glutamine uptake	nmol (10^6 cells)$^{-1}$ h^{-1}	75 ± 2.7	66 ± 5.0	156 ± 7.1	167 ± 1.7	114 ± 3.8	184.2 ± 8.04
Arginine uptake	nmol (10^6 cells)$^{-1}$ h^{-1}	6 ± 0.2	37 ± 1.0	15 ± 1.2	9 ± 0.5	4 ± 0.6	63.76 ± 1.26
Valine uptake	nmol (10^6 cells)$^{-1}$ h^{-1}	-2 ± 1.3	2 ± 2.9	17.8 ± 2.4	7.09 ± 0.5	-1 ± 0.8	24.6 ± 3.9
Leucine uptake	nmol (10^6 cells)$^{-1}$ h^{-1}	9 ± 0.5	13 ± 2.7	28.2 ± 2.7	17.4 ± 0.7	4 ± 0.8	36.5 ± 4.1
Isoleucine uptake	nmol (10^6 cells)$^{-1}$ h^{-1}	7.4 ± 0.5	6.0 ± 1.9	28.4 ± 2.8	15.5 ± 1.2	-0.6 ± 0.3	33.2 ± 4.1
Lactate excretion	nmol (10^6 cells)$^{-1}$ h^{-1}	313 ± 7	1014 ± 6	1734 ± 29	1146 ± 41	762 ± 10	993 ± 7.18
Glutamate excretion	nmol (10^6 cells)$^{-1}$ h^{-1}	13 ± 1	16 ± 1	18 ± 1	41 ± 2	53 ± 2	43.7 ± 0.51
Alanine excretion	nmol (10^6 cells)$^{-1}$ h^{-1}	27.0 ± 0.6	22.0 ± 0.8	20.9 ± 0.9	0.0 ± 0.0	22.8 ± 0.2	19.75 ± 1.11
Proline excretion	nmol (10^6 cells)$^{-1}$ h^{-1}	3.5 ± 0.2	4.0 ± 0.2	3.1 ± 0.3	12.1 ± 0.1	4.4 ± 0.1	11.83 ± 0.286
Ornithine excretion	nmol (10^6 cells)$^{-1}$ h^{-1}	2.9 ± 0.1	43.0 ± 1.4	4.0 ± 1.2	4.69 ± 0.1	23.4 ± 0.4	59.7 ± 1.5

Table S2. Results of ¹³C-MFA of 12 cell lines.

	Caki-1	DLD1	HCT116	HepG2	HT-1080	Huh-1	LS174T	MCF7	MiaPaCa2	Panc1	WiDr	YKG1
Fitting results (χ^2 test)												
RSS	22.3	11.4	33.8	3.5	35.5	16.3	16.4	46.2	23.8	33.6	14.6	20.3
Threshold	71.0	66.3	62.8	33.9	52.2	51.0	62.8	51.7	59.3	55.8	51.0	41.3
p_value	0.99	0.99	0.90	0.99	0.54	0.99	0.99	0.12	0.99	0.75	0.99	0.85
Metabolic flux (nmol $(10^6$ cells$)^{-1}$ h^{-1})												
r1_Glc_uptake	1038	729	417	713	761	342	219	565	878	567	445	493
r2_PGI_fw	2814	6509	1473	3145	4888	2265	1138	5014	4158	2144	4355	1671
r3_PGI_rv	1820	5829	1083	2531	4206	1937	954	4489	3314	1614	3986	1200
r4_PFK	1060	706	403	675	731	335	202	546	860	552	416	482
r5_FBA_fw	2381	6078	2050	1550	3043	524	1452	4178	1478	577	977	1411
r6_FBA_rv	1321	5372	1647	875	2312	190	1250	3632	617	24	562	929
r7_TPI_fw	3067	6092	4002	5238	3491	2584	2080	2164	8672	2266	4254	3121
r8_TPI_rv	2008	5388	3600	4564	2761	2251	1880	1620	7813	1715	3840	2640
r9_GAPDH_fw	2936	2915	1393	1818	2526	2378	1891	4620	9407	1624	1934	2761
r10_GAPDH_rv	784	1493	582	438	1042	1707	1479	3520	7680	509	1081	1792
r11_PEPH_fw	3007	6469	1401	1523	2323	903	1826	4846	2814	1966	3666	3401
r12_PEPH_rv	861	5057	597	152	844	236	1424	3756	1097	854	2821	2438
r13_PK	2146	1411	804	1372	1478	667	403	1091	1717	1112	846	963
r16_PDH	121	93	65	31	56	73	80	66	50	62	67	56
r17_CS	121	93	65	31	56	73	80	66	50	62	67	56
r18_IDH_fw	145	59	42	15	34	81	73	74	33	72	53	42
r19_IDH_rv	48	12	9	18	4	27	30	48	25	24	16	11
r20_AKGDH	238	97	94	37	140	176	81	48	127	158	58	149

r21_SDH_fw	437	4095	229	7226	967	3255	664	80	6444	191	1547	922
r22_SDH_rv	199	3998	135	7189	827	3079	584	33	6318	33	1488	773
r23_FH_fw	6008	6205	337	1575	1401	1171	871	3266	10139	4201	4791	475
r24_FH_rv	5769	6107	243	1538	0	995	790	3218	10012	4043	4733	326
r25_MDH_fw	225	197	85	90	66	1652	2216	106	102	62	512	181
r26_MDH_rv	116	111	28	78	15	1680	2142	62	63	0	446	138
r27_PC	13	6	8	19	5	100	7	22	11	0	1	13
r28_MAE	147	44	61	49	108	217	34	33	118	106	13	125
r29_G6PDH	102	43	23	94	75	11	31	36	30	35	73	19
r30_6PGDH	102	43	23	94	75	11	31	36	30	35	73	19
r31_RPI_fw	6058	5880	2519	5254	3705	2977	1060	4532	10310	4389	2367	2001
r32_RPI_rv	6022	5863	2509	5221	3679	2972	1047	4518	10297	4376	2341	1993
r33_RPE_fw	5778	4351	2849	64	4634	3515	2220	4931	8754	4811	1992	3673
r34_RPE_rv	5712	4325	2836	3	4585	3509	2202	4910	8737	4788	1945	3662
r35_TKL1_fw	346	44	2188	6742	439	10	21	42	33	85	3682	27
r36_TKL1_rv	313	31	2181	6712	415	7	12	32	25	73	3659	21
r37_TAL1_fw	100	81	32	7533	94	22	1336	5070	4653	50	37	59
r38_TAL1_rv	67	68	25	7503	70	19	1327	5059	4644	39	14	53
r39_TKL2_fw	33	13	7	5435	24	3	9	11	9	11	23	5
r40_TKL2_rv	0	0	0	5405	0	0	0	0	0	0	0	0
r41_LDH	2154	1321	775	1282	1511	707	313	1003	1743	1153	761	992
r42_Lac_excretion	2154	1321	775	1282	1511	707	313	1003	1743	1153	761	992
r46_Gln_uptake	329	84	102	251	277	165	72	71	154	166	114	185
r47_GLC	325	78	98	247	273	163	67	65	148	164	110	182
r49_GDH_fw	6059	254	382	39	604	703	222	445	1031	538	481	119

r50_GDH_rv	5918	204	321	0	494	580	185	422	912	428	460	0
r51_Glu_excretion	143	1	22	136	133	43	13	16	18	41	53	44
r52_Pro	38	15	6	25	16	8	8	9	9	14	8	15
r53_Pro_excretion	35	9	2	21	12	6	4	4	3	12	4	12
r54_Asp	7	13	9	10	8	5	11	12	12	4	8	7
r55_Asp_excretion	0	0	0	0	0	0	0	0	0	0	0	0
r56_Asn	3	5	4	4	3	2	4	5	5	2	3	3
r57_Asn_excretion	0	0	0	0	0	0	0	0	0	0	0	0
r58_GPT	6	34	17	88	14	4	36	32	31	4	30	26
r59_Ala_excretion	0	23	9	80	7	0	27	22	21	0	23	20
r71_Leu_uptake	6	11	7	8	0	4	8	9	10	3	7	6
r72_Ile_uptake	3	6	4	4	4	2	5	5	6	2	4	3
r75_Val_uptake	4	8	5	6	4	3	6	7	7	2	5	4
r77_Arg_uptake	22	25	26	83	57	22	6	37	15	10	5	64
r78_Arg_degradation	18	18	21	78	53	19	0	31	8	7	0	60
r79_Orn_excretion	18	22	24	120	63	6	3	42	4	5	23	60
r82_ACYL	24	46	32	33	26	19	37	41	43	15	30	26
r83_AcCOA_BIO	24	46	32	33	26	19	37	41	43	15	30	26
r84_G6P_BIO	3	5	4	4	3	2	4	5	5	2	3	3
r85_R5P_BIO	2	4	3	3	2	2	3	4	4	1	3	2
r86_DHAP_BIO	1	2	2	2	1	1	2	2	2	1	1	1
r87_Ser_BIO	6	11	8	8	6	4	9	10	10	4	7	6
r89_Ala_BIO	6	11	8	8	6	4	9	10	10	4	7	6
r90_Asp_BIO	4	8	5	6	4	3	6	7	7	2	5	4
r91_Asn_BIO	3	5	4	4	3	2	4	5	5	2	3	3
r92_Glu_BIO	4	7	5	5	4	3	6	6	7	2	5	4

r93_Gln_BIO	3	6	4	4	3	2	5	5	6	2	4	3
r94_Pro_BIO	3	6	4	4	3	2	5	5	5	2	4	3
r95_Arg_BIO	4	7	5	5	4	3	6	6	7	2	5	4
r96_Leu_BIO	6	11	7	8	0	4	8	9	10	3	7	6
r97_Ile_BIO	3	6	4	4	4	2	5	5	6	2	4	3
r98_Val_BIO	4	8	5	6	4	3	6	7	7	2	5	4
r100_CO2_in	6060	6631	3610	7546	6039	3970	2220	5070	10310	556	4810	366
r101_CO2_ex	5367	6313	3343	7356	5635	3539	1958	4884	9989	147	4562	0
r104_MAL_transport_fw	69	6758	3103	71	3495	74	2190	1512	8912	30	4789	3634
r105_MAL_transport_rv	86	6791	3126	95	3514	87	2216	1541	8942	41	4810	3653
r106_MDH_cyt_fw	5041	81	599	5804	7	3861	2193	5041	9421	66	1880	3297
r107_MDH_cyt	5058	114	622	5828	26	3874	2219	5070	9451	77	1901	3316
r108_Net_ATP_production	4086	2618	1674	1817	2396	1571	1460	1806	2455	2205	1731	1908
r109_OXPHOS	688	451	317	174	316	333	397	280	254	376	334	327
r110_Net_NADPH_production	351	130	107	238	258	238	96	104	178	177	158	163
PGI	994	680	390	615	683	329	184	525	843	530	369	471
FBA	1060	706	403	675	731	335	202	546	860	552	416	482
TPI	1059	704	402	674	730	334	200	544	858	552	414	481
GAPDH	2152	1423	812	1380	1485	671	412	1101	1727	1115	853	969
PEPH	2146	1411	804	1372	1478	667	403	1091	1717	1112	846	963
IDH	97	47	32	-2	30	54	43	25	7	47	37	30
SDH	238	97	94	37	140	176	81	48	127	158	58	149
MDH	108	87	56	12	51	-27	73	44	39	62	66	43
RPI	36	17	10	34	27	5	13	14	13	13	26	8
RBE	66	26	13	61	48	6	18	21	17	23	46	11

TKT1	33	13	7	30	24	3	9	11	9	11	23	5
TAL	33	13	7	30	24	3	9	11	9	11	23	5
TKT2	33	13	7	30	24	3	9	11	9	11	23	5
FH	238	97	94	37	140	176	81	48	127	158	58	149
MDHcyt	-17	-33	-23	-24	-19	-13	-26	-29	-31	-11	-21	-19
GDH	141	51	62	39	110	122	37	22	119	110	21	119

Table S3. Constraints of medium components for flux balance analysis.

Constraint	Medium components
Uptake: Forbidden, Excretion: Allowed	M_nh4_b, M_co2tot_b, M_lac_L_b, M_urea_b, M_gudac_b, M_biomass_b
Uptake: Allowed, Excretion: Allowed	M_glc_D_b, M_gln_L_b, M_ile_L_b, M_leu_L_b, M_lys_L_b, M_phe_L_b, M_thr_L_b, M_trp_L_b, M_tyr_L_b, M_val_L_b, M_his_L_b, M_met_L_b, M_chol_b, M_h_b, M_h2o_b, M_pi_b, M_o2_b, M_biomass_other_c, ent_ex_
Fixed at measured value	M_arg_L_b, M_orn_b, M_glu_L_b, M_pro_L_b, M_ala_L_b, M_asn_L_b, M_asp_L_b
Uptake: Forbidden, Excretion: Forbidden	M_13_cis_oretn_b, M_13_cis_retn_b, M_Asn_X_Ser_Thr_b, M_avite1_b, M_avite2_b, M_bvite_b, M_core5_b, M_core7_b, M_core8_b, M_datp_b, M_datp_b, M_dctp_b, M_dctp_b, M_dem2emgacpail_prot_hs_b, M_dgpi_prot_hs_b, M_dgtp_b, M_dgtp_b, M_dttp_b, M_dttp_b, M_ethamp_b, M_gncore2_b, M_gpi_sig_b, M_hretn_b, M_mem2emgacpail_prot_hs_b, M_n5m2masn_b, M_oretn_b, M_Ser_Thr_b, M_Ser_Gly_Ala_X_Gly_b, M_sprm_b, M_T_antigen_b, M_yvite_b, M_citr_L_b, M_pre_prot_b, M_c81coa_b, M_dd2coa_b, M_decdicoa_b, M_octdececoa_b, M_tetdece1coa_b, M_1a25dhvitd3_b, M_4abut_b, M_5HPET_b, M_taur_b, M_pe_hs_b, M_pmtcoa_b, M_5HPET_b, M_4hpro_LT_b, M_Lcystin_b, M_anth_b, M_btn_b, M_fol_b, M_ncam_b, M_pnto_R_b, M_10fthf_b, M_11_cis_retfa_b, M_13_cis_retnglc_b, M_1glyc_hs_b, M_1mncam_b, M_2425dhvitd2_b, M_2425dhvitd3_b, M_24nph_b, M_25hvitd2_b, M_25hvitd3_b, M_2hb_b, M_2mcit_b, M_34dhoxpeg_b, M_34dhphe_b, M_35cgmp_b, M_3aib_b, M_3aib_D_b, M_3mlda_b, M_4abut_b, M_4hdebrisoquine_b, M_4hphac_b, M_4mptnl_b, M_4mtolbutamide_b, M_4nph_b, M_4nphsf_b, M_4pyrdx_b, M_5adtststerone_b, M_5adtststeroneglc_b, M_5fthf_b, M_5homeprazole_b, M_5htrp_b, M_5mthf_b, M_6htststerone_b, M_9_cis_retfa_b, M_abt_b, M_ac_b, M_acac_b, M_acald_b, M_acetone_b, M_acgalfucgalacgalfuc12gal14acglcgalgluside_hs_b, M_acgalfucgalacgalfucgalacglcgal14acglcgalgluside_hs_b, M_acgam_b, M_ach_b, M_ade_b, M_adn_b, M_adp_b, M_adprbp_b, M_adprib_b, M_adrn_b, M_adrnl_b, M_aflatoxin_b, M_ahandrostanglc_b, M_ak2lgchol_hs_b, M_akg_b, M_ala_B_b, M_ala_D_b, M_aldstrn_b, M_amp_b, M_andrstrn_b, M_andrstrnglc_b, M_antipyrene_b, M_apnnox_b, M_appnn_b, M_aprgstrn_b, M_aqcobal_b, M_arab_L_b,

M_arach_b, M_arachd_b, M_ascb_L_b, M_asp_D_b, M_atp_b,
M_avite1_b, M_avite2_b, M_bhb_b, M_bildglcur_b,
M_bilglcur_b, M_bilirub_b, M_biocyt_b, M_btn_b, M_but_b,
M_bvite_b, M_bz_b, M_ca2_b, M_camp_b, M_caro_b,
M_carveol_b, M_cgly_b, M_cholate_b, M_chsterol_b, M_chtn_b,
M_cit_b, M_clpnd_b, M_cmp_b, M_co_b, M_coumarin_b,
M_creat_b, M_crmp_hs_b, M_crn_b, M_crtsl_b, M_crtstrn_b,
M_crvnc_b, M_csn_b, M_cyan_b, M_cys_L_b, M_cytd_b,
M_dad_2_b, M_dag_hs_b, M_dcsptn1_b, M_dcyt_b,
M_debrisoquine_b, M_dgchol_b, M_dgsn_b, M_dhdascb_b,
M_dheas_b, M_dhf_b, M_digalsgalside_hs_b, M_din_b,
M_dlnlcg_b, M_dmantipyrine_b, M_dmhptcrn_b, M_dopa_b,
M_dopasf_b, M_drib_b, M_duri_b, M_eaflatoxin_b,
M_ebastine_b, M_ebastineoh_b, M_eicostet_b, M_elaid_b,
M_estradiol_b, M_estradiolglc_b, M_estriolglc_b,
M_estroneglc_b, M_estrones_b, M_etoh_b, M_fe2_b, M_fe3_b,
M_fol_b, M_for_b, M_fru_b,
M_fuc13galacglcgal14acglcgalgluside_hs_b,
M_fuc14galacglcgalgluside_hs_b,
M_fucacgalfucgalacglcgalgluside_hs_b,
M_fucacngal14acglcgalgluside_hs_b,
M_fucfuc12gal14acglcgalgluside_hs_b,
M_fucfuc132galacglcgal14acglcgalgluside_hs_b,
M_fucfucfucgalacglc13galacglcgal14acglcgalgluside_hs_b,
M_fucfucfucgalacglcgal14acglcgalgluside_hs_b,
M_fucfucgalacglcgalgluside_hs_b,
M_fucgal14acglcgalgluside_hs_b,
M_fucgalfucgalacglcgalgluside_hs_b, M_fucgalgbside_hs_b,
M_fuc_L_b, M_gal_b, M_galacglcgalgbside_hs_b,
M_galfuc12gal14acglcgalgluside_hs_b,
M_galfucgalacglcgal14acglcgalgluside_hs_b,
M_galgalfucfucgalacglcgalacglcgal14acglcgalgluside_hs_b,
M_galgalgalthcrm_hs_b, M_gam_b, M_gbside_hs_b,
M_gchola_b, M_gdp_b, M_gluala_b, M_gly_b, M_glyb_b,
M_glyc_b, M_glyc_S_b, M_glygn2_b, M_glygn4_b,
M_glygn5_b, M_gmp_b, M_gsn_b, M_gt1a_hs_b, M_gthox_b,
M_gthrd_b, M_gtp_b, M_gua_b, M_h_b, M_h2o2_b, M_ha_b,
M_ha_pre1_b, M_hco3_b, M_hcoumarin_b, M_hdca_b,
M_hdcea_b, M_hestratriol_b, M_hexc_b, M_hista_b,
M_hom_L_b, M_hpdca_b, M_htaxol_b, M_hxan_b, M_i_b,
M_idp_b, M_imp_b, M_inost_b, M_ins_b, M_lac_D_b, M_lcts_b,
M_Lcystin_b, M_leuktrA4_b, M_leuktrB4_b, M_leuktrC4_b,
M_leuktrD4_b, M_leuktrE4_b, M_leuktrF4_b, M_lgnc_b,
M_limnen_b, M_lipoate_b, M_lnlc_b, M_lnlnca_b, M_lnlncg_b,
M_lpchol_hs_b, M_mag_hs_b, M_malt_b, M_malttr_b,
M_man_b, M_meoh_b, M_mepi_b, M_mthgxl_b,
M_n2m2nmasn_b, M_nac_b, M_nad_b, M_nadp_b, M_ncam_b,
M_no_b, M_npthl_b, M_nrpphr_b, M_nrpphrsf_b, M_nrvnc_b,

M_o2s_b, M_ocdca_b, M_ocdcea_b, M_octa_b,
M_omeprazole_b, M_onpthl_b, M_oxa_b, M_paf_hs_b,
M_pchol_hs_b, M_pe_hs_b, M_peplys_b, M_perillyl_b,
M_pglyc_hs_b, M_pheacgln_b, M_pheme_b, M_phyQ_b,
M_phyt_b, M_pnto_R_b, M_ppa_b, M_prgstrn_b, M_pro_D_b,
M_prostgd2_b, M_prostge1_b, M_prostge2_b, M_prostgf2_b,
M_ps_hs_b, M_ptdca_b, M_pydam_b, M_pydx_b, M_pydxn_b,
M_pyr_b, M_rbt_b, M_retfa_b, M_retinol_b, M_retinol_9_cis_b,
M_retinol_cis_11_b, M_retn_b, M_retnglc_b, M_rib_D_b,
M_ribflv_b, M_Rtotal_b, M_sarcs_b, M_sel_b, M_ser_D_b,
M_ser_L_b, M_sl_L_b, M_so4_b, M_spc_hs_b, M_sph1p_b,
M_sphs1p_b, M_srtn_b, M_strch1_b, M_strch2_b, M_strdnc_b,
M_succ_b, M_sucr_b, M_tag_hs_b, M_tagat_D_b, M_taur_b,
M_taxol_b, M_tchola_b, M_tcynt_b, M_tdchola_b, M_tethex3_b,
M_tetpent3_b, M_thf_b, M_thm_b, M_thmmp_b, M_thmtp_b,
M_thym_b, M_thymd_b, M_thyox_L_b, M_tmndnc_b,
M_tolbutamide_b, M_tre_b, M_triodthy_b, M_triodthysuf_b,
M_tststerone_b, M_tststeroneglc_b, M_tsul_b, M_ttdca_b,
M_txa2_b, M_tymsf_b, M_Tyr_ggn_b, M_udp_b, M_ump_b,
M_ura_b, M_urate_b, M_uri_b, M_utp_b, M_vacc_b, M_vitd2_b,
M_vitd3_b, M_whddca_b, M_whhdca_b, M_whtststerone_b,
M_xolest2_hs_b, M_xoltri24_b, M_xoltri25_b, M_xoltri27_b,
M_xyl_D_b, M_xylt_b, M_yvite_b, M_4abutn_b, M_acmana_b,
M_ahdt_b, M_ctp_b, M_dgmp_b, M_dgtp_b, M_dha_b,
M_dhap_b, M_dtmp_b, M_dttp_b, M_fad_b, M_fald_b,
M_g1p_b, M_HC00229_b, M_HC00250_b, M_HC01104_b,
M_HC01361_b, M_HC01440_b, M_HC01441_b,
M_HC01444_b, M_HC01446_b, M_HC01577_b,
M_HC01609_b, M_HC01610_b, M_HC01700_b,
M_HC02160_b, M_HC02161_b, M_itp_b, M_orot_b, M_prpp_b,
M_ptrc_b, M_pydx5p_b, M_spmd_b, M_udpg_b, M_no2_b,
M_so3_b, M_sprm_b, M_prostgh2_b, M_prostgi2_b, M_ppi_b,
M_cdp_b, M_dtdp_b, M_HC00955_b, M_citr_L_b,
M_HC01787_b, M_C02470_b, M_HC00822_b, M_C02528_b,
M_HC02192_b, M_CE5560_b, M_HC02195_b, M_HC02196_b,
M_HC02220_b, M_HC02154_b, M_ddca_b, M_HC02199_b,
M_HC02200_b, M_HC02201_b, M_HC02172_b,
M_HC02191_b, M_HC02194_b, M_HC02197_b,
M_HC02198_b, M_HC02187_b, M_HC02180_b,
M_wharachd_b, M_HC02202_b, M_HC02203_b,
M_HC02204_b, M_HC02205_b, M_HC02206_b,
M_HC02207_b, M_HC02208_b, M_HC02210_b,
M_HC02213_b, M_HC02214_b, M_HC02216_b,
M_HC02217_b, M_malcoa_b, M_arachcoa_b, M_coa_b,
M_CE2250_b, M_CE1940_b, M_CE2011_b, M_maltttr_b,
M_maltpt_b, M_malthx_b, M_malthp_b, M_CE2839_b,
M_CE2838_b, M_CE1950_b, M_cynt_b, M_23cump_b,
M_3ump_b, M_CE5797_b, M_CE5798_b, M_CE4633_b,

M_CE4881_b, M_glcur_b, M_udpgal_b, M_crm_hs_b,
M_galside_hs_b, M_CE0074_b, M_cdpea_b, M_12HPET_b,
M_C04849_b, M_3bcrn_b, M_3ddcrn_b, M_3deccrn_b,
M_3hdececrn_b, M_3hexdcrn_b, M_3ivcrn_b, M_3octdec2crn_b,
M_3octdeccrn_b, M_3octdece1crn_b, M_3tdcrn_b,
M_3tetd7ecoacrn_b, M_3thexddcoacrn_b, M_3ttetddcoacrn_b,
M_c101crn_b, M_c10crn_b, M_c4crn_b, M_c51crn_b,
M_c5dc_b, M_c6crn_b, M_c81crn_b, M_c8crn_b,
M_ddece1crn_b, M_ddeccrn_b, M_decdicrn_b, M_doco13ac_b,
M_docosac_b, M_ivcrn_b, M_ttdcrn_b, M_tetdec2crn_b,
M_tetdece1crn_b, M_dca_b, M_4hpro_LT_b, M_adocbl_b,
M_alaala_b, M_bglc_b, M_carn_b, M_dchac_b, M_glgchlo_b,
M_gltcho_b, M_glygly_b, M_glyleu_b, M_glyphe_b,
M_glypro_b, M_glysar_b, M_gum_b, M_gumgchol_b,
M_gumtchol_b, M_leugly_b, M_leuleu_b, M_pectingchol_b,
M_pectintchol_b, M_progly_b, M_psyl_b, M_psylchol_b,
M_psyltchol_b, M_sbt_D_b, M_slfcys_b, M_tdechola_b,
M_cysam_b, M_dpcoa_b, M_fmn_b, M_hyptaur_b, M_oh1_b,
M_pan4p_b, M_ptth_b, M_q10_b, M_q10h2_b, M_34hpp_b,
M_3mob_b, M_3mop_b, M_4mop_b, M_5mta_b, M_5oxpro_b,
M_ahcys_b, M_aicar_b, M_anth_b, M_cbasp_b, M_mal_L_b,
M_idour_b, M_fum_b, M_xmp_b, M_dcmp_b, M_HC00342_b,
M_glyc3p_b

Table S4. Comparison of thermodynamic parameters.

	Glucose	O_2	Lactic acid	Total CO_2	Generation per mol ATP	
					Aerobic glycolysis (Glucose => 2 Lactic acid)	OXPHOS (Glucose + 6 O_2 => 6 Total CO_2)
Enthalpy change (heat),$\Delta_r H'°$ kJ mol^{-1} [1]	-1266.0	-11.7	-688.0	-693.0	-55.0	-88.2
Entropy, S_m J K^{-1} mol^{-1} [4]	212	204.8	192	213 (CO_2) + 70 (H_2O)	86.0	8.0
Gibbs free energy change, $\Delta_r G'°$ kJ mol^{-1} [1]	-429.0	0.0	-314.0	-547.0	-99.5	-89.2
Gibbs free energy change, $\Delta_r G'°$ kJ mol^{-1} [2]	-429.0	16.4	-311.0	-548.0	-96.5	-92.4
Gibbs free energy change, $\Delta_r G'^m$ kJ mol^{-1} [2]	-446.0	-0.7	-328.0	-565.0	-105.0	-91.9
Gibbs free energy change, $\Delta_r G'$ kJ mol^{-1} [3]	-437.6	-5.0	-324.7	-553.5	-105.9	-89.2
Concentration in medium (mM)	16	0.24	3.0	44 (3.7 g/L sodium bicarbonate)		
Arbitrarily selected irrelevant parameters						
Molecular Weight	180	32	90	44	0.0	0.0
Melting point [5]	146	-218	18	-55.6	-55.0	25.9
Boiling point [5]	999	-182	122	-78.5	-377.5	6.8
Topological Polar Surface Area [6]	110	34.1	57.5	34.1	2.5	-3.3
logp [6]	-2.6	-1.1	-0.7	0.9	0.6	0.4

1) At 298.15 K, 1 Bar, pH 7, ionic strength 0.1M retrieved from [22].

2) Retrieved from eQuilibrator pH 7, ionic strength 0.1M on 24th March 2021 (https://equilibrator.weizmann.ac.il/).

3) Calculated from $\Delta_r G'^m$ kJ mol^{-1} at 310.15 K given concentration.

4) Retrieved from Calculla (https://calculla.com/standard_entropy).

5) Retrieved from Wikipedia (https://ja.wikipedia.org/).

6) Retrieved from PubChem (https://pubchem.ncbi.nlm.nih.gov/).

Table S5. Standard enthalpy of formation.

	$\Delta_r H^{\prime \circ}$ kJ mol^{-1} [1]
Glucose	-1266
CO_2 total	-693
H_2O	-286
Lactate	-688
$NH_4^+(NH_3(aq))$	-133
Glutamine	-808
Glutamate	-982
Urea	-319
O_2	-11.7
H^+	0
Alanine	-557
Proline	-515 [4]
Phosphate	-1300
Arginine	-621 [4]
Isoleucine	-638 [4]
Leucine	-637 [4]
Lysine	-756 [5]
Phenylalanine	-467 [4]
Threonine	-807 [4]
Tryptophan	-415 [4]
Tyrosine	-685 [4]
Valine	-617 [4]
Histidine	-467 [4]
Methionine	-578 [4]
Ornithine	-694 [2]
Aspartate	-945
Asparagine	-789.4 [4]
Na+	-240 [4]
Cl-	-167 [4]
Glycine	-525
Serine	-732.7 [4]
Citrate	-1514
Biomass	-5.5 kJ/g [3]

1) At 298.15 K, 1 Bar, pH 7, ionic strength 0.1M retrieved from [22].
2) Collected from [40].
3) Collected from [35]. Means of yeast data were used in this study.
4) Collected from textbook (Atkins et al. Physical Chemistry for the Life Sciences)
5) Collected from [41].

Table S6. RNAseq analysis of HCT116 and MCF-7 cells cultured under different temperatures.

(A) Summary of RNAseq analysis

SampleName	NumOfReads	reference	mapping_rate	NumOfGeneCountZero
MCF-7_36°C	15,559,985	hg19	95.20%	10,029/26,256
MCF-7_37°C	19,105,357	hg19	95.30%	9,681/26,256
MCF-7_38°C	18,858,384	hg19	95.40%	9,576/26,256
HCT116_36°C	19,749,645	hg19	94.60%	9,769/26,256
HCT116_37°C	21,646,230	hg19	94.80%	9,714/26,256
HCT116_38°C	18,735,250	hg19	94.20%	9,821/26,256

(B) Comparison of gene expression level

	HCT166	MCF-7	HCT166 & MCF-7
36°C < 37°C <38°C	3606	3975	619
36°C > 37°C <38°C	4906	3793	959
36°C > 37°C >38°C	3154	4533	686
36°C < 37°C >38°C	5208	4576	1144
Others (ex. Zero expression)	9382	9379	not determined

Table S7. Gene enrichment analysis using the KEGG 2019 human dataset by Enrichr.

Term	Adjusted P-value	Genes
Cell cycle	1.43E-05	PCNA;PLK1;CDC6;CDC25A;CCNA2;PTTG1;CCNE2;ORC1;TFDP2;ESPL1;ORC3;MDM2;CDK1;MCM3;ANAPC4;ANAPC5;E2F3;MCM6
Proteasome	1.43E-05	PSMA5;PSMA6;PSMD12;PSMD6;PSMA3;PSMA2;PSMC1;POMP;PSMB1;PSME4;PSMD1
Spliceosome	2.10E-05	TCERG1;HNRNPA3;RBM8A;DDX46;SF3B6;SRSF1;HNRNPU;WBP11;LSM3;U2SURP;HNRNPK;PRPF3;TRA2B;SRSF3;SNRPF;SRSF6;SMNDC1;RBMX
DNA replication	8.40E-05	RFC5;PCNA;RNASEH1;RFC4;LIG1;RPA1;MCM3;MCM6;DNA2
Citrate cycle (TCA cycle)	1.54E-04	PDHA1;MDH2;IDH3G;IDH2;IDH3B;OGDHL;DLAT;SDHD
Oocyte meiosis	7.20E-03	PLK1;PPP2CA;PPP1CC;RPS6KA3;PPP3CB;PTTG1;CCNE2;ESPL1;CDK1;ANAPC4;ANAPC5;CALM1;CAMK2G
Mismatch repair	2.34E-02	RFC5;PCNA;RFC4;LIG1;RPA1
Pyruvate metabolism	4.13E-02	ALDH3A2;LDHA;PDHA1;MDH2;DLAT;ALDH7A1

Table S8. Metabolic model for ^{13}C-MFA.

Reaction ID	Stoichiometry	Atom mapping
r1_Glc_uptake	SubsGlc + ATP --> G6P	ABCDEF --> ABCDEF
r2_PGI_fw	G6P --> F6P	ABCDEF --> ABCDEF
r3_PGI_rv	F6P --> G6P	ABCDEF --> ABCDEF
r4_PFK	F6P + ATP --> FBP	ABCDEF --> ABCDEF
r5_FBA_fw	FBP --> DHAP + GAP	ABCDEF --> CBA + DEF
r6_FBA_rv	DHAP + GAP --> FBP	CBA + DEF --> ABCDEF
r7_TPI_fw	DHAP --> GAP	ABC --> ABC
r8_TPI_rv	GAP --> DHAP	ABC --> ABC
r9_GAPDH_fw	GAP --> PGA + NADH + ATP	ABC --> ABC
r10_GAPDH_rv	PGA + NADH + ATP --> GAP	ABC --> ABC
r11_PEPH_fw	PGA --> PEP	ABC --> ABC
r12_PEPH_rv	PEP --> PGA	ABC --> ABC
r13_PK	PEP --> Pyr + ATP	ABC --> ABC
r16_PDH	Pyr --> AcCOAmit + CO2in + NADH	ABC --> BC + A
r17_CS	AcCOAmit + Oxa --> Cit	AB + CDEF --> FEDBAC
r18_IDH_fw	Cit --> aKG + CO2in + NADH	ABCDEF --> ABCDE + F
r19_IDH_rv	aKG + CO2in + NADH --> Cit	ABCDE + F --> ABCDEF
r20_AKGDH	aKG --> Suc + CO2in + ATP + NADH	ABCDE --> BCDE + A
r21_SDH_fw	Suc-->Fum+{0.6}NADH	ABCD --> ABCD
r22_SDH_rv	Fum+{0.6}NADH-->Suc	ABCD --> ABCD
r23_FH_fw	Fum-->Mal	ABCD --> ABCD
r24_FH_rv	Mal-->Fum	ABCD --> ABCD
r25_MDH_fw	Mal --> Oxa+NADH	ABCD --> ABCD
r26_MDH_rv	Oxa+NADH --> Mal	ABCD --> ABCD
r27_PC	Pyr + CO2in + ATP --> Oxa	ABC + D --> ABCD
r28_MAE	Mal --> Pyr + CO2in + NADPH	ABCD --> ABC + D
r29_G6PDH	G6P --> m6PG + NADPH	ABCDEF --> ABCDEF
r30_6PGDH	m6PG --> Ru5P + CO2in + NADPH	ABCDEF --> BCDEF + A
r31_RPI_fw	Ru5P --> R5P	ABCDE --> ABCDE
r32_RPI_rv	R5P --> Ru5P	ABCDE --> ABCDE
r33_RPE_fw	Ru5P --> Xu5P	ABCDE --> ABCDE
r34_RPE_rv	Xu5P --> Ru5P	ABCDE --> ABCDE

r35_TKL1_fw	R5P + Xu5P --> S7P + GAP	ABCDE + FGHIJ --> FGABCDE + HIJ
r36_TKL1_rv	GAP + S7P --> Xu5P + R5P	HIJ + FGABCDE --> FGHIJ + ABCDE
r37_TAL1_fw	GAP + S7P --> F6P + E4P	ABC + DEFGHIJ --> DEFABC + GHIJ
r38_TAL1_rv	E4P + F6P --> S7P + GAP	GHIJ + DEFABC --> DEFGHIJ + ABC
r39_TKL2_fw	E4P + Xu5P --> F6P + GAP	ABCD + EFGHI --> EFABCD + GHI
r40_TKL2_rv	GAP + F6P --> Xu5P + E4P	GHI + EFABCD --> EFGHI + ABCD
r41_LDH	Pyr + NADH --> Lac	ABC --> ABC
r42_Lac_excretion	Lac --> LacEx	nd
r46_Gln_uptake	SubsGln --> Gln	ABCDE --> ABCDE
r47_GLC	Gln --> Glu	ABCDE --> ABCDE
r49_GDH_fw	Glu --> aKG	ABCDE --> ABCDE
r50_GDH_rv	aKG --> Glu	ABCDE --> ABCDE
r51_Glu_excretion	Glu --> GluEx	nd
r52_Pro	Glu --> Pro	ABCDE --> ABCDE
r53_Pro_excretion	Pro --> ProEx	nd
r54_Asp	Oxacyt --> Asp	ABCD --> ABCD
r55_Asp_excretion	Asp --> AspEx	nd
r56_Asn	Asp --> Asn	ABCD --> ABCD
r57_Asn_excretion	Asn --> AsnEx	nd
r58_GPT	Pyr --> Ala	ABC --> ABC
r59_Ala_excretion	Ala --> AlaEx	nd
r71_Leu_uptake	SubsLeu --> Leu	ABCDEF --> ABCDEF
r72_Ile_uptake	SubsIle --> Ile	ABCDEF --> ABCDEF
r75_Val_uptake	SubsVal --> Val	ABCDE --> ABCDE
r77_Arg_uptake	SubsArg --> Arg	ABCDEF --> ABCDEF
r78_Arg_degradation	Arg --> Glu	ABCDEF --> ABCDE
r79_Orn_excretion	Glu --> OrnEx	nd
r82_ACYL	Cit --> AcCOAcyt + Oxacyt	FEDBAC --> AB + CDEF
r83_AcCOA_BIO	AcCOAcyt --> AcCOABiomass	nd
r84_G6P_BIO	G6P --> G6PBiomass	nd
r85_R5P_BIO	R5P --> R5PBiomass	nd
r86_DHAP_BIO	DHAP --> DHAPBiomass	nd
r87_Ser_BIO	PGA --> SerBiomass	nd
r89_Ala_BIO	Ala --> AlaBiomass	nd
r90_Asp_BIO	Asp --> AspBiomass	nd
r91_Asn_BIO	Asn --> AsnBiomass	nd
r92_Glu_BIO	Glu --> GluBiomass	nd

r93_Gln_BIO	Gln --> GlnBiomass	nd
r94_Pro_BIO	Pro --> ProBiomass	nd
r95_Arg_BIO	Arg --> ArgBiomass	nd
r96_Leu_BIO	Leu --> LeuBiomass	nd
r97_Ile_BIO	Ile --> IleBiomass	nd
r98_Val_BIO	Val --> ValBiomass	nd
r100_CO2_in	CO2in --> CO2Ex	A --> A
r101_CO2_ex	SubsCO2 --> CO2in	A --> A
r104_MAL_transport_fw	Mal --> Malcyt	ABCD --> ABCD
r105_MAL_transport_rv	Malcyt --> Mal	ABCD --> ABCD
r106_MDH_cyt_fw	Malcyt --> Oxacyt + NADH	ABCD --> ABCD
r107_MDH_cyt	Oxacyt + NADH --> Malcyt	ABCD --> ABCD
r108_Net_ATP_production	ATP --> ATPex	nd
r109_OXPHOS	NADH --> 2.5 ATP	nd
r110_Net_NADPH_production	NADPH --> NADPHex	nd

Supplementary references

36.   Okahashi, N. *et al.* Sugar phosphate analysis with baseline separation and soft ionization by gas chromatography-negative chemical ionization-mass spectrometry improves flux estimation of bidirectional reactions in cancer cells. *Metab. Eng.* (2018). doi:10.1016/J.YMBEN.2018.08.011

37.   Antoniewicz, M. R., Kelleher, J. K. & Stephanopoulos, G. Determination of confidence intervals of metabolic fluxes estimated from stable isotope measurements. *Metab. Eng.* **8**, 324–337 (2006).

38.   Bianconi, E. *et al.* An estimation of the number of cells in the human body. *Ann. Hum. Biol.* **40**, 463–471 (2013).

39.   Abas, L., Bogoyevitch, M. A. & Guppy, M. Mitochondrial ATP production is necessary for activation of the extracellular-signal-regulated kinases during ischaemia/reperfusion in rat myocyte-derived H9c2 cells. *Biochem. J.* **349**, 119–126 (2000).

40.   Lutkin, A. I., Chernikov, V. V., Krutova, O. N. & Skvortsov, I. A. Standard enthalpies of formation of l-lysine and the products of its dissociation in aqueous solutions. *J. Therm. Anal. Calorim. 2017 1301* **130**, 457–460 (2017).

41.   Lytkin, A. I., Chernikov, V. V. & Krutova, O. N. Standard Enthalpies of Formation of L-Ornithine and Products of Its Dissociation in Aqueous Solutions. *Russ. J. Inorg. Chem. 2018 638* **63**, 1112–1115 (2018).

715
720
725
730