Stimulation of immunity-linked genes by membrane disruption is linked to Golgi function

and the ARF-1 GTPase

Christofer M. Welsh[1], Lorissa J. Smulan[2], Matthew J. Fanelli[1], Dominique S. Lui[1], and

Amy K. Walker[1]

[1]Program in Molecular Medicine, UMASS Medical School, Worcester, MA

[2]Department of Medicine, UMASS Medical School, Worcester, MA

Contact: amy.walker@umassmed.edu

Summary

Immune-linked genes (ILGs) are activated in response to pathogens but can also be activated by lipid imbalance. Why pathogen attack and metabolic changes both impact ILG activation is unclear. Organelles in the secretory pathway have distinct protein and lipid components and genetically separable stress programs. These stress pathways activate restorative transcriptional programs when lipid ratios become unbalanced or during dysregulated protein folding and trafficking. We find that ILGs are specifically activated when membrane phosphatidylcholine ratios change in the secretory pathway. Consistent with this result, disruption of Golgi function after RNAi targeting the ADP-ribosylation factor ARF-1 also activates ILG expression. Since increased protein secretion is altered by metabolic changes and pathogen responses, our data argue that ILG upregulation is a conserved, coordinated response to changes in trafficking resulting from intrinsic cues (lipid changes) or extrinsic stimulation (during the immune response). These findings uncover important and previously unexplored links between metabolism and the stress response.

Introduction

Cellular stress responses both react to extrinsic insults by upregulating genes that defend the cell. Stress responses also restore organelle function in response to intrinsic changes. Several of these intrinsic stress response pathways are also linked to metabolic processes. For example, overnutrition can lead to metabolic changes that induce cellular stress through an increase in reactive oxygen species (Wellen and Thompson, 2010). Several signaling pathways play a dual role, functioning as coordinators of nutrient responses as well as stress responses. For example, the nutritional sensor mTORC1 can influence stress-responsive transcription (Aramburu et al., 2014). In addition, insulin signaling pathways are strongly linked to stress. For example, mutations in *daf-2*, the *Caenorhabditis elegans* insulin receptor/insulin-like growth factor ortholog, produce highly stress-resistant animals (Son et al., 2019). Finally, changes in ER lipids have a profound effect on stress pathways (Gianfrancesco et al., 2018; Halbleib et al., 2017; Koh et al., 2018). Specialized ER sensors detect both the accumulation of unfolded proteins and overload of the secretory pathway. These sensors then initiate gene expression programs that reduce protein load or increase lipid production (reviewed in: Hotamisligil and Davis, 2016; Radanović et al., 2018).

While stress responses and metabolism are clearly intertwined, the biological advantages of linking these processes are less clear. For example, in a previous study, we found that RNAi-mediated knockdown of two different lipid synthesis modulators, *sams-1* and *sbp-1*, caused upregulation of pathogen-response genes in *C. elegans* (Ding et al., 2015). One of these lipid modulators, *sams-1/MAT1A,* encodes an S-

adenosylmethionine synthase important for phosphatidylcholine (PC) production

(Walker et al., 2011). The Troemel lab also identified *sams-1* as a regulator of the

infection response gene *irg-1* in an RNAi screen (Dunbar et al., 2012). The other lipid

modulator identified in our previous screen, *sbp-1/SREBPF1,* codes for a master

transcriptional regulator of lipid synthesis genes (Horton et al., 2002). Reduction of *sbp-*

1/SREBPF1 causes a decrease in total lipid stores (McKay et al., 2003). Notably,

mammalian SREBPF1 knockdown also results in the enrichment of immunity-linked

gene expression in human cells (Wu and Näär, 2019). Recent studies from other labs

have found that ILGs are upregulated after mutation or RNAi of genes affecting PC

synthesis. ILG activation was seen upon disruption of *pmt-2* (Ho et al., 2020) or *lpin-1* in

the presence of excessive glucose (Jung et al., 2020). These immune-linked genes

were also upregulated in *skn-1* mutants, a transcriptional regulator of the stress

response and metabolic genes (Nhan et al., 2019). In addition, dietary restriction

results in the upregulation of similar gene sets (Wu et al., 2019). These results together

suggest that multiple types of lipid imbalance seem to impact immunity-linked gene

expression and that these links may be conserved across species, as indicated by

results in human cells cited above .

Exposure of *C. elegans* to bacterial pathogens stimulates the expression of a diverse

set of genes, including antimicrobial peptides, enzymes for detoxifying xenobiotics, and

neuromodulatory peptides to coordinate inter-organ defenses (Engelmann et al., 2011;

Fletcher et al., 2019; Mallo et al., 2002; Troemel et al., 2006). However, many of the

genes upregulated in response to either bacteria or fungi do not fall into these clearly

defensive pathways (Estes et al., 2010) or have clear functional roles and are instead

defined by pathogen-responsive expression. In order to understand the relationship

between ILG regulation and changes in lipid levels, we compared lipidomics from *sams-*

1 and *sbp-1* animals with results from a targeted RNAi screen for activation of the

immunity-linked reporter, p*sym-1::GFP* (Shivers et al., 2009). We found that *sams-1*

and sbp-1 knockdown animals both had broad changes in their lipidome. However,

fractionated *sbp-1* extracts showed low PC in the ER/Golgi fraction, as we previously

found in *sams-1* animals (Smulan et al., 2016). Our targeted RNAi screen argues that

synthesis of two lipid classes, PC and sphingomyelin (SM), increase p*sym-1::GFP*

expression. In addition, RNAi of Golgi/ER transport regulators, including the GTPase

arf-1/ARF1, activated the immunity-linked reporter. Notably, some ILGs were also

upregulated after *arf-1*/ARF1 RNAi, suggesting that disrupting Golgi function regulates

immunity-linked genes. Finally, we find that RNAi of one ILG, *hpo-6*, disrupts the

trafficking of two secreted proteins. Taken together, our results suggest that some ILGs

act to balance the increased secretory load in pathogen-stimulated cells. While immune

function is critical for host defenses, immune activation in non-pathogenic conditions,

such as in metabolic disease, may have deleterious effects (Zmora et al., 2017). Thus,

our studies provide critical links between shifts in lipid metabolism and stimulation of

immunity-linked gene expression.

Results

Gene set enrichment shows upregulation of pathogen-response genes in multiple models of membrane dysfunction

Levels of lipids within membranes are tightly monitored, and imbalances may induce cellular stress pathways that regulate genes to restore lipid ratios (Hotamisligil and Davis, 2016). We previously found that *sams-1*, *lpin-1,* and *sbp-1* are part of a feedback loop that responds to shifts in membrane PC levels and activates immunity-linked genes (Fig. 1A; Smulan et al., 2016; Walker et al., 2011). In order to test why immunity-linked genes are activated after disruption of lipid-synthesis regulators we first conducted a bioinformatics analysis using our recently developed *C. elegans* specific gene ontology tool, WormCat 2.0. This computational analysis allows combinatorial graphing of gene enrichment scores based on detailed annotation of all *C. elegans* genes (Higgins et al., 2021; Holdorf et al., 2019). This tool uses a nested annotation strategy (Category1/Category2/Category3) to provide broader or more detailed information on genome-wide datasets. In order to better characterize these ILGs, we used WormCat to directly compare genes upregulated in the *sams-1, sbp-1,* and *lpin-1* mutants (**Fig1A**). The Lee lab generously shared data of genes that are upregulated when *lpin-1* is knocked down in the absence of the stress-triggering glucose (Jung et al., 2020) **Table S1**). WormCat analysis indicates that all three knockdowns result in enrichment in the broad Category1(Cat1) STRESS RESPONSE category, and more specifically in Pathogen at the Cat2 level, similar to results from GO analysis (**Fig1A-C**). Notably, the strongest enrichments at the Cat3 level were in STRESS RESPONSE: Pathogen: *other*. This category contains genes that increase upon pathogen exposure

in *C. elegans* but do not have defined functions as antimicrobial peptides or within

pathogen-stimulated signaling pathways such as MAPK signaling (Higgins et al., 2021;

Holdorf et al., 2019). This class of genes is not explicitly defined by a traditional GO

analysis. We also noted that the landscape of upregulated genes was complex, with

STRESS RESPONSE: Detoxification enrichment scores increasing in both *sams-1* and

lpin-1 but not *sbp-1* (**Fig1B**). In addition, METABOLISM: Lipid was increased after

sams-1 and *lpin-1* RNAi (**Fig1B**). These data are consistent with our previous studies

showing that SBP-1 transcriptional targets are upregulated in these animals (Walker et

al., 2011), as are immunity-linked categories after SREBP-1 knockdown in human

melanoma cell lines (Wu and Näär, 2019). Finally, WormCat analysis of microarray

data from the Thibault lab (Koh et al., 2018) finds that STRESS RESPONSE: Pathogen

is increased after knockdown of *pmt-2*, a methyltransferase essential in SAM-dependent

PC production (**FigS1A**). Thus, genes linked to immunity are upregulated across a

diverse set of lipid synthesis modulators in *C. elegans* and mammalian cells.

SBP-1/SREBP1 is a basic helix-loop-helix transcription factor required to express a

suite of lipid metabolic genes, including *fat-7* in *C. elegans* and its ortholog SCD1 in

mammals (HORTON et al., 2002; McKay et al., 2003). It is also necessary for the

upregulation of *fat-7* in lipid-replete *sams-1(lof)* animals ((Walker et al., 2011), and

Fig1D). Since SBP-1 upregulates lipid synthesis genes, we tested the possibility that it

directly functions in ILG upregulation by using *sams-1(RNAi)* to increase the expression

of multiple SBP-1 target genes (Walker et al., 2011). Whereas *fat-7* depends on *sbp-1*

in both the control and *sams-1(RNAi)* background, *hpo-6* is expressed at high levels in

both *sbp-1(RNAi)* and *sams-1(lof); sbp-1(RNAi)* animals (**Fig1D**). This suggests that

the effects of SBP-1 on ILG upregulation are indirectly related to its function as a

transcriptional regulator.

Many pathogen-responsive genes in *C. elegans* act downstream of signals from the

p38/MAPK14 kinase PMK-1 (Troemel et al., 2006). In previous studies in *C .elegans*,

we found that low PC in *sams-1(lof)* animals leads to constitutive phosphorylation of

PMK-1 and that ILG induction was dependent on both PC and *pmk-1* (Ding et al., 2015).

In order to confirm this effect in mammalian cells, we used siRNA to knock down the

rate-limiting enzyme for PC production, *PCYT1,* in human hepatoma cells (**FigS1B**)*.* We

found that p38/MAPK14 phosphorylation is induced to levels similar to

lipopolysaccharide (LPS). In *C. elegans*, knockdown of *sbp-1* also induces PMK-1

phosphorylation (**FigS1C**). Taken together, the results suggest that upregulation of

ILGs in response to lipid perturbations may share regulatory link to pathogen-stimulated

pathways.

Loss of sbp-1 decreases PC levels in ER/Golgi extracts

Immunity-linked gene expression and PMK-1 phosphorylation both occur after reduction

in *sams-1* and *sbp-1*, although these genes act at different points in lipid synthesis

regulation. Next, we sought to identify any shared signatures in the total lipidomes of

these mutants by LC/MS. Because animals were lysed directly into lipid extraction

buffer, levels were normalized to total lipid as in (Smulan et al., 2016). The entire

lipidome was broadly altered in both cases, with 27% of lipid species showing significant

changes in *sams-1(RNAi)* animals and 30% after *sbp-1(RNAi)* (**Fig2A, B; Table S2**).

We next examined lipid levels at the class and species level. As in our previous studies

(Cell 2011), *sams-1* animals showed lower PC and higher TAG (**Fig2C**), and we also

noted lower phosphatidylserine (PS) and SM levels in comparison to other lipid classes

(**Fig2C, FigS1C**). TAG was lower after *sbp-1* RNAi, as expected from previous studies

(Yang et al., 2006). We also found that while reductions in PC did not reach

significance, PE increased, and Cer levels dropped (**Fig2C, D**). In short, *sbp-1* RNAi

induces broad changes in the lipidome across multiple classes of lipids.

The properties of lipid classes can be altered when distributions of acyl chains change

significantly. We find that there are significant differences in the populations of species

within phospholipids (PC, phosphatidylethanolamine (PE), lyso-

phosphatidylethanolamine (LPE), and phosphatidylglycerol (PG), triglycerides (TAG),

diacylglycerides (DG), and sphingomyelins (SM) after *sams-1(RNAi)*. Changes in

species distribution in lipid classes after *sbp-1* RNAi were limited to TAG, lyso-

phosphatidylcholine (LPC), and ceramides (Cer) in agreement with previous studies

showing lower TAGs in *sbp-1(RNAi)* animals.   We asked if *sams-1* and *sbp-1* RNAi

caused similar shifts in animals and found overlapping changes in 180 shared lipid

species (**TableS2**). 57 of these species were decreased in both, 17 increased in both,

and 102 lipid species changed in opposite directions out of 1405 total lipid species

detected. Lipid species diverged most strongly within the TG class, with increases in

sams-1(RNAi) animals and decreases after *sbp-1* RNAi (**TableS2**). This agreed with

total class levels. For lipid species that increased in both animals, PE species were the

highest, with 15 changing at significant levels. 74 PC species changed in both *sams-1*

and *sbp-1* animals, with 35 decreasing in both (**TableS2**). Finally, three SM species

decreased in both animals. Thus, while *sams-1* or *sbp-1* knockdown each lead to broad

changes in the total lipidome, relatively few overlapping species changed in both

instances. Of those, PC species were the best represented among lipid species that

decreased upon knockdown of either *sams-1* or *sbp-1* in animals.

The total lipidome contains multiple organelles, including the plasma membrane,

mitochondria, nuclear membrane, and Golgi/ER. However, lipid ratios in specific

organelles may be reflective of particular functions; for example, we previously found

that PC ratios in ER/Golgi fractions were associated with activation of SBP-1 in *sams-1*

and *lpin-1* animals (Smulan et al., 2016). Therefore, we performed LC/MS lipidomics on

microsomes from *sbp-1(RNAi)* animals. Because the lipidomics was performed on

fractionated animals, we used total protein for normalization. We found a broader

fraction of the ER/Golgi lipidome was affected after *sbp-1(RNAi)* than in the total lipid

samples. Nearly 40% of lipid species were altered in these fractions (**Fig2E; Table S2**),

with significant changes in species across most classes (**FigS1C**). Similar to the

unfractionated extracts, levels of TAG and Cer decreased in the ER/Golgi (**Fig2F**).

Notably, significant ER/Golgi-specific decreases occurred in PC and DG (**Fig2F, G;**

FigS1D). This demonstrates that *sbp-1(RNAi)* has distinct effects on the ER/Golgi

membranes and shows decreases in PC levels at the sub-compartment level, similar to

sams-1 RNAi.

Targeted RNAi Screen Reveals roles for Ceramide/Sphingomyelin synthesis and COP I transport in immune gene expression

Low PC levels are associated with ILG expression in *sams-1* (Ding et al., 2015) and *pmt-2(RNAi)* animals (Koh et al., 2018). The low PC in the ER/Golgi compartment of both *sams-1* (Smulan et al., 2016) and *sbp-1* RNAi animals suggests this could be a shared signature linked to ILG upregulation. To understand how these lipids might act, we next sought to identify genes important for ILG expression. We screened for upregulation of an immune reporter, p*sysm-1*::GFP (Shivers et al., 2009), with an RNAi sub-library that targets genes involved in complex lipid metabolism, lipid signaling and selected ER/Golgi transport regulators. We previously used a lipid-function RNAi sub-library to identify low-PC regulators of SBP-1/SREBP-1 (Smulan et al., 2016). Notably, we constructed RNAi clones for genes not represented in the Arhinger or Orfeome RNAi libraries. The resulting library includes all predicted phospholipases and enzymes for synthesis of complex lipids (Smulan et al., 2016). For the current screen, we expanded the sub-library to include more genes involved in ceramide/sphingomyelin synthesis and confirmed all clone identities by sequencing. Thus, this library has a broader representation of genes linked to complex lipid synthesis than the commercially available RNAi libraries. p*sysm-1*::GFP is strongly induced by *sbp-1(RNAi)* (**Fig3A**) and has been used by multiple labs as a robust marker for immune gene induction (Jeong et al., 2017; Pender and Horvitz, 2018; Wu et al., 2019).

In order to identify genes in lipid synthesis important for ILG upregualion, we screened this library in quadruplicate in p*sysm-1*::GFP animals and identified a list of 20

candidates for retesting (**Fig3B; Table S3**). Retesting to identify the strongest

candidates occurred in 3 stages: visual rescreening (4X), imaging of candidates from

the rescreen, and qRT-PCR to assess changes in GFP expression. We also included

the *sbp-1* target genes *fat-5*, *fat-6,* and *fat-7* (McKay et al., 2003; Yang et al., 2006) in

the visual retesting. These genes are orthologs of mammalian Stearoyl CoA

desaturases (SCDs) (Watts and Browse, 2002) and can change membrane fluidity

(Bodhicharla et al., 2018; Ruiz et al., 2018) or induce lipid bilayer stress by changing

ratios of saturated/unsaturated acyl chains within lipid classes (Ho et al., 2018, 2020;

Koh et al., 2018).

Candidates from the targeted screen fell into two major groups. First, enzymes linked to

PC and SM synthesis were found (**Fig3C-G; FigS3A, B; TableS3**). This included two

isoforms of the rate-limiting enzyme for PC production, *pcyt-1,* and *sptl-1*, which initiates

the first committed step in sphingolipid synthesis. Thus, our genetic data supports the

notion that changes in PC or SMs are a shared signature linked to ILG activation in

sams-1 and *sbp-1* RNAi animals. Notably, *fat-5, -6,* or *fat-7* RNAi, which would change

acyl chains saturation within each lipid class, had modest effects on the p*sysm-1*::GFP

immune reporter (**Fig3H, FigS3A, B; Table S3**), suggesting changes in membrane

fluidity downstream of these enzymes is not a major contributor to the immunity-linked

gene upregulation.

Second, genes involved in Golgi/ER trafficking activated the reporter in both strains

(**Fig3C, D; Fig43A, B; TableS3**). Golgi/ER trafficking genes included four genes

forming the COPI complex (which also caused developmental delays) and ARF1

guanine activating factor (*agef-1*). The ARF-1 GTPase, a key regulator of these factors,

was a false negative in the original screen, but strongly activated the reporter in the

rescreening. (**Fig3I**; **Table S3**). The coatomer proteins work in a complex (Spang, 2002)

and are required for viability (Fraser et al., 2000). Therefore, we chose one candidate,

copa-1, and performed RNAi with diluted bacteria, allowing animals to develop fully.

We found that the immune activation reporter was strongly expressed in the *copa-1*

knockdown (**Fig3I**). Stimulation of the reporter after the loss of multiple parts of the

COP I machinery and its key regulator *arf-1* strongly argues that ER/Golgi dynamics

signal to induce immunity-linked genes.

The p*sysm-1*::GFP reporter represents a single gene in the immune response program.

To more broadly survey ILGs in our screen validation, we used qRT-PCR with primers

specific to the endogenous *sysm-1* gene, *irg-1, irg-2,* and *hpo-6*, and found robust

activation of each of our candidate reporters after knockdown of *arf-1* or *copa-1*, but

more modest effects with the SM genes or SCDs (**FigS4C-G**). Integration of results

from this targeted RNAi screen with our lipidomics studies shows overlap in two main

areas, PC and SM synthesis, which are themselves connected. We previously linked

changes in PC levels in *sams-1* microsomes to the inactivation of ARF-1, which drove

proteolytic processing of SBP-1/SREBP-1 (Smulan et al., 2016). These results suggest

that blocking ARF-1 function could also be linked to ILG activation.

Disruption of ER/Golgi activates immunity-linked genes

The upregulation of *sysm-1* and other immune genes after *arf-1* RNAi prompted us to

examine genome-wide changes in the transcriptome that could occur in response to the

loss of ARF-1 function. Using RNA-seq, we found that 705 genes were significantly

upregulated after *arf-1* RNAi (**Fig4A, FigS3A; Table S4**) and that 40% of these genes

were also upregulated in *sams-1* or *sbp-1* RNAi animals (**FigS4A**). Downregulated

genes in *sams-1, sbp-1,* or *arf-1* animals had low levels of overlap (**FigS4B**). WormCat

gene set enrichment shows that STRESS RESPONSE: Pathogen: *Other* gene sets are

significantly upregulated in *arf-1(RNAi)* animals as they are after *sbp-1, sams-1,* or *lpin-*

1 RNAi (**Fig4B**). Next, we examined WormCat enrichment within the genes that were

shared between these three RNAi sets and found that shared genes retained strong

enrichment for STRESS RESPONSE: Pathogen (**FigS4C-D**). Thus, *arf-1* GTPase

knockdown causes similar or overlapping effects in immune gene expression to those

seen after broad lipid disruption by *sbp-1* RNAi. Cycling of the ARF1 GTPase requires

interaction with the Golgi membrane (Spang, 2002). In order to study the localization of

ARF-1, we obtained a strain where endogenous ARF-1 is C-terminally tagged by

CRISPR. Next, we asked if ARF-1 localization was altered in *sbp-1(RNAi)* animals by

examining expression in the ARF-1::RFP animals. In intestinal cells, ARF-1::RFP

formed a punctate pattern (**Fig4C**) similar to that of the Golgi ministacks characteristic

of *C. elegans* cells (Broekhuis et al., 2013) (See also **Fig4D**).   Strikingly, ARF-1::RFP

appeared to form larger, more irregular punctae in *sbp-1(RNAi)* animals as well as

having additional diffuse localization, similar to patterns seen with RNAi of the coatomer

protein COPA-1 (**Fig4C**). This suggests that ARF-1 functions at the Golgi are altered in

sbp-1(RNAi) animals.

Loss of human ARF1 or blocking ARF1 cycling with the fungal toxin BrefeldinA disrupts

Golgi integrity (Klausner et al., 1992). In addition, our previous studies found that

lowering PC levels through RNAi of *sams-1* or knockdown of *PCYT1* (the rate-limiting

enzyme for PC production) in mammalian cells blocked ARF1 GTPase activity and

disrupted Golgi structure (Smulan et al., 2016; Walker et al., 2011). Because PC levels

in *sbp-1(RNAi)* ER/Golgi fractions decreased (**Fig2C**) and ARF-1::RFP was

mislocalized, we next examined Golgi structure in *sbp-1(RNAi)* animals using a reporter,

MANS::GFP, driven by an intestinal reporter (Chen et al., 2006). MANS::GFP is a

fusion of *aman-2* (alpha-mannosidase, a conserved Golgi specific protein) and marks

the mini-Golgi stacks characteristic of *C. elegans* (Rolls et al., 2002). Consistent with

previous studies (Ackema et al., 2013; Sato et al., 2006), knockdown of *arf-1*

dramatically shifts the Golgi puncta to diffuse localization across the cytoplasm (**Fig4D**).

copa-1 RNAi also diminished Golgi stacks along with altering ARF-1::RFP localization

(**Fig4C, D**). *sbp-1* knockdown affects Golgi structure, but results in a different pattern

(**Fig4E**). Golgi stacks are more numerous and smaller with increases in diffuse

cytoplasmic localization (**Fig4E**). Notably, *sptl-1(RNAi)* did not noticeably alter Golgi

marked by MANS::GFP and ARF-1::RFP puncta were visible, suggesting that *sptl-1*

knockdown may mediate effects on immune genes through and SM pathways through

different cellular membranes. Taken together, we find that ARF-1 activity (Smulan et

al., 2016) and localization (this study) can be affected by *sams-1* or *sbp-1* RNAi and that

targeting *arf-1* or the coatomer components can activate ILGs. Thus, blocking or

limiting ARF-1 function at the Golgi may be part of the signal to activate these genes.

Immunity-linked gene upregulation in models of trafficking dysfunction

The Golgi apparatus accepts proteins from the ER that are destined for secretion,

processing them by glycosylation before secretory vesicles are loaded (Boncompain

and Weigel, 2018). Stimulation of the innate immune system may significantly impact

trafficking load, as antimicrobial peptides are shuttled through the membrane trafficking

system (Taguchi and Mukai, 2019). Indeed activation of the ER stress response has

been noted in multiple systems when large numbers of proteins need to be produced

and secreted (Gardner et al., 2013). Interestingly, previous data suggest that induction

of membrane stress may also be a sensor of infection (Lamitina and Chevet, 2012).

Based on work by the Ewbank lab, they also suggested that some pathogen-response

genes act to support the trafficking of antimicrobial peptides (AMPs) (Couillault et al.,

2012; Lamitina and Chevet, 2012). We noted that the immunity-linked genes

upregulated at low PC are largely outside of the antimicrobial categories and are

comprised of gene sets defined by their shared expression upon pathogen exposure

rather than function (Estes et al., 2010; Holdorf et al., 2019). We hypothesized that

some of these genes might be responding to direct stress on the trafficking system.

To explore this idea, we turned to two commonly used systems for studying trafficking in

C. elegans, visualization of VIT-2::GFP (Grant and Sato, 2006) and _myo-3_::GFP (Fares

and Greenwald, 2001). _vit-2_ is vitellogenin produced and secreted from intestinal cells

and taken up by the germline (Grant and Hirsh, 1999). _myo-3_::ssGFP has a signal

sequence on the fluorophore and is expressed in body wall muscle (Fares and

Greenwald, 2001) Two published studies examine genome-wide mRNA expression

patterns when *vit-2* is overexpressed or misregulated. First, we used WormCat to

examine category enrichment data from germline-less *glp-1* mutants obtained by the

Blackwell lab, where they showed that *vit-2*::GFP builds up near its production site in

the intestines (Steinbaugh et al., 2015)and found strong enrichment patterns in

STRESS RESPONSE: Pathogen: other (Fig5A). Next, we used WormCat to examine

RNA-seq data from VIT-2::GFP (Singh and Aballay, 2017) overexpressing animals

produced by the Aballay group and also fund increased stress response genes (**Fig5B).**

In order to more directly test the hypothesis that increased protein load could lead to

activation of these ILGs, we compared two strains with a differing copy number of *vit-2*;

RT130, made by microparticle bombardment and containing the GFP array in addition

to wild type copies (Balklava et al., 2016) and BCN9071 which is a CRISPR-generated

allele integrated into the endogenous locus (Rompay et al., 2015) (**Fig4E**). Van

Rompay reported that total yolk protein levels were higher in strains harboring *vit-*

2::GFP derived from RT130 (Rompay et al., 2015). Where we found by RT-PCR that

GFP expression levels were also increased (**Fig4F**). Thus, increasing the amount of

VIT-2::GFP produced was also associated with ILG upregulation.

We found that ILGs become upregulated in the absence of pathogen response when

lipid levels become unbalanced, ARF-1 functions are compromised, or when protein

production is increased. We next tested whether increased ILG expression was part of

a response to help stimulate passage through the secretory pathway when trafficking

load is high. One of the most strongly upregulated genes is *hpo-6*, which is not present

in any of the commercially available RNAi libraries. To study its function more closely in

this context, we made a cDNA construct to allow RNAi in VIT-2::GFP RT130 in wild type

and *sams-1(lof)* animals (Ding et al., 2015). *hpo-6* was originally identified in a screen

for genes whose loss increased sensitivity to a pore-forming toxin (Kao et al., 2011). It

has a glycoprotein domain and may occur in membrane rafts (Rao et al., 2011), but has

no apparent homology to human proteins . *hpo-6(RNAi)* causes a slight increase in

VIT-2::GFP in wild-type animals (**Fig6S**). However, in the low PC *sams-1(lof)*

background, VIT-2::GFP pooling and aggregation suggested a broad effect on

trafficking patterns (**Fig6S)**. Yolk in *C. elegans* consists of both lipids and proteins

(Perez and Lehner, 2019). Because low lipid levels in *sbp-1(RNAi)* animals makes

results with VIT-2::GFP difficult to interpret, we used the *myo-3*::ssGFP reporter model

(Fares and Greenwald, 2001), which expresses and secretes GFP in body wall muscle

cells. We used this reporter to examine the effects of *hpo-6* on secretion when *sbp-1*

function is reduced. First, we confirmed that knockdown of *arf-1* or *sbp-1* affected

secretion of the reporter, finding broad pooling and aggregation after *arf-1(RNAi)* and

aggregation with *sbp-1(RNAi)* (**Fig5H**). As in VIT-2::GFP, *hpo-6(RNAi)* had modest

effects in the wild-type background, with the appearance of small aggregates. (**Fig5H**).

In order to test the effects of blunting *hpo-6* upregulation in sbp-1 animals, we crossed

myo-3::ssGFP into a strain harboring a hypomorphic *sbp-1* allele *sbp-1(ep79)* (Brock et

al., 2006). We found that *hpo-6* knockdown increased the pooling and aggregation in

myo-3::ssGFP; *sbp-1(ep79)* animals (**Fig5I**). This suggests that some of ILGs respond

to changes in secretory function that could occur with lipid imbalance, trafficking load, or

pathogen exposure and support secretory function.

Discussion

Cellular stress arises when essential functions are limited or become unbalanced. Membranous organelles are sensitive to both changes in intrinsic protein activity or alterations of lipids within their bilayers and elicit gene expression programs when stressed. While some stress-responsive genes have clear functions, many genes are defined solely as being responsive to specific stresses (Holdorf et al., 2019; Nadal et al., 2011). It is striking that altering membrane lipids activates a gene expression program shared with pathogen stress in both *C. elegans* (Ding et al., 2015; Ho et al., 2020; Jung et al., 2020) and mammals (Wu and Näär, 2019). Mammalian SREBP-1a has also been implicated as a regulator of innate immune gene expression in macrophages (Im et al., 2011). The secretory pathway is relevant to both membrane stress and immune function about might therefore be regulated by both responses. Membrane organelles such as the ER and Golgi are part of a more extensive trafficking system where proteins and lipids are created, modified, then sent to proper intercellular locations or secreted (Yarwood et al., 2020). Innate immune responses challenge trafficking systems as antimicrobial peptides are produced in mass in the ER, sent in lipid vesicles to the Golgi, where many are glycosylated and then packaged into lipid vesicles for secretion (Bednarska et al., 2017). For example, the *C. elegans* genome contains more than 300 genes that are produced in response to specific pathogens (Engelmann et al., 2011) and share similarity to known antimicrobial effectors (Dierking et al., 2016). In addition, pathogen-responses require an intact ER stress pathway to allow adjustment to the increased trafficking load (Lamitina and Chevet, 2012; Richardson et al., 2010).

The ER contains multiple pathways that respond to changes in proteostasis and lipid

levels (Gardner et al., 2013). These processes are orchestrated by ER-intrinsic proteins

that sense changes in protein folding or alterations in the lipid bilayer and have

important roles in basal conditions, as well as when the secretory system is overloaded

(Hotamisligil and Davis, 2016; Safra et al., 2013). ER-linked events are also important

during pathogen exposure in *C. elegans*; changes in ribosomal function at the ER have

a well-defined impact on pathogen responses (Dunbar et al., 2012; McEwan et al.,

2012) and pathogen responses depend on ER stress pathways (Richardson et al.,

2010; Tillman et al., 2018). Interestingly, the Lee lab also found that ILGs are

upregulated when trafficking is inhibited through blocking glycosylation in the ER (Jeong

et al., 2020). Other organelles in the secretory pathway, such as the Golgi, have their

own stress sensors and response pathways (Machamer, 2015). Our results suggest

that disruption of trafficking outside the ER by targeting ARF-1 in the Golgi can also

initiate a stress response stimulating ILG induction.

Genes upregulated by pathogen exposure include both antimicrobial effectors and

many genes of unknown function (Estes et al., 2010; Troemel et al., 2006). Notably, the

ILGs enriched after membrane stress are primarily those of unknown function. Several

lines of evidence prompted us to hypothesize that some of these genes might be part of

a response linked to stress in membranes of the secretory pathway rather than a direct

response to an extrinsic pathogen. First, the involvement of secretory organs was

implicated by alterations in PC ratios in the ER/Golgi in both *sams-1* and *sbp-1* RNAi

lipidomes. Second, our targeted RNAi screen for regulators of p*sysm-1::GFP* identified

the *arf-1* GTPase and coatomer proteins, which are critical Golgi/ER transport.

Importantly, knockdown of *arf-1* was sufficient to induce ILG upregulation.

Strengthening this link between disruption of lipid levels and ARF-1 function, we also

found *sbp-1* RNAi altered ARF-1::RFP localization and Golgi morphology.

ARF-1 is a critical regulator of trafficking, coordinating retrograde traffic from the Golgi

to the ER and regulating secretory function in the trans-Golgi (Donaldson and Jackson,

2011). Cycling of ARF-1 GTPases depends on membrane localization of the ARF

GTPase, GEF, and GAP (GTPase activating protein) (Donaldson and Jackson, 2011).

We previously found that knockdown of PC synthesis enzymes blocked ARF1 cycling

and limited membrane association of the ARF GEF GBF1 in cultured human cells

(Smulan et al., 2016). This suggests low PC levels affect ARF1 activity by limiting the

ability of the GTPase, GEF, and GAP to associate at the Golgi membrane and initiate

GDP-GTP cycling. However, PC could be linked to ARF1 through other mechanisms.

ARF1 and phospholipase D (PLD) function have been linked by multiple labs (Riebeling

et al., 2009). PLD cleaves PC molecules to produce choline and phosphatidic acid,

which in turn stimulates vesicle formation (McDermott et al., 2004). This regulatory loop

requires PC, which is low in ER/Golgi membranes in *sams-1* and *sbp-1* RNAi animals.

DAG kinases, which could be stimulated by changes in DAGs in the *sbp-1* or *sams-*

1(RNAi) animals, can also affect ARF activity and trafficking (Huijbregts et al., 2000).

Finally, Protein Kinase D has an important role in maintaining Golgi structure, and its

ortholog has been linked to immunity in *C. elegans* (Malhotra and Campelo, 2011; Ren

et al., 2009). However, none of these components activated the p*sysm-1::GFP* reporter

in our RNAi screen, suggesting they could be important in other regulatory contexts.

Indeed, the breadth of change in the lipidomes after *sams-1* or *sbp-1* RNAi suggests

that other inputs could link lipids and ILG expression. In addition, other cellular

processes may also affect ILG expression either in basal conditions or during pathogen

infection. For example, there is a complex relationship between pathogen infection,

proteotoxic stress, and induction of ILGs (Reddy et al., 2017)

In conclusion, our data suggest that the ILG induction after both broad disruption in

membrane lipids and pathogen exposure occurs because both processes stress the

trafficking system. We also find that one the upregulated immune-linked genes, *hpo-6*,

can facilitate trafficking function. Our results suggest ILG induction may represent a

"multi-membrane" stress response encompassing the roles of both the ER and the

Golgi in trafficking. Many mammalian studies on the effect of lipids on innate immune

functions have focused on specific lipids as stimulators of signaling pathways (Barnett

and Kagan, 2019). Taken together, these results suggest that levels of specific

membrane lipids can initiate stress responses sensitive to trafficking defects, which

could impact immune function as well as other secretion-dependent processes. Cells

contain complex systems for managing stress in the initial organelle of the secretory

process, the ER, which may respond to disturbances in protein synthesis, aggregation,

or lipid imbalances with specific programs of gene expression (Costa-Mattioli and

Walter, 2020; Hotamisligil and Davis, 2016). Thus, the expression of ILGs may reflect a

broader stress response that affects multiple secretory organelles, as processes beyond

translation and folding are affected.

Experimental Model and Subject Details

C. elegans **strains, RNAi constructs, and screening**.

N2 (wild type), p*sysm-1::GFP,* and OP50 were obtained from the *Caenorhabditis*

Genetics Center. CRISPR-tagged ARF-1, p*arf-1::arf-1::RFP,* was obtained from *In vivo*

Biosystems and outcrossed 3 times. Normal growth media (NGM) was used unless

otherwise noted. For RNAi, gravid adults were bleached onto NGM plates

supplemented with ampicillin, tetracycline, and 6mM IPTG and 10X concentrated

bacterial cultures. *C. elegans* were allowed to develop until the young adult stage

before harvesting. Because of larval arrest, *copa-1* RNAi bacteria were diluted 1:10 in

control RNAi bacteria before plating. For the RNAi screen, L1 larvae were plated into

96 well plates spotted with RNAi bacteria, and L4/young adults were scored from -3 to

+3 with 0 as no change in 4 independent replicates.

Cell culture and siRNA transfection

HepG2 cells (ATCC, HB-8065) were grown in Minimum Essential Medium (Invitrogen)

plus 10% FBS (Invitrogen), Glutamine (Invitrogen), and Sodium Pyruvate (Invitrogen).

siRNA oligonucleotides transfections were done for 48 hours using Lipofectamine

RNAiMAX Transfection Reagent (Invitrogen, 13778100), and cells were held for 16

hours in 1% Lipoprotein Deficient Serum (LDS) (Biomedical Technologies, BT907) and

25 µg/ml ALLN (Calbiochem) for 30 min prior to harvesting.

Immunoblotting:

HepG2 cells: Syringe passage was used to lyse cells in High Salt RIPA (50 mM Tris,

pH 7.4; 400 mM NaCl; 0.1% SDS; 0.5% NaDeoxycholate; 1% NP-40; 1 mM DTT, 2.5

µg/mL ALLN, Complete protease inhibitors (Roche). Invitrogen NuPage gels (4- 12%)

were used for protein separation before transfer to nitrocellulose. Blots were probed

with antibodies to phosphorylated p38 MAP Kinase, total p38, and STAT1 were used as

a control. Immune complexes were visualized with Luminol Reagent (Millipore).

Densitometry was performed by scanning the film, then analysis of pixel intensity with

ImageJ software. Graphs show the average of at least three independent experiments

with control values normalized to one.

C. elegans: Young adult animals were lysed by sonication in High Salt RIPA, and

immunoblotting was performed as above.

Microscopy

P*sysm*-1::GFP *C. elegans* strains were grown until the L4/young adult transition, and

images were acquired on a Zeiss Dissecting Scope with an Axiocam camera. ARF-

1::RFP and MANS-1::GFP images were acquired on a Leica SPE confocal, and

projections of confocal slices were produced. All images were taken at identical gain

settings within experimental sets, and Adobe Photoshop was used for corrections to

levels across experimental sets.

Lipidomics

C. elegans total and microsomal lipidomics, including fractionation protocols, were

performed at the Whitehead Metabolomics core as in Smulan et al. 2016. Statistical

analysis was performed in Graphpad prism.

Gene expression analysis

Lysis of young adult *C. elegans* was performed in 0.5% SDS, 5% β-ME, 10 mM EDTA,

10 mM Tris-HCl ph7.4, 0.5 mg/ml Proteinase K, before purification of RNA by TRI-

Reagent (Sigma). cDNA was produced with Transcriptor First-strand cDNA kits

(Roche), and RT-PCR was performed using Kappa SYBR Green 2X Mastermix.

RNA for sequencing was purified using RNAeasy columns (Qiagen). RNA sequencing

(including library construction was performed by BGI (Hong Kong). Reads were

analyzed through the Dolphin analysis platform (https://dolphin.umassmed.edu/), using

DeSeq to identify differentially expressed genes (Yukselen et al., 2020). Gene set

enrichment was performed using WormCat (www.wormcat.com) (Holdorf et al., 2019).

552
553

Acknowledgments

We would like acknowledge Drs. Seung-Jae Lee, Sujeong Kwon (Korea Advanced Institute of Science and Technology) for providing the *lpin-1* RNA seq data. We would also like to thank Drs. Seung-Jae Lee, Emily Troemel (USD) and members of the Walker lab for comments on the manuscript as well as Dr. Marian Walhout (UMASS Chan School of Medicine) for helpful discussion. We would like to acknowledge Wei Ding for technical support and Caroline Lewis (MIT) for metabolomics screening at the Whitehead Metabolomics core. We also thank Life Science Editors and Dr. Sabbi Lall for manuscript editing. Funding was from the NIA at the NIH: R01AG053355 and R56AG068670. Some strains were provided by the CGC, which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440).

Author Contributions

Conceptualization; AKW, CMW, LJS

Verification: AKW, CMW, LJS

Formal Analysis: AKW, CMW, LJS

Investigation: AKW, CMW, LJS, DL, MF

Resources: AKW

Data Curation: AKW

Writing: AKW

Writing/Editing: AKW, CMW, LJS,

Visualization: AKW

586   Supervision: AKW
587
588   Funding: AKW
589

Declaration of Interests

The authors declare no competing interests

Figure Legends

Figure 1: Upregulation of immunity-linked genes occurs downstream of multiple

types of membrane lipid disruption. WormCat gene set enrichment shows STRESS

RESPONSE (**A**) and STRESS RESPONSE: Pathogen (**B**) are upregulated in *lpin-1*

RNAi animals, as in *sams-1* and *sbp-1(RNAi)* animals (Ding et al., 2015). Legend is in

C. qRT-PCR shows that upregulation of an immunity-linked gene, *hpo-6*, in *sams-1(lof)*

animals is not dependent on *sbp-1* (**D**).

Figure 2: Broad lipidomic changes in total and microsomal lipid compartments

after *sams-1* or *sbp-1* RNAi. LC/MS lipidomics show that nearly a third of total lipid

species change after *sams-1* or *sbp-1* RNAi (**A, B**). Significant changes in major lipid

classes are mostly distinct in *sams-1* and *sbp-1* RNAi (**C, D**). See also **Figure S1** (**B,**

D) for lipid species distributions and additional lipid class analysis. The microsomal

(ER/Golgi) lipidome is also broadly altered after *sbp-1 RNAi* (**E**) and shows changes in

different lipid classes than in the total lipidome (**F, G**). Error bars show standard

deviation. Lipid class data is calculated by students T-Test. See Figure S1 for lipid

species distributions and additional lipid class analysis.

Figure 3: Targeted RNAi screen reveals candidates linked to Golgi/ER trafficking

and ceramide/sphingomyelin synthesis impact immunity gene expression. (**A**)

Immune gene reporter p*sysm-1*::GFP is increased after *sbp-1* RNAi. (**B**) Screen

schematic. (**C**) Schematic showing RNAi library coverage and screen candidates.

Epifluorescence imaging showing validation of reporter activity after RNAi of lipid PC

synthesis genes (**D**), SM synthesis genes (**E**) and the SCD *fat-7* (**F**). Epifluorescence

images showing GFP levels after RNAi of the *arf-1* GTPase or coatomer component

copa-1 (**I**). Because of poor larval development, bacterial for *copa-1* RNAi was diluted

to 1:10. **J**. Epifluorescence images showing GFP levels genes linked to Golgi function.

For qRT-PCR, see **Figure S4.**

Figure 4: Disruption of Golgi/ER trafficking increases expression of immunity-

related genes. A. Scatter plot shows changes in gene expression after *arf-1* RNAi. **B.**

WormCat gene set enrichment chart shows an increase in stress response and

pathogen linked genes. **C.** Confocal projections of CRISPR tagged ARF-1::RFP reveal

that that Golgi-specific pattern is disrupted after *sbp-1* RNAi. Scale bar is 10 microns.

D. Confocal projections of the *C. elegans* Golgi marker MANS::GFP (Chen et al., 2006)

in Control, *sbp-1, arf-1,* and *copa-1 (1:10)* RNAi intestines. Scale bar is 10 microns.

Figure 5: Immunity-linked genes are linked to trafficking disruption. WormCat

gene enrichment of upregulated genes from *glp-1*(AA2735) (Steinbaugh et al., 2015)

and *vit-2(ac3)* compared to SJ4005 (Singh and Aballay, 2017)for Category 1 (**A**) and

Category 2 (**B**). Legend is in **C**. qRT-PCR comparing gene expression in wild type,

single copy *vit-2::GFP* (BCN9071, s) or multi copy (RT130, m) animals. **D** and **E** show

increase in GFP or *vit-2*. Immune response genes are shown in **F-I**. **J**. Confocal

images of body wall muscle cells expressing of the secretion marker *myo-3*::ssGFP. **K**.

Body wall muscle cells in imaged by confocal microscopy in *myo-3*::ssGFP; *sbp-1(ep79)*

animals.

Supplemental information

Figure S1: Activation of Stress response pathways when PC synthesis is disrupted. A. WormCat analysis of microarray data from Koh, et al. 2018. **B.** The p38 MAP kinase MAPK14 is phosphorylated after siRNA of *PCYT1* in human hepatoma cells. **C.** Constitutive phosphorylation of the *C. elegans* MAPK14 ortholog, PMK-1, occurs after both *sams-1* and *sbp-1(RNAi)*.

Figure S2: Broad lipidomic changes in total and microsomal lipid compartments after *sams-1* or *sbp-1* RNAi. **A**. Schematic showing effects of PC producing pathways on ARF-1. Tables showing numbers lipid species within each class in the total lipidome (**B**) or ER/Golgi lipidome in *sbp-1(RNAi)* animals (**D**). Green boxes denote classes in which the distribution of the species within each RNAi significantly differed from controls using the Wilcoxen test for non-parametric data. Levels of lipid classes in total *sams-1* or *sbp-1* total lipidomics (**C**) or *sbp-1* ER/Golgi lipidomics (**E**).

Figure S3: Candidates from lipid sub-library RNAi screen. Table showing candidates from quadruplicate screening of the RNAi sub-library along with other genes selected as controls for the rescreen.

Figure S4: Regulation of immunity-linked genes in conditions of membrane disruption. qRT PCR quantitating GFP upregulation after knockdown of Golgi trafficking regulators **(A)** or lipid synthesis genes **(B)**. qRT-PCR from *psysm-1*::GFP (A-

H) or p*sysm-1*::GFP; *sbp-1(ep79)* (I-K) shows effects of disrupting *arf-1* or membrane

lipids on ILGs. Error bars show standard deviation.

Figure S5: Shared upregulation of immunity linked genes in *sams-1, sbp-1* and

***arf-1* transcriptomic studies.** Venn diagrams showing limited overlap of up- (A) or

downregulated (**B**) genes after *sbp-1, sams-1* or *arf-1* RNAi. **C-D.** WormCat analysis

comparing overlapping gene sets from *arf-1, sbp-1* and *sams-1* upregulated genes.

Data for *arf-1* alone is the same as **Figure 4B**. Legend is in **E**. Confocal micrographs

showing minimal changes after *sptl-1* RNAi in of MANS::GFP (**F**) or ARF-1::RFP (**G**).

Bar show 10 microns.

Figure S6: Reduction in *hpo-6* exacerbates *sams-1(lof)* trafficking defects.

Confocal projections of *C. elegans* intestinal cells expressing *vit-2*::GFP (RT130) in wild

type or *sams-1(lof)* backgrounds.

Table S1: Transcriptomics and gene set enrichment from *lpin-1(RNAi)* animals. 2

fold upregulated genes in *lpin-1(RNAi)* animals. Gene set enrichment was performed

using WormCat (Holdorf et al., 2019).

Table S2: **Total lipidomics of *sams-1* and *sbp-1(RNAi)* animals and microsomal**

lipidomics of *sbp-1(RNAi)* adults. Datasets from two lipidomic studies. The first is

LCMS data normalized to total lipids compared between Control, *sbp-1,* and *sams-*

1(RNAi) adults. Tabs are labeled sams_sbp_Lipid class. The second dataset contains

LCMS from Control and *sbp-1(RNAi)* microsome preps from adult animals. Significance

of fold changes in individual lipids determined by students T-test.

Table S3: Results from the screen of the "lipid sub-library." Tabs 1 and 2 contain

averaged results from quadruplicate screening of *sysm-1*::GFP or *sysm-1::GFP; sbp-*

1(ep79). The secondary screen contains averages from candidates selected from the

first screen, plus *fat-5, fat-6,* and *fat-7* as additional test RNAis. The secondary screen

was conducted in quadruplicate. ND is not done.

Table S4: Transcriptomic responses to trafficking disruption. RNAseq from control vs.

arf-1/arf-1.2(RNAi) animals (arf_1_all, arf_2Xup) and gene set enrichment analysis

produced in WormCat for Category 1, Category2 and Category3). WormCat analysis of

vit-2::GFP (Singh and Aballay, 2017)and *glp-1* (Steinbaugh et al., 2015) datasets.

References

Ackema, K.B., Sauder, U., Solinger, J.A., and Spang, A. (2013). The ArfGEF GBF-1 Is Required for ER Structure, Secretion and Endocytic Transport in *C. elegans*. Plos One *8*, e67076.

Aramburu, J., Ortells, M.C., Tejedor, S., Buxadé, M., and López-Rodríguez, C. (2014). Transcriptional regulation of the stress response by mTOR. Sci Signal *7*, re2–re2.

Balklava, Z., Rathnakumar, N.D., Vashist, S., Schweinsberg, P.J., and Grant, B.D. (2016). Linking Gene Expression in the Intestine to Production of Gametes Through the Phosphate Transporter PITR-1 in *Caenorhabditis elegans*. Genetics *204*, 153 162.

Barnett, K.C., and Kagan, J.C. (2019). Lipids that directly regulate innate immune signal transduction. Innate Immun *26*, 4–14.

Bednarska, N.G., Wren, B.W., and Willcocks, S.J. (2017). The importance of the glycosylation of antimicrobial peptides: natural and synthetic approaches. Drug Discov Today *22*, 919–926.

Bodhicharla, R., Devkota, R., Ruiz, M., and Pilon, M. (2018). Membrane Fluidity Is Regulated Cell Non-autonomously by *Caenorhabditis elegans* PAQR-2 and Its Mammalian Homolog AdipoR2. Genetics *210*, 189 201.

Boncompain, G., and Weigel, A.V. (2018). Transport and sorting in the Golgi complex: multiple mechanisms sort diverse cargo. Curr Opin Cell Biol *50*, 94–101.

Brock, T.J., Browse, J., and Watts, J.L. (2006). Genetic Regulation of Unsaturated Fatty Acid Composition in *C. elegans*. Plos Genet *2*, e108.

Broekhuis, J.R., Rademakers, S., Burghoorn, J., and Jansen, G. (2013). SQL-1, homologue of the Golgi protein GMAP210, modulates intraflagellar transport in *C. elegans*. J Cell Sci *126*, 1785–1795.

Chen, C.C.-H., Schweinsberg, P.J., Vashist, S., Mareiniss, D.P., Lambie, E.J., and Grant, B.D. (2006). RAB-10 Is Required for Endocytic Recycling in the *Caenorhabditis elegans* Intestine. Mol Biol Cell *17*, 1286 1297.

Costa-Mattioli, M., and Walter, P. (2020). The integrated stress response: From mechanism to disease. Science *368*, eaat5314.

Couillault, C., Fourquet, P., Pophillat, M., and Ewbank, J.J. (2012). A UPR-independent infection-specific role for a BiP/GRP78 protein in the control of antimicrobial peptide expression in *C. elegans* epidermis. Virulence *3*, 299 308.

Dierking, K., Yang, W., and Schulenburg, H. (2016). Antimicrobial effectors in the
nematode *Caenorhabditis elegans*: an outgroup to the Arthropoda. Philosophical
Transactions Royal Soc B Biological Sci *371*, 20150299.

Ding, W., Smulan, L.J., Hou, N.S., Taubert, S., Watts, J.L., and Walker, A.K. (2015). s-
Adenosylmethionine Levels Govern Innate Immunity through Distinct Methylation-
Dependent Pathways. Cell Metab *22*, 633 645.

Donaldson, J.G., and Jackson, C.L. (2011). ARF family G proteins and their regulators:
roles in membrane transport, development and disease. Nat Rev Mol Cell Bio *12*, 362
375.

Dunbar, T.L., Yan, Z., Balla, K.M., Smelkinson, M.G., and Troemel, E.R. (2012). *C.
elegans* detects pathogen-induced translational inhibition to activate immune signaling.
Cell Host Microbe *11*, 375 386.

Engelmann, I., Griffon, A., Tichit, L., Montañana-Sanchis, F., Wang, G., Reinke, V.,
Waterston, R.H., Hillier, L.W., and Ewbank, J.J. (2011). A Comprehensive Analysis of
Gene Expression Changes Provoked by Bacterial and Fungal Infection in *C. elegans*.
Plos One *6*, e19055.

Estes, K.A., Dunbar, T.L., Powell, J.R., Ausubel, F.M., and Troemel, E.R. (2010). bZIP
transcription factor *zip-2* mediates an early response to Pseudomonas aeruginosa
infection in *Caenorhabditis elegans*. Proc National Acad Sci *107*, 2153 2158.

Fares, H., and Greenwald, I. (2001). Genetic Analysis of Endocytosis in *Caenorhabditis
elegans* : Coelomocyte Uptake Defective Mutants. Genetics *159*, 133–145.

Fletcher, M., Tillman, E.J., Butty, V.L., Levine, S.S., and Kim, D.H. (2019). Global
transcriptional regulation of innate immunity by ATF-7 in *C. elegans*. Plos Genet *15*,
e1007830.

Fraser, A.G., Kamath, R.S., Zipperlen, P., Martinez-Campos, M., Sohrmann, M., and
Ahringer, J. (2000). Functional genomic analysis of *C. elegans* chromosome I by
systematic RNA interference. Nature *408*, 325–330.

Gardner, B.M., Pincus, D., Gotthardt, K., Gallagher, C.M., and Walter, P. (2013).
Endoplasmic Reticulum Stress Sensing in the Unfolded Protein Response. Csh
Perspect Biol *5*, a013169.

Gianfrancesco, M.A., Paquot, N., Piette, J., and Legrand-Poels, S. (2018). Lipid bilayer
stress in obesity-linked inflammatory and metabolic disorders. Biochem Pharmacol *153*,
168–183.

Grant, B., and Hirsh, D. (1999). Receptor-mediated endocytosis in the *Caenorhabditis
elegans* oocyte. Mol Biol Cell *10*, 4311 4326.

Grant, B.D., and Sato, M. (2006). Intracellular trafficking. Wormbook Online Rev C
Elegans Biology 1–9.

Halbleib, K., Pesek, K., Covino, R., Hofbauer, H.F., Wunnicke, D., Hänelt, I., Hummer,
G., and Ernst, R. (2017). Activation of the Unfolded Protein Response by Lipid Bilayer
Stress. Mol Cell *67*, 673 684.e8.

Higgins, D.P., Weisman, C.M., Lui, D.S., D'Agostino, F.A., and Walker, A.K. (2021).
WormCat 2.0 defines characteristics and conservation of poorly annotated genes in
Caenorhabditis elegans. Biorxiv. https://doi.org/10.1101/2021.11.11.467968

Ho, N., Xu, C., and Thibault, G. (2018). From the unfolded protein response to
metabolic diseases – lipids under the spotlight. J Cell Sci *131*, jcs199307.

Ho, N., Yap, W.S., Xu, J., Wu, H., Koh, J.H., Goh, W.W.B., George, B., Chong, S.C.,
Taubert, S., and Thibault, G. (2020). Stress sensor Ire1 deploys a divergent
transcriptional program in response to lipid bilayer stress. J Cell Biol *219*.

Holdorf, A.D., Higgins, D.P., Hart, A.C., Boag, P.R., Pazour, G.J., Walhout, A.J.M., and
Walker, A.K. (2019). WormCat: An Online Tool for Annotation and Visualization of
Caenorhabditis elegans Genome-Scale Data. Genetics genetics.302919.2019.

HORTON, J.D., GOLDSTEIN, J.L., and BROWN, M.S. (2002). SREBPs: Transcriptional
Mediators of Lipid Homeostasis. Cold Spring Harb Sym *67*, 491–498.

Hotamisligil, G.S., and Davis, R.J. (2016). Cell Signaling and Stress Responses. Csh
Perspect Biol *8*, a006072.

Huijbregts, R.P.H., Topalof, L., and Bankaitis, V.A. (2000). Lipid Metabolism and
Regulation of Membrane Trafficking. Traffic *1*, 195–202.

Im, S.-S., Yousef, L., Blaschitz, C., Liu, J.Z., Edwards, R.A., Young, S.G., Raffatellu, M.,
and Osborne, T.F. (2011). Linking Lipid Metabolism to the Innate Immune Response in
Macrophages through Sterol Regulatory Element Binding Protein-1a. Cell Metab *13*,
540 549.

Jeong, D., Lee, D., Hwang, S., Lee, Y., Lee, J., Seo, M., Hwang, W., Seo, K., Hwang,
A.B., Artan, M., et al. (2017). Mitochondrial chaperone HSP-60 regulates anti-bacterial
immunity via p38 MAP kinase signaling. Embo J *36*, 1046–1065.

Jeong, D.-E., Lee, Y., Ham, S., Lee, D., Kwon, S., Park, H.-E.H., Hwang, S.-Y., Yoo, J.-
Y., Roh, T.-Y., and Lee, S.-J.V. (2020). Inhibition of the oligosaccharyl transferase in
Caenorhabditis elegans that compromises ER proteostasis suppresses p38-dependent
protection against pathogenic bacteria. Plos Genet *16*, e1008617.

Jung, Y., Kwon, S., Ham, S., Lee, D., Park, H.H., Yamaoka, Y., Jeong, D., Artan, M.,
Altintas, O., Park, S., et al. (2020). *Caenorhabditis elegans* Lipin 1 moderates the
lifespan-shortening effects of dietary glucose by maintaining ω-6 polyunsaturated fatty
acids. Aging Cell *19*.

Kao, C.-Y., Los, F.C.O., Huffman, D.L., Wachi, S., Kloft, N., Husmann, M., Karabrahimi,
V., Schwartz, J.-L., Bellier, A., Ha, C., et al. (2011). Global Functional Analyses of
Cellular Responses to Pore-Forming Toxins. Plos Pathog *7*, e1001314.

Klausner, R.D., Donaldson, J.G., and Lippincott-Schwartz, J. (1992). Brefeldin A:
insights into the control of membrane traffic and organelle structure. The Journal of Cell
Biology *116*, 1071 1080.

Koh, J.H., Wang, L., Beaudoin-Chabot, C., and Thibault, G. (2018). Lipid bilayer stress-
activated IRE-1 modulates autophagy during endoplasmic reticulum stress. J Cell Sci
131, jcs217992.

Lamitina, T., and Chevet, E. (2012). To UPR… and beyond!: A new role for a
BiP/GRP78 protein in the control of antimicrobial peptide expression in *C. elegans*
epidermis. Virulence *3*, 238–240.

Machamer, C.E. (2015). The Golgi complex in stress and death. Front Neurosci-Switz *9*,
421.

Malhotra, V., and Campelo, F. (2011). PKD Regulates Membrane Fission to Generate
TGN to Cell Surface Transport Carriers. Csh Perspect Biol *3*, a005280 a005280.

Mallo, G.V., Kurz, C.L., Couillault, C., Pujol, N., Granjeaud, S., Kohara, Y., and Ewbank,
J.J. (2002). Inducible Antibacterial Defense System in *C. elegans*. Curr Biol *12*, 1209–
1214.

McDermott, M., Wakelam, M.J.O., and Morris, A.J. (2004). Phospholipase D. Biochem
Cell Biol *82*, 225–253.

McEwan, D.L., Kirienko, N.V., and Ausubel, F.M. (2012). Host translational inhibition by
Pseudomonas aeruginosa Exotoxin A Triggers an immune response in *Caenorhabditis*
elegans. Cell Host Microbe *11*, 364 374.

McKay, R.M., McKay, J.P., Avery, L., and Graff, J.M. (2003). *C. elegans* A Model for
Exploring the Genetics of Fat Storage. Dev Cell *4*, 131–142.

Nadal, E. de, Ammerer, G., and Posas, F. (2011). Controlling gene expression in
response to stress. Nat Rev Genet *12*, 833–845.

Nhan, J.D., Turner, C.D., Anderson, S.M., Yen, C.-A., Dalton, H.M., Cheesman, H.K.,
Ruter, D.L., Naresh, N.U., Haynes, C.M., Soukas, A.A., et al. (2019). Redirection of

SKN-1 abates the negative metabolic outcomes of a perceived pathogen infection. Proc
National Acad Sci *116*, 22322–22330.

Pender, C.L., and Horvitz, H.R. (2018). Hypoxia-inducible factor cell non-autonomously
regulates *C. elegans* stress responses and behavior via a nuclear receptor. Elife *7*,
e36828.

Perez, M.F., and Lehner, B. (2019). Vitellogenins - Yolk Gene Function and Regulation
in *Caenorhabditis elegans*. Front Physiol *10*, 1067.

Radanović, T., Reinhard, J., Ballweg, S., Pesek, K., and Ernst, R. (2018). An Emerging
Group of Membrane Property Sensors Controls the Physical State of Organellar
Membranes to Maintain Their Identity. Bioessays *40*, 1700250.

Rao, W., Isaac, R.E., and Keen, J.N. (2011). An analysis of the *Caenorhabditis elegans*
lipid raft proteome using geLC-MS/MS. J Proteomics *74*, 242 253.

Reddy, K.C., Dror, T., Sowa, J.N., Panek, J., Chen, K., Lim, E.S., Wang, D., and
Troemel, E.R. (2017). An Intracellular Pathogen Response Pathway Promotes
Proteostasis in C. elegans. Curr Biol *27*.

Ren, M., Feng, H., Fu, Y., Land, M., and Rubin, C.S. (2009). Protein Kinase D Is an
Essential Regulator of C. elegans Innate Immunity. Immunity *30*, 521–532.

Richardson, C.E., Kooistra, T., and Kim, D.H. (2010). An essential role for XBP-1 in host
protection against immune activation in *C. elegans*. Nature *463*, 1092 1095.

Riebeling, C., Morris, A.J., and Shields, D. (2009). Phospholipase D in the Golgi
apparatus. Biochimica Et Biophysica Acta Bba - Mol Cell Biology Lipids *1791*, 876 880.

Rolls, M.M., Hall, D.H., Victor, M., Stelzer, E.H.K., and Rapoport, T.A. (2002). Targeting
of Rough Endoplasmic Reticulum Membrane Proteins and Ribosomes in Invertebrate
Neurons. Mol Biol Cell *13*, 1778–1791.

Rompay, L.V., Borghgraef, C., Beets, I., Caers, J., and Temmerman, L. (2015). New
genetic regulators question relevance of abundant yolk protein production in *C. elegans*.
Sci Rep-Uk *5*, 1 16.

Ruiz, M., Bodhicharla, R., Svensk, E., Devkota, R., Busayavalasa, K., Palmgren, H.,
Ståhlman, M., Boren, J., and Pilon, M. (2018). Membrane fluidity is regulated by the C.
elegans transmembrane protein FLD-1 and its human homologs TLCD1/2. Elife *7*,
1309.

Safra, M., Ben-Hamo, S., Kenyon, C., and Henis-Korenblit, S. (2013). The *ire-1* ER
stress-response pathway is required for normal secretory-protein metabolism in *C.*
elegans. J Cell Sci *126*, 4136–4146.

Sato, K., Sato, M., Audhya, A., Oegema, K., Schweinsberg, P., and Grant, B.D. (2006).
Dynamic Regulation of Caveolin-1 Trafficking in the Germ Line and Embryo of
Caenorhabditis elegans. Mol Biol Cell *17*, 3085 3094.

Shivers, R.P., Kooistra, T., Chu, S.W., Pagano, D.J., and Kim, D.H. (2009). Tissue-
Specific Activities of an Immune Signaling Module Regulate Physiological Responses to
Pathogenic and Nutritional Bacteria in *C. elegans*. Cell Host Microbe *6*, 321–330.

Singh, J., and Aballay, A. (2017). Endoplasmic Reticulum Stress Caused by Lipoprotein
Accumulation Suppresses Immunity against Bacterial Pathogens and Contributes to
Immunosenescence. Mbio *8*, e00778-17.

Smulan, L.J., Ding, W., Freinkman, E., Gujja, S., Edwards, Y.J.K., and Walker, A.K.
(2016). Cholesterol-Independent SREBP-1 Maturation Is Linked to ARF1 Inactivation.
Cell Reports *16*, 9 18.

Son, H.G., Altintas, O., Kim, E.J.E., Kwon, S., and Lee, S.V. (2019). Age-dependent
changes and biomarkers of aging in *Caenorhabditis elegans*. Aging Cell *18*, e12853.

Spang, A. (2002). ARF1 regulatory factors and COPI vesicle formation. Curr Opin Cell
Biol *14*, 423–427.

Steinbaugh, M.J., Narasimhan, S.D., Robida-Stubbs, S., Mazzeo, L.E.M., Dreyfuss,
J.M., Hourihan, J.M., Raghavan, P., Operaña, T.N., Esmaillie, R., and Blackwell, T.K.
(2015). Lipid-mediated regulation of SKN-1/Nrf in response to germ cell absence. Elife
4, 466.

Taguchi, T., and Mukai, K. (2019). Innate immunity signalling and membrane trafficking.
Curr Opin Cell Biol *59*, 1–7.

Tillman, E.J., Richardson, C.E., Cattie, D.J., Reddy, K.C., Lehrbach, N.J., Droste, R.,
Ruvkun, G., and Kim, D.H. (2018). Endoplasmic Reticulum Homeostasis Is Modulated
by the Forkhead Transcription Factor FKH-9 During Infection of *Caenorhabditis
elegans*. Genetics *210*, genetics.301450.2018.

Troemel, E.R., Chu, S.W., Reinke, V., Lee, S.S., Ausubel, F.M., and Kim, D.H. (2006).
p38 MAPK regulates expression of immune response genes and contributes to
longevity in *C. elegans*. Plos Genet *2*, e183.

Walker, A.K., Jacobs, R.L., Watts, J.L., Rottiers, V., Jiang, K., Finnegan, D.M., Shioda,
T., Hansen, M., Yang, F., Niebergall, L.J., et al. (2011). A Conserved SREBP-
1/Phosphatidylcholine Feedback Circuit Regulates Lipogenesis in Metazoans. Cell *147*,
840 852.

Watts, J.L., and Browse, J. (2002). Genetic dissection of polyunsaturated fatty acid
synthesis in *Caenorhabditis elegans*. Proc National Acad Sci *99*, 5854 5859.

Wellen, K.E., and Thompson, C.B. (2010). Cellular Metabolic Stress: Considering How
Cells Respond to Nutrient Excess. Mol Cell *40*, 323 332.

Wu, S., and Näär, A.M. (2019). SREBP1-dependent de novo fatty acid synthesis gene
expression is elevated in malignant melanoma and represents a cellular survival trait.
Sci Rep-Uk *9*, 10369.

Wu, Z., Isik, M., Moroz, N., Steinbaugh, M.J., Zhang, P., and Blackwell, T.K. (2019).
Dietary Restriction Extends Lifespan through Metabolic Regulation of Innate Immunity.
Cell Metab *29*, 1192-1205.e8.

Yang, F., Vought, B.W., Satterlee, J.S., Walker, A.K., Sun, Z.-Y.J., Watts, J.L.,
DeBeaumont, R., Saito, R.M., Hyberts, S.G., Yang, S., et al. (2006). An ARC/Mediator
subunit required for SREBP control of cholesterol and lipid homeostasis. Nature *442*,
700 704.

Yarwood, R., Hellicar, J., Woodman, P.G., and Lowe, M. (2020). Membrane trafficking
in health and disease. Dis Model Mech *13*, dmm043448.

Yukselen, O., Turkyilmaz, O., Ozturk, A.R., Garber, M., and Kucukural, A. (2020).
DolphinNext: a distributed data processing platform for high throughput genomics. Bmc
Genomics *21*, 310.

Zmora, N., Bashiardes, S., Levy, M., and Elinav, E. (2017). The Role of the Immune
System in Metabolic Health and Disease. Cell Metab *25*, 506–521.

A

WormCat Category 1 GSEA

B

WormCat Category 2/3 GSEA

C

D

A *sysm-1*::GFP

Control

sbp-1 (RNAi)

B Complex Lipid RNAi sublibrary 2.0

230 genes

sysm-1::GFP

C Screen Candidates

Phospho-lipids

Trigly-cerides

Ceramides Sphingomyelins

Lipid Signaling

Selected COPI/ COPII, Trafficking Components

Selected *sams-1* or *sbp-1* upregulated genes

pcyt-1
pyct-2.2
ckb-3

sptl-1
sms-3

nape-1

arf-1   *copa-1*   *sar-1*
agef-1 *copb-1*   *syx-1*
copz-1 *copg-1*
F32D8.4 (Yip1p)

pmp-5

D Control   *sbp-1*   *pcyt-1*   *pcyt-2*   *pcyt-2.2*

sysm-1::GFP

E Control   *sbp-1*   *ckb-3*

sysm-1::GFP

F Control   *sbp-1*   *sptl-1*

sysm-1::GFP

G Control   *sbp-1*   *sms-3*

sysm-1::GFP

H Control   *sbp-1*   *fat-6*

sysm-1::GFP

I Control   *sbp-1*   *arf-1*   *copa-1 (d)*

sysm-1::GFP

J Control   *sbp-1*   *F32D8.4*   *syx-1*

sysm-1::GFP

A

arf-1(RNAi)

Mean Log$_{10}$ counts (y-axis)

Mean Log$_{10}$ counts (x-axis)

B

CATEGORY 1

arf-1

Category 2

STRESS RESPONSE
Pathogen
Other
UNKNOWN
PROT PROTEASOME
PROT GENERAL

10^{-40}
10^{-20}
10^{-10}
10^{-5}
0.001
0.05
NS

C

ARF-1::RFP

Control

sbp-1

copa-1

D

MANS::GFP(Golgi)

Control   *arf-1*

sbp-1   *copa-1*

A

WormCat Category 1 GSEA

glp-1 *vit-2*

STRESS RESPONSE
ECM
PROT GENERAL
METABOLISM
TM TRANSPORT
UNKNOWN
MUSCLE FUN
NEURONAL FUN
CYTOSKELETON

B

WormCat Category 2 GSEA

glp-1 *vit-2*

STRESS RESPONSE
Pathogen
Detoxification
GST
C-type Lectin

C

10^{-40}
10^{-20}
10^{-10}
10^{-5}
0.001
0.05
NS

D GFP

E *vit-2*

F *hpo-6*

G *irg-1*

H *irg-2*

I T24C4.4

J *myo-3*::ssGFP

Control   *arf-1(RNAi)*
sbp-1(RNAi)   *hpo6—1(RNAi)*

K *myo-3*::ssGFP; *sbp-1(ep79)*

Control
arf-1
hpo-6