Stimulation of immunity-linked genes by membrane disruption is linked to Golgi function

and the ARF-1 GTPase

Matthew J. Fanelli[1], Christofer M. Welsh[1,2], Dominique S. Lui[1], Lorissa J. Smulan[3] and *Amy

K. Walker[1]

[1]Program in Molecular Medicine, UMASS Chan Medical School, Worcester, MA

[2]Morningside School of Biomedical Sciences, UMASS Chan Medical School, Worcester, MA

[3]Department of Medicine, UMASS Chan Medical School, Worcester, MA

Corresponding author: amy.walker@umassmed.edu

Abstract

Immunity-linked genes (ILGs) are activated by pathogens but also may respond to imbalances in

lipids. Why pathogen attack and metabolic changes both impact ILG activation is unclear. We

find that ILGs are activated when membrane phosphatidylcholine ratios change in secretory

organelles in *C. elegans*. RNAi targeting of the ADP-ribosylation factor ARF-1, which disrupts

the Golgi, also activates ILG expression, suggesting that activation of this membrane stress

response could occur outside the ER. Our data argue that ILG upregulation is a coordinated

response to changes in trafficking resulting from intrinsic cues (changes in membrane lipids) or

extrinsic stimulation (increased secretion during immune response). Indeed, a focused RNAi

screen of ILGs uncovered defects in secretion of two GFP reporters as well as accumulation of a

pathogen-responsive CUB-domain fusion protein. These results also suggests that genes shared

between the classical pathogen responses and lipid stress may act to counteract stress on

secretory function.

Teaser: Pathogen response genes are also activated by lipid imbalances, which we suggest

occurs because both processes put stress on the secretory pathway.

Introduction

Introduction

Cellular stress responses react to extrinsic insults by upregulating genes that defend the cell.

Stress responses also restore organelle function in response to intrinsic changes. Several of these

intrinsic stress response pathways are also linked to metabolic processes. For example,

overnutrition can lead to metabolic changes that induce cellular stress through an increase in

reactive oxygen species (*1*). Several signaling pathways play a dual role, functioning as

coordinators of nutrient responses as well as stress responses. For example, the nutritional

sensor mTORC1 can influence stress-responsive transcription (*2*). In addition, insulin signaling

pathways are strongly linked to stress. In addition, mutations in *daf-2*, the *Caenorhabditis*

elegans insulin receptor/insulin-like growth factor ortholog, produce highly stress-resistant

animals (*3*). Finally, changes in ER lipids have a profound effect on stress pathways (*4–6*).

Specialized ER sensors detect both the accumulation of unfolded proteins and overload of the

secretory pathway. These sensors then initiate gene expression programs that reduce protein

load or increase lipid production (*7, 8*).

While stress responses and metabolism are clearly intertwined, the biological advantages of

linking these processes are less clear. For example, in a previous study, we found that RNAi-

mediated knockdown of two different lipid synthesis modulators, *sams-1* and *sbp-1*, caused

upregulation of pathogen-response genes in *C. elegans* (*9*). One of these lipid modulators,

sams-1/MAT1A, encodes an S-adenosylmethionine synthase important for phosphatidylcholine

(PC) production (*10*). The Troemel lab also identified *sams-1* as a regulator of the infection

response gene *irg-1* in an RNAi screen (*11*). The other lipid modulator identified in our previous

screen, *sbp-1/SREBPF1,* codes for a master transcriptional regulator of lipid synthesis genes

(*12*). Reduction of *sbp-1/SREBPF1* causes a decrease in total lipid stores (*13*). Notably,

mammalian *SREBPF1* knockdown also results in the enrichment of immunity-linked gene

expression in human cells (*14*). Recent studies from other labs have found that ILGs are

upregulated after mutation or RNAi of genes affecting PC synthesis. ILG activation was seen

upon disruption of *pmt-2* (*15*) or *lpin-1* in the presence of excessive glucose (*16*). These

immunity-linked genes were also upregulated in *skn-1* mutants, a transcriptional regulator of the

stress response and metabolic genes (*17*). In addition, dietary restriction results in the

upregulation of similar gene sets (*18*). These results together suggest that multiple types of lipid

imbalance seem to impact immunity-linked gene expression and that these links may be

conserved across species, as indicated by results in human cells cited above.

Exposure of *C. elegans* to bacterial pathogens stimulates the expression of a diverse set of

genes, including antimicrobial peptides, enzymes for detoxifying xenobiotics, and

neuromodulatory peptides to coordinate inter-organ defenses (*19–22*). However, many of the

genes upregulated in response to either bacteria or fungi do not fall into these clearly defensive

pathways (*23*) or have clear functional roles. Instead, these genes are defined by pathogen-

responsive expression. In order to understand why knockdown of genes acting in lipid

metabolism might activate genes linked to immune responses when no pathogens were present,

we took two approaches. First, we looked for lipid signatures that could signal this response and

were shared between *sams-1* and *sbp-1* by comparing lipidomics of whole animals and

fractioned extracts. Second, we performed a RNAi screen of genes functioning in complex lipid

synthesis and ER/Golgi dynamics for activation of the immunity-linked reporter, p*sysm-1::GFP*

(*24*). In the lipidomics studies, we found that *sams-1 and sbp-1* knockdown animals had broad

and distinct changes in their lipidome, but both showed lowering of PC levels in the ER/Golgi

fraction which we previously linked to SBP-1/SREBP1 maturation in *sams-1* animals (*25*).

Strengthening the links to PC, our targeted RNAi screen showed that synthesis of PC, increased

p*sysm-1::GFP* expression. RNAi of Golgi/ER transport regulators, including the GTPase *arf-*

1/ARF1, activated the immunity-linked reporter. Notably, some ILGs were also upregulated

after *arf-1*/ARF1 RNAi, suggesting that disrupting Golgi function regulates immunity-linked

genes. Finally, we performed a targeted screen, and found that interfering with ILGs expression

in *sbp-1(ep79)* animals disrupted trafficking of a secreted reporter. Taken together, our results

suggest that some ILGs act to balance the increased secretory load in pathogen-stimulated cells

or when membrane lipid balance is perturbed. While immune function is critical for host

defenses, immune activation in non-pathogenic conditions, such as in metabolic disease, may

have deleterious effects (*26*). Therefore, it is critical to define cellular roles for pathogen-

responsive genes. Our studies suggest that genes upregulated in response to both pathogens and

lipid perturbation in *C. elegans* could act to support secretory processes altered in both

biological contexts.

Results

Gene set enrichment shows upregulation of pathogen-response genes in multiple models of membrane dysfunction

Levels of lipids within membranes are tightly monitored, and imbalances may induce cellular stress pathways that regulate genes to restore lipid ratios (*8*). We previously found that *sams-1*, *lpin-1,* and *sbp-1* are part of a feedback loop that responds to shifts in membrane PC levels and activates immunity-linked genes (**Fig. 1A**)(*9*, *25*). In order to test why immunity-linked genes are activated after disruption of lipid-synthesis regulators we first conducted a bioinformatics analysis using our recently developed *C. elegans* specific gene ontology tool, WormCat 2.0. This computational analysis allows combinatorial graphing of gene enrichment scores based on detailed annotation of all *C. elegans* genes (*27*, *28*) and uses a nested annotation strategy (Category1/Category2/Category3) to provide broader or more detailed information on genome-wide datasets. The WormCat annotation strategy also includes a specialized category, Unassigned, to include genes cannot be classified using our strict criteria. Genes that are generally characterized are placed in UNASSIGNED at the Category 1 level in WormCat 2.0 ("other" in the 1.0 version), but we also use unassigned to denote genes within larger, well described category whose specific roles were less clear. For example, in WormCat, genes with roles in immunity are classified at the Cat1 and Cat2 level in STRESS RESPONSE: Pathogen. This category is broken down into Cat3 classifications for transcriptional or signaling regulators, neuropeptide-like proteins, proteins containing common antimicrobial motifs (saposin, caenacin, CUB, and others).

Figure 1: Upregulation of immunity-linked genes occurs downstream of multiple types of

membrane lipid disruption. A. Schematic diagram of lipid metabolic genes linked to immune

gene expression. Meta-analysis of published RNA seq data from linking lipid perturbation to

immune gene expression using WormCat 2.0 for Category 1 (Cat1) (**B**) and for the Category 2

(Cat2) level under STRESS RESPONSE (**C**).

Legend is in **D.**

This category also includes 97 genes upregulated during the pathogen response, but lacking

molecular characterization, which were designated STRESS RESPONSE: Pathogen: *unassigned*

(*27*, *28*). This sub-category had the highest enrichment after RNAi of *sbp-1* or *sams-1* (*27–29*),

suggesting that immunity linked genes upregulated after lipid unbalance could function in area

common to immune response and lipid stress.

In order to better characterize these ILGs, we used WormCat to perform a meta-analysis of

published data comparing categories enriched in genes upregulated by the pathogen PA14 (*22*)

with multiple models of disrupted lipogenesis (**Fig1A**). We found enrichment in

Category1(Cat1) STRESS RESPONSE, and more specifically in Pathogen at the Cat2 level in

genes upregulated after the lipogenic transcriptional regulators *sbp-1* (*9*), *mdt-15* (*30*) or *nhr-49*

(*31*) or enzymes acting in lipid synthesis, *sams-1* (*9*), *lpin-1* ((*16*); **DataS1**), *pmt-2* (*6*), and *hyl-2*

(*32*) were knocked down as well as when exogenous lipids such as P80 (oleic acid mimic) (*33*)

or cholesterol were added (*34*) (**Fig1B-C**). Notably, the strongest enrichments at the Cat3 level

were in STRESS RESPONSE: Pathogen: *unassigned*. Genes in this category are defined by

solely upregulation upon pathogen exposure in *C. elegans* but do not have defined functions as

antimicrobial peptides or within pathogen-stimulated signaling pathways such as MAPK

signaling (*27*, *35*). Importantly, this class of genes is not explicitly defined by a traditional GO

analysis. Genes upregulated in *pmt-2(RNAi)* animals were also linked to the Lipid Bilayer Stress

induced Unfolded Protein Response (UPR[LBS]), a UPR[ER] mechanism (*6*), therefore, we compared

WormCat enrichment of genes increasing after tunicamycin treatment (*36*). We found that while

STRESS RESPONSE: Pathogen: *unassigned* genes were enriched, the category enrichment was

independent of the ER stress regulators *xbp-1* or *ire-1 (***FigS1A)**, suggesting potential for a

distinct regulatory mechanism.

We also noted that the landscape of upregulated genes was complex across other categories, with

STRESS RESPONSE: Detoxification enrichment scores increasing in both *sams-1* and *lpin-1* but

not *sbp-1* (**Fig1C**). In addition, METABOLISM: Lipid was increased after *lpin-1* RNAi

(**Fig1C**). These data are consistent with our previous studies showing that SBP-1 transcriptional

targets are upregulated in these animals (*10*). Interestingly, immunity-linked categories after

SREBP-1 knockdown in human melanoma cell lines (*14*). Thus, genes linked to immunity are

upregulated across a diverse set of lipid synthesis modulators in *C. elegans* and mammalian cells

and occur in the absence of pathogen stimulation.

SBP-1/SREBP1 is a basic helix-loop-helix transcription factor required to express a suite of lipid

metabolic genes, including *fat-7* in *C. elegans* and its ortholog SCD1 in mammals (*12*, *13*). It is

also necessary for the upregulation of *fat-7* in lipid-replete *sams-1(lof)* animals ((*10*) (**FigS1B**).

Since SBP-1 upregulates lipid synthesis genes, we tested the possibility that it directly functions

in ILG upregulation by using *sams-1(RNAi)* to increase the expression of multiple SBP-1 target

genes (*10*) and assessing the need for *sbp-1*. As we previously reported in Walker, 2011, *fat-7*

depends on *sbp-1* in both the control and *sams-1(RNAi)* background. However, the ILG *hpo-6* is

expressed at high levels when induced by reduction in *sbp-1*, *sams-1*, or in *sams-1(lof); sbp-*

1(RNAi) animals (**FigS1C**), suggesting that the effects of SBP-1 on ILG upregulation are

indirectly related to its function as a transcriptional regulator.

Many ILGs in *C. elegans* act downstream of signals from the p38/MAPK14 kinase PMK-1 (*22*).

In previous studies in *C .elegans*, we found that low PC in *sams-1(lof)* animals leads to

constitutive phosphorylation of PMK-1 and that ILG induction was dependent on both PC and

pmk-1 (*9*). We confirmed that knockdown of *sbp-1* also induces PMK-1 phosphorylation

(**FigS1D**). Since ILGs can be induced in mammalian cells in response to reduction in *SREBPF1*,

we used siRNA to knock down the rate-limiting enzyme for PC production, *PCYT1,* in human

hepatoma cells (**FigS1E**). We found that p38/MAPK14 phosphorylation is induced to levels

similar to lipopolysaccharide (LPS). Taken together, the results suggest that upregulation of

ILGs in response to lipid perturbations may share regulatory link to pathogen-stimulated

pathways.

Loss of sbp-1 decreases PC levels in ER/Golgi extracts

Immunity-linked gene expression and PMK-1 phosphorylation both occur after reduction in

sams-1 and *sbp-1*, although these genes act at different points in lipid synthesis regulation. In

order to identify shared lipid signatures that might stimulate this process,   we analyzed the total

lipidomes of these mutants by LC/MS. Because animals were lysed directly into lipid extraction

buffer, levels were normalized to total lipid as in (*25*). The entire lipidome was broadly altered

in both cases, with 27% of lipid species showing significant changes in *sams-1(RNAi)* animals

and 30% after *sbp-1(RNAi)* (**Fig2A, B; DataS2**). We next examined lipid levels at the class and

species level. As in our previous studies (*10*), *sams-1* animals showed lower PC and higher TAG

(Triacylglyceride) (**Fig2C**), and we also noted lower phosphatidylserine (PS) and sphingomyelin

(SM) levels in comparison to other lipid classes (**Fig2C, FigS2C**). TAG was lower after *sbp-1*

RNAi, as expected from previous studies (*37*). We also found that while reductions in PC did

not reach significance, PE (Phosphatidylethanolamine) increased, and Cer (Ceramide) levels

dropped (**Fig2C, D**). In short, *sbp-1* RNAi induces broad changes in the lipidome across

multiple classes of lipids.

The properties of lipid classes can be altered when distributions of acyl chains alter membrane

properties (*38*). We find that there are significant differences in the populations of species within

phospholipids (PC, PE, lyso-phosphatidylethanolamine (LPE), and phosphatidylglycerol (PG),

TAG, diacylglycerides (DG), and SMs after *sams-1(RNAi)*. Changes in species distribution in

lipid classes after *sbp-1* RNAi were limited to TAG, lyso-phosphatidylcholine (LPC), and Cer in

agreement with previous studies showing lower TAGs in *sbp-1(RNAi)* animals. We asked if

sams-1 and *sbp-1* RNAi caused similar shifts in animals and found overlapping changes in 180

shared lipid species (**DataS2**). Fifty-seven of these species were decreased in both, 17 increased

in both, and 102 lipid species changed in opposite directions out of 1405 total lipid species

detected. Lipid species diverged most strongly within the TG class, with increases in *sams-*

1(RNAi) animals and decreases after *sbp-1* RNAi (**DataS2**). This agreed with total class levels.

For lipid species that increased in both animals, PE species were the highest, with 15 changing at

significant levels. Seventy-four PC species changed in both *sams-1* and *sbp-1* knockdown

animals, with 35 decreasing in both (**TableS3**). Finally, three SM species decreased in both

animals. Thus, while *sams-1* or *sbp-1* knockdown each lead to broad changes in the total

lipidome, relatively few overlapping species changed in both instances. Of those, PC species

were the best represented among lipid species that decreased upon knockdown of either *sams-1*

or *sbp-1* in animals.

Figure 2: Broad lipidomic changes in total and microsomal lipid compartments after *sams-1 or sbp-1* RNAi. LC/MS lipidomics show that nearly a third of total lipid species change after *sams-1* or *sbp-1* RNAi (**A, B**). Significant changes in major lipid classes are mostly distinct in *sams-1* and *sbp-1* RNAi (**C, D**). See also **Figure S1** (**B, D**) for lipid species distributions and additional lipid class analysis and **Data S2** for all values. The microsomal (ER/Golgi) lipidome is also broadly altered after *sbp-1 RNAi* (**E**) and shows changes in different lipid classes than in the total lipidome (**F, G**). Error bars show standard deviation. Lipid class data is calculated by students T-Test. See Figure S1 for lipid species distributions and additional lipid class analysis.

The total lipidome contains multiple organelles, including the plasma membrane, mitochondria,

nuclear membrane, and Golgi/ER. However, lipid ratios in specific organelles may be reflective

of particular functions; for example, we previously found that PC ratios in ER/Golgi fractions

were associated with activation of SBP-1 in *sams-1* and *lpin-1* animals (*25*). Therefore, we

performed LC/MS lipidomics on microsomes from *sbp-1(RNAi)* animals to assess lipids in the

ER/Golgi fraction. Because the lipidomics was performed on fractionated animals, we used total

protein for normalization. We found a broader fraction of the ER/Golgi lipidome was affected

after *sbp-1(RNAi)* than in the total lipid samples. Nearly 40% of lipid species were altered in

these fractions (**Fig2E; Data23**), with significant changes in species across most classes

(**FigS2C**). Similar to the unfractionated extracts, levels of TAG and Cer decreased in the

ER/Golgi (**Fig2F**). Notably, significant ER/Golgi-specific decreases occurred in PC and DG

(**Fig2F; FigS2D**). This demonstrates that *sbp-1(RNAi)* has distinct effects on the ER/Golgi

membranes and shows decreases in PC levels at the sub-compartment level, similar to *sams-1*

RNAi.

Targeted RNAi Screen Reveals roles for COP I transport in immune gene expression

Low PC levels are associated with ILG expression in *sams-1* (*9*) and *pmt-2(RNAi)* animals (*6*).

The low PC in the ER/Golgi compartment of both *sams-1* (*25*) and *sbp-1* RNAi animals suggests

this could be a shared signature linked to ILG upregulation. To understand how these lipids

might act, we next sought to identify genes important for ILG expression. We screened for

upregulation of an immune reporter, p*sysm-1*::GFP (*24*), with an RNAi sub-library that targets

genes involved in complex lipid metabolism, lipid signaling and selected ER/Golgi transport

regulators. We previously used a lipid-function RNAi sub-library to identify low-PC regulators

of SBP-1/SREBP-1 (*25*). Notably, we constructed RNAi clones for genes not represented in the

Arhinger or Orfeome RNAi libraries. The resulting library includes all predicted phospholipases

and enzymes for synthesis of complex lipids (*25*). For the current screen, we expanded the sub-

library to include more genes involved in ceramide/sphingomyelin synthesis and confirmed all

clone identities by sequencing. Thus, this library has a broader representation of genes linked to

complex lipid synthesis than the commercially available RNAi libraries. p*sym-1*::GFP is

strongly induced by *sbp-1(RNAi)* (**Fig3A**) and has been used by multiple labs as a robust marker

for immune gene induction (*18*, *39*, *40*).

In order to identify genes in lipid synthesis important for ILG upregulation, we screened this

library in quadruplicate in p*sym-1*::GFP animals and identified a list of 20 candidates for

retesting (**Fig3B; DataS3**). Retesting to identify the strongest candidates occurred in 3 stages:

visual rescreening (4X), imaging of candidates from the rescreen, and qRT-PCR to assess

changes in GFP expression. We also included the *sbp-1* target genes *fat-5*, *fat-6,* and *fat-7* (*13*,

37) in the visual retesting. These genes are orthologs of mammalian Stearoyl CoA desaturases

(SCDs) (*41*) and can change membrane fluidity (*42*, *43*) or induce lipid bilayer stress by

changing ratios of saturated/unsaturated acyl chains within lipid classes (*6*, *15*, *44*).

Candidates from the targeted screen fell into two major groups. First, enzymes linked to PC and

SM synthesis were found (**Fig3C-G; FigS3A, B; DataS3**). This included two isoforms of the

rate-limiting enzyme for PC production, *pcyt-1,* and *sptl-1*, which initiates the first committed

step in sphingolipid synthesis. Thus, our genetic data supports the notion that changes in PC or

SMs are a shared signature linked to ILG activation in *sams-1* and *sbp-1* RNAi animals.

A

sysm-1::GFP

Control

sbp-1 (RNAi)

B

Complex Lipid RNAi sublibrary 2.0

230 genes

sysm-1::GFP

C

Screen Candidates

Phospho-lipids

pcyt-1
pyct-2.2
ckb-3

Trigly-cerides

Ceramides
Sphingomyeins

sptl-1
sms-3

Lipid Signaling

nape-1

Selected COPI/COPII, Trafficking Components

arf-1   *copa-1*   *sar-1*
agef-1 *copb-1*   *syx-1*
copz-1 *copg-1*

Selected *sams-1* or *sbp-1* upregulated genes

pmp-5

D

Control | *sbp-1* | *pcyt-1* | *pcyt-2* | *pcyt-2.2*

sysm-1::GFP

E

Control | *sbp-1* | *ckb-3*

sysm-1::GFP

F

Control | *sbp-1* | *sptl-1*

sysm-1::GFP

G

Control | *sbp-1* | *sms-3*

sysm-1::GFP

H

Control | *sbp-1* | *fat-6*

sysm-1::GFP

I

Control | *sbp-1* | *arf-1* | *copa-1 (d)*

sysm-1::GFP

J

Control | *sbp-1* | *syx-5*

sysm-1::GFP

Figure 3: Targeted RNAi screen reveals candidates linked to Golgi/ER trafficking and

ceramide/sphingomyelin synthesis impact immunity gene expression. (**A**) Immune gene

reporter p*sysm-1*::GFP is increased after *sbp-1* RNAi. (**B**) Screen schematic. (**C**) Schematic

showing RNAi library coverage and screen candidates. Epifluorescence imaging showing

validation of reporter activity after RNAi of lipid PC synthesis genes (**D, E**), SM synthesis genes

(**F, G**) and the stearoyl CoA desaturase *fat6/7* (**H**). Epifluorescence images showing GFP levels

after RNAi of the *arf-1* GTPase or coatomer component *copa-1* or proteins acting at the Golgi

such as *syx-1*/Syntaxin5 (**I, J**). Because of poor larval development, bacteria for *copa-1* RNAi

was diluted to 1:10. Internal control for effect of *sbp-1(RNAi)* included in each panel and DIC

image included as an inset for each RNAi. Quantitation of images is in **Figure S4, A, B**.

Notably, *fat-5, fat-6,* or *fat-7* RNAi, which would change acyl chain saturation within each lipid

class, had modest effects on the p*sysm-1*::GFP immune reporter (**Fig3H, FigS3A, B; DataS3**),

suggesting changes in membrane fluidity downstream of these enzymes is not a major

contributor to the immunity-linked gene upregulation.

Second, Golgi/ER trafficking genes included four genes forming the COPI complex (which also

caused developmental delays) and ARF1 guanine activating factor (*agef-1*). The ARF-1

GTPase, a key regulator of these factors, was a false negative in the original screen, but strongly

activated the reporter in the rescreening. (**Fig3I**; **DataS3**). The coatomer proteins work in a

complex (*45*) and are required for viability (*46*). Therefore, we chose one candidate, *copa-1,* and

performed RNAi with diluted bacteria, allowing animals to develop fully. We found that the

immune activation reporter was strongly expressed in the *copa-1* knockdown (**Fig3I**).

Stimulation of the reporter after the loss of multiple parts of the COP I machinery and its key

regulator *arf-1* strongly argues that ER/Golgi dynamics signal to induce immunity-linked genes.

The p*sysm-1*::GFP reporter represents a single gene in the immune response program. To more

broadly survey ILGs in our screen validation, we used qRT-PCR with primers specific to *irg-1,*

irg-2, and *hpo-6*, and found robust activation of each of our candidate reporters after knockdown

of *arf-1* or *copa-1*, but more modest effects with the SM genes or SCDs (**FigS4C-H),** implicating

ARF1 and Golgi function in ILG activation.

Disruption of ER/Golgi activates immunity-linked genes

Cycling of the ARF1 GTPase requires interaction with the Golgi membrane (*45*). Alteration of

membrane lipid ratios can affect interactions with ARF1 regulatory proteins, limiting GTPase

cycling and blocking ARF1 activity (*25*, *47*). We previously linked changes in PC levels in

sams-1 microsomes to the inactivation of ARF-1, which drove proteolytic processing of SBP-

1/SREBP-1 (*25*). Therefore, we reasoned that multiple modes of membrane lipid disruption

could signal secretory stress through ARF-1 and activate ILGs. The upregulation of *sysm-1* and

other immune genes after *arf-1* RNAi prompted us to examine genome-wide changes in the

transcriptome that could occur in response to the loss of ARF-1 function. Using RNA-seq, we

found that 705 genes were significantly upregulated after *arf-1* RNAi (**Fig4A, DataS4**) and that

40% of genes were also upregulated in *sams-1* or *sbp-1* RNAi animals (**Fig4B**). Downregulated

genes in *sams-1, sbp-1,* or *arf-1* animals had low levels of overlap (**Fig4C**). WormCat gene set

enrichment shows that STRESS RESPONSE: Pathogen: *Other* gene sets are significantly

upregulated in *arf-1(RNAi)* animals as they are after *sbp-1, sams-1,* or *lpin-1* RNAi (**Fig4D-F**).

Next, we examined WormCat enrichment within the genes that were shared between these three

RNAi sets and found that shared genes retained strong enrichment for STRESS RESPONSE:

Pathogen (**Fig4D-F**). Thus, *arf-1* GTPase knockdown causes similar or overlapping effects in

immune gene expression to those seen after broad lipid disruption by *sbp-1* RNAi.

In order to study the localization of ARF-1, we obtained a strain where endogenous ARF-1 is C-

terminally tagged with mCherry by CRISPR (WAL10). Next, we asked if ARF-1 localization

was altered in *sbp-1(RNAi)* animals by examining expression in the ARF-1::mCherry animals.

In intestinal cells, ARF-1::mCherry formed a punctate pattern (**Fig4G-H**) characteristic of the

Golgi ministacks occurring in *C. elegans* cells (*48*) and which localize adjacent to a Golgi

marker, MANS::GFP (*49*), expressed in the intestine. MANS::GFP is a fusion of *aman-2* (alpha-

mannosidase, a conserved Golgi specific protein) and marks the mini-Golgi stacks characteristic

Figure 4: Disruption of Golgi/ER trafficking increases expression of immunity-related

genes. A. Scatter plot shows changes in gene expression after *arf-1* RNAi. Venn diagrams

showing limited overlap of up- (**B**) or downregulated (**C**) genes after *sbp-1, sams-1* or *arf-1*

RNAi. WormCat gene set enrichment chart shows an increase in stress response and pathogen

linked genes in Category1 (**D**) and Category2 levels (**E**). Legend is in **F. G.** Confocal

projections of endogenously tagged ARF-1::mCherry reveal that that Golgi-like pattern is

disrupted after *sbp-1* RNAi and quantitation in **Figure S5A**. **H**. Confocal images of intestinal

cells from ARF-1::mCherry; ges-1::MANS-1::GFP animals. Scale bar is 10 microns. **I**.

Confocal projections of the *C. elegans* Golgi marker MANS::GFP (*49*) in Control, *sbp-1, arf-1,*

and *copa-1 (1:10)* RNAi intestines with quantitation of puncta intensity and number in **Figure**

S5B, C. Scale bar is 10 microns. Significance is determined by the Mann-Whitney test.

346 of *C. elegans* (*50*). This suggests that, as in other eukaryotes, *C. elegans* ARF-1 acts at the

347 Golgi apparatus. Strikingly, ARF-1::mCherry appeared to form larger, more irregular punctae in

348 *sbp-1(RNAi)* animals as well as having additional diffuse localization, similar to patterns seen

349 with RNAi of the coatomer protein COPA-1 (**Fig4C**). This suggests that ARF-1 functions at the

350 Golgi are altered in *sbp-1(RNAi)* animals.

352 Loss of human ARF1 or blocking ARF1 cycling with the fungal toxin Brefeldin A disrupts Golgi

353 integrity (*51*). In addition, our previous studies found that lowering PC levels through RNAi of

354 *sams-1* or knockdown of *PCYT1* (the rate-limiting enzyme for PC production) in mammalian

355 cells blocked ARF1 GTPase activity and disrupted Golgi structure (*10*, *25*). Because PC levels

356 in *sbp-1(RNAi)* ER/Golgi fractions decreased (**Fig2C**) and ARF-1::mCherry was mis-localized,

357 we next examined Golgi structure after *sbp-1(RNAi)* in animals where MANS::GFP was driven

358 by an intestinal reporter (*49*). Consistent with previous studies (*52*, *53*), knockdown of *arf-1*

359 dramatically shifts the Golgi puncta to diffuse localization across the cytoplasm (**Fig4I; Fig**

360 **S5A**). *copa-1* RNAi also diminished Golgi stacks along with altering ARF-1::mCherry

361 localization. *sbp-1* knockdown affects Golgi structure, but results in a different pattern, in which

362 Golgi stacks are smaller and more numerous with increases in diffuse cytoplasmic localization

363 (**Fig4I, Fig5A**). We also examined MANS::GFP patterns in other candidates from our screen

364 and noted that like *arf-1*, *copa-1* and *sar-1* were required for MANS-1::GFP puncta (**FigS5B, C**),

365 as expected for core constituents of COP I and COP II transport. Contributors to PC synthesis

366 also broadly abrogated MANS::GFP puncta, however, change in fatty acid desaturation, which

367 induces UPR[LBS] (*6*), caused a slight increase in MANS::GFP puncta intensity, suggesting that

368 membrane stress initiated by Golgi misfunction has separable defects. Our screen candidates

also included genes from ceramide synthesis, which can also affect membrane function (*54*).

Notably, *sptl-1(RNAi)* did not noticeably alter Golgi marked by MANS::GFP and ARF-

1::mCherry puncta were visible (**FigS5A, D**), suggesting that *sptl-1* knockdown may mediate

effects on immune genes through distinct cellular membranes. Taken together, we find that

ARF-1 activity (*25*) and localization (this study) can be affected by *sams-1* or *sbp-1* RNAi and

that targeting *arf-1* or the coatomer components can activate ILGs. Thus, blocking or limiting

ARF-1 function at the Golgi may be part of the signal to activate these genes.

Immunity-linked gene upregulation in models of trafficking dysfunction

The Golgi apparatus accepts proteins from the ER that are destined for secretion, processing

them by glycosylation before secretory vesicles are loaded (*55*). Stimulation of the innate

immune system may significantly impact trafficking load, as antimicrobial peptides are shuttled

through the membrane trafficking system (*56*). Indeed, activation of the ER stress response has

been noted in multiple systems when large numbers of proteins need to be produced and secreted

(*57*). Interestingly, previous data suggest that induction of membrane stress may also be a sensor

of infection (*58*). Based on work by the Ewbank lab, they also suggested that some pathogen-

response genes act to support the trafficking of antimicrobial peptides (AMPs) (*58*, *59*). We

noted that the immunity-linked genes upregulated at low PC are largely outside of the

antimicrobial categories and are comprised of gene sets defined by their shared expression upon

pathogen exposure rather than function (*23*, *27*). We hypothesized that some of these genes

might be responding to direct stress on the trafficking system.

To explore this idea, we turned to two commonly used systems for studying trafficking in *C.*

elegans, comparing ILG gene expression in strains with increasing copy number of

VIT-2::GFP (*60*) and visualization of trafficking in both VIT-2::GFP and *myo-3*::ssGFP (*61*).

vit-2 is a vitellogenin produced and secreted from intestinal cells then taken up by the germline

(*62*). *myo-3*::ssGFP has a signal sequence on the fluorophore and is expressed in body wall

muscle (*61*). First, we used WormCat to examine category enrichment data from two published

studies examining genome-wide mRNA expression patterns when *vit-2* is overexpressed or

misregulated, placing stress on the secretory system. The Blackwell lab showed that germline-

less *glp-1* mutants exhibit *vit-2*::GFP buildup near its production site in the intestines (*63*).

WormCat analysis of their RNA seq data found strong enrichment patterns in STRESS

RESPONSE: Pathogen. (**Fig5A**, **C; DataS4**). Next, we used WormCat to examine RNA-seq

data from VIT-2::GFP (*64*) overexpressing animals produced by the Aballay group and also

found increases in the Stress Response category enrichment (**Fig5B, C; DataS4**). As two

models sharing VIT-2::GFP secretory overload show increases in STRESS RESPONSE:

Pathogen genes, we sought to directly test this model by comparing ILG expression in two

strains with a differing copy number of *vit-2*; RT130, made by microparticle bombardment and

containing the GFP array in addition to wild type copies (*65*) and BCN9071 which is a CRISPR-

generated allele integrated into the endogenous locus (*66*) (**Fig5D-F**). While VIT-2::GFP

intestinal puncta appeared similar in control conditions, RNAi of *arf-1* produced larger

aggregates in RT130 (**Fig5D**), suggesting that this strain is sensitized to stress in the secretory

system. Strikingly, expression of ILGs increase with VIT-2::GFP copy number (**Fig5G-J**),

demonstrating that ILG expression can be induced in the absence of pathogens by altering

increasing expression of proteins destined for the secretory pathway.

Figure 5: Immunity-linked gene expression is linked to trafficking disruption. WormCat

gene enrichment of upregulated genes from *glp-1*(AA2735) (*63*) and *vit-2(ac3)* compared to

SJ4005 (*64*) for Category 1 (**A**) and Category 2 (**B**). Legend is in **C**. **D.** Confocal projections of

C. elegans intestinal cells expressing *vit-2*::GFP (RT130) or *vit-2*::GFP (*BCN9701*) in wild type

or *arf-1(RNAi)* animals. qRT-PCR comparing gene expression in wild type, single copy *vit-*

2::GFP (BCN9071, s) or multi copy (RT130, m) animals. **E** and **F** show increase in GFP or *vit-*

2. Immune response genes are shown in **F-J**. Significance is determined by Students T Test.

Knockdown of STRESS RESPONSE: Pathogen genes disrupt a trafficking reporter

We found that ILGs become upregulated in the absence of pathogen response when lipid levels become unbalanced, ARF-1 functions are compromised, or when production of secreted proteins increases. This prompted us to ask if some ILGs might function as part of a response to help stimulate passage through the secretory pathway when trafficking load is high, as when pathogen responses drive antimicrobial protein production, or when the structures of the secretory system are compromised. As a pilot experiment, we focused on one of the most strongly upregulated genes, *hpo-6*, which is not present in any of the commercially available RNAi libraries. To study its function more closely in this context, we made a cDNA construct to allow RNAi in VIT-2::GFP RT130 in wild type and *sams-1(lof)* animals (*9*). *hpo-6* was originally identified in a screen for genes whose loss increased sensitivity to a pore-forming toxin (*67*). It has a glycoprotein domain and may occur in membrane rafts (*68*), but has no apparent homology to human proteins. *hpo-6(RNAi)* causes a slight increase in VIT-2::GFP in wild-type animals (**FigS6A**). However, in the low PC *sams-1(lof)* background, VIT-2::GFP pooling and aggregation suggested a broad effect on trafficking patterns. Yolk in *C. elegans* consists of both lipids and proteins (*69*). Because low lipid levels in *sbp-1(RNAi)* animals makes results with VIT-2::GFP difficult to interpret, we used the *myo-3*::ssGFP reporter model (*61*), which expresses and secretes GFP in body wall muscle cells.

We conducted a small scale screen of 19 other STRESS RESPONSE: Pathogen: unassigned genes on secretion in *myo-3*::ssGFP and the sensitized *sbp-1(ep79)*; *myo-3::GFP* backgrounds, imaging body wall muscle cells in triplicate and manually scoring aggregation in blinded

samples (**Fig6A**). *hpo-6* and two other candidate genes, *irg-2* and ZK6.11/*irg-8* were validated

by confocal imaging, blinded visual scoring and quantitation of puncta size and number with

Cell Profiler (*70*). We found that as in *arf-1* RNAi, reduction in *hpo-6, irg-2* and *irg-8*/ZK6.11

increased puncta size (**Fig6B-D; FigS6; D-E**), with related decrease in aggregate number in both

wild type and the *sbp-1(ep79)* background, reflecting aggregation and disruption of normal

secretory processes.

Many of the STRESS RESPONSE: Pathogen genes are expressed in the intestine; therefore, we

also examined a ssGFP expressed under the *glo-1* promoter (*71*) (**Fig6E**). Using epifluorescence

imaging, we found that intestinal ssGFP accumulated after RNAi of *sbp-1, arf-1, hpo-6, irg-2*

and *irg-8* (**Fig6F, G**). Confocal imaging shows that ssGFP puncta aggregate in the intestine of

*sbp-1(RNAi)*animals, as in muscle tissue and that this also occurs with *hpo-6* RNAi (**Fig6G, H**).

Thus, in a variety of cell types, we find that interference with *sbp-1*, or candidate STRESS

RESPONSE: Pathogen genes, causes aggregation and pooling of ssGFP in the absence of

pathogen stimulation, suggesting a role in secretory processes.

Figure 6: ILG loss can enhance aggregation of secreted proteins.

A. Schematic diagram of small-scale screen of STRESS RESPONSE: Pathogen: *other* genes for

aggregation of a secreted reporter in body wall muscle (*myo-3*::ssGFP). Confocal projections of

ssGFP in body wall muscle in control or the *sbp-1(ep79)* background (**B**). Quantification of

three independent replicates by Cell Profiler determining number of puncta (**C, D**) and puncta

area (**Figure S6D, E**). **E.** Epifluorescence images of ssGFP expressed from an intestinal

reporter with DIC images as insets. Quantitation of pixel intensity with ImageJ is in **F**.

Confocal projections showing ssGFP aggregation in **G**, with puncta quantitation by Cell Profiler

in **H**. Significance is determined by the Mann-Whitney test.

475 **Knockdown of STRESS RESPONSE: Pathogen genes affects an endogenous PA14**

476 **stimulated peptide**

477 The secretory pathway carries out many basal functions and must increase capacity during

478 stresses which require increases in protein production, such as pathogen challenge. The ER

479 stress pathway plays a key role during this increase (*72, 73*) and our results suggest that

480 alterations in the Golgi may also be sensed and result in upregulation of ILGs. Our studies

481 ssGFP reporters allowed isolation of effects of lipid imbalances on basal secretory processes

482 from the complex response to pathogens. However, ILGs are defined by high level expression

483 upon pathogen exposure and could also affect secretion of antimicrobial peptides (AMPs) or

484 other defensive molecules. *C. elegans* contains a large number of proteins with domains

485 common to AMPs such as ShKT, Sushi and CUB domains (*74*). In order to test effects of the

486 ILGs on an endogenous secreted protein in the pathogen response, we selected C32H11.9/*cld-1*

487 (cub-like domain 1). In our previous RNA seq experiments, it was the highest upregulated gene

488 after PA14 exposures (**Fig7A**) and was also strongly induced during other early timepoints (*22*),

489 but not induced by lipid imbalances in *sbp-1(RNAi)* animals (**Fig7B**). It contains a signal

490 sequence, suggesting that is secreted and a CUB domain (**Fig7C**). We obtained an allele with

491 GFP fused to the C terminus of *cld-1*/C32H11.6 using CRISPR

492 (PHX2081(C32H11.6::GFP(*sybIS7080*))), which accumulated in intestine after PA14 exposure

493 (**Fig7D**), consistent with disruption with secretion. Notably, RNAi of *sbp-1* disrupted CLD-

494 1::GFP production, causing increases total levels, aggregation and accumulation in the

495 pseudocoelom, consistent with defects in secretory processes resulting from lipid imbalances

496 (**Fig7D**). Knockdown of with *arf-1* also produced similar effects, strengthening these links.

497 Interference with *hpo-6*, a candidate ILG, also caused accumulation and aggregation of CLD-

498   1::GFP during PA14 exposure, suggesting that some ILG genes act to support secretory function

499   during stress, which may be initiated by lipid imbalances or pathogen exposure.

Figure 7: ILG loss can enhance intestinal accumulation of a Pseudomonas-induced CUB

domain peptide.

Expression levels of C32H9.11 in response to Pseudomonas aeruginosa (PA14) from Ding, et al.

2018 (**A**) or after *sbp-1* RNAi from Ding, et al. 2015 (**B**). **C.** Domain structure of C32H11.9

showing signal sequence and CUB domain. **D.** Spinning disc confocal micrographs showing

intestinal accumulation of *cld-1*/C32H11.9::GFP in response to PA14 after *sbp-1*, *arf-1* or ILG

RNAi taken at 488mn. Gut granules are shown with an overlay of a micrograph of the same

section from taken at 564nm. Quantitation is in **E**.

Discussion

Pathogen elicited gene expression programs are part of a specialized stress response where specific sets of antimicrobial peptides are produced, then secreted in response to pathogens. This type of innate immune response initiates as pathogens are recognized by interaction with cellular proteins or cellular processes are disturbed (*75*). In *C. elegans*, evidence for microbe/pathogen associated molecular patterns (MAMPS/PAMPS) has been less clear, however, surveillance of cellular processes such as translation (*11*), initiating the unfolded protein response or impacting lipid bilayer function activates pathogen responsive signaling and transcription of ILGs (*9*, *15*, *16*, *34*). While some studies have suggested changing levels of specific lipids can prime pathogen responses (*34*), lipid perturbations activating pathogen-responsive gene expression do not necessarily confer pathogen resistance (*9*), and ILG expression levels may be much lower than what occurs in a pathogen response. Links between membrane lipid levels and immune gene activation also occur in other organisms. In *Drosophila*, synthesis of PC is critical to immune function, balancing stored lipids with need to maintain membranes for secretory processes (*76*). In mammals, interference with a master transcriptional regulator of lipid biosynthetic enzymes, *sbp-1/SREBP-1* activates a gene expression program shared with pathogen stress in both *C. elegans* and mammals (*14*). This raises the larger the question of why ILG expression increases when membrane lipids are unbalanced in the absence of pathogens.

Membrane lipids form the organelles of the secretory system, which can sense changes in protein homeostasis or lipids levels and elicit gene expression programs when stressed. Multiple well-studied pathways are linked to changes in proteostasis or lipid levels within the ER (*57*). These processes are orchestrated by ER-intrinsic proteins that sense changes in protein folding or

alterations in the lipid bilayer (*6*, *8*, *15*, *77*). These ER-linked events are also important during

pathogen exposure in *C. elegans*; changes in ribosomal function at the ER have a well-defined

impact on pathogen responses (*11*, *78*) and pathogen responses depend on ER stress pathways to

protect this organelle from the demands of the immune response (*73*, *79*). Interestingly, the Lee

lab also found that ILGs are upregulated when trafficking is inhibited through blocking

glycosylation in the ER (*80*). In addition, pathogen-responses require an intact ER stress

pathway to allow adjustment to the increased trafficking load (*58*, *73*), as antimicrobial peptides

are produced in mass in the ER, sent in lipid vesicles to the Golgi, where many are glycosylated

and then packaged into lipid vesicles for secretion (*81*). For example, the *C. elegans* genome

contains more than 300 genes that are produced in response to specific pathogens and which

share similarity to known antimicrobial effectors (*82*). However, many of the genes in the *C.*

elegans innate immune response do not have obvious similarity to antimicrobial effectors and are

defined solely as being responsive pathogens (*27*, *83*). It is this class that are largely shared

between pathogen-induced and membrane stress-induced stress programs. Several lines of

evidence prompted us to hypothesize that some of these genes might be part of a response linked

to stress in membranes of the secretory pathway rather than a direct response to an extrinsic

pathogen. First, the involvement of secretory organs was implicated by alterations in PC ratios

in the ER/Golgi in both *sams-1* and *sbp-1* RNAi lipidomes. Second, our targeted RNAi screen

for regulators of p*sysm-1::GFP* identified the *arf-1* GTPase and coatomer proteins, which are

critical Golgi/ER transport. Importantly, knockdown of *arf-1* was sufficient to induce ILG

upregulation. Strengthening this link between disruption of lipid levels and ARF-1 function, we

also found *sbp-1* RNAi altered ARF-1::mCherry localization and Golgi morphology.

Other organelles in the secretory pathway, such as the Golgi, have their own stress sensors and

response pathways (*84*), some of which act by impacting the ARF1 GTPase (*85*). ARF-1 is a

critical regulator of trafficking, coordinating retrograde traffic from the Golgi to the ER and

regulating secretory function in the trans-Golgi (*86*). Cycling of ARF-1 GTPases depends on

membrane localization of the ARF GTPase, GEF (Guanine exchange factor), and GAP (GTPase

activating protein) (*86*). We previously found that knockdown of PC synthesis enzymes blocked

ARF1 cycling and limited membrane association of the ARF GEF GBF1 in cultured human cells

(*25*). This suggests low PC levels affect ARF1 activity by limiting the ability of the GTPase,

GEF, and GAP to associate at the Golgi membrane and initiate GDP-GTP cycling. However, PC

could be linked to ARF1 through other mechanisms. ARF1 and phospholipase D (PLD) function

have been linked by multiple labs (*87*). PLD cleaves PC molecules to produce choline and

phosphatidic acid, which in turn stimulates vesicle formation (*88*). This regulatory loop requires

PC, which is low in ER/Golgi membranes in *sams-1* and *sbp-1* RNAi animals. DAG kinases,

which could be stimulated by changes in DAGs in the *sbp-1* or *sams-1(RNAi)* animals, can also

affect ARF activity and trafficking (*89*). Finally, Protein Kinase D has an important role in

maintaining Golgi structure, and its ortholog has been linked to immunity in *C. elegans* (*90*, *91*).

However, DAG kinase or PKD (Protein Kinase D) loss failed to activate the p*sysm-1::GFP*

reporter in our RNAi screen, suggesting they could be important in other regulatory contexts.

While the lipidomic changes in *sbp-1* and *sams-1* animals are broad and individual lipids could

act through multiple independent mechanisms, our results show that decreasing PC is sufficient

to limit ARF-1 function (*25*) and activate ILG expression in the absence of pathogen.

Stress-responsive genes, such as ILGs, may contain several classes of factors with distinct

functions. Many of the ILGs, for example, those in the STRESS RESPONSE: Pathogen:

unassigned WormCat categories (*27*, *28*), lack domains associated with AMP function. While

some could have antimicrobial properties with distinct motifs, the activation of these genes

through intrinsic stress in the Golgi prompted us to ask if others acted to support secretory

function. Our data showing that reduction of *hpo-6, irg-2* and *irg-8* increased aggregation and

pooling of a secreted reporter in control and *sbp-1(ep79)* backgrounds suggests they could

protect or support secretory function occurring after broad disruption in membrane lipids or after

pathogen exposure occurs because both processes stress the trafficking system. Thus, some

ILGs may function generally a "multi-membrane" stress response encompassing the roles of both

the ER and the Golgi in trafficking. Taken together, these results illustrate mechanisms linking

lipid metabolism to genes activated in the immune response through effects on the secretory

system.

Materials and Methods

C. elegans strains, RNAi constructs, and screening.

N2 (wild type), p*sysm-1::GFP (*AU78*), myo-3*::GFP (GS1912), MANS::GFP (RT1315) and

OP50-1 were obtained from the *Caenorhabditis* Genetics Center. Exglo-1::ssGFP (GH639) was

a gift from Dr. Greg Hermann (Lewis and Clark University). *myo-3::ssGFP*;*sbp-*

1(ep79)(WAL510) was constructed in this study. CRISPR-tagged ARF-1, p*arf-1::ARF-*

1::mCherry (knu418), was obtained from the In Vivo Biosystems (COP1415) and then

outcrossed 3 times (WAL10). Imaging of MANS::GFP was done from a strain also harboring a

CRISPR tagged warf-1::RFP (ker4) (WAL12). (*cld-1*/C32H11.6::GFP was constructed by

SUNY Biotech using CRISPR (PHX2081(*sybIS7080*)). Normal growth media (NGM) was used

unless otherwise noted. For RNAi, gravid adults were bleached onto NGM plates supplemented

with ampicillin, tetracycline, and 6mM IPTG and 10X concentrated bacterial cultures. *C.*

elegans were allowed to develop until the young adult stage before harvesting. Because of larval

arrest, *copa-1* RNAi bacteria were diluted 1:10 in control RNAi bacteria before plating. For the

RNAi screen of *psysm-1*::GFP, L1 larvae were plated into 96 well plates spotted with RNAi

bacteria, and L4/young adults were scored from -3 to +3 with 0 as no change in 4 independent

replicates on a Zeiss GFP dissecting scope with an Axiocam camera. Candidates that were

positive in ¾ replicates were tested an additional four times, imaging 3 sets of 10 animals per

RNAi. Images quantitation is described below. For RNAi screen of *myo-3*::ssGFP and *myo-*

3::ssGFP; *sbp-1(ep79)*, 4 independent replicates were subjected to RNAi, placed in channeled

agarose pads and confocal images were collected on a Leica SP6.

Cell culture and siRNA transfection

HepG2 cells (ATCC, HB-8065) were grown in Minimum Essential Medium (Invitrogen) plus

10% FBS (Invitrogen), Glutamine (Invitrogen), and Sodium Pyruvate (Invitrogen). siRNA

oligonucleotides transfections were done for 48 hours using Lipofectamine RNAiMAX

Transfection Reagent (Invitrogen, 13778100), and cells were held for 16 hours in 1%

Lipoprotein Deficient Serum (LDS) (Biomedical Technologies, BT907) and 25 μg/ml ALLN

(Calbiochem) for 30 min prior to harvesting.

Immunoblotting:

HepG2 cells: Syringe passage was used to lyse cells in High Salt RIPA (50 mM Tris, pH 7.4;

400 mM NaCl; 0.1% SDS; 0.5% NaDeoxycholate; 1% NP-40; 1 mM DTT, 2.5 μg/mL ALLN,

Complete protease inhibitors (Roche). Invitrogen NuPage gels (4- 12%) were used for protein

separation before transfer to nitrocellulose. Blots were probed with antibodies to phosphorylated

p38 MAP Kinase, total p38, and STAT1 were used as a control. Immune complexes were

visualized with Luminol Reagent (Millipore). Densitometry was performed by scanning the

film, then analysis of pixel intensity with ImageJ software. Graphs show the average of at least

three independent experiments with control values normalized to one.

C. elegans: Young adult animals were lysed by sonication in High Salt RIPA, and

immunoblotting was performed as above.

Microscopy and Image quantitation

ARF-1::mCherry, MANS-1::GFP and ssGFP images were acquired on a Leica SPE confocal,

and projections of confocal slices were produced. All images were taken at identical gain

settings within experimental sets, and Adobe Photoshop was used for corrections to levels across

experimental sets. Image quantitation of GFP fluorescence was done using FIJI/ImageJ2.

Briefly, each photo was processed individually, beginning by splitting the image into three color

channels (Red/Green/Blue) and choosing the green channel for analysis. This image was then

converted to an 8-bit grey image and processed using the function "Enhance Local Contrast

(CLAHE)" (Blocksize: 250, Histogram bins: 256, Max Slope: 5.00, Mask: none, Fast (less

accurate): no (unchecked)). Using the circle selection tool, four sections (w: 55 h: 25) of the

brightest areas across each worm's intestine were selected from this processed image and

measured for intensity (integrated intensity). One additional section was taken from beside each

worm to account for background fluorescence. The background reading was subtracted from

each individual intestinal reading, then the four corrected readings were added to get a sum of

fluorescent intensity. Imaging of C32H11.9/*cld-1*::GFP was performed on a W1

Yokogawa spinning disk confocal microscope equipped with a Photometrics Prime BSI Express

sCMOS camera. Quantitation was performed with Nikon Imaging System software. Statistical

tests performed include Normality and Lognormality tests (Anderson-Darling, D'Agostino &

Pearson, Shapiro-Wilk, Kolmogorov-Smirnov) and Mann-Whitney tests.

Lipidomics

C. elegans total and microsomal lipidomics, including fractionation protocols, were performed

at the Whitehead Metabolomics core as in Smulan et al. 2016. Statistical analysis was

performed in Graphpad prism.

Gene expression analysis

Lysis of young adult *C. elegans* was performed in 0.5% SDS, 5% β-ME, 10 mM EDTA, 10 mM

Tris-HCl ph7.4, 0.5 mg/ml Proteinase K, before purification of RNA by TRI-Reagent (Sigma).

cDNA was produced with Transcriptor First-strand cDNA kits (Roche), and RT-PCR was

performed using Kappa SYBR Green 2X Mastermix.

RNA for sequencing was purified using RNAeasy columns (Qiagen). RNA sequencing

(including library construction was performed by BGI (Hong Kong). Reads were analyzed

through the Dolphin analysis platform (https://dolphin.umassmed.edu/), using DeSeq to identify

differentially expressed genes (*92*). Gene set enrichment was performed using WormCat

(www.wormcat.com) (*27*, *28*).

675
676

Acknowledgments

We would like acknowledge Drs. Seung-Jae Lee, Sujeong Kwon (Korea Advanced Institute of Science and Technology) for providing the *lpin-1* RNA seq data. In addition, Dr. Greg Hermann provided the *glo-1*::ssGFP strain (GH639). We would also like to thank Drs. Seung-Jae Lee, Emily Troemel (USD), Cole Haynes (UMASS Chan School of Medicine), Alexandra Byrne (UMASS Chan School of Medicine) and members of the Walker lab for comments on the manuscript in addition to Dr. Marian Walhout (UMASS Chan School of Medicine) for helpful discussion. We would like to acknowledge Wei Ding for technical support and Caroline Lewis (MIT) for metabolomics screening at the Whitehead Metabolomics core. We also thank Life Science Editors and Dr. Sabbi Lall for manuscript editing. Funding was from the NIA at the NIH: R01AG053355, R56AG068670 and R01AG068670. Some strains were provided by the CGC, which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440).

Author Contributions

Conceptualization; AKW, MJF, CMW, LJS

Verification: AKW, , MJF, CMW, LJS

Formal Analysis: AKW, MJF, CMW, LJS, DML

Investigation: AKW, MJF,CMW, LJS, DL, MF

Resources: AKW

Data Curation: AKW

Writing: AKW

Writing/Editing: AKW, MJF, CMW, DSL, LJS

708 Visualization: AKW
709
710 Supervision: AKW
711
712 Funding: AKW
713

Declaration of Interests

715 The authors declare no competing interests.

Data and materials availability:
718 All data are available in the main text or the supplementary materials will be shared. GEO

719 submission of RNA seq data is in progress.

References

1. K. E. Wellen, C. B. Thompson, A two-way street: reciprocal regulation of metabolism and signalling *Mol Cell*, (2010).

2. J. Aramburu, M. C. Ortells, S. Tejedor, M. Buxadé, C. López-Rodríguez, Transcriptional regulation of the stress response by mTOR. *Sci Signal*. 7, re2–re2 (2014).

3. H. G. Son, O. Altintas, E. J. E. Kim, S. Kwon, S. V. Lee, Age-dependent changes and biomarkers of aging in *Caenorhabditis elegans*. *Aging Cell*. 18, e12853 (2019).

4. K. Halbleib, K. Pesek, R. Covino, H. F. Hofbauer, D. Wunnicke, I. Hänelt, G. Hummer, R. Ernst, Activation of the Unfolded Protein Response by Lipid Bilayer Stres. *Mol Cell*, (2017).

5. M. A. Gianfrancesco, N. Paquot, J. Piette, S. Legrand-Poels, Lipid bilayer stress in obesity-linked inflammatory and metabolic disorders. *Biochem Pharmacol*. 153, 168–183 (2018).

6. J. H. Koh, L. Wang, C. Beaudoin-Chabot, G. Thibault, Lipid bilayer stress-activated IRE-1 modulates autophagy during endoplasmic reticulum stress. *J Cell Sci*. 131, jcs217992 (2018).

7. T. Radanović, J. Reinhard, S. Ballweg, K. Pesek, R. Ernst, An Emerging Group of Membrane Property Sensors Controls the Physical State of Organellar Membranes to Maintain Their Identity. *Bioessays*. 40, 1700250 (2018).

8. G. S. Hotamisligil, R. J. Davis, Cell Signaling and Stress Responses. *Csh Perspect Biol*. 8, a006072 (2016).

9. W. Ding, L. J. Smulan, N. S. Hou, S. Taubert, J. L. Watts, A. K. Walker, s-Adenosylmethionine Levels Govern Innate Immunity through Distinct Methylation-Dependent Pathways *Cell Metab*, (2015).

10. A. K. Walker, R. L. Jacobs, J. L. Watts, V. Rottiers, K. Jiang, D. M. Finnegan, T. Shioda, M. Hansen, F. Yang, L. J. Niebergall, D. E. Vance, M. Tzoneva, A. C. Hart, A. M. Näär, A Conserved SREBP-1/Phosphatidylcholine Feedback Circuit Regulates Lipogenesis in Metazoans. *Cell*, (2011).

11. T. L. Dunbar, Z. Yan, K. M. Balla, M. G. Smelkinson, E. R. Troemel, *C. elegans* detects pathogen-induced translational inhibition to activate immune signaling. *Cell Host Microbe*, (2012).

12. J. D. Horton, J. L. GOLDSTEIN, M. S. BROWN, SREBPs: Transcriptional Mediators of Lipid Homeostasis. *Cold Spring Harb Sym*. 67, 491–498 (2002).

13. R. M. McKay, J. P. McKay, L. Avery, J. M. Graff, C. elegans A Model for Exploring the
Genetics of Fat Storage. *Dev Cell*. 4, 131–142 (2003).

14. S. Wu, A. M. Näär, SREBP1-dependent de novo fatty acid synthesis gene expression is
elevated in malignant melanoma and represents a cellular survival trait. *Sci Rep-uk*. 9, 10369
(2019).

15. N. Ho, W. S. Yap, J. Xu, H. Wu, J. H. Koh, W. W. B. Goh, B. George, S. C. Chong, S.
Taubert, G. Thibault, Stress sensor Ire1 deploys a divergent transcriptional program in response
to lipid bilayer stress. *J Cell Biol*. 219 (2020), doi:10.1083/jcb.201909165.

16. Y. Jung, S. Kwon, S. Ham, D. Lee, H. H. Park, Y. Yamaoka, D. Jeong, M. Artan, O.
Altintas, S. Park, W. Hwang, Y. Lee, H. G. Son, S. W. A. An, E. J. E. Kim, M. Seo, S. V. Lee,
Caenorhabditis elegans Lipin 1 moderates the lifespan-shortening effects of dietary glucose by
maintaining ω-6 polyunsaturated fatty acids. *Aging Cell*. 19 (2020), doi:10.1111/acel.13150.

17. J. D. Nhan, C. D. Turner, S. M. Anderson, C.-A. Yen, H. M. Dalton, H. K. Cheesman, D. L.
Ruter, N. U. Naresh, C. M. Haynes, A. A. Soukas, R. Pukkila-Worley, S. P. Curran, Redirection
of SKN-1 abates the negative metabolic outcomes of a perceived pathogen infection. *Proc*
National Acad Sci. 116, 22322–22330 (2019).

18. Z. Wu, M. Isik, N. Moroz, M. J. Steinbaugh, P. Zhang, T. K. Blackwell, Dietary Restriction
Extends Lifespan through Metabolic Regulation of Innate Immunity. *Cell Metab*. 29, 1192-
1205.e8 (2019).

19. I. Engelmann, A. Griffon, L. Tichit, F. Montañana-Sanchis, G. Wang, V. Reinke, R. H.
Waterston, L. W. Hillier, J. J. Ewbank, A Comprehensive Analysis of Gene Expression Changes
Provoked by Bacterial and Fungal Infection in C. elegans. *Plos One*. 6, e19055 (2011).

20. G. V. Mallo, C. L. Kurz, C. Couillault, N. Pujol, S. Granjeaud, Y. Kohara, J. J. Ewbank,
Inducible Antibacterial Defense System in *C. elegans*. *Curr Biol*. 12, 1209–1214 (2002).

21. M. Fletcher, E. J. Tillman, V. L. Butty, S. S. Levine, D. H. Kim, Global transcriptional
regulation of innate immunity by ATF-7 in *C. elegans*. *Plos Genet*. 15, e1007830 (2019).

22. E. R. Troemel, S. W. Chu, V. Reinke, S. S. Lee, F. M. Ausubel, D. H. Kim, p38 MAPK
regulates expression of immune response genes and contributes to longevity in *C. elegans*. *Plos*
Genet. 2, e183 (2006).

23. K. A. Estes, T. L. Dunbar, J. R. Powell, F. M. Ausubel, E. R. Troemel, bZIP transcription
factor *zip-2* mediates an early response to Pseudomonas aeruginosa infection in *Caenorhabditis*
elegans. *Proc National Acad Sci*, (2010).

24. R. P. Shivers, T. Kooistra, S. W. Chu, D. J. Pagano, D. H. Kim, Tissue-Specific Activities of
an Immune Signaling Module Regulate Physiological Responses to Pathogenic and Nutritional
Bacteria in *C. elegans*. *Cell Host Microbe*. 6, 321–330 (2009).

25. L. J. Smulan, W. Ding, E. Freinkman, S. Gujja, Y. J. K. Edwards, A. K. Walker, Cholesterol-
Independent SREBP-1 Maturation Is Linked to ARF1 Inactivation. *Cell Reports*, (2016).

26. N. Zmora, S. Bashiardes, M. Levy, E. Elinav, The Role of the Immune System in Metabolic
Health and Disease. *Cell Metab*. 25, 506–521 (2017).

27. A. D. Holdorf, D. P. Higgins, A. C. Hart, P. R. Boag, G. J. Pazour, A. J. M. Walhout, A. K.
Walker, WormCat: An Online Tool for Annotation and Visualization of *Caenorhabditis elegans*
Genome-Scale Data. *Genetics*, (2019).

28. D. P. Higgins, C. M. Weisman, D. S. Lui, F. A. D'Agostino, A. K. Walker, Defining
characteristics and conservation of poorly annotated genes in *Caenorhabditis elegans* using
WormCat 2.0. *Genetics* (2022), doi:10.1093/genetics/iyac085.

29. W. Ding, D. P. Higgins, D. K. Yadav, A. A. Godbole, R. Pukkila-Worley, A. K. Walker,
Stress-responsive and metabolic gene regulation are altered in low S-adenosylmethionine. *Plos*
Genet. 14, e1007812 (2018).

30. S. Taubert, M. R. V. Gilst, M. Hansen, K. R. Yamamoto, A Mediator subunit, MDT-15,
integrates regulation of fatty acid metabolism by NHR-49-dependent and -independent pathways
in *C. elegans*. *Gene Dev*. 20, 1137–1149 (2006).

31. G. Y. S. Goh, J. J. Winter, F. Bhanshali, K. R. S. Doering, R. Lai, K. Lee, E. A. Veal, S.
Taubert, NHR-49/HNF4 integrates regulation of fatty acid metabolism with a protective
transcriptional response to oxidative stress and fasting. *Aging Cell*. 17, e12743 (2018).

32. M. L. Ladage, S. D. King, D. J. Burks, D. L. Quan, A. M. Garcia, R. K. Azad, P. A. Padilla,
Glucose or Altered Ceramide Biosynthesis Mediate Oxygen Deprivation Sensitivity Through
Novel Pathways Revealed by Transcriptome Analysis in *Caenorhabditis elegans*. *G3 Genes*
Genomes Genetics, (2016).

33. Y. J. Liu, G. E. Janssens, R. L. McIntyre, M. Molenaars, R. Kamble, A. W. Gao, A.
Jongejan, M. van Weeghel, A. W. MacInnes, R. H. Houtkooper, Glycine promotes longevity in
Caenorhabditis elegans in a methionine cycle-dependent fashion. *Plos Genet*. 15, e1007633
(2019).

34. N. D. Peterson, J. D. Icso, J. E. Salisbury, T. Rodríguez, P. R. Thompson, R. Pukkila-
Worley, Pathogen infection and cholesterol deficiency activate the *C. elegans* p38 immune
pathway through a TIR-1/SARM1 phase transition. *Elife*. 11, e74206 (2022).

35. D. P. Higgins, C. M. Weisman, D. S. Lui, F. A. D'Agostino, A. K. Walker, WormCat 2.0
defines characteristics and conservation of poorly annotated genes in *Caenorhabditis elegans*.
Genetics (2022), doi:10.1093/genetics/iyac085.

36. X. Shen, R. E. Ellis, K. Sakaki, R. J. Kaufman, Genetic Interactions Due to Constitutive and
Inducible Gene Regulation Mediated by the Unfolded Protein Response in C. elegans. *Plos*
Genet. 1, e37 (2005).

37. F. Yang, B. W. Vought, J. S. Satterlee, A. K. Walker, Z.-Y. J. Sun, J. L. Watts, R.
DeBeaumont, R. M. Saito, S. G. Hyberts, S. Yang, C. Macol, L. Iyer, R. Tjian, S. van den
Heuvel, A. C. Hart, G. Wagner, A. M. Näär, *Nature* (2006).

38. J. C. M. Holthuis, A. K. Menon, *Nature*, Lipid landscapes and pipelines in membrane
homeostasis (2014).

39. D. Jeong, D. Lee, S. Hwang, Y. Lee, J. Lee, M. Seo, W. Hwang, K. Seo, A. B. Hwang, M.
Artan, H. G. Son, J. Jo, H. Baek, Y. M. Oh, Y. Ryu, H. Kim, C. M. Ha, J. Yoo, S. V. Lee,
Mitochondrial chaperone HSP-60 regulates anti-bacterial immunity via p38 MAP kinase
signaling. *Embo J*. 36, 1046–1065 (2017).

40. C. L. Pender, H. R. Horvitz, Hypoxia-inducible factor cell non-autonomously regulates C.
elegans stress responses and behavior via a nuclear receptor. *Elife*. 7, e36828 (2018).

41. J. L. Watts, J. Browse, Genetic dissection of polyunsaturated fatty acid synthesis in
Caenorhabditis elegans. *Proc National Acad Sci*, (2006).

42. M. Ruiz, R. Bodhicharla, E. Svensk, R. Devkota, K. Busayavalasa, H. Palmgren, M.
Ståhlman, J. Boren, M. Pilon, Membrane fluidity is regulated by the C. elegans transmembrane
protein FLD-1 and its human homologs TLCD1/2. *Elife*. 7, 1309 (2018).

43. R. Bodhicharla, R. Devkota, M. Ruiz, M. Pilon, *Genetics*, Membrane Fluidity Is Regulated
Cell Non-autonomously by Caenorhabditis elegans PAQR-2 and Its Mammalian Homolog
AdipoR2 (2018).

44. N. Ho, C. Xu, G. Thibault, From the unfolded protein response to metabolic diseases – lipids
under the spotlight. *J Cell Sci*. 131, jcs199307 (2018).

45. A. Spang, ARF1 regulatory factors and COPI vesicle formation. *Curr Opin Cell Biol*. 14,
423–427 (2002).

46. A. G. Fraser, R. S. Kamath, P. Zipperlen, M. Martinez-Campos, M. Sohrmann, J. Ahringer,
Functional genomic analysis of *C. elegans* chromosome I by systematic RNA interference.
Nature. 408, 325–330 (2000).

47. M. A. Gustafson, J. C. Fromme, Regulation of Arf activation occurs via distinct mechanisms
at early and late Golgi compartments. *Mol Biol Cell*, (2017).

48. J. R. Broekhuis, S. Rademakers, J. Burghoorn, G. Jansen, SQL-1, homologue of the Golgi
protein GMAP210, modulates intraflagellar transport in *C. elegans*. *J Cell Sci*. 126, 1785–1795
(2013).

49. C. C.-H. Chen, P. J. Schweinsberg, S. Vashist, D. P. Mareiniss, E. J. Lambie, B. D. Grant,
RAB-10 Is Required for Endocytic Recycling in the Caenorhabditis elegans Intestine *Mol Biol*
Cell, (2006).

50. M. M. Rolls, D. H. Hall, M. Victor, E. H. K. Stelzer, T. A. Rapoport, Targeting of Rough
Endoplasmic Reticulum Membrane Proteins and Ribosomes in Invertebrate Neurons. *Mol Biol*
Cell. 13, 1778–1791 (2002).

51. R. D. Klausner, J. G. Donaldson, J. Lippincott-Schwartz, Brefeldin A: insights into the
control of membrane traffic and organelle structure. *The Journal of cell biology*, 1992.

52. K. Sato, M. Sato, A. Audhya, K. Oegema, P. Schweinsberg, B. D. Grant, Dynamic
Regulation of Caveolin-1 Trafficking in the Germ Line and Embryo of Caenorhabditis elegans
Mol Biol Cell, (2006).

53. K. B. Ackema, U. Sauder, J. A. Solinger, A. Spang, The ArfGEF GBF-1 Is Required for ER
Structure, Secretion and Endocytic Transport in *C. elegans*. *Plos One*. 8, e67076 (2013).

54. I. D. Zelnik, A. E. Ventura, J. L. Kim, L. C. Silva, A. H. Futerman, The role of ceramide in
regulating endoplasmic reticulum function. *Biochimica Et Biophysica Acta Bba - Mol Cell*
Biology Lipids (2019), doi:10.1016/j.bbalip.2019.06.015.

55. G. Boncompain, A. V. Weigel, Transport and sorting in the Golgi complex: multiple
mechanisms sort diverse cargo. *Curr Opin Cell Biol*. 50, 94–101 (2018).

56. T. Taguchi, K. Mukai, Innate immunity signalling and membrane trafficking. *Curr Opin Cell*
Biol. 59, 1–7 (2019).

57. B. M. Gardner, D. Pincus, K. Gotthardt, C. M. Gallagher, P. Walter, Endoplasmic Reticulum
Stress Sensing in the Unfolded Protein Response. *Csh Perspect Biol*. 5, a013169 (2013).

58. T. Lamitina, E. Chevet, To UPR… and beyond!: A new role for a BiP/GRP78 protein in the
control of antimicrobial peptide expression in *C. elegans* epidermis. *Virulence*. 3, 238–240
(2012).

59. C. Couillault, P. Fourquet, M. Pophillat, J. J. Ewbank, A UPR-independent infection-specific
role for a BiP/GRP78 protein in the control of antimicrobial peptide expression in *C. elegans*
epidermis *Virulence*, 2012.

60. B. D. Grant, M. Sato, Intracellular trafficking. *Wormbook Online Rev C Elegans Biology*, 1–
9 (2006).

61. H. Fares, I. Greenwald, Genetic Analysis of Endocytosis in *Caenorhabditis elegan*s :
Coelomocyte Uptake Defective Mutants. *Genetics*. 159, 133–145 (2001).

62. B. Grant, D. Hirsh, Evidence that RME-1, a conserved *C. elegans* EH-domain protein,
functions in endocytic recycling *Mol Biol Cell*, (2001).

63. M. J. Steinbaugh, S. D. Narasimhan, S. Robida-Stubbs, L. E. M. Mazzeo, J. M. Dreyfuss, J.
M. Hourihan, P. Raghavan, T. N. Operaña, R. Esmaillie, T. K. Blackwell, Lipid-mediated
regulation of SKN-1/Nrf in response to germ cell absence. *Elife*. 4, 466 (2015).

64. J. Singh, A. Aballay, Endoplasmic Reticulum Stress Caused by Lipoprotein Accumulation
Suppresses Immunity against Bacterial Pathogens and Contributes to Immunosenescence. *Mbio*.
8, e00778-17 (2017).

65. Z. Balklava, N. D. Rathnakumar, S. Vashist, P. J. Schweinsberg, B. D. Grant, *Genetics*,
Genome-wide analysis identifies a general requirement for polarity proteins in endocytic traffic
(2016).

66. L. V. Rompay, C. Borghgraef, I. Beets, J. Caers, L. Temmerman, New genetic regulators
question relevance of abundant yolk protein production in *C. elegans Sci Rep-uk*, (2015).

67. C.-Y. Kao, F. C. O. Los, D. L. Huffman, S. Wachi, N. Kloft, M. Husmann, V. Karabrahimi,
J.-L. Schwartz, A. Bellier, C. Ha, Y. Sagong, H. Fan, P. Ghosh, M. Hsieh, C.-S. Hsu, L. Chen, R.
V. Aroian, Global Functional Analyses of Cellular Responses to Pore-Forming Toxins. *Plos*
Pathog. 7, e1001314 (2011).

68. W. Rao, R. E. Isaac, J. N. Keen, An analysis of the *Caenorhabditis elegans* lipid raft
proteome using geLC-MS/MS *J Proteomics*, (2011).

69. M. F. Perez, B. Lehner, Vitellogenins - Yolk Gene Function and Regulation in
Caenorhabditis elegans. *Front Physiol*. 10, 1067 (2019).

70. A. E. Carpenter, T. R. Jones, M. R. Lamprecht, C. Clarke, I. H. Kang, O. Friman, D. A.
Guertin, J. H. Chang, R. A. Lindquist, J. Moffat, P. Golland, D. M. Sabatini, CellProfiler: image
analysis software for identifying and quantifying cell phenotypes. *Genome Biol*. 7, R100 (2006).

71. G. J. Hermann, L. K. Schroeder, C. A. Hieb, A. M. Kershner, B. M. Rabbitts, P. Fonarev, B.
D. Grant, J. R. Priess, Genetic Analysis of Lysosomal Trafficking in *Caenorhabditis elegans*.
Mol Biol Cell. 16, 3273–3288 (2005).

72. A. Shaheen, Effect of the unfolded protein response on ER protein export: a potential new
mechanism to relieve ER stress. *Cell Stress Chaperones*. 23, 797–806 (2018).

73. C. E. Richardson, T. Kooistra, D. H. Kim, An essential role for XBP-1 in host protection
against immune activation in C. elegans. *Nature*, (2010).

74. J. J. Ewbank, O. Zugasti, *C. elegans*: model host and tool for antimicrobial drug discovery.
Dis Model Mech. 4, 300–304 (2011).

75. R. Medzhitov, Recognition of microorganisms and activation of the immune response.
Nature. 449, 819–826 (2007).

76. B. A. Martínez, R. G. Hoyle, S. Yeudall, M. E. Granade, T. E. Harris, J. D. Castle, N.
Leitinger, M. L. Bland, Innate immune signaling in *Drosophila* shifts anabolic lipid metabolism
from triglyceride storage to phospholipid synthesis to support immune function. *Plos Genet*. 16,
e1009192 (2020).

77. M. Safra, S. Ben-Hamo, C. Kenyon, S. Henis-Korenblit, The *ire-1* ER stress-response
pathway is required for normal secretory-protein metabolism in *C. elegans*. *J Cell Sci*. 126,
4136–4146 (2013).

78. D. L. McEwan, N. V. Kirienko, F. M. Ausubel, *Cell Host Microbe*, (2010).

79. E. J. Tillman, C. E. Richardson, D. J. Cattie, K. C. Reddy, N. J. Lehrbach, R. Droste, G.
Ruvkun, D. H. Kim, *Genetics*, (2018).

80. D.-E. Jeong, Y. Lee, S. Ham, D. Lee, S. Kwon, H.-E. H. Park, S.-Y. Hwang, J.-Y. Yoo, T.-Y.
Roh, S.-J. V. Lee, Inhibition of the oligosaccharyl transferase in *Caenorhabditis elegans* that
compromises ER proteostasis suppresses p38-dependent protection against pathogenic bacteria.
Plos Genet. 16, e1008617 (2020).

81. N. G. Bednarska, B. W. Wren, S. J. Willcocks, The importance of the glycosylation of
antimicrobial peptides: natural and synthetic approaches. *Drug Discov Today*. 22, 919–926
(2017).

82. K. Dierking, W. Yang, H. Schulenburg, Antimicrobial effectors in the nematode
Caenorhabditis elegans: an outgroup to the Arthropoda. *Philosophical Transactions Royal Soc B*
Biological Sci. 371, 20150299 (2016).

83. E. de Nadal, G. Ammerer, F. Posas, Controlling gene expression in response to stress. *Nat*
Rev Genet. 12, 833–845 (2011).

84. C. E. Machamer, The Golgi complex in stress and death. *Front Neurosci-switz*. 9, 421
(2015).

85. J. H. Reiling, A. J. Olive, S. Sanyal, J. E. Carette, T. R. Brummelkamp, H. L. Ploegh, M. N.
Starnbach, D. M. Sabatini, *Nat Cell Biol*, (2013).

86. J. G. Donaldson, C. L. Jackson, *Nat Rev Mol Cell Bio*, (2011).

87. C. Riebeling, A. J. Morris, D. Shields, *Biochimica Et Biophysica Acta Bba - Mol Cell*
Biology Lipids, (2009).

88. M. McDermott, M. J. O. Wakelam, A. J. Morris, Phospholipase D. *Biochem Cell Biol*. 82,
225–253 (2004).

89. R. P. H. Huijbregts, L. Topalof, V. A. Bankaitis, Lipid Metabolism and Regulation of
Membrane Trafficking. *Traffic*. 1, 195–202 (2000).

90. M. Ren, H. Feng, Y. Fu, M. Land, C. S. Rubin, Protein Kinase D Is an Essential Regulator of
C. elegans Innate Immunity. *Immunity*. 30, 521–532 (2009).

91. V. Malhotra, F. Campelo, *Csh Perspect Biol*, (2011).

92. O. Yukselen, O. Turkyilmaz, A. R. Ozturk, M. Garber, A. Kucukural, DolphinNext: a
distributed data processing platform for high throughput genomics. *Bmc Genomics*. 21, 310
(2020).

959