Type I interferon shapes brain distribution and tropism of tick-borne flavivirus

Nunya Chotiwan[1,2,3*,#], Ebba Rosendal[1,2,#], Stefanie M. A. Willekens[1,2,4,#], Erin

Schexnaydre[1,2,5,6], Emma Nilsson[1,2], Richard Lindqvist[1,2] , Max Hahn[4], Ionut Sebastian

Mihai[2,7,10], Federico Morini[4], Jianguo Zhang[1,2,5,6], Gregory D. Ebel[8], Lars-Anders

Carlson[2,5,6], Johan Henriksson[2,7] , Ulf Ahlgren[4], Daniel Marcellino[9], Anna K. Överby[1,2*]

[1] Department of Clinical Microbiology, Umeå University; 90185 Umeå, Sweden

[2] The Laboratory for Molecular Infection Medicine Sweden (MIMS), Umeå University; 90187 Umeå,

Sweden

[3] Chakri Naruebodindra Medical Institute, Faculty of Medicine Ramathibodi Hospital, Mahidol University,

Bangpla, Bangpli, Samutprakarn, 10540, Thailand

[4] Umeå Centre for Molecular Medicine (UCMM), Umeå University; 90187 Umeå, Sweden

[5] Wallenberg Centre for Molecular Medicine, Umeå University; 90187 Umeå, Sweden

[6] Department of Medical Biochemistry and Biophysics, Umeå University; 90187 Umeå, Sweden

[7] Department of Department of Molecular biology, Umeå University; 90187 Umeå, Sweden

[8] Department of Microbiology, Immunology and Pathology, Colorado State University, USA

[9] Department of Integrative Medical Biology, Umeå University; 90187 Umeå, Sweden

[10] Företagsforskarskolan, Umeå University, 90187 Umeå, Sweden

*Corresponding authors: Anna K Överby and Nunya Chotiwan

[#] Contributed equally

Email: anna.overby@umu.se and nunya.chotiwan@umu.se

Abstract

Viral tropism within the brain and the role(s) of vertebrate immune response to neurotropic flaviviruses

infection is largely understudied. We combined multimodal imaging (cm-nm scale) with single nuclei RNA-

sequencing to study Langat virus in wildtype and interferon alpha/beta receptor knockout ($Ifnar^{-/-}$) mice to

visualize viral pathogenesis and define molecular mechanisms. Whole brain viral infection was imaged by

Optical Projection Tomography coregistered to *ex vivo* MRI. Infection was limited to grey matter of

sensory systems in wildtype mice, but extended into white matter, meninges and choroid plexus in $Ifnar^{-/-}$

mice. Cells in wildtype displayed strong type I and II IFN responses, likely due to *Ifnb* expressing

astrocytes, infiltration of macrophages and *Ifng*-expressing CD8+NK cells, whereas in $Ifnar^{-/-}$, the

absence of this response contributed to a shift in cellular tropism towards non-activated resident

microglia. Multimodal imaging-transcriptomics exemplifies a powerful way to characterize mechanisms of

viral pathogenesis and tropism.

Introduction

Viruses from multiple families are able to cross the physical brain barriers, infect the central nervous system (CNS) and cause disease [1,2]. These include several neurotropic viruses from the genus *Flavivirus*, such as West Nile virus (WNV), Zika virus (ZIKV), Japanese encephalitis virus (JEV) and tick-borne encephalitis virus (TBEV). These flaviviruses frequently cause severe diseases in humans, including encephalitis and meningitis, and lead to significant mortality and long-term neurological sequelae among survivors [3-5]. While global disease burden remains high, no direct treatment for these viral infections is currently available. Consequently, patients rely solely on symptomatic treatment to moderate clinical disease and innate and acquired immune responses to respond to infection, ultimately resulting in virus clearance. Therefore, a better understanding of viral pathogenesis and the triggered immune response is required for the development of additional diagnostic tools, effective therapies, and preventive measures [6,7]

Induction of type I interferon (IFN-I) is a critical part of the host innate immune response, an important first-line defense against viral infection [8]. The IFN-I response can be activated locally in the CNS and is essential to protect the brain against neurotropic flaviviruses [9-11]. *In vitro* studies have demonstrated that the IFN-I response is rapidly induced in astrocytes upon TBEV infection and this response protects neurons from infection [12,13]. In models with a deficient or compromised IFN-I response, neurons and astrocytes were more susceptible *in vitro* [12,14]. In rodent models, this response limits the spread of viral CNS infection and protects mice from lethal encephalitis by Langat virus (LGTV), a less-virulent model for TBEV [9,15]. Furthermore, mice with a deficient IFN-I response showed increased inflammation, blood-brain barrier (BBB) breakdown and increased immune cell infiltration compared to wildtype (WT) mice after LGTV infection [9]. In addition to viral receptors and entry factors, the innate immune response is known to determine cell susceptibility to viral infection [14,16,17], yet how the IFN-I response influences this tropism and shapes global distribution of tick-borne flavivirus infection in the brain is not well understood.

Therefore, we infected WT mice and mice lacking the IFN-I response (*Ifnar$^{-/-}$*) intracranially with LGTV,

monitored virus distribution in the brain using advanced imaging techniques and measured cellular

responses to infection by single nuclei RNA sequence (snRNAseq). For a holistic view of viral brain

infection, we devised an imaging approach that combines viral distribution in the whole brain (~0.5 cm^3),

obtained by Optical Projection Tomography (OPT) [18], with detailed anatomic information provided by an

ex vivo magnetic resonance imaging (MRI)-based brain template. We found sensory grey matter (GM)

areas infected in WT mice and this distribution expanded to white matter (WM) in the absence of IFN-I

response. Using additional imaging modalities, zooming in to nm scale, we found that the IFN-I response

restricts viral replication in meninges and epithelial cells of choroid plexus (ChP). Interestingly, the cellular

tropism of LGTV in cerebrum shifted from neurons in WT towards Iba1-positive cells in *Ifnar$^{-/-}$* mice. We

then used snRNAseq of cerebral cortex to clarify the transcriptional changes in different cell populations

to identify the molecular mechanism of viral pathogenesis. We found that infection in WT mice induced a

strong inflammatory response with expression of both type I and type II IFNs (IFN-II), which protected

microglia from infection. We also observed infiltration of several distinct immune cell populations,

including a large population of peripheral macrophages (MØs). Together, our data provide an

unprecedented view of virus distribution and tropism within the brain and highlight the importance of local

IFN-I induction in influencing virus distribution, tropism and pathogenesis.

Results

Whole brain 3D imaging and coregistration of LGTV infection with MRI in WT and Ifnar$^{-/-}$ mice

Local IFN-I restricts neurotropic flavivirus infection in rodent brain as demonstrated by nucleic acid

amplification and immunohistochemistry assays [9,12,15,19]. However, these techniques do not provide a

comprehensive understanding of viral distribution. Therefore, we set out to establish an imaging approach

for neurotropic virus infection in whole mouse brain using OPT, to specifically investigate the effect of

IFN-I on viral distribution. To surpass brain barriers, we infected mice intracranially, and brains were

collected at humane endpoint, 6-7 days and 4-5 days post-infection for WT and *Ifnar$^{-/-}$*, respectively

(Supplementary Fig. 1a). Whole brains were bleached to reduce auto-fluorescence, immunolabeled

against LGTV nonstructural protein 5 (NS5), mounted in agarose, optically cleared, and scanned using

OPT (Fig. 1a,b). Tomographic reconstruction enabled visualization of viral infection throughout the entire

brain in 3D (Supplementary Fig. 1b). Applying this method, we observed that viral infection in WT brains

was predominantly localized in cerebral cortex, with weak signals in the olfactory bulb and with no

detectable infection in lateral ventricles nor the fourth ventricle (Fig. 1c inset *ii*, Supplementary Fig.1c, and

Supplementary Movie 1). In contrast, in *Ifnar*$^{-/-}$ brains, we observed weak infection throughout the

cerebral cortex but a pronounced infection in the meninges, third ventricle, fourth ventricle, and the

interior wall of the lateral ventricles, spanning into the anterior cerebrum and olfactory bulbs, a pattern

closely resembling the rostral migratory stream (Fig. 1c inset *iii*, Supplementary Fig. 1c and

Supplementary Movie 1). We also detected some low-level unspecific signal in uninfected brains (mock

group) (Fig. 1c inset *i* and Supplementary Fig. 1c), indicating possible ventricular antibody trapping or a

low, unspecific antibody absorption in meninges during whole brain immunolabeling.

Although OPT enables visualization of LGTV distribution in the whole brain with great sensitivity and high

spatial resolution (Fig. 1c), the anatomical information obtained from tissue autofluorescence using this

technique is insufficient to provide detailed anatomical information, especially in the cerebrum. As the

brain is divided into multiple structures [20], linked to different physiological functions, it would be of great

value to improve the anatomical reference frame, to allow precise identification of infected brain areas. To

address this, we acquired structural *ex vivo* MR images from brains after chemical preprocessing and

OPT acquisition. These images were coregistered with viral OPT signal, which resulted in fusion images

with good spatial alignment (Fig. 1d). In some infected brains, we also observed hyperintense lesions on

the T1-weighted images, suggesting virus-induced damages (Fig. 1d inset *i*). To obtain a reference frame

with improved anatomical detail and tissue contrast and to overcome high MRI-scanning costs of

individual brains in the future, we used the high-resolution Optically Cleared UMeå (OCUM) brain

template, which was generated from MR images acquired after clearing for optical imaging [21]. In

optimized OPT-OCUM fusion images, viral signal is displayed within a detailed anatomical context, in 3D,

where neuronal pathways and trajectories can be observed, *e.g.*, the rostral migratory stream in an *Ifnar*$^{-}$

$^{/-}$ infected brain (Fig. 1e).

IFN-I response restricts viral infection in meninges and ChP

In WT brains, we observed weak viral signal in the meninges and the interior wall of third ventricle on

OPT (Fig. 2 and Supplementary Fig. 1c). However, we did not detect viral antigen in these regions by

confocal microscopy (Fig. 2b, d), indicating non-specific antibody trapping in WT mice. In *Ifnar*$^{-/-}$ brains,

on the other hand, we observed intense viral OPT signal within meninges, which was confirmed by

confocal microscopy (Fig. 2b). Additionally, we observed strong viral signal located in all ventricles (Fig.

1c inset *iii*, 2a). To complement OPT data, we used light sheet fluorescence microscopy (LSFM) to obtain

higher resolution images of viral signal in the fourth ventricle and found that ChP, the secretory tissue that

resides within ventricles, was highly infected (Fig. 2c and Supplementary Movie 2). Confocal analysis

confirmed that ChP of all ventricles, as well as the ependymal cells lining the ventricular walls, were highly

infected in *Ifnar*$^{-/-}$ brains, but not in WT (Fig. 2d). To further evaluate cellular tropism within ChP, we

stained *Ifnar*$^{-/-}$ brains with different cellular markers and the viral antigen (NS5) and found that viral

infection colocalized with the epithelial cell marker aquaporin-1 (AQPI) in ChP (Fig. 2e). This finding is

different from what was recently shown in ZIKV infected mouse brains, where ZIKV specifically targeted

pericytes [22].

To further investigate the subcellular localization of virus particles within ChP epithelial cells, we imaged

fourth ventricle ChP of *Ifnar*$^{-/-}$ brain using electron microscopy. Transmission electron microscopy (TEM)

of ChP epithelial cells, identified by the presence of cilia, revealed an extreme distortion of endoplasmic

reticulum (ER) membranes in infected tissue (Fig. 2f), in line with previous findings for other flaviviruses in

cell culture [23-26]. Higher resolution imaging revealed virus particles in the ER lumen (Fig. 2g, inset *i*) and

formation of viral replication complexes as bud-like invaginations of the ER membrane (Fig. 2g, inset *ii*).

Segmentation of volumetric images, taken by focused ion beam milling-scanning electron microscopy

(FIB-SEM), depict the 3D architecture of replication complexes (Fig. 2h-i). Replication vesicles were

clustered together within a dilated ER, and virions were detected both within the infected cell and

between two cells (visualized by green and blue; Fig. 2j), indicating active viral replication and viral

spread within the tissue. TEM and FIB-SEM support earlier observations, confirming productive infection

of epithelial cells of ChP in *Ifnar*$^{-/-}$ brains and provide insight into cytoarchitectural changes induced by

viral infection *in vivo.* Taken together, the observed differences between WT and *Ifnar*$^{-/-}$ mice indicate that

the IFN-I response efficiently restricts viral replication in the meninges, ventricles and ChP.

Neuronal circuit mapping and shift in cellular tropism of viral infection

Whole brain OPT-MRI enabled us to visualize, map and quantify viral brain distribution in distinct areas

(Fig. 3). Using the OCUM atlas, with 336 distinct anatomical regions (Fig. 3a), we first analyzed the viral

signal in the fourth ventricle (Fig. 3b-c) as proof of principle and saw that the viral signal in *Ifnar*$^{-/-}$ is

higher compared to WT (p=0.002). Next, we analyzed the viral signal in cortex and olfactory bulb. In WT

mice, viral signal was visually detected in entorhinal cortex, piriform area, and primary auditory cortex,

which are parts of the olfactory and auditory systems. These regions reside in GM (Fig. 3d, left panel),

which consists of neuronal cell bodies. In contrast, *Ifnar*$^{-/-}$ mice displayed a more widespread distribution

of viral infection, including GM and WM (Fig. 3d). In GM, we observed viral signal within the olfactory

bulb, entorhinal cortex, dorsal endopiriform nucleus and piriform area. In WM, we detected viral signal

within the olfactory limb of the anterior commissure, the lateral olfactory tract, anterior forceps of corpus

callosum, and supra-callosal WM. These WM infected regions are also part of the olfactory systems. Of

note, the lateral ventricle, subependymal layer of olfactory ventricle, and the olfactory limb of the anterior

commissure are all parts of the rostral migratory stream, which suggests specific LGTV infection of the

ventriculo–olfactory neurogenic system in *Ifnar*$^{-/-}$ brains. It should be noted, however, that large individual

differences in viral OPT signal were observed in both WT and *Ifnar*$^{-/-}$ mice (Supplementary Fig. 1c),

which, in combination with the relatively small sample size, was also reflected in the viral OPT

quantification. In that sense, not all brain regions displaying visual viral signal survived thresholding in

quantification (Supplementary Table 1). Nevertheless, OPT quantification confirmed the differences in

viral infection between WT and *Ifnar*$^{-/-}$ mice (Fig. 3e-l). In the cortex, WT mice were highly infected in the

primary auditory cortex (p=0.003), ventral secondary auditory cortex (p=0.002), temporal association area

(p=0.047) and composite entorhinal cortical region (p=0.033) (Fig. 3e-g and supplementary Fig. 2a).

However, when taking a closer look at the individual VOIs in the composite entorhinal region, we only

observe a significant difference in the medial entorhinal cortex (adjusted p=0.037) after multiple

comparison correction (Supplementary Fig. 2b). While *Ifnar*$^{-/-}$ mice were significantly higher infected in

the olfactory peduncle (p=0.016) (Fig.3h). *Ifnar*$^{-/-}$ mice displayed more extensive infection in the olfactory

bulb (p=0.001), subependymal zone of the olfactory ventricle (p=0.037), and the following WM tracts:

olfactory limb of anterior commissure (p=0.037) and lateral olfactory tract (p=0.015) (Fig. 3i-l). In the

composite hippocampal region, no infection could be detected in WT nor *Ifnar*$^{-/-}$ mice, and only one

hippocampal region CA2Or showed infection in WT mice (Supplementary Fig. 2c, d).

Cellular tropism in these infected brain regions was further investigated by confocal microscopy. In

sections containing areas of rostral migratory stream, we observed viral infection in immature neurons,

identified by doublecortin (DCX) immunoreactivity in both WT and *Ifnar*$^{-/-}$ brains (Fig. 4a). This indicates

that immature neurons are susceptible to viral infection irrespective of type I IFN signaling of the host. On

the other hand, no infection was found in astrocytes of either genotype (Fig. 4b). This is in contrast with

primary astrocytes in cell culture, which were found to be highly infected by several neurotropic

flaviviruses in the absence of IFNAR [12]. Focusing on the cerebrum the total number of infected cells was

similar between WT (1,631 ±526) and *Ifnar*$^{-/-}$ (1,947 ±213). The majority of infected cells in WT brains

were neurons, which have round-cell body morphology and reside in GM, whereas only 3.8 ± 4.8% of all

infected cells in cerebrum were positive for the microglial marker Iba1 (Fig. 4c-e). In *Ifnar*$^{-/-}$ brains,

surprisingly, we observed a shift in cellular tropism from neurons to microglia in which half (50 ± 5.3%) of

infected cells were Iba1-positive (Fig. 4 c-e). To further investigate this phenomenon in another system,

we isolated primary microglia from neonatal WT and *Ifnar*$^{-/-}$ brains and infected them *in vitro*.

Independently of IFNAR protein expression, microglia exhibited low susceptibility to LGTV infection *in*

vitro (Supplementary Fig. 3).

Taken together, while viral infection is restricted predominantly to neuronal cell bodies of sensory systems

of GM in WT, viral tropism expands to WM in the absence of IFN-I signaling. *In vivo,* cellular tropism was

also found to shift towards Iba1-positive cells in the absence of IFN-I signaling, however this tropism shift

is not observed in primary microglia from *Ifnar*$^{-/-}$ mice *in vitro*.

Single nuclei RNA sequencing analysis of virus-induced cellular responses

The increased susceptibility of *Ifnar*$^{-/-}$ Iba1-positive microglia *in vivo* compared to primary microglia *in*

vitro indicates that additional factors of the cellular milieu in the brain influence the tropism of LGTV

infection in the absence of IFN-I signaling. Alternatively, this tropism shift might be a result of Iba1

expressing infiltrating MØ that have been detected in the brain after LGTV infection [9]. To investigate

these hypotheses, we performed droplet-based single nuclei transcriptomic analysis (10x) of cerebral

cortex isolated from LGTV infected WT and *Ifnar*$^{-/-}$ mice as well as for uninfected controls (Fig. 5a). After

data processing and quality control, we obtained a total of 30,403 nuclei with a median of 1,173 genes

detected per nucleus. Using graph-based clustering and analysis of established marker genes, we

identified cells of each major cell type expected in cerebral cortex; *i.e.*, neurons, astrocytes, microglia,

oligodendrocytes, oligo progenitor cells (OPCs), endothelial cells, vascular leptomeningeal cells (VLMCs)

and pericytes (Fig. 5b-d, Supplementary Table 2). We also detected a small population of *Ttr* expressing

cells, most likely representing residual ChP epithelial cells [27]. Interestingly, we also identified cells

corresponding to CD8+NK cells in infected WT brains, but not in *Ifnar*$^{-/-}$ or uninfected controls (Fig. 5c).

We confirmed the prominent infiltration of CD8+ T-cells, but not CD4 T-cells or B-cells, in WT mice by

qPCR of whole cortex from infected mice (Supplementary Fig. 4a).

Viral RNA in our snRNA data was analyzed by alignment to the TP21 LGTV genome (NC_003690). In

WT mice, we detected most viral RNA in excitatory neurons and pericytes, while in *Ifnar*$^{-/-}$ mice, the

LGTV reads were mainly found in Microglia/machrophage (Micro/MØ) and ChP, and to a lesser extent

astrocytes (Fig. 5e). This supports our immunofluorescence data (Fig. 2, 4) and indicates that the removal

of ER-derived viral replication complexes from nuclei was not complete using the standard nuclear

isolation protocol for snRNAseq.

IFN-I response increases inflammation and immune cell infiltration

To understand the fundamental differences between WT and *Ifnar*[-/-] mice, we next analyzed differential

gene expression in all cell types at baseline (uninfected controls). Although we previously reported that

primary cell cultures of astrocytes from WT and *Ifnar*[-/-] mice display major gene expression differences at

baseline [12], our snRNAseq analysis presented very little differences *in vivo* (Supplementary Fig. 4b). This

suggests that baseline expression of entry factors or restriction factors in microglia is not driving the shift

in cellular tropism seen in *Ifnar*[-/-] mice. Subsequently, we evaluated the differentially expressed genes

(DEGs) in response to LGTV infection and performed gene set enrichment analysis (GSEA) on the

Reactome pathways [28]. In WT brains, we identified a large number of DEGs (2,846), with the majority of

expression changes in microglia, astrocytes, oligodendrocytes, VLMC and endothelial cells (Fig. 5e,

Supplementary Table 2). Although we detected a similar viral load in cerebral cortex of WT and *Ifnar*[-/-]

mice at endpoint (Supplementary Fig. 4c), the response to viral infection was markedly different in WT

and *Ifnar*[-/-], and a lower number of DEGs were identified (1,039) in *Ifnar*[-/-] (Fig. 5f, Supplementary Table

2). The majority of DEGs for both WT and *Ifnar*[-/-] were found to be upregulated (Supplementary Table 2).

We compared the overlap of upregulated DEGs amongst the five most responsive cell types based upon

number of DEGs (astro, micro/MØ, oligo, VLMCs and endo) (Fig. 5g). In WT mice, 73 genes,

corresponding to 9% of all upregulated DEGs, were commonly upregulated in these cells showing a

partial overlap and uniform cell response to infection. Of note, 57% of the genes were not shared, and

specific to each cell type and for astro 23%, micro/MØ 48%, oligo 21%, VLMCs 12% and endo 18% of the

DEGs were uniquely upregulated, emphasizing the advantage of a snRNAseq approach. In *Ifnar*[-/-],

however, only nine genes, corresponding to 3% of all upregulated DEGs were commonly upregulated,

indicating that the majority of responses are unique to each cell type.

We found a greater number of significant pathways in WT compared to *Ifnar*[−/−] (Fig. 5h, Supplementary

Table 2). In WT mice, there was a great overlap between the pathways upregulated in the different cells

and, as expected, the IFN signaling pathway was the most upregulated pathway in all cell types (Fig. 5i).

It has been reported that some cells are able to mount an IFN-independent upregulation of ISGs [29,30]. In

line with this, we also detected an IFN-independent upregulation of IFN stimulated genes in astrocytes

and VLMCs (Fig. 5i) in *Ifnar*[−/−] mice. Astrocytes upregulated 18 ISGs previously shown to be IFN

independently upregulated [30], and some with known antiviral activity e.g. *Rsad2* (viperin) [14,31], possibly

contributing to lower the susceptibility of these cells *in vivo* (Fig. 4b and Fig. 5e). However, this antiviral

IFN-independent response and were not detected in IFNAR deficient microglia.

IFNγ is known to compensate for lack of IFN-I, and can exert non-cytolytic antiviral activity against WNV,

Sindbis virus and measles virus infection in neurons [32-35]. Moreover, brains infected with VSV have been

shown to induce more *Ifng* in the absence of IFNAR signaling compared to WT [29]. Therefore, we

specifically evaluated expression changes of IFN-II (*Ifng*) in whole cortex of mice with qPCR. We found

high upregulation of *Ifng* in WT but not in *Ifnar*[−/−] after infection (Supplementary Fig. 4d). Analyzing the

cell types expressing *Ifng* and IFN-Is in WT, we found that particularly CD8+NK cells strongly expressed

Ifng while ChP, astrocytes and microglia were the main producers of *Ifnb* (Fig. 5j). One of the main

functions of IFN signaling is the upregulation of antiviral ISGs to limit viral replication and spread. We

detected *Rsad2* and *Ifitm3,* two ISGs with confirmed antiviral activity against TBEV [31,36], in all cell types in

WT mice, with the strongest expression in microglia, astrocytes, VLMCs, Endo and ChP, whereas only

the ChP expressed *Rsad2* and both ChP and Endo expressed *Ifitm3* in *Ifnar*[−/−] mice (Fig. 5k).

Since *Ifng* expressing CD8+NK cells might contribute to the increased inflammation in infected WT mice,

we evaluated the expression of known T cell chemoattractants (Supplementary Fig. 4e) and observed an

increase in *Ccl2, Ccl5 and Cxcl10* levels in infected WT compared to *Ifnar*[−/−] mice. Furthermore, VLMCs,

pericytes and endothelial cells of WT mice expressed high levels of *Cxcl9,* a chemokine dependent on

IFNγ for cerebral expression [37]. This further confirms the role of both IFN-I and IFN-II signaling in the

brain of these mice. A high expression of corresponding receptors *Ccr2* and *Ccr5* were found in micro/MØ

and CD8+NK cells and *Ccr1* in micro/MØ (Supplementary Fig. 4e). We performed sub-clustering of the

infiltrating CD8+NK cell population but were unable to obtain a clear separation between CD8+ and NK

cells (Fig. 6a), while flow cytometry confirmed two distinct populations of CD8+ T-cells (CD8$^+$) and NK

cells (CD335$^+$) in infected WT mice (Supplementary Fig. 5a). We found granzyme A (*Gzma*), granzyme B

(*Gzma*) and perforin (*Prf1*) to be highly upregulated in these cells in our data set (Supplementary Table

2), which are used by both CD8+ T cells and NK cells to kill target cells [38] and may contribute to

immunopathology.

Microglial activation and large infiltration of peripheral macrophages

We detected a major increase of microglia/MØ in WT mice upon infection (from 3.4% to 25%) (Fig. 5c).

This suggests that the inflammatory milieu in infected WT mice induce either proliferation of resident

microglia and/or infiltration of peripheral MØs. By a sub-clustering of the microglia/MØ population, we

identified, in addition to microglia (*Cxcr1*) and MØ (*Slfn4)*, a cluster of monocyte-like cells (*Ccl3*, *Ccl4,*

Il12b and *Tnf)* and a small number of neutrophils (*S100a9*, *S100a8* and *Clec4d*) (Fig. 6b, Supplementary

Table 2). In *Ifnar*$^{-/-}$ mice, we found predominantly microglia (Fig. 6c, left panel), while in WT, we observed

a large MØ population during infection (Fig. 6c, right panel). The cellular identity of microglia and MØ was

confirmed by expression of cell type specific marker genes [39] (Supplementary Fig. 5b) as well as

expression of *Tmem119* (Fig. 6d), a highly specific marker for resident microglia [40]. We used flow

cytometry to confirm the presence of infiltrating MØ (CD11b$^+$ CD45^{high}) in LGTV infected WT mice, which

were not seen in untreated controls or infected *Ifnar*$^{-/-}$ mice (Fig. 6e). Interestingly, although we only

detected a small population microglia in infected WT mice in our snRNAseq dataset, the numbers of

microglia did not change after infection, as detected with FACS analysis (Fig. 6e). Flow cytometry further

confirmed TMEM119 as a microglia specific marker (Fig. 6f) and confocal microscopy showed the

presence of infiltrating CD45$^+$ cells in infected WT (Fig. 6g).

We used immunohistochemistry to further investigate the activation and infection status of the

microglia/MØ in WT and *Ifnar*$^{-/-}$ mice. In WT mice, multiple activation states of uninfected microglia were

identified by microglial morphology (Fig. 6h). In *Ifnar*$^{-/-}$ mice, the morphology showed non-activated

microglia and the majority of LGTV infected cells were positive for TMEM119 (Fig. 6h). However, Iba1-

positive TMEM119-negative infected cells were occasionally detected (Fig. 6h). To better understand

what drives the differences in microglial activation between WT and *Ifnar*$^{-/-}$, we analyzed the cell-cell

communication in the snRNA dataset by Cellchat [41]. To look more specifically at microglia, the cells

annotated as MØ were separated before analysis. We found fewer overall interactions in infected *Ifnar*$^{-/-}$

mice compared to infected WT mice (Fig. 6i), partially explained by the absence of MØ and CD8+NK

cells. Looking at the specific numbers of unique ligand-receptor interactions in WT and *Ifnar*$^{-/-}$ going into

microglia cells during infection, the total number of interactions was reduced by 43% in *Ifnar*$^{-/-}$ (Fig. 6j and

Supplementary Table 2). These interactions are however very different and only 12% of interactions in

WT are shared with *Ifnar*$^{-/-}$. Together this suggests that the absence of the local inflammatory milieu (Fig.

5) and the specific cues from other cells that should activate the microglia are lost in *Ifnar*$^{-/-}$ thus

rendering them more susceptible to infection.

Discussion

While it is clear that the IFN-I response limits viral replication in the brain, viral distribution and cellular

tropism on a global whole-brain level have not been addressed, nor their underlying molecular

mechanisms. In this study, using multiple imaging modalities (ranging from cm to nm) and single nuclei

transcriptomics technology, we evaluated the impact of the IFN-I response on viral distribution within the

whole brain with high resolution MRI-guided anatomical precision and the transcriptional changes to

identify the molecular mechanism of viral pathogenesis. We show that type I IFN response specifically

limits LGTV replication in ChP and cerebral WM. We also show that the specific interplay between cells

and the local milieu *in vivo*, determine the susceptibility to viral infection. In the absence of IFN-I signaling,

the complete response to infection is hampered, microglia does not become activated thus rendering

them susceptible to infection.

To date, the details regarding viral brain infection and pathogenesis have been partially concealed by an inability to visualize viral infection on the whole organ level. OPT and LSFM are 3D imaging techniques for transparent mesoscopic-sized (mm-cm) tissue samples; and are therefore suitable for rodent whole brain imaging. These methods provide information at cellular resolution while still capturing the entire tissue in 3D and have been used in various applications for targeted imaging (see for example Alanentalo et al. 2007, Hahn et al. 2021 and Dodt et al. 2007 [42-44]). However, anatomical information obtained from optical imaging, based on tissue autofluorescence, is limited. MRI, conversely, is widely used for anatomical brain imaging since it has exquisite contrast and resolution [45]. Structural MRI is not suitable for imaging pathogens, but when combined with OPT, one could gain information on both viral distribution and its anatomical location. Several mouse brain templates are available. However, they are either based on histology [20,46] or, *in vivo* or *ex vivo* MRI *in situ* [47-49]. OPT images of whole brain viral signal are acquired *ex vivo* after tissue bleaching, de-hydration and optical clearing. These chemical treatments result in a certain degree of tissue shrinkage, which leads to misalignment when coregistering brain OPT signals to existing templates. Therefore, we used the OCUM brain template [21], specifically created from *ex vivo* MR images acquired after OPT pre-processing, yielding improved alignment and precise anatomical mapping of virus distribution in the brain.

Here, we studied the effect of the local IFN-I response on LGTV neurovirulence in the brain after intracranial infection. This model was chosen to eliminate all peripheral IFN effects and to provide equal amounts of virus and full access to the brain. OPT-MRI fusion images, together with LSFM, revealed widespread viral infection in olfactory bulb, rostral migratory stream, ventricles, ChP, GM and WM, and showed that whole brain infection patterns are influenced by the IFN-I response. Interestingly, at endpoint, meninges and ChP were only infected in *Ifnar*[−/−] brains. This is similar to Herpes Simplex virus I (HSV1) and ZIKV, where viral infection is detected in ChP of *Ifnar*[−/−] brains but not in WT mice [22,50]. This strongly suggests that the IFN-I response plays a major role in preventing viral infection in ChP. Of note, while we detected productive LGTV infection in ChP epithelial cells, a previous study found that ZIKV

mainly infects pericytes in ChP [22], indicating tropism differences exist among neurotropic flaviviruses. In

cerebrum, infection in both WT and *Ifnar*$^{-/-}$ was predominantly restricted to regions processing sensory

information, which has been reported previously in animal experiments with other neurotropic flaviviruses,

such as TBEV and JEV [51,52]. When comparing *Ifnar*$^{-/-}$ and WT visually, we observed that the olfactory

system was most infected and widespread in *Ifnar*$^{-/-}$ mice, with extensive virus signal in areas processing

sensory input from the olfactory bulb such as the granular cell layer of the olfactory bulb. These visual

observations were also confirmed by the atlas-based quantification of viral signal in the olfactory bulb.

Furthermore, VOI-based quantification revealed, in addition to the granular cell layer, clear viral infection

of other cell layers as well as in the lateral olfactory tract (Supplementary Table 1), which underlines the

advantage of atlas-based OPT analysis over sole-visual observations. While anosmia (loss of smell) is

not a common symptom in tick-borne encephalitis patients, nor has it been tested in animal models, a

case with anosmia post TBEV vaccination has been reported [53]. Although we could observe viral signal in

cortical areas involved in processing and integrating olfactory and gustatory information (piriform cortex,

endopiriform nucleus and agranular insula, all located in the orbitofrontal cortex) in some mice. These

regions did not survive thresholded quantification, most likely due to the high inter-individual differences in

both groups and the relatively small sample size. Nevertheless, endopiriform nucleus consists of

specialized cells with long axons projecting towards, among others, the entorhinal cortex, located at the

caudal end of the temporal lobe, another cortical area identified to be infected in our study, both visually

and quantitatively. Interestingly, OPT quantification showed higher infection in the composite entorhinal

region in WT when compared to *Ifnar*$^{-/-}$, which can be mainly attributed to the contribution of the medial

entorhinal cortex in the composite entorhinal region (Supplementary Fig. 2a, Supplementary Table 1),

and in the primary and secondary auditory cortex of WT mice. Dysfunction of the auditory processing

pathway in CNS can lead to sensorineural hearing loss and has been reported, although very rarely, after

HSV1 and WNV encephalitis in combination with other severe neurological complications [54-56]. Other

flaviviruses such as ZIKV, WNV, and JEV were described to target hippocampal regions, affecting

learning and memory [57-59]. We did not see hippocampal infection of LGTV visually, however, hippocampal

area CA20r showed infection in WT. It has also been shown by others that LGTV can cause impaired

memory function and anxiety–like behavior in WT mice [60].

Interestingly, we observed hyperintense lesions on T1-weighted images of infected brains, which is in line

with imaging data obtained from TBE patients, which predominantly show lesions in the basal ganglia,

thalami and cerebellum [61]. Interestingly, post-mortem IHC of TBEV brains showed the presence of virus in

these regions [62]. However, in our study, the cortical lesions detected on T1 images appeared to be much

more widespread than the viral signal detected by OPT. These observations suggest that CNS damage

cannot be explained only by the presence of viral infection in a particular area and suggest

neuroinflammation-induced lesions.

Single cell techniques are useful in studying the transcriptional response in specific cell types upon

infection or inflammation. In contrast to bulk RNA sequencing or microarray analysis which measures the

combined response of all cells, single cell techniques better capture cell heterogenicity and genes

expressed exclusively by small cell populations. In theory, the single nuclei isolation should only isolate

the nuclei from cells. However, we could detect enrichment of LGTV reads in specific populations of cells

(excitatory neurons and pericytes in WT and microglia and ChP in $Ifnar^{-/-}$). We show with

immunofluorescence that neurons are the main target in WT and microglia and ChP are infected in $Ifnar^{-/-}$

, the detected LGTV reads are likely not contamination from free-floating RNA. It is more likely that the

ER, which contain viral replication vesicles, is not completely detached from the nuclei during processing.

Similar to what has previously been shown for ZIKV [22] and JEV [63,64] we found LGTV reads in pericytes in

WT mice. For both ZIKV and JEV the infection of pericytes has been suggested to be important for the

virus crossing the BBB [65], as pericytes are crucial for the interaction between astrocytes and the

endothelia and thus BBB integrity [66]. In this study we are unable to draw conclusions about how LGTV

crosses the brain barriers to enter the CNS as we use intracranial route of infection. However, we do see

that depending on the IFN response the virus targets either pericytes in WT or ChP epithelial cells in the

$Ifnar^{-/-}$.

Several reports have shown IFN-I independent upregulation of ISGs upon infection [29,30,67], In our study

this was limited to only a few cell types. In contrast to what we previously showed for astrocytes cultured

in vitro [12], we did not detect a difference in baseline gene expression levels between WT and *Ifnar*[-/-] *in*

vivo. On the same note, we detected a strong tropism shift towards microglia in the absence of IFNAR

expression in mice, however, IFNAR deficient microglia in monoculture remained resistant to infection *in*

vitro. This might be explained by the inhibitory signals that microglia receive from other cells *in vivo* [68],

and the absence of these in primary monocultures might lead to the "priming" of the microglia to be in an

activated state. Others have shown that in order for microglia in the olfactory bulb to become activated

they need specific cues from neurons and astrocytes which are dependent of IFNAR expression[69],

however, the identity of these cues are not known. What we see in our data from the cortex is that; cells

in WT mice upregulate cytokines and chemokines that seem to specifically attract CD8+T cells and NK

cells but not CD4+ T cells to the brain. These cells secrete of IFNγ which is known to upregulate CCR1

on microglia [70]. In accordance with this we observe CCR1 upregulation in WT but not in *Ifnar*[-/-] microglia.

The CCL5 (RANTES)-CCR1 interaction is important for microglia activation [71]. Most cells of both

genotypes upregulate CCL5 after infection, which may also interact with the receptor CCR5 present on

microglia from both WT and *Ifnar*[-/-]. However, CCL5-CCR5 does not seem to be enough to activate

microglia as the majority of infected microglia in *Ifnar*[-/-] remained in a morphological non-activated state.

In contrast, the uninfected WT microglia show different stages of activation, which might protect them

from infection. Interestingly, the observed tropism shift and increase in Iba1-positive cellular infection by

LGTV in cortex is similar to what was observed in STING-deficient mice, infected with HSV-I [72].

We observed infection in both WM and GM in *Ifnar*[-/-] brains, while infection in WT brains was mostly

restricted to the latter. This restriction to GM might relate to the surprising shift in cellular tropism from

neurons in WT to Iba1-positive cells in *Ifnar*[-/-] mice. Interestingly, we did not detect Iba1-positive infected

cells in cerebellum nor the brain stem in *Ifnar*[-/-] mice, indicating location specific differences of Iba1-

positive cell populations. Interestingly, we observed an increase in the proportion of microglia/MØ in the

cortex of WT mice upon infection using snRNAseq. Sub-clustering of these cells showed a dominance of

infiltrating MØ. FACS analysis confirmed the presence of infiltrating MØ (CD11b$^+$, CD45^{high}) and resident

microglia (CD11b$^+$, CD45^{int}) in WT mice. We can only speculate why this population of microglia in WT

was relatively small the snRNAseq data set. One possibility is that activated microglia become fragile and

rupture during sample processing. In *Ifnar*$^{-/-}$ mice the microglia population remained constant and very

few MØ were detected. The majority of the infected Iba1-positive cells were TMEM119 positive, indicating

that resident microglia are targeted by the virus. Proper activation of microglia in olfactory bulb requires

IFNAR signaling in neurons and astrocytes but not in microglia [69]. In line with this, infection of cultured

primary microglia from cortex showed low susceptibility to LGTV infection, independent of IFNAR

expression. This discrepancy between *in vitro* and *in vivo* experimentation strongly indicates that cellular

communication between neurons, astrocytes and microglia, which produce a local milieu within the brain,

dictates the outcome and susceptibility of infection. Therefore, one should be cautious when extrapolating

in vitro findings to the *in vivo* setting.

Here, we investigated the effect of the local IFN-I response on tick-borne flavivirus neurotropism. Using

multimodal imaging, spanning a range from cm to nm, we were able to anatomically map viral distribution

and identify target cells where we could visualize active replication and the release of virions. We found

that the IFN-I response shapes viral distribution in various neuronal pathways and brain regions as well

as its cellular tropism. Together, these results suggest that cell types or cell populations that belong to

different brain areas may respond to viral infection in different ways, and that the response is highly

region- and cell-specific. Additionally, we used snRNAseq to analyze the cellular response following

infection and define the molecular mechanisms of pathogenesis. In WT mice, we found a strong IFN-I

signaling and inflammatory response with IFNγ expression, infiltration of MØ, CD8+ T cells and NK cells,

and activated microglia expressing *Ccr1*. In the absence of IFN-I signaling the cells are unable to respond

to viral infection and only astrocytes showed limited IFN-independent response. The lack of IFNγ

expression within the brain and absence of CCR1 on microglia may prevent their activation, thus

contributing to the shift in viral tropism to resident microglia which consequently became the main target

of infection. Collectively, our approach provides unprecedented insight into the outcome of viral infection,

the importance of IFNs and the molecular mechanisms of viral pathogenesis. Future studies are required

to systematically explore the temporal, spatial and cellular responses to viral infection to understand the

effects of the IFN response in full, and to improve our understanding of viral neuropathogenesis.

Methods

Animals

C57BL/6 WT mice and *Ifnar*[−/−] mice in C57BL/6 background were kindly provided by N.O. Gekara [8] Mice

were bred as homozygotes and maintained under specific pathogen-free conditions Animal experiments

were approved by the regional Animal Research Ethics Committee of Northern Norrland and by the

Swedish Board of Agriculture (ethical permits: A9-2018 and A41-2019), and all procedures were

performed according to their guidelines.

Viruses

LGTV strain TP21 was a kind gift from G. Dobler. LGTV stock was produced in VeroB4 cells, a kind gift

from G. Dobler [26] and harvested on day 3 post infection when cytopathic effects were apparent. Virus

supernatant was harvested on day 7 post-infection and titrated on VeroB4 cells using focus-forming

assay [26].

Virus infection model in the mouse

Animals (7- to 13-week-old, mixed gender) were either left untreated, mock-treated (PBS) or infected with

LGTV. After sedation with ketamine (100 µg/g body weight) and xylazine (5 µg/g body weight) or

Isoflurane, animals were intracranially inoculated with LGTV suspended in 20 µL PBS, 1,000 focus

forming units for Fig. 1-3 and 10,000 for Fig. 4-5. Infected mice were euthanized using oxygen deprivation

when they developed one severe symptom, such as: >20% weight loss, bilateral eye infection, diarrhea or

hind-limb paralysis; or when they developed three milder symptoms, such as: >15% weight loss,

unilateral eye infection, facial edema, ruffled fur or overt bradykinesia, and/or development of

stereotypies.

Whole-mount immunohistochemistry (IHC) and OPT

Following euthanasia, cardiac perfusion was performed using 20 mL PBS followed by 20 mL 4% w/v

paraformaldehyde (PFA) in PBS, after perfusion, brain was removed and further immersed in PFA for 2 h

before thoroughly washed by PBS. PFA-fixed brains were fluorescently immunolabeled with an antibody

against viral NS5 and processed for OPT imaging, as previously described [42,73]. Briefly, the brain was

dehydrated in a stepwise gradient of methanol (MeOH), permeabilized by repeated freeze–thawing in

MeOH at −80 °C and bleached in a solution (MeOH:H_2O_2:DMSO at 2:3:1) to quench tissue

autofluorescence. Specimen were rehydrated in TBST (50 mM Tris-HCl, pH 7.4, 150 mM NaCl, and 0.1%

v/v TritonX-100), blocked with 10% v/v normal goat serum (NGS) (#CL1200-100, Nordic Biosite,

Sweden), 5% v/v DMSO, and 0.01% w/v sodium azide in TBST at 37 °C for 48 h, and labeled with

primary (chicken anti-NS5; Supplementary Table 4)) and secondary (goat anti-chicken Alexa Fluor 680;

Supplementary Table 4) antibodies diluted in blocking buffer. They were incubated at 37 °C for 4 days at

each staining step. The stained tissue was mounted in 1.5% w/v low melting point SeaPlaque™ agarose

(#50101, Lonza, Switzerland) and optically cleared using a BABB solution (benzyl alcohol

(#1.09626.1000, Supelco, USA): benzyl benzoate (#10654752, Thermo Fisher Scientific, USA) at 1:2).

OPT imaging was performed with an in-house developed near-infrared OPT system (described in detail in

Eriksson et al. 2013 [73]), with a zoom factor of 1.6 or 2.0, that resulted in an isotropic voxel dimension of

16.5 μm^3 and 13.2 μm^3, respectively. To obtain specific fluorescent viral (NS5) signal (coupled with Alexa

Fluor 680), and tissue autofluorescence signals, OPT images were acquired at the following filter settings:

Ex: 665/45 nm, Em: 725/50 nm (exposure time: 7000 ms) and Ex: 425/60 nm, Em: 480 nm (exposure

time: 500 ms), respectively.

To increase the signal-to-noise ratio for NS5, the pixel intensity range of all images was adjusted to

display the minimum and maximum, and a contrast-limited adaptive histogram equalization (CLAHE)

algorithm with a tile size of 64 × 64 was applied to projection images acquired in the NS5 channel [74].

Tomographic reconstruction was performed using NRecon software (v.1.7.0.4, Skyscan microCT, Bruker,

Belgium) with additional misalignment compensation and ring artifact reduction. Image files were

converted to Imaris files (.ims) using the Imaris file converter (v9.5.1, Bitplane, UK). NS5 signal from all

imaged brains was adjusted to display at min = 0, max = 200, and gamma = 1.2, and colored using red

glow color scheme. The signal was superimposed onto the corresponding tissue autofluorescence image

using 3D iso-surface rendering in Imaris software (v9.5.1, Bitplane).

Light sheet fluorescent imaging

High-resolution images of ChP in the fourth ventricle of individual *Ifnar*$^{-/-}$ brains, previously scanned using

OPT, were acquired by LSFM. To compensate for potential photobleaching effects due to homogeneous

sample illumination during OPT acquisition, the sample was relabeled using both primary and secondary

antibodies (see above) and cleared in BABB without agarose mounting. The brain was then scanned

using an UltraMicroscope II (Miltenyi Biotec, Germany) with a 1× Olympus objective (PLAPO 2XC,

Olympus, Japan), coupled to an Olympus zoom body providing 0.63–6.3× magnification with a lens-

corrected dipping cap MVPLAPO 2×DC DBE objective (Olympus). For image acquisition, left and right

light sheets were merged with a 0.14 numerical aperture, which resulted in a light sheet z-thickness of

3.87 μm and 60% width, while using a 10–15 step blending dynamic focus across the field of view. Image

sections were generated by Imspector Pro software (v7.0124.0, LaVision Biotec Gmbh, Germany) and

stitched together using the TeraStitcher script (v9), implemented in Imspector Pro. Stitched images were

then converted into Imaris files (*.ims files) using the Imaris file converter (v9.5.1, Bitplane).

MRI acquisition

After OPT, two brains were rehydrated in TBST. T1-weighted images were then acquired at 9.4 Tesla

using a preclinical MR system (Bruker BioSpec 94/20, Bruker, Germany) equipped with a cryogenic RF

coil (MRI CryoProbe, Bruker) running Paravision 7.0 software (Bruker). Specifically, Modified Driven

Equilibrium Fourier Transform (MDEFT) sequence with 5 repetitions (TR: 3000 ms; TE: 3 ms; TI: 950 ms;

voxel dimension: $40 \times 40 \times 40mm^3$) was performed. Postprocessing of images involved the realignment

and averaging of individual repetitions using statistical parametric mapping (SPM8) (the Wellcome Trust

Centre for Neuroimaging, UCL, UK; www.fil.ion.ucl.ac.uk/spm) implemented in Matlab (R2014a, The

MathWorks Inc., USA).

Creation of OPT-MRI fusion images

OPT images with viral signal and autofluorescence signal were reconstructed in DICOM format using

NRecon software (v.1.7.0.4, Bruker) followed by their conversion into NifTi using PMOD VIEW tool (v.4.2,

PMOD Technologies Inc., Switzerland) or the dcm2nii tool in MRIcron software for OPT and MR images,

respectively. Coregistration and normalization of OPT with the OCUM MRI template was performed using

the toolbox SPMmouse in SPM8. All transformation matrices were calculated using individual tissue

autofluorescence from OPT images and applied to the corresponding viral OPT images [21]. The accuracy

of all transformations were assessed visually by two individual readers in the check registration tool of

SPM8. Fusion images of viral OPT signal were created for each individual brain using its own MRI and

with the OCUM template using the PMOD VIEW tool or Amira-Avizo software (v6.3.0, Thermo Fisher

Scientific) for 3D renderings. Finally, brain areas with viral signal were identified according to the OCUM

atlas.

Atlas-based OPT quantification

To assess the differences in viral infection between WT and *Ifnar*$^{-/-}$ mice in more detail and to provide a

metric for viral infection in distinct brain regions, a Volume of Interest (VOI)-based analysis was performed

using VOI analysis in the PMOD VIEW tool and the OCUM atlas, resulting in 336 automatically delineated

brain VOIs. Individual VOI metrics were calculated after voxel intensities were thresholded to remove

background and the highest pixel intensities were normalized to the average of the highest pixel

intensities for each subject to obtain relative VOI intensities. Afterwards, all individual VOI metrics of all

individual images (MOCK, WT and *Ifnar*[−/−]) were normalized to the individual VOI averages of the MOCK

group to obtain an arbitrary unit relative to MOCK. Thereafter, larger bilateral composite regions of

interest (ROIs) were created and the arbitrary units of all individual VOIs in a composite ROI were

summed to provide the total arbitrary unit in the composite ROI. Finally, unpaired t-tests were performed

in GraphPad Prism after F-test analysis for equal variances between groups. To analyze individual VOIs

within a composite ROI, multiple unpaired t-test with Holm-Sidak multiple comparison correction (alpha =

0.05) were performed.

IHC for brain slices

Following euthanasia, cardiac perfusion with 4% PFA and PBS was performed as described above

followed by immersion in 4% PFA at 4 °C overnight. PFA-fixed brain was washed in PBS, dehydrated

overnight in a 30% w/v sucrose solution, snap-frozen on dry ice in Optimal Cutting Temperature

medium (#361603E, VWR, USA), and stored at –80 °C until cryosectioning. The brain was sectioned

along the sagittal plane at 10 μm thickness using a rotatory microtome cryostat (Microm Microtome

Cryostat HM 500M, Microm, USA). Brain sections were permeabilized and blocked in 10% v/v NGS,

0.2% v/v TritonX-100, and 1% w/v bovine serum albumin in PBS for 1 h at room temperature (RT),

immunolabelled with primary and secondary antibodies (Supplementary Table 4) diluted in 2% v/v

NGS and 0.5% v/v TritonX-100 in PBS for overnight at 4 °C and 1 h at RT in the dark, respectively.

Confocal fluorescence microscopy was performed using a Zeiss 710 confocal microscope (Zeiss,

Germany) controlled by Zeiss Zen interface (v.14.0.19.201) with Axio Observer inverted microscope

equipped with Plan Apochromat 20×0.8, C-Apochromat 40×/1.2, and Plan Apochromat 63×/1.4

objective lens or a Leica SP8 Laser Scanning Confocal Microscope equipped with HC PL APO

20x/0.75 (Leica), HC PL APO 40x/1.3 or HC PL APO 63x/1.40 and Leica Application Suit X software

(LAS X, v.3.5.5, Leica). For quantification of microglial infection, a Leica SP8 Laser Scanning Confocal

Microscope equipped with HC PL APO 20x/0.75 (Leica) and Leica Application Suit X software (LAS X,

v.3.5.5, Leica) was used.

Quantification of microglial infection

Sagittal brain slices (2 *Ifnar*[−/−] and 3 WT mice, 6-8 sections per animal) were co-stained for NS5 and Iba1

and the infected microglia were quantified in Imaris (v. 9.7.2). Therefore, stitched confocal images of the

cerebrum were converted to .ims files using the Imaris file converter (v. 9.8.0). After background

subtraction, a colocalization channel, displaying only cells positive for both NS5 and Iba1, was built using

the "coloc" tool. Then, the "cell" function was applied to the NS5 and the colocalization channel to

calculate the percentage of infected microglia, using a cell background subtraction width of 3 μm. The

percentage of infected microglia was calculated for all slices and expressed as mean ± standard deviation

(SD). Statistical analysis was performed using an unpaired two-tailed t-test with Welch´s correction in

GraphPad Prism v9.3.1 (GraphPad Software Inc., San Diego, CA, USA). The level of significance was set

at $p<0.05$.

Fixation, resin embedding, and staining of tissue for electron microscopy

ChP from the brain of LGTV-infected *Ifnar*[−/−] mice were prepared for electron microscopy by cardiac

perfusion with 20 mL of 0.1 M phosphate buffer, followed by 20 mL of 2.5% w/v glutaraldehyde and 4%

w/v PFA in 0.1 M phosphate fixative solution. Fourth ventricle ChP was dissected and further immersed in

the same fixative solution for an additional 24 hours. The tissue was stained and subsequently embedded

in resin using the rOTO (reduced osmium tetroxide, thiocarbohydrazide, osmium tetroxide) method

{Deerinck, 2010 #194}. The tissue samples were placed in a solution of 1.5% w/v potassium ferrocyanide

and 2% w/v osmium tetroxide (OsO4), and then incubated in Pelco Biowave Pro+ (Pelco, Fresno, USA),

a microwave tissue processor ("the processor"), under vacuum for 14 min. After two rinses with MilliQ

water on the bench, the samples were washed twice with MilliQ water in the processor without vacuum

pressurization. Then, the samples were incubated in 1% w/v thiocarbohydrazide solution for 20 min. After

another MilliQ water rinse on the bench, the samples were again washed twice in the processor (no

vacuum). Next, the samples were placed in 2% w/v OsO_4 solution and run in the processor under vacuum

for 14 min, followed by an additional water and processor wash. The samples were placed in 1% w/v

uranyl acetate solution and run in the processor under vacuum for 12 min, followed by another water and

processor wash. The samples were then dehydrated in a stepwise ethanol gradient series:30%, 50%,

70%, 80%, 90%, 95%, and 100%, twice in the processor without vacuum. The dehydrated samples were

infiltrated with an increasing concentration of Durcupan ACM resin (Sigma-Aldrich) using the following

stepwise ratios of ethanol to Durcupan resin: 1:3, 1:1, and 3:1 in the processor for 3 min under vacuum.

The two final infiltration steps were performed in 100% resin. Finally, the samples were transferred to

tissue molds, and placed at 60 °C for 48 h for complete polymerization of the resin.

TEM

Using a Reichert UltraCut S ultramicrotome (Leica, Germany), resin-embedded samples were trimmed,

and 50 nm sections were cut using a diamond knife and placed onto copper slot grids. Resin-embedded

sections were imaged using a 120 kV Talos L120C transmission electron microscope (Thermo Fischer

Scientific) fitted with a LaB6 filament and Ceta 4k × 4k CMOS camera sensor. Images were acquired at

2600×, 8500×, and 36,000× magnification corresponding to a pixel size of 54.3, 17.0, and 4.1 Å/px,

respectively, at the specimen level. TEM images were analyzed by ImageJ software (NIH).

Volume imaging using FIB-SEM

Resin-embedded tissue blocks were trimmed, mounted on SEM stubs, and then coated with a 5 nm

platinum layer using a Q150T-ES sputter coater (Quorum Technologies, UK) before FIB-SEM volume

imaging. Data was acquired using Scios Dual beam microscope (FIB-SEM) (Thermo Fischer Scientific).

Electron beam imaging was acquired at 2 kV, 0.1 nA current, 1.9 × 1.9 nm pixel spacing, 7 mm working

distance, 10 μs acquisition time, and 3072 × 2048 resolution using a T1 detector. SEM images were

acquired every 20 nm. The working voltage of gallium ion beam was set at 30 kV, and 0.5 nA current was

used for FIB milling. The specimens were imaged at 5 × 5 μm block face and 5 μm depth. FIB milling and

SEM imaging were automated using the Auto Slice and View software (Thermo Fischer Scientific). SEM

volume images were aligned and reconstructed using ImageJ (NIH) with linear stack alignment, with SIFT

and MultiStackRegistration plugins [75,76]. Analysis and segmentation of SEM volume images was done

using Amira-Avizo software (v2020.3.1, Thermo Fisher Scientific).

Single nuclei isolation and sequencing

After euthanasia (day 5 for WT and day 3 for *Ifnar*$^{-/-}$), brains were collected in cold Hibernate A GlutaMAX medium (Thermo Fisher) without cardiac perfusion. For each condition one male and one female mouse were pooled and single nuclei were isolated from the cerebral cortex using the protocol modified from Gaublomme et al 2019 [77]. Cerebral cortexes from the hemisphere that did not receive needle injection from one male and one female mouse were manually dissected and homogenized in lysis buffer (10 mM Tris pH 7.4, 146 mM NaCl, 1 mM CaCl2, 21 mM MaCl2, 0.1% NP40 and 40 U/ml RNase inhibitor) using Dounce homogenizer for 10-20 times. Brain homogenates were washed with wash buffer (10mM Tris pH 7.4, 146 mM NaCl, 1mM CaCl2, 21 mM MaCl2, 0.01% BSA and 40 U/ml RNase inhibitor) and filter through 20 μm pre-separation filters (Miltenyi Biotec, 130-041-407) and were centrifuged with a swing bucket at 500xg, 4^{o}C for 5 min. The supernatant was removed and resuspended in the stain buffer (10 mM Tris pH 7.4, 146 mM NaCl, 1 mM CaCl$_2$, 21 mM MaCl$_2$, 2% BSA and 40 U/ml RNase inhibitor). Nuclei were stained with 10 μg/ml DAPI and subjected to sorting from cell debris using FACS (BD FACSAria III). The gate was set to sort for DAPI positive population (FSC 302, SSC 313 and BV421 313(log). The flow rate was adjusted to maintain about 10 nl/nuclei and with 80% efficiency. Sorted nuclei were transferred into low binding tubes and centrifuged using a swing bucket at 500xg, 4^{o}C for 5 min. The overall quality of sorted nuclei was assessed under a microscope. Overall, 80% viable nuclei were then diluted for snRNAseq processing, using 10x Genomics 3' v2 kit according to manufacturer's instructions. The target recovery cells for WT and *Ifnar*$^{-/-}$ conditions were 8,800 cells. Reverse transcription, cDNA amplification and library construction were performed, and libraries were sequenced on an Illumina 6000 platform according to the manufacturer's instruction.

Data processing and quality control snRNAseq

A custom reference genome was made by combining cellranger-arc-GRCh38-2020-A-2.0.0, NC_003690 (LGTV), and the neomycin gene sequence from pMC1neo (Addgene https://www.addgene.org/vector-database/3549/, from XhoI to BamHI, based on design in Müller *et al* [8]. Demultiplexed reads were processed using CellRanger v6.1.2 with options expect-cells = 8000. All data sets were filtered prior to

further processing. First, genes detected in fewer than 3 cells or nuclei with < 300 features were removed.

A threshold of < 10% mitochondrial reads were applied and reads for the lncRNA Malat1 were removed.

Potential doublets were addressed by i) removing cells with high number (> 5000) of detected features, ii)

removing nuclei with co-expression of female and male genes (*Xist* and *Eif2s3y*) and iii) following cell

annotation and subclustering of large cell types, removing distinct clusters showing co-expression of

multiple cell type specific markers (e.g. neuronal and glial markers). The relative expression levels of the

Neo gene present in *Ifnar*$^{-/-}$ [8] and LGTV, were used to quantitatively verify the sample identities.

Cell annotation

Data analysis was performed in R version 4.2.1 using the Seurat package version 4.1.0 [78]. Data were

normalized   (NormalizeData),   top   2000   variable   features   identified   for   each   data   set

(FindVariableFeatures) and data integrated using FindIntegrationAnchors and IntegrateData in Seurat.

The integrated expression matrix was scaled (ScaleData), subjected to principal component analysis

(RunPCR) and the first 30 dimensions used as input for RunUMAP and to construct a shared nearest

neighbor graph (SNN, FindNeighbors). Clustering was performed using FindClusters at resolution 1.

Following data integration and dimensional reduction/clustering, gene expression data (assay "RNA") was

scaled and centered (ScaleData). The positive differential expression of each other cluster compared to

all others (FindAllMarkers, test "wilcox") in combination with expression of established marker genes [79,80],

Allen Brain Atlas, https://portal.brain-map.org/) was used to assign preliminary identities (excitatory

neurons; neuron.ex, inhibatotry neurons; neuron.in, microglia/macrophages; micro/MØ, astrocytes; astro,

oligodendrocytes; oligo, oligodendrocyte progenitor cells; OPC, vascular leptomeningeal cells; VLMCs,

pericytes; peri, endothelial cells; endo, CD8+ T-cells and NK cells; CD8+NK, choroid plexus epithelial

cells; ChP). Low-quality cells were identified as clusters with a combination of i) few significant

differentially expressed genes and ii) high fraction of transcripts from mitochondrial and ribosomal genes

and removed manually. To distinguish between microglia and other infiltrating immune cell population,

this population was subsetted and the steps for dimension reduction, clustering and differential gene

expression repeated.

DEG analysis, GSEA and CellChat

Differentially expressed genes were identified using FindMarkers in Seurat, test MAST, applying a

threshold of $\log_2$ fold change ($\log_2$FC) >1 and adjusted p values (padj) <0.05. GSEA was performed using

the fgsea package version 1.22.0 in R [81]. Genes were ordered by average $\log_2$FC on the the MSigDB

Reactome pathways obtained using the misgdbr package version 7.5.1. Pathways with a Benjamini-

Hochberg corrected p-value <0.05 were considered significant. A comparison of mock (PBS) treated

samples with untreated samples showed little transcriptional response to the injection procedure as such.

Therefore, all further DEG analysis and GSEA was made comparing infected animals with uninfected

(mock and untreated). Cell-cell communication within infected data sets was analyzed by CellChat

version 1.6.1 [41]. A cutoff of 25% gene expression within group was applied and only cell types with more

than 10 cells were included. To specifically characterize communications involving microglia, cells

annotated as MΦ were separated from the micro/MΦ cluster prior to analysis.

Isolation and infection of primary cells

Microglia were isolated from the astrocyte cultures as previously described [82] and grown in DMEM,

supplemented with 10 % FBS, 100 U/mL of penicillin and 100 μg/mL streptomycin (Gibco), and mixed

glial-conditioned medium (1:1). Monolayers of microglia were infected with LGTV (MOI 1) for 1 h at 37°C

and 5% CO_2 before the inoculum was removed and replaced with fresh medium. Cell supernatant was

harvested at 72 h post infection and viral titers determined by focus forming assay as previously

described [26], and normalized to input control 3 hours post infection.

RNA extraction and RT-qPCR

Total RNA was extracted from cerebral cortexes from one hemisphere of infected mice were and viral

replication quantified by qPCR as previously described [9]. In short, the cerebral cortex was manually

dissected and homogenized in QIAzol Lysis Reagent (Quiagen) using 1.3 mm Chrome-Steel Beads

(BioSpec) and the tissue homogenizer FastPrep-24 (MP). The RNA was extracted using the Nucleo-Spin

RNA II kit (Macherey-Nagel). 1,000 ng of total RNA was used as input for cDNA synthesis using High-

capacity cDNA Reverse Transcription kit (Thermo Fisher). LGTV RNA was quantified using qPCRBIO

probe mix Hi-ROX (PCR Biosystems) and primers recognizing NS3 [15]. GAPDH was used as

housekeeping gene, detected by QuantiTect primer assay (QT01658692, Qiagen) and the qPCRBIO

SyGreen mix Hi-ROX (PCR Biosystems). All experiments were run on a StepOnePlus real-time PCR

system (Applied Biosystems).

Flow cytometry

After euthanasia (day 5 for WT and day 3 for *Ifnar*[-/-]) mice were perfused with 15 mL cold PBS and the

cerebral cortex collected in cold Hank's Balanced Salt Solution (HBSS). The tissue was homogenized

through a 70 μm cell strainer, digested with a collagenase solution (PBS, 500 μg/mL collagenase D, 10

μg/mL trypsin inhibitor TLCK, 10 μg/mL DNase I, 10 mM HEPES) for 20 min at 37°C. Cells were washed

in FACS buffer (PBS, 2% FBS, 2 mM EDTA), 1200 rpm, 5 min at 4°C and purified on a discontinuous 70

to 30% Percoll gradient by centrifugation at 800 xg for 20 min at room temperature. Cells were stained

with directly conjugated antibodies (Supplementary Table 4) diluted in FACS buffer for 30 min on ice,

washed twice in FACS buffer and analyzed on a ZE5 Cell Analyzer (BioRad). Analysis was performed

using FlowJo software (v10.0.7r2, BD)*.*

Acknowledgments

We thank G. Dobler (Bundeswehr Institute of Microbiology, Munich, Germany) for providing a stock of

LGTV strain TP21; A. Pletnev (Laboratory of Infectious Diseases, NIAID, NIH, Bethesda, MD, USA) for

providing the GFP-expressing LGTV infectious clone; and N.O. Gekara (Department of Molecular

Biosciences, Stockholm University, Stockholm, Sweden) for providing the *Ifnar*[-/-] mouse colony. We

thank M. Eriksson and C. Nord (UCMM, Umeå University, Umeå, Sweden) for OPT training, and J.

Gilthorpe (Department of Integrative Medical Biology, Umeå University, Umeå, Sweden) for invaluable

discussions. We also acknowledge Umeå Center for Microbial Research (UCMR); the Biochemical

Imaging Center at Umeå University (BICU), and the National Microscopy Infrastructure for microscopy

support; the Umeå Centre for Electron Microscopy (UCEM), a SciLifeLab National Cryo-EM facility for EM

support; the Small Animal Research and Imaging Facility (SARIF), and the Flow@CliMi platform at Umeå

University for their flow cytometry support. The Single Cell Core Facility @ Flemingsberg campus

(SICOF), Karolinska Institute, their technical advice is gratefully acknowledged. Illustration (Fig. 1a, 2b

and 4a) were made with BioRender.com

Funding

NC and ES were supported by the MIMS Excellence by Choice Postdoctoral Program under the

patronage of Emmanuelle Charpentier from the Kempe Foundation and Knut and Alice Wallenberg

Foundation, grant SMK-1532 and KAW2015.0284, respectively. LAC was supported by the Swedish

Research Council grant 2018-05851 and Wallenberg Centre for Molecular Medicine Umeå. U.A. was

supported by the Swedish Research Council grant 2017-01307, Umeå University Medical Faculty and the

Kempe foundations. DM was supported by Umeå University Medical Faculty, and AKÖ was supported by

the Swedish Research Council grants 2020-06224 and 2018-05851, and the Laboratory for Molecular

Infection Medicine Sweden (MIMS). JH is supported by the Swedish Research Council grant 2021-06602.

ISM is supported by the Umeå University Industrial Doctoral School.

Data collection with confocal microscopy and EM were also supported by the Biochemical Imaging Center

at Umeå University (BICU), and the National Microscopy Infrastructure (NMI; VR-RFI 2019-00217) and

the Umeå Centre for Electron Microscopy (UCEM), a SciLifeLab National Cryo-EM facility and part of NMI

(VR-RFI 2016-00968) grants from the Knut and Alice Wallenberg Foundation and Kempe Foundation,

respectively. The computations were performed using resources provided by the Swedish National

Infrastructure for Computing (SNIC) through the Uppsala Multidisciplinary Center for Advanced

Computational Science (UPPMAX) under project SNIC 2022/5-18.

Author's contributions

789 NC, ER, SMAW, ES, JZ, LAC, UA, DM, AKÖ designed the experiment, NC, SMAW, ER and EN

790 performed animal experiments, NC, SMAW, MH and EN acquired images and analyzed data from OPT,

791 LSFM and confocal microscopy, SMAW, FM and DM acquired images and analyzed data from MRI and

792 ES, JZ and LAC acquired images and analyzed data from EM. RL isolation, infection and analysis of

793 microglia *in vitro*. NC, JH, IMS processed samples for snRNAseq, ER, JH and GE analyzed snRNAseq

794 data. ER processed samples, ran the flow cytometry and analyzed the data. NC and AKÖ generated final

795 figures. LAC, UA, DM, JH and AKÖ supervised the experiments. NC, ER, SMAW, ES, LAC, UA, DM,

796 AKÖ, wrote and edited the manuscript. All authors revised the manuscript.

798 ***Competing interests***

799 Authors declare that they have no competing interests

801 ***Materials and Correspondence***

802 All data generated or analyzed during this study are included in this published article and its

803 supplementary information file. Raw image data can be requested from the corresponding authors with

804 reasonable means to transfer large data files. Raw single nuclei RNAseq data is available at

805 ArrayExpress #E-MTAB-12131. For the analysis, open-source algorithms were used as described in

806 STAR Methods and all computer codes used for the analyses are available at

807 https://github.com/ERosendal/LGTV_cortex_sn.

810 **Figures**

Fig. 1. Whole brain 3D imaging of viral infection. a A workflow for viral infection and whole brain imaging. **b** Mouse brain appearance in aqueous and in optical clearing solvent, BABB. **c** Volumetric 3D-render of the brain of mock and LGTV infected brains immunolabeled with anti-NS5 antibody (virus marker; red glow). The signal intensity was normalized within an individual brain and adjusted to identical minimum and maximum. For each image pair, the top and lateral views of the same specimen are shown. Five mice per group were analyzed. Representative images are shown here, and the remaining brains (n = 4) are shown in Supplementary Fig. 1b. Scale bar = 2,000 μm. **d** OPT-MRI fusion image created using viral OPT signal (red) with own MR scan of WT and *Ifnar*−/− brains. **e** 3D mapping of viral OPT signal with the OCUM template.

Fig. 2. IFN-I response restricts viral infection in meninges and ChP. a OPT-MRI mapping viral infection from OPT signal (red) onto the OCUM template (grey), focusing at the lateral, third and fourth ventricles. Images were shown in three representative coronal plains as indicated in the bottom panel. Representative images are shown, from n=5. **b** Confocal images of cortical slices showing meninges (yellow arrowheads) and endothelial cells (red arrowheads) immunolabeled with antibodies against viral NS5. Scale bars = 100 µm. **c** LSFM showing high-resolution image of the fourth ventricle ChP infected with LGTV, from a representative brain. (*i*) Orientation of the brain in the LSFM during image acquisition. (*ii*) Volumetric 3D render of LSFM of LGTV infected fourth ventricle ChP, immunolabeled with anti-NS5 antibodies (red glow). ChP was imaged in 3 tiles at 2.5× magnification and stitched together with 20% overlap. z = 2000 µm; scale bar = 300 µm. Solid-line square indicates the area imaged and dashed-line square shows the orientation of (*iii*). (*iii*) Tomographic section viewed from the YZ plane at 4× magnification. **d** Confocal images of infected cells in the fourth ventricle ChP. ChP (indicated by asterisk)

of WT and *Ifnar*$^{-/-}$ brains, were immunolabeled with antibodies against NS5, mature neuron (TUBB3) and

nucleus (DAPI; blue). Scale bars = 100 µm. **e** Maximum intensity projection of confocal z stack capturing

fourth ventricle ChP, immunolabeled with antibodies against NS5 and ChP epithelial cell (AQPI) Scale

bars = 20 µm. **f** Ultrastructure of virus replication complex in ChP of *Ifnar*$^{-/-}$ brain by electron microscopy.

Low-magnification TEM image of ChP epithelial cells with a heavily infected cell indicated by dilated ER

membrane. Black arrows indicate cilia on the apical side of ChP epithelial cells. Scale bar = 5 µm. **g** High-

magnification TEM of ChP epithelial cell in (e). Scale bar = 1 µm. Insets show high magnification images

of (*i*) virus particles and (*ii*) replication complexes within dilated ER. Scale bar = 200 nm. **h** FIB-SEM

volume imaging shows viral replication complexes within dilated ER. Scale bar = 500 nm. **i** Segmentation

image created from the 3D volume images in (g). Animation is shown in the Supplementary Movie 3.

Scale bar = 500 nm. **j** FIB-SEM slice-through volume image of two infected cells (blue and green). Scale

bar = 1 µm. Inset shows different z depths of the same volume. Boxes *i-iii* (left) show apparent released

virus particles between the cilia, also displayed as enlarged images (right, arrowheads). Scale bar = 100

nm. The confocal images represent n=3/genotype; 2 slices/brain, and the EM images represent n=2

brains; 3-5 technical replicates.

Fig. 3. Imaging and quantification of viral infection revealed a shift in infected regions shaped by the IFN-I response. a Axial and coronal representation of the OCUM brain template with its according atlas. **b** Typical example of an OPT-OCUM fusion displaying signal in the fourth ventricle (upper panel) and the VOI delineation based on the OCUM-atlas (lower panel). **c** Quantification of viral OPT signal in fourth ventricle of WT and *Ifnar*$^{-/-}$ mice. All values were normalized to the MOCK group and are expressed in arbitrary units (A.U.). **d** Anatomical mapping of infected cortical and olfactory bulb after coregistration of viral OPT signal (red) with OCUM brain template (grey). Images of WT and *Ifnar*$^{-/-}$

infected brains are shown in two representative axial plains. **e-l** Quantification of viral OPT signal in

representative VOIs or composite ROIs. * p=0.05, *** p=0.005, **** p=0.001

Fig. 4. Lack of IFNAR expression shifts the cellular tropisms from neurons to microglia in the

cortex of mice. a-c Maximum-intensity projection of confocal z-stack. The images were taken from

sagittal brain sections (10 µm) using confocal microscope (n =3 per genotype). Scale bars = 20 µm. **a**

Captured at the granule cell layer of olfactory bulb. The sections were immunolabeled using antibodies

against viral NS5, doublecortin (DCX) for immature neurons in the rostral migratory stream and nucleus

(DAPI; blue). **b** Captured within entorhinal cortex and piriform area. The sections were immunolabeled

with antibodies against viral NS5, GFAP for astrocyte and nucleus. **c** Captured within entorhinal cortex

and piriform area. The sections were immunolabeled using antibodies against viral NS5, Iba1 for

microglia, and nucleus (DAPI; blue). **d** Low magnification images show representative fields of the

sagittal slices of cerebral cortex stained with NS5 and Iba1 in WT and *Ifnar*$^{-/-}$ brains. Scale bars = 100

μm. **e** Percent of Iba1-positive infected cells of total number of infected cells in cortical slices. Statistical

analysis showed a significant difference (****, p < 0.0001) between the percentage infected Iba1-positive

cells in *Ifnar*$^{-/-}$ mice as compared to WT (n=2-3 per genotype and 6-8 slices per brain).

Fig. 5. Single nuclei RNA sequencing (snRNAseq) analysis of LGTV infection show strong IFN

response in the cortex of WT mice. a Schematic overview of the workflow. For each condition one male

and one female mouse were pooled. **b** Uniform Manifold Approximation and Projection (UMAP) of 30,403

nuclei captured by droplet based snRNAseq (10x) colored by assigned identities; excitatory neurons;

neuron.ex, inhibitory neurons; neuron.in, microglia/macrophages; micro/MØ, astrocytes; astro,

oligodendrocytes; oligo, oligodendrocyte progenitor cells; OPC, vascular leptomeningeal cells; VLMCs,

pericytes; peri, endothelial cells; endo, natural killer and CD8+ T-cells; CD8+NK, choroid plexus epithelial

cells; ChP. **c** Relative proportion of each cell type in uninfected and LGTV infected samples. **d** Dotplot

showing expression of canonical marker genes used to identify major cell types. **e** Dotplot showing LGTV

read counts for each cell type. **f** Number of DEGs ($log_2FC>1$, $p_{adj} < 0.05$) induced by LGTV infection in

WT or *Ifnar$^{-/-}$* mice. **g** Overlap in upregulated DEGs between five most reactive cell types, shown as a

modified UpSet plot for WT and *Ifnar$^{-/-}$* separately. **h** Number of significant Reactome pathways following

LGTV infection. **i** DotPlot showing examples of Reactome pathways altered by infection, color

corresponding to Normalized Enrichment Score and size corresponding to $-log_{10}$ adjusted p-value.

DotPlot showing expression of IFN-I and IFN-II (**j**) and ISGs (**k**) in infected data sets.

Fig. 6. IFN signaling is important for infiltration of immune cells and activation of microglia after

LGTV infection.

a Sub clustering of 144 nuclei of CD8+NK cell subset found in infected WT, colored by gene signatures

for CD8+ T cells (*Cd3g, Cd3d, Cd28, Cd8a, Cd8b1*) of NK cells (*Klrblc, Klrd1, Nrc1, Nkg7*). **b**

Subclustering of 2,363 nuclei of micro/MØ cell subset belonging to all datasets, colored by assigned

identities visualized by UMAP; Microglia, Macrophages (MØ), Monocyte-like cells and Neutrophils. **c**

UMAP from (b) showing nuclei from each condition separately. **d** DotPlot showing expression of *Aif1*

(Iba1) and *Tmem119* in cell subtypes. **e** Flow cytometry of brain leukocytes from the cerebral cortex of

WT and *Ifnar*$^{-/-}$ mice. Representative scatterplots and percentage microglia (CD11b$^+$, CD45^{int}) and MΦ

(CD11b$^+$, CD45^{high}). Data represent the mean and standard error of the mean (SEM), n=5 mice. **f**

Percentage of microglia and MΦ amongst TMEM119$^+$ cells, n = 20 (microglia) and 5 (MΦ). Maximum-

intensity projection of confocal z-stack captured within the sagittal slices of cortex (n=3 per group). The

sections were immunolabeled with antibodies against CD45 and TMEM119 (**g**) or viral NS5 (green), Iba1

(red) and TMEM119 (white) (**h**) in WT and *Ifnar*$^{-/-}$ brains. ¤ resting state microglia, # different activated

sates of resident microglia and * infiltrating macrophage. Scale bar = 60 µm (g) or 20 µm (h). **i** The

communication probability of interactions in infected WT and *Ifnar*$^{-/-}$ data sets and **j** number of unique

ligand-receptor interactions going from the different cells to microglia, analyzed by CellChat.

References

References

1   Sips, G. J., Wilschut, J. & Smit, J. M. Neuroinvasive flavivirus infections. *Rev Med Virol*

22, 69-87, doi:10.1002/rmv.712 (2012).

2   Gosztonyi, G. & Koprowski, H. The concept of neurotropism and selective vulnerability

("pathoclisis") in virus infections of the nervous system--a historical overview. *Curr Top*

Microbiol Immunol **253**, 1-13, doi:10.1007/978-3-662-10356-2_1 (2001).

3   Sejvar, J. J. The long-term outcomes of human West Nile virus infection. *Clin Infect Dis*

44, 1617-1624, doi:10.1086/518281 (2007).

4   Turtle, L., Griffiths, M. J. & Solomon, T. Encephalitis caused by flaviviruses. *QJM* **105**,

219-223, doi:10.1093/qjmed/hcs013 (2012).

5   Gould, E. A. & Solomon, T. Pathogenic flaviviruses. *Lancet* **371**, 500-509,

doi:10.1016/S0140-6736(08)60238-X (2008).

6   Ruzek, D. *et al.* Tick-borne encephalitis in Europe and Russia: Review of pathogenesis,

clinical features, therapy, and vaccines. *Antiviral Res* **164**, 23-51,

doi:10.1016/j.antiviral.2019.01.014 (2019).

7   Kubinski, M. *et al.* Tick-Borne Encephalitis Virus: A Quest for Better Vaccines against a

Virus on the Rise. *Vaccines (Basel)* **8**, doi:10.3390/vaccines8030451 (2020).

8   Muller, U. *et al.* Functional role of type I and type II interferons in antiviral defense.

Science **264**, 1918-1921, doi:10.1126/science.8009221 (1994).

9   Weber, E. *et al.* Type I interferon protects mice from fatal neurotropic infection with

Langat virus by systemic and local antiviral responses. *J Virol* **88**, 12202-12212,

doi:10.1128/JVI.01215-14 (2014).

10   Lazear, H. M., Pinto, A. K., Vogt, M. R., Gale, M., Jr. & Diamond, M. S. Beta interferon

950   controls West Nile virus infection and pathogenesis in mice. *J Virol* **85**, 7186-7194,

951   doi:10.1128/JVI.00396-11 (2011).

952   11   Samuel, M. A. & Diamond, M. S. Alpha/beta interferon protects against lethal West Nile

953   virus infection by restricting cellular tropism and enhancing neuronal survival. *J Virol* **79**,

954   13350-13361, doi:10.1128/JVI.79.21.13350-13361.2005 (2005).

955   12   Lindqvist, R. *et al.* Fast type I interferon response protects astrocytes from flavivirus

956   infection and virus-induced cytopathic effects. *J Neuroinflammation* **13**, 277,

957   doi:10.1186/s12974-016-0748-7 (2016).

958   13   Fares, M. *et al.* Pathological modeling of TBEV infection reveals differential innate

959   immune responses in human neurons and astrocytes that correlate with their

960   susceptibility to infection. *J Neuroinflammation* **17**, 76, doi:10.1186/s12974-020-01756-x

961   (2020).

962   14   Lindqvist, R., Kurhade, C., Gilthorpe, J. D. & Overby, A. K. Cell-type- and region-specific

963   restriction of neurotropic flavivirus infection by viperin. *J Neuroinflammation* **15**, 80,

964   doi:10.1186/s12974-018-1119-3 (2018).

965   15   Kurhade, C. *et al.* Type I Interferon response in olfactory bulb, the site of tick-borne

966   flavivirus accumulation, is primarily regulated by IPS-1. *J Neuroinflammation* **13**, 22,

967   doi:10.1186/s12974-016-0487-9 (2016).

968   16   Dionne, K. R., Galvin, J. M., Schittone, S. A., Clarke, P. & Tyler, K. L. Type I interferon

969   signaling limits reoviral tropism within the brain and prevents lethal systemic infection. *J*

970   *Neurovirol* **17**, 314-326, doi:10.1007/s13365-011-0038-1 (2011).

971   17   Zhao, L., Rose, K. M., Elliott, R., Van Rooijen, N. & Weiss, S. R. Cell-type-specific type I

972   interferon antagonism influences organ tropism of murine coronavirus. *J Virol* **85**, 10058-

973   10068, doi:10.1128/JVI.05075-11 (2011).

974 18 Sharpe, J. *et al.* Optical projection tomography as a tool for 3D microscopy and gene

975   expression studies. *Science* **296**, 541-545, doi:10.1126/science.1068206 (2002).

976 19 Maffioli, C., Grandgirard, D., Engler, O. & Leib, S. L. A tick-borne encephalitis model in

977   infant rats infected with langat virus. *J Neuropathol Exp Neurol* **73**, 1107-1115,

978   doi:10.1097/NEN.0000000000000131 (2014).

979 20 Wang, Q. *et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D

980   Reference Atlas. *Cell* **181**, 936-953 e920, doi:10.1016/j.cell.2020.04.007 (2020).

981 21 Willekens, S. M. A. *et al.* An MR-based brain template and atlas for optical projection

982   tomography and light sheet fluorescence microscopy. *BioRxiv.*,

983   doi:https://doi.org/10.1101/2022.11.14.516420 (2022).

984 22 Kim, J. *et al.* Zika virus infects pericytes in the choroid plexus and enters the central

985   nervous system through the blood-cerebrospinal fluid barrier. *PLoS Pathog* **16**,

986   e1008204, doi:10.1371/journal.ppat.1008204 (2020).

987 23 Welsch, S. *et al.* Composition and three-dimensional architecture of the dengue virus

988   replication and assembly sites. *Cell Host Microbe* **5**, 365-375,

989   doi:10.1016/j.chom.2009.03.007 (2009).

990 24 Miorin, L. *et al.* Three-dimensional architecture of tick-borne encephalitis virus replication

991   sites and trafficking of the replicated RNA. *J Virol* **87**, 6469-6481,

992   doi:10.1128/JVI.03456-12 (2013).

993 25 Mackenzie, J. M. & Westaway, E. G. Assembly and maturation of the flavivirus Kunjin

994   virus appear to occur in the rough endoplasmic reticulum and along the secretory

995   pathway, respectively. *J Virol* **75**, 10787-10799, doi:10.1128/JVI.75.22.10787-

996   10799.2001 (2001).

26   Overby, A. K., Popov, V. L., Niedrig, M. & Weber, F. Tick-borne encephalitis virus delays

interferon induction and hides its double-stranded RNA in intracellular membrane

vesicles. *J Virol* **84**, 8470-8483, doi:10.1128/JVI.00176-10 (2010).

27   Herbert, J. *et al.* Transthyretin: a choroid plexus-specific transport protein in human

brain. The 1986 S. Weir Mitchell award. *Neurology* **36**, 900-911,

doi:10.1212/wnl.36.7.900 (1986).

28   Gillespie, M. *et al.* The reactome pathway knowledgebase 2022. *Nucleic Acids Res* **50**,

D687-D692, doi:10.1093/nar/gkab1028 (2022).

29   Mishra, A. R., Byrareddy, S. N. & Nayak, D. IFN-I Independent Antiviral Immune

Response to Vesicular Stomatitis Virus Challenge in Mouse Brain. *Vaccines (Basel)* **8**,

doi:10.3390/vaccines8020326 (2020).

30   Cui, D., Espinola, E. E., Arora, K. & Brinton, M. A. Two Interferon-Stimulated Response

Elements Cooperatively Regulate Interferon-Stimulated Gene Expression in West Nile

Virus-Infected IFNAR(-/-) Mouse Embryo Fibroblasts. *J Virol* **95**, e0104021,

doi:10.1128/JVI.01040-21 (2021).

31   Panayiotou, C. *et al.* Viperin Restricts Zika Virus and Tick-Borne Encephalitis Virus

Replication by Targeting NS3 for Proteasomal Degradation. *J Virol* **92**,

doi:10.1128/JVI.02054-17 (2018).

32   Zhang, M. *et al.* Anti-West Nile virus activity of in vitro expanded human primary natural

killer cells. *BMC Immunol* **11**, 3, doi:10.1186/1471-2172-11-3 (2010).

33   Patterson, C. E., Lawrence, D. M., Echols, L. A. & Rall, G. F. Immune-mediated

protection from measles virus-induced central nervous system disease is noncytolytic

and gamma interferon dependent. *J Virol* **76**, 4497-4506, doi:10.1128/jvi.76.9.4497-

4506.2002 (2002).

34   Burdeinick-Kerr, R., Govindarajan, D. & Griffin, D. E. Noncytolytic clearance of sindbis

virus infection from neurons by gamma interferon is dependent on Jak/STAT signaling. *J*

Virol **83**, 3429-3435, doi:10.1128/JVI.02381-08 (2009).

35   Kang, S., Brown, H. M. & Hwang, S. Direct Antiviral Mechanisms of Interferon-Gamma.

Immune Netw **18**, e33, doi:10.4110/in.2018.18.e33 (2018).

36   Chmielewska, A. M. *et al.* The Role of IFITM Proteins in Tick-Borne Encephalitis Virus

Infection. *J Virol* **96**, e0113021, doi:10.1128/JVI.01130-21 (2022).

37   Ochiai, E. *et al.* CXCL9 is important for recruiting immune T cells into the brain and

inducing an accumulation of the T cells to the areas of tachyzoite proliferation to prevent

reactivation of chronic cerebral infection with Toxoplasma gondii. *Am J Pathol* **185**, 314-

324, doi:10.1016/j.ajpath.2014.10.003 (2015).

38   Trapani, J. A. Granzymes: a family of lymphocyte granule serine proteases. *Genome*

Biol **2**, REVIEWS3014, doi:10.1186/gb-2001-2-12-reviews3014 (2001).

39   DePaula-Silva, A. B. *et al.* Differential transcriptional profiles identify microglial- and

macrophage-specific gene markers expressed during virus-induced neuroinflammation.

J Neuroinflammation **16**, 152, doi:10.1186/s12974-019-1545-x (2019).

40   Satoh, J. *et al.* TMEM119 marks a subset of microglia in the human brain.

Neuropathology **36**, 39-49, doi:10.1111/neup.12235 (2016).

41   Jin, S. *et al.* Inference and analysis of cell-cell communication using CellChat. *Nat*

Commun **12**, 1088, doi:10.1038/s41467-021-21246-9 (2021).

42   Alanentalo, T. *et al.* Tomographic molecular imaging and 3D quantification within adult

mouse organs. *Nat Methods* **4**, 31-33, doi:10.1038/nmeth985 (2007).

43   Hahn, M. *et al.* 3D imaging of human organs with micrometer resolution - applied to the

endocrine pancreas. *Commun Biol* **4**, 1063, doi:10.1038/s42003-021-02589-x (2021).

44   Dodt, H. U. *et al.* Ultramicroscopy: three-dimensional visualization of neuronal networks

in the whole mouse brain. *Nat Methods* **4**, 331-336, doi:10.1038/nmeth1036 (2007).

45   Bocan, T. M., Panchal, R. G. & Bavari, S. Applications of in vivo imaging in the

evaluation of the pathophysiology of viral and bacterial infections and in development of

countermeasures to BSL3/4 pathogens. *Mol Imaging Biol* **17**, 4-17, doi:10.1007/s11307-

014-0759-7 (2015).

46   Sunkin, S. M. *et al.* Allen Brain Atlas: an integrated spatio-temporal portal for exploring

the central nervous system. *Nucleic Acids Res* **41**, D996-D1008,

doi:10.1093/nar/gks1042 (2013).

47   Hikishima, K. *et al.* In vivo microscopic voxel-based morphometry with a brain template

to characterize strain-specific structures in the mouse brain. *Sci Rep* **7**, 85,

doi:10.1038/s41598-017-00148-1 (2017).

48   Dorr, A. E., Lerch, J. P., Spring, S., Kabani, N. & Henkelman, R. M. High resolution

three-dimensional brain atlas using an average magnetic resonance image of 40 adult

C57Bl/6J mice. *Neuroimage* **42**, 60-69, doi:10.1016/j.neuroimage.2008.03.037 (2008).

49   Barriere, D. A. *et al.* Brain orchestration of pregnancy and maternal behavior in mice: A

longitudinal morphometric study. *Neuroimage* **230**, 117776,

doi:10.1016/j.neuroimage.2021.117776 (2021).

50   Wilcox, D. R., Folmsbee, S. S., Muller, W. J. & Longnecker, R. The Type I Interferon

Response Determines Differences in Choroid Plexus Susceptibility between Newborns

and Adults in Herpes Simplex Virus Encephalitis. *mBio* **7**, e00437-00416,

doi:10.1128/mBio.00437-16 (2016).

51   Stadtbaumer, K., Leschnik, M. W. & Nell, B. Tick-borne encephalitis virus as a possible

cause of optic neuritis in a dog. *Vet Ophthalmol* **7**, 271-277, doi:10.1111/j.1463-

5224.2004.04030.x (2004).

52   Fu, T. L., Ong, K. C., Tan, S. H. & Wong, K. T. Japanese Encephalitis Virus Infects the

Thalamus Early Followed by Sensory-Associated Cortex and Other Parts of the Central

and Peripheral Nervous Systems. *J Neuropathol Exp Neurol* **78**, 1160-1170,

doi:10.1093/jnen/nlz103 (2019).

53   Vodicka, J., Jelinkova, H. & Chrobok, V. Smell impairment after tick-borne encephalitis

vaccination: case report. *Vaccine* **28**, 886-888, doi:10.1016/j.vaccine.2009.11.012

(2010).

54   Westerberg, B. D., Atashband, S. & Kozak, F. K. A systematic review of the incidence of

sensorineural hearing loss in neonates exposed to Herpes simplex virus (HSV). *Int J*

Pediatr Otorhinolaryngol **72**, 931-937, doi:10.1016/j.ijporl.2008.03.001 (2008).

55   Kaga, K., Kaga, M., Tamai, F. & Shindo, M. Auditory agnosia in children after herpes

encephalitis. *Acta Otolaryngol* **123**, 232-235, doi:10.1080/00016480310000958 (2003).

56   Casetta, I. *et al.* West Nile virus neuroinvasive disease presenting with acute flaccid

paralysis and bilateral sensorineural hearing loss. *J Neurol* **258**, 1880-1881,

doi:10.1007/s00415-011-6011-3 (2011).

57   Figueiredo, C. P. *et al.* Zika virus replicates in adult human brain tissue and impairs

synapses and memory in mice. *Nat Commun* **10**, 3890, doi:10.1038/s41467-019-11866-

7 (2019).

58   Vasek, M. J. *et al.* A complement-microglial axis drives synapse loss during virus-

induced memory impairment. *Nature* **534**, 538-543, doi:10.1038/nature18283 (2016).

59   Chauhan, P. S., Khanna, V. K., Kalita, J. & Misra, U. K. Japanese Encephalitis Virus

Infection Results in Transient Dysfunction of Memory Learning and Cholinesterase

Inhibition. *Mol Neurobiol* **54**, 4705-4715, doi:10.1007/s12035-016-9963-6 (2017).

60 Cornelius, A. D. A. *et al.* Langat virus infection affects hippocampal neuron morphology

and function in mice without disease signs. *J Neuroinflammation* **17**, 278,

doi:10.1186/s12974-020-01951-w (2020).

61 Pichler, A. *et al.* Magnetic resonance imaging and clinical findings in adults with tick-

borne encephalitis. *J Neurol Sci* **375**, 266-269, doi:10.1016/j.jns.2017.02.003 (2017).

62 Gelpi, E. *et al.* Visualization of Central European tick-borne encephalitis infection in fatal

human cases. *J Neuropathol Exp Neurol* **64**, 506-512, doi:10.1093/jnen/64.6.506 (2005).

63 Butsabong, T., Felippe, M., Campagnolo, P. & Maringer, K. The emerging role of

perivascular cells (pericytes) in viral pathogenesis. *J Gen Virol* **102**,

doi:10.1099/jgv.0.001634 (2021).

64 Chang, C. Y. *et al.* Interplay of inflammatory gene expression in pericytes following

Japanese encephalitis virus infection. *Brain Behav Immun* **66**, 230-243,

doi:10.1016/j.bbi.2017.07.003 (2017).

65 Cle, M. *et al.* Zika Virus Infection Promotes Local Inflammation, Cell Adhesion Molecule

Upregulation, and Leukocyte Recruitment at the Blood-Brain Barrier. *mBio* **11**,

doi:10.1128/mBio.01183-20 (2020).

66 Armulik, A. *et al.* Pericytes regulate the blood-brain barrier. *Nature* **468**, 557-561,

doi:10.1038/nature09522 (2010).

67 Raaben, M., Groot Koerkamp, M. J., Rottier, P. J. & de Haan, C. A. Type I interferon

receptor-independent and -dependent host transcriptional responses to mouse hepatitis

coronavirus infection in vivo. *BMC Genomics* **10**, 350, doi:10.1186/1471-2164-10-350

(2009).

68 Timmerman, R., Burm, S. M. & Bajramovic, J. J. An Overview of in vitro Methods to

Study Microglia. *Front Cell Neurosci* **12**, 242, doi:10.3389/fncel.2018.00242 (2018).

69   Chhatbar, C. *et al.* Type I Interferon Receptor Signaling of Neurons and Astrocytes

Regulates Microglia Activation during Viral Encephalitis. *Cell Rep* **25**, 118-129 e114,

doi:10.1016/j.celrep.2018.09.003 (2018).

70   Bonecchi, R. *et al.* Up-regulation of CCR1 and CCR3 and induction of chemotaxis to CC

chemokines by IFN-gamma in human neutrophils. *J Immunol* **162**, 474-479 (1999).

71   Yan, J. *et al.* CCR1 Activation Promotes Neuroinflammation Through

CCR1/TPR1/ERK1/2 Signaling Pathway After Intracerebral Hemorrhage in Mice.

Neurotherapeutics **17**, 1170-1183, doi:10.1007/s13311-019-00821-5 (2020).

72   Reinert, L. S. *et al.* Sensing of HSV-1 by the cGAS-STING pathway in microglia

orchestrates antiviral defence in the CNS. *Nat Commun* **7**, 13348,

doi:10.1038/ncomms13348 (2016).

73   Eriksson, A. U. *et al.* Near infrared optical projection tomography for assessments of

beta-cell mass distribution in diabetes research. *J Vis Exp*, e50238, doi:10.3791/50238

(2013).

74   Hornblad, A., Cheddad, A. & Ahlgren, U. An improved protocol for optical projection

tomography imaging reveals lobular heterogeneities in pancreatic islet and beta-cell

mass distribution. *Islets* **3**, 204-208, doi:10.4161/isl.3.4.16417 (2011).

75   Saalfeld, S. *Linear stack alignment with SIFT. https://imagej.net/imagej-wiki-*

static/Linear_Stack_Alignment_with_SIFT, 2008).

76   Busse, B. & Miura, K. *MultiStackReg. https://biii.eu/multistackreg*, 2016).

77   Gaublomme, J. T. *et al.* Nuclei multiplexing with barcoded antibodies for single-nucleus

genomics. *Nat Commun* **10**, 2907, doi:10.1038/s41467-019-10756-2 (2019).

78   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587

e3529, doi:10.1016/j.cell.2021.04.048 (2021).

79   McKenzie, A. T. *et al.* Brain Cell Type Specific Gene Expression and Co-expression

Network Architectures. *Sci Rep* **8**, 8868, doi:10.1038/s41598-018-27293-5 (2018).

80   Vanlandewijck, M. *et al.* A molecular atlas of cell types and zonation in the brain

vasculature. *Nature* **554**, 475-480, doi:10.1038/nature25739 (2018).

81   Sergushichev, A. A. An algorithm for fast preranked gene set enrichment analysis using

cumulative statistic calculation. *BioRxiv*, doi:https://doi.org/10.1101/060012 (2016).

82   Saura, J., Tusell, J. M. & Serratosa, J. High-yield isolation of murine microglia by mild

trypsinization. *Glia* **44**, 183-189, doi:10.1002/glia.10274 (2003).

1149