Influence network model uncovers relations between biological processes and mutational signatures

Bayarbaatar Amgalan, Damian Wojtowicz, Yoo-Ah Kim*, Teresa M. Przytycka*

National Center for Biotechnology Information, National Library of Medicine, National Institutes of Health, Bethesda, MD 20894, USA

Abstract. There is a growing appreciation that mutagenic processes can be studied through the lenses of mutational signatures: characteristic mutation patterns attributed to individual mutagens. However, the link between mutagens and observed mutation patterns is not always obvious. In addition, mutational processes interact with each other, as well as with other cellular processes, in ways that are not fully understood. To gain insights into these relationships we developed a new approach, GenSigNet, that leverages a sparse partial correlation technique and a novel confounder-free higher moment-based approach for the construction of a directed network containing nodes representing genes and mutational signatures. When applied to cancer data, GenSigNet allowed to uncover important relations between mutational signatures and cellular processes including a relation between APOBEC hypermutation and activation of regulatory T Cells (Tregs), a relation between APOBEC mutations and changes in DNA conformation, and impact of homologous recombination deficiency on a clustered APOBEC mutations in breast cancer. GenSigNet also exposed a relation between the SBS8 signature of unknown aetiology and the nucleotide excision repair pathway.

GeneSigNet is freely available at https://github.com/ncbi/GeneSigNet.

Keywords: Mutational Signatures, networks, causality inference, mutational processes in cancer, APOBEC

1   Introduction

Traditionally, research in cancer genomics has been focused on the identification of cancer driving mutations that confer a growth advantage to cancer cells. However, since cancer cells emerge as a result of various mutagenic processes such as UV light or a faulty DNA repair mechanism, cancer genomes also accumulate numerous mutations with seemingly no direct roles in carcinogenesis. Nevertheless, these so-called passenger mutations can provide valuable information about mutagenic processes that cells have undergone. Different mutagenic processes leave characteristic mutation imprints on the cancer genome, called *mutational signatures*, facilitating the identification of specific mutagenic processes. In addition, different cancer genomes can experience different levels of exposure to a particular mutagenic process. This leads to a different number of mutations caused by such process. Hence, the *exposure* of a cancer genome to a given mutagenic process is measured by the number of mutations in the given genome that are attributed to the corresponding mutational signature. Starting from the pioneering work of Alexandrov et al. [1], several computational methods have been developed to decompose a cancer genome's mutation catalog (the set of all mutations in the genome) into mutational signatures and their exposures.

Elucidating mutagenic processes associated with mutational signatures is of fundamental importance for a better understanding of carcinogenesis and the design of novel cancer treatments. Some of these mutational signatures have already been linked to specific mutagenic processes [2,3]. For example, mutational processes can be caused by both intrinsic (e.g., DNA repair deficiency) and extrinsic (e.g., UV radiation, tobacco smoking) factors. However, the etiology of many signatures still remains unknown or not fully understood. In addition, there may be complicated interactions between mutagenic processes and other cellular processes. For instance, tobacco smoking is not only mutagenic itself but it is also believed to activate an immune response [4]. Conversely, a perturbation of some cellular processes such as DNA replication or repair pathways can also be mutagenic. Furthermore, mutagenic processes themselves have been known to interact with each other: homologous recombination deficiency (HRD) is typically accompanied by a mutational signature related to APOBEC activity [5,6]. Untangling the interactions between mutagenic processes as well as their interactions with cellular pathways is critical to fully understand biological processes underlying mutational signatures.

Gene expression provides the most accessible measurement of the activities of cellular processes. Therefore, interrogating the relation between gene expression and the exposure of a mutational signature can provide important clues on the etiology of the signature. For example, deficiencies in the activities of several genes such as MUTYH [7], ERCC2 [8], MSH6 [9], and FHIT [10] have been linked to specific signatures. In addition, a correlation between the expression of the APOBEC family of genes and the exposures of signatures SBS2 and SBS13 (so-called APOBEC mutational signatures) has frequently been observed [2,11]. Indeed, applying a clustering-based method to genome-wide expression and mutational signature data, Kim et al. identified coherently expressed groups of genes associated with specific combinations of mutational signatures providing interesting clues about these signatures [6]. However, such cluster-based analysis can only provide broadly defined associations between signature exposures and the activities of biological processes, lacking more mechanistic explanations.

To fill this gap, we introduce a network-based method, named GENESIGNET (Gene and Signature Influence Network Model), aiming to *uncover interactions among mutagenic and cellular processes associated with mutational signatures.* Utilizing gene expression and mutational signature data from cancer patients, GENESIGNET constructs a Gene-Signature Network (GSN) which provides directed relationships among two types of nodes – MutStates (Mutagenic States) and genes. MutState nodes are in one-to-one correspondence with mutational signatures and each MutState represents an ab-

stract (directly unobserved) cell state associated with the emergence of the corresponding muta-
tional signature (see Section 2.1 for a detailed description). Both types of nodes have patient-specific
activities: gene expression for nodes corresponding to genes and signature exposure for nodes cor-
responding to MutStates. These activities are utilized by GeneSigNet for inferring GSN's edges
(Fig. 1).

Inferring causal relationships between the activities of biological entities has been considered
in other biological applications, most prominently in gene regulatory network inference. However,
these approaches are typically based on perturbation data or prior information such as transcription
factors binding, protein-protein interaction networks, among others [12,13,14,15]. For example, gene
regulatory networks assume that causal relations originate from transcription factors to target genes.
In our setting, no perturbation data or prior knowledge is available, requiring GeneSigNet to
rely solely on the observed activities of nodes for inferring edges and their directions. To achieve
deeper mechanistic insights about relations between mutational signatures and cellular processes,
GeneSigNet focuses on inferring most informative edges by performing two complementary steps:
(i) a sparse partial correlation technique (SPCS) to obtain an initial weighted-directed network from
gene expression and exposure of mutational signatures and (ii) a higher moment-based strategy to
revise the initial network by orienting bidirectional edges. We note that some methods that infer gene
regulatory networks from gene expression data [16,17,18,19] may be applicable in our setting even
though they are typically employed in the context where directionality can be inferred from prior
knowledge of regulatory directions. Accordingly, we compared our method with GENIE3 which is
considered to be the best method in this class [18,20,21] and found that GeneSigNet outperforms
it.

We applied GeneSigNet to two cancer datasets, breast cancer and lung adenocarcinoma, se-
lected based on the number of patients with gene expression data and the complexity of their
mutation landscape. The relations inferred by the GeneSigNet model are overall consistent with
current knowledge but also include several interesting novel findings. In particular, the model sug-
gests a causative relation from the homologous recombination deficiency signature (SBS3) to a
clustered APOBEC mutation signature, and also linked Signature 8 (SBS8) to the NER pathway.
The latter connection is consistent with the recent findings based on an experimental study in mouse
[22]. In addition, GeneSigNet identified a novel relation between APOBEC hypermutation and
the activation of regulatory T cells which presents an important implication in immunotherapy, and
captured a relation of APOBEC signature (SBS2) with DNA conformation changes among other

Fig. 1. Gene-Signature Network (GSN). (A) GSN is a weighted-directed network consisting of two types of
nodes: genes (green circles) and, associated with signatures, MutStates (orange ovals). Edges of a GSN represent
inferred influences and might be either positive (red) or negative (blue). **(B)** Edges of a GSN are inferred based on
the activities of nodes (gene expression for genes and signature exposures for MutStates (top panel)). A directed
edge from a gene to a MutState represents a putative causal relation from the gene to the MutState. Analogously, a
directed edge from a MutState to a gene represents an inferred influence of the MutState on the gene.

findings. Our results demonstrate that GENESIGNET provides novel and important insights that were not accessible with existing methods.

2   Results

2.1   The GENESIGNET method

Gene-Signature Network.   The main idea behind GENESIGNET is a construction of a Gene-Signature Network (GSN), consisting of two types of nodes: nodes corresponding to genes and nodes corresponding to mutational signatures (Fig. 1A). The fact that expression of a gene influences expression of other genes is well accepted. Similarly, the exposure of a mutational signature can be seen as a measure of the activity of an endogenous mutagenic process or the activity of a biological process activated by an exogenous mutagen. As discussed in the introduction, such processes interact with other cellular processes influencing gene expression. This motivates the concept of *MutState* defined as an abstract representation of cellular state associated with a specific mutational signature. The activity of this state is measured as the exposure of the corresponding mutational signature. Statistical relations of such MutState with gene expression and other MutStates are expected to shed light on the genes and pathways contributing to or being influenced by the processes associated with the corresponding mutational signature. To this end, GENESIGNET infers directed edges between both types of nodes, utilizing patient-specific gene expression (for genes) and exposures of mutational signatures (for MutStates) as described below (Fig. 1B). Other than the difference in the definitions of node activities (and subsequent interpretation), the network inference algorithm does not distinguish between the two types of nodes.

A high level description of the GENESIGNET method.   GENESIGNET method constructs GSN in two main steps: (i) constructing a preliminary directed network using a sparse partial correlation selection (SPCS) and (ii) revising the initial network by resolving the directions of its bidirectional edges using a higher moment-based strategy. The workflow of the GENESIGNET method is shown in Fig. 2.

In the first step, given the input matrix describing the activities of genes and MutStates across samples, GENESIGNET constructs an initial weighted-directed graph using a sparse partial correlation selection (SPCS) (Fig. 2B). Specifically, considering each node as a target GENESIGNET uses SPCS to compute the weights of incoming effects from the other $n-1$ nodes. This SPCS step builds on the fact that partial correlations can be approximated by regression coefficients and utilizes a constraint on the l_1 norm enforcing the weights of the incoming edges to provide a sparse solution while avoiding over-fitting issues (see Supplementary Sections S1.1 and S1.2 for a detailed description). We note that since SPCS is applied to each node separately, GENESIGNET ensures a locally sparse solution rather than constraining the network edges globally.

In the second step, GENESIGNET refines the initial network by reducing the number of bidirectional edges. Specifically, the initial network obtained in the first step might still contain bidirectional edges due to the presence of confounding effects on the observational data. In this step, GENESIGNET attempts to resolve the directionality of bidirectional edges using a new *confounder-free higher moment-based strategy*. For any two nodes having potential effects on each other in the initial network (endpoints of a bidirectional edge), we first remove confounding effects due to the presence of the other $n-2$ variables from the observed activities of the two nodes obtaining residuals representing the remaining dependencies between the pair [23]. Under the confounder-free assumption, the causal variable can be distinguished from the affected variable by comparing the higher moments of the two residual distributions. Specifically, the affected variable is expected to be closer to normality than the causal factor, and the skewness and kurtosis are the higher moment statistics

Fig. 2. Workflow of GENESIGNET. (A) Input matrix X of node activities constructed by concatenating gene expression values (for genes) and signature exposures (for MutStates) across p samples (patients). **(B)** Given the input matrix X, we infer a network of n nodes. For each node, Sparse Partial Correlation Selection (SPCS) is used to simultaneously estimate the weights of incoming effects from the other $n-1$ nodes. **(C)** For each bidirectional edge (j, f), the residual vectors r_j and r_f corresponding to nodes j and f are obtained by removing effects of the $n-2$ nodes other than the two nodes of the considered edge. The non-zero weight w_{kf} obtained by SPCS denotes the strength of the confounding effect on node f coming from node k. The direction of a causal effect between the pair of nodes is determined based on higher moment statistics, skewness and kurtosis of residuals r_j and r_f. If both moments support the same direction then the edge corresponding to the opposite direction is removed, otherwise, both edges remain in the network. **(D)** Edge weights are normalized using an alternative scaling algorithm, and the final weighted-directed network is obtained as the output.

Fig. 3. Basic higher moment statistics (A) versus the confounder-free higher moment-based strategy (B). The experimentally confirmed regulatory relation is from CEBPB to MBD2. Without removing confounding effects from expressions of the two genes, the two higher moment statistics, skewness and kurtosis, provide contradictory decisions. After removing the confounding effects, the higher moment statistics provide the correct direction. The values in brown highlight the indication of causal factors.

used to measure the close-normality of distributions. If both moments support the same direction consistent with the stronger weight of the influence then the edge corresponding to the opposite direction is removed. Otherwise, both edges remain in the network (see Supplementary Section S1.3 for details). Fig. 3 provides a real example illustrating the necessity of removing confounding effects before applying the higher moment statistics to correctly indicate the causality direction. Before removing confounding effects the two higher moment statistics provided contradictory directions. (Fig. 3A). After removing the confounding effects, both statistics correctly indicate the causal direction (Fig. 3B). More examples based on experimentally confirmed regulatory directions [24] are provided in Fig. S2 Finally, a matrix normalization algorithm, alternate scaling [25], is used to bring the total incoming and outgoing effects of each node to the same range (see Supplementary Section S1.4 for details). A detailed description of GENESIGNET is provided in Supplementary Section S1.

2.2   Evaluation of the directionality inference

Inferring directionality solely based on node activities without perturbation data or additional information about the regulatory mechanism in biological processes is fundamentally challenging. As mentioned earlier, various methods have been developed for inferring gene regulatory networks (GRN) based on gene expression data. GENIE3 is considered one of the best performing method in this class [18,20,21]. The problem of inferring GRN is simpler than the problem of inferring GSN since the direction of edges in GRN can be decided based on the information about which genes are transcription factors and which are target genes. However, GENIE3 infers, for every node (a gene) in the network, weights (influence scores) from all other genes in the network to this gene. Thus, it constructs a fully connected, weighted, directed graph. Given this property of GENIE3 approach, we tested whether utilizing edge weights inferred by GENIE3 and selecting the direction corresponding to the heavier of the two opposing edges could outperform the strategy employed by GENESIGNET. We compared the performance of the methods on breast cancer (BRCA) and lung cancer (LUAD) data sets (see Supplementary Section S2 for details). To evaluate the inferred directionality, we used protein-DNA interactions from the ChEA database [24] as a gold standard after excluding all bidirectional edges and self-interactions, and only retaining the remaining directed edges. The performance was measured by computing accuracy and p-values were computed using a binomial test. GENESIGNET clearly outperformed GENIE3 (Table 1). GENIE3 approach resulted in a worse than random performance. We hypothesized that this behavior is due to selecting higher weight edge irrespective of the inferred interaction strength. This might create a noisy assignment if the assigned weights are very small in both directions. Subsequently, we modified this approach and considered only edges that are above a certain threshold. The threshold (0.0037 and 0.0017 for BRCA and LUAD, respectively) was selected to maximize the performance of GENIE3 on the test set to give it maximum benefit. The so-optimized GENIE3 performed significantly better than its original version but still worse than GENESIGNET, despite being optimized using the test set. A similar optimization of GENESIGNET provided only a modest improvement since the version without the additional learned parameter had already good performance. A detailed description for learning the threshold value is provided in Supplementary Section S1.3.

In addition, we tested the robustness of the confounder-free higher moment-based strategy by performing a bootstrap sampling which is described in Supplementary Section S1.3. The approach was highly reproducible as summarized in the section and Fig. S1 in Supplementary Information.

	Methods	GeneSigNet	GENIE3	GeneSigNet optimized	GENIE3 optimized
BRCA	Correct fraction	0.597	0.455	0.614	0.583
	p value	7.05e-06	1	3.1e-07	3.4e-03
	Correct directions	300	9,240	316	162
	Incorrect directions	202	11,052	202	116
LUAD	Correct fraction	0.559	0.505	0.571	0.557
	p value	4.2e-03	0.0731	5.4e-04	1.8e-03
	Correct directions	282	11,049	310	370
	Incorrect directions	222	10,833	233	294

Table 1. The evaluation of directionality inference. Using ChEA database as a gold standard, a binomial test was performed to evaluate the directionality inference by comparing the number of correctly assigned directions to the number of incorrect directions. 'Correct directions' is the number of inferred directions consistent with the ChEA database, whereas 'Incorrect directions' denotes the number of inconsistent inferences.

2.3   Insights from the analysis of breast cancer data

Mutational signatures in breast cancer and construction of the GSN.   We utilized breast cancer (BRCA) data collection obtained from ICGC which includes 266 cancer samples providing both whole-genome sequencing data and gene expression data (for details, see Supplementary Section S2.1). The breast cancer genomes harbor mutations mainly contributed by 6 COSMIC mutational signatures – SBS1, 2, 3, 5, 8, and 13. We further refined the mutational signatures based on mutation density and sample correlations. The mutations in BRCA are characterized by occurrences of short highly mutated regions whose origin is believed to be different than sparse mutations [6,26,11,27,28]. The information available from whole-genome sequencing allows for distinguishing these two types of mutation patterns and to treat such dense and sparse mutation regions differently. The post-processing of mutational signatures resulted in 6 signature groups that we use for subsequent analysis to construct the GSN – SBS1, APOBEC-C (clustered SBS2 and SBS13 corresponding to APOBEC hypermutation), APOBEC-D (SBS2 corresponding to disperse APOBEC mutations), DSB (SBS3 and clustered SBS8), SBS5, and SBS8D (dispersed SBS8). In addition to gene expressions and exposures of mutational signatures, we included a node indicating the binary status of homologous recombination deficiency (HRD) as it is assumed to lead to specific patterns of mutational signatures in BRCA [29]. We applied GeneSigNet to construct a GSN for genes, mutational signatures, and HRD status, and to find relations between these features.

Consistency of GeneSigNet results with current knowledge.   Many relations uncovered with GeneSigNet are consistent with our current knowledge on mutational signatures, confirming the validity of our method. In particular, it is well appreciated that homologous recombination (HR) plays an important role in the double-strand break (DSB) repair mechanism and that HR deficiency is associated with the DSB signature [30]. Indeed, our network correctly predicted a strong positive influence from HRD status to the DSB signature (Fig. 4). In addition, GeneSigNet identified the known negative impact of BRCA1 expression on the DSB signature which is also consistent with the role of BRCA1 in HRD [30]. Furthermore, GeneSigNet captured the impact of HRD on chromosome separation, reflecting the role of homologous recombination in maintaining genomic stability [31,32], and identified the association of APOBEC-D with telomere maintenance, consistent with the well recognized role of APOBEC mutagenesis in replication [33,34].

Interestingly, our method linked SBS8 to the nucleotide excision repair (NER) pathway (Fig. 4). The etiology of this signature has remained unknown until a recent experimental study linked it to the NER pathway as well [22]. This demonstrates the power of the GeneSigNet method to uncover non-obvious relationships.

Fig. 4. Subnetworks of GSN for BRCA centered on (induced by) MutStates associated with Homologous Recombination, APOBEC and SBS8. Edge and node colors are as in Fig. 1. In boxes, there are the names of the genes adjacent to a given MutState with edge weight cut-off ($|w_{ij}| \geq 0.01$). The genes in bold are discussed in more detail in the text. If the adjacent genes are enriched with specific GO pathways ($q-value < 0.01$) then only the pathway genes are provided in the box. The HRD status dominantly contributes to the mutation strength in the DSB repair MutState. Increased DSB exposure leads to an increase of the exposures of APOBEC-C and (indirectly) APOBEC-D MutStates. SBS8 mutations are linked to the deficiency of nucleotide excision repair. An extended network including all MutStates and an extended list of genes is provided in Fig. S3 in Supplementary Information.

Untangling the interactions between APOBEC and DSB processes. Previous studies speculated that APOBEC related mutational signatures can arise in multiple different scenarios. First, double-strand breaks (DSB) created by the homologous recombination deficiency (HRD) provide mutational opportunities for APOBEC enzymes to act on the ssDNA regions, resulting in clustered APOBEC mutations [35,36,28]. In another scenario, a recent study attributed APOBEC-mediated hypermutations to the normal activity of mismatch repair which also involves creating ssDNA regions, generating "fog" APOBEC mutations [27]. The complex interplay between APOBEC activities and other DNA repair mechanisms is yet to be elucidated.

Focusing on the interactions of APOBEC signatures with the other MutStates and genes, we observe that GENESIGNET supports a positive influence of the DSB on APOBEC-C MutState, consistent with the assumption that double-strand breaks provide an opportunity for APOBEC mutations. Additionally, our analysis reveals that the expression level of the APOBEC3B enzyme is associated with the strength of the DSB signature. Indeed, a previous study proposed that APOBEC3 proteins are recruited to DSB sites to participate in the DSB repair process [5]. Thus, DSB contributes to an increase in APOBEC-C strength by two different mechanisms: (i) increased mutation opportunity due to ssDNA created by DSB and (ii) increased mutation probability due to increased APOBEC3B expression. Note that increased APOBEC expression would also increase APOBEC mutations in the "fog" regions proposed in [27].

On the other hand, the activity of APOBEC-D is positively influenced by APOBEC-C activity, without direct relation to DSB. In fact, GENESIGNET inferred a negative influence from HR status

to APOBEC-D MutState, confirming that different mutagenic processes are involved in clustered and dispersed APOBEC mutations (Fig. 4).

APOBEC hypermutation activates regulatory T cells – implications for immunotherapy. Interestingly, GO enrichment analysis of the genes associated with APOBEC mutational signatures (genes influenced by APOBEC-C MutState) revealed significant enrichment in positive regulation of regulatory T cell differentiation (Fig. 4). Tumor cells with mutated DNA are likely to produce tumor-associated neoantigens (mutated peptides presented at their surface) that allow the immune system to recognize and destroy tumor cells. However, cells carrying a high mutation burden often develop mechanisms of immune tolerance involving activation of regulatory T cells (Tregs) to protect themselves from the destruction [37,38]. Tregs, a subtype of T cells that suppress the immune response, are important for maintaining cell homeostasis and self-tolerance but can also interfere with anti-tumor immune response [39]. The top three genes (FOXP3, BCL6, and LILRB2) positively influenced by APOBEC-C signature are all related to such inhibitory mechanism to immune response [40,41,42]. FOXP3 is a transcriptional regulator playing a crucial role in the inhibitory function of Tregs. BCL6 is also essential for the stability of Tregs that promotes tumor growth. LILRB2 is a receptor for class I MHC antigens and is involved in the down-regulation of the immune response and the development of immune tolerance.

Patients with cancers displaying a high mutation burden can benefit from immunotherapy [43]. In particular, the APOBEC mutational signature was identified as a potential predictive marker for immunotherapy response in some cancers [44,45]. However, an increased number of Tregs in a tumor may lead to resistance to immune checkpoint inhibitors [46,47]. Thus, our finding suggests that a combined strategy targeting Tregs in addition to immune checkpoint inhibitors would be most beneficial for a better outcome in APOBEC hypermutated breast cancer tumors.

2.4   Insights from the analysis of lung adenocarcinoma data

We next analyzed lung adenocarcinoma (LUAD) data using 466 cancer samples from the TCGA project. The exposure levels of 6 COSMIC mutational signatures (SBS1, 2, 4, 5, 13, and 40) present in the exome sequencing data were integrated with the RNAseq expression data of 2433 genes belonging to the DNA metabolic and immune system processes in Gene Ontology terms to uncover influence between signatures and genes (for details, see Supplementary Section S2.2).

GeneSigNet uncovers immune response due to smoking. Two prominent mutational signatures in LUAD, SBS4 and SBS5, are assumed to result from exogenous causes [6]. SBS4 is associated specifically with exposure to cigarette smoking in lungs. SBS5 is known to accompany the smoking signature but it is also present in many other cancer types. Previous studies suggested that cigarette smoking stimulates an inflammatory response [4]. Consistent with these findings, the genes identified by GeneSigNet as influenced by SBS4 and SBS5 MutStates are indeed enriched with immune response genes (Fig. 5).

GeneSigNet also identified the influence of the signatures SBS4 and SBS5 on two APOBEC signatures – SBS2 and SBS13. The APOBEC signatures are associated with immune response and this relationship is consistent with the previously proposed immune activation due to smoking exposure [48]. In addition, GeneSigNet correctly captured the association of SBS13 (consequently SBS2) with the expressions of APOBEC3B and APOBEC3A enzymes, and also identified the association of SBS13 with pyrimidine related catabolic processes, potentially reflecting the fact that SBS13 involves a pyrimidine to pyrimidine mutation (Fig. 5).

Finally, tobacco smoking is known to induce GPR15-expressing T cells; although the exact role of GPR15 in response to smoking is yet to be elucidated [49]. Therefore, we investigated

Fig. 5. Subnetwork of LUAD GSN centered on MutStates known to be related to smoking and APOBEC. The meaning of colors and boxes is the same as in Fig. 1 and 4. An extended network including all MutStates and an extended list of genes is provided in Fig. S4 in Supplementary Information.

Fig. 6. Subnetwork of Lung Adenocarcinoma GSN centered on the node representing GPR15 gene. Expression of the GPR15 gene contributes to the activation of immune responses. The meaning of edge and node colors, and boxes is the same as in Fig. 1 and 4. An extended network including all MutStates and an extended list of genes is provided in Fig. S4 in Supplementary Information.

whether the results of GENESIGNET provide additional insights into it. Consistently with previous studies, GENESIGNET inferred a strong association between GPR15 and SBS4 (without resolving the direction, see also Fig. S4 in Supplementary Information). Next, we analyzed the influence that GPR15 has on other nodes of the GSN network. The results of GENESIGNET suggest that GPR15 is involved in the negative regulation of several genes related to chemotaxis, including IL10, a cytokine with potent anti-inflammatory properties, and has a positive impact on lymphocyte migration and leukocyte mediated cytotoxicity (Fig. 6).

GENESIGNET points to the role of DNA geometric changes for APOBEC signature SBS2. As discussed earlier, APOBEC can only act on single-stranded DNA (ssDNA). Interestingly, one of the GO terms associated with SBS2 MutState identified by GENESIGNET is DNA geometric change (Fig. 5). DNA geometric changes are local changes of DNA conformation such as bulky DNA adducts (a type of DNA damage due to exposure to cigarette smoke) or DNA secondary structures such as ZDNA, cruciform, or quadruplex. Indeed, these structures often involve the formation of ssDNA regions which, in turn, provide mutation opportunities for APOBEC enzymes [50,51,52]. The formation of DNA secondary structures is often associated with DNA supercoiling - a form of

DNA stress that is resolved by Topoisomerase 1 (TOP1). Interestingly, GeneSigNet identified a negative influence of TOP1 expression on one of the genes (XPA) contributing to this GO term. This suggests a relation between DNA stress mediated by TOP1 and APOBEC activity.

3   Discussion

Elucidating the nature of mutagenic processes and their interactions with cellular processes is of fundamental importance for understating cancer etiology and guiding cancer therapy. Here, we propose GeneSigNet, a new network-based approach that infers the relation between gene expression and the strength of mutation patterns (signature exposures) allowing us to uncover the relations between signatures and processes involved in DNA repair and immune response among other cellular processes. Recognizing the limitations of the previous approaches, GeneSigNet relies on a construction of a sparse directed network. For each node (gene or MutState), it selects a sparse set of incoming edges representing dominating incoming effects so that their combination explains the activity of the node. Aiming to capture the most dominant influences, the method utilizes sparse partial correlation coefficients. In general, the inference of direction of influential relations from statistical dependencies without any support from prior knowledge about regulatory mechanisms is highly challenging. However, it has a potential advantage of capturing the real dependencies in the investigated condition instead of constraining the outcomes subject to the general prior information. GeneSigNet provides an important step toward this direction that is independent of the specific application considered in this study.

Overall the relations discovered by GeneSigNet are consistent with the current knowledge, boosting the confidence in the method's applicability. In addition, GeneSigNet provided several new biological insights concerning the relation between mutagenic processes and other cellular processes. For example, the uncovered relation between APOBEC hypermutation and activation of regulatory T-Cell can have an important implication in immunotherapy. We note that focusing on a sparse set of edges reduces the power of GO enrichment analysis and requires more specific biological knowledge for interpreting the results. Yet, this potential disadvantage is compensated by the compelling mechanistic insights provided by the method.

Acknowledgements

This study was supported by the Intramural Research Programs of the National Library of Medicine (NLM), National Institutes of Health, USA. We thank Jan Hoinka for his helpful comments on the manuscript.

References

1. L. B. Alexandrov, S. Nik-Zainal, D. C. Wedge, P. J. Campbell, and M. R. Stratton. Deciphering signatures of mutational processes operative in human cancer. *Cell Rep*, 3(1):246–259, Jan 2013.
2. L. B. Alexandrov, J. Kim, N. J. Haradhvala, M. N. Huang, A. W. Tian Ng, Y. Wu, A. Boot, K. R. Covington, D. A. Gordenin, E. N. Bergstrom, S. M. A. Islam, N. Lopez-Bigas, L. J. Klimczak, J. R. McPherson, S. Morganella, R. Sabarinathan, D. A. Wheeler, V. Mustonen, G. Getz, S. G. Rozen, M. R. Stratton, L. B. Alexandrov, E. N. Bergstrom, A. Boot, P. Boutros, K. Chan, K. R. Covington, A. Fujimoto, G. Getz, D. A. Gordenin, N. J. Haradhvala, M. N. Huang, S. M. A. Islam, M. Kazanov, J. Kim, L. J. Klimczak, N. Lopez-Bigas, M. Lawrence, I. Martincorena, J. R. McPherson, S. Morganella, V. Mustonen, H. Nakagawa, A. W. Tian Ng, P. Polak, S. Prokopec, S. A. Roberts, S. G. Rozen, R. Sabarinathan, N. Saini, T. Shibata, Y. Shiraishi, M. R. Stratton, B. T. Teh, I. Vázquez-García, D. A. Wheeler, Y. Wu, F. Yousif, and W. Yu. The repertoire of mutational signatures in human cancer. *Nature*, 578(7793):94–101, 02 2020.
3. Y. A. Kim, M. D. M. Leiserson, P. Moorjani, R. Sharan, D. Wojtowicz, and T. M. Przytycka. Mutational Signatures: From Methods to Mechanisms. *Annu Rev Biomed Data Sci*, 4:189–206, 07 2021.

4. L. B. Alexandrov, Y. S. Ju, K. Haase, P. Van Loo, I. Martincorena, S. Nik-Zainal, Y. Totoki, A. Fujimoto, H. Nakagawa, T. Shibata, P. J. Campbell, P. Vineis, D. H. Phillips, and M. R. Stratton. Mutational signatures associated with tobacco smoking in human cancer. *Science*, 354(6312):618–622, 11 2016.

5. R. Nowarski and M. Kotler. APOBEC3 cytidine deaminases in double-strand DNA break repair and cancer promotion. *Cancer Res*, 73(12):3494–3498, Jun 2013.

6. Y. A. Kim, D. Wojtowicz, R. Sarto Basso, I. Sason, W. Robinson, D. S. Hochbaum, M. D. M. Leiserson, R. Sharan, F. Vadin, and T. M. Przytycka. Network-based approaches elucidate differences within APOBEC and clock-like signatures in breast cancer. *Genome Med*, 12(1):52, 05 2020.

7. A. Viel, A. Bruselles, E. Meccia, M. Fornasarig, M. Quaia, V. Canzonieri, E. Policicchio, E. D. Urso, M. Agostini, M. Genuardi, E. Lucci-Cordisco, T. Venesio, A. Martayan, M. G. Diodoro, L. Sanchez-Mete, V. Stigliano, F. Mazzei, F. Grasso, A. Giuliani, M. Baiocchi, R. Maestro, G. Giannini, M. Tartaglia, L. B. Alexandrov, and M. Bignami. A Specific Mutational Signature Associated with DNA 8-Oxoguanine Persistence in MUTYH-defective Colorectal Cancer. *EBioMedicine*, 20:39–49, Jun 2017.

8. J. Kim, K. W. Mouw, P. Polak, L. Z. Braunstein, A. Kamburov, D. J. Kwiatkowski, J. E. Rosenberg, E. M. Van Allen, A. D'Andrea, and G. Getz. Somatic ERCC2 mutations are associated with a distinct genomic signature in urothelial tumors. *Nat Genet*, 48(6):600–606, 06 2016.

9. Xueqing Zou, Michel Owusu, Rebecca Harris, Stephen P Jackson, Joanna I Loizou, and Serena Nik-Zainal. Validating the concept of mutational signatures with isogenic cell models. *Nature communications*, 9(1):1–16, 2018.

10. S. Volinia, T. Druck, C. A. Paisie, M. S. Schrock, and K. Huebner. The ubiquitous 'cancer mutational signature' 5 occurs specifically in cancers with deleted FHIT alleles. *Oncotarget*, 8(60):102199–102211, Nov 2017.

11. D. Wojtowicz, I. Sason, X. Huang, Y. A. Kim, M. D. M. Leiserson, T. M. Przytycka, and R. Sharan. Hidden Markov models lead to higher resolution maps of mutation signature activity in cancer. *Genome Med*, 11(1):49, 07 2019.

12. D. Silverbush and R. Sharan. Network orientation via shortest paths. *Bioinformatics*, 30(10):1449–1455, May 2014.

13. A. Gitter, J. Klein-Seetharaman, A. Gupta, and Z. Bar-Joseph. Discovering pathways by orienting edges in protein interaction networks. *Nucleic Acids Res*, 39(4):e22, Mar 2011.

14. A. Vinayagam, U. Stelzl, R. Foulle, S. Plassmann, M. Zenkner, J. Timm, H. E. Assmus, M. A. Andrade-Navarro, and E. E. Wanker. A directed protein interaction network for investigating intracellular signal transduction. *Sci Signal*, 4(189):rs8, Sep 2011.

15. D. Silverbush and R. Sharan. A systematic approach to orient the human protein-protein interaction network. *Nat Commun*, 10(1):3015, 07 2019.

16. N. Friedman, M. Linial, I. Nachman, and D. Pe'er. Using Bayesian networks to analyze expression data. *J Comput Biol*, 7(3-4):601–620, 2000.

17. Y. Yuan, C. T. Li, and O. Windram. Directed partial correlation: inferring large-scale gene regulatory network through induced topology disruptions. *PLoS One*, 6(4):e16835, Apr 2011.

18. V. A. Huynh-Thu, A. Irrthum, L. Wehenkel, and P. Geurts. Inferring regulatory networks from expression data using tree-based methods. *PLoS One*, 5(9), Sep 2010.

19. R. Opgen-Rhein and K. Strimmer. From correlation to causation networks: a simple approximate learning algorithm and its application to high-dimensional plant gene expression data. *BMC Syst Biol*, 1:37, Aug 2007.

20. Y. Kang, D. Thieffry, and L. Cantini. Evaluating the Reproducibility of Single-Cell Gene Regulatory Network Inference Algorithms. *Front Genet*, 12:617282, 2021.

21. V. A. Huynh-Thu and P. Geurts. dynGENIE3: dynamical GENIE3 for the inference of gene networks from time series expression data. *Sci Rep*, 8(1):3384, 02 2018.

22. M. Jager, F. Blokzijl, E. Kuijk, J. Bertl, M. Vougioukalaki, R. Janssen, N. Besselink, S. Boymans, J. de Ligt, J. S. Pedersen, J. Hoeijmakers, J. Pothof, R. van Boxtel, and E. Cuppen. Deficiency of nucleotide excision repair is associated with mutational signature observed in cancer. *Genome Res*, 29(7):1067–1077, 07 2019.

23. Yasunori Fujikoshi, Vladimir V Ulyanov, and Ryoichi Shimizu. *Multivariate statistics: High-dimensional and large-sample approximations*, volume 760. John Wiley & Sons, 2011.

24. A. Lachmann, H. Xu, J. Krishnan, S. I. Berger, A. R. Mazloom, and A. Ma'ayan. ChEA: transcription factor regulation inferred from integrating genome-wide ChIP-X experiments. *Bioinformatics*, 26(19):2438–2444, Oct 2010.

25. Richard Sinkhorn and Paul Knopp. Concerning nonnegative matrices and doubly stochastic matrices. *Pacific Journal of Mathematics*, 21(2):343–348, 1967.

26. F. Supek and B. Lehner. Clustered Mutation Signatures Reveal that Error-Prone DNA Repair Targets Mutations to Active Genes. *Cell*, 170(3):534–547, Jul 2017.

27. D. Mas-Ponte and F. Supek. DNA mismatch repair promotes APOBEC3-mediated diffuse hypermutation in human cancers. *Nat Genet*, 52(9):958–968, 09 2020.

28. B. J. Taylor, S. Nik-Zainal, Y. L. Wu, L. A. Stebbings, K. Raine, P. J. Campbell, C. Rada, M. R. Stratton, and M. S. Neuberger. DNA deaminases induce break-associated mutation showers with implication of APOBEC3B and 3A in breast cancer kataegis. *Elife*, 2:e00534, Apr 2013.

29. H. Davies, D. Glodzik, S. Morganella, L. R. Yates, J. Staaf, X. Zou, M. Ramakrishna, S. Martin, S. Boyault, A. M. Sieuwerts, P. T. Simpson, T. A. King, K. Raine, J. E. Eyfjord, G. Kong, A. Borg, E. Birney, H. G. Stunnenberg, M. J. van de Vijver, A. L. Børresen-Dale, J. W. Martens, P. N. Span, S. R. Lakhani, A. Vincent-Salomon, C. Sotiriou, A. Tutt, A. M. Thompson, S. Van Laere, A. L. Richardson, A. Viari, P. J. Campbell, M. R. Stratton, and S. Nik-Zainal. HRDetect is a predictor of BRCA1 and BRCA2 deficiency based on mutational signatures. *Nat Med*, 23(4):517–525, Apr 2017.

30. R. Prakash, Y. Zhang, W. Feng, and M. Jasin. Homologous recombination and human health: the roles of BRCA1, BRCA2, and associated proteins. *Cold Spring Harb Perspect Biol*, 7(4):a016600, Apr 2015.

31. M. E. Moynahan and M. Jasin. Mitotic homologous recombination maintains genomic stability and suppresses tumorigenesis. *Nat Rev Mol Cell Biol*, 11(3):196–207, Mar 2010.

32. X. Li and W. D. Heyer. Homologous recombination in DNA repair and DNA damage tolerance. *Cell Res*, 18(1):99–113, Jan 2008.

33. V. B. Seplyarskiy, R. A. Soldatov, K. Y. Popadin, S. E. Antonarakis, G. A. Bazykin, and S. I. Nikolaev. APOBEC-induced mutations in human cancers are strongly enriched on the lagging DNA strand during replication. *Genome Res*, 26(2):174–182, Feb 2016.

34. N. Kanu, M. A. Cerone, G. Goh, L. P. Zalmas, J. Bartkova, M. Dietzen, N. McGranahan, R. Rogers, E. K. Law, I. Gromova, M. Kschischo, M. I. Walton, O. W. Rossanese, J. Bartek, R. S. Harris, S. Venkatesan, and C. Swanton. DNA replication stress mediates APOBEC3 family mutagenesis in breast cancer. *Genome Biol*, 17(1):185, 09 2016.

35. C. J. Sakofsky, N. Saini, L. J. Klimczak, K. Chan, E. P. Malc, P. A. Mieczkowski, A. B. Burkholder, D. Fargo, and D. A. Gordenin. Repair of multiple simultaneous double-strand breaks causes bursts of genome-wide clustered hypermutation. *PLoS Biol*, 17(9):e3000464, 09 2019.

36. K. Chan and D. A. Gordenin. Clusters of Multiple Mutations: Incidence and Molecular Mechanisms. *Annu Rev Genet*, 49:243–267, 2015.

37. T. A. Chan, M. Yarchoan, E. Jaffee, C. Swanton, S. A. Quezada, A. Stenzinger, and S. Peters. Development of tumor mutation burden as an immunotherapy biomarker: utility for the oncology clinic. *Ann Oncol*, 30(1):44–56, 01 2019.

38. S. Venkatesan, R. Rosenthal, N. Kanu, N. McGranahan, J. Bartek, S. A. Quezada, J. Hare, R. S. Harris, and C. Swanton. Perspective: APOBEC mutagenesis in drug resistance and immune escape in HIV and cancer evolution. *Ann Oncol*, 29(3):563–572, 03 2018.

39. A. Facciabene, G. T. Motz, and G. Coukos. T-regulatory cells: key players in tumor immune escape and angiogenesis. *Cancer Res*, 72(9):2162–2171, May 2012.

40. A. Y. Rudensky. Regulatory T cells and Foxp3. *Immunol Rev*, 241(1):260–268, May 2011.

41. Y. Chung, S. Tanaka, F. Chu, R. I. Nurieva, G. J. Martinez, S. Rawal, Y. H. Wang, H. Lim, J. M. Reynolds, X. H. Zhou, H. M. Fan, Z. M. Liu, S. S. Neelapu, and C. Dong. Follicular regulatory T cells expressing Foxp3 and Bcl-6 suppress germinal center reactions. *Nat Med*, 17(8):983–988, Jul 2011.

42. H. M. Chen, W. van der Touw, Y. S. Wang, K. Kang, S. Mai, J. Zhang, D. Alsina-Beauchamp, J. A. Duty, S. K. Mungamuri, B. Zhang, T. Moran, R. Flavell, S. Aaronson, H. M. Hu, H. Arase, S. Ramanathan, R. Flores, P. Y. Pan, and S. H. Chen. Blocking immunoinhibitory receptor LILRB2 reprograms tumor-associated myeloid cells and promotes antitumor immunity. *J Clin Invest*, 128(12):5647–5662, 12 2018.

43. L. M. Sholl, F. R. Hirsch, D. Hwang, J. Botling, F. Lopez-Rios, L. Bubendorf, M. Mino-Kenudson, A. C. Roden, M. B. Beasley, A. Borczuk, E. Brambilla, G. Chen, T. Y. Chou, J. H. Chung, W. A. Cooper, S. Dacic, S. Lantuejoul, D. Jain, D. Lin, Y. Minami, A. Moreira, A. G. Nicholson, M. Noguchi, M. Papotti, G. Pelosi, C. Poleri, N. Rekhtman, M. S. Tsao, E. Thunnissen, W. Travis, Y. Yatabe, A. Yoshida, J. B. Daigneault, A. Zehir, S. Peters, I. I. Wistuba, K. M. Kerr, and J. W. Longshore. The Promises and Challenges of Tumor Mutation Burden as an Immunotherapy Biomarker: A Perspective from the International Association for the Study of Lung Cancer Pathology Committee. *J Thorac Oncol*, 15(9):1409–1424, 09 2020.

44. S. Wang, M. Jia, Z. He, and X. S. Liu. APOBEC3B and APOBEC mutational signature as potential predictive markers for immunotherapy response in non-small cell lung cancer. *Oncogene*, 37(29):3924–3936, 07 2018.

45. D. L. Faden, F. Ding, Y. Lin, S. Zhai, F. Kuo, T. A. Chan, L. G. Morris, and R. L. Ferris. APOBEC mutagenesis is tightly linked to the immune landscape and immunotherapy biomarkers in head and neck squamous cell carcinoma. *Oral Oncol*, 96:140–147, 09 2019.

46. R. Saleh and E. Elkord. Treg-mediated acquired resistance to immune checkpoint inhibitors. *Cancer Lett*, 457:168–179, 08 2019.

47. D. R. Principe, L. Chiec, N. A. Mohindra, and H. G. Munshi. Regulatory T-Cells as an Emerging Barrier to Immune Checkpoint Inhibition in Lung Cancer. *Front Oncol*, 11:684098, 2021.

48. P. A. Patriarca, W. H. Foege, and T. A. Swartz. Progress in polio eradication. *Lancet*, 342(8885):1461–1464, Dec 1993.

49. S. Kõks and G. Kõks. Activation of GPR15 and its involvement in the biological effects of smoking. *Exp Biol Med (Maywood)*, 242(11):1207–1212, 06 2017.

50. F. Kouzine, D. Wojtowicz, L. Baranello, A. Yamane, S. Nelson, W. Resch, K. R. Kieffer-Kwon, C. J. Benham, R. Casellas, T. M. Przytycka, and D. Levens. Permanganate/S1 Nuclease Footprinting Reveals Non-B DNA Structures with Regulatory Potential across a Mammalian Genome. *Cell Syst*, 4(3):344–356, 03 2017.

51. X. Zou, S. Morganella, D. Glodzik, H. Davies, Y. Li, M. R. Stratton, and S. Nik-Zainal. Short inverted repeats contribute to localized mutability in human somatic cells. *Nucleic Acids Res*, 45(19):11213–11221, Nov 2017.

52. W. M. Guiblet, M. A. Cremona, R. S. Harris, D. Chen, K. A. Eckert, F. Chiaromonte, Y. F. Huang, and K. D. Makova. Non-B DNA: a major contributor to small- and large-scale variation in nucleotide substitution frequencies across the genome. *Nucleic Acids Res*, 49(3):1497–1516, 02 2021.

53. M Tsagris. Bayesian network learning with the PC algorithm: an improved and correct variation. *Applied Artificial Intelligence*, 33(2):101–123, Oct 2019.

54. M. Scutari, C. Vitolo, and A. Tucker. Learning Bayesian networks from big data with greedy search: computational complexity and efficient implementation. *Statistics and Computing*, 9(5):1095–1108, Feb 2019.

55. Finn V Jensen and Thomas Dyhre Nielsen. *Bayesian networks and decision graphs*, volume 2. Springer, 2007.

56. J. Peng, P. Wang, N. Zhou, and J. Zhu. Partial Correlation Estimation by Joint Sparse Regression Models. *J Am Stat Assoc*, 104(486):735–746, Jun 2009.

57. D. Yu, J. Lim, X. Wang, F. Liang, and G. Xiao. Enhanced construction of gene regulatory networks using hub gene information. *BMC Bioinformatics*, 18(1):186, Mar 2017.

58. Robert Tibshirani. Regression shrinkage and selection via the lasso. *Journal of the Royal Statistical Society: Series B (Methodological)*, 58(1):267–288, 1996.

59. Hirotugu Akaike. A new look at the statistical model identification. *IEEE transactions on automatic control*, 19(6):716–723, 1974.

60. Judea Pearl. The seven tools of causal inference, with reflections on machine learning. *Communications of the ACM*, 62(3):54–60, 2019.

61. T. Chu, C. Glymour, R. Scheines, and P. Spirtes. A statistical problem for inference to regulatory structure from associations of gene expression measurements with microarrays. *Bioinformatics*, 19(9):1147–1152, Jun 2003.

62. Clark Glymour, Kun Zhang, and Peter Spirtes. Review of causal discovery methods based on graphical models. *Frontiers in genetics*, 10:524, 2019.

63. Yadolah Dodge and Iraj Yadegari. On direction of dependence. *Metrika*, 72(1):139–150, 2010.

64. W. Wiedermann and J. Sebastian. Direction Dependence Analysis in the Presence of Confounders: Applications to Linear Mediation Models Using Observational Data. *Multivariate Behav Res*, 55(4):495–515, 2020.

65. S. Nik-Zainal, H. Davies, J. Staaf, M. Ramakrishna, D. Glodzik, X. Zou, I. Martincorena, L. B. Alexandrov, S. Martin, D. C. Wedge, P. Van Loo, Y. S. Ju, M. Smid, A. B. Brinkman, S. Morganella, M. R. Aure, O. C. Lingjærde, A. Langerød, M. Ringnér, S. M. Ahn, S. Boyault, J. E. Brock, A. Broeks, A. Butler, C. Desmedt, L. Dirix, S. Dronov, A. Fatima, J. A. Foekens, M. Gerstung, G. K. Hooijer, S. J. Jang, D. R. Jones, H. Y. Kim, T. A. King, S. Krishnamurthy, H. J. Lee, J. Y. Lee, Y. Li, S. McLaren, A. Menzies, V. Mustonen, S. O'Meara, I. Pauporté, X. Pivot, C. A. Purdie, K. Raine, K. Ramakrishnan, F. G. Rodríguez-González, G. Romieu, A. M. Sieuwerts, P. T. Simpson, R. Shepherd, L. Stebbings, O. A. Stefansson, J. Teague, S. Tommasi, I. Treilleux, G. G. Van den Eynden, P. Vermeulen, A. Vincent-Salomon, L. Yates, C. Caldas, L. van't Veer, A. Tutt, S. Knappskog, B. K. Tan, J. Jonkers, Å. Borg, N. T. Ueno, C. Sotiriou, A. Viari, P. A. Futreal, P. J. Campbell, P. N. Span, S. Van Laere, S. R. Lakhani, J. E. Eyfjord, A. M. Thompson, E. Birney, H. G. Stunnenberg, M. J. van de Vijver, J. W. Martens, A. L. Børresen-Dale, A. L. Richardson, G. Kong, G. Thomas, and M. R. Stratton. Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature*, 534(7605):47–54, 06 2016.

66. S. et al Nik-Zainal. Author Correction: Landscape of somatic mutations in 560 breast cancer whole-genome sequences. *Nature*, 566(7742):E1, 02 2019.

67. Xiaoqing Huang, Damian Wojtowicz, and Teresa M Przytycka. Detecting presence of mutational signatures in cancer with confidence. *Bioinformatics*, 34(2):330–37, 2018.

Supplementary Information

S1   Methods

GeneSigNet constructs a sparse network representing the causal dependencies among nodes in a Gene-Signature Network (GSN). The nodes corresponding to genes and MutStates (mutational signatures) are considered random variables and the activities of the variables are denoted as expressions of genes and exposures of mutational signatures across patients (samples). Given an input matrix describing the activities of variables, our network construction method consists of two complementary steps: (i) a sparse partial correlation technique (SPCS) to obtain an initial weighted-directed network by estimating statistical dependencies (edges) between variables (nodes) and (ii) a higher moment-based strategy to refine the initial network by orienting bidirectional edge. A high-level description of GeneSigNet is provided in Section 2.1 in the main manuscript, and the detailed description and related literature are presented below.

S1.1   A sparse partial correlation selection (SPCS)

In the proposed model, the network inference algorithm does not distinguish between the two types of nodes (genes and MutStates) and infers causal dependencies between variables based on their observed activities. In this setting, correlation networks are widely used to explore and visualize dependencies in high-dimensional data. However, without assuming prior knowledge, an ordinary correlation itself provides no means to distinguish between causal and affected factors in underlying causal processes [19].

Bayesian networks can be used to infer causal relations of nodes representing their local conditional dependencies via a directed acyclic graph (DAG) [16,53,54]. Alternative to constructing a DAG, directed partial correlation (DPC) [17] and regression tree based GENIE3 [18] methods have been proposed to uncover conditional dependencies in observed data. However, learning the structure of Bayesian networks from large data is computationally challenging [54] and the inferred networks are always acyclic and thus they do not support feedback loops [55]. On the other hand, DPC and GENIE3 return a complete list of interactions with non-zero weights of connectivity strengths, hence generating fully-connected networks in which the choice of an optimal confidence threshold is left open. Other methods such as the sparse partial correlation estimation (SPACE) [56] and its extension (ESPASE) [57], consider a penalized regression approach to construct the gene regulatory network. Specifically, their joint loss function requires predetermined penalty weights to combine the regression losses over all the response variables. Although the two methods utilize sparse variable selections, both estimations provide symmetric weight matrices representing weighted-undirected networks.

In this work, we modeled causal dependencies as sparse partial correlation coefficients which are obtained by minimizing a least square error subject to the unit l_1 norm ball. Inspired by the theoretical foundations for approximating partial correlations (Box 1), SPCS selects the best combination of a small number of explanatory factors that, under the conditional dependency assumption, explains the activity of each node in the GSN (Fig. 2B).

> **Box 1: Partial correlations can be approximated by multivariate regression coefficients**
>
> Consider the ordinary correlation ρ_{12} between two random variables v_1 and v_2. If v_1 and v_2 are correlated with $n-2$ other variables $v_3, v_4, \ldots, v_n$, we may regard ρ_{12} as a mixture of a direct correlation between v_1 and v_2 and an indirect portion due to the presence of other variables correlating with v_1 and v_2. The partial correlation measuring the direct portion of the total correlation can be represented as a correlation between v_1 and v_2 after removing effects due to other variables by a linear regression and the least square linear regression coefficients are proportional to the partial correlation coefficients [23].

The entire network is represented as a weighted-directed graph, $\mathbb{G} = (\mathbb{V}, \mathbb{E})$, where a set of nodes $\mathbb{V}$ represents genes and MutStates, and a set of edges $\mathbb{E}$ represents the relationships among the nodes. As mentioned in the Section 2.1 the network inference algorithm does not distinguish between the two types of nodes and assumes the nodes as random variables.

Let $\mathbb{I}$ denote the index set of n variables representing the nodes in $\mathbb{V}$. The nodes have observational activities over samples, and a $p \times n$ matrix $X = \{x_{ij}\}$ represents the data consisting of expressions of m genes and exposures of $n-m$ mutational signatures across p samples (patients). Assuming the incoming effects on a given variable f from its dominating covariates, the observed value x_{if}, corresponding to i-th sample, can be approximated as the following affine combination

$$x_{if} \approx \sum_{k \in \mathbb{I}\setminus\{f\}} x_{ik} w_{kf} + w_{0f} \tag{S1}$$

where $\mathbb{I}\setminus\{f\}$ denotes the index set of all variables except for the response variable f, and $w_{*f} \in R^{n-1}$ denotes the weights of incoming effects on f from the other $n-1$ variables and w_{0f} denotes the intercept adjusting the fitness between the response variable and its prediction.

Thus, our goal is to find the minimum of the least square error function subject to a unit l_1 norm constraint on w_{*f} as following

$$\begin{aligned} \underset{w_{*f} \in R^{n-1}, w_{0f} \in R}{\text{minimize}} \quad & \sum_{i=1}^{p} \left(x_{if} - \sum_{k \in \mathbb{I}\setminus\{f\}} x_{ik} w_{kf} - w_{0f} \right)^2 \\ \text{subject to} \quad & \sum_{k \in \mathbb{I}\setminus\{f\}} |w_{kf}| \leq 1 \end{aligned} \tag{S2}$$

For a response node (affected) f, a vector $w_{*f} = (w_{1f}, w_{2f}, \ldots w_{f-1f}, w_{f+1f}, \ldots w_{nf})^T$, the solution to the problem in Equation (S2), represents the weights of incoming effects from the other $n-1$ nodes (causal factors) in the network. The l_1 norm constraint pushes the weights of insignificant effects towards zero and allows selecting only dominant influences on the given response variable. Hence, this regularization acts to avoid over-fitting issues. A non-zero w_{kf} ($k \neq f$), selected for the node f, denotes the partial correlation coefficient representing the potential effect of the node k on the node f. In this setting, focusing on the activity of every single node in the network, the optimization problem in Equation (S2) considers all possible combinations of incoming effects from the other $n-1$ nodes and selects the best combination with their optimal influence weights to explain the given response activity under conditional dependency assumption. Therefore, the causality relationship between nodes k and f is estimated in the presence of the other $n-2$ variables.

S1.2   Solving SPCS model

An accurate solution of the problem in Equation (S2) is critical for the robust estimation of dependencies in the GSN. Although the least square error function is convex, the l_1 norm sparsity constraint is non-differentiable and derivative-based techniques such as Lagrange multipliers and Karush-Kuhn-Tucker (KKT) conditions are not directly applicable here due to the non-smoothness. Another attempt to resolve such an issue is to decompose the inequality of the l_1 norm into 2^n inequality constraints [58]. However, biological networks are often large in scale and it is practically difficult to accurately minimize such a large scale objective function over the exponential number of constraints within a reasonable time. Thus, to approximate the non-smooth constrained optimization, we rewrite the initial formulation in Equation (S2) as an unconstrained form with a penalty term

$$\underset{w_{*f}\in R^{n-1}, w_{0f}\in R}{\text{minimize}} \quad \sum_{i=1}^{p} \left(x_{if} - \sum_{k\in\mathbb{I}\setminus\{f\}} x_{ik}w_{kf} - w_{0f} \right)^2 + \lambda_f \cdot \sum_{k\in\mathbb{I}\setminus\{f\}} |w_{kf}| \tag{S3}$$

where a tuning parameter λ_f controls the strength of the penalty term, chosen for the given variable f to provide a balance between the least square error term and the l_1 norm constraint in the formulation in Equation (S2). In general, describing the criterion to define a reasonable value of λ_f is not trivial, due to the incompleteness of information linked to biological relevance. In fact, the exact relationship between the radius of the l_1 norm ball in Equation (S2) and the tuning parameter λ_f in Equation (S3) is assumed to be data-dependent. Therefore, it is reasonable to use a data-driven strategy for choosing λ_f. Akaike information criterion (AIC) is a statistical technique that provides the relative quality of statistical models for a given data by combining the maximum likelihood estimation of fitness with the number of parameters for inference [59]. The AIC is used in this work to decide the value of λ_f providing a solution with reasonable total incoming effect on f from its dominating factors.

S1.3   Higher moment-based strategy for causality direction

The solution to the problem in Equation (S2) may provide similar weights in both directions for some pairs of variables due to the presence of effects from confounding factors and noise in addition to their real dependencies. This uncertainty may require a complimentary analysis to decide the causality directions on these dependencies.

One way to decide causality relations is to perform perturbation experiments [60]. However, optimization of experimental design to predict which combination of perturbations allows to discover causality flows in a given network topology is often challenging and costly [61]. Hence, revealing causality directions by analyzing purely observational data has become a special focus of network biology [62]. Under a confounder-free assumption, higher moment statistics [63] indicate causality direction between two dependent variables from purely observational data. Alternative to the directionality decision for bivariate distributions, a confounder model [64] was recently designed to assign causality directions for several factors under a standard dependency assumption. By combining the key ideas of the two methods, we propose a higher moment-based strategy to decide causality directions for the bidirected edges in the initial network. The idea is to generate a bivariate distribution for two correlated variables by removing confounding effects from their observed activities, and then decide the causality direction between them using higher moment statistics on the corresponding residual activities (Fig. 2C). Specifically, for two variables having similar effects on each other (bidirectional edge) in the initial GSN, we first calculate their residual activities by removing effects due to the presence of the other $n-2$ variables [23]. Upon the removal, the corresponding residuals are expected to follow a bivariate distribution and only the dependency between

the focused pair remains in the residual activities. Thus, the causal variable can be distinguished from the affected by comparing the higher moments of the two residual distributions. Under the confounder-free assumption, the affected variable is closer to normality than the causal factor and the skewness and kurtosis are the higher moment statistics used to measure close-normality of the residual distributions.

Let x_{*j} and x_{*f} be j-th and f-th columns of the given data matrix X representing the observed activities of the variables j and f respectively. Then, the residual activities corresponding to the variables j and f can be obtained as follows

$$
\begin{aligned}
r_j &= x_{*j} - \sum_{k \in \mathbb{I}\backslash\{j,f\}} x_{*k} w_{kj} \\
r_f &= x_{*f} - \sum_{k \in \mathbb{I}\backslash\{f,j\}} x_{*k} w_{kf}
\end{aligned}
\tag{S4}
$$

where r_j and r_f are column vectors of size p, representing the residual activities after removing the confounding effects from the observed activities x_{*j} and x_{*f} respectively. For a bivariate distribution of two correlated variables, the affected variable is closer to the normal distribution than the causal factor, and the higher moment statistics, skewness and kurtosis, can be used to measure the close-normality [63]. As the result of the confounding effect removal, the residual activities r_j and r_f are assumed to follow a bivariate distribution, the causal direction between variables j and f can be identified by comparing the distribution shapes of r_j and r_f. We assign a causality direction between j and f as following

$$
Edge(j,f) = \begin{cases} j \rightarrow f \text{ and } w_{fj} := 0, & \text{if } |w_{jf}| > |w_{fj}| \text{ and } |\gamma_j| > |\gamma_f| \text{ and } |\delta_j| > |\delta_f| \\ j \leftarrow f \text{ and } w_{jf} := 0, & \text{if } |w_{jf}| < |w_{fj}| \text{ and } |\gamma_j| < |\gamma_f| \text{ and } |\delta_j| < |\delta_f| \\ j \leftrightarrow f, & \text{otherwise} \end{cases}
\tag{S5}
$$

where $\gamma_j = E[(r_j - \mu_{r_j})^3]/\sigma_{r_j}^3$ and $\gamma_f = E[(r_f - \mu_{r_f})^3]/\sigma_{r_f}^3$ describe the skewnesses of r_j and r_f while $\delta_j = E[(r_j - \mu_{r_j})^4]/\sigma_{r_j}^4 - 3$ and $\delta_f = E[(r_f - \mu_{r_f})^4]/\sigma_{r_f}^4 - 3$ describe the kurtosises respectively, where $\mu_.$ and $\sigma_.$ are the mean and standard deviation of the respective variables. We first decided dirctionality for every bidirectional edge in the GSN using the strategy S5. The performance is shown in first column of Table 1. Then, we explored different thresholds for edge weight cut-off τ to obtain the best set of directed edges. In particular, we accept the direction with a stronger weight if one direction provides a stronger weight than the threshold τ while the opposite direction provides a weaker weight compared to τ. The edge with the weaker weight is consequently removed from the network. The higher moment-based strategy in Equation (S5) is used to decide the direction if $|w_{fj}| \geq \tau$ and $|w_{jf}| \geq \tau$. The optimal threshold value was chosen ($\tau = 0.0391$ for BRCA and $\tau = 0.0521$ for LUAD) to maximize the fraction of correct directions in the set of recovered edges. The performance evaluation is provided in the third column (GENESIGNET optimized) of Table 1.

Complementary to the partial correlation selection, the higher moment-based strategy provides a causality direction between two correlated variables based on their distribution shapes if the dependency direction is not resolved by the SPCS. Particularly, this strategy is proposed to remove false directed-edges from the initial partial correlation network. To evaluate the robustness of our directionality inference, we tested the reproductivity of the inferred directions by performing a bootstrapping analysis. This analysis was done by repeatedly taking 80% of samples at random without replacement and assigning a direction for each edge using the higher moment-based strategy. Of 3,987 (BRCA) and 2,685 (LUAD) edge directions that were resolved in the presenting results, the reproductivity of each edge direction is tested 100 times during the random sampling. We

obtained 83.06% consistent, 2.7% inconsistent and 14.24% unsolved directions for total 3,968×100 decisions in the bootstrapping analysis on BRCA, and 84.04% consistent, 2.8% inconsistent and 13.17% unsolved directions for 2,685×100 decisions in the same analysis on LUAD. These results are summarized in Fig. S1.

Fig. S1. The reproductivity of the inferred directions assigned by the higher moment-based strategy. The numbers of consistent and inconsistent assignments during 100 times resampling are shown in the histograms **(A)** BRCA and **(B)** LUAD. The horizontal axis represents the percentages of resolved directions during the resampling while the vertical axis represents the numbers of consistently and inconsistently assigned directions. For example, out of 3,987 edge directions were tested in the bootstrap analysis for BRCA, the consistent percentages of 1,295 directed edges fall in the interval [97%,100%] (the last bin of blue histogram) while the inconsistent percentages of 2,916 directed edges fall in the interval [0%,2%] (the first bin of green histogram). 'Consistent directions' represents the percentage of the total consistent decisions for overall the directed edges during the 100 times resampling while 'Inconsistent directions' presents the percentage of the total inconsistent decisions. 'Unsolved directions' denotes the percentage of the total unsolved decisions that the two higher moment statistics provide contradictory decisions.

S1.4   Normalization of incoming and outgoing effects

The magnitudes of the causal effects in the network may provide valuable information to prioritize the candidate associations of genes and MutStates with underlining biological processes because the edge weights denote the contribution scores from causal factors to their affected targets. Hence, it is reasonable to bring the total incoming and outgoing effects of nodes to the same range in the GSN. The total incoming effect on each node was attempted to be normalized into the l_1 norm constraint in Equation (S2). However, the total outgoing effects are free from normalization. Moreover, an additional update described in Equation (S5) was performed to remove edges from the initial network obtained by the SPCS model in Equation (S2).

  We adopted a matrix normalization technique, alternate scaling [25], to rescale columns and rows of the weight matrix W into the unit l_1 norm ball. This procedure begins with rows in which each is mapped into the unit ball. Then do the same operation on columns, then on rows, and so on, until the sequence of matrices converges. The absolute difference of two consequence updates, by rows (W_{rows}) and by columns ($W_{columns}$), is used as the convergence criterion such that $\|W_{rows} - W_{columns}\|_F < 10^{-15}$. In the cases of the BRCA and LUAD analysis, the convergence was achieved after only 5 and 6 iterations, respectively. Note that this normalization increases the sparsity of the GSN since every column and row of W is iteratively mapped onto the l_1 norm space ($\|w_{k*}\|_{l_1} \leq 1$ for k-th row

and $\|w_{*k}\|_{l_1} \leq 1$ for k-th column of W) which rescales the weights to lower values, even assigning zero weights to weak associations during the iterative procedure.

S2 Materials

S2.1 Breast cancer data

The normalized gene expression data for 266 breast cancer (BRCA) patients were downloaded from Table S7 in [65,66]. Gene expression profiles for 2,204 genes involved in either DNA metabolic or immune response processes of the Gene Ontology (GO) database were selected for the analysis.

For mutational signatures, somatic mutation data were downloaded from the ICGC data portal (https://daco.icgc.org, release 22). The 3,479,652 point mutations were assigned to mutational signatures using SigMa [11]. SigMa divided all mutations into two groups, close-by **C**lustered and **D**ispersed mutations, and assigned each of these mutations to one of 12 COSMIC v2 signatures which were previously identified as active in BRCA (Signatures 1, 2, 3, 5, 6, 8, 13, 17, 18, 20, 26 and 30; https://cancer.sanger.ac.uk/cosmic/signatures). From the signatures classified by SigMa as described above, signature phenotype profiles 1D, 2C/D, 3C/D, 5D, 8C/D, and 13C/D that had exposure levels of at least 10% within each group were selected for further analysis (the numbering refers to the COSMIC signature index and C/D denotes signatures attributed to clustered and dispersed mutations). Examining their correlation patterns among patients, some of the signatures were grouped as follows: Signatures 3C/D and 8D were combined into DSB (double-stranded DNA break repair) related signatures, and Signatures 2C and 13C/D into APOBEC related signatures. The remaining signatures are treated separately, resulting in Signature 1, 2D, 5, APOBEC, DSB. A log transformation was consequently performed on exposures of each signature to make its distribution shape closer to a bell curve of normality.

Furthermore, we included binary information of homologous recombination deficiency as an additional variable in the analysis. The binary alteration information was obtained by aggregating functional inactivation information for BRCA1/BRCA2 and 16 other HR genes as provided in Supplementary Tables 4a and 4b of Davies *et al.* [29]. The positive entries were assigned a real value of 4.218 in the SPCS model (Section S2 in Supplementary Information) with the hyperparameter search for the best performance in terms of the means of minimum least square errors and maximum Pearson correlation between responses and predictions over all nodes.

S2.2 Lung adenocarcinoma data

The expression data (RNA-seq) of the lung adenocarcinoma (LUAD) from The Cancer Genome Atlas (TCGA) project were downloaded from the Genomic Data Commons Data Portal on 2020-06-05 (https://portal.gdc.cancer.gov/). Normalization and variance-stabilizing transformation (vst) of HTSeq count data were performed using DESeq2. Tumor and normal samples were split into different groups and only one sample per donor was kept in each group.

The TCGA LUAD exome mutation spectra were downloaded from Synapse (accession number: syn11801889) and decomposed into COSMIC v3 signatures SBS1, SBS2, SBS4, SBS5, SBS13, SBS40, and SBS45 using the quadratic programming (QP) approach available in the R package SignatureEstimation [67]. Only signatures predominantly active in lung cancer (signatures that were present in at least 5% of samples and were responsible for at least 1% of mutations) were considered based on the initial sample decomposition provided by Alexandrov et al. [2] (Synapse accession number: syn11804065). Signature SBS45 is likely a sequencing artifact so it was omitted from further analyses presented in this study. The same log transformation used in BRCA analysis was performed on signature exposure data as well.

We analyzed 466 tumor samples that had both gene expression and mutational signature exposure data available. We analyzed 2,433 genes belonging to the DNA metabolic process and immune system process in GO terms (genes that are not expressed in at least 10% of the samples were omitted). The gene expression and mutational signature exposure data were combined to form an input data matrix.

S3   Supplementary Figures

Fig. S2. The removal of confounding effects allows the higher moment statistics to distinguish causal factor from its own target. For two random variables in each panel, the joint distribution of their observational activities is shown on the left while that of corresponding residual activities is on the right. The experimentally confirmed regulatory relations [24] indicate the correct directions. The values in brown highlight the causal factors. **(A)** Without removing confounding effects from expressions of EGR1 and JUN, both moments provided an incorrect direction. After removing the confounding effects, the two moments indicate the correct direction. **(B)** Before removing the effects, the skewness provide the direction from CEBPB to MBD2 while the kurtosis supports the opposite direction. However, the proposed higher moment-based strategy indicates the correct direction **(C)** Either without or with the effect removal, the two moments provide the experimentally confirmed influence in WAS from SPI1. Although the higher moment-based strategy did not suggest any changes on the directionality decision, the effect removal uncovers a stronger association between the two genes. **(D)** The two moments provide contradictory decisions without removing the confounding effects from the exposures of the two mutational signatures. However, with the removal, the inferred direction is from SBS4 to SBS40.

Fig. S3. Subnetworks of GSN for BRCA centered on MutStates Edge and node colors are as in Fig. 1. If the genes adjacent to a given MutState are enriched with specific GO pathways ($q - value < 0.01$), the names of the enriched genes are provided under the pathway names in a box and the other adjacent genes with edge weight cut-off ($|w_{ij}| \geq 0.01$) are titled as 'Other genes' in the same box. If the adjacent genes are not enriched with any specific GO pathways ($q - value < 0.01$), the names of the genes with edge weight cut-off ($|w_{ij}| \geq 0.01$) are provided in a box. The complete lists of upstream and downstream sets for MutStates and their enriched GO terms obtained in the analysis on the BRCA data are available as a resource at the GENESIGNET GitHub repository (https://github.com/ncbi/GeneSigNet).

Fig. S4. Subnetworks of GSN for LUAD centered on MutStates The meaning of edge and node colors, and boxes is the same as in Fig. S3. The complete lists of upstream and downstream sets for MutStates and their enriched GO terms obtained in the analysis on the LUAD data are available as a resource at the GENESIGNET GitHub repository (https://github.com/ncbi/GeneSigNet).