A general model of the dynamics of genes with dual σ factor preference

Ines S. C. Baptista[1,+], Vinodh Kandavalli[1,2,+], Vatsala Chauhan[1], Mohammed Bahurudeen[1], Bilena Almeida[1], Cristina Palma[1], Suchintak Dash[1], and Andre S. Ribeiro[1,3, *]

[1] Laboratory of Biosystem Dynamics, Faculty of Medicine and Health Technology, Tampere University, Tampere, 33520, Finland.

[2] Department of Cell and Molecular Biology, Uppsala University, Uppsala, 752 37, Sweden.

[3] CA3 CTS/UNINOVA. Faculty of Science and Technology, NOVA University of Lisbon, Quinta da Torre, 2829-516, Caparica, Portugal.

[+] Equal contributions.

* To whom correspondence should be addressed. E-mail: andre.sanchesribeiro@tuni.fi.

Present Address: Andre S. Ribeiro, Arvo Ylpön katu 34, P.O. Box 100, 33014, Tampere University, Finland.

Abstract

Escherichia coli uses the ability of σ factors to recognize specific DNA sequences in order to quickly control large gene cohorts. While most genes respond to only one σ

factor, approximately 5% have dual σ factor preference. The ones in significant numbers are 'σ^{70+38} genes', responsive to σ^{70}, which controls housekeeping genes, as well as to σ^{38}, which activates genes during stationary growth and stresses. We show that σ^{70+38} genes are almost as upregulated in stationary growth as genes responsive to σ^{38} alone. Also, their response strengths to σ^{38} are predictable from their promoter sequences. Next, we propose a sequence- and σ^{38} level-dependent, analytical model of σ^{70+38} genes applicable in the exponential, stationary, and in the transition period between the two growth phases. Finally, we propose a general model, applicable to other σ factors as well. This model can guide the design of synthetic circuits with sequence-dependent sensitivity and plasticity to transitions between the exponential and stationary growth phases.

Author Summary

Present challenges in Synthetic Biology include the design of genetic circuits that are robust to growth phase transitions and whose responsiveness is sequence-dependent, and, thus predictable prior to design. We present and validate an empirical-based, sequence-dependent analytical model of *E. coli* genes with dual responsiveness to the regulators σ^{70} and σ^{38}. These genes, supported by our sequence-dependent model, could become building blocks for synthetic genetic circuits functional in both the exponential and the stationary growth phases.

1. Introduction

In *Escherichia coli (E. coli)*, genes are expressed by RNA polymerase (RNAP), which has 5 subunits ($\alpha_2\beta\beta'\omega$) and can synthesize RNA by starting and ending at sequence-specific DNA locations, with the former being known as 'promoter' regions [1]. Transcription is constantly regulated, mostly at the promoter regions, which typically harbor transcription factor (TF) binding sites (TFBS) and other regulatory sequence motifs [2–7]. Such regulation is an essential survival skill of adaptability to transitions in both internal as well as external conditions [8,9]

One mechanism of large-scale, relatively synchronous gene expression regulation is executed by σ factors [4,10–13]. *E. coli* has seven different σ factors [14]. During exponential growth in optimal conditions, RNAP mostly transcribes genes with preference for σ^{70}, responsible for basic cell functions [15]. Other σ factors are expressed under specific stresses [16], and activate a specific gene cohort. For example, after growing exponentially at the cost of environment components, *E. coli* usually switches to stationary growth. This is triggered, among other, by RpoS (σ^{38}), whose appearance activates ~10% of the genome [17,18], which leads to key phenotypic modifications [18–24].

During this adaptation, the concentration of RNAP remains relatively constant [25] and, since the number of RNAP core enzymes is limited, σ factors compete for them [10,12,14,20,26,27]. Thus, when σ^{38} numbers increase, not only are the genes responsive to σ^{38} activated, but also other genes, previously active, are indirectly negatively regulated, due to the reduction in RNAPs carrying σ^{70} [10,26,28].

For this regulatory system to be efficient, promoters need to have high specificity to one and only one σ factor. Nevertheless, there is a small fraction of promoters that can recognize more than one σ factor [6,29,30]. The most common (84%) are the

promoters responsive to both σ⁷⁰ and σ³⁸ [31]. Other cases are too rare to obtain

sufficient statistics on their kinetics (Table S3 in S3 Appendix).

Here we investigate how the sequences of promoters recognized by both σ^{70} and σ^{38}

relate to the dynamics of genes that they control (here named 'σ^{70+38} genes'). From

flow cytometry and RNA-seq measurements, along with sequence analysis, we

establish a relationship between RNA and protein levels in each growth phase with

the promoter sequence (logos of positions -41 to -1 shown in Fig S1 in S2 Appendix).

From this, we propose a model of the dynamics σ^{70+38} genes that fits their behavior

prior, during, and following the transition to stationary growth. The model is dependent

on both the promoter-sequence as well as the levels of σ factors in the cell.

2. Results

2.1. RNA fold changes when shifting to stationary growth

To study how σ^{70+38} genes respond to stationary growth, we first identified when are

cells in exponential and stationary growth. We used both wild type (WT, control) cells

and a MGmCherry strain [19] carrying fluorescently tagged σ^{38} to measure its levels

prior to, during, and after shifting from exponential to stationary growth (Section 4.2).

Cells at 150 min after being placed in fresh medium are used as representatives of

cells in mid-exponential growth (Fig 1A). Meanwhile, at 500 min and onwards, cells

exhibit stationary growth (Fig 1A).

From the measurements, σ^{38} levels are small at 150 min, while from 500 min onwards

they are stably high (Fig 1B). Considering that the cell areas are ~15% smaller in the

stationary phase (Figs S3A and S3B in S2 Appendix), at 700 min the absolute

concentration of σ^{38} is ~32 times higher than at 150 min (Fig 1C and Fig S3C in S2

Appendix). Meanwhile, RNAP concentration is largely stable (only ~7% higher during

stationary growth) (Fig 1C and Fig S3D in S2 Appendix).

Next, we measured changes in RNA levels, by RNA-seq, between moments 150 min

and 500 minutes by calculating the $\log_2$ fold changes in RNA levels (LFC$_{RNA}$) (Sections

4.2 and 4.4). Gaussian fits to the LFC$_{RNA}$ distributions (Fig 1D) of σ^{70+38} genes, σ^{70}

genes, and σ^{38} genes, as well as all other genes, have high coefficient of determination

(R^2) values of the fits suggesting that they capture well the shapes of the empirical

distributions.

In general, σ^{70+38} genes are nearly as upregulated as σ^{38} genes (Fig 1D). On the other

hand, σ^{70} genes are weakly downregulated, likely due to indirect negative regulation

caused by the increase in σ^{38} levels combined with the limited pool of RNAP core

enzymes [10,12,14,20,26,27]. Finally, most other genes (~2737 out of the 4308 genes)

are relatively unresponsive.

Fig 1. Cell growth, RNAP, RpoS and genome-wide RNA levels when changing from exponential to stationary growth. (A) Optical densities 'O.D.$_{600}$' of wild-type MG1655 (control) and MGmCherry (*rpoS::mcherry*) strains (Section 4.1). **(B)** Mean population fluorescence of WT and MGmCherry strains. Black vertical bars show the timing of measurements in exponential (150) and stationary (500 minutes onwards) growth. At 0 minutes, cells were moved to fresh medium (Section 4.2). **(C)** Fold-change between the exponential and stationary growth phases of mean cellular fluorescence relative to cell area (FC(μ/Cell Area)) due to changes in RNAP-GFP and in RpoS-mCherry, respectively. Error bars are the standard error of the mean (SEM). **(D)** Gaussian fits to the distributions of LFC$_{RNA}$ of gene cohorts. Vertical lines mark the mean**.**

2.2. Propagation of shifts in RNA numbers to protein numbers

We used a YFP strain library to measure single-cell protein levels (Section 4.5) of 9 out of the 64 σ^{70+38} genes (only 15 of the 64 are represented in this library and some exhibited too weak signals to be detected by the flow cytometer). These 9 genes, according to their LFC$_{RNA}$, cover well the state space of response strengths of σ^{70+38} genes (Section S1.2 in S1 Appendix).

From the single-cell distributions of protein levels in exponential and stationary growth, we extracted means, μ_P, and squared coefficient of variations, CV^2_P. We also calculated the log$_2$ fold changes in μ_P, LFC$_P$. Since LFC$_P$ and the corresponding LFC$_{RNA}$ are linearly positively correlated, with a high R^2 (Fig 2A), we conclude that changes in RNA levels of σ^{70+38} genes during the growth phase transition propagate to their protein levels.

We then analyzed the noise in the dynamics of these genes, prior and after the growth phase transition. In Figs 2B1 and 2B2, CV^2_P decreases quickly with μ_P for small μ_P,

but only weakly for high μ_P. This is well described by a function of the form:

$CV_P^2 = C \times \mu_P^{-1} + N$ [32] (where N is the noise floor, a lower bound on the cell-to-cell

variability of protein levels in clonal populations due to extrinsic factors [33]), in

agreement with past reports [32,34,35].

Fig 2. Single-cell protein levels of σ^{70+38} genes in the exponential and stationary growth

phases. (A) $\log_2$ fold-change of protein levels, LFC_P, plotted against the corresponding

LFC_{RNA}, during the shifting from exponential to stationary growth. **(B1)** and **(B2)** CV^2_P plotted

against μ_P of σ^{70+38} genes in the exponential and stationary growth phases, respectively. Error

bars (small) are the standard error of mean. All figures show the best fitting curves and their

95% prediction bounds (shadow areas). Horizontal dashed lines are the noise floors.

2.3. Full sequence-dependent model of the shift in the transcription dynamics of σ^{70+38} genes

Here, we present a sequence-dependent model of transcription of σ^{70+38} genes. Based

on past works, we assume that only σ^{70} is present in high numbers (in holoenzyme

form) in the exponential phase and that only σ^{38} increases significantly in numbers

when shifting to stationary growth [10,14,20,26,36–38].

The model therefore accounts for the competition between σ^{70} and σ^{38} for binding to

RNAP core enzymes, since these exist in limited numbers in *E. coli*

[10,12,14,20,26,27] and since σ factor numbers differ between the two growth phases.

For this, we set reactions for binding and unbinding of σ^{70} and σ^{38} to free floating RNAP

core enzymes, (R1a) and (R1b), where $K^{\sigma 70}$ and $K^{\sigma 38}$ are the ratios between the

association and dissociation rate constants:

$$\text{RNAP} + \sigma^{70} \xleftrightarrow{\quad K^{\sigma 70} \quad} \text{RNAP}.\sigma^{70} \tag{R1a}$$

$$\text{RNAP} + \sigma^{38} \xleftrightarrow{\quad K^{\sigma 38} \quad} \text{RNAP}.\sigma^{38} \tag{R1b}$$

Given (R1a) and (R1b), RNA production dynamics depends on the ratio between σ^{70}

and σ^{38} numbers if, and only if, $K^{\sigma 70}$ and $K^{\sigma 38}$ differ. Meanwhile, the limited pool of

RNAPs is accounted for by setting:

$$\left[RNAP_{total} \right] = \left[RNAP.\sigma^{70} \right] + \left[RNAP.\sigma^{38} \right]. \tag{1}$$

Next, to introduce the dual responsiveness to σ^{70} and σ^{38}, we define two competing,

sequence-dependent reactions of transcription, (R2a) and (R2b), differing in which

holoenzyme binds to the promoter. Further, to also introduce sequence dependence,

we define two sequence-based distances: $D_{\sigma 38}$ and $D_{\sigma 70}$. $D_{\sigma 38}$ is the distance, in

number of nucleotides, of a promoter sequence differing from the consensus (average)

sequence of promoters with σ^{38} dependency. $D_{\sigma 70}$ is the distance in number of

nucleotides differing from the consensus sequence of promoters with σ^{70} dependency

(Section 4.8.3):

$$\text{Pro} + \text{RNAP}.\sigma^{70} \xrightarrow{\quad k_t^{\sigma 70}(D_{\sigma 70}) \quad} \text{Pro} + \text{RNA} + \text{RNAP} + \sigma^{70} \tag{R2a}$$

$$\text{Pro} + \text{RNAP}.\sigma^{38} \xrightarrow{\quad k_t^{\sigma 38}(D^{\sigma 38}) \quad} \text{Pro} + \text{RNA} + \text{RNAP} + \sigma^{38} \tag{R2b}$$

The rate constants of these reactions, which control the affinities to the holoenzymes,

are expected to differ between genes, as they depend on the promoter sequences

(and potentially which transcription factors are acting on the promoters, here not

represented for simplicity). Nevertheless, here, also for simplicity, $kt^{\sigma70}$ and $kt^{\sigma38}$ are

assumed to be a function of only $D_{\sigma70}$ and $D_{\sigma38}$ respectively.

Together, the set of reactions (R1a), (R1b), (R2a) and (R2b), model the transcription

kinetics of σ^{70+38} genes before, during and, after shifting from exponential to stationary

growth. The rates $kt^{\sigma70}(D_{\sigma70})$ and $kt^{\sigma38}(D_{\sigma38})$ are dissected below.

2.4. Reduced model of the shift in transcription kinetics

To reduce the model, we assume that the numbers of RNAP.σ^{70} and RNAP.σ^{38} in the

cells are significantly larger than 1, which is expected. Also, we assume that the rate-

limiting steps in transcription initiation (during which a promoter is occupied and thus

new events cannot start [39,40]) are shorter in time-length than the intervals between

transcription events. If so, for any σ^{70+38} gene:

$$\text{Pro} \xrightarrow{\left[RNAP.\sigma^{70}\right] \cdot k_t^{\sigma70}(D_{\sigma70})} \text{Pro} + \text{RNA} + \text{RNAP} + \sigma^{70} \tag{R3a}$$

$$\text{Pro} \xrightarrow{\left[RNAP.\sigma^{38}\right] \cdot k_t^{\sigma38}(D_{\sigma38})} \text{Pro} + \text{RNA} + \text{RNAP} + \sigma^{38} \tag{R3b}$$

We then merge (R3a) and (R3b) into a single transcription process:

$$\text{Pro} \xrightarrow{\left[RNAP.\sigma^{70}\right] \cdot k_t^{\sigma70}(D_{\sigma70}) + \left[RNAP.\sigma^{38}\right] \cdot k_t^{\sigma38}(D_{\sigma38})} \text{Pro} + \text{RNA} \tag{R4}$$

Finally, to complete the model, we included a reaction for translation of RNAs into

proteins (R5), as well as events of RNA (R6) and protein (R7) decay due to

degradation and dilution by cell division:

$\quad \text{RNA} + \text{Rib} \xrightarrow{k_{tr}} \text{RNA} + \text{Rib} + \text{P}$ (R5)

$\quad \text{RNA} \xrightarrow{\gamma_{RNA}} \varnothing$ (R6)

$\quad \text{P} \xrightarrow{\gamma_P} \varnothing$ (R7)

2.5. Analytical solution of the reduced model

193 Based on the reduced model, we obtained an analytical solution for the expected fold

194 change in RNA numbers of a gene whose promoter has preference for both σ^{70} as

195 well as σ^{38}. From (R4), (R5), (R6), and (R7) we derived the expected RNA numbers

196 produced by a σ^{70+38} gene:

$$197 \quad RNA = \frac{\left[RNAP.\sigma^{70} \right] \cdot k_t^{\sigma 70}(D_{\sigma 70}) + \left[RNAP.\sigma^{38} \right] \cdot k_t^{\sigma 38}(D_{\sigma 38})}{\gamma_{RNA}} \tag{2}$$

198 For simplicity, the rate constants are not expected to differ between the growth phases

199 (as they depend on biophysical parameters that should not change significantly, such

200 as binding affinities, etc.). Consequently, the fold-change in RNA numbers between

201 the two growth phases should equal:

$$202 \quad FC_{RNA} = \frac{RNA_{sta}}{RNA_{exp}} = \frac{\left[RNAP.\sigma^{70} \right]_{sta} \cdot k_t^{\sigma 70}(D_{\sigma 70}) + \left[RNAP.\sigma^{38} \right]_{sta} \cdot k_t^{\sigma 38}(D_{\sigma 38})}{\left[RNAP.\sigma^{70} \right]_{exp} \cdot k_t^{\sigma 70}(D_{\sigma 70}) + \left[RNAP.\sigma^{38} \right]_{exp} \cdot k_t^{\sigma 38}(D_{\sigma 38})} \cdot \Gamma \tag{3}$$

203 where γ is the ratio between the RNA decay rates in the two growth phases:

$$204 \quad \Gamma = \frac{\gamma_{RNA_{exp}}}{\gamma_{RNA_{sta}}} \tag{4}$$

Since [RNAP_total] is similar in the two growth phases (Fig 1C):

$$FC_{RNA} = \cfrac{\cfrac{\left[RNAP.\sigma^{70}\right]_{sta}}{\left[RNAP_{total}\right]} \cdot k_t^{\sigma70}(D_{\sigma70}) + \cfrac{\left[RNAP.\sigma^{38}\right]_{sta}}{\left[RNAP_{total}\right]} \cdot k_t^{\sigma38}(D_{\sigma38})}{\cfrac{\left[RNAP.\sigma^{70}\right]_{exp}}{\left[RNAP_{total}\right]} \cdot k_t^{\sigma70}(D_{\sigma70}) + \cfrac{\left[RNAP.\sigma^{38}\right]_{exp}}{\left[RNAP_{total}\right]} \cdot k_t^{\sigma38}(D_{\sigma38})} \Gamma \quad (5)$$

Also, considering equation (1), then:

$$\frac{\left[RNAP.\sigma^{70}\right]}{\left[RNAP_{total}\right]} + \frac{\left[RNAP.\sigma^{38}\right]}{\left[RNAP_{total}\right]} \approx 1 \tag{6}$$

For simplicity, let ρ_{exp} and ρ_{sta} be:

$$\rho_{exp} = \frac{\left[RNAP.\sigma^{38}\right]_{exp}}{\left[RNAP_{total}\right]} = \frac{\left[RNAP_{total}\right] \cdot \left[\sigma^{38}\right]_{exp} \cdot K^{\sigma38}}{\left[RNAP_{total}\right] \cdot \left(\left[\sigma^{70}\right]_{exp} \cdot K^{\sigma70} + \left[\sigma^{38}\right]_{exp} \cdot K^{\sigma38}\right)} \tag{7a}$$

$$\rho_{sta} = \frac{\left[RNAP.\sigma^{38}\right]_{sta}}{\left[RNAP_{total}\right]} = \frac{\left[RNAP_{total}\right] \cdot \left[\sigma^{38}\right]_{sta} \cdot K^{\sigma38}}{\left[RNAP_{total}\right] \cdot \left(\left[\sigma^{70}\right]_{sta} \cdot K^{\sigma70} + \left[\sigma^{38}\right]_{sta} \cdot K^{\sigma38}\right)} \tag{7b}$$

From (5):

$$FC_{RNA} = \frac{(1-\rho_{sta}) \cdot k_t^{\sigma70}(D_{\sigma70}) + \rho_{sta} \cdot k_t^{\sigma38}(D_{\sigma38})}{(1-\rho_{exp}) \cdot k_t^{\sigma70}(D_{\sigma70}) + \rho_{exp} \cdot k_t^{\sigma38}(D_{\sigma38})} \cdot \Gamma \tag{8}$$

Finally, since, in the exponential growth phase, $\left[RNAP.\sigma^{70}\right] \gg \left[RNAP.\sigma^{38}\right]$, then

$\rho_{exp} \approx 0$. Thus, (8) can be simplified:

$$FC_{RNA} = \gamma \cdot (1-\rho_{sta}) + \rho_{sta} \cdot \Gamma \cdot \frac{k_t^{\sigma38}(D_{\sigma38})}{k_t^{\sigma70}(D_{\sigma70})} \tag{9}$$

All parameters in (9) can be measured directly, except $k_t^{\sigma70}(D_{\sigma70})$ and $k_t^{\sigma38}(D_{\sigma38})$.

Meanwhile, in the stationary phase: $\left[\sigma^{38}\right]^{\text{sta}} = 0.3 \cdot \left[\sigma^{70}\right]$ [20,26,37]. Also, $K^{\sigma70} = 5K^{\sigma38}$

[41]. Thus:

$$\rho^{sta} = \frac{\left[RNAP.\sigma^{38}\right]_{\text{sta}}}{\left[RNAP_{total}\right]} \approx \frac{0.3 \cdot \left[\sigma^{70}\right]_{\text{sta}} \cdot \dfrac{K^{\sigma70}}{5}}{\left[\sigma^{70}\right]_{\text{sta}} \cdot K^{\sigma70} + 0.3 \cdot \left[\sigma^{70}\right]_{\text{sta}} \cdot \dfrac{K^{\sigma70}}{5}} = 0.0566 \qquad (10)$$

2.6. Promoter sequence affects the expression of σ^{70+38} genes

To model the influence of the promoter sequence (i.e., $D_{\sigma70}$ and $D_{\sigma38}$) on the response

strength of σ^{70+38} genes to σ^{38}, we studied three models. In all, we assumed the basal

levels of transcription rates, '$k_{t0}^{\sigma70}$' and '$k_{t0}^{\sigma38}$'. The overall transcription rates are tuned

by a function f (Table 1) of the influence of $D_{\sigma70}$ and $D_{\sigma38}$ on the basal transcription

rate:

$\quad k_t^{\sigma70}(D_{\sigma70}) = k_{t0}^{\sigma70} \cdot f\left(D_{\sigma70}\right)$ \hfill (11a)

$\quad k_t^{\sigma38}(D_{\sigma38}) = k_{t0}^{\sigma38} \cdot f\left(D_{\sigma38}\right)$ \hfill (11b)

Consider that, in general, as $D_{\sigma i}$ of a promoter increases, the promoter should become

more distinct from the 'average' promoter with preference for σ^i. Consequently, its

affinity to σ^i is expected to decrease. If this holds true, then the promoter consensus

sequence of genes with preference for a given σ factor should have strong affinity to

that σ factor. We hypothesize, for simplicity, it has the strongest affinity. If true, it

follows that as $D_{\sigma38}$ increases, the transcription rate by RNAP.σ^{38} should decrease.

Having this in account, to model how a promoter sequence controls the transcription

kinetics of promoters recognized by σ^{70} and by σ^{38}, we considered a linear model

(model I in Table 1), a rational model (model II), where the transcription rate is

inversely proportional to $D_{\sigma i}$, and an exponential model (model II), similar to the one in

[6,29,30] for σ^{70} genes.

Table 1. Models of how a sequence's p-distance, D_σ, affects a promoter's response to

changes in the concentration of its preferred σ factor. Here, m is an empirical-based constant.

I. Linear model	$f(D_{\sigma i}) = 1 - m_i \cdot D_{\sigma i}$, $m_i > 0$ & $m_i \times D_{\sigma i} < 1$, $i = 38$ or 70
II. Rational model	$f(D_{\sigma i}) = \dfrac{1}{1 + m_i \cdot D_{\sigma i}}$, $m_i > 0$, $i = 38$ or 70
III. Exponential model	$f(D_{\sigma i}) = e^{-m_i \cdot D_{\sigma i}}$, $m_i > 0$, $i = 38$ or 70

We fitted the models (Fig 3) to the empirical fold changes (FC) of σ^{70+38} genes (FC$_{RNA}$)

with known input TFs (Table S4 and Table S6) (FC rather than LFC were used,

because this allowed for simpler equations for the fitting surfaces). From the R^2 values

of the best fitting models (Table 2), we find that the model III best describes how $D_{\sigma 70}$

and $D_{\sigma 38}$ affect $k_t^{\sigma 70}$ and $k_t^{\sigma 38}$.

Next, we validated the fitted models by quantifying how well they predict the dynamics

(Fig 3) of σ^{70+38} genes without input TFs (Table S4 and Table S6). Given the R^2 values,

we conclude that the tuned model III predicts well their dynamics. Further, the fact that

the model obtained from genes with TF regulation fits well (with a higher R^2) the genes

without TF regulation is suggestive that, in our measurement conditions, such TF

regulation is not exerting a significant role.

Table 2. Goodness of fit, measured by the R^2 of the surface fitting of FC_{RNA} as a function of

$D_{\sigma 38}$ and $D_{\sigma 70}$. Shown are the models and best fitting parameter values, where $K_0 = \dfrac{k_{t0}^{\sigma 38}}{k_{t0}^{\sigma 70}}$. The

surfaces were fitted to σ^{70+38} genes without input TFs and validated on genes σ^{70+38} genes with

input TFs.

I. Linear model	Surface Equation	$FC_{RNA} = \gamma \cdot (1 - \rho_{\text{sta}}) + \gamma \cdot \rho_{\text{sta}} \cdot K_0 \cdot \left(\dfrac{1 - m_{38} \times D_{\sigma 38}}{1 - m_{70} \times D_{\sigma 70}} \right)$ (12a)	
	Coefficients	m_{38}	-834.80
		m_{70}	-1.01×10^4
		Γ	85.60
		K_0	2.39×10^{-4}
		ρ_{sta}	0.06 (Equation (10))
	R^2	σ^{70+38} genes with input TFs	<0
		σ^{70+38} genes without input TFs	<0
II. Rational model	Surface Equation	$FC_{RNA} = \gamma \cdot (1 - \rho_{\text{sta}}) + \gamma \cdot \rho_{\text{sta}} \cdot K_0 \cdot \left(\dfrac{1 + m_{70} \cdot D_{\sigma 70}}{1 + m_{38} \cdot D_{\sigma 38}} \right)$ (12b)	
	Coefficients	m_{38}	-0.38
		m_{70}	26.51
		Γ	0.21
		K_0	241.70
		ρ_{sta}	0.06 (Equation (10))
	R^2	σ^{70+38} genes with input TFs	0.04

		σ^{70+38} genes without input TFs	0.00
III. Exponential model	Surface Equation	$FC_{RNA} = \gamma \cdot (1 - \rho_{\text{sta}}) + \gamma \cdot \rho_{\text{sta}} \cdot K_0 \cdot e^{-(m_{38} \cdot D_{\sigma 38} - m_{70} \cdot D_{\sigma 70})}$ (12c)	
	Coefficients	m_{38}	30.78
		m_{70}	56.90
		Γ	28.20
		K_0	3.841×10^{-9}
		ρ_{sta}	0.06 (Equation (10))
	R^2	σ^{70+38} genes with input TFs	0.55
		σ^{70+38} genes without input TFs	0.98

Fig 3. Fold changes in RNA levels of σ^{70+38} genes plotted against their promoter sequence *p*-distances to the consensus sequence of σ^{70} dependent promoters ($D_{\sigma 70}$) and of σ^{38} dependent promoters ($D_{\sigma 38}$). (A1-A3) show the best fitting surfaces to FC(µ$_{RNA}$)

assuming **(A1)** a linear function, **(A2)** a rational function, and **(A3)** an exponentially decreasing

function. Only σ^{70+38} genes with input TFs and FDR < 0.05 are included. Light red points are

above the surface, while dark ones are below. **(B1-B3)** Same plots but the surfaces are those

obtained from (A1-A3) and applied to σ^{70+38} genes without input TFs and FDR < 0.05. The

dashed black lines depict the vertical distances between the estimated and measured

FC(μ_{RNA}).

Finally, from Fig 2A, the LFC of protein numbers (LFC_P) correlates linearly with

LFC_{RNA}. The same is predictable from the model (reactions R5 to R7) (Section S1.3

in S1 Appendix). Here, to obtain the best fitting model for protein log2 fold changes,

we fitted a line to the data in Fig 2A, and extracted from it a scaling factor, α, between

LFC_{RNA} and LFC_P. From this:

$$LFC_P = \log_2\left[\Gamma \cdot (1-\rho_{sta}) + \Gamma \cdot \rho_{sta} \cdot K_0 \cdot e^{-(m_{38} \cdot D_{\sigma 38} - m_{70} \cdot D_{\sigma 70})} \right] \cdot \alpha + \beta \qquad (13)$$

where β equals -0.67 and is the intercept between the y axis and the best fitting line,

while α is the scaling factor between RNA and protein $\log_2$ fold changes and equals

0.1 (Fig 2A).

2.7. σ^{70+38} genes without known input transcription factors

Next, we fitted the exponential model III directly to σ^{70+38} genes without input TFs (not

to mistake with when they were used to validate the surface created from σ^{70+38} genes

with TF inputs). The best fitting model (R^2 of 1) has similar m_{38} and m_{70} (49.04 and

52.48, respectively). Thus, for this restricted set of genes, since $\Delta = D_{\sigma 38} - D_{\sigma 70}$

(Section 4.8.3), the surface equation (12c) in Table 2 can be approximated to:

$$FC_{RNA} \approx \Gamma \cdot (1-\rho_{sta}) + \Gamma \cdot \rho_{sta} \cdot K_0 \cdot e^{-r \times D} \qquad \text{, where } r = \frac{m_{38} + m_{70}}{2}$$

Applying the $\log_2$ to both sides, one has:

$LFC_{RNA} \approx \log_2\left(\Gamma\cdot(1-\rho_{sta})+\Gamma\cdot\rho_{sta}\cdot K_0\cdot e^{-r\times D}\right)=C_1+\log_2\left(\Gamma\cdot\rho_{sta}\cdot K_0\right)+\log_2\left(e^{-r\cdot D}\right)=C_1+C_2+\log_2\left(e^{-r\cdot D}\right)$

If (C_1+C_2) is much smaller than $\log_2\left(e^{-r\cdot D}\right)$, we expect that

$LFC_{RNA} \propto \log_2\left(e^{-r\cdot D}\right)\approx -\dfrac{r}{\ln(2)}\cdot D$ (14)

Therefore, and even though $D_{\sigma38}$ and $D_{\sigma70}$ are not significantly correlated (Fig 4A), one

finds a linear correlation between LFC_{RNA} and Δ (Fig 4B).

Fig 4. Log$_2$ fold changes in RNA levels of σ^{70+38} genes and their promoter p-distances

to the consensus sequence of σ^{70} dependent promoters ($D_{σ70}$) and of σ^{38} dependent

promoters ($D_{σ38}$). (A) Scatter plot of $D_{σ38}$ plotted against $D_{σ70}$. **(B)** Scatter plot of $\Delta = D_{σ38}$ -

$D_{σ70}$ plotted against LFC_{RNA}. The shadows of the best fitting lines are the 95% prediction

bounds.

For comparison, we produced similar figures in search for linear correlations between

Δ and LFC_{RNA} in the cohorts of genes whose promoters have preference for σ^{70}, σ^{38},

σ^{24}, σ^{28}, σ^{32}, and σ^{54} (Figs 5A-5F). No significant correlation with a high value of R^2

was found. We further studied the two small cohorts of genes whose promoters have

preference for two σ factors other than σ^{70+38}. Here, correlations are found, but they

are not statistically significant, likely due to the small numbers of genes (Figs 5G and

5H).

Fig 5. Fold changes in RNA levels and Δ. Genes whose promoters have preference for **(A)**

σ^{70} **(B)** σ^{38} **(C)** σ^{24} **(D)** σ^{28} **(E)** σ^{32} **(F)** σ^{54} **(G)** σ^{70+24} and **(H)** σ^{70+32} are plotted against $\Delta = D_{\sigma38}$ -

$D_{\sigma70}$. The shadows of the best fitting lines are the 95% prediction bounds.

2.8.   Expanding the model to the growth phase transition period

We expanded the model to include the transition period between the growth phases.

For this, we collected temporal data on the protein numbers of three σ^{70+38} genes,

specifically pstS, aidB and asr. These genes are used as representative of σ^{70+38}

genes with, strong, mild, and weak response strengths to the phase transition,

respectively (Fig 6A). All data was well fitted by Hill functions (Fig 1B and Fig 6A).

Moreover, in all three genes, there is a linear relationship between the fold changes in

protein levels and in σ^{38} levels over time (Fig 6B).

Fig 6. Temporal changes in the fold-change of protein levels as σ^{38} changes. (a) Protein

levels of three σ^{70+38} genes prior to, during, and after the transition from exponential to

stationary growth phases. The balls are the empirical mean fold changes (FC_P) in protein

expression levels relative to the first moment. The lines are the corresponding best fitting Hill

functions (parameter values in Table S5 in S3 Appendix). **(b)** Scatter plot of FC_P against the

corresponding σ^{38} levels (data from Fig 1B) over time. The shadows are the 95% prediction

bounds.

The dependency on σ^{38} levels is accounted for in model (ρ_{sta} is a function of σ^{38} levels

in equation (7b), which are time dependent). Consequently, given the goodness of fit

of the Hill functions, we propose the following time-dependent model:

$$FC_P(t) = \frac{\left(FC_P^{\max} - 1\right) \cdot t^s}{h^s + t^s} + b \,,$$ (15a)

where:

$$LFC_P^{\max} = \left(\log_2 \left[\Gamma \cdot (1 - \rho_{sta}^{\max}) + \Gamma \cdot \rho_{sta}^{\max} \cdot K_0 \cdot e^{-(m_{38} \cdot D_{\sigma 38} - m_{70} \cdot D_{\sigma 70})} \right] \right) \cdot \alpha - \beta \qquad (15b)$$

$$FC_P^{\max} = \frac{\left[\Gamma \cdot (1 - \rho_{sta}^{\max}) + \Gamma \cdot \rho_{sta}^{\max} \cdot K_0 \cdot e^{-(m_{38} \cdot D_{\sigma 38} - m_{70} \cdot D_{\sigma 70})} \right]^{\alpha}}{2^{\beta}} \qquad (15c)$$

Here, $\rho_{sta}^{\max}$ is the final (thus, maximum) concentration of RNAP.σ^{38} relative to the total

concentration of bound RNAPs. Meanwhile, $FC_P^{\max}$ is the expected fold change in

protein numbers of a σ^{70+38} gene, which is reached after the transition to stationary

growth is complete. Finally, *b* (the intercept) is the expression level controlled by the

promoter of interest, when in exponential growth phase. Meanwhile, *s* (the slope) is

the corresponding response strength, and *h* (the half-activation coefficient) is a

measure of response time to changes in σ^{38} levels.

2.9. Model generalization

While here we only found a correlation between the promoter sequence and the gene's

response in σ^{70+38} genes, it may be that genes whose promoters have different σ

preferences exhibit similar sequence-dependent behaviors during the stresses that

they are responsive to. Thus, we generalized the model to be applicable to any stress

and responsive gene cohort. As an example, we set a model for genes responsive to

all seven σ factors of *E. coli*. For this, first, we expand reactions (R1a) and (R1b), to

include all σ factors as follows:

$\text{RNAP} + \sigma^i \xleftrightarrow{K^{\sigma i}} \text{RNAP}.\sigma^i$, where i = 70, 54, 38, 32, 28, 24 or 19 \qquad (R8a-R8g)

Next, we generalize eq. (1) to account for the σ factors in holoenzyme form:

$$\left[RNAP_{total} \right] = \sum_{i} \left[RNAP.\sigma^i \right],$$ where i = 70, 54, 38, 32, 28, 24 or 19   (16)

Given this, we generalize (R4) as follows:

$$\text{Pro} \xrightarrow{\sum_{i} \left[RNAP.\sigma^i \right] \times k_t^{\sigma i}(D_{\sigma i})} \text{Pro} + \text{RNA},$$ where i = 70, 54, 38, 32, 28, 24 or 19   (R9)

This general model can be tuned based on the numbers of each σ factor present in

the conditions considered, and the consensus sequences to each σ factor (Table S5).

In addition, following the findings in Section 2.8, it should be feasible to introduce

factors to account for the timing of the changes in the transition period.

359   3. Discussion

From information on the promoters' sequence and dynamics, we proposed and

validated a sequence-dependent, kinetic model of genes controlled by promoters that

are responsive to both σ^{70} and σ^{38}. This model of σ^{70+38} genes, which accurately

predicts how their dynamics change for exponential and stationary growth, is an

expansion of a past model of promoters with preference for σ^{70} alone [26], but has two

competing reactions for transcription by RNAP when bound to σ^{70} and when bound to

σ^{38}. In addition, these reactions' rate constants are sequence-dependent, in line with

the hypothesis that the consensus sequence of promoters with preference for a given

σ factor should provide the strongest affinity to that σ factor. The model also accounts

for the period of the growth phase transition of the cell, which is linked to σ^{38} levels

(from lowest to highest). Finally, we proposed (yet to validate) a general model,

applicable to promoters with any given σ factor(s) preference.

Relevantly, we identified a simple correlation between the promoter sequence of σ^{70+38}

genes and their response strength to the growth phase transition, making it

predictable. So far, this predictability has not been reported for any cohort of natural

promoters of *E. coli*, even in stable growth conditions. Similar relationships have only

been reported for synthetic libraries of promoter variants under stable growth

[5,6,30,42].

Since it remains challenging to predict if a sequence can act as a promoter and, if so,

with which strength and under what regulatory mechanisms [5], this predictability of

the dynamics from the sequence for the natural cohort of σ^{70+38} genes is of interest for

three main reasons. First, these promoters and their variants (with varying p-distances

from the consensus sequences of σ^{70} and σ^{38} genes) could become key components

of a toolbox for engineering circuits whose response kinetics is predictable from the

promoter sequences. Further, such circuits should be functional in both exponential

and stationary growth phases, with tunable responsiveness to the phase transition.

Finally, their model could be a starting point for broader models of sequence-

predictable adaptability of natural promoters to various stresses.

It may prove difficult to find similar relationships between the sequences of other

natural promoters and their responsiveness to other σ factors (or global regulators).

E.g., we failed to find a relationship between the sequences of promoters responsive

to σ^{38} alone and their response strength to σ^{38} (even for genes without TF inputs (Fig

5b)). Also, it remains to be proved whether promoters responsive to σ^{70}, σ^{54}, σ^{32}, σ^{28},

σ^{24}, or dual combinations, have similar behaviors when those σ factors change.

It also remains to be studied whether the sequence-dynamics relationship in σ^{70+38}

genes is causal or coincidental. While the large numbers of promoters studied here

support the causality, a definitive proof likely requires producing synthetic promoters

similar to the natural ones but differing in p-distances to the consensus sequences of

promoters with preference for σ^{38} and for σ^{70}, respectively.

Also, in the future, it should be investigated if the sequence-dependent

responsiveness of σ^{70+38} genes to σ^{38} levels contributes to why their promoters (from

positions -41 to -1) are highly conserved (Fig S1 in S2 Appendix) and/or why the TFs

that they code for commonly serve as input TFs to essential genes [43] (2.5 times

more than average, Fisher test p-value < 0.05).

Finally, over-representation tests of the ontology [44,45] of σ^{70+38} genes suggest that

they are commonly involved in respiration (Fig S4 in S2 Appendix, Table S4 in S3

Appendix). In agreement, this process is highly affected when changing from

exponential to stationary growth [23,46], since aerobic respiration is reduced, while

fermentation and anaerobic respiration are enhanced [18]. Moreover, we observed

that σ^{70+38} genes are amongst the most responsive to the change from growth phase

out of the genes associated with these functions (Fig S4 in S2 Appendix). This

suggests that they may control the processes that need most alterations during the

adaptation. Therefore, externally regulating these genes may thus give significant

control over these processes. This is particularly appealing since this control could be

exerted by tuning their promoter sequences (i.e., their p-distances) making the effects

of their integration on a genetic circuit's response to the growth phase shift largely

predictable as well. This strategy may thus contribute to the engineering of synthetic

circuits with tailored responses to growth phase transitions.

4. Materials and Methods

4.1. Bacterial strains and media

E. coli strains and plasmids are listed in Tables S1 and S2 in S3 Appendix. In short,

we used YFP strains from the genetic stock center (CGSC) of Yale University, U.S.A

[32] and, as support, a low-copy plasmid fusion library of fluorescent reporter strains

using GFP to track promoter activity [47]. In both, one measures the levels of

fluorescent proteins under the control of the promoters of our genes of interest which

are a good proxy for the native protein levels [35][39]. For simplicity, we refer to these

fluorescence levels as 'protein levels'.

We used a RL1314 strain (rpoC::GFP) generously provided by Robert Landick,

University of Wisconsin-Madison [48]), to measure RNA Polymerase levels. Their

rpoC gene codes for β' sub-unit endogenously tagged with GFP. Since *rpoC* codes for

the β' subunit, a limiting factor in the assembly of RNAP holoenzyme [49,50], its

numbers serve as a good proxy for RNAP numbers. For simplicity, [RNAP] refers to

the sum of RNAP core and holoenzymes in a cell. Further, we used a MG*mCherry*

(*rpoS*::mCherry) strain to measure RpoS levels (kind gift from James Locke [19]). Their

rpoS gene codes for σ^{38} endogenously tagged with mCherry. Finally, we used wild-

type K12 MG1655 strain for control.

We used M9 medium (1xM9 Salts, 2 mM MgSO4, 0.1 mM CaCl2; 5xM9 Salts 34 g/L

Na2HPO4, 15 g/L KH2PO4, 2.5 g/L NaCl, 5 g/L NH4Cl) supplemented with 0.2%

Casamino acids and 0.4% glucose, and Luria-Bertani (LB) medium with 10 g peptone,

10 g NaCl, and 5 g yeast extract in 1000 ml distilled water. We used the antibiotics

kanamycin and chloramphenicol from Sigma Aldrich, U.S.A.

4.2. Growth rates and growth phases

Growth rates were measured by spectrophotometry (BioTek Synergy HTX Multi-Mode

Microplate Reader). From a glycerol stock (-80 °C), cells were streaked on LB agar

plates (2%) and incubated at 37 °C, overnight. Next, a single colony was picked,

inoculated in LB medium with antibiotics (Section 4.1), and incubated at 30°C

overnight with shaking. Overnight cultures were further diluted into fresh medium to

an optical density of 600 nm (O.D.$_{600}$) of 0.01 and incubated for growth by shaking at

250 rpm at 37°C. OD$_{600}$ was recorded every 20 min, for 800 min. Cells were extracted

at 150 min and at 700 min after inoculation into fresh medium to represent cells in

exponential and stationary growth phases, respectively (Fig 1A).

4.3. Microscopy and Image analysis

To collect microscopy data, cells were placed between a coverslip and agarose gel

pad (2%) and visualized by a confocal laser-scanning system, using a 100× objective.

Green fluorescence images were captured using a 488 nm laser and a 514/30 nm

emission filter. Phase-contrast images were simultaneously acquired for purposes of

segmentation and to assess health, morphology, and physiology.

Images were analyzed using "CellAging" software [51]. After automatically segmenting

cells and manually correcting errors, we applied 2D Gaussian filters to remove

measurement noise, and extracted each cell's total fluorescent intensity.

4.4. RNA-seq

4.4.1. Sample preparation

Cells were grown until reaching either the exponential or stationary growth phases, respectively. At each of these moments, they were treated with a double volume of RNA protect bacteria reagent (Qiagen, Germany) for 5 min, at room temperature, to prevent RNA degradation. Cells were then pelleted and frozen immediately at -80 ºC. After unfreezing the cells, total RNA was extracted using RNeasy mini-kit (Qiagen) according to the instructions. RNA was treated twice with DNase (Turbo DNA-free kit, Ambion, Life Technologies, U.S.A.) and quantified using Qubit 2.0 Fluorometer RNA assay (Invitrogen, Carlsbad, CA, USA). Quality of total RNA was determined by gel electrophoresis, using 1% agarose gel stained with SYBR safe (Invitrogen, U.S.A.). RNA was detected using UV with a Chemidoc XRS imager (BioRad, U.S.A.).

4.4.2. Sequencing

Sequencing was performed by GENEWIZ, Inc. (Leipzig, Germany). The RNA integrity number (RIN) of the samples was obtained with Agilent 4200 TapeStation (Agilent Technologies, Palo Alto, CA, USA). Ribosomal RNA depletion was performed using Ribo-Zero Gold Kit (Bacterial probe) (Illumina, San Diego, CA, USA). RNA-seq libraries were constructed using NEBNext Ultra RNA Library Prep Kit (NEB, Ipswich, MA, USA). Sequencing libraries were multiplexed and clustered on 1 lane of a flow-cell. Samples were sequenced using single-index, 2x150bp paired-end (PE) configuration on an Illumina HiSeq instrument. Image analysis and base calling were conducted with HiSeq Control Software (HCS). Raw sequence data (.bcl files) were converted into fastq files and de-multiplexed using Illumina bcl2fastq v.2.20. One mismatch was allowed for index sequence identification.

4.4.3. Data analysis

RNA-seq data analysis pipeline: i) RNA sequencing reads were trimmed to remove

adapter sequences and nucleotides with poor quality with Trimmomatic [52] v.0.39. ii)

Trimmed reads were then mapped to the reference genome (*E. coli* MG1655,

NC_000913.3) using STAR v.2.5.2b aligner [53]. iii) *featureCounts* from Rsubread R

package (v.1.34.7) was used to calculate unique gene hit counts [54]. Genes with less

than 5 counts in more than 3 samples, and genes whose mean counts are less than

10, were removed. iv) Next, the unique gene hit counts were used for subsequent

differential expression analysis, using DESeq2 R package (v.1.24.0) [55] to compare

gene expression between samples and calculate p-values and log2 of fold changes

(LFC) using Wald tests (function *nbinomWaldTest*). P-values were adjusted for

multiple hypotheses testing (Benjamini–Hochberg, BH procedure [56]) and genes with

adjusted p-values (False discovery rate (FDR)) smaller than 0.05 were classified as

differentially expressed (D.E.).

Due to filtering (iii), only 783 (777 D.E.) out of the 931 σ^{70} genes, 87 (all D.E.) out of

the 93 σ^{38} genes, 57 (56 D.E.) out of the 64 σ^{70+38} genes, and 3257 (2737 D.E.) out of

3607 remaining genes were assessed by RNA-seq. Other genes did not produce

enough reads.

4.5. Flow cytometry

We used a ACEA NovoCyte Flow Cytometer equipped with yellow (561 nm) and blue

lasers (488 nm), controlled by Novo Express (V1.50). Cells were diluted 1:10000 into

1 ml of PBS vortexed for 10 sec. For each gene and condition, we performed 3

biological replicates, acquiring 50,000 events in each. For GFP and YFP, we used the

blue laser for excitation and the FITC-H channel (520/20 nm filter) for emission. For

mCherry, we used a yellow laser for excitation and PE-Texas Red (615/20 nm filter)

for emission. We collected events at a flow rate of 14 µl/minute, a core diameter of

7.7 µM, and adjusted PMT voltage for each parameter. We have set the FSC-H

detection threshold to 5000.

We collected single-cell fluorescence distributions. To remove outliers due to debris,

cell doublets, and other undesired events, we performed gating by setting the

maximum values of SSC-H to $5{\times}10^4$ and of FSC-H to 10^5. These limits allowed

observing the whole distributions, without cutting their natural tails. The results did not

differ widely from using unsupervised gating as in [57,58], if setting α to 0.95.

Finally, we discarded "far out" events using Tukey's fences [59]. This had little effect

on the results. Furthermore, background fluorescence was not subtracted since doing

so did not affect the results (Section S1.2 in S1 Appendix and Fig S2 in S2 Appendix).

4.6. Spectrophotometry

Time-lapse protein fluorescence of MG*mCherry* cells was measured by a BioTek

Synergy HTX Multi-Mode Microplate Reader. Overnight cultured cells were diluted

1:1000 times into fresh M9 medium, aliquoted into 24 well dark bottom microplates,

and kept at 37°C with shaking of 250 rpm. Fluorescence intensities were recorded for

14 hours, every 20 min, using excitation (590/20 nm) and emission (645/20 nm) filters.

Temporal profiles of fluorescence intensity were extracted using Gen5 software, based

on 6 biological repeats.

4.7. Statistical analyses of the data

Data points from microscopy, flow cytometry, etc. are based on at least 3 biological

replicates each. Meanwhile, to assess if genes measured by flow cytometry were a

good representative of the set of all σ^{70+38} genes, we used two-sample Kolmogorov-

Smirnov tests (KS tests) to compare their sequences ($D_{\sigma 38}$ and $D_{\sigma 70}$) and dynamics

(LFC_{RNA}, μ, and CV^2). Further, gaussian fittings were made by the Distribution Fitter

app (MATLAB). To evaluate its fitting, we calculated the coefficient of determination

(R^2) between the PDF of the distributions and the PDF of the fitting. Finally, linear

correlations were estimated by least-squares fitting (FITLM of MATLAB) and

considered significant if the *p*-value of an *F*-test on the fitted regression line was

smaller than 0.05. Other curves were fitted using MATLAB's curve fitting toolbox (apart

from the fitting of Hill functions, which was done using the Hill function [60]).

As for gene ontology representations, we perform overrepresentation tests using

PANTHER Classification [61] for finding significant overrepresentations by Fisher's

exact tests. For *p*-values < 0.05, the null hypothesis that there are no associations

between the gene cohort and the corresponding GO of the biological process is

rejected. This *p*-value is corrected by calculating if the overall False Discovery Rate

(FDR) is < 0.05. FDR is calculated by the Benjamini-Hochberg procedure [56].

4.8.   Features of the promoter sequences

4.8.1. σ factor preference

Table S3 in S3 Appendix informs on the σ factor preference of the genes. From

Regulon DB v10.5 (August 14, 2019), we obtained lists of all transcription units (TUs),

promoters (including σ factor preferences), and genes of *E. coli* [62]. Recently, we

compared our lists with information on July 1st, 2021 and found no changes that would

affect the conclusions. TUs only differed by ~1%, promoters by ~0.5%, and genes by

~1%.

From the 3548 TUs (gene(s) transcribed from a single promoter), we extracted 2179

with known promoters, containing 2713 genes in total. To minimize interference to the

classification of σ factor preferences arising from transcription by multiple promoters,

we narrowed our list to 1824 genes transcribed by only one promoter and, of those, to

the 1328 genes with known σ factor preference. From those, 1242 have a preference

for only one σ factor, including 931 with a preference for σ^{70}, and 93 with a preference

for σ^{38}. Conversely, 76 genes have promoters with a preference for two σ factors. Out

of these, 64 are transcribed by only one promoter with a preference for σ^{70} and σ^{38}.

4.8.2. Promoter sequence logos

Promoter sequence logos were created using WebLogo [63]. In each position (from -

41 to -1), we counted how many times a nucleotide is present in all promoters

considered. Then, we stacked all nucleotides (A, C, T, G) on top of each other, and

sorted from the least found one in the bottom to the most present one on the top. For

each position, we quantified its 'bit', as the difference between the maximum

information possible (entropy given the 4 nucleotides) and the information considering

the variability of the nucleotides (sum of the entropy for each of the 4 nucleotides) in

that position (observed entropy): $\text{Bit} = \log_2 4 - \sum_{n=1}^{4} f_n \times \log_2(f_n)$. From this, the height

of each letter is set to be proportional to its frequency of occurrence. Finally, the height

of each position was normalized, so as to equal its corresponding amount of

information (with the more conserved positions having more bits) [64].

4.8.3. Promoter sequence p-distance

The p-distance D of a promoter [65] is the fraction of its nucleotides between positions

-41 to -1 (assuming that transcription start site starts at position +1) that differ from the

consensus (most common) nucleotide in that position of a cohort of genes (here,

genes with preference for a given σ factor). We extracted the consensus sequences

related to each σ from RegulonDB [62]. To measure the D of promoters with

preference for both σ^{38} and σ^{70}, we calculated the p-distances $D_{\sigma 38}$ and $D_{\sigma 70}$, and then

an overall p-distance, Δ, defined as: $\Delta = D_{\sigma 38} - D_{\sigma 70}$.

5. Supporting Information

S1 Appendix. Extended Methods and Materials. (Word)

S2 Appendix. Supporting Figure. (Word)

S3 Appendix. Supporting Tables. (Word)

S4 Table. List of genes classified as σ^{70}, and σ^{38} or as both σ^{70} and σ^{38} dependent

(Excel)

S5 Table. Consensus sequences of promoters with preference for one σ factor.

(Excel)

S6 Table. Fold changes in RNA levels of genes with a promoter with preference for

σ^{70} and σ^{38}. Measurements by RNA-seq in the exponential and stationary growth

phases. (Excel)

S7 Table. Statistics of single-cell distributions of fluorescence of cells measured by

flow cytometry. (Excel)

6. References

1.   Chen J, Chiu C, Gopalkrishnan S, Chen AY, Olinares PDB, Saecker RM, et al.
Stepwise Promoter Melting by Bacterial RNA Polymerase. Mol Cell. 2020;78: 275-
288.e6. doi:10.1016/j.molcel.2020.02.017

2.   Browning DF, Busby SJW. Local and global regulation of transcription initiation in
bacteria. Nat Rev Microbiol. 2016;14: 638–650. doi:10.1038/nrmicro.2016.103

3.   Browning DF, Busby SJW. The regulation of bacterial transcription initiation. Nat
Rev Microbiol. 2004;2: 57–65. doi:10.1038/nrmicro787

4.   Cho BK, Kim D, Knight EM, Zengler K, Palsson BO. Genome-scale reconstruction
of the sigma factor network in Escherichia coli: Topology and functional states.
BMC Biol. 2014;12: 1–11. doi:10.1186/1741-7007-12-4

5.   Urtecho G, Tripp AD, Insigne KD, Kim H, Kosuri S. Systematic Dissection of
Sequence Elements Controlling σ70 Promoters Using a Genomically Encoded
Multiplexed Reporter Assay in Escherichia coli. Biochemistry. 2019;58: 1539–
1551. doi:10.1021/acs.biochem.7b01069

6.   Einav T, Phillips R. How the avidity of polymerase binding to the –35/–10 promoter
sites affects gene expression. Proceedings of the National Academy of Sciences.
2019;116: 13340 LP – 13345. doi:10.1073/pnas.1905615116

7.   Jones DL, Brewster RC, Phillips R. Promoter architecture dictates cell-to-cell

variability   in   gene   expression.   Science.   2014;346:   1533–1536.

doi:10.1126/science.1255301

8.   Stoebel DM, Hokamp K, Last MS, Dorman CJ. Compensatory evolution of gene

regulation in response to stress by Escherichia coli lacking RpoS. PLoS Genet.

2009;5: e1000671. doi:10.1371/journal.pgen.1000671

9.   Kannan G, Wilks JC, Fitzgerald DM, Jones BD, Bondurant SS, Slonczewski JL.

Rapid acid treatment of Escherichia coli: transcriptomic response and recovery.

BMC Microbiol. 2008;8: 37. doi:10.1186/1471-2180-8-37

10.   Farewell A, Kvint K, Nyström T. Negative regulation by RpoS: A case of sigma

factor   competition.   Mol   Microbiol.   1998;29:   1039–1051.   doi:10.1046/j.1365-

2958.1998.00990.x

11.   Rouvière PE, De Las Peñas A, Mecsas J, Lu CZ, Rudd KE, Gross CA. rpoE, the

gene encoding the second heat-shock sigma factor, sigma E, in Escherichia coli.

EMBO J. 1995;14: 1032–1042. doi:10.1002/j.1460-2075.1995.tb07084.x

12.   Dong T, Schellhorn HE. Global effect of RpoS on gene expression in pathogenic

Escherichia coli O157:H7 strain EDL933. BMC Genomics. 2009;10: 349.

doi:10.1186/1471-2164-10-349

13.   Rahman M, Hasan MR, Oba T, Shimizu K. Effect of rpoS gene knockout on the

metabolism of Escherichia coli during exponential growth phase and early

stationary phase based on gene expressions, enzyme activities and intracellular

metabolite   concentrations.   Biotechnol   Bioeng.   2006;94:   585–595.

doi:10.1002/bit.20858

14. Maeda H, Fujita N, Ishihama A. Competition among seven Escherichia coli σ

subunits: Relative binding affinities to the core RNA polymerase. Nucleic Acids

Res. 2000;28: 3497–3503. doi:10.1093/nar/28.18.3497

15. Feklístov A, Sharon BD, Darst SA, Gross CA. Bacterial sigma factors: A historical,

structural, and genomic perspective. Annu Rev Microbiol. 2014;68: 357–376.

doi:10.1146/annurev-micro-092412-155737

16. Helmann JD, Chamberlin MJ. Structure and function of bacterial sigma factors.

Annu   Rev   Biochem.   1988;57:   839–872.

doi:10.1146/annurev.bi.57.070188.004203

17. Hengge R. Proteolysis of σS (RpoS) and the general stress response in

Escherichia   coli.   Research   in   Microbiology.   2009.   pp.   667–676.

doi:10.1016/j.resmic.2009.08.014

18. Weber H, Polen T, Heuveling J, Wendisch VF, Hengge R. Genome-wide analysis

of the general stress response network in Escherichia coli: sigmaS-dependent

genes, promoters, and sigma factor selectivity. J Bacteriol. 2005;187: 1591–1603.

doi:10.1128/JB.187.5.1591-1603.2005

19. Patange O, Schwall C, Jones M, Villava C, Griffith DA, Phillips A, et al.

Escherichia coli can survive stress by noisy growth modulation. Nat Commun.

2018;9. doi:10.1038/s41467-018-07702-z

20. Jishage M, Iwata A, Ueda S, Ishihama A. Regulation of RNA polymerase sigma

subunit synthesis in Escherichia coli: Intracellular levels of four species of sigma

subunit under various growth conditions. J Bacteriol. 1996;178: 5447–5451.

doi:10.1128/jb.178.18.5447-5451.1996

21. Lange R, Hengge-Aronis R. The cellular concentration of the σ(S) subunit of RNA

polymerase in Escherichia coli is controlled at the levels of transcription,

translation, and protein stability. Genes and Development. 1994;8: 1600–1612.

doi:10.1101/gad.8.13.1600

22. Gruber TM, Gross CA. Multiple Sigma Subunits and the Partitioning of Bacterial

Transcription　　Space.　　Annu　　Rev　　Microbiol.　　2003;57:　　441–466.

doi:10.1146/annurev.micro.57.030502.090913

23. Riedel TE, Berelson WM, Nealson KH, Finkel SE. Oxygen consumption rates of

bacteria under nutrient-limited conditions. Appl Environ Microbiol. 2013;79: 4921–

4931. doi:10.1128/AEM.00756-13

24. Abantika Ganguly DC. A Comparative Kinetic and Thermodynamic Perspective

of the σ-Competition Model in Escherichia coli. Biophys J. 2012;103: 1325.

doi:10.1016/j.bpj.2012.08.013

25. Dennis PP, Bremer H. Modulation of Chemical Composition and Other

Parameters of the Cell at Different Exponential Growth Rates. EcoSal Plus.

2008;3. doi:10.1128/ecosal.5.2.3

26. Kandavalli VK, Tran H, Ribeiro AS. Effects of σ factor competition are promoter

initiation kinetics dependent. Biochim Biophys Acta. 2016;1859: 1281–1288.

doi:10.1016/j.bbagrm.2016.07.011

27. Grigorova IL, Phleger NJ, Mutalik VK, Gross CA. Insights into transcriptional

regulation and sigma competition from an equilibrium model of RNA polymerase

binding to DNA. Proc Natl Acad Sci U S A. 2006;103: 5332–5337.

doi:10.1073/pnas.0600828103

28. Mauri M, Klumpp S. A Model for Sigma Factor Competition in Bacterial Cells.

PLoS Comput Biol. 2014;10: e1003845. doi:10.1371/journal.pcbi.1003845

29. Kinney JB, Murugan A, Callan CG, Cox EC. Using deep sequencing to

characterize the biophysical mechanism of a transcriptional regulatory sequence.

Proc Natl Acad Sci U S A. 2010;107: 9158–9163. doi:10.1073/pnas.1004290107

30. Brewster RC, Jones DL, Phillips R. Tuning Promoter Strength through RNA

Polymerase Binding Site Design in Escherichia coli. PLoS Comput Biol. 2012;8:

e1002811. doi:10.1371/journal.pcbi.1002811

31. Tanaka K, Takayanagi Y, Fujita N, Ishihama A, Takahashi H. Heterogeneity of

the principal sigma factor in Escherichia coli: the rpoS gene product, sigma 38, is

a second principal sigma factor of RNA polymerase in stationary-phase

Escherichia coli. Proc Natl Acad Sci U S A. 1993;90: 3511–3515.

doi:10.1073/pnas.90.8.3511

32. Taniguchi Y, Choi PJ, Li G-W, Chen H, Babu M, Hearn J, et al. Quantifying E. coli

Proteome and Transcriptome with Single-Molecule Sensitivity in Single Cells.

Science. 2010;329: 533–538. doi:10.1126/science.1188308

33. Hausser J, Mayo A, Keren L, Alon U. Central dogma rates and the trade-off

between precision and economy in gene expression. Nat Commun. 2019;10: 1–

15. doi:10.1038/s41467-018-07391-8

34. Bar-Even A, Paulsson J, Maheshri N, Carmi M, O'Shea E, Pilpel Y, et al. Noise in

protein expression scales with natural protein abundance. Nat Genet. 2006;38:

636–643. doi:10.1038/ng1807

35. Newman JRS, Ghaemmaghami S, Ihmels J, Breslow DK, Noble M, DeRisi JL, et

al. Single-cell proteomic analysis of S. cerevisiae reveals the architecture of

biological noise. Nature. 2006;441: 840–846. doi:10.1038/nature04785

36. Ishihama A. Functional modulation of Escherichia coli RNA polymerase. Annu

Rev Microbiol. 2000;54: 499–518. doi:10.1146/annurev.micro.54.1.499

37. Jishage M, Ishihama A. Regulation of RNA polymerase sigma subunit synthesis

in Escherichia coli: intracellular levels of sigma 70 and sigma 38. J Bacteriol.

1995;177: 6832–6835. doi:10.1128/jb.177.23.6832-6835.1995

38. Piper SE, Mitchell JE, Lee DJ, Busby SJW. A global view of Escherichia coli Rsd

protein and its interactions. Mol Biosyst. 2009;5: 1943–1947.

doi:10.1039/B904955J

39. McClure WR. Mechanism and Control of Transcription Initiation in Prokaryotes.

Annu   Rev   Biochem.   1985;54:   171–204.

doi:10.1146/annurev.bi.54.070185.001131

40. Muthukrishnan AB, Martikainen A, Neeli-Venkata R, Ribeiro AS. In vivo

transcription kinetics of a synthetic gene uninvolved in stress-response pathways

in   stressed   Escherichia   coli   cells.   PLoS   One.   2014;9.

doi:10.1371/journal.pone.0109005

41. Colland F, Fujita N, Ishihama A, Kolb A. The interaction between sigmaS, the

stationary phase sigma factor, and the core enzyme of Escherichia coli RNA

polymerase.   Genes   Cells.   2002;7:   233–247.   doi:10.1046/j.1365-

2443.2002.00517.x

42. Liu M, Tolstorukov M, Zhurkin V, Garges S, Adhya S. A Mutant Spacer Sequence

between -35 and -10 Elements Makes the P_{lac} Promoter Hyperactive and cAMP

Receptor Protein-Independent. Proc Natl Acad Sci U S A. 2004;101: 6911–6916.

Available: http://www.jstor.org/stable/3372027

43. Baba T, Ara T, Hasegawa M, Takai Y, Okumura Y, Baba M, et al. Construction of

Escherichia coli K-12 in-frame, single-gene knockout mutants: the Keio collection.

Mol Syst Biol. 2006;2: 2006.0008-2006.0008. doi:10.1038/msb4100050

44. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene

ontology: tool for the unification of biology. The Gene Ontology Consortium. Nat

Genet. 2000;25: 25–29. doi:10.1038/75556

45. Gene Ontology Consortium. The Gene Ontology resource: enriching a GOld

mine. Nucleic Acids Res. 2021;49: D325–D334. doi:10.1093/nar/gkaa1113

46. Andersen KB, von Meyenburg K. Are growth rates of Escherichia coli in batch

cultures   limited   by   respiration?   J   Bacteriol.   1980;144:   114–123.

doi:10.1128/JB.144.1.114-123.1980

47. Zaslaver A, Bren A, Ronen M, Itzkovitz S, Kikoin I, Shavit S, et al. A

comprehensive library of fluorescent transcriptional reporters for Escherichia coli.

Nat Methods. 2006;3: 623–628. doi:10.1038/nmeth895

48. Bratton BP, Mooney RA, Weisshaar JC. Spatial distribution and diffusive motion

of RNA polymerase in live Escherichia coli. J Bacteriol. 2011;193: 5138–5146.

doi:10.1128/JB.00198-11

49. Izard J, Gomez Balderas CDC, Ropers D, Lacour S, Song X, Yang Y, et al. A

synthetic growth switch based on controlled expression of RNA polymerase. Mol

Syst Biol. 2015;11: 840. doi:10.15252/msb.20156382

50. Chamberlin MJ. The selectivity of transcription. Annu Rev Biochem. 1974;43:

721–775. doi:10.1146/annurev.bi.43.070174.003445

51. Häkkinen A, Muthukrishnan A-B, Mora A, Fonseca JM, Ribeiro AS. CellAging: a

tool to study segregation and partitioning in division in cell lineages of Escherichia

coli. Bioinformatics. 2013;29: 1708–1709. doi:10.1093/bioinformatics/btt194

52. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina

sequence data. Bioinformatics. 2014;30: 2114. doi:10.1093/bioinformatics/btu170

53. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat

Methods. 2012;9: 357–359. doi:10.1038/nmeth.1923

54. Liao Y, Smyth GK, Shi W. The R package Rsubread is easier, faster, cheaper

and better for alignment and quantification of RNA sequencing reads. Nucleic

Acids Res. 2019;47: e47. doi:10.1093/nar/gkz114

55. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion

for RNA-seq data with DESeq2. Genome Biol. 2014;15: 550. doi:10.1186/s13059-

014-0550-8

56. Benjamini Y, Hochberg Y. Controlling the False Discovery Rate: A Practical and

Powerful Approach to Multiple Testing. J R Stat Soc Series B Stat Methodol.

1995;57: 289–300. Available: http://www.jstor.org/stable/2346101

57. Startceva S, Kandavalli VK, Visa A, Ribeiro AS. Regulation of asymmetries in the

kinetics and protein numbers of bacterial gene expression. Biochimica et

Biophysica Acta - Gene Regulatory Mechanisms. 2019;1862: 119–128.

doi:10.1016/j.bbagrm.2018.12.005

58. Razo-Mejia M, Barnes SL, Belliveau NM, Chure G, Einav T, Lewis M, et al. Tuning

Transcriptional Regulation through Signaling: A Predictive Theory of Allosteric

Induction. Cell Systems. 2018;6: 456-469.e10. doi:10.1016/j.cels.2018.02.004

59. Tukey JW. Schematic Summaries. In: Mosteller F, editor. Exploratory Data

Analysis. Reading, Massachusetts: Addison-Wesley Publishing Company; 1977.

pp. 27–56.

60. Olsen T. Hill function. In: MATLAB Central File Exchange [Internet]. 27 Sep 2018

[cited   27   Oct   2021].   Available:

https://www.mathworks.com/matlabcentral/fileexchange/68935-hill-function)

61. Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. PANTHER version 14:

more genomes, a new PANTHER GO-slim and improvements in enrichment

analysis   tools.   Nucleic   Acids   Res.   2019;47:   D419–D426.

doi:10.1093/nar/gky1038

62. Santos-Zavaleta A, Salgado H, Gama-Castro S, Sánchez-Pérez M, Gómez-

Romero L, Ledezma-Tejeida D, et al. RegulonDB v 10.5: tackling challenges to

unify classic and high throughput knowledge of gene regulation in E. coli K-12.

Nucleic Acids Res. 2018;47: D212–D220. doi:10.1093/nar/gky1077

63. Crooks G, Hon G, Chandonia J, Brenner S. NCBI GenBank FTP Site\nWebLogo:

a   sequence   logo   generator.   Genome   Res.   2004;14:   1188–1190.

doi:10.1101/gr.849004.1

64. Schneider TD, Stephens RM. Sequence logos: A new way to display consensus

sequences. Nucleic Acids Res. 1990;18: 6097–6100. doi:10.1093/nar/18.20.6097

65. Nei M, Zhang J. Evolutionary Distance: Estimation. Encyclopedia of Life

Sciences. 2006; 1–4. doi:10.1038/npg.els.0005108