In vitro culture of bovine fibroblasts using select serum-free

media supplemented with *Chlorella vulgaris* extract

Authors

Galileo Defendi-Cho (ORCID: 0000-0001-8747-8143), g_defendicho@coloradocollege.edu

*Timothy M. Gould (ORCID: 0000-0002-6858-1265), tgould@coloradocollege.edu

*Corresponding author

Affiliation

Colorado College, Department of Chemistry and Biochemistry, Colorado Springs, CO, USA.

Abstract

Standard cell culture practices require addition of animal-derived serum to culture media to

achieve adequate cell growth. Typically, 5-10% by volume of fetal bovine serum (FBS) is used,

which accounts for a vast majority of the cost of media while also imposing environmental and

ethical concerns associated with the use of animal serum. Here we tested the efficacy of

culturing cells by replacing serum in the media with algae extract and select additives. Using

LC-MS, we compared molecular signatures of FBS to *Chlorella* algae extracts and identified

NAD(H)/NADP(H) as common and relatively abundant features in their characteristic profiles.

Bovine fibroblasts, cultured in serum-free media supplemented with *C. vulgaris* extract and just

two growth factors plus insulin, showed significant growth with enhanced viability compared to

control cells cultured without serum, albeit still lower than that of controls cultured with 10%

FBS. Moreover, *C. vulgaris* extract enhanced cell viability beyond that of cells cultured with the

two growth factors and insulin alone. These results suggest that key components in serum

which are essential for cell growth may also be present in *C. vulgaris* extract, demonstrating

that it may be used at least as a partial alternative to serum for cell culture applications.

Keywords

Serum replacement, serum-free media, algae extract, bovine fibroblast, cell cultured meat

Acknowledgements

This work was supported by the Figge-Bourquin Award and by Student-Faculty Collaborative

Grants via the Creativity and Innovation and Dean's Offices at Colorado College. The authors

thank Rachel Wonciar and Tanya Cervantes in the Department of Chemistry and Biochemistry,

and Carrie Moon and Darrell Killian in the Department of Molecular Biology at Colorado

College, for their technical assistance.

Author Contributions

G.D.C. and T.M.G. designed the study, performed experiments, analyzed data, and wrote the

paper, with both authors contributing equally.

Statements and Declarations

There are no competing interests.

Introduction

In 2013 it was estimated that raising livestock contributed roughly 14% to total greenhouse gas emissions, more than half of which was attributable to meat production (Gerber et al. 2013). Cell cultured meat has the potential to offer substantial environmental and social advantages over traditional agriculture practices using livestock (Tuomisto and Teixeira De Mattos 2011; Ladak and Anthis 2021), yet analysis on the economics of cultured versus conventional meat production shows high cost barriers, particularly those associated with the nutrients required (Humbird 2021). Cell culture media is commonly supplemented with fetal bovine serum (FBS) because animal serum promotes cell growth (Penttinen and Saxén 1959). Based on current commercial prices, the cost of culture-grade FBS is estimated at around $1.20 per mL and that of other media components at $0.04 per mL; thus, for typical complete growth media containing 10% FBS, most of its total cost is attributed to FBS, by roughly three-fold. Therefore, low cost and environmentally favorable alternatives to serum are attractive for cell culture applications, particularly for the cultured meat industry.

Because animal-derived serums like FBS are very complex, including thousands of different molecules that can be highly varied in their relative abundance due to many dynamic physiological factors, it is unlikely that synthetic approaches to serum alternatives will prove cost effective if technically feasible. Researchers have studied the culture of cells in serum-free conditions for many decades with varying degrees of success, yet most cell culture practices continue using FBS. Following studies which identified key components necessary to culture cells in the absence of serum (Murakami and Masui 1980; Chen et al. 2011), there are now chemically-defined serum-free media formulations that are commercially available (e.g., E8™).

However, the cost of these serum-free formulations is still quite high, primarily due to the

protein growth factors required. Such formulations were developed mainly for the purpose of

consistently culturing human induced pluripotent stem cells (iPSCs); they were not originally

intended for cultured meat applications although this does typically involve culturing stem cells.

Nevertheless, determining the minimal components and concentrations at which they are

needed to achieve adequate and consistent cell growth remains an important undertaking for

the cultured meat industry (Specht 2020).

We were intrigued by a recent study in which the authors reported growth effects of

algae extract on mammalian cells (Ng et al. 2020). Ng and colleagues report that extract

derived from *Chlorella vulgaris* promoted growth of Chinese hamster ovary (CHO) and

mesenchymal stem cells (MSC) under certain conditions, some of which did not contain any

serum. In another study, researchers reported high levels of glucose and amino acids present

in various microalgae extracts, with particularly high levels of aspartate and glutamate detected

in extract from *C. vulgaris* (Okamoto et al. 2020); these authors proceeded to show that they

could use such algae extracts as a substitute to culture cells in media deficient in glucose and

amino acids. More recently, this same group showed that *C. vulgaris* extract can be used as a

replacement for DMEM in the culture of primary bovine myoblasts (Okamoto et al. 2022).

Here we tested the efficacy of culturing cells by replacing serum in the media with algae

extract and found that we could achieve appreciable growth and viability of bovine fibroblasts

using select serum-free media supplemented with *C. vulgaris* extract (CVE). We characterized

molecular features of algae extracts, compared them to that of FBS, and found evidence

suggesting that NAD(H)/NADP(H) are common and relatively abundant components. This study

establishes how algae extract may be used to culture cells in serum-free conditions while

providing further insight into essential components required for cell growth.

Materials and Methods

<u>Chemicals and Reagents</u>

Liquid chlorella growth factor (SKU 504) and *Chlorella vulgaris* extract capsules (SKU 583) were

purchased from BioPure® Healing Products (Woodinville, WA). Pierce™ universal nuclease for

cell lysis was purchased from Thermo Fisher Scientific Corporation (Waltham, MA).

Recombinant human fibroblast growth factor (FGF-2, catalog no. 100-18B) and transforming

growth factor beta (TGF-β, catalog no. 100-21) were purchased from PeproTech, Inc. (Cranbury,

NJ). Bovine insulin, sodium L-ascorbate and selenium dioxide were from Sigma-Aldrich. Cell

titer 96® aqueous one solution cell proliferation (MTS) assay (catalog no. G3580) was purchased

from Promega Corporation (Madison, WI).

<u>Fractionation of Algae Extracts</u>

Algae extracts termed *Chlorella vulgaris* extract (CVE) and chlorella growth factor (CGF) were

derived from two different strains, *Chlorella vulgaris* (CVE) and *Chlorella pyrenoidosa* (CGF).

CVE was obtained from capsules in dry powder form and dissolved directly in serum-free

DMEM. CGF stock solution was purchased as a concentrated liquid hot-water extract from

which all solutions were made on a percent volume basis by diluting CGF directly in serum-free

DMEM. For fractionation experiments, crude solutions of 10 g/L CVE and 10% CGF were first

prepared in serum-free DMEM. Crude CVE and CGF were then either sterile-filtered through

0.22 μm nylon filters or centrifuged at 10,000 x *g* for 10 minutes in a benchtop microcentrifuge.

After centrifugation, supernatants were removed, and the pellets were resuspended directly in

serum-free DMEM.

<u>LC-MS</u>

Sample separation was performed by liquid chromatography (LC) using a UPLC BEH C18 column

(2.1 x 50 mm, 1.7 μm) fitted to an Acquity UPLC system with PDA detector (Waters).

Chromatographic profiles were acquired at wavelengths 200-400 nm. Injection volume was 7.5

μL after equilibration, and the column temperature was held at 30°C. Samples were eluted

from the column using a gradient mobile phase that consisted of phase A (0.1% formic acid in

water) and phase B (0.1% formic acid in acetonitrile). The linear gradient elution procedure

was as follows: (step 1) 100% A from 0-3 minutes, (step 2) 100% to 60% A from 3-20 minutes,

(step 3) 60% to 0% A from 20-21 minutes, (step 4) 0% A from 21-24 minutes, (step 5) 0% to

100% A from 24-25 minutes. The column flow rate was set at 0.2 mL/min from 0-21 minutes,

increased to 0.6 mL/min from 21-24 minutes, and then returned to the initial rate from 24-25

minutes. Electrospray ionization mass spectrometry (ESI-MS) was performed using an Acquity

QDa detector (Waters). Capillary and cone voltages were 0.8 kV and 10 V, respectively. MS

measurements were collected in positive ion mode with a mass scan range from 100-1200 m/z.

Cell Culture

The EBTr (NBL-4) cell line was purchased from ATCC (CCL-44[TM]). Cells were routinely cultured at

37°C with 5% CO_2 in complete Dulbecco's Modified Eagle's Medium (DMEM) supplemented

with 10% FBS (Gibco), 100 U/mL penicillin and 100 μg/mL streptomycin (Thermo Fisher

Scientific, catalog no. 15140148); serum-free DMEM included antibiotics but no FBS. Passaging

was routinely done as follows: cells were first washed with 1X phosphate-buffered saline (PBS),

then placed in an incubator with 0.25% Trypsin-EDTA solution (Gibco) for approximately 2

minutes immediately after which complete DMEM was added to quench trypsin activity; cells

were collected and centrifuged at 300 x g for 5 minutes, and then the cell pellet was

resuspended in fresh media. EBTr cells were passaged every 3-4 days at a 1:3 passage ratio.

Excess cells were stored in liquid nitrogen at 1-2x10^6 cells/mL in freezing media containing

90:10% FBS:DMSO. The number of passages for experiments was limited to less than twenty.

<u>Viability (MTS) Assays</u>

EBTr cells were first harvested, counted, and resuspended at a density of 2.5x10^5 cells/mL in

serum-free DMEM. Cells were then seeded in clear, flat-bottom 96-well plates to achieve a

target density of approximately 2x10^4 cells (80 μL) per well. Seeded cells were incubated for 2-

4 hours to allow sufficient time for cells to adhere to plates prior to experiments. For all

viability experiments/assays, each condition corresponded to a single column in a 96-well plate.

Test media conditions were achieved by delivering 20 μL of 5X test solutions (prepared in

serum-free DMEM) to the appropriate column wells, bringing total well volumes to 100 μL.

Columns 1 and 12 as well as the top and bottom rows of all 96-well plates were treated as mock

wells and excluded from data analysis to control for plate edge effects (Burt et al. 1979).

Immediately after adding 5X test solutions to plates, cells were returned to the incubator for

the experiment duration (typically 3 days or approximately 72 hours) after which 20 μL of MTS

reagent was added to each plate well. Plates were then incubated for another 4 hours. (For the

experiment shown in Fig. 2d, only 10 μL of MTS reagent was added to plate wells because there

was not enough reagent to add 20 μL to all 96 wells.) End-point absorbance measurements

were taken with a plate reader at 490 nm (BioTek Instruments, Agilent Technologies).

<u>Cell Counting and Imaging</u>

Cells were harvested, pelleted, and resuspended in serum-free DMEM. Two replicate 50 μL

samples of cell suspensions were each diluted 1:1 in trypan blue dye, from which 10 μL was

sampled and pipetted onto a hemocytometer (improved Neubauer ruling pattern). Live and

dead/blue (non-intact) cell counts were performed in duplicate (i.e., one count per sample).

Brightfield images were obtained using an inverted tissue culture microscope (Hund Wetzlar)

fitted with a fixed microscope adapter connected to a 5.1-megapixel digital camera (AmScope).

Original grayscale image tiles were merged into a single multi-panel image representing each

experiment. Multi-panel images were processed with Fiji/ImageJ2 open-source software (NIH).

Image processing involved pseudo flat field correction to balance uneven illumination across

images, Fast Fourier Transform (FFT) bandpass filtering to reduce image scan lines, and

brightness/contrast adjustments.

Data Analyses and Statistics

LC-MS datasets, including both matrix LC and MS data from 3 replicate injections of the same

sample type, were first averaged and normalized; mean data matrices were then visualized

using R software (v4.1.0) with viridis and ggplot2 packages. Principal component analysis (PCA)

on averaged MS data was performed in R using the prcomp function. Potential parent species

were identified in a mass spectral database (MassBank.eu) by searching the m/z value and

average relative intensity of peaks shared between samples. All other data were analyzed

and/or visualized using Prism software (v9.2.0, GraphPad), in which two-sided t tests with

Welch's correction were used to determine statistical significance between compared groups.

Results

Molecular features of FBS and algae extracts show both distinct and common components.

Given their potential growth effects, we sought to compare algae extracts (CVE and CGF,

each of which was derived from a different *Chlorella* strain) to each other and FBS using LC-MS.

Due to limited scope and the molecular complexity of these samples, our aim was not to look

for specific compounds but rather to gain a qualitative profile for each sample type. Fig. 1a

shows LC chromatograms of FBS compared to CVE and CGF samples. Large broad peaks around

260 nm for species that eluted off the column within the first few minutes were observed in

both CVE and CGF but not FBS samples; these large peaks did not correspond to any

appreciable signal(s) detected by the mass spectrometer at such early elution times (Fig. 1b).

All samples showed both substantial absorbance (between 200-250 nm) and mass signals at

retention times from approximately 18-21 minutes; the relative magnitude of this cluster of

signals was comparable for FBS and CVE samples, whereas that of CGF samples was lower. Fig.

1c shows the results of a principal component analysis (PCA) comparing total ion

chromatograms (TICs) of all FBS, CVE and CGF samples (i.e., replicates averaged in Fig. 1b).

Replicate injections of the same sample type clustered together and separated in PC1 but not

PC2 values; moreover, CVE and FBS samples were more similar in PC4 values than CGF samples.

We also visualized how signals at each *m/z* value and retention time compared across samples

(Fig. 1d). Each sample type showed a common pattern with a cluster of relatively high signal

peaks corresponding to *m/z* values ranging from roughly 800-1200 and retention times from

about 15-21 minutes. Both CVE and CGF samples also shared signal features at lower *m/z*

values (mainly over the 400-800 range) and retention times from about 5-15 minutes; this

broad cluster of signals was less apparent if not absent in FBS samples. We further examined

the most striking MS feature found in all samples, i.e., the cluster of signals at m/z values from

about 800-1200 and retention times from 15-21 minutes. Fig. 1e highlights five relatively

abundant and shared peaks within this cluster, noting their m/z whole integer value, relative

abundance, and corresponding potential parent ion identity(s). The top candidate identified by

a mass spectral database search (MassBank.eu) for the most abundant shared peak (no. 3 in

Fig. 1e) was nicotinamide adenine dinucleotide (NAD^+) and/or NAD phosphate ($NADP^+$).

<u>Algae extracts alone in the absence of FBS slightly improve cell viability but not growth.</u>

We initially set out to test if algae extracts have growth effects on cells cultured in the

absence of any FBS, using MTS assays as an initial readout and proxy for potential cell growth.

Fig. 2 shows the relative viability of EBTr cells cultured over a 10,000-fold range of

concentrations of CVE or CGF. Media supplemented with CVE and CGF alone significantly

increased EBTr cell viability compared to serum-free controls; such effects were more

pronounced for CVE than CGF and for relatively high concentrations of CVE. However,

increases in relative cell viability due to CVE supplementation alone did not exceed beyond

roughly 15% in any condition; moreover, and importantly, upon examining CVE-treated cells

under the microscope immediately prior to performing MTS assays, we did not observe any

convincing level of growth or cell doubling(s).

Given the partial but limited growth shown in Fig. 2a, we wondered if there may be

components in the CVE which were acting to inhibit cell growth. We decided to separate the

different components of CVE through filtration, centrifugation, and treatment with nuclease,

using MTS assays to compare head-to-head the extract in its crude, filtered, supernatant, and

pellet forms, each with and without nuclease treatment. Fig. S1a shows the relative viability of

EBTr cells cultured with these different CVE fractions. No obvious differences were seen

between crude, filtered and supernatant forms of extract, with viabilities of such fractions all

falling near the range of 15-20%. The viability of cells treated with the resuspended pellet was

notably lower than that of all other fractions, falling to nearly 0%, similar to that of no serum

controls. Further, CVE fractions treated with nuclease showed mostly reduced viability

compared to the corresponding fractions not treated with nuclease.

Fig. S1 (panels b and c) show the LC-MS characterization of the supernatant and pellet

forms of CVE used in the experiment described in Fig. S1a. UV absorbance data shows sharp

peaks around 20 minutes in the 200-250 nm range for both the pellet and the supernatant. The

supernatant also shows a pair of broader peaks at around 200 and 260 nm that appear to have

eluted within the first minute, neither of which was present in the pellet form. TICs of mass

spectrometry data for m/z values between 400 and 1200 for these two forms of extract show

similar footprints. Both show similarly the highest intensity spectra between 15 and 20

minutes, with relatively smaller spectra between 0 and 15 minutes. The spectra of the pellet

form show slightly greater relative signals in the 0-6-minute range. Notably, the large UV peaks

in both sample types correspond to increased signals in their respective mass spectra at around

20-21 minutes, however the large UV signal eluting early and only in the supernatant did not

correspond to any appreciably large signal(s) in its mass spectra.

Select serum-free media supplemented with *C. vulgaris* extract enhances cell viability.

After seeing limited results from algae extract alone, we decided to move our research

in a different direction, focusing on growth factors, which have previously been used to

successfully culture stem cells in an "Essential 8" (E8^{TM}) serum-free medium (Chen et al. 2011).

Fig. 2b shows percent viability of EBTr cells treated with different concentrations of basic

fibroblast growth factor (FGF) and transforming growth factor β1 (TGF). Neither growth factor

alone improved viability beyond 5-10% relative to serum-free controls, yet viability increased to

around 15% for EBTr cells treated with a combination of 10 ng/mL FGF and 10 ng/mL TGF.

Thus, we decided to use a combination of FGF and TGF for our next experiment in which we

tested these growth factors together with bovine insulin, vitamin C (in the form of L-ascorbate),

and selenium (in the form of selenium dioxide), all of which are listed as ingredients of the E8^{TM}

media. Fig. 2c shows insulin as a critical factor in real cell growth, in conjunction with growth

factors; treatments with both insulin and growth factors improved EBTr cell viability to around

35% relative to controls cultured with FBS. Despite their inclusion in E8^{TM} media, adding L-

ascorbate, selenium, or both, slightly reduced EBTr cell viability compared to that of cells

cultured with growth factors and insulin alone. We decided next to bring algae extracts back

into our experiments, combining them with insulin and growth factors and/or L-ascorbate and

selenium. Fig. 2d shows that EBTr cell viability reached approximately 75% (relative to FBS

controls) when treated with a combination of CVE, growth factors and insulin. On the other

hand, the viability of cells treated with CGF, growth factors and insulin, was actually lower than

that of cells treated with growth factors and insulin alone. Once again, the addition of L-

ascorbate and selenium had little if any beneficial effect on cell viability.

<u>Select serum-free media supplemented with *C. vulgaris* extract promotes cell growth and</u>

<u>expansion.</u>

After seeing promising viability results (i.e., Fig. 2d), we decided the next step was to

test if EBTr cells would expand when treated with insulin, growth factors and CVE in a cell count

experiment. Fig. 3a shows cell counts of CVE and CGF with supplements (i.e., growth factors,

insulin, L-ascorbate, and selenium) compared to controls with and without serum (10% FBS).

The number of EBTr cells increased in cultures with CVE or CGF plus supplements by day 2,

showing definitive growth, yet still not as much growth as control cells cultured with 10% FBS.

By day 4, the number of cells cultured with CGF plus supplements fell to a level near that of no

serum controls, but cells cultured with CVE plus supplements continued growth to around half

that of controls cultured with 10% FBS. On day 8, the number of cells cultured with CVE plus

supplements was comparable to that of controls cultured with 10% FBS, the latter of which had

decreased since day 4. Fig. 3b shows the corresponding cell death percentages for the

experiment described in Fig. 3a. Cell death percentages were largely in the 10-15% range,

slightly increasing over time, with the most death occurring for cells cultured with CGF plus

supplements on day 4 jumping to about 40%, corroborating the loss of growth shown for such

cultures between day 2 and day 4 shown in Fig. 3a. Fig. 3c shows images captured of the

experiment described in Fig. 3a and 3b. EBTr cells cultured with 10% FBS show sustained

growth through day 4 until reaching confluency, at which point growth plateaued between days

4 and 8. Cells cultured with CVE plus supplements also appeared to grow through day 4,

reaching a relatively high level of confluency; on day 8 the level of confluency appears about

the same as day 4 (perhaps slightly more) with more dead cells in the image. Cells cultured

with CGF plus supplements appear to show growth until day 2, though notably less than cells

cultured with CVE plus supplements; however, by day 4 most if not all remaining cells appear

dead, and by day 8, only dead cells remain in the image. (Note that EBTr cells in wells treated

with algae extract appear larger than cells in wells where algae extract was not present.) The

no serum control shows some cells with normal morphology on day 2, with many dead cells in

the image as well; by days 4 and 8, hardly any cells remain besides a few dead cells.

Given that EBTr cells showed growth and markedly improved viability when cultured

with CVE and select supplements, we decided next to attempt to expand EBTr cells from a 1 mL

culture (12-well plate) to a 2 mL culture (6-well plate) under these conditions (Fig. S2). On day

1, cells cultured with 10% FBS or with CVE plus supplements attached effectively, while the no

serum control already shows many dead cells. By day 4, the no serum control well looks nearly

identical to that on day 1, with some cells still attached but many dead with no visible growth.

By day 4, cells cultured with 10% FBS or CVE plus supplements show obvious growth, nearing

confluency. As observed in Fig. 3c, cells cultured with CVE plus supplements appear slightly

bigger than cells not cultured with CVE, meaning that fewer cells were necessary to reach

similar levels of confluency. After the day 4 pictures were taken, cells were detached and

removed from the 1 mL cultures, then pelleted and resuspended into fresh 2 mL cultures. The

following day, after cells attached to the plate (day 5), the relative amount of cells in the 10%

FBS control well was notably greater than that in the well with CVE plus supplements. By day 8,

cells cultured with 10% FBS reached near-confluency, while cells cultured with CVE plus

supplements remained at nearly 50% confluency. On both days 5 and 8, the no serum control

wells were essentially devoid of live cells, with only a few dead cells remaining.

Discussion

Regarding our characterization of algae extracts compared to FBS, the most striking and

obvious feature of the CVE and CGF samples that was not present in FBS samples were large

absorbance peaks around 260 nm, which eluted off the column with the aqueous mobile phase

within the first couple minutes. Such an absorbance profile is a characteristic property of

nucleic acid (Setlow and Boyce 1960), and not surprising as a relatively abundant component

present in crude cellular extracts. Moreover, since no appreciable mass signal(s) at this

retention period was/were observed, it supports the notion that the species producing this

absorbance spectra are likely nucleic acid species (e.g., DNA/RNA fragments containing at least

4 bases) with m/z values greater than that which could be detected (i.e., >1200). However,

because the sensing of foreign nucleic acid often triggers cell death responses (Schlee and

Hartmann 2016), we did not expect that foreign nucleic acid would be beneficial to cell

growth/viability, rather on the contrary. Yet our results suggest that this putative nucleic acid

at least did not harm cells and may have even been beneficial because nuclease-treated

extracts, as well as the insoluble pellet with presumably very little nucleic acid (as evidenced by

loss of the characteristic absorbance peak at 260 nm in Fig. S2a), both decreased cell viability as

compared to cells given extract that was not pre-treated with nuclease or cells treated with

soluble supernatant.

Despite an increase in EBTr cell viability observed for cells cultured with different

concentrations of CVE and GCF alone, when such plates were observed under the microscope,

no convincing amount of cell division was visible. When algae extract (CVE) was combined with

growth factors, insulin and other additives, a similar marginal increase in cell viability of around

20% was observed in an environment where cells were already dividing. These observations of

an increase in viability caused by CVE in both diving and non-dividing environments are

consistent with other studies (Okamoto et al. 2022), suggesting that CVE increases cell viability

independent of effects on proliferation. The possibility that algae extract enhances the cellular

metabolic potential without promoting cell division requires further investigation. Cell viability

assays using tetrazolium salts like MTS are often used as a proxy for cell growth, yet what such

assays actually measure is the activity of mitochondrial NADH and NADPH-dependent

dehydrogenase enzymes which reduce the reagent to a formazan product (Dunigan et al. 1995).

In most cases, the more cells there are the more mitochondria are available to reduce the MTS

reagent, but it is possible that the algae extracts have a positive effect on such metabolic

activity without promoting cell growth. Given that a common and relatively abundant signal

feature found in the mass spectra of algae extract samples mapped to $NAD^+/NADP^+$ as the

parent species, it is possible that enriched levels of NAD(H)/NADP(H) present in the algae

extracts may have caused enhanced activity of the dehydrogenase enzymes in the cells thus

explaining the observed increase in viability. Further, it was previously shown that CVE also

contains high levels of glutamate (Okamoto et al. 2020), as *Chlorella* algae use NADPH-

dependent glutamate dehydrogenase enzymes to control their cell cycle (Talley et al. 1972).

Glutamate has been shown to increase oxidation of NADH (Kannurpatti and Joshi 1999;

Maalouf et al. 2007), and increase cancer cell viability and growth (Reiman et al. 1981; Takano

et al. 2001; Gelb et al. 2015; Yamaguchi et al. 2020). Moreover, in fibroblasts, glutamate

reduced toxicity induced by high levels of cysteine (Bannai and Ishii 1982), and glutamate

dehydrogenase deficiency reduced cell viability (Tatsumi et al. 1989). Therefore, it is possible

that enriched levels of glutamate in algae extracts may also be responsible for the increased

viability observed.

Our experiments with components of the E8^{TM} media clearly showed that insulin, basic-

FGF and TGF-β are necessary for cell growth, and that these components are roughly additive in

terms of their benefits to cell viability. Early studies examining serum-free culture conditions

showed FGF and insulin as key determinants for growth (Mcfarland et al. 1991). Insulin causes

cells to import glucose, providing energy and nucleotide building blocks needed for replication.

In viability assays, addition of CVE also proved to be roughly additive, with CVE in addition to

basic-FGF, TGF-β, insulin, and other additives combined yielding the highest viability reading of

any serum-free condition tested. We demonstrated that EBTr cells can grow and expand into a

new culture vessel when cultured in serum-free media that included at minimum basic-FGF,

TGF-β, insulin and algae extract in the form of CVE, although such growth was not as robust as

that of cells cultured with 10% FBS. Recent studies have shown that mitochondrial NADP(H)

levels are essential for cell growth (Tran et al. 2021), without which cells cannot make proline

required to support nucleotide and protein synthesis. Thus, it is plausible that enriched levels

of NAD(H)/NADP(H) in CVE extract may provide a true benefit to cell growth; however, whether

the additive increase in viability from CVE translates to any real benefit in growth rate beyond

that from basic-FGF, TGF-β and insulin alone remains unclear. Regardless of whether algae

extract is truly beneficial to cell growth, it is important to note that bovine fibroblasts could be

cultured with only a subset of the E8^{TM} components present. This may at least suggest some

potential economic benefit due to lower cost serum-free formulations requiring fewer

components. Since growth factors and insulin directly interact with their respective receptors

through a relatively small range of residues across the entire protein, further experiments could

test whether smaller peptide fragments mapping to such regions within these proteins remain

actionable enough to sustain cell growth. Such substitutes may simplify and further lower the

cost of serum-free formulations in which cells can be effectively cultured. Similarly, small

molecules that act as agonists on basic-FGF and/or TGF-β receptors could also be tested for the

same purpose.

Some limitations of our studies include aspects of the growth experiments in which we

did not anticipate how differences in cell size could impact the results. In these experiments, it

appears from the images as though cells cultured with algae extract were larger in size as

compared to those not cultured with algae extract. Notably, increases in cell viability correlate

with increased cell size (Dreesen and Fussenegger 2011), which could at least partially explain

this observation since we also found that algae extract typically increased cell viability.

However, differences in cell size observed with and without algae extract in the culture make

the growth and expansion experiments difficult to interpret. For instance, during the expansion

experiment, it appears that more cells were transferred from the 1 mL culture with 10% FBS

compared to the culture with CVE plus supplements despite these two groups appearing to

have a similar level of confluency on day 4. While both conditions promoted growth, the cells

cultured in 10% FBS not only grow faster but require less space per cell. As a result, in the 2 mL

cultures, the cells cultured in 10% FBS had a higher base count from which to start growing and

were thus much more confluent by day 8. Despite the observed increase in viability, it is also a

possibility that this increase in cell size corresponds to replicative senescence. Cells increase in

size when they enter a permanent cell-cycle arrest and senescence (Biran et al. 2017), so

further experiments examining the extent to which proliferation is affected by CVE are

necessary. In a separate case, it should be noted that the day 8 FBS control result in our cell

count experiment (Fig. 3c) shows a sharp decrease in cell number, which can be explained by

over-confluence of the culture dish, resulting in cell death. As a result, the day 4 image and cell

count more accurately display the efficacy of CVE plus supplements, since it would be expected

that in a larger dish, the FBS control condition would continue to promote cell growth through

day 8. Furthermore, we must acknowledge limitations in counting the fraction of dead cells

recorded in our cell count experiments. As noted in the methods section, old media was first

removed and the cells were briefly washed, which likely removed a large amount of previously

dead and floating cells. Thus, the dead cell count more accurately represents the percentage of

cells which were nearly dead, or dead, but not yet detached and floating. Finally, it is important

to note that algae extracts may be subject to batch-to-batch variation depending on the growth

and extraction conditions used to generate them, thus further study of such variation in *C.*

vulgaris extract is important to confirm the reproducibility of results.

Taken together, our studies show that algae extract derived from *Chlorella vulgaris*

enhances viability of bovine fibroblasts, possibly due at least in part to its enriched levels of

NAD(H)/NADP(H) and/or glutamate, and that CVE can be used together with other supplements

(basic-FGF, TGF-β, and insulin) to grow cells in the absence of serum. Further studies are

needed to determine the extent to which CVE promotes cell growth and whether the

supplementary components can be further modified to reduce their cost.

References

Bannai S, Ishii T (1982) Transport of Cystine and Cysteine and Cell Growth in Cultured Human

Diploid Fibroblasts: Effect of Glutamate and Homocysteate. Journal of Cellular Physiology

112:265–272

Biran A, Zada L, Abou Karam P, et al (2017) Quantitative identification of senescent cells in

aging and disease. Aging Cell 16:661–671. https://doi.org/10.1111/acel.12592

Burt SM, Carter TJN, Kricka LJ (1979) Thermal characteristics of microtitre plates used in

immunological assays. Journal of Immunological Methods 31:3-4.

https://doi.org/10.1016/0022-1759(79)90135-2

Chen G, Gulbranson DR, Hou Z, et al (2011) Chemically defined conditions for human iPSC

derivation and culture. Nature Methods 8:424–429. https://doi.org/10.1038/nmeth.1593

Dreesen I, Fussenegger M (2011) Ectopic expression of human mTOR increases viability,

robustness, cell size, proliferation, and antibody production of chinese hamster ovary cells.

Biotechnology and Bioengineering 108:853–866. https://doi.org/10.1002/bit.22990

Dunigan D, Waters S, Owen T (1995) Aqueous soluble tetrazolium/formazan MTS as an

indicator of NADH- and NADPH-dependent dehydrogenase activity. Biotechniques 19:640–

649

Gelb T, Pshenichkin S, Rodriguez OC, et al (2015) Metabotropic glutamate receptor 1 acts as a

dependence receptor creating a requirement for glutamate to sustain the viability and

growth of human melanomas. Oncogene 34:2711–2720.

https://doi.org/10.1038/onc.2014.231

Gerber PJ, Steinfield H, Henderson B, et al (2013) Tackling climate change through livestock - a

global assessment of emissions and mitigation opportunities. Food and Agriculture

Organization of the United Nations

Humbird D (2021) Scale-up economics for cultured meat. Biotechnology and Bioengineering

118:3239–3250. https://doi.org/10.1002/bit.27848

Kannurpatti SS, Joshi NB (1999) Energy metabolism and NAD-NADH redox state in brain slices in

response to glutamate exposure and ischemia. Metabolic Brain Disease 14:33-43

Ladak A, Anthis JR (2021) Animals, Food, and Technology (AFT) Survey: 2020 Update. Sentience

Institute

Maalouf M, Sullivan PG, Davis L, et al (2007) Ketones inhibit mitochondrial production of

reactive oxygen species production following glutamate excitotoxicity by increasing NADH

oxidation. Neuroscience 145:256–264.

https://doi.org/10.1016/j.neuroscience.2006.11.065

Mcfarland DC, Ane J, Pesall E, et al (1991) Proliferation of the turkey myogenic satellite cell in a

serum-free medium. Comp Biochem Physiol 99A:163–167

Murakami H, Masui H (1980) Hormonal control of human colon carcinoma cell growth in serum-

free medium. Proc Natl Acad Sci USA 77:3464–3468

Ng JY, Chua ML, Zhang C, et al (2020) Chlorella vulgaris Extract as a Serum Replacement That

Enhances Mammalian Cell Growth and Protein Expression. Frontiers in Bioengineering and

Biotechnology 8:564667. https://doi.org/10.3389/fbioe.2020.564667

Okamoto Y, Haraguchi Y, Sawamura N, et al (2020) Mammalian cell cultivation using nutrients

extracted from microalgae. Biotechnology Progress 36:e2941.

https://doi.org/10.1002/btpr.2941

Okamoto Y, Haraguchi Y, Yoshida A, et al (2022) Proliferation and differentiation of primary

bovine myoblasts using Chlorella vulgaris extract for sustainable production of cultured

meat. Biotechnology Progress. https://doi.org/10.1002/btpr.3239

Penttinen K, Saxén E (1959) Growth-Controlling Action of Human Serum in Cell Culture. Nature

184:1570–1571

Reiman RE, Huvos AG, Benua RS, et al (1981) Quotient Imaging with N-13 L-Glutamate in

Osteogenic Sarcoma: Correlation with Tumor Viability. Cancer 48:1976–1981

Schlee M, Hartmann G (2016) Discriminating self from non-self in nucleic acid sensing. Nature

Reviews Immunology 16:566–580

Setlow R, Boyce R (1960) The ultraviolet light inactivation of phiX174 bacteriophage at different

wave lengths and pH's. Biophysical Journal 1:29–41

Specht L (2020) An analysis of culture medium costs and production volumes for cultivated

meat. The Good Food Institute

Takano T, Lin JHC, Arcuinoo G, et al (2001) Glutamate release promotes growth of malignant

gliomas. Nature Medicine 7:1010–1015

Talley DJ, White LH, Schmidt RR (1972) Evidence for NADH-and NADPH-specific isozymes of

glutamate dehydrogenase and the continuous inducibility of the NADPH-specific isozyme

throughout the cell cycle of the Eucaryote Chlorella*. Journal of Biological Chemistry

247:7927–7935

Tatsumi C, Yorifuji S, Takahashi M, Tarui S (1989) Decreased viability of skin fibroblasts from

patients with glutamate dehydrogenase deficiency. Neurology 39:451-452.

https://doi.org/10.1212/WNL.39.3.451

Tran DH, Kesavan R, Rion H, et al (2021) Mitochondrial NADP+ is essential for proline

biosynthesis during cell growth. Nature Metabolism 3:571–585.

https://doi.org/10.1038/s42255-021-00374-y

Tuomisto HL, Teixeira De Mattos MJ (2011) Environmental impacts of cultured meat

production. Environmental Science and Technology 45:6117–6123.

https://doi.org/10.1021/es200130u

Yamaguchi I, Yoshimura SH, Katoh H (2020) High cell density increases glioblastoma cell viability

under glucose deprivation via degradation of the cystine/glutamate transporter xCT

(SLC7A11). Journal of Biological Chemistry 295:6936–6945.

https://doi.org/10.1074/jbc.RA119.012213

Figure Legends

Fig. 1

Comparative characterization of *Chlorella* extracts versus FBS by LC-MS. (**a**) Perspective 3D

surface plots are shown comparing UV chromatograms of FBS, CVE and CGF samples

(wavelengths 200-400 nm); each plot shows the mean normalized absorbance spectra of three

replicate injections of the same sample type. Arrows indicate large, broad peaks around 260

nm present in CVE and CGF samples. (**b**) Normalized total ion chromatograms (TICs) are shown

as the sum of signals of all m/z values from 400-1200 for FBS, CVE and CGF samples;

representative data shown are averaged spectra from three injections of the same sample type.

(**c**) Scatter plot shows results of principal component analysis (PCA) on panel b MS TIC data;

sample types are indicated by different colors with PC2 (circles) and PC4 (diamonds) values

plotted on the same y-axis scale. (**d**) Perspective 3D surface plots show normalized mass

spectra of FBS, CVE and CGF samples; data shown are averages of three replicate injections.

Dashed box indicates region of interest where signal location/intensity appears similar across

all three sample types (i.e., retention times from 15-21 minutes and m/z values from 800-1200).

(**e**) Aligned mass spectra show the sum of all signals for retention times from 15-21 minutes at

each whole integer m/z value from 800-1200; data are shown as mean normalized spectra of

three replicate injections of the same sample. Red vertical lines indicate signals of relatively

high abundance shared across all three sample types; table lists potential parent species for

corresponding signals based on top candidates identified by database (MassBank.eu) search

Fig. 2

Select growth factor and insulin supplemented *C. vulgaris* extract improves cell viability in the

absence of FBS. (**a**) Bar plots show viability of EBTr cells cultured with CVE and CGF relative to

controls cultured in 10% FBS at day 3. Panel (**b**) shows relative viability of EBTr cells cultured in

serum-free media supplemented with 10 or 100 ng/mL fibroblast growth factor 2 (FGF), or 1 or

10 ng/mL transforming growth factor β1 (TGF), or combinations thereof, relative to controls

cultured in 10% FBS after 3 days. Panel (**c**) shows relative viability of EBTr cells cultured in

serum-free media supplemented with growth factors (GFs), or 3 μM bovine insulin, or 0.2 mM

L-ascorbate (LA), or 80 nM selenium dioxide (Se), or combinations thereof, relative to controls

cultured in 10% FBS. Growth factors were used at 30 ng/mL (FGF) and 10 ng/mL (TGF). Panel

(**d**) shows relative viability of EBTr cells cultured in serum-free media supplemented with 10 g/L

CVE or 10% CGF alone, or GFs and insulin with and without LA and Se, or combinations thereof,

compared to controls cultured in 10% FBS. Concentrations of GFs, insulin, LA and Se were the

same as that for panel c. Statistical significance was determined by *t*-test with Welch's

correction; significance levels are indicated as *$p < 0.05$, ***$p < 0.001$, ****$p < 0.0001$, ns=not

significant, n=6

Fig. 3

Growth factor and insulin supplemented *C. vulgaris* extract stimulates cell growth in the

absence of FBS. Line plots show live (**a**) and percent dead/non-intact (**b**) EBTr cells after 0, 2, 4

and 8 days in culture. Different media conditions are indicated by different colors; all counts

were performed in duplicate. (**c**) Brightfield images show representative EBTr cell density and

morphology for each condition and time point; white scale bar indicates length of ~100 μm.

CVE was used at 10 g/L and CGF at 10% by volume. Supplements include 10 ng/mL TGF, 30

ng/mL FGF, 3 μM bovine insulin, 0.2 mM L-ascorbate, and 80 nM selenium

Figure 1

Peak	*m/z*	Mean shared signal	Top MassBank candidate
1	934	0.22	Epigallocatechin gallate (EPCG)
2	1021	0.52	DSGEGDFXAEGGGVR (Fibrinopeptide A derivative)
3	1076	0.90	NAD$^+$/NADP$^+$
4	1151	0.68	Ecdysterone 22-O-benzoate
5	1173	0.50	ADpSGEGDFXAEGGGVR (Fibrinopeptide A derivative)

Figure 2

a

b

c

	10% FBS	CVE + supplements	CGF + supplements	No serum
Day 2				
Day 4				
Day 8				

~100 μm