Human Cytomegalovirus modifies placental small extracellular vesicle secretion and composition towards a proviral phenotype to enhance infection of fetal recipient cells

Mathilde Bergamelli [1], Hélène Martin [1], Yann Aubert [1], Jean-Michel Mansuy [2], Marlène Marcellin[3], Odile Burlet-Schiltz [3], Ilse Hurbain [4,5], Graça Raposo [4,5], Jacques Izopet [1,2], Thierry Fournier [6], Alexandra Benchoua [7], Mélinda Bénard [1,8], Marion Groussolles [1,9,10], Géraldine Cartron [11], Yann Tanguy le Gac [11], Nathalie Moinard [12,13], Gisela D'Angelo [4] and Cécile E. Malnou [1]*

[1] Institut Toulousain des Maladies Infectieuses et Inflammatoires (Infinity), Université de Toulouse, INSERM, CNRS, UPS, Toulouse, France.
[2] CHU Toulouse, Hôpital Purpan, Laboratoire de Virologie, Toulouse, France.
[3] Institut de Pharmacologie et de Biologie Structurale, IPBS, Université de Toulouse, CNRS, UPS, Toulouse, France. Infrastructure nationale de protéomique, ProFI, FR 2048, Toulouse, France.
[4] Institut Curie, CNRS UMR144, Structure et Compartiments Membranaires, Université Paris Sciences et Lettres, Paris, France.
[5] Institut Curie, CNRS UMR144, Plateforme d'imagerie cellulaire et tissulaire (PICT-IBiSA), Université Paris Sciences et Lettres, Paris, France.
[6] Université de Paris, Inserm, UMR-S1139, Physiopathologie & pharmacotoxicologie du placenta humain, microbiote pré & postnatal (3PHM), F-75006 Paris, France.
[7] CECS, I-STEM, AFM, Evry, France.
[8] CHU Toulouse, Hôpital des Enfants, Service de Néonatalogie, Toulouse, France.
[9] CHU Toulouse, Hôpital Paule de Viguier, Service de Diagnostic Prénatal, Toulouse, France.
[10] Equipe SPHERE Epidémiologie et Analyses en Santé Publique: Risques, Maladies chroniques et handicaps, Université de Toulouse, INSERM UMR1027, UPS, Toulouse, France.
[11] CHU Toulouse, Hôpital Paule de Viguier, Service de Gynécologie Obstétrique, Toulouse, France.
[12] Développement Embryonnaire, Fertilité, Environnement (DEFE), INSERM UMR 1203, Université de Toulouse et Université de Montpellier, France.
[13] CECOS, Service médecine de la reproduction, CHU Toulouse, Hôpital Paule de Viguier, Toulouse, France.
* corresponding author: Cécile E. Malnou
Email: cecile.malnou@univ-tlse3.fr

Author Contributions:
MBer and HM designed and performed most experiments and analyzed data. JMM and JI were in charge of the viral titrations. MM, OBS and YA performed the proteomic experiments and data analysis. MBén, MG, GC and YTLG collected the clinical samples and data. NM from Germethèque was in charge of the ethical issues

and received women authorizations. IH, GDA and GR supervised the electron
microscopy experiments and data analysis. TF and AB generated the HIPECs and
NSCs respectively, and gave scientific expertise concerning their culture. CEM got
grants for this study, conceived and designed the study, analyzed data and wrote the
paper, with the help of MBer, YA and OBS. All authors provided critical feedback
and helped shape the research, analysis and manuscript.
50

51 **Competing Interest Statement:**
52 The authors declare that the research was conducted in the absence of any commercial
53 or financial relationships that could be construed as a potential conflict of interest.
54
Keywords: hCMV, congenital infection, extracellular vesicles, placenta,
cytotrophoblast

Abstract

Although placental small extracellular vesicles (sEVs) are extensively studied in the
context of pregnancy, little is known about their role during human cytomegalovirus
(hCMV) congenital infection, especially at the beginning of pregnancy. In this study,
we examined the consequences of hCMV infection on sEVs production, composition
and function using an immortalized human cytotrophoblast cell line derived from first
trimester placenta. By combining complementary approaches of biochemistry,
electron microscopy and quantitative proteomic analysis, we showed that hCMV
infection increases the yield of sEVs produced by cytotrophoblasts and modifies their
protein content towards a proviral phenotype. We further demonstrate that sEVs
secreted by hCMV-infected cytotrophoblasts potentiate infection in naive recipient
cells of fetal origin, including human neural stem cells. Importantly, these functional
consequences are also observed with sEVs prepared from either an *ex vivo* model of
infected histocultures from early placenta or from the amniotic fluid of patients
naturally infected by hCMV at the beginning of pregnancy. Based on these findings,
we propose that placental sEVs could be key actors favoring viral dissemination to the
fetal brain during hCMV congenital infection.

Significance Statement
Human cytomegalovirus (hCMV) infection is a major issue during pregnancy,
affecting 1% of births in western countries. Despite extensive research, the
pathophysiology of this congenital infection remains unclear. Recently, increasing
evidence point to the key role of placental small extracellular vesicles (sEVs) in
materno-fetal communication during pregnancy. Here, we examined the impact of
hCMV infection on the protein composition and function of placental sEVs. We
observe that hCMV infection leads to major changes in placental sEV protein content.
Functional studies show the ability of sEVs produced by placental infected cells to
facilitate further infection of naive recipient fetal cells, notably human neural stem
cells. Our study demonstrates that placental sEVs are key players of hCMV
pathophysiology during congenital infection.

86
87 **Main Text**

88
89 **Introduction**

90
Human cytomegalovirus (hCMV) belongs to the *Herpesviridae* family and its
prevalence is of 50 to 90 % in global human population. Most of the hCMV infections
occurring among immunocompetent adults induce an asymptomatic acute replication
phase followed by a lifelong persistent latent state. However, hCMV primo-infection,
reinfection and/or reactivation may severely compromise the health of
immunocompromised people, and is a major issue during pregnancy (1, 2).
Congenital infection by hCMV affects 1% of live births in western countries, making
hCMV the most frequently transmitted virus *in utero* (2, 3), causing placental and
fetal impairments of variable severity. The most severe consequences are observed
when transmission occurs during peri-conceptional period or first trimester (4).
Infection of the placenta itself allows the virus to actively replicate and enable its
further access to the fetus (3, 5, 6). The infected placenta can develop a pathology that
may lead to miscarriage, premature delivery, *intra* uterine growth retardation or even
fetal death (2, 4). On the other side, the infection of the fetus causes various visceral
and central nervous system damage, congenital hCMV infection being the most
common cause of brain malformations and deafness of infectious origin (2, 7, 8).
Despite the extensive research conducted so far, the pathophysiology of hCMV
infection remains unclear, especially concerning potential factors which may explain
the wide variety of clinical manifestations and their severity (2).
In the course of hCMV congenital infection, the placenta is a central key
organ, i.e. the target of viral replication allowing further vertical transmission towards
the fetus. Amongst the numerous placental functions, a recently described and
extensively studied mode of communication between both maternal and fetal sides
consists in the production of placental extracellular vesicles (EVs) (9, 10). EVs are
membranous vesicles secreted by cells in both physiological and pathological
situations, which main subtypes can be distinguished depending on their biogenesis
and size into small EVs (sEVs) and large EVs (lEVs). They are specifically composed
of various molecules such as proteins, lipids and coding and non-coding RNAs (11,
12). Once released into the extracellular space, EVs can be internalized by other cells,
in their immediate environment or within long distances, wherein they exert
regulatory roles (13). For example, placental EVs can be uptaken by Natural Killer
cells (14, 15) or by primary placental fibroblasts (16). Although the understanding of
the biological relevance of placental EVs *in vivo* remains limited, recent findings
highlight their roles in cell-cell communication underlying the feto-placenta-maternal
dialogue during pregnancy (17, 18). Interestingly, placental EV content is altered
upon gestational diseases such as preeclampsia, preterm birth or gestational diabetes
mellitus, and recent literature points towards a putative role of dysregulated placental
EVs during pathological pregnancies (19-24).

Although placental sEVs are extensively studied in the context of pregnancy
diseases, little is currently known about their role during hCMV congenital infection,
especially at the very beginning of pregnancy where the most severe sequelae take
their origin. Our recent data described a dysregulation of the surface expression of
placental sEV markers upon hCMV infection in an *ex vivo* model of first trimester
placental histocultures, suggesting a putative role for viral dissemination (25). In the
present study, we used immortalized cytotrophoblasts derived from first trimester
placenta, named HIPEC (26), to comprehensively examine the consequences of
hCMV infection on sEV production, content and function in recipient cells. We show
that hCMV increases the sEV production by HIPECs and alters their protein content
towards a proviral phenotype. Finally, we observe that sEVs secreted by hCMV-
infected HIPECs potentiate infection in naive recipient cells, including human neural
stem cells (NSCs). Importantly, this enhancement of hCMV infection is also observed
with sEVs from *ex vivo* early placental histocultures and with sEVs purified from
amniotic fluid of patient infected by hCMV during first trimester of pregnancy. We
propose that placental sEVs could be key players favoring viral dissemination towards
fetal brain during hCMV congenital infection.
146
147 **Results**
148
HIPEC infection by hCMV leads to an increase of sEV secretion without
modifying their general features
To study the consequences of hCMV infection on placental sEV secretion,
composition and function in early pregnancy, HIPEC conditioned media were
collected between 48-72 h post-infection before sEV preparation, times at which 50-
80 % of cells were infected (Supplementary Figure S1B). To exclude any viral
contamination of sEVs, an infectivity test was systematically carried out for each
preparation. No infection was detected in cells incubated with sEVs alone
(Supplementary Figure S1C). Moreover, no structure evoking viral particles was
observed by transmission electron microscopy (TEM) on sEVs from infected HIPECs
(Supplementary Figure S2, and data not shown).
Despite a decrease of cell number upon hCMV infection compared to non-
infected (NI) HIPECs (Supplementary Figure S1D), the quantification of sEVs
isolated per cell showed a significant higher yield upon infection, with an increase of
around 40 % (Figure 1A). No difference in either the mean size or the mode size of
the sEVs was observed upon infection when analyzed by nanoparticle tracking
analysis (NTA; Figure 1B) and sEVs from both NI or infected HIPECs exhibited the
same typical structure and shape as evidenced by TEM (Figure 1C and
Supplementary Figure S2).
We next assessed the impact of hCMV infection on the expression of sEV
canonical markers. By combining western-blotting, immunolabeling electron
microscopy (IEM) and multiplex bead-based flow cytometry, we observed that sEV
preparations expressed specific vesicular markers including CD9, CD81, Alix and
Tsg101, but no endoplasmic reticulum or mitochondrial markers (Figure 1D, E and

supplementary Figure S3), attesting the purity of the sEV preparations (27). No
drastic differences were observed in their expression levels upon infection as
compared to NI cells, except for CD63 protein which was not detected in whole cell
lysates, but enriched in sEVs from hCMV-infected HIPECs (Figure 1E). By IEM, at
the level of individual vesicles, we noticed that this increase was correlated to the
presence of a small proportion of sEVs highly positive for CD63 (between 1-5 %),
while the others remained negative (Figure 1D and supplementary Figure S4). By
multiplex bead-based flow cytometry assay, no significant increase in CD63
expression was detected between sEVs from NI or infected HIPECs upon infection,
certainly due to the low proportion of positive vesicles (supplementary Figure S3).
Hence, hCMV infection increased HIPEC-sEV production but did not globally impact
on their global features, except for CD63 that is detected in a subpopulation of
vesicles upon infection.
186
sEVs secreted by infected HIPECs harbor a proviral protein cargo
To extend the molecular characterization of sEVs secreted by hCMV-infected
HIPECs, a comprehensive proteomic approach that allows for a deeper analysis of
sEV composition was carried out. The analysis by mass spectrometry-based
quantitative proteomics of equivalent amounts of sEVs from NI or hCMV-infected
HIPECs led to the identification of 3079 proteins across all samples (3048 human and
31 viral; Supplementary Table S1). Among the 3048 human proteins identified, the
gProfiler2 R package was able to interrogate 2936 proteins, for which the term
"extracellular exosome" (Gene ontology GO:0070062) appeared as the most
significantly enriched (false discovery rate (FDR) $= 1.087859e^{-259}$, R package
gProfiler2), with 962 of them (32.8 %) associated with the "extracellular exosome"
GO term. Conversely, these 962 proteins constituted 44.1 % of the proteins which
define the GO term, and the 94 of the top 100 most frequently identified exosomal
proteins, as defined by the Exocarta database
(http://exocarta.org/exosome_markers_new), were detected in the sEV preparations.
Proteomics data indicated that 37 proteins, including 31 viral proteins, were
significantly over-represented and 15 human proteins under-represented in sEVs from
hCMV-infected HIPECs (Figure 2A). Interestingly, the Thy-1 cellular protein, known
to play an important role to facilitate hCMV entry into cells *via* macropinocytosis (28,
29), was significantly enriched in sEVs upon infection (Supplementary Figure S5A).
Moreover, by using Ingenuity Pathway Analysis (IPA) tool, several biological
functions were found to be significantly over- or under-represented in sEVs upon
infection (Figure 2B). The most modulated pathway identified was "autophagy", with
several actors showing altered expression in sEVs from infected HIPECs, leading to
an overall pattern of autophagy activation (Figure 2C). Since the EV content primarily
reflects the composition of the cells they are derived from, this may also reflect the
activation of the autophagy pathway induced during early hCMV infection of host
cells (30-32). On the other hand, two pathways related to mitochondrial functions
(*i.e.,* "consumption of oxygen" and "ATP synthesis ") were found to be lower in sEVs
from hCMV-infected HIPECs than in NI HIPECs (Figure 2B). This may also be the

consequence of hCMV-induced mitochondrial dysfunctions (33-36), likely *via*
interference with the antiviral Viperin protein, which leads to decreased cellular ATP
levels (37).
Importantly, the proteomic data also revealed that 31 proteins were of viral
origin (Figure 2D and Supplementary Table S2). They are involved in different
aspects of hCMV infection, from viral entry to egress, quiescence, as well as
pathogenicity and immune evasion, and are mainly immediate early or late proteins
(Supplementary Figure S5B). Interestingly, although sEV preparations were devoid of
viral particles as assessed by infectivity assays and TEM (supplementary Figures S1C
and S2), some of the viral proteins in sEV from hCMV-infected HIPECs are
structural proteins (Figure 2D and Supplementary Table S2), i.e. the envelope
proteins gB, gH and gM, as it has already been described in other studies (38-40).
Most of the other proteins identified were capsid and tegument proteins that are
delivered to host cells upon infection, or proteins immediately expressed after virus
entry such as IE1 and IE2, which play a role in early transcriptions (41-43). We also
identified pp65, which participates to the transactivation of viral major immediate
early genes (44, 45), as well as pp71, which stimulates viral immediate early
transcription and inhibits the host innate response by targeting STING (46, 47).
Finally, IRS1 and TRS1, which inhibit the establishment of an antiviral state in
infected cells, in particular by antagonizing the autophagy pathway induced upon
hCMV infection (30, 31), were also detected. Altogether, analysis of the proteomic
data indicates that sEVs secreted from infected HIPECs carry a protein cargo with
proviral properties. The incorporation of these viral proteins into sEVs could enhance
the viral spread by providing the recipient cells with elements that may facilitate the
early steps of a further hCMV infection.
242
sEVs are efficiently uptaken by recipient MCR5 cells
We next examined whether MRC5 cells could uptake sEVs. PKH67-stained
sEVs were incubated with MRC5 cells for different times and the internalization of
labeled-sEVs was evaluated both by confocal fluorescence (Figures 3A and
supplementary S6) and flow cytometry (Figure 3B, C). As early as 2 h post-
incubation, some MRC5 cells already showed numerous cytoplasmic puncta for both
sEVs produced by NI and infected HIPECs (Supplementary Figure S6). At 16 h,
many cells showed bright green puncta in their cytoplasm (Figure 3A), with a
percentage of positive cells evaluated to 13% by flow cytometry, which was not
altered up to 24h (Figure 3B-C). Moreover, these puncta were visible in the same
confocal plan as actin (revealed by phalloidin staining), corroborating the
intracytoplasmic localization of PKH67 fluorescence, and excluding therefore a
possible binding of sEVs to the cell surface (Figure 3A, Supplementary Figures S6B
and C and Supplementary Video S1). However, in comparison to the fluorescence
data (Figure 3A), this percentage may be somewhat underestimated, as subtraction of
cell autofluorescence background may mask a significant amount of uptaken sEVs,
which fluoresce poorly, given the small size of the vesicles and thus, the small
number of PKH67 molecules incorporated. Altogether, these data indicate that the

sEVs secreted by HIPECs are largely uptaken by recipient cells, and also suggest that
the internalized sEVs may thereafter exert a biological function.
263

sEVs from hCMV-infected HIPECs potentialize further infection of recipient
MRC5 cells
Based on the proteomic data suggesting a proviral activity of sEVs, we
assessed their ability to modulate hCMV infection. As sEV content was composed of
proteins prone to act on early steps of infection - entry and immediate early
transcriptions - we reasoned that any putative action of sEVs should take place
immediately after delivery of their cargo into recipient cells. In this regard, different
amounts of sEVs from NI or hCMV-infected HIPECs were incubated with MRC5
cells either concomitantly or 2 h before hCMV infection (Figure 4A). 24 h later, cells
were subjected to an anti-IE1/2 immunofluorescence (Figure 4B). When added alone,
sEVs from infected HIPECs did not lead to any detectable expression of IE1/2 in
MRC5 cells (Figure 4Ba), confirming that they were not contaminated by residual
infectious viral particle. This also indicates that the detection of IE proteins upon
infection was due to viral gene expression and not to the presence of IE proteins
carried by sEV from hCMV-infected HIPECs, even if IE proteins 1 et 2 were detected
in proteomic analysis. As shown in Figure 4C, when added simultaneously with
hCMV, sEVs did not influence the level of MRC5 cell infection, whatever their origin
and quantity, as compared to non-treated cells. In contrast, the addition of increasing
doses of sEV from infected HIPECs 2 h before infection led to a potentiation of the
infection, when compared to cells treated with sEV from NI HIPECs. This increase
was significant for the two highest doses of sEVs applied, with a stimulation of
infection of around 17 % and 30 % when 50 and 200 sEVs were added per cell,
respectively (Figure 4D).
287

sEVs from hCMV-infected HIPECs, placental tissues and amniotic fluids
enhance infection of human neural stem cells
To examine the potential proviral role of placental sEVs on hCMV
transmission towards the fetal brain, similar experiments using human neural stem
cells (NSCs) with different sources of sEVs were performed. NSCs, which are
permissive for hCMV infection, constitute a particularly relevant model for studying
the consequences of hCMV infection on fetal neural progenitors (7, 48). As observed
with MRC5 cells, sEVs from hCMV-infected HIPECs promote a significant
enhancement of hCMV infection of NCSs as compared to sEVs from NI HIPECs,
with a significant mean increase of 42 % (Figure 5A). To get closer to physiological
conditions, an *ex vivo* model of first trimester placental histocultures, that we have
previously developed (49, 50), was next infected by hCMV and used for sEV
isolation. Recent data obtained from this model showed a modification of sEV surface
markers upon hCMV infection and suggested a proviral role for placental sEVs (25).
Again, sEVs produced by infected placental histocultures significantly potentiated
hCMV infection of NSCs in comparison to sEVs secreted by NI histocultures, with a
mean of 48 % of increase (Figure 5B).

Finally, we assessed the impact of sEVs prepared from amniotic fluid (AF),
that contain an important proportion of placental sEVs (51), obtained from naturally
infected pregnant women (Figure 5C). AF was collected from 3 women, whose
clinical data are summarized in Supplementary Table S3. The first one (NP19) had no
history of infection or other maternal disease during pregnancy. The second one
(NP6) presented a first trimester hCMV seroconversion with her AF negative for
hCMV by qPCR and no fetal defects reported by imaging. The third one (NP12) also
seroconverted during first trimester of pregnancy but her AF was positive for hCMV
by qPCR and the fetus presented major brain lesions. A control experiment verifying
that no infectious particle was present in sEVs preparations was systematically
performed in parallel (data not shown). As shown in Figure 5C, level of infection of
NSCs incubated with sEVs prepared from the non-infected patient NP19 or from NP6
(infected patient but with AF negative for hCMV) remained similar. Remarkably,
when sEVs were prepared from NP12, a case of hCMV positive pregnancy with
severe impairment of fetal development, a significant increase of infection of around
20 % was observed in comparison to incubation with sEVs from NP6 or NP19,
indicating that NP12 sEVs favored the enhancement of hCMV infection in recipient
NSCs. Hence, our data establish the existence of a tight correlation between the
clinical severity of hCMV-complicated pregnancy and the proviral effect of sEVs
prepared from patient AF.
325
326   **Discussion**
327
Although the role of placental sEVs during normal and pathological
pregnancy is extensively studied (18, 20, 23, 24), the impact and consequences of
hCMV infection on placental sEVs are far to be deciphered, in particular at the
beginning of pregnancy. We recently described that infection of first trimester *ex vivo*
placental histocultures modified sEV surface markers, suggesting a potential proviral
role of the sEVs (25). However, the difficulty of purifying high sEV amounts from
this model hampered the possibility to conduct a study combining exhaustive analysis
of sEV composition and biological function. In this work, we used a combination of a
cytotrophoblast cell line deriving from first trimester placenta (26), *ex vivo* placental
histocultures and patient amniotic fluids to isolate sEVs and examine the impact of
hCMV infection on their composition and function.
Despite a large number of publications describing a role of EVs on pregnancy
regulation, the nature of examined EVs is often not precisely stated, and confusion
remains about the subtypes of EVs studied. Here, we focused our study on sEVs,
considering their importance in communication between mother and fetus and the fact
that they are often dysregulated during pregnancy pathologies (18, 21, 23). Using the
gold-standard method based on differential ultracentrifugation followed by density
gradient ultracentrifugation, sEVs devoid of viral particles were systematically
isolated in a rigorous manner (52). Their characterization by a combination of
complementary approaches revealed that they harbor exosome-like features in terms
of size, structure and presence of canonical EV markers (27), but their endosomal

origin cannot formally be stated, since we did not examine the mechanisms
underlying the biogenesis of sEVs in the present study. Instead, we demonstrated that
sEVs from hCMV infected cells, histocultures and prepared from amniotic fluids of
infected pregnant women present an altered protein content that facilitate the infection
of naive recipient fetal cells.

We found that hCMV infection of HIPECs increased the yield of sEV
production. However, this increase was counterbalanced by a reduced cell growth
upon infection. Hence, the global quantity of sEVs harvested from hCMV- or mock-
infected HIPECs remained similar between both conditions. As hCMV mainly
replicates in discrete foci in placenta *in vivo*, it is unlikely that hCMV infection would
impact on the global quantity of sEVs secreted by the placenta, but we cannot exclude
the possibility that local increase of sEV secretion induced by infection may modify
placental micro-environment.

Interestingly, most of the viral proteins identified in our proteomic study have
not been described in the other proteomic study done of sEVs secreted from hCMV-
infected MRC5 cells (39). Conversely, some viral proteins identified by Turner *et al*,
such as IR11, IRL12, US14, were not found in our study. One plausible explanation is
that, at relatively early time points within the replication cycle (48 h to 72 h upon
infection in our present study) the content of sEVs secreted - highly enriched in viral
envelop, capsid, tegument and immediate early proteins - may prime the neighboring
cells for viral dissemination and spread. In contrast, at latter time point (5 days post
infection in Turner *et al* study), the sEV protein cargo could more serve immuno-
evasion functions, by expressing for example the viral Fc-gamma receptor homologue
IR11/gp34 (39). Hence, it seems that composition of sEV secreted by hCMV-infected
cells evolves with time, depending on their state of infection and the step of hCMV
replication cycle.

In addition to the viral proteins carried by HIPEC sEVs, sEV composition for
proteins of cellular origin was also altered upon hCMV infection. By using IPA to
analyze biological pathway, the term "autophagy" was the first over-represented in
sEVs from infected HIPECs, likely reflecting the autophagy activation induced by
hCMV in host cells at the very early times of infection (30-32). Importantly, TRS1
and IRS1, described to antagonize autophagy at latter time points of the viral cycle
(30, 31), were also found in sEVs, which may consequently inhibit the induction of
autophagy in recipient cells upon sEV uptake. Thy-1, which plays an important role
for favoring hCMV entry into cells by macropinocytosis (28, 29), was also highly
over-represented in sEVs from infected cells. Hence, all the elements brought by the
proteomic analysis of sEVs from infected HIPECs indicate that they were prone to
facilitate viral infection of recipient cells.

By examining the function of sEVs from NI or hCMV-infected HIPECs, we
observed that sEVs hCMV enhanced further infection of MRC5 cells. Such proviral
properties of sEVs produced by infected cells have already been described for other
Herpesviridae (53, 54) and very recently for hCMV (40) - although not in a placental
context - and confirm a general role of sEVs in modulating viral transmission for
many viral families (55, 56). Since placental sEVs are found both in maternal and

fetal sides at high quantities as soon as the early beginning of pregnancy (57, 58), we
hypothesized that they may facilitate hCMV dissemination not only in the placental
tissues but also towards the fetus, notably in the fetal brain, by enhancing hCMV
infectivity in fetal neural cells. Indeed, we observed that sEVs prepared from HIPECs
or first trimester placental explants enhanced hCMV infection of human NSCs.
Finally, our observations were confirmed using clinical samples. Strikingly, a
significant enhancement of infection was observed with sEVs isolated from the
amniotic fluid of an infected woman (with amniotic fluid positive for hCMV) whose
fetus showed severe neurodevelopmental impairments, in contrast to sEVs isolated
from the amniotic fluid of a non-infected pregnant woman, or an infected woman with
amniotic fluid negative for hCMV whose fetus showed no detectable developmental
defects. Hence, placental sEVs seem to be key players of hCMV dissemination
towards the fetus during congenital infection and may be one of the missing links
explaining the variety and severity of sequelae observed during viral congenital
infection.
408
409 **Materials and Methods**
410
Human ethic approval
The use of NSCs from human embryonic stem cells was approved by the French
authorities (Agence de la Biomédecine, authorization number SASB0920178S).
The biological resource center Germethèque (BB-0033-00081; declaration: DC-2014-
2202; authorization: AC-2015-2350) obtained the written consent from each patient
(CPP.2.15.27) for the use of human samples and their associated data. For first
trimester placenta explants, the steering committee of Germethèque gave its approval
for the realization of this study on Feb 5[th], 2019. The hosting request made to
Germethèque bears the number 20190201 and its contract is referenced under the
number 19 155C. For amniotic fluid, approval was obtained on July 12[th], 2019, the
hosting request bears the number 20190606 and the contract is referenced under the
number DIR-20190823022.
423
Cell lines
HIPECs were obtained from Dr T. Fournier (Inserm, Paris; Transfer agreement
n°170448). The expression of the cytokeratin 7 specific cytotrophoblastic marker was
verified (data not shown). They were cultured in DMEM / F12 medium (Gibco) at
50/50 ratio (v/v), in the presence of 10% fetal bovine serum (FBS, Sigma-Aldrich),
100 U/ml penicillin - 100 µg/ml streptomycin (Gibco) and 100 µg/ml normocin
(Invivogen).
MRC5 cells (RD-Biotech) were cultured in Dulbecco's Modified Eagle Medium
(DMEM with Glutamax, Gibco) with the same supplementation as HIPECs.
NSCs were obtained from Dr A. Benchoua (I-Stem, Evry, France), they were
produced from ES human cells (SA001, I-STEM, UMR861 France) (48) and
maintained in growth medium as described (7). Stem character of NSCs was

systematically assessed by immunofluorescence against Nestin and SOX2 proteins
(data not shown).
Cell cultures were checked for the absence of mycoplasma (Plasmotest, Invivogen,
Toulouse, France).
440

Virus production, titration and infection
442 The endotheliotropic VHL/E strain of hCMV - a gift from Dr C. Sinzger, University
443 of Ulm, Germany - was used in this study (59). Viral stocks were obtained upon
444 amplification of the virus on MRC5 cells, concentrated by ultracentrifugation, and
445 titrated by indirect immunofluorescence as described (7). In some experiments, virus
446 titration was also performed by qPCR from cell culture supernatants (60).
447

sEV preparation
Culture medium was previously depleted from EVs to obtain "Exofree" medium:
DMEM supplemented with 20 % FBS was ultracentrifuged at 100,000 g for 16 hours
at 4 °C and filtered at 0.22 µm. Exofree medium was then obtained by a 1:1 dilution
with F12 to reach 10 % FBS.
4 million HIPECs were seeded in 150 cm^2 flask, with 6 flasks per condition. 24 h
later, cells were infected or not by hCMV at multiplicity of infection (MOI) of 10
(Supplementary Figure S1A). Culture medium was replaced by Exofree medium 24 h
post-infection, after having previously ensured that this did not affect the cell growth
(Supplementary Figure S1E). Culture supernatants were collected at 48 h and 72 h
post-infection. Medium were pooled for each condition (non-infected or infected) and
submitted to sEV preparation protocol. Cell number at the end of the experiment was
determined by counting upon trypsinization (Supplementary Figure 1D) and cell
viability evaluated by trypan blue.
Procedures for sEV preparation were realized as described previously (25), according
to ISEV guidelines (27). All relevant data were submitted to the EV-TRACK
knowledgebase (EV-TRACK ID: EV210154) and obtained an EV-METRIC score of
100 % for HIPEC and placental explant EVs (61).
466

Nanoparticle tracking analysis (NTA)
468 sEV preparations were tracked using a NanoSight LM10 (Malvern Panalytical)
469 equipped with a 405 nm laser. Videos were recorded three times for each sample at
470 constant temperature (22 °C) during 60 s and analyzed with NTA Software 2.0
471 (Malvern instruments Ltd). Data were analyzed with Excel and GraphPad Prism (v8)
472 softwares.
473

Transmission electron microscopy and immunolabeling electron microscopy
Procedures were performed essentially as described (25, 62, 63). Immunodetection
was carried out with the following primary antibodies: mouse anti-human CD63
(Abcam ab23792), mouse anti-human CD9 or mouse anti-human CD81 (both from Dr
E. Rubinstein, Université Paris-Sud, Institut André Lwoff, Villejuif, France).
Secondary incubation was performed with a rabbit anti mouse Fc fragment (Dako

Agilent Z0412), then grids were incubated with Protein A-Gold 10 nm (Cell
Microscopy Center, Department of Cell Biology, Utrecht University). All samples
were observed with a Tecnai Spirit electron microscope (FEI, Eindhoven, The
Netherlands), and digital acquisitions were made with a numeric 4k CCD camera
(Quemesa, Olympus, Münster, Germany). Images were analysed with iTEM software
(EMSIS) and statistical studies were done with GraphPad Prism software (v8).
486
Western blot
Western blots were realized as previously described (25), by using the following
primary antibodies: mouse anti-CD81 (200 ng/ml, Santa-Cruz), mouse anti-CD63
(500 ng/ml, BD Pharmingen), mouse anti-CD9 (100 ng/ml, Millipore), rabbit anti-
Tsg101 (1 µg/ml, Abcam), rabbit anti-Alix (1 µg/ml, Abcam), mouse anti-Thy1 (0.5
µg/ml, Biolegend), rabbit anti-Tom20 (1/500, Sigma) or goat anti-Calnexin (2 µg/ml,
Abcam). After incubation with the secondary antibody, membranes were visualized
using the Odyssey Infrared Imaging System (LI-COR Biosciences).
495
Quantitative proteomic analysis
Sample preparation
Protein samples in Laemmli buffer (3 biological replicates of sEVs preparation from
non-infected and hCMV-infected cytotrophoblasts cells) were submitted to reduction
and alkylation (30 mM DTT and 90 mM iodoacetamide, respectively). Protein
samples were digested with trypsin on S-trap Micro devices (Protifi) according to
manufacturer's protocol, with the following modifications: precipitation was
performed using 545 µl S-Trap buffer and 1 µg Trypsin was added per sample for
digestion.
NanoLC-MS/MS analysis
Peptides were analyzed by nanoLC-MS/MS using an UltiMate 3000 RSLCnano
system coupled to a Q-Exactive-Plus mass spectrometer (Thermo Fisher Scientific,
Bremen, Germany). Five µL of each sample were loaded on a C-18 precolumn (300
µm ID x 5 mm, Dionex) in a solvent made of 5 % acetonitrile and 0.05 % TFA and at
a flow rate of 20 µL/min. After 5 min of desalting, the precolumn was switched
online with the analytical C-18 column (75 µm ID x 15 cm, Reprosil C18)
equilibrated in 95 % solvent A (5 % acetonitrile, 0.2 % formic acid) and 5 % solvent
B (80 % acetonitrile, 0.2 % formic acid). Peptides were eluted using a 5 to 50 %
gradient of solvent B over 105 min at a flow rate of 300 nL/min. The Q-Exactive-Plus
was operated in a data-dependent acquisition mode with the XCalibur software.
Survey scan MS were acquired in the Orbitrap on the 350-1500 m/z range with the
resolution set to a value of 70000. The 10 most intense ions per survey scan were
selected for HCD fragmentation. Dynamic exclusion was employed within 30 s to
prevent repetitive selection of the same peptide. At least 3 injections were performed
for each sample.
Bioinformatics data analysis of mass spectrometry raw files
Raw MS files were processed with the Mascot software for database search and with
Proline (64) for label-free quantitative analysis. Data were searched against *Human*

herpesvirus 5 and Human entries of the UniProtKB protein database (Human
betaherpesvirus 5 clone VHL-E-BAC19 and release Uniprot Swiss-Prot February
2018). Carbamidomethylation of cysteines was set as a fixed modification, whereas
oxidation of methionine was set as variable modification. Specificity of trypsin/P
digestion was set for cleavage after K or R, and two missed trypsin cleavage sites
were allowed. The mass tolerance was set to 10 ppm for the precursor and to 20 mmu
in tandem MS mode. Minimum peptide length was set to 7 amino acids, and
identification results were further validated in Proline by the target decoy approach
using a reverse database at both a PSM and protein false-discovery rate of 1%. After
mean of replicate injections, the abundance values were log2 transformed and missing
values were replaced by random numbers drawn from a normal distribution with a
width of 0.3 and down shift of 1.8 using the Perseus toolbox (version 1.6.7.0). For
statistical analysis, a Student *t*-test (two-tailed *t*-test, equal variances) was then
performed on log2 transformed values to analyse differences in protein abundances in
all biologic group comparisons. Significance level was set at $p = 0.05$, and log2
ratios were considered relevant if higher than 1 or lower than -1. The mass
spectrometry proteomics data have been deposited to the ProteomeXchange
Consortium via the PRIDE (65) partner repository with the dataset identifier
PXD029146.
543
544 Functional proteomic data analysis
Volcano plot was established for proteins whose mean abundance exhibited a log2
ratio higher than 1 or lower than -1 and when Student's *t*-test *p*-values were <= 0.05
between the infected and the non-infected conditions. The list of human proteins
exhibiting a normalized mean protein abundance log2 ratio > 1 or < -1 between sEVs
from non-infected or hCMV-infected samples was used as an input for analysis with
QIAGEN Ingenuity Pathway Analysis (IPA) (66). Results from IPA biological
functions and diseases analysis were filtered to retrieve annotations having an
absolute activation z-score > 1 and defined by less than 150 molecules. The resulting
annotations were manually curated to remove redundant annotations sharing identical
genes, keeping annotations defined by the greater number of molecules.
555
556 Flow cytometry analysis
After incubation of cells with PKH67-stained sEVs, cells were washed twice with
PBS and trypsinized, before proceeding to flow cytometry analysis. PKH67 positive
cells were analyzed on a Macsquant VYB Flow Cytometer (Miltenyi Biotec), by
using FCS and FITC fluorescence parameters, and by subtracting cell
autofluorescence background. Data were analyzed with FlowJo (BD) and GraphPad
Prism (v8) software.
563
564 Immunofluorescence
Cells were fixed using 4 % PFA (Electron microscopy Sciences) at room temperature
for 20 min. Permeabilization was performed with PBS 0.3 % Triton-X100
(Thermofisher scientific) for 10 min, followed by 1 h incubation in blocking buffer

(PBS with 5 % FBS). Incubation with primary antibodies diluted in blocking buffer
was carried out overnight at 4 °C, against hCMV immediate early protein 1 and 2 (1
µg/ml; Abcam IE1/IE2 CH160 ab53495), nestin (4 µg/ml; Abcam 10C2 ab 22035), or
SOX2 (1/500 of stock; Cell Signaling D6D9 #3579). Secondary antibody incubation
(Goat anti mouse or rabbit - Alexa-fluor 488 or 594 (2 µg/ml; Thermo Fischer
Scientific)) was performed at room temperature for 1 h. For actin staining, Alexa-
fluor 568 phalloidin (5 µg/ml; Thermo Fischer Scientific A12380) was incubated on
cells overnight at 4°C. DAPI staining (1 µg/ml; Sigma) was performed for 10 min at
room temperature. ProLong Gold without DAPI (Thermo Fischer Scientific) was used
for coverslip mounting.
Widefield acquisitions were realized using Apotome microscope (Zeiss) and confocal
acquisitions were made on SP8-STED microscope (Leica). Image processing was
performed using ImageJ. GraphPad Prism (v8) software was used to perform data
statistical analysis.
582
Placental histoculture
Placental histocultures were carried out as described (25, 49) on first trimester
placentas (4 placentas; mean = 13.11 ± 0.49 (SEM) weeks of amenorrhea, *i.e.,* 11.11 ±
0.49 weeks of pregnancy; age of the women: mean = 23 ± 1.5 (SEM) year-old).
Briefly, trophoblastic villi were dissected in small explants and infected or not by
hCMV overnight before deposition on gelatin sponges (Gelfoam, Pfizer) in Exofree
medium. Conditioned medium was collected and renewed every 3 to 4 days. At 14
days of culture, collected medium was pooled for each condition and used to perform
sEV preparation. Placental explants were weighed for normalization of resuspension
volume.
593
594 **Amniotic fluid collection**
595 Remaining amniotic fluid (AF) collected during classic patient care was saved for the
596 present study. It was subjected to a 10 min centrifugation at 3000 rpm in order to
597 remove the cells and large debris before its storage at -80 °C. The sEV preparation
598 was carried out according to the procedure described above.
599
600
601 **Acknowledgments**
602
We greatly thank C. Chouquet, B. Rauwel, M. Mouysset, L. Battut, C.
Mengelle, S. Lhomme, F. Chauvrier, P. Verdy, N. Kopf, as well as the whole
ViNeDys team, for their technical assistance and their numerous advice and
discussions which allowed the progress of this work. We thank also AL. Iscache, V.
Duplan-Eche and F. L'Faqihi-Olive, from the cytometry facility of Infinity, as well as
S. Allart and D. Daviaud, from the imaging facility of Infinity, and finally the GeT
technical service of Infinity. We thank the medical and paramedical staff of the
gynecology unit at Paule de Viguier Hospital, the patients who agreed to participate in
the study, as well as L. Bujan and M. Aubry, from the Germethèque. We finally

warmly thank E. Haanappel from the Institute of Pharmacology and Structural Biology, who granting us access to the Nanosight device, and Audrey Esclatine and Daniel Gonzalez-Dunia for critical reading of the manuscript and insightful comments.

This project has received financial support from the French Biomedicine Agency, institutional grants from Inserm, CNRS and Toulouse 3 University. This project is part of the doctorate thesis of Mathilde Bergamelli, who was funded by the Ministry of Education and Research (MESR). The TEM experiments were performed on PICT-IBiSA, Institut Curie, Paris, member of the France-BioImaging national research infrastructure, and were supported by the French National Research Agency through the "Investments for the Future" program (France-BioImaging, ANR-11-INSB-04), supported by the CelTisPhyBio Labex (N° ANR-11-LB0038) part of the IDEX PSL (N°ANR-10-IDEX-0001-02 PSL). The proteomic part of this project was supported in part by the Région Occitanie, European funds (Fonds Européens de Développement Régional, FEDER), Toulouse Métropole, and by the French Ministry of Research with the Investissement d'Avenir Infrastructures Nationales en Biologie et Santé program (ProFI, Proteomics French Infrastructure project, ANR-10-INBS-08).

References

1. Cannon MJ, Schmid DS, & Hyde TB (2010) Review of cytomegalovirus seroprevalence and demographic characteristics associated with infection. *Reviews in medical virology* 20(4):202-213.

2. Leruez-Ville M, Foulon I, Pass R, & Ville Y (2020) Cytomegalovirus infection during pregnancy: state of the science. *American journal of obstetrics and gynecology* 223(3):330-349.

3. Pereira L, Maidji E, Fisher SJ, McDonagh S, & Tabata T (2007) HCMV persistence in the population: potential transplacental transmission. *Human Herpesviruses: Biology, Therapy, and Immunoprophylaxis*, eds Arvin A, Campadelli-Fiume G, Mocarski E, Moore PS, Roizman B, Whitley R, & Yamanishi KCambridge).

4. Pereira L, Tabata T, Petitt M, & Fang-Hoover J (2017) Congenital cytomegalovirus infection undermines early development and functions of the human placenta. *Placenta* 59 Suppl 1:S8-S16.

5. Tabata T, *et al.* (2012) Cytomegalovirus impairs cytotrophoblast-induced lymphangiogenesis and vascular remodeling in an in vivo human placentation model. *Am J Pathol* 181(5):1540-1559.

6. Tabata T, Petitt M, Fang-Hoover J, Zydek M, & Pereira L (2016) Persistent Cytomegalovirus Infection in Amniotic Membranes of the Human Placenta. *Am J Pathol* 186(11):2970-2986.

7. Rolland M, *et al.* (2016) PPARgamma Is Activated during Congenital Cytomegalovirus Infection and Inhibits Neuronogenesis from Human Neural Stem Cells. *PLoS Pathog* 12(4):e1005547.

8. Rolland M, *et al.* (2021) Human cytomegalovirus infection is associated with increased expression of the lissencephaly gene PAFAH1B1 encoding LIS1 in neural stem cells and congenitally infected brains. *J Pathol* 254(1):92-102.

9. Sarker S, *et al.* (2014) Placenta-derived exosomes continuously increase in maternal circulation over the first trimester of pregnancy. *Journal of translational medicine* 12:204.

10. Luo SS, *et al.* (2009) Human villous trophoblasts express and secrete placenta-
specific microRNAs into maternal circulation via exosomes. *Biol Reprod* 81(4):717-
729.
11. Colombo M, Raposo G, & Thery C (2014) Biogenesis, secretion, and intercellular
interactions of exosomes and other extracellular vesicles. *Annual review of cell and*
developmental biology 30:255-289.
12. van Niel G, D'Angelo G, & Raposo G (2018) Shedding light on the cell biology of
extracellular vesicles. *Nature reviews. Molecular cell biology* 19(4):213-228.
13. Kalluri R & LeBleu VS (2020) The biology, function, and biomedical applications of
exosomes. *Science* 367(6478).
14. Ishida Y, *et al.* (2015) Maternal peripheral blood natural killer cells incorporate
placenta-associated microRNAs during pregnancy. *Int J Mol Med* 35(6):1511-1524.
15. Chaiwangyen W, *et al.* (2020) MiR-519d-3p in Trophoblastic Cells: Effects, Targets
and Transfer to Allogeneic Immune Cells via Extracellular Vesicles. *Int J Mol Sci*
21(10).
16. Li H, *et al.* (2020) Internalization of trophoblastic small extracellular vesicles and
detection of their miRNA cargo in P-bodies. *J Extracell Vesicles* 9(1):1812261.
17. Salomon C, *et al.* (2014) Extravillous trophoblast cells-derived exosomes promote
vascular smooth muscle cell migration. *Front Pharmacol* 5:175.
18. Sadovsky Y, *et al.* (2020) Placental small extracellular vesicles: Current questions and
investigative opportunities. *Placenta* 102:34-38.
19. Chiarello DI, *et al.* (2018) Foetoplacental communication via extracellular vesicles in
normal pregnancy and preeclampsia. *Mol Aspects Med* 60:69-80.
20. Mitchell MD, *et al.* (2015) Placental exosomes in normal and complicated pregnancy.
American journal of obstetrics and gynecology 213(4 Suppl):S173-181.
21. Salomon C & Rice GE (2017) Role of Exosomes in Placental Homeostasis and
Pregnancy Disorders. *Prog Mol Biol Transl Sci* 145:163-179.
22. Salomon C, *et al.* (2016) Gestational Diabetes Mellitus Is Associated With Changes in
the Concentration and Bioactivity of Placenta-Derived Exosomes in Maternal
Circulation Across Gestation. *Diabetes* 65(3):598-609.
23. Jin J & Menon R (2018) Placental exosomes: A proxy to understand pregnancy
complications. *Am J Reprod Immunol* 79(5):e12788.
24. Malnou EC, Umlauf D, Mouysset M, & Cavaille J (2019) Imprinted MicroRNA Gene
Clusters in the Evolution, Development, and Functions of Mammalian Placenta.
Front Genet 9:706.
25. Bergamelli M, *et al.* (2021) Human Cytomegalovirus Infection Changes the Pattern of
Surface Markers of Small Extracellular Vesicles Isolated From First Trimester
Placental Long-Term Histocultures. *Front Cell Dev Biol* 9:689122.
26. Pavan L, *et al.* (2003) Human invasive trophoblasts transformed with simian virus 40
provide a new tool to study the role of PPARgamma in cell invasion process.
Carcinogenesis 24(8):1325-1336.
27. Thery C, *et al.* (2018) Minimal information for studies of extracellular vesicles 2018
(MISEV2018): a position statement of the International Society for Extracellular
Vesicles and update of the MISEV2014 guidelines. *J Extracell Vesicles* 7(1):1535750.
28. Li Q, Fischer E, & Cohen JI (2016) Cell Surface THY-1 Contributes to Human
Cytomegalovirus Entry via a Macropinocytosis-Like Process. *J Virol* 90(21):9766-
9781.
29. Li Q, Wilkie AR, Weller M, Liu X, & Cohen JI (2015) THY-1 Cell Surface Antigen (CD90)
Has an Important Role in the Initial Stage of Human Cytomegalovirus Infection. *PLoS*
Pathog 11(7):e1004999.

30. Chaumorcel M, *et al.* (2012) The human cytomegalovirus protein TRS1 inhibits
autophagy via its interaction with Beclin 1. *J Virol* 86(5):2571-2584.
31. Mouna L, *et al.* (2016) Analysis of the role of autophagy inhibition by two
complementary human cytomegalovirus BECN1/Beclin 1-binding proteins.
Autophagy 12(2):327-342.
32. Taisne C, *et al.* (2019) Human cytomegalovirus hijacks the autophagic machinery and
LC3 homologs in order to optimize cytoplasmic envelopment of mature infectious
particles. *Scientific reports* 9(1):4560.
33. Karniely S, *et al.* (2016) Human Cytomegalovirus Infection Upregulates the
Mitochondrial Transcription and Translation Machineries. *mBio* 7(2):e00029.
34. Federspiel JD, *et al.* (2020) Mitochondria and Peroxisome Remodeling across
Cytomegalovirus Infection Time Viewed through the Lens of Inter-ViSTA. *Cell reports*
32(4):107943.
35. Combs JA, *et al.* (2020) Human Cytomegalovirus Alters Host Cell Mitochondrial
Function during Acute Infection. *J Virol* 94(2).
36. Monk CH & Zwezdaryk KJ (2020) Host Mitochondrial Requirements of
Cytomegalovirus Replication. *Curr Clin Microbiol Rep* 7(4):115-123.
37. Seo JY, Yaneva R, Hinson ER, & Cresswell P (2011) Human cytomegalovirus directly
induces the antiviral protein viperin to enhance infectivity. *Science* 332(6033):1093-
1097.
38. Zicari S, *et al.* (2018) Human cytomegalovirus-infected cells release extracellular
vesicles that carry viral surface proteins. *Virology* 524:97-105.
39. Turner DL, Korneev DV, Purdy JG, de Marco A, & Mathias RA (2020) The host
exosome pathway underpins biogenesis of the human cytomegalovirus virion. *Elife*
9.
40. Streck NT, Zhao Y, Sundstrom JM, & Buchkovich NJ (2020) Human Cytomegalovirus
Utilizes Extracellular Vesicles To Enhance Virus Spread. *J Virol* 94(16).
41. Mocarski ES, Kemble GW, Lyle JM, & Greaves RF (1996) A deletion mutant in the
human cytomegalovirus gene encoding IE1(491aa) is replication defective due to a
failure in autoregulation. *Proceedings of the National Academy of Sciences of the*
United States of America 93(21):11321-11326.
42. Nevels M, Paulus C, & Shenk T (2004) Human cytomegalovirus immediate-early 1
protein facilitates viral replication by antagonizing histone deacetylation.
Proceedings of the National Academy of Sciences of the United States of America
101(49):17234-17239.
43. Wiebusch L & Hagemeier C (1999) Human cytomegalovirus 86-kilodalton IE2 protein
blocks cell cycle progression in G(1). *J Virol* 73(11):9274-9283.
44. Cristea IM, *et al.* (2010) Human cytomegalovirus pUL83 stimulates activity of the
viral immediate-early promoter through its interaction with the cellular IFI16
protein. *J Virol* 84(15):7803-7814.
45. Arcangeletti MC, *et al.* (2011) Cell-cycle-dependent localization of human
cytomegalovirus UL83 phosphoprotein in the nucleolus and modulation of viral gene
expression in human embryo fibroblasts in vitro. *J Cell Biochem* 112(1):307-317.
46. Lukashchuk V, McFarlane S, Everett RD, & Preston CM (2008) Human
cytomegalovirus protein pp71 displaces the chromatin-associated factor ATRX from
nuclear domain 10 at early stages of infection. *J Virol* 82(24):12543-12554.
47. Fu YZ, *et al.* (2017) Human Cytomegalovirus Tegument Protein UL82 Inhibits STING-
Mediated Signaling to Evade Antiviral Immunity. *Cell host & microbe* 21(2):231-243.
48. Boissart C, Nissan X, Giraud-Triboult K, Peschanski M, & Benchoua A (2012) miR-125
potentiates early neural specification of human embryonic stem cells. *Development*
139(7):1247-1257.

49. Lopez H, *et al.* (2011) Novel model of placental tissue explants infected by cytomegalovirus reveals different permissiveness in early and term placentae and inhibition of indoleamine 2,3-dioxygenase activity. *Placenta* 32(7):522-530.

50. Benard M, *et al.* (2014) Human cytomegalovirus infection induces leukotriene B4 and 5-lipoxygenase expression in human placentae and umbilical vein endothelial cells. *Placenta* 35(6):345-350.

51. Dixon CL, *et al.* (2018) Amniotic Fluid Exosome Proteomic Profile Exhibits Unique Pathways of Term and Preterm Labor. *Endocrinology* 159(5):2229-2240.

52. Thery C, Amigorena S, Raposo G, & Clayton A (2006) Isolation and characterization of exosomes from cell culture supernatants and biological fluids. *Curr Protoc Cell Biol* Chapter 3:Unit 3 22.

53. Bello-Morales R & Lopez-Guerrero JA (2018) Extracellular Vesicles in Herpes Viral Spread and Immune Evasion. *Front Microbiol* 9:2572.

54. Cone AS, York SB, & Meckes DG, Jr. (2019) Extracellular Vesicles in Epstein-Barr Virus Pathogenesis. *Curr Clin Microbiol Rep* 6(3):121-131.

55. Lenassi M, *et al.* (2010) HIV Nef is secreted in exosomes and triggers apoptosis in bystander CD4+ T cells. *Traffic* 11(1):110-122.

56. Grunvogel O, *et al.* (2018) Secretion of Hepatitis C Virus Replication Intermediates Reduces Activation of Toll-Like Receptor 3 in Hepatocytes. *Gastroenterology* 154(8):2237-2251 e2216.

57. Liu H, *et al.* (2018) Estimation of the burden of human placental micro- and nano-vesicles extruded into the maternal blood from 8 to 12 weeks of gestation. *Placenta* 72-73:41-47.

58. Miranda J, *et al.* (2018) Placental exosomes profile in maternal and fetal circulation in intrauterine growth restriction - Liquid biopsies to monitoring fetal growth. *Placenta* 64:34-43.

59. Stegmann C, Rothemund F, Laib Sampaio K, Adler B, & Sinzger C (2019) The N Terminus of Human Cytomegalovirus Glycoprotein O Is Important for Binding to the Cellular Receptor PDGFRalpha. *J Virol* 93(11).

60. Mengelle C, *et al.* (2016) Performance of a completely automated system for monitoring CMV DNA in plasma. *J Clin Virol* 79:25-31.

61. Consortium E-T, *et al.* (2017) EV-TRACK: transparent reporting and centralizing knowledge in extracellular vesicle research. *Nat Methods* 14(3):228-232.

62. Hurbain I, Romao M, Bergam P, Heiligenstein X, & Raposo G (2017) Analyzing Lysosome-Related Organelles by Electron Microscopy. *Methods Mol Biol* 1594:43-71.

63. Raposo G, *et al.* (1996) B lymphocytes secrete antigen-presenting vesicles. *The Journal of experimental medicine* 183(3):1161-1172.

64. Bouyssie D, *et al.* (2020) Proline: an efficient and user-friendly software suite for large-scale proteomics. *Bioinformatics* 36(10):3148-3155.

65. Perez-Riverol Y, *et al.* (2019) The PRIDE database and related tools and resources in 2019: improving support for quantification data. *Nucleic acids research* 47(D1):D442-D450.

66. Thomas S & Bonchev D (2010) A survey of current software for network analysis in molecular biology. *Hum Genomics* 4(5):353-360.

     Figures and Tables

     Figure 1: Characterization of sEVs from NI and hCMV infected HIPECs.

811
812
A) Yield of sEV recovered from non-infected (NI) or infected (hCMV) HIPECs. *, p
= 0.0464 by paired t-test for 7 independent experiments. B) Comparison of mean size
(left histogram) and mode size (right histogram) between sEVs from non-infected
(NI) or infected (hCMV) HIPECs. Histograms show the mean ± SEM of three
independent experiments. ns: non-significant by Mann Whitney test. C) Electron

microscopy images of sEVs (indicated by an arrow) prepared from non-infected (NI)
or infected (hCMV) HIPECs. Magnification = 26000 X. Scale bar = 100 nm. Images
are representative of at least three independent experiments. D) TEM observation of
sEV - isolated from non-infected (NI) or infected (hCMV) HIPECs - which were
immunogold-labelled for CD9, CD81 or CD63, and revealed with Protein A-gold
particle of 10 nm diameter. Scale bar = 100 nm. Magnification = 26000 X. In CD63
IEM, only one example of positive vesicle, representing around 1-5 % of sEVs
isolated upon infection, is shown, the other being negative (see Supplementary Figure
3 for wide field image). E) Western-blot realized on either whole cell lysates (left
wells) or purified sEVs (right wells), from non-infected (NI) or infected (hCMV)
HIPECs. Proteins of interest and their corresponding molecular weight are indicated
on the right of the Figure, with a smear for CD63 due to the non-reducing conditions
of the western blot, which preserve its rich glycosylated pattern.
831

Figure 2: Proteomic analysis of sEV composition upon infection of HIPECs by hCMV.

A) Volcano-plot representing differences in normalized mean protein abundance in sEVs hCMV *versus* sEVs NI. Human and viral proteins exhibiting significant differences between the two conditions are represented by circles and triangles, respectively (Student T-test *p*-value <=0.05 and log2 ratio >=1 or <=-1). Red: over-represented proteins; Blue: under- represented proteins. B) Dot plot representation of the top diseases and biological functions associated with human proteins exhibiting an absolute normalized mean abundance log2 ratio greater than 1 or lower than -1, in

sEVs hCMV *versus* sEVs NI. Top diseases and biological functions associated with
changes in the protein content of sEVs upon hCMV-infection were identified using
QIAGEN Ingenuity Pathway Analysis (IPA). Size of the circles depends on the
number of the proteins identified in the corresponding pathway; level of blue intensity
depends on the *p*-value. C) Human proteins associated with the predicted increased
activation state of autophagy pathway as determined by IPA. D) Heatmap
representation of the biological functions associated with hCMV viral proteins
expressed in sEVs hCMV.
851

852 **Figure 3: Internalization of sEVs isolated from non-infected HIPECs in fetal**
853 **MRC5 cells.**

854
A) Confocal images of fluorescence microscopy carried out on MRC5 cells after 16h
incubation with PKH67-labelled sEVs. Blue: DAPI; Red: Phalloidin; Green: PKH67.
Scale bar: 100 µm. Magnification = 63 X. The right image corresponds to the
orthogonal projection of the cell z-axis. B) Histogram representing the percentage of
PKH67 positive cells along time, upon incubation of MRC5 cells with sEVs. Bars
represent the mean ± SEM of three independent experiments. C) Monitoring of
PKH67-labeled sEVs internalization by MRC5 cells by flow cytometry. Dot plots
represent MRC5 cell fluorescence upon incubation with PKH67-stained sEVs (200
sEVs/cell) for cells that have not been incubated with sEVs (NT, non-treated), or
upon 2 h or 16 h of incubation. X-axis: PKH-67 fluorescence intensity; Y-axis: FSC.
Gate indicates cells positive for PKH67.
866
867

868 **Figure 4: Effect of HIPEC sEVs on MRC5 cell permissiveness for hCMV.**

869
A) Experimental procedure (NT=non-treated). B) Immunofluorescence performed on
MRC5 cells against IE viral antigen (blue: DAPI; red: IE1/2). a- Non-infected control
MRC5 cells upon 24 h incubation with sEVs isolated from hCMV-infected HIPECs.
b,c,d- MRC5 cells were either non-treated (b- NT) or incubated during 2 h with sEVs
prepared from non-infected (c- sEV NI) or infected HIPECs (d- sEV hCMV) with 50
sEV per cell, then infected during 24 h with hCMV at a MOI of 0.5 before proceeding
to immunofluorescence. Scale bar: 200 μm. C-D) MRC5 cells were incubated with
sEVs prepared from non-infected (sEV NI) or hCMV-infected (sEV hCMV) HIPECs
and infected by hCMV at a MOI of 0.5, concomitantly (C) or 2 h after sEV incubation
(D). Three increasing doses of sEVs were used in these experiments (20, 50 or 200
sEV per MRC5 cell, from left to right). 24 h later, expression of IE antigen was
assessed by immunofluorescence. Quantification of the percentage of IE positive cells
was carried out and normalized by the percentage of cells infected by hCMV without
any sEV (NT). Each dot is an independent experiment and corresponds to the mean of
the counting of 10 fields, with around 70 cells counted, *i.e.,* around 700 cells per dot.
n = 4 to 10 independent experiments. Since sEVs used in infection assays were

prepared each time in parallel between non-infected or hCMV-infected HIPECs from
a given batch, statistical analysis was done by pairing the results between sEV NI and
sEV hCMV for each independent experiment. ns, non-significant; *, $p < 0.05$; *** $p <$
0.001 by paired t-test.
890

891 **Figure 5: Effect of sEVs from different sources on neural stem cells**
892 **permissiveness for hCMV.**

893
NSCs were incubated with sEVs (200 sEVs per cell) prepared from HIPECs (A), *ex*
vivo first trimester placental histoculture (B), or amniotic fluids from naturally
infected patients (C), then infected by hCMV at a MOI of 3. 24 h upon infection,
expression of IE antigen was assessed by immunofluorescence. Each dot is an
independent experiment and corresponds to the mean of the counting of 10 fields,
with around 70 cells counted, *i.e.* around 700 cells per dot. n = 3 to 7 independent
experiments. For (A) and (B), HIPECs or placental explants were either non-infected
(sEV NI) or hCMV-infected (sEV hCMV). Quantification of the percentage of IE
positive cells was carried out and normalized by the percentage of infection of cells
infected by hCMV without any sEV (NT). Since sEVs used in functional assays were
prepared each time in parallel between non-infected or hCMV-infected
cytotrophoblasts or placental explants from a given batch, statistical analysis was
done by pairing the results between sEV NI and sEV hCMV for each independent
experiment. ns, non-significant; *, $p < 0.05$ by paired *t*-test. For (C), NSCs were
incubated with sEVs prepared from amniotic fluid (AF) from patient NP19 (non-

infected), NP6 (hCMV infected, AF negative for hCMV) or NP12 (hCMV infected,
AF positive for hCMV). Histograms represents the mean ± SEM of the percentage of
infected cells in each condition. The result of the ordinary one-way ANOVA test
performed is $p = 0.0006$ (***). Statistical comparison between the conditions were
done by Tukey's multiple comparisons test upon one-way ANOVA test. ns, non-
significant; ** $p < 0.005$; *** $p < 0.001$.
915

Supplementary Figure S1: Model system.

A) Timeline representing the HIPEC seeding, infection, medium renewal and collection before sEV preparation. D: days; MOI: Multiplicity of infection. B) HIPECs are permissive to hCMV infection. Immunofluorescence against Immediate Early 1/2 (IE) antigens was done on HIPECs either non-infected (NI) or at 48 h or 72 h post-infection (hpi). Green: IE; Blue: DAPI; Magnification: 20 x. C) Immunofluorescence realized against hCMV IE1/2 antigens in non-treated MRC5 cells (Mock), or upon incubation during 24 h with either hCMV at MOI 3 (hCMV), sEVs isolated from non-infected (sEV NI) or from hCMV-infected HIPECs (sEV hCMV). Magnification = 20 x. Blue (upper panel): DAPI; red (lower panel): IE. Below images are indicated the results of RT-qPCR realized against hCMV UL55 mRNAs on RNA extracted from MRC5 cells at 48 h post-incubation with hCMV or sEV preparations (+: amplification; -:

no amplification). Data are representative for at least three independent experiments. D) Total cell number counted at the time of sEV preparation, at 72 hpi, for non-infected (NI) or hCMV infected HIPECs. Results are representative of 15 independent experiments. *p<0,0001* by paired *t*-test. E) Comparison of cell growth between complete and Exofree culture medium, by counting of HIPEC number in 12-well plates in which 10^5 cells were initially seeded. Symbols represent mean ± SEM for three independent experiments. A 2-way ANOVA statistical test indicated no significant difference between complete and Exofree medium culture conditions (*p=0.4414*).

Supplementary Figure S2: Wide-field transmission electron microscopy.

NI

hCMV

TEM micrographs showing sEV from HIPEC either non-infected (NI; upper panel) or infected (hCMV; lower panel). Scale bar: 500 nm.

Supplementary Figure S3: Multiplex bead-based flow cytometry comparison of sEV prepared from non-infected or hCMV-infected HIPECs.

Bead-based multiplex analysis using the MACSPlex Exosome Kit, human (Miltenyi Biotec) were realized on sEV preparations by flow cytometry according to the manufacturer's instructions. This allowed the quantification of 39 different EV markers, distinguishable by flow cytometry by a specific PE and FITC labeling. The MACSQuant Analyzer 10 flow cytometer (Miltenyi Biotec) was used for analysis. The tool MACSQuantify was used to analyze data (v2.11.1746.19438). GraphPad Prism (v8) software was used to perform statistical analysis of the data. The heat-map represents the mean of 3 independent experiments, for different sEV markers indicated on the left column, for sEV prepared from non-infected of hCMV-infected HIPECs. Blue intensity is proportional to the level of expression calculated in Median Fluorescence Intensity, indicated on the right of the heat-map. ns, non-significant by two-way ANOVA.

Supplementary Figure S4: Wide field immuno-electron microscopy anti CD63.

NI

hCMV

TEM micrographs showing sEV from HIPECs either non-infected (NI; upper panel) or infected (hCMV; lower panel), after immunogold labelling against CD63. Arrow in the lower panel indicates a CD63 positive sEV. Scale bar: 500 nm.

Supplementary Figure S5: Complement to proteomic analysis.

A) Western-blot showing the enrichment of the cellular Thy-1 protein in sEVs from infected HIPECs (NI: non-infected). This result is representative of three independent experiments. B) Heat-map representing the different viral protein found in sEV isolated from infected HIPECs, classified depending on their expression timeline during hCMV infection (Immediate early, early or late).

Supplementary Figure S6: sEV internalization by MRC5 at 2 h post-incubation.

MCR5 cells, observed by confocal fluorescence microcopy, upon 2h of incubation with PKH67 stained sEVs (200 sEVs/cell) prepared from non-infected (panels B), hCMV-infected (panels C) or without sEV incubation (panels A). Green: PKH67 staining of sEV; red: actin staining by phalloidin; blue: DAPI. Scale bar: 10 μm.

Supplementary Video S1: MCR5 cells after 16 h incubation with PKH67 stained sEVs from NI HIPECs (50 sEVs/cells). Observation by confocal fluorescence microscopy. Blue: DAPI; Red: Phalloidin; Green: PKH67. Magnification = 63 X.

Supplementary Table S1: Excel file.

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD029146.

Supplementary Table S2: Bibliography references for establishment of Figure 2D and supplementary Figure S5B

Protein name	Gene name	Reference	Link
Capsid scaffolding protein	UL80	Loveland et al. J. Virol. 2007	https://doi.org/10.1128/JVI.01903-06
Capside smallest protein	UL48/49	Yu et al. J. Virol 2005	https://doi.org/10.1128/JVI.79.2.1327-1332.2005
DNA polymerase processivity factor	UL44	Silva et al. Virology 2011	https://doi.org/10.1016/j.virol.2011.06.008
Early phosphoprotein p84	UL112	Schommartz et al. J. Virol. 2017	https://doi.org/10.1128/JVI.00254-17
Glycoprotein B	GB / UL55	Isaacson et al. J. Virol 2009 Halary et al. Immunity 2002	https://doi.org/10.1128/JVI.01251-08 https://doi.org/10.1016/S1074-7613(02)00447-8
Glycoprotein H	UL75	Yurochko et al. J.Virol 1997	https://doi.org/10.1128/jvi.71.7.5051-5059.1997
Glycoprotein M	GM	Mach et al. J.Virol. 2020 Liu et al. Sci. Adv. 2021	https://doi.org/10.1128/JVI.79.4.2160-2170.2005 https://doi.org/10.1126/sciadv.abf3178
Glycoprotein N	UL73	Mach et al. J. Virol. 2005 Shimamura et al. J. Virol. 2006	https://doi.org/10.1128/JVI.79.4.2160-2170.2005 https://doi.org/10.1128/JVI.80.9.4591-4600.2006
Glycoprotein UL16	UL16	Dunn et al. JEM 2003 Müller et al. PloS Pathog. 2010	https://dx.doi.org/10.1084%2Fjem.20022059 https://doi.org/10.1371/journal.ppat.1000723
IE1	UL123	Mocarski et al. PNAS 1993 Nevels et al. PNAS 2004	https://dx.doi.org/10.1073%2Fpnas.93.21.11321 https://doi.org/10.1073/pnas.0407933101
IE2	UL122	Castillo et al. J. Virol. 2000 Paulus et al. Viruses 2009	https://doi.org/10.1128/JVI.74.17.8028-8037.2000 https://doi.org/10.3390/v1030760
IRS1	IRS1/US22	Child et al. J. Virol 2004 Marshall et al. J. Virol 2009	https://doi.org/10.1128/JVI.78.1.197-205.2004 https://doi.org/10.1128/JVI.02489-08
Large tegument protein deneddylase	UL48	Kim et al. J. Virol. 2016	https://doi.org/10.1128/JVI.02766-15
Major capsid protein	UL86	Furhmann et al. J. Infect. Dis. 2008	https://doi.org/10.1086/587692
Nuclear egress protein 1	UL53	Salm et al. J. Virol. 2009	https://doi.org/10.1128/JVI.02441-08
pp150	UL32	Bogdanow et al. PNAS 2013 Tandon et al. J. Virol 2008	https://dx.doi.org/10.1073%2Fpnas.1312235110 https://doi.org/10.1128/JVI.00533-08
pp28	UL99	Silva et al. J. Virol. 2003	https://doi.org/10.1128/JVI.77.19.10594-10605.2003
pp65	UL83	Cristea et al. J. Virol. 2010 Tomtishen et al. Virol. J. 2012 Arcangeletti et al. J. Cell Biochem. 2011	https://doi.org/10.1128/JVI.00139-10 https://dx.doi.org/10.1186%2F1743-422X-9-22 https://doi.org/10.1002/jcb.22928
pp71	UL82	Kalejta et al. Mol. Cell Biol. 2003 Tomtishen et al. Virol. J. 2012	https://dx.doi.org/10.1128%2FMCB.23.6.1885-1895.2003 https://dx.doi.org/10.1186%2F1743-422X-9-22
pp85	UL25	Nobre et al. eLife 2019	https://doi.org/10.7554/eLife.49894
Protein UL88	UL88	Kumar et al. J. Virol. 2020 Kalejta MMBR 2008	https://doi.org/10.1128/JVI.00474-20 https://doi.org/10.1128/MMBR.00040-07
RIR1	UL45	Patron et al. J. Gen. Virol. 2003	https://doi.org/10.1099/vir.0.19452-0
Tegument protein UL23	UL23	Feng et al. PloS Pathog. 2018 Feng et al. Front. Microbiol. 2021	https://doi.org/10.1371/journal.ppat.1006867 https://dx.doi.org/10.3389%2Ffmicb.2021.692515
Tegument protein UL51	UL51	Borst et al. J. Virol 2013	https://doi.org/10.1128/JVI.01955-12
Tegument protein US22	US22	Kalejta MMBR 2008	https://doi.org/10.1128/MMBR.00040-07
Tegument serine/threonine kinase	UL97	Krosky et al. J. Virol 2003 Hertel et al. PloS Pathog. 2007	https://doi.org/10.1128/JVI.77.2.905-914.2003 https://doi.org/10.1371/journal.ppat.0030006
Tegument UL35	UL35	Salsman et al. J. Virol. 2012 Fabits et al. Microorganisms 2020	https://doi.org/10.1128/JVI.05442-11 https://doi.org/10.3390/microorganisms8060790
Triplex capsid protein 1	UL46	Varnum et al. J. Virol 2004	https://doi.org/10.1128/JVI.78.20.10960-10966.2004
Triplex capsid protein 2	UL85	Varnum et al. J. Virol 2004	https://doi.org/10.1128/JVI.78.20.10960-10966.2004
TRS1	TRS1	Child et al. J. Virol 2004 Adamo et al. J. Virol 2004	https://doi.org/10.1128/JVI.78.1.197-205.2004 https://doi.org/10.1128/JVI.78.19.10221-10229.2004
UL37 immediate early glycoprotein	UL37	McCormick et al. J. Virol. 2003	https://doi.org/10.1128/JVI.77.1.631-641.2003

Supplementary Table S3

Patient	NP19	NP6	NP12
Age (y)	34	25	31
Time of AF punction (WA)	33	24	31
hCMV seroconversion	No (old immunity)	Yes	Yes
Time of seroconversion (WA)	NA	Between 3-5	<8
PCR hCMV of AF	Negative	Negative	Positive
Log PCR hCMV on AF	NA	NA	8 log
Fetal damage (ultrasound)	Chylothorax Bilateral hydrothorax Idiopathic hydramnios No infectious background or genetic abnormalities detected	20th percentile	Brain damage (periventricular hyperechoic halo, short corpus callosum, occipital horn cyst, germinal zone abnormalities) - Intrauterine growth retardation (3rd percentile)
Gestation end	Vaginal delivery at 36 WA	Vaginal delivery at term	Medical abortion at 32 WA

y: year; WA: week of amenorrhea; AF: amniotic fluid.