Functional divergence and spectral tuning of microbial rhodopsins from an ancestral proton pump

Cathryn D. Sephus[1], Evrim Fer[2,3], Amanda K. Garcia[3], Zachary R. Adam[4,5], Edward W. Schwieterman[5,6], Betül Kaçar[3*]

1. Scripps Institution of Oceanography, University of California at San Diego, La Jolla, CA, USA

2. Microbiology Doctoral Training Program, University of Wisconsin-Madison, Madison, WI, USA

3. Department of Bacteriology, University of Wisconsin-Madison, Madison, WI, USA

4. Department of Geoscience, University of Wisconsin-Madison, Madison, WI, USA

5. Blue Marble Space Institute of Science, Seattle, WA, USA

6. Department of Earth and Planetary Sciences, University of California, Riverside, CA, USA

*To whom correspondence should be addressed: bkacar@wisc.edu

Running title: Functional divergence and spectral tuning of microbial rhodopsins from an ancestral proton pump

Keywords: early Earth; phototrophy; rhodopsin; radiative transfer; origins of life; biosignature; habitability

ABSTRACT

For billions of years, life has continuously adapted to dynamic physical conditions near the Earth's surface. Fossils and other preserved biosignatures in the paleontological record are the most direct evidence for reconstructing the broad historical contours of this adaptive interplay. However, biosignatures dating to Earth's earliest history are exceedingly rare. Here, we combine phylogenetic inference of primordial rhodopsin proteins with modeled spectral features of the Precambrian Earth environment to reconstruct the paleobiological history of this essential family of photoactive transmembrane proteins. Our results suggest that ancestral microbial rhodopsins likely acted as light-driven proton pumps and were spectrally tuned toward the absorption of green light, which would have enabled their hosts to occupy depths in a water column or biofilm where UV wavelengths were attenuated. Subsequent diversification of rhodopsin functions and peak absorption frequencies track the diversification of surface ecological niches induced by the accumulation of atmospheric oxygen. Inferred ancestors retain distinct associations between extant functions and peak absorption frequencies. Our findings suggest that novel information encoded by biomolecules can be used as paleosensors for conditions of ancient, inhabited niches of host organisms not represented elsewhere in the paleontological record. The coupling of functional diversification and spectral tuning of this pervasive protein family underscores the utility of rhodopsins as universal testbeds for inferring remotely detectable biosignatures on inhabited planetary bodies.

INTRODUCTION

Decoding the complex relationships between life and the environments it inhabits is central to reconstructing the contours of our planet's history of habitability. These contours are critical for constraining the physical limits of biological functionality, identifying possibly 'universal' biological behavioral modes, and aiding the search for life elsewhere in our universe[1,2]. Planetary habitability is strongly influenced by solar insolation and photon radiance flux, which couple planetary and stellar evolution to the pervasiveness of life[3,4]. The Earth and sun, however, each possess idiosyncratic internal structures, stability points, and observational limitations that obfuscate the reconstruction of long-term habitability parameters.

The information encoded in life itself may provide novel insights into how our planet has maintained planetary habitability where geologic and stellar inferences fall short. The earliest

paleontological record lacks abundant, direct evidence of major metabolic and morphogenetic
features[5,6], but the contemporary bioinformatic record is incredibly rich and includes several
environment-sensitive biomolecular components. Reconstructing the ancestral sequences of
biomolecules that are strongly coupled to habitability parameters may provide a means of
extrapolating ancient biological phenotypes—and the environmental conditions that shaped
them—back to the Last Universal Common Ancestor (LUCA)[7-9].

Photosensitive enzymes are key intermediaries that connect intracellular chemical states,
extracellular substrate availability, and solar irradiance. These biomolecules constitute a
promising system for tracking ancient physical parameters not recorded in the geologic record. All
known phototrophic metabolisms on Earth rely on one of three energy-converting pigments which
transform light energy into chemical energy. These pigments include chlorophylls,
bacteriochlorophylls, and retinal[10]. Retinal-based pigment proteins, known as rhodopsins, have
been found in Archaea, Bacteria, Eukarya, and giant viruses[11,12]. Rhodopsin pervasiveness was
initially attributed to high rates of gene transfer between microbial organisms[13], but subsequent
analyses indicate that a rhodopsin-like functionality may have been present in LUCA[14].

Retinal-based phototrophy utilizes a light-capture system that is simpler than its chlorophyll-based
counterparts[15]. Rhodopsin pigments are composed of one seven-transmembrane opsin protein
and one retinal chromophore[16]. Photoisomerization of the retinal chromophore within the opsin
active site (also referred to as the retinal binding pocket) triggers a series of cyclical
conformational changes that can directly drive ion transport across the cellular membrane,
generating a proton-motive force that may be harnessed by ATP synthase to generate ATP[17].
The simple bipartite architecture of rhodopsins has also been harnessed by biology for uses
beyond ion transport, including both phototaxis and photophobia, substrate uptake, and starvation
prevention[18-21]. Rhodopsins can be tuned to absorb photons across much of the visible light
spectrum[22] and have been sampled from exposed terrestrial surfaces to the sub-photic marine
zone[23]. The structural simplicity and functional diversity of microbial (type I) rhodopsins may have
promoted their spread to global econiches[24]. Today, type I rhodopsins are among the dominant
energy-transducing agents that harvest solar energy in the surface ocean[10]. The combination of
cladistic pervasiveness, phylogenetic tractability, and niche ubiquity indicates that ancestral type
I rhodopsins are excellent candidate phylogenetic systems for characterizing long-term conditions
of habitability and paleoecology where geologic indicators are unavailable[25-27].

We reconstructed the evolutionary history of microbial type I rhodopsins to explore ancestral

properties that might serve as a guide for understanding early Earth paleoecology and habitability.

Protein sequence analyses of both extant and phylogenetically inferred ancestral rhodopsins were

used to constrain possible ancestral rhodopsin functions. In addition, maximum absorbance

wavelength of ancestral rhodopsins was predicted by a machine-learning algorithm that models

the relationship between spectral absorbance peaks and amino acid sequence[28]. Our analyses

suggest proton-pump functionality and green wavelength absorption for the last common ancestor

of ancestral type I rhodopsins. Green wavelength absorption is parsimonious with our modeled

spectral features of the Precambrian Earth environment, providing an adaptive advantage to

shallow coastal marine niches with abundant nutrient and metabolically useful light availability but

attenuated phototoxic UV irradiation. The proliferation of specialized functions and the spectral

tuning of rhodopsins tracks ecological upheaval wrought by the rise of atmospheric oxygen, with

significant accumulation marked by the interval known as the Great Oxidation Event (GOE)

approximately 2.3 billion years ago[29,30]. The pattern of niche proliferation imprinted upon the

rhodopsin protein network is broadly consistent with trends in oxygen-sensitive stable isotope

signatures and mineral distributions throughout the Precambrian[29,31]. The convergence of our

results from reconstructed, ancestral biomolecular characteristics and indirect physical models

constrains the manner by which phototrophic organisms have adapted to changing habitability

parameters throughout Earth's macroevolutionary history.

RESULTS AND DISCUSSION

Conservation and statistical support for ancestral rhodopsin residues important for

function and spectral tuning

We assembled a dataset of representative protein sequences with which to reconstruct the

evolutionary history and ancestral traits of microbial rhodopsins. BLASTp[32] was used to identify

452 homologous type I rhodopsin sequences from the National Center for Biotechnology

Information (NCBI) non-redundant and UniProt protein databases. We additionally identified 116

homologs belonging to heliorhodopsins, a recently discovered rhodopsin class that exhibits an

inverted membrane topology relative to type I rhodopsins and constitutes a distinct outgroup[33-35].

Animal, or type II, rhodopsins, though exhibiting a similar seven transmembrane architecture,

possess little sequence homology with microbial type I rhodopsins and were thus excluded from

the dataset[13,35]. Our type I rhodopsin sequence dataset comprises four major archaeal rhodopsin

groups: 1) bacteriorhodopsin, outward proton pumps; 2) halorhodopsin, inward chloride pumps; 3) sensory rhodopsin I, positive phototaxis mediators; and 4) sensory rhodopsin II, negative phototaxis mediators[36]. The dataset also includes two major bacterial rhodopsin groups: 1) proteorhodopsin and 2) xanthorhodopsin, both outward proton pumps, with the latter distinguished by the ability to bind a second carotenoid chromophore[36]. These taxonomically and functionally distinct lineages were resolved into their respective clades by maximum likelihood phylogenetic reconstruction (Fig. 1), producing a tree topology largely in agreement with previous analyses[11,37,38]. Alternate phylogenetic trees, generated by incorporating next-best-fit evolutionary models, reproduced this general topology (see Methods; Supplementary Fig. 1). Ancestral amino acid sequences were inferred by maximum likelihood from internal nodes of our rhodopsin phylogeny. Five of the oldest internal nodes were selected as targets for further phenotypic analysis: 1) Anc1, ancestral to bacteriorhodopsins and halorhodopsins, 2) Anc2, ancestral to sensory rhodopsins, 3) Anc3, ancestral to archaeal rhodopsins, 4) Anc4, ancestral to bacterial rhodopsins, and 5) Anc5, ancestral to all type I rhodopsins (Fig. 1, Supplementary Fig. 2).

Due to their deep position in the phylogenetic tree, mean site posterior probabilities for ancestors Anc1-Anc5 range between 0.55 and 0.76 (Fig. 2a; Supplementary Table 1). However, statistical support is enriched for 23 ancestral binding pocket residues (<5 Å from the retinal chromophore), constituting the most conserved element of microbial rhodopsin structure[39]. Mean posterior probabilities for binding pocket residues fall between 0.90 and 0.97 (Supplementary Table 1, 2). Nine of these residues have ≥95% mean pairwise identity across all extant sequences included in our dataset (Y83, W86, P91, G122, W182, Y185, W189, D212, and K216 in the bacteriorhodopsin of *Halobacterium salinarum*). Most have significant, experimentally determined contributions to rhodopsin function[39-43], including the strictly conserved K216 residue that is covalently bonded to retinal via a Schiff base linkage[40]. Other functions include maintenance of the internal charge environment for ion-pumping rhodopsins[39,44] or mediation of transducer interactions required for phototactic response in sensory rhodopsins[45]. Finally, binding pocket residues have been implicated as having outsized influence in rhodopsin color tuning[46,47]. Thus, the enriched statistical support for ancestral binding pocket residues permits sequence-based inferences of historical rhodopsin functionality.

Ancestral type I rhodopsins likely acted as light-driven proton pumps

Comparative analysis of extant variants provides compelling circumstantial evidence that primordial rhodopsins functioned as transporters but the evolutionary progression of functions or of photon absorptivity have not been explicitly demonstrated to date[37]. As an initial, global approach to constraining early rhodopsin functionality, 3D protein structures of maximum likelihood ancestral sequences Anc1-Anc5 were predicted with Phyre2[51] and used for functional annotation by the COFACTOR algorithm within the I-TASSER software pipeline[52]. Modeled ancestral rhodopsin structures preserve the characteristic seven-transmembrane alpha-helical fold, otherwise known as the G-protein coupled receptor fold[16]—of extant microbial type I homologs (Fig. 3). Kyte and Doolittle hydropathy plots for Anc1-Anc5 reveal seven highly hydrophobic regions, consistent with the conservation of seven transmembrane domains (Supplementary Fig. 3). Membrane localization of ancestral rhodopsins was also predicted by the SOSUI tool[53] (Supplementary Table 3). Additionally, the ~21-residue-long C-terminal sequence of these ancestral rhodopsins form a distinct hydrophilic region, indicating an orientation towards the cytoplasm as observed with extant type I rhodopsins[33]. Finally, the structure-based functional predictions are consistent with early type I rhodopsins that carried out phototransduction, formed protein-chromophore linkages, and acted as proton transmembrane transporters (Supplementary Table 4).

Given the comparatively high statistical support and functional significance of ancestral binding pocket residues, we interrogated these specific sequence features to provide complementary insights into plausible functionality of ancient rhodopsins. For extant, uncharacterized rhodopsins, functional annotation is often guided by a characteristic, three-residue motif that includes sites homologous to the D85, T89, and D96 residues (a.k.a., "DTD" motif) in bacteriorhodopsin (*H. salinarum*)[35,39,54,55]. These sites are located within a section of the third transmembrane helix ("Helix C"; Fig. 3c, Fig. 4) that constitutes a portion of the retinal binding pocket. Variation in this motif is strongly correlated with functional differentiation in rhodopsins. For example, "DTD" or "DTE" motifs (glutamic acid replaces the D96 residue in the latter) are characteristic of bacteriorhodopsins, xanthorhodopsins, and proteorhodopsins, which all function as light-driven proton pumps and exhibit similar photosensitive mechanisms[39]. By contrast, chloride-pumping halorhodopsins exhibit a "TSA" motif and sensory rhodopsins instead exhibit a "DTY" of "DTF" motif.

The functional correlation of the three-residue motif can be explained by the role these sites play

during ion binding and transfer. In proton-pumping rhodopsins, the negatively charged residues

of the DTD/DTE motifs are protonated during intermediate steps of the bacteriorhodopsin,

xanthorhodopsin, and proteorhodopsin photocycles, permitting unidirectional proton transport out

of the cytoplasm[39]. By contrast, the TSA motif typical of chloride-pumping rhodopsins lacks a

negative charge relative to the DTD/DTE motifs, which is compensated by the binding of the

chloride ion[55,56]. The functional significance of these motifs has been underscored by experiments

demonstrating the conversion of proton-pumping bacteriorhodopsins into chloride pumps by a

D85T mutation at the first site of the DTD motif[57]. In certain cases, functional predictions of extant

rhodopsins based on these characteristic motifs have been shown to prove reliable upon

experimental characterization[54].

At the deepest nodes in the rhodopsin phylogeny, the most likely ancestral motifs are: 1) Anc1,

DTD, 2) Anc2, DTY, 3) Anc3, DTD, 4) Anc4, DTE, and 5) Anc5, DTE (Fig. 3). These motifs suggest

a proton-pumping function for all ancestors bearing the DTD/DTE motifs, except for Anc2, which,

as the common ancestor to sensory rhodopsins I and II, exhibits the DTY motif consistent with

extant sensory rhodopsins. Most inferred residues at these three sites have posterior probabilities

>0.90 for Anc1-Anc5 (Supplementary Table 2). The exceptions are in Anc1, Anc3, and Anc5, for

which the third site in the motif has a posterior probability of 0.71, 0.54, or 0.61, respectively.

However, the next most likely ancestral states for these sites are either aspartic acid or glutamic

acid, maintaining the negatively charged residue observed in extant proton transporters. Thus, a

motif consistent with an ancestral proton-pumping function is well supported for Anc1, Anc3,

Anc4, and Anc5, which together suggests that other microbial functions such as other forms of

ion transport and phototaxis may be derived from an ancestral, light-driven proton pump

functionality. This model is compatible with experimental findings that sensory rhodopsins, when

expressed in alternate microbial hosts in the absence of associated transducer proteins, retain

the ability to transport protons, suggesting that the photosensory function is more derived[49].

The early evolution of proton transport by rhodopsins or rhodopsin-like proteins would likely have

provided an adaptive, simplistic means of maintaining the intracellular environment of primitive

cells. Life on early Earth likely faced several hurdles related to membrane permeability that

implicate the presence of relatively simple, membrane-spanning motive proteins proximal to

LUCA[58-60]. One of these hurdles was circumventing the disruptive cellular tendency to expand

and lyse due to the influx of salts from their environment passing across their semi-permeable

protocell membrane[61]. The simplistic membrane-spanning ion-pump infrastructure of an early

rhodopsin could have provided cells inhabiting photic environments with the means to counteract

diffusive flux with a mechanism of active transport[62,63]. Biological preference for transporting

cations in lieu of anions is consistent with the proposed acidic conditions of the early Archaean

ocean, where a high concentration of chloride-based salts such as H^+, Na^+, K^+, Ca^+, and Mg^+

would have been prevalent[62,64]. Acquiring a means of agency over intercellular ion gradient

dynamics would have permitted the maintenance of transmembrane potential and chemiosmotic

coupling for energy generation, which may have been present in LUCA[14,65,66]. From a bioenergetic

perspective, our results support the premise that a rhodopsin-like proton pump could have

enabled chemiosmotic coupling and simplistic energy production in primordial cells.

The type I rhodopsin ancestor was likely tuned to absorb green light

Extant rhodopsin variants exhibit absorption capacity across much of the visible spectrum, and it

is unclear which portion of the spectrum the earliest rhodopsins would have attenuated, nor

whether changes in absorption correlated with the diversification of new functions. To understand

the evolution of rhodopsin spectral tuning, we used the recently developed machine-learning

algorithm, GrplassoSeq[28]. This method approximates the maximum absorbance wavelength

(λ_{max}) of rhodopsin protein sequences by producing a statistical model that describes the

relationship between amino acids within the seven transmembrane helices and absorbance

features of experimentally characterized homologs. The model was demonstrated to predict λ_{max}

of rhodopsins with an average error of ±7.8 nm. Here, λ_{max} values were predicted for extant

sequences in the phylogenetic dataset, inferred maximum likelihood ancestors, and alternative

ancestors for nodes Anc1-An5 generated to test the robustness of wavelength prediction to the

statistical uncertainty of ancestral sequence inference (see Methods; Fig. 2b,c)

The first-order spectral patterns predicted for ancestral rhodopsins indicate lineage-, and,

potentially function-specific, color tuning. λ_{max} for all ancestors range between ~480 to 585 nm,

occupying the cyan-to-yellow range of the visible light spectrum (Fig. 1). Different rhodopsin

lineages, in addition to being associated with varying functions, also exhibit characteristic λ_{max}

values. For example, our results suggest that halorhodopsins evolved to absorb yellow light

(~565-585 nm) early, with secondary shifts toward green light absorption in a specific sublineage.

The bacteriorhodopsin, sensory rhodopsin I, and xanthorhodopsin clades largely exhibit green

spectral tuning (~500-565 nm), whereas shifts towards cyan (~480-500 nm) absorption can be

observed within the sensory rhodopsin II and proteorhodopsin clades. Notable are the so-called blue-light and green-light absorbing members of the proteorhodopsin clade, whose differential spectral tuning has been suggested to correlate with habitat depth of host microbes in the marine environment[67]. Because green light is attenuated at shallower depths, it has been argued that blue-absorbing rhodopsins are hosted by organisms that occupy comparatively deeper portions of the water column. Our λ_{max} predictions suggest that blue-light absorbing proteorhodopsins are derived from a green-light absorbing ancestor which would thus have been better adapted to a shallow marine environment.

All oldest, maximum likelihood ancestral sequences (Anc1-Anc5) from our rhodopsin phylogeny have predicted λ_{max} values between 537 and 543 nm (Fig. 1). Anc5 is the deepest ancestral node, having both the greatest ancestral sequence uncertainty and, by extension, the greatest variation (~20 nm range) in predicted λ_{max} values across alternate ancestral reconstructions. Even accounting for alternate ancestral sequence reconstructions at these same nodes, the predicted λ_{max} range for early ancestors is only expanded to ~525 to 545 nm. This range is still well within the green portion of the visible light spectrum, even if additionally accounting for the ±7.8 nm average error associated with the wavelength prediction algorithm[28]. Together, our spectral tuning analyses support a model wherein extant type I rhodopsins diversified from an ancestor tuned toward the absorption of green light.

Indicators of early rhodopsin host niche habitability and coevolution with Earth environment

Correlations between tuning of photon absorption and cellular function across modern variants might present a logical basis to infer the most likely niche that hosts of primordial, microbial rhodopsins would have occupied. Paleoecological and paleoenvironmental inferences based on organism-level associations and anatomical attributes are well-established in paleobiology, particularly for ecosystems that existed billions of years ago and for which sedimentological indicators are poor, absent or contested[68-70]. Inferences of 'phylogenetic paleoecology' based on molecule-level attributes constitute a relatively new approach to these problems[25,27].

Color tuning over microbial rhodopsin evolution has likely been shaped by a complex interplay of both functional and environmental constraints[15,22,46,71]. Regarding the former, it is not unexpected that both spectral absorbance patterns and protein function are evolutionarily correlated, given

the observation that many of the same rhodopsin binding pocket sites strongly determine both

features[22,28,71]. Our analysis likewise indicates that λ_{max} values correlate with cellular function

throughout rhodopsin's history. The oldest common ancestors of each clade possess comparable

λ_{max} values to modern counterparts, which themselves appear generally binned according to

functionally distinct lineages (Fig. 1).

Spectral tuning may be further segregated by the photon availability of the host environment and

by selective pressures to enable simultaneous optical control of distinct functions by different

colors of light. Host organisms containing rhodopsin variants are found almost everywhere near

Earth's surface, occupying a variety of different niches. For example, chloride-pumping

halorhodopsins are mostly limited to halophilic Archaea[72] and maximally absorb yellow light,

matching the spectral insolation peak experienced in the terrestrial salt ponds in which they

inhabit. By contrast, sensory rhodopsin II and proteorhodopsin absorbances hosted in shallow-

water photic zones are shifted towards blue light, where peak spectral irradiance is attenuated by

water transmissivity. Sensory rhodopsin I (SRI) tuning is unique because it appears to have been

shaped both by its dual roles in phototactic and photophobic responses, as well as by the

absorption profiles of other rhodopsin proteins in the same host. In a host halobacterium*,* orange

light interaction with SRI produces electrons that drive flagella motion toward light, but absorption

of extra photons in the UV triggers another cascade that stops flagella motion to avoid

phototoxicity[73,74], thereby guiding the host to wavelengths where both halorhodopsin and

bacteriorhodopsin absorb maximally[21].

Inferring the environmental constraints on early rhodopsin evolution is complicated by the fact

that Earth's atmospheric evolution has drastically altered the surface photon irradiance profile

over billions of years. Thus, we reconstructed photon irradiance profiles at various water depths

for both modern and haze-free Archean Earth scenarios, informed by an updated prescription for

the composition of Earth's early atmosphere with high carbon dioxide and methane, but no

oxygen[75] (Fig. 5). These models coupled an advanced atmospheric radiative transfer model

SMART[76] and the absorption profile of liquid water[77]. We found that the level of shielding enjoyed

at the surface of Earth today in a haze-free Archean atmosphere is only possible at depths of tens

of meters or deeper, and that the equivalent protective depth level is a strong function of

wavelength. At visible wavelengths, red and infrared photons also become strongly attenuated

with depth, leading to a wavelength of maximum transparency at ~485 nm in the blue-green

portion of the visible spectrum (assuming no impurities, particulate matter will preferentially scatter

blue light over green light). This contrasts markedly with a peak spectral irradiance at Earth's surface without water attenuation, which ranges from roughly 500-530 nm with a corresponding color of green-yellow.

In addition to reconstructing historical changes in Earth surface irradiation, we explored the impact that the GOE[29], one of the most significant environmental transitions in Earth-life coevolution, may have had in the evolutionary diversification of microbial type I rhodopsins. Light and oxygen both destroy the rhodopsin-associated retinal easily[78], so the various functions of rhodopsins as sensors of these external parameters are intimately tied to sequence changes throughout the phylogenetic family. The last common ancestor of haloarchaea is thought to have possessed the halorhodopsin (and perhaps a bacteriorhodopsin) gene[48]. As haloarchaea are oxygen-consuming heterotrophs, they would likely have originated only after the rise of oxygenic photosynthesis[24], constraining the maximum age of the halorhodopsin clade to at least the GOE at ~2.3 Ga, and perhaps as far back as the first indicators of oxygenation around 3.0 Ga[79,80]. In addition, sensory rhodopsin II further drives a photophobic response under conditions of abundant oxygen to avoid photooxidative damage, a function that likely evolved after the emergence of a well oxygenated post-GOE environment. These observations would indicate that early rhodopsins (i.e., Anc1-Anc5) reflect ancestral functions carried out within anoxic hosts of great (likely early Precambrian) age, regardless of the placement of the root of the rhodopsin phylogenetic network or the exact timing of primordial rhodopsin origination.

Together, our results indicate that ancestral rhodopsins transported protons across the cellular membrane and exhibited spectral tuning toward green photons over a span of time encompassing much of Earth's early history of inhabitation. An assessment of the pervasiveness of light-driven sodium/proton export pumps among prokaryotes has been interpreted as indicating proximity of rhodopsins to the shared bacterial-archaeal superfamily ancestor[14]. This would place the deepest proton-pumping ancestral nodes of our phylogenetic tree proximal to LUCA, with inferred ages of greater than ~3.5 Ga. Such early, green-absorbing rhodopsins would have been well adapted to the water transmission peak at blue-green wavelengths in the marine early Earth surface environment, even at moderate depths in the ocean such that their host organisms could escape damaging UV rays prior to the accumulation of protective atmospheric ozone (Fig. 5a,b, Fig. 6). These spectral regions notably occupy the window excluded by the absorption peaks of later-evolved chlorophylls and bacteriochlorophylls (Fig. 5d, Fig. 6b). Rhodopsin terrestrial and surface water host niches following the GOE were later expanded by reduced UV irradiance, but shallow

water niches were simultaneously more sharply discriminated by the increase in dissolved oxygen
concentrations, contributing to the extant pattern of functional and sequence divergence of the
rhodopsin protein family. This is consistent with the presumed antiquity of the rhodopsin enzyme,
and broadly compatible with trends in mid-to-late Precambrian microfossil abundance, which
(questions of taphonomic bias aside) consistently indicate significant microfossil diversity and
abundance in near-shore marine deposits[81,82]. Host microorganisms residing in these photic
zones would have been able to access available nutrients (e.g., phosphorous and nitrate), and
avoid harmful UV radiation while simultaneously harnessing beneficial regions of the visible
electromagnetic spectrum via the selective attenuation of light (and selective avoidance of
dissolved oxygen) within the water column.
373
374   **Implications for planetary habitability and the search for life beyond Earth**
375
The coevolution of environment and life early in Earth's history serves as a model for predicting
universal, detectable biosignatures that might be generated on a microbe-dominated planet
beyond our solar system. The current approach to constraining the possible suite of remotely
detectable biosignatures leverages the aggregate spectral features of photosystems and pigment
proteins which are sensitive to different portions of the Sun's emission spectrum (Fig. 5)[24].
Photosystems, however, are complicated, specific structures with a relatively limited capacity for
functional variability or spectral tunability. The dual roles of active site residues as determinants
of both function and peak photon absorptivity demonstrate that opsin pigment proteins generate
a suite of biosignatures that are tunable to our Sun's peak irradiance profile, to long-term changes
in the Earth surface chemical system, and to ecological niche diversification over billions of years.
Our phylogenetic analysis indicates that rhodopsin's structural simplicity, functional variability and
spectral tunability make it an ideal testbed for assessing universal properties of biosignature
production on candidate microbe-dominated planets that may differ from our referent biological
(Earth-Sun) system.
390
A generic characteristic of life is that it is 'tuned' to the conditions found on our planet that enable
its existence. These conditions refer to the chemical and physical characteristics that circumscribe
habitability, namely substrate availability, ambient temperature, pH, atmospheric pressure and
composition, salt concentration, and solar insolation, and irradiance, among other geochemical
and atmospheric attributes. As demonstrated here, enzymes can be far more tuned to these
external conditions than other molecules or anatomical features that compose organisms,

precisely because many enzymes serve as intermediaries for channelizing photons and substrates into metabolic activity. Our results show that enzyme families (and to some extent, the niches occupied by their hosts) can be amenable to paleobiological reconstruction in ways that cannot be accomplished using conventional molecular biosignatures alone.

Utilizing enzymes as paleosensors of geologic and stellar conditions in deep time, leveraging the informatic characteristics of enzyme composition, and the tight coupling between internal (cellular) and external (environmental) conditions, permits future studies exploring circumstances of habitability and biosignature production over Earth's deep and varied history. Light absorbance profiles produced by microbial pigments can be used to prime future disk-integrated observations of exoplanets, provide surficial albedo parameters for complex atmospheric radiative transfer models, and directly aid the development of instrumentation for the detection and characterization of life in the universe both within and outside our Solar System. Forthcoming integrative studies that reconcile experimentally validated biological photon absorbances with more sophisticated planetary system analyses would thus be beneficial, for instance, in applying microbial spectral properties to assess the potential habitability or inhabitation of extrasolar planets[83-85]. Overall, understanding biomolecular adaptations may benefit planetary habitability assessments when planetary and stellar indicators are poorly constrained.

METHODS

Phylogenetic reconstruction and ancestral sequence inference

A type I microbial rhodopsin protein sequence dataset was created by identifying homologs from the National Center for Biotechnology Information non-redundant protein database in December 2020. Homologous sequences were identified and obtained via BLASTp[32] using query amino acid sequences acquired from UniProt[86] (Supplementary Table 5) and an expect value cutoff of <1e-5. The sequence dataset was curated to remove duplicate and partial sequences. The final dataset consisted of 568 sequences: 96 bacteriorhodopsin, 144 halorhodopsin, 83 proteorhodopsin, 29 sensory rhodopsin I, 56 sensory rhodopsin II, 44 xanthorhodopsin, and 116 outgroup heliorhodopsin sequences. The dataset was aligned by MAFFT (v7.427)[87] with a gap open penalty of 1.53 and offset value of 0.0. Fungal and algal type I rhodopsin sequences were excluded from the dataset due to long-branch-attraction effects observed within the bacterial rhodopsin clade.

The rhodopsin protein sequence alignment was used to select the best-fit evolutionary model by
ModelFinder[88] implemented in the W-IQ-TREE web server[89]. Models accounting for the proportion
of invariant sites ("+I") were not included in model selection due to issues optimizing rate
heterogeneity parameters independently[90]. The three top-ranking models scored by the Akaike
and Bayesian Information Criterion, LG+G+F, WAG+G+F, and VT+G+F, were used for maximum
likelihood phylogenetic inference by RAxML (v8)[91]. Branch support was evaluated using 100 rapid
bootstrap replicates. The type I rhodopsin clade was rooted with heliorhodopsin sequences, a
protein group known to share a distant homology with type I rhodopsins[33,92]. Because all tested
evolutionary models yielded comparable topologies, the best-fit model, LG+G+F, was
implemented for ancestral sequence inference by PAML (v4)[93]. Ancestral insertion/deletion sites
were reconstructed using a binary likelihood model following Aadland et al.[94].

Homology modeling and functional predictions of ancestral rhodopsin sequences

Inferred maximum likelihood ancestral sequences for the five deepest phylogenetic nodes (Anc1-
Anc5) were submitted to the Phyre2 web server[51] for structural homology modeling. Selected
structural templates for modeling and associated statistics are listed in Supplementary Table 6.
Predicted ancestral rhodopsin structures were used for structure-based functional annotation by
COFACTOR implemented in the I-TASSER software suite[52], which predicts protein function
based on global and local similarities to template proteins with known structures and functions.

Alternative ancestral sequence library generation

For ancestral nodes Anc1-Anc5, 50 alternate protein sequences were generated from site-wise
posterior probabilities to test the robustness of downstream analyses to statistical uncertainty of
ancestral inference. For each ancestral node, regularly increasing percentages (>0% to 100%) of
ambiguously reconstructed sites in the maximum likelihood sequence—defined here as having
posterior probabilities <0.7—were replaced with the next most probable residue. Thus, the least
probable ancestral sequence explored here by this approach had all ambiguous residues
replaced by an alternate residue, likely representing the fringe of plausible ancestral sequence
space[95].

Wavelength Prediction

Ancestral maximum absorbance wavelength (λ_{max}) values were predicted using the machine
learning model, GrplassoSeq[28] implemented in R. This statistical model describes the relationship
between primary amino acid sequence and rhodopsin color tuning based on a training rhodopsin
sequence dataset with experimentally validated λ_{max} values. Maximum likelihood and alternate
ancestral rhodopsin sequences, as well as all extant rhodopsin sequences used for phylogenetic
inference, were aligned to the training dataset for λ_{max} prediction.
471
472 **Radiative transfer and pigment spectra**
473
Surface photon irradiance spectra were calculated with the Spectral Mapping Atmospheric
Radiative Transfer (SMART) model developed by D. Crisp[76]. Calculation assumed cloud-free
conditions with a solar zenith angle of 60 degrees and summed both direct and diffuse radiation
at the surface. The modern Earth atmospheric profile was assumed to be a 1962 U.S. Standard
Atmosphere employed by the InterComparison of Radiation Codes in Climate Models (ICRCCM)
project[96], whereas the Archean scenario was assumed to be a haze-free anoxic atmosphere from
Arney et al.[75] with a CH_4/CO_2 ratio of 0.1. The absorption coefficients of Segelstein[77] and the Beer-
Lambert Law were used to calculate the transmissivity and the attenuation of the photon
irradiances as a function of depth.
483
484 **Data availability**
485
486 All code and phylogenetic datasets can be found at https://github.com/kacarlab/Rhodopsin.
487

488 **ACKNOWLEDGEMENTS**
489
We thank I. Takeuchi, M. Karasuyama, and K. Inoue for access to the machine learning algorithms
used to predict rhodopsin wavelength absorbances used in this work and for the supportive
assistance, and S. DasSarma for the insightful conversations. This work was supported by the
NASA Arizona Space Grant (C.D.S.), the NASA Astrobiology Postdoctoral Fellowship, (A.K.G.),
the NASA Interdisciplinary Astrobiology Research Consortium (ICAR) MUSE, a member of the
NASA Early Cell to Multicellularity Research Coordination Network funded via the Astrobiology
Research Program No 80NSSC21K0592 (B.K.), the NASA Early Career Faculty Award No.
80NSSC19K1617 (B.K.) and an NSF Emerging Frontiers Program Award No. 1724090 (B.K.),
NASA's Alternative Earths team funded by the NASA Astrobiology Institute under Cooperative

Agreement Number NNA15BB03A and the NASA ICAR program, and the Virtual Planetary
Laboratory, a member of the NASA Nexus for Exoplanet System Science (NExSS) funded via
NASA Astrobiology Program Grant No. 80NSSC18K0829 (E.W.S.).

AUTHOR CONTRIBUTIONS

B.K. conceived, designed, and supervised the overall study. C.D.S. designed and performed the
phylogenetic and modeling experiments, collected, and analyzed the data. E.F. generated the
ancient protein library and wavelength prediction pipeline with input from A.K.G. E.W.S.
conducted the radiative transfer calculations. All authors discussed and interpreted the results.
B.K. and C.D.S wrote the manuscript with input and/or text contributions from all authors.

COMPETING INTERESTS STATEMENT

The authors declare no competing interests.

REFERENCES

McKay, C. P. Requirements and limits for life in the context of exoplanets. *Proceedings of the National Academy of Sciences* **111**, 12628-12633, doi:https://doi.org/10.1073/pnas.1304212111 (2014).

Harrison, J. P., Gheeraert, N., Tsigelnitskiy, D., Cockell, C.S. The limits for life under multiple extremes. *Trends in microbiology* **21**, 204-212, doi:https://doi.org/10.1016/j.tim.2013.01.006 (2013).

Lunine, J. I. Physical conditions on the early Earth. *Philosophical Transactions of the Royal Society B* **361**, 1721-1731, doi:https://doi.org/10.1098/rstb.2006.1900 (2006).

Newman, M. J., Rood, R.T. Implications of Solar Evolution for the Earth's Early Atmosphere. *Science* **198**, 1035-1037, doi:https://doi.org/10.1126/science.198.4321.1035 (1977).

Garcia, A. K., Cavanaugh, C. M. & Kacar, B. The curious consistency of carbon biosignatures over billions of years of Earth-life coevolution. *ISME J*, doi:10.1038/s41396-021-00971-5 (2021).

Butterfield, N. J. Macroevolution and Macroecology through Deep Time. *Palaeontology* **50**, 41-55, doi:10.1111/j.1475-4983.2006.00613.x (2007).

Doolittle, W. F. The nature of the universal ancestor and the evolution of the proteome. *Current opinion in structural biology* **10**, 355–358, doi:https://doi.org/10.1016/s0959-440x(00)00096-8 (2000).

Benner, S. A., Caraco, M.D., Thomson, J.M., Gaucher, E.A. Planetary biology—paleontological, geological, and molecular histories of life. *Science* **296**, 864-868, doi:https://doi.org/10.1126/science.1069863 (2002).

Garcia, A. K., Kaçar, B. How to resurrect ancestral proteins as proxies for ancient biogeochemistry. *Free Radical Biology and Medicine* **140**, 260-269, doi:https://doi.org/10.1016/j.freeradbiomed.2019.03.033 (2019).

Gómez-Conarnau, L., Raven, J.A., Levine, N.M., Cutter, L.S., Wang, D., Seegers, B., Arístegui, J., Fuhrman, J.A., Gasol, J.M., Sañudo-Wilhelmy, S.A. . Microbial rhodopsins are major contributors to the solar energy captured in the sea. *Science Advances* **5**, eaaw8855, doi:https://doi.org/10.1126/sciadv.aaw8855 (2019).

Pinhassi, J., DeLong, E.F., Béjà, O., González, J.M., Pedrós-Alió, C. Marine Bacterial and Archaeal Ion-Pumping Rhodopsins: Genetic Diversity, Physiology, and Ecology. *Microbiology and Molecular Biology Reviews* **80**, 929–954, doi:https://doi.org/10.1128/MMBR.00003-16 (2016).

Bryant, D. A. & Frigaard, N.-U. Prokaryotic photosynthesis and phototrophy illuminated. *Trends in Microbiology* **14**, 488-496, doi:10.1016/j.tim.2006.09.001 (2006).

Sharma, A. K., Spudich, J. L., Doolittle, W.F. Microbial rhodopsins: functional versatility and genetic mobility. *Trends in microbiology* **14**, 463–469, doi:https://doi.org/10.1016/j.tim.2006.09.006 (2006).

Shalaeva, D. N., Galperin, M.Y., Mulkidjanian, A.Y. Eukaryotic G protein-coupled receptors as descendants of prokaryotic sodium-translocating rhodopsins. *Biology Direct* **10**, 1-12, doi:https://doi.org/10.1186/s13062-015-0091-4 (2015).

DasSarma, S. & Schwieterman, E. W. Early evolution of purple retinal pigments on Earth and implications for exoplanet biosignatures. *International Journal of Astrobiology*, 1-10, doi:10.1017/s1473550418000423 (2018).

Mackin, K. A., Roy, R.A., Theobald, D.L. An empirical test of convergent evolution in rhodopsins. *Molecular biology and evolution* **31**, 85–95, doi: https://doi.org/10.1093/molbev/mst171 (2014).

17 Racker, E., Stoeckenius, W. Reconstitution of purple membrane vesicles catalyzing light-
driven proton uptake and adenosine triphosphate formation. *The Journal of Biological*
Chemistry **249**, 662–663, doi:https://doi.org/10.1016/S0021-9258(19)43080-9 (1974).
18 Akram, N., Palovaara, J., Forsberg, J., Lindh, M.V., Milton, D.L., Luo, H., González, J.M.,
Pinhassi, J. Regulation of proteorhodopsin gene expression by nutrient limitation in the
marine bacterium Vibrio sp. AND4. *Environmental microbiology* **15**, 1400–1415,
doi:https://doi.org/10.1111/1462-2920.12085 (2013).
19 Govorunova, E., Sineshchekov, O.A., Li, H., Spudich, J.L. Microbial Rhodopsins:
Diversity, Mechanisms, and Optogenetic Applications. *Annual Review of Biochemistry* **86**,
845-872, doi: https://doi.org/10.1146/annurev-biochem-101910-144233 (2017).
20 Steindler, L., Schwalbach, M.S., Smith, D.P., Chan, F., Giovannoni, S.J. . Energy starved
Candidatus Pelagibacter ubique substitutes light-mediated ATP production for
endogenous carbon respiration. *PloS one* **6**, e19725,
doi:https://doi.org/10.1371/journal.pone.0019725 (2011).
21 Sharma, A. K. *Insights into the ecology and evolution of the light-harvesting microbial*
rhodopsins, Dalhousie University, (2009).
22 Engqvist, M. K. M. *et al.* Directed Evolution of Gloeobacter violaceus Rhodopsin Spectral
Properties. *Journal of Molecular Biology* **427**, 205-220,
doi:https://doi.org/10.1016/j.jmb.2014.06.015 (2015).
23 Inoue, K. *et al.* Spectroscopic Study of Proton-Transfer Mechanism of Inward Proton-
Pump Rhodopsin, Parvularcula oceani Xenorhodopsin. *The Journal of Physical Chemistry*
B **122**, 6453-6461, doi:https://doi.org/10.1021/acs.jpcb.8b01279 (2018).
24 Schwieterman, E. W. *et al.* Exoplanet Biosignatures: A Review of Remotely Detectable
Signs of Life. *Astrobiology* **18**, 663-708, doi:https://doi.org/10.1089/ast.2017.1729 (2018).
25 Lamsdell, J. C., Congreve, C. R., Hopkins, M. J., Krug, A. Z. & Patzkowsky, M. E.
Phylogenetic Paleoecology: Tree-Thinking and Ecology in Deep Time. *Trends in Ecology*
& Evolution **32**, 452-463, doi:10.1016/j.tree.2017.03.002 (2017).
26 Finkel, O. M., Béjà, O. & Belkin, S. Global abundance of microbial rhodopsins. *The ISME*
Journal **7**, 448-451, doi:10.1038/ismej.2012.112 (2012).
27 Kacar, B., Guy, L., Smith, E. & Baross, J. Resurrecting ancestral genes in bacteria to
interpret ancient biosignatures. *Philos Trans R Soc A* **375**, doi:10.1098/rsta.2016.0352
(2017).
28 Karasuyama, M., Inoue, K., Nakamura, R., Kandori, H. & Takeuchi, I. Understanding
Colour Tuning Rules and Predicting Absorption Wavelengths of Microbial Rhodopsins by
Data-Driven Machine-Learning Approach. *Sci Rep* **8**, 15580, doi:10.1038/s41598-018-
33984-w (2018).
29 Lyons, T. W., Reinhard, C. T. & Planavsky, N. J. The rise of oxygen in Earth's early ocean
and atmosphere. *Nature* **506**, 307-315, doi:10.1038/nature13068 (2014).
30 Sánchez-Baracaldo, P., Bianchini, G., Wilson, J. D. & Knoll, A. H. Cyanobacteria and
biogeochemical cycles through Earth history. *Trends in Microbiology*,
doi:10.1016/j.tim.2021.05.008 (2021).
31 Hazen, R. M. *et al.* Mercury (Hg) mineral evolution: A mineralogical record of
supercontinent assembly, changing ocean geochemistry, and the emerging terrestrial
biosphere. *American Mineralogist* **97**, 1013-1042, doi:10.2138/am.2012.3922 (2012).
32 Camacho, C. *et al.* BLAST+: architecture and applications. *BMC Bioinformatics* **10**, 421,
doi:10.1186/1471-2105-10-421 (2009).
33 Pushkarev, A., Inoue, K., Larom, S., Flores-Uribe, J., Singh, M., Konno, M., Tomida, S.,
Ito, S., Nakamura, R., Tsunoda, S. P., Philosof, A., Sharon, I., Yutin, N., Koonin, E. V.,
Kandori, H., Béjà, O. . A distinct abundant group of microbial rhodopsins discovered using
functional metagenomics. *Nature* **558**, 595–599, doi:https://doi.org/10.1038/s41586-018-
0225-9 (2018).

34 Lu, Y. *et al.* Crystal structure of heliorhodopsin 48C12. *Cell Research* **30**, 88-90,
doi:10.1038/s41422-019-0266-0 (2019).
35 Rozenberg, A., Inoue, K., Kandori, H. & Béjà, O. Microbial Rhodopsins: The Last Two
Decades. *Annual Review of Microbiology* **75**, 427-447, doi:10.1146/annurev-micro-
031721-020452 (2021).
36 Ernst, O. P., Lodowski, D.T., Elstner, M., Hegemann, P., Brown, L.S., Kandori, H.
Microbial and Animal Rhodopsins: Structures, Functions, and Molecular Mechanisms.
Chemical Reviews **114**, 126–163, doi:https://doi.org/10.1021/cr4003769 (2014).
37 Ihara, K., Umemura, T., Katagiri, I., Kitajima-Ihara, T., Sugiyama, Y., Kimura, Y., &
Mukohata, Y. Evolution of the archaeal rhodopsins: evolution rate changes by gene
duplication and functional differentiation. *Journal of molecular biology* **285**, 163–174,
doi:https://doi.org/10.1006/jmbi.1998.2286 (1999).
38 Shibata, M., Inoue, K., Ikeda, K., Konno, M., Singh, M., Kataoka, C., Abe-Yoshizumi, R.,
Kandori, H., & Uchihashi, T. . Oligomeric states of microbial rhodopsins determined by
high-speed atomic force microscopy and circular dichroic spectroscopy. *Scientific reports*
8, 8262, doi:https://doi.org/10.1038/s41598-018-26606-y (2018).
39 Kandori, H. Ion-pumping microbial rhodopsins. *Front Mol Biosci* **2**, 52,
doi:10.3389/fmolb.2015.00052 (2015).
40 Babitzki, G., Denschlag, R., Tavan, P. Polarization effects stabilize bacteriorhodopsin's
chromophore
binding pocket: a molecular dynamics study. *The journal of physical chemistry B* **113**, 10483–
10495, doi:https://doi.org/10.1021/jp902428x (2009).
41 Greenhalgh, D. A., Farrens, D.L., Subramaniam, S., Khorana, H.G. Hydrophobic Amino
Acids in the Retinal-binding Pocket of Bacteriorhodopsin. *The Journal of Biological*
Chemistry **268**, 20305-20311, doi:https://doi.org/10.1016/S0021-9258(20)80729-7
(1993).
42 Ding, X., Sun, C., Cui, H., Chen, S., Gao, Y., Yang, Y., Wang, J., He, X., Iuga, D., Tian,
F., Watts, A., Zhao, X. Functional roles of tyrosine 185 during the bacteriorhodopsin
photocycle as revealed by in situ spectroscopic studies. *Biochimica et biophysica acta:*
bioenergetics **1859**, 1006–1014, doi:https://doi.org/10.1016/j.bbabio.2018.05.011 (2018).
43 Kusnetzow, A., Singh, D.L., Martin, C.H., Barani, I.J., Birge, R.R. Nature of the
chromophore binding site of bacteriorhodopsin: the potential role of Arg82 as a principal
counterion. *Biophysical journal* **76**, 2370–2389, doi:https://doi.org/10.1016/S0006-
3495(99)77394-7 (1999).
44 Song, Y. & Gunner, M. R. Halorhodopsin pumps Cl- and bacteriorhodopsin pumps protons
by a common mechanism that uses conserved electrostatic interactions. *Proc Natl Acad*
Sci USA **111**, 16377-16382, doi:10.1073/pnas.1411119111 (2014).
45 Sudo, Y., Furutani, Y., Kandori, H. & Spudich, J. L. Functional importance of the
interhelical hydrogen bond between Thr204 and Tyr174 of sensory rhodopsin II and its
alteration during the signaling process. *J Biol Chem* **281**, 34239-34245,
doi:10.1074/jbc.M605907200 (2006).
46 Man, D. Diversification and spectral tuning in marine proteorhodopsins. *EMBO J* **22**, 1725-
1731, doi:10.1093/emboj/cdg183 (2003).
47 Inoue, K., Karasuyama, M., Nakamura, R., Konno, M., Yamada, D., Mannen, K., Nagata,
T., Inatsu, Y., Yawo, H., Yura, K., Béjà, O., Kandori, H., Takeuchi, I. . Exploration of natural
red-shifted rhodopsins using a machine learning-based Bayesian experimental design.
Communications biology **4**, 362, doi:https://doi.org/10.1038/s42003-021-01878-9 (2021).
48 Sharma, A. K., Walsh, D.A., Bapteste, E., Rodriguez-Valera, F., Ford Doolittle, W., Papke,
R.T. Evolution of rhodopsin ion pumps in haloarchaea. *BMC evolutionary biology* **7**, 1-13,
doi:https://doi.org/10.1186/1471-2148-7-79 (2007).

Bogomolni, R. A. *et al.* Removal of transducer HtrI allows electrogenic proton translocation by sensory rhodopsin I. *Proc Natl Acad Sci USA* **91**, 10188-10192, doi:10.1073/pnas.91.21.10188 (1994).

Spudich, J. L., Sineshchekov, O. A. & Govorunova, E. G. Mechanism divergence in microbial rhodopsins. *Biochimica et Biophysica Acta (BBA) - Bioenergetics* **1837**, 546-552, doi:10.1016/j.bbabio.2013.06.006 (2014).

Kelley, L. A., Mezulis, S., Yates, C. M., Wass, M. N. & Sternberg, M. J. E. The Phyre2 web portal for protein modeling, prediction and analysis. *Nature Protocols* **10**, 845-858, doi:10.1038/nprot.2015.053 (2015).

Roy, A., Kucukural, A., Zhang, Y. I-TASSER: a unified platform for automated protein structure and function prediction. *Nature Protocols* **5**, 725–738, doi:https://doi.org/10.1038/nprot.2010.5 (2010).

Hirokawa, T., Boon-Chieng, S. & Mitaku, S. SOSUI: classification and secondary structure prediction system for membrane proteins. *Bioinformatics* **14**, 378-379, doi:10.1093/bioinformatics/14.4.378 (1998).

Hasegawa, M. *et al.* A unique clade of light-driven proton-pumping rhodopsins evolved in the cyanobacterial lineage. *Sci Rep* **10**, 16752, doi:10.1038/s41598-020-73606-y (2020).

Inoue, K. Shining light on rhodopsin selectivity: How do proteins decide whether to transport H+ or Cl–? *J Biol Chem* **295**, 14805-14806, doi:10.1074/jbc.H120.016032 (2020).

Besaw, J. E. *et al.* The crystal structures of a chloride-pumping microbial rhodopsin and its proton-pumping mutant illuminate proton transfer determinants. *J Biol Chem* **295**, 14793-14804, doi:10.1074/jbc.RA120.014118 (2020).

Sasaki, J. *et al.* Conversion of Bacteriorhodopsin into a Chloride Ion Pump. *Science* **269**, 73-75, doi:10.1126/science.7604281 (1995).

Mulkidjanian, A. Y., Galperin, M. Y. & Koonin, E. V. Co-evolution of primordial membranes and membrane proteins. *Trends Biochem Sci* **34**, 206-215, doi:10.1016/j.tibs.2009.01.005 (2009).

Deamer, D. W. & Dworkin, J. P. in *Prebiotic Chemistry Topics in Current Chemistry* Ch. Chapter 136806, 1-27 (2005).

Pohorille, A., Wilson, M. A. & Chipot, C. *Origins of Life and Evolution of the Biosphere* **33**, 173-197, doi:10.1023/a:1024627726231 (2003).

Pohorille, A., Deamer, D. . Self-assembly and function of primitive cell membranes. *Research in Microbiology* **160**, 449-456, doi:https://doi.org/10.1016/j.resmic.2009.06.004 (2009).

Damer, B., Deamer, D. . The Hot Spring Hypothesis for an Origin of Life. *Astrobiology* **20**, 429-452, doi:https://doi.org/10.1089/ast.2019.2045 (2020).

Maloney, P. C., Wilson, T.H. The Evolution of Ion Pumps. *BioScience* **35**, 43-48, doi:https://doi.org/10.2307/1310083 (1985).

Ueda, H., Shibuya, T. Composition of the primordial ocean just after its formation: Constraints from the reactions between the primitive crust and a strongly acidic, CO2-rich fluid at elevated temperatures and pressures. *Minerals* **11**, doi:https://doi.org/10.3390/min11040389 (2021).

Penny, D., Sojo, V., Pomiankowski, A. & Lane, N. A Bioenergetic Basis for Membrane Divergence in Archaea and Bacteria. *PLoS Biology* **12**, doi:10.1371/journal.pbio.1001926 (2014).

Lane, N., Allen, J. F. & Martin, W. How did LUCA make a living? Chemiosmosis in the origin of life. *Bioessays* **32**, 271-280, doi:10.1002/bies.200900131 (2010).

Ozaki, Y., Kawashima, T., Abe-Yoshizumi, R. & Kandori, H. A color-determining amino acid residue of proteorhodopsin. *Biochemistry* **53**, 6032-6040, doi:10.1021/bi500842w (2014).

Adam, Z. R., Skidmore, M. L., Mogk, D. W. & Butterfield, N. J. A Laurentian record of the earliest fossil eukaryotes. *Geology* **45**, 387-390, doi:10.1130/g38749.1 (2017).

Butterfield, N. J. & Rainbird, R. H. Diverse organic-walled fossils, including "possible dinoflagellates," from the early Neoproterozoic of arctic Canada. *Geology* **26**, doi:10.1130/0091-7613(1998)026<0963:Dowfip>2.3.Co;2 (1998).

Hofmann, H. J. & Jackson, G. D. Shale-Facies Microfossils from the Proterozoic Bylot Supergroup, Baffin Island, Canada. *J Paleontol* **68**, 1-35, doi:10.1017/s0022336000062314 (1994).

Tsujimura, M. & Ishikita, H. Insights into the Protein Functions and Absorption Wavelengths of Microbial Rhodopsins. *The Journal of Physical Chemistry B* **124**, 11819-11826, doi:10.1021/acs.jpcb.0c08910 (2020).

Ugalde, J. A., Podell, S., Narasingarao, P. & Allen, E. E. Xenorhodopsins, an enigmatic new class of microbial rhodopsins horizontally transferred between archaea and bacteria. *Biology Direct* **6**, doi:10.1186/1745-6150-6-52 (2011).

Wilde, A. & Mullineaux, C. W. Light-controlled motility in prokaryotes and the problem of directional light perception. *FEMS Microbiology Reviews* **41**, 900-922, doi:10.1093/femsre/fux045 (2017).

Spudich, J. L. & Bogomolni, R. A. Mechanism of colour discrimination by a bacterial sensory rhodopsin. *Nature* **312**, 509-513, doi:10.1038/312509a0 (1984).

Arney, G., Domagal-Goldman, S.D., Meadows, V.S., Wolf, E.T., Schwieterman, E., Charnay, B., Claire, M., Hébrard, E., Trainer, M.G. The Pale Orange Dot: The Spectrum and Habitability of Hazy Arhean Earth. *Astrobiology* **16**, 873-899, doi:https://doi.org/10.1089/ast.2015.1422 (2016).

Meadows, V. S., Crisp, D. Ground-based near-infrared observations of the Venus nightside: The thermal structure and water abundance near the surface. *Journal of Geophysical Research: Planets* **101**, 4595-4622, doi:https://doi.org/10.1029/95JE03567 (1996).

Segelstein, D. J. The complex refractive index of water. *University of Missouri*, doi:http://hdl.handle.net/10355/11599 (1981).

Hampp, N. & Oesterhelt, D. in *Nanobiotechnology* (eds Christof M. Niemeyer & Chad A. Mirkin) 146-167 (2004).

Cardona, T. Early Archean origin of heterodimeric Photosystem I. *Heliyon* **4**, doi:10.1016/j.heliyon.2018.e00548 (2018).

Olson, S. L., Kump, L. R. & Kasting, J. F. Quantifying the areal extent and dissolved oxygen concentrations of Archean oxygen oases. *Chem Geol* **362**, 35-43, doi:10.1016/j.chemgeo.2013.08.012 (2013).

Adam, Z. R., Skidmore, M.L., Mogk, D.M. Paleoenvironmental implications of an expanded microfossil assemblage from the Chamberlain Formation, ower Belt Supergroup, Montana. *The Geological Society of America Special Paper* **522** (2016).

Javaux, E. J., Knoll, A.H. Micropaleontology of the lower Mesoproterozoic Roper Group, Australia, and implications for early eukaryotic evolution. *Journal of Paleontology* **91**, 199-229, doi:https://doi.org/10.1017/jpa.2016.124 (2017).

DasSarma, S., DasSarma, P., Laye, V.J., Schwieterman, E.W. Extremophilic models for astrobiology: haloarchaeal survival strategies and pigments for remote sensing. *Extremophiles* **24**, 31–41, doi:https://doi.org/10.1007/s00792-019-01126-3 (2020).

Hegde, S., Paulino-Lima, I.G., Kent, R., Kaltenegger, L., Rothschild, L. Surface biosignatures of exo-Earths: Remote detection of extraterrestrial life. *Proceedings of the National Academy of Sciences* **112**, 3886–3891, doi:https://doi.org/10.1073/pnas.1421237112 (2015).

Schwieterman, E. W., Cockell, C.S., Meadows, V.S. Nonphotosynthetic Pigments as Potential Biosignatures. *Astrobiology* **15**, 341–361, doi:https://doi.org/10.1089/ast.2014.1178 (2015).

Uniprot Consortium. UniProt: a worldwide hub of protein knowledge. *Nucleic Acids Research* **47**, D506-D515, doi:10.1093/nar/gky1049 (2019).

Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software version 7: improvements in performance and usability. *Mol Biol Evol* **30**, 772-780, doi:10.1093/molbev/mst010 (2013).

Kalyaanamoorthy, S., Minh, B. Q., Wong, T. K. F., von Haeseler, A. & Jermiin, L. S. ModelFinder: fast model selection for accurate phylogenetic estimates. *Nature Methods* **14**, 587-589, doi:10.1038/nmeth.4285 (2017).

Trifinopoulos, J., Nguyen, L.-T., von Haeseler, A. & Minh, B. Q. W-IQ-TREE: a fast online phylogenetic tool for maximum likelihood analysis. *Nucleic Acids Research* **44**, W232-W235, doi:10.1093/nar/gkw256 (2016).

Yang, Z. *Computational Molecular Evolution*. 374 (Oxford University Press, 2006).

Stamatakis, A. RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **30**, 1312-1313, doi:10.1093/bioinformatics/btu033 (2014).

Shibukawa, A., Kojima, K., Nakajima, Y., Nishimura, Y., Yoshizawa, S., Sudo, Y. . Photochemical Characterization of a New Heliorhodopsin from the Gram-Negative Eubacterium Bellilinea caldifistulae (BcHeR) and Comparison with Heliorhodopsin-48C12. *Biochemistry* **58**, 2934–2943, doi:https://doi.org/10.1021/acs.biochem.9b00257 (2019).

Yang, Z. PAML 4: Phylogenetic Analysis by Maximum Likelihood. *Molecular Biology and Evolution* **24**, 1586–1591, doi:https://doi.org/10.1093/molbev/msm088 (2007).

Aadland, K., Pugh, C. & Kolaczkowski, B. High-Throughput Reconstruction of Ancestral Protein Sequence, Structure, and Molecular Function. *Methods Mol Biol* **1851**, 135-170, doi:10.1007/978-1-4939-8736-8_8 (2019).

Eick, G. N., Bridgham, J. T., Anderson, D. P., Harms, M. J. & Thornton, J. W. Robustness of Reconstructed Ancestral Protein Functions to Statistical Uncertainty. *Mol Biol Evol* **34**, 247-261, doi:10.1093/molbev/msw223 (2017).

Luther, F. M., Ellingson, R.G., Fouquart, Y., Fels, S., Scott, N.A., Wiscombe, W.J. Intercomparison of Radiation Codes in Climate Models (ICRCCM): Longwave Clear-Sky Results—A Workshop Summary. *Bulletin of the American Meteorological Society* **69**, 40-48, doi:https://doi.org/10.1175/1520-0477-69.1.40 (1988).

Chen, M., Schliep, M., Willows, R.D., Cai, Z.-L., Neilan, B.A., Scheer, H. A Red-Shifted Chlorophyll. *Science* **329**, 1318–1319, doi:https://doi.org/10.1126/science.1191127 (2010).

Chen, M., Blankenship, R.E. Expanding the solar spectrum used by photosynthesis. *Trends in Plant Science* **16**, 427–431, doi:https://doi.org/10.1016/j.tplants.2011.03.011 (2011).

Frigaard, N., Larsen, Kim, L., Cox, R.P. Spectrochromatography of photosynthetic pigments as a fingerprinting technique for microbial phototrophs. *FEMS Microbiology Ecology* **20**, 69–77, doi:https://doi.org/10.1016/0168-6496(96)00005-0 (1996).

Jeffrey, S. W. Purification and properties of chlorophyll c from Sargassum flavicans. *Biochemical Journal* **86**, 313-318, doi:https://doi.org/10.1042/bj0860313 (1963).

FIGURES

Fig. 1 Evolution of spectral tuning mapped to a maximum likelihood phylogeny of microbial type I rhodopsin protein sequences. Nodes are colored according to the predicted maximum absorption wavelength (λ_{max}) of the associated maximum likelihood (ML) ancestral sequence (color scale shown in plot, top left). Boxplot shows the predicted λ_{max} distribution among alternate ancestral sequences for nodes Anc1-Anc5 (n = 50 per node; see Methods and Fig. 2b,c). Branch support values are derived from 100 bootstrap replicates. Branch scale indicates 1.0 amino acid substitutions per site. Heliorhodopsin outgroup clade is collapsed. BR: bacteriorhodopsin; HR, halorhodopsin; SR II, sensory rhodopsin II; SR I, sensory rhodopsin I; PR, proteorhodopsin; XR, xanthorhodopsin; HeR, heliorhodopsin.

Fig. 2 Incorporation of statistical uncertainty of ancestral rhodopsins inference into wavelength prediction. a Site-wise posterior probabilities of maximum likelihood (ML) ancestral rhodopsin sequences. Sites labeled "functional motif" are homologous to the three-residue DTD motif in extant *H. salinarum* bacteriorhodopsin. Helices are labeled above. **b** Generation of alternative ("alt") ancestral sequences. For each node, alternate sequences were generated by replacing 0% to 100% of ambiguous sites (defined as having a posterior probability < 0.7) with the next most probable residue. **c** Methodology for maximum absorbance wavelength (λ_{max}) prediction. λ_{max} values of extant and ancestral (both ML and alternative) rhodopsin sequences were predicted by a machine-learning model (GrplassoSeq), trained on a dataset of experimentally characterized rhodopsin proteins with known λ_{max} values.

Fig. 3 Structural conservation between the inferred, type I rhodopsin last common

ancestor, Anc5, and extant bacteriorhodopsin. **a** Homology model of Anc5 aligned to the

extant *H. salinarum* bacteriorhodopsin ("BR"; PDB: 1C3W). **b** Retinal binding pocket of ancestral

and extant rhodopsins. Conserved residues that interact with the retinal chromophore are labeled.

c Residues homologous to the *H. salinarum* DTD motif correlative with rhodopsin function. Extant

bacteriorhodopsin exhibits a DTD motif, compared to the DTE motif in Anc5. **b,c** All displayed

Anc5 residues are identical to those of extant bacteriorhodopsin except for site 96 (D96E, boxed).

Fig. 4 Rhodopsin binding pocket residues in extant and inferred ancestral sequences. a Secondary structure of extant *H. salinarum* bacteriorhodopsin, indicating the binding pocket residues as well as the functionally correlative DTD motif. **b** Protein sequence alignment of Helix C residues for representative, extant type I rhodopsins and ancestral rhodopsins. Binding pocket residues are indicated by bold font. **a,b** Sites labeled "functional motif" are homologous to the three-residue DTD motif in extant *H. salinarum* bacteriorhodopsin.

Fig. 5 Diversity of biological pigment absorbance profiles in the context of changing surface photon irradiance over Earth history. **a,b** Photon irradiance profiles at varying water depths, given a modern (**a**) or Archean (**b**) atmospheric composition. **c** Water transmission profile at varying water depths. Transmission was calculated from the coefficients of Segelstein[77] and the Beer-Lambert Law. **a,b,c** Vertical dashed line indicates the maximum water transmission wavelength of 485 nm. **d** Normalized absorbance for bacteriorhodopsin ("BR"), chlorophylls ("Chl"), and bacteriochlorophylls ("Bchl"). Shaded purple region indicates the extant and ancestral range of predicted rhodopsin absorption peaks in Fig. 1. Chlorophyll and bacteriochlorophyll spectra are sourced from the Virtual Planetary Laboratory's Biological Pigments Database (http://vplapps.astro.washington.edu/pigments), with contributions from refs [97-100]. The bacteiorhodopsin spectrum was sourced from DasSarma and Schwieterman[15].

Fig. 6 Coupling between surface irradiance, spectral tuning, and functional diversity over microbial rhodopsin evolution. a Microbial hosts of ancestral type I rhodopsins would likely have occupied the marine environment where high Archean UV irradiation was attenuated and where they could absorb green wavelengths to which they were maximally tuned. These ancestral rhodopsins would likely have functioned as light-driven proton pumps, providing both a simplistic means of energy production and maintenance of the intracellular environment. **b** Color tuning of microbial rhodopsins explored the spectral window not already occupied by other biological pigments such as chlorophylls and bacteriochlorophylls (grey, adapted from Fig. 5d). Spectral diversification would have accompanied functional specialization, which itself was likely responsive to expanding ecological niches driven by the rise of atmospheric oxygen.