scCODE: an R package for personalized differentially expressed gene detection on single-cell RNA-sequencing data

Authors: Jiawei Zou[1,2], Miaochen Wang[4], Zhen Zhang[4], Zheqi Liu[4], Xiaobin Zhang[6,7], Rong Hua[6], Ke Chen[5]*, Xin Zou[3]*, Jie Hao[2]*.

Affiliations:

[1] School of Life Sciences and Biotechnology, Shanghai Centre for Systems Biomedicine, Shanghai Jiao Tong University, Shanghai, China.

[2] Institute of Clinical Science, Zhongshan Hospital, Fudan University, Shanghai, China.

[3] Jinshan Hospital Center for Tumor Diagnosis & Therapy, Jinshan Hospital, Fudan University, Shanghai, 201508, China.

[4] Department of Oral and Maxillofacial-Head & Neck Oncology, Shanghai Ninth People's Hospital, Shanghai Jiao Tong University School of Medicine; College of Stomatology, Shanghai Jiao Tong University; National Center for Stomatology; National Clinical Research Center for Oral Diseases; Shanghai Key Laboratory of Stomatology.

[5] Shanghai Key Laboratory of Plant Functional Genomics and Resources, Shanghai Chenshan Botanical Garden, Shanghai, 201602, China

[6] Department of Thoracic Surgery, Shanghai Chest Hospital, Shanghai Jiao Tong University, Shanghai, China.

[7] Department of Cardiovascular Surgery, Shanghai Chest Hospital, Shanghai Jiao Tong University, Shanghai, China.

* Corresponding authors.

Email: Jie Hao: hao.jie@zs-hospital.sh.cn; Jiawei Zou: jiawZou@163.com; Xin Zou: x.zou@hotmail.co.uk; Ke Chen: chenke@csnbgsh.cn; Miaochen Wang: miaochenW@126.com; Zhen Zhang: 326197290@qq.com; Zheqi Liu: 734672500@sjtu.edu.cn; Xiaobin Zhang: zxb5212@163.com; Rong Hua: huagang9205@163.com.

Abstract

Differential expression (DE) gene detection in single-cell RNA-seq (scRNA-seq) data is a key step to understand the biological question investigated. We find that DE methods together with gene filtering have profound impact on DE gene identification, and different datasets will benefit from personalized DE gene detection strategies. Existing tools don't take gene filtering into consideration, and couldn't evaluate DE performance on real datasets without prior knowledge of true results. Based on two new metrics, we propose scCODE (single cell Consensus Optimization of Differentially Expressed gene detection), an R package to automatically optimize DE gene detection for each experimental scRNA-seq dataset.

Keywords: bioinformatics, scRNA-seq data, differentially expressed gene detection.

Background

In the past few years, stupendous development in single-cell RNA sequencing (scRNA-seq) has made it a powerful technology in transcriptomic biology. scRNA-seq data can provide unprecedented transcriptomic profile of individual cells. For the interpretation of scRNA-seq dataset where comparison scenarios occur, for example, cell development[1], cancer research[2, 3] and immune research[4, 5], etc., the detection of DE gene is critical for downstream researches. Accurate and reliable DE gene detection can accelerate the discovery of novel regulator genes and biomarkers[6, 7].

In order to detect the true DE genes in scRNA-seq data, various DE methods and algorithms have been developed, including methods particularly designed for scRNA-seq data, such as MAST[8], BPSC[9] and scDD[10], methods previously used in bulk data, such as edgeR[11], DESeq2[12] and the more common methods such as t-test[13] and Wilcoxon-test[13] et al. And there also exists method[14] integrating these methods.

Although there exist several tools to assess the performance of DE methods on cohort of scRNA-seq data[15, 16], they only evaluate each method based on the average performance on many scRNA-seq datasets. However, the method selected this way may not always be suitable for every dataset, there's still no solution to optimize DE gene detection performance for a specific

50   experimental scRNA-seq dataset. In addition, there is lack of systematic evaluation on gene

filtering methods, regardless its profound influence in scRNA-seq data interpretation[17, 18].

Here, we first systematically evaluated the performance of combinations of different gene filtering

and DE methods based on the metrics used in conquer[16], we exhibited the true influence of gene

filtering and that different analysis strategy varies a lot in DE gene detection results. Besides, we

55   noticed that each dataset requires specific method to achieve its optimal result, but there is lack of

means to determine the most suitable method for a given dataset without prior knowledge of the

true DE genes. Thus, we constructed two novel metrics, CDO (Consistent DE genes Order) and

AUCC (Area under consistency curve). Principally, the two metrics rate the performance of DE

gene detection strategies by evaluating their efficiency of capturing common genes. The reliability

60   of the two metrics was justified on simulated datasets with known true DE genes, and both metrics

showed significant correlation with the AUROC (Area Under Receiver Operating Characteristic

Curve) which was regarded as the ground truth. Based on the new metrics, we constructed

scCODE (single cell Consensus Optimization of Differentially Expressed gene detection), an R

package to provide optimized DE gene results based on the consensus genes obtained by the top

65   suitable analysis strategies. We found that scCODE demonstrated better and more robust

performance, while other analysis strategies showed unstable performance on different simulated

datasets. The scCODE was further justified on a mouse scRNA-seq data[19] comparing activated

CD4+ T cells with naïve CD4+ T cells. The obtained biological processes associated with T cell activation highly agreed with the experimental design of the origin study[19].

Methods

In this article, we focused on the DE gene detection of two-group comparison. First, systematic evaluation of various analysis strategies by combining gene filtering (conquer[16], OGFSC (Optimal Gene filtering for Single-cell data)[18], scmap[20] and no-filter) and DE methods was performed based on the following metrics, i.e., FPR (False positive Rate), SNR (Signal-Noise Ratio), FDP (False discovery Proportion), TPR (True Positive Rate) and AUROC (Figure. 1a). However, the above metrics are not applicable in real scenarios as they all require prior knowledge of true DE genes. To tackle this issue, two new metrics CDO and AUCC were constructed for evaluating DE gene detection performance in experimental scenarios. Using the new metrics, we designed scCODE platform to optimize the DE gene detection performances (Figure. 1b).

Single cell RNA-seq data

We collected 20 scRNA-seq datasets from the conquer[16] database, including 3 datasets of UMI protocol and 17 datasets of full-length ones. These datasets varied in organism (mouse or human), cell type, sample size and other aspects. The scRNA-seq data used in this article was categorized into 3 types: simulated data without true DE genes (null data), simulated data with true DE genes and real experimental datasets.

Simulated data without true DE genes (null data): We generated multiple null instances from each experimental dataset by sampling from one type of cell, where no true DE gene was expected. One type of cells from each real dataset was selected as the source data where the null data sampled from. A stratification strategy was applied on the sample size, the number of cells in each sampling varied from 20%, 40%, 60%, 80% to 100% of the total number of the cells of the source dataset, and each sampling was replicated 5 times, thus we got 25 null instances from each experimental dataset.

Simulated data with true DE genes: Using the principle similar as above, we first generated null simulate data. Then, all cells were repeatedly resampled into two groups, while the gene expression of one cell group was multiplied by a fold-change value. In the simulated datasets, 2% of the genes were randomly selected as DE genes, the fold change of which depended on a gamma distribution with shape 4 and base varied in 2,3,4,5,6,7,8,10.

Experimental data: As mentioned before, we collected 20 real datasets, some of which contain more than two types of cells. And only two types of cells were selected as origin experimental data for DE gene detection between two groups of cells.

More information is detailed in the data information table (Supplementary table 1).

Evaluation of DE gene detection performance

Gene filtering methods

Although filtering genes with high-level of noise is already applied in many researches[21-23], there is no golden standard for gene filtering in scRNA-seq data. The methods used in most studies are hard-thresholding methods based on expression level, i.e., genes with expression value below threshold would be excluded. Conquer was included in this study as an option for hard thresholding. Another approach, OGFSC[18], which builds multiple candidate thresholding curves between CV^2 (coefficient of variation) and gene expression level μ, and provides an optimal thresholding curve to preserve biological signal and reduce technical noise. We also included feature selection function from scmap[20] as an alternative gene filtering method here, which ranks the information contained in each gene. The 3 gene filtering methods were benchmarked against no-filter strategy.

Conquer: conquer[16] is a representational filtering method of hard thresholding, which has been widely applied in scRNA-seq data analysis. Conquer filters genes with an expression above 1 count in less than 25% of the cells. In this article, we made it filter genes with an auto-adjusted threshold (5% - 25%) in order to maintain sufficient number of genes (>=2000).

OGFSC: OGFSC[18] is a tool developed particularly for filtering genes in scRNA-seq data, which distinguishes signal from noise based on a soft thresholding curve of expression level and variance. In this article, the thresholding curve is automatically determined by OGFSC.

Scmap: The feature selection of scmap[20] is not designed particularly for filtering genes, but it's power of sorting the informative genes is similar to that we concerned about. In this article, the function of feature selection in scmap is treated as an alternative gene filtering method, and the number of remained genes (top features) is set to 5000.

DE methods

There are plenty of DE methods for scRNA-seq data, here we examined 10 methods (Table 1). Some methods are usually used in bulk data, including edgeR[11], DESeq2[12] and limma[24]. For edgeR, we applied likelihood ratio test (LRT)[25] with default TMM normalization and default dispersion estimation. DESeq2 was run on rounded data by default settings. Voom-limma was applied on counts data by default settings. We also included 2 nonparametric methods, Wilcoxon test and SAMseq[26]. The Wilcoxon test were applied on log-transformed data with default settings. SAMseq was run on count data with permutations = 1000. Besides, 4 methods particularly designed for scRNA-seq data were applied, including MAST[8], BPSC[9], scDD[10] and DEsingle[27]. We applied MAST to log-transformed counts data without considering the cellular detection rate. BPSC and DEsingle were applied to count data with default settings. scDD

was run on count data, with prior parameter alpha=0.01, mu0=0, s0=0.01, a0=0.01, b0=0.01.

Finally, we also applied a non-paired t-test[13] on log-transformed data.

Table 1. DE methods and algorithm used in this article.

DE methods	Input data*	Software version
BPSC[9]	Count	0.99.2
DESeq2[12]	Count	1.30.1
edgeR[11]	Count	3.32.1
t-test[13]	log2(count+1)	stats 4.0.2
MAST[8]	log2(count+1)	1.16.0
SAMseq[26]	Count	samr 3.0
scDD[10]	Count	1.14.0
Wilcoxon-test[13]	log2(count+1)	stats 4.0.2
Voom-limma [24]	Count	3.46.0

*Input data: data type used in each DE methods, counts refers to gene read counts.

140 *Metrics*

The evaluation of the DE gene detection was based on several criteria commonly used before. FPR was calculated on null data, and genes detected with p-value ≤ 0.05 were recognized as false positive results. FDP, TPR and AUROC were calculated from simulated data with true DE genes, whereas genes with adjusted p-value ('fdr') ≤ 0.05 were assigned as positive DE genes. SNR was

145 calculated on both simulated data with true DE genes and experimental data by comparing the four characteristics between positive genes detected and non-positive ones.

Table 2. Metrics used in evaluation of analysis strategies for DE gene detection.

Metrics	Calculate from data	Description
FPR	null data	False positive rate
FDP	Simulated data with	False discovery proportion
TPR	Simulated data with	True positive rate
AUROC	Simulated data with	Area under ROC curve

SNR*	Simulated data with true DE genes/real	Signal-Noise Ratio

*SNR: signal-noise ratio was exhibited in 4 aspects of expression level, $\log 2(average\ expression + 1)$, $\log 2(variance + 1)$, coefficient of variation (CV) and the fraction of zero. And the SNR is calculated similar to conquer[16] like:

$SNR = \frac{\mu_p - \mu_{Np}}{\sigma_p - \sigma_{Np}}$; μ_p (μ_{Np}) and σ_p (σ_{Np}) represent the mean and s.d of the characteristic of the positive (not -positive) genes.

scCODE

scCODE metrics

The above metrics are highly dependent on the prior knowledge of true DE genes, which is not always available. AUROC is the golden standard widely applied, and it's defined as below:

$$AUROC = P\left(r(xp_i) > r(xn_j)\right) = 1 - \frac{\sum_M r(xp_i) - M(M+1)/2}{MN} \tag{1}$$

where $r(x)$ represents the rank of instance x, xp_i represents a randomly selected true positive instance, xn_j represents a randomly selected true negative instance; M represents the number of true positive instance, N represents the number of true negative ones.

To approximate AUROC in real cases, two metrics CDO and AUCC were used here. The basic concept of CDO and AUCC is similar to AUROC but requires no prior knowledge. CDO regards the most frequently identified positive genes as seed DE genes. It further reflects the ability of each method to detect these seed genes as positive results before other genes by calculate their

165   relative rank sorted by p-value in the DE results of each analysis strategy. AUCC is not a new metric designed by us, it's used to calculate the consistency between two pair of results. We assume that positive DE genes detected by more methods as more reliable, and the method with a higher CDO and AUCC is also considered as more trustworthy.

CDO: Consistent DE gene Order, is a novel parameter proposed here, range from 0 to 1.

170   Basically, the common DE genes detected by no less than 75% of the different analysis strategies are considered as seed DE genes. For DE gene detection in experimental data, we consider the seed DE genes as a part of the true positive results. if we could identify $M1$ seed DE genes, we calculate CDO like below:

$$CDO = P\left(r(xp_{i1}) > r(xn_{j1})\right) = 1 - \frac{\sum_{M1} r(xp_{i1}) - \frac{M1(M1+1)}{2}}{M1N1} \qquad (2)$$

175   where $M1 < M; N1 > N$.

We find that:

$$CDO = \frac{N}{N1} P\big(r(xp_{i1}) > r(xn_{j1})\big)_E |xn_{kj1} \in True\ negative\ +$$

$$\frac{M-M1}{N1} P\big(r(xp_{i1}) > r(xn_{k1})\big)_E |xn_{k1} \in False\ negative \tag{3}$$

While

$$P\big(r(xp_{i1}) > r(xn_{j1})\big) |xn_{kj1} \in True\ negative = AUROC \tag{4}$$

$$P\big(r(xp_{i1}) > r(xn_{k1})\big) |xn_{k1} \in False\ negative = c \tag{5}$$

180   where c is a constant which depends on the bias of seed DE genes, and $0 \le c \le 1$.

So

$$CDO = \frac{N}{N1} AUROC + c\frac{M-M1}{N1}$$

$$\Leftrightarrow CDO \propto AUROC \tag{6}$$

There is a liner relationship between CDO and $AUROC$.

In practice, we make the number of seed DE genes $M1$ no less than 5 in order to increase the

185   robustness of seed DE genes.

AUCC: AUCC (Area under Concordance Curve) [16], a measurement of concordance. In this

article, we calculated the AUCC value between different DE methods but with the procession of

the same gene filtering method. We build concordance curve for each pair of DE methods, for top

n genes, and the AUCC value of each DE method is the average AUCC value of itself with the

190   other 9 DE methods. AUCC can be calculated like below:

$$AUCC = \int_1^n y_x dx \tag{7}$$

where y_x is the number of common results between the two strategies， n represents the top n

DE gene results.

Consensus optimization

195   Using CDO and AUCC, we are able to identify optimal strategies for the specific scRNA-seq

dataset. Based on the DE genes detected by various analysis strategies, scCODE produces

consensus DE gene results. scCODE summarizes the top (default as all) DE genes from each of

the strategy selected. The principle of consensus optimization is that the DE genes with higher

frequency of observation by different analysis strategies are more reliable. We rank the genes by

200   their number of observations. For the genes with same number of observations, they are sorted by

absolute fold changes. Finally, the top n (median number of the DE genes detected from each

analysis strategy) genes are used as final result from scCODE.

Results

For evaluation of DE gene detection, we adopted 4 gene filtering approaches and 10 DE methods,

205 which lead to 40 analysis strategies (see Methods). Each data instance was first filtered with the

number of retained genes no less than 2000. From the remaining genes, DE ones were detected

using various DE methods. The results of 40 analysis strategies for each data instance were

systematically evaluated by the above metrics. In this article, while calculating the CDO and

AUCC metrics, the top n (default as 500) DE genes detected by each analysis strategy were

210 considered. Based on CDO and AUCC, scCODE provided personally optimized DE gene results

for any given scRNA-seq dataset.

The evaluation of FPR on null dataset without true DE genes

Using null datasets, we evaluated the type I error control power of different analysis strategies by

FPR. We recorded the proportion of genes that were assigned with a p-value lower than 0.05.

215 Similar to the evaluation in conquer[16], we found that most of these DE methods could keep the

FPR near to the imposed level. Some DE methods resulted an FPR higher than the imposed level

such as edgeR and limma, while other methods seemed to achieve an FPR value under the

theoretical level (Figure. 2). Furthermore, the FPRs after gene filtering procedure were generally

lower than those without filtering in most cases. For the DE methods that could keep the FPR

220 value well under 0.05, such as scDD, MAST and t-test, FPR seemed to be optimized by coupling

with OGFSC gene filtering. Overall, the FPR evaluation shows that gene filtering before DE gene detection does help to decrease the FPR. We also demonstrated the relationship between FPR and cell sample size. A cubic regression for each method shows that the cubic curve of FPR waved up when sample size increased (Supplementary Figure. 1). And we observed that the curves of the

225   same DE method were very similar in shape with various filtering methods. Therefore, the relation between FPR and sample size depends more on DE methods.

The evaluation of SNR on data containing DE genes

In order to evaluate the DE gene detection bias, we compared the characteristics of genes detected

230   as DE (adjusted p-value < 0.05) with that non-DE ones. The characteristics include the $\log 2(average\ expression + 1),\ \log 2(variance + 1)$, coefficient of variation (CV) and the fraction of zero (the expression level) among all cells. We calculated SNR on both simulated data and experimental datasets with known true DE genes. And only the results with more than 5 genes detected as DE were recorded as valid results. In fact, the results on experimental datasets were

235   largely similar with that of simulated data with true DE genes (Figure. 3, Supplementary Figure. 2). We noticed that the average expression level, variance and CV of the DE genes detected by either filtering methods and DE methods were significantly higher than that of non-DE ones (Figure. 3a, b, c), while the fraction of zeros was mostly lower (Figure. 3d). Furthermore,

considering the performance of different filtering methods, we observed that the SNR of each

240 characteristic after filtering by scmap and conquer was closer to 0. It indicated that the DE genes

detected by any analysis strategies with scmap or conquer shared a more similar characteristic

with those assigned as non-DE.

The evaluation of FDP, TPR and AUROC on simulated data with true signal

Using the simulated data with a proportion of true DE genes, we investigated the performance of

245 FDP, TPR and AUROC of different analysis strategies (Figure. 4). The FDP values of different

methods varied in a wide range, some methods such as SAMseq, BPSC and scDD could keep a

low FDP level in DE gene detection, but other methods including DESeq2 and limma, trended to

result in a higher FDP (Figure. 4a). By comparing the performance of different filtering methods,

we observed that all of the three filtering methods generally achieved lower FDP. For the DE

250 methods which achieved low FDP like MAST and BPSC, false positive error could be alleviated

by coupling with OGFSC. Filtering genes by conquer and scmap significantly improved the TPR

value for all DE methods (Figure. 4b).

AUROC can be considered as a direct and gold-standard metric to assess the performance of

methods. We observed that a lower FDP did not necessarily mean a higher AUROC, for example,

255 although MAST and scDD achieve lower FDP than edgeR and limma, they had similar AUROC

value (Figure. 4c). Gene filtering had a significant impact on AUROC values. The performances

of DE gene detection with gene filtering method varied a lot. If no-filter was performed, the AUROC value would range between only 0.6 to 0.8. It's remarkable that filtering genes by conquer and scmap could boost the AUROC up to 0.95 for DE methods like edgeR and scDD, and the average level for most DE methods would range between 0.8 to 0.9. The evaluation of TPR and AUROC indicated that the overall performances of most DE methods were generally similar, while filtering method seemed to be the one making the key difference (Figure. 4b, c).

Metrics for personalized assessment in scCODE

The previous sections have demonstrated that filtering out genes containing much noise could be of great beneficial to DE gene detection in scRNA-seq data, therefore, the performance of DE gene detection depends on both gene filtering methods and DE methods. However, we found that the AUROC values for different datasets were divergent, which suggests that the appropriate analysis strategy for different dataset may differ (Figure. 5a). It implies there needs a personalized analyzing strategy for each specific scRNA-seq dataset. However, existing metrics, such as AUROC, which requires known true results, could not be used on real scRNA-seq dataset prior knowledge of truth DE genes. Thus, we developed CDO to imitate AUROC in real scRNA-seq datasets, and applied AUCC with the concept that consensus results could be more reliable.

We validated the performance of CDO and AUCC on 19 simulated datasets with 2% of genes randomly selected as true DE genes. We recorded the results of each dataset individually, and the

275 corresponding AUROC results were used as reference. The "good" methods suggested by AUROC

were largely divergent for different datasets (Figure. 5a). We further investigated the correlation

between CDO, AUCC and AUROC for each simulated dataset. For most datasets, the correlation

between CDO and the reference AUROC was significant (Figure. 5b, Supplementary figure. 3).

The AUCC was also highly correlated with AUROC for most simulated datasets (Figure. 5C,

280 Supplementary figure. 4). Consistent to the mathematic derivation presented in the Methods

section, we here demonstrate CDO and AUCC are reliable approximates of AUROC in real

scenarios, and therefore we propose to use the two metrics in scCODE, which makes it practical

for personalized analysis strategy.

scCODE benchmarked on simulated datasets

285 Based on the metrics CDO and AUCC, scCODE could access the performance of different

analysis strategies on experimental scRNA-seq datasets. As mentioned before, we consider the DE

genes detected by majority of analysis strategies as more reliable, and make a further consensus

optimization of the DE gene results detected by the "suitable" analysis strategies assigned by

scCODE. Finally, scCODE provides the consensus optimized DE gene results as DE gene output.

290 Here, we benchmarked the performance of scCODE against other analysis strategies on several

simulated datasets. The sensitivity and specificity of the DE results were considered, a higher TPR

of the top DE genes would indicate better performance. The same analysis strategy applied to

different datasets obtained divergent performances. scCODE achieved higher TPR than most analysis strategies, which indicated that scCODE could identify DE genes more accurately for these simulated datasets (Figure. 6). Besides, the performance of scCODE was more stable on different simulated datasets.

scCODE validated on a real dataset

In this article, we also applied scCODE on a real scRNA-seq dataset from immune-stimulation research of mouse[19], containing stimulated CD4+ T cells (by antibodies to CD3ε and CD28) and naive ones (Figure. 7). The heatmap represented the normalized results of CDO and AUCC, which exhibited the appropriateness of various analysis strategy for the data. We selected 5 suitable analysis strategies (i.e., OGFSC-Wilcoxon-test, conquer-limma, etc.), and 5 unsuitable analysis strategies (i.e., OGFSC-MAST, no-filter-DEsingle, etc.) (Figure. 7a). Then, for each strategy, top DE genes with the smallest p-values (all with $p<0.05$) were identified as candidate DE genes. The results from suitable analysis strategies were compared to the unsuitable ones. The suitable analysis strategies detected a total of 1055 unique DE genes, while the unsuitable ones detected 1546 unique genes (Figure. 7b). Besides, the suitable analysis strategies had more overlapped genes (more than 50% genes were detected $\geq$ 3 times) than that of unsuitable ones (only 1.3% genes were detected $\geq$ 3 times). It indicated that the suitable strategies trend to achieve more consistent results.

Furthermore, we also compared the results after consensus ranking and selection (Figure. 7 c, d). The up-regulated genes of stimulated CD4+ T cells were processed by gene set enrichment analysis (Gene Ontology (GO)-biological process (BP) and KEGG pathway), and top 5 results were exhibited (Figure. 7e, f). We found that all of the enriched GOBP functions uncovered by suitable analysis strategies, such as RIBOSOME_BIOGENESIS, NCRNA_PROCESSING and RRNA_METABOLIC_PROCESS, were in profound agreement with the conclusion of the original article[19]; while that was not the case for the unsuitable results. And the KEGG results of suitable analysis strategies also detected a pathway (T_CELL_RECEPTOR_SIGNALING_PATHWAY) closely associated with T cell activation. The consensual DE gene results from the suitable analysis strategies of methods given by scCODE were more closely matched with the experimental question studied.

Discussion

DE gene detection in scRNA-seq data is one of the most concerned steps in data analyses. Gene filtering is generally applied before DE gene detection, but the influence of gene filtering is not fully recognized before. In this article, we first presented a systematic benchmark on both gene filtering methods and DE methods for scRNA-seq data. We exhibited the results from the aspects of FPR, FDP, SNR and AUROC. Without considering the influence of gene filtering, we got results in agreement with previous researches[15, 16]. Some methods like MAST and Wilcoxon-test performed well in specificity, and trended to detect less false DE genes (low FPR

330   and FDP). But not a single DE method showed a superiority in sensitivity (TPR), and most DE

methods achieved similar performance by AUROC. The performances with gene filtering methods

were divergent, and the improvement it brought was significant. We have shown that gene filtering

does help DE gene detection in all aspects. We also summarized the evaluation on multiple

simulated datasets (Supplementary figure. 5). The summary categorized each analysis strategy into

335   "Good", "intermediate" and "Poor" on each metric. It also showed that the performance of DE

gene detection depended on both gene filtering and DE processing.

Previous evaluation tools were mainly based on evaluating performance with cohort of datasets.

We have shown that the performance of DE gene detection on different datasets varies, and

personalized analytical strategy on the specific dataset studied is necessary. To assess different

340   methods in practical DE detection studies, two novel metrics CDO and AUCC were proposed in

this study. Theoretically, the new metrics could faithfully assess a DE gene detection result; and it

was also approved on simulated datasets by demonstrating significant correlation between the

proposed metrics and AUROC as reference. In fact, the reliability of CDO is based on the

hypothesis that the seed genes are true DE ones. We considered the common DE genes detected by

345   different analysis strategies as seed genes, and investigated the threshold to guarantee its reliability.

We validated the relationship between the threshold of seed gene selection (genes detected in how

many percent of strategies) and significance of correlation between CDO and AUROC

(Supplementary figure. 6). The results on simulated data created by splatter[28] showed that the

reliability of CDO was highly dependent on the threshold (Supplementary figure. 6). A low

350   threshold tended to result in high false-positive rate in seed gene selection, while a too high

threshold would sharply reduce the number of seed genes and its stability. Thus, we suggested and

applied a threshold of seed genes higher than 75%, and restricted the number of seed genes no less

than 5 in this article.

The R package based on the two metrics, scCODE, answered the question of which analysis

355   strategies are appropriate for a scRNA-seq dataset. More importantly, it can provide optimized DE

gene detection results of experimental datasets. We validated scCODE on both simulated datasets

and a real scRNA-seq dataset. Figure. 6 showed that scCODE achieved better and more robust

results than most analysis strategies on each dataset, as it judged and applied personalized optimal

strategies on these datasets, while the other analysis strategies performed unstably on different

360   datasets. Besides, the validation on the experimental scRNA-seq dataset exhibited that scCODE

could provide reliable assessment of these strategies on real scRNA-seq dataset, and could further

provide reliable DE gene results for the users.

scCODE currently contains 4 filtering methods and 10 DE methods, it automatically produces the

optimal DE gene results for the given dataset (A demo output of scCODE could be seen in

365   Supplementary table. 2). A limitation of scCODE is that it needs a comparison of multiple

methods to find the seed DE genes, thus it may be computationally demanding. We also provided

a light version of scCODE including only a few reliable methods, such as scDD, MAST and

DESeq2 (Supplementary figure. 5a). The light version provided very similar result on the experimental data tested (Supplementary figure. 5b). We believe that scCODE is convenient to conduct optimized DE gene detection for specific scRNA-seq data. In the future, more methods will be incorporated in scCODE. Besides, there may be other preprocessing steps which may be included in the evaluation framework of scCODE, such as normalization[29, 30], etc.

Conclusion

In this study, we systematically evaluated the true influence of gene filtering on DE gene detection. The performance of DE gene detection depends on both gene filtering and DE processing. And we found that personalized analysis strategy is needed to achieve optimal DE gene results for each experimental scRNA-seq dataset. However, existing metrics requiring prior truth DE genes were not suitable for experimental datasets. We established two novel metrics CDO and AUCC for it, and observed significant correlation between the new metrics and AUROC (regarded as ground truth) on multiple simulated datasets independently. Finally, we developed scCODE, a R package for optimizing DE gene detection of scRNA-seq dataset. scCODE performed better and more robust on different simulated datasets. The consensus optimized DE gene results suggested by scCODE better match the experimental design of the test data. We hope our user-friendly tool scCODE could help the analysis of each scRNA-seq dataset, and could be beneficial to maintain more reliable DE gene results.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Availability of data and materials

All of the code were based on R version 4.02.

The origin data used in evaluation are available in the conquer repository, at http://imlspenticton.uzh.ch:3838/conquer/.

The experimental data used to test scCODE is available in the reference paper[19], and also can be found at https://github.com/XZouProjects/scCODE/tree/main/data.

The R code for evaluation of gene filtering and DE methods is available at https://github.com/XZouProjects/scCODE/evaluation.

The installable version of the scCODE R package is available at https://github.com/XZouProjects/scCODE.

Competing interests

The authors have declared no competing interests.

Funding

This work was supported in part by the National Natural Science Foundation of China [31601070 to JH, 31800253 to KC and JH]; the Translational Medicine Cross Research Fund of Shanghai Jiao Tong University [ZH2018QNB29 to JH, RH, XBZ]; the Natural Science Foundation of Shanghai [grant number 16ZR1417900 to XZ]; the Shanghai Pujiang program [grant numbers 16PJ1405200 to XZ, 16PJ1405100 to JH]; the Shanghai Sailing Program [grant number 17YF1410400 to KC].

Authors' contributions

JWZ, JH, XZ, KC conceived and designed the algorithm. JWZ developed the software. JH, XZ, KC, MCW, ZZ and ZQL provided data interpretation and biological explanation. JWZ, JH, XZ and KC wrote the manuscript, JH, XZ, KC, JWZ and MCW prepared figures. RH and XBZ provided critical suggestions. The manuscript was approved by all authors.

Acknowledgements

We thank Li Xuejing for her help of Figure. 2.

The data analysis was supported by GorgeousAI (Gu Ai) Co. Ltd.

Reference

1.   La Manno G, Gyllborg D, Codeluppi S, Nishimura K, Salto C, Zeisel A, Borm LE, Stott SRW, Toledo EM, Villaescusa JC, et al: **Molecular Diversity of Midbrain Development in Mouse, Human, and Stem Cells.** *Cell* 2016, **167:**566-580 e519.

2.   Young MD, Mitchell TJ, Vieira Braga FA, Tran MGB, Stewart BJ, Ferdinand JR, Collord G, Botting RA, Popescu DM, Loudon KW, et al: **Single-cell transcriptomes from human kidneys reveal the cellular identity of renal tumors.** *Science* 2018, **361:**594-599.

3.   Litzenburger UM, Buenrostro JD, Wu B, Shen Y, Sheffield NC, Kathiria A, Greenleaf WJ, Chang HY: **Single-cell epigenomic variability reveals functional cancer heterogeneity.** *Genome Biol* 2017, **18:**15.

4.   Satpathy AT, Granja JM, Yost KE, Qi Y, Meschi F, McDermott GP, Olsen BN, Mumbach MR, Pierce SE, Corces MR, et al: **Massively parallel single-cell chromatin landscapes of human immune cell development and intratumoral T cell exhaustion.** *Nat Biotechnol* 2019, **37:**925-936.

5.   Azizi E, Carr AJ, Plitas G, Cornish AE, Konopacki C, Prabhakaran S, Nainys J, Wu K, Kiseliovas V, Setty M, et al: **Single-Cell Map of Diverse Immune Phenotypes in the Breast Tumor Microenvironment.** *Cell* 2018, **174:**1293-1308 e1236.

6.   Lang C, Campbell KR, Ryan BJ, Carling P, Attar M, Vowles J, Perestenko OV, Bowden R, Baig F, Kasten M, et al: **Single-Cell Sequencing of iPSC-Dopamine Neurons Reconstructs Disease Progression and Identifies HDAC4 as a Regulator of Parkinson Cell Phenotypes.** *Cell Stem Cell* 2019, **24:**93-106 e106.

7.   Pollen AA, Nowakowski TJ, Shuga J, Wang X, Leyrat AA, Lui JH, Li N, Szpankowski L, Fowler B, Chen P, et al: **Low-coverage single-cell mRNA sequencing reveals cellular heterogeneity and activated signaling pathways in developing cerebral cortex.** *Nat Biotechnol* 2014, **32:**1053-1058.

8.   Finak G, McDavid A, Yajima M, Deng J, Gersuk V, Shalek AK, Slichter CK, Miller HW, McElrath MJ, Prlic M, et al: **MAST: a flexible statistical framework for assessing**

445   **transcriptional changes and characterizing heterogeneity in single-cell RNA sequencing data.** *Genome Biol* 2015, **16:**278.

9.   Vu TN, Wills QF, Kalari KR, Niu N, Wang L, Rantalainen M, Pawitan Y: **Beta-Poisson model for single-cell RNA-seq data analyses.** *Bioinformatics* 2016, **32:**2128-2135.

10.   Korthauer KD, Chu LF, Newton MA, Li Y, Thomson J, Stewart R, Kendziorski C: **A**
450   **statistical approach for identifying differential distributions in single-cell RNA-seq experiments.** *Genome Biol* 2016, **17:**222.

11.   Robinson MD, McCarthy DJ, Smyth GK: **edgeR: a Bioconductor package for differential expression analysis of digital gene expression data.** *Bioinformatics* 2010, **26:**139-140.

12.   Love MI, Huber W, Anders S: **Moderated estimation of fold change and dispersion for**
455   **RNA-seq data with DESeq2.** *Genome Biol* 2014, **15:**550.

13.   Robinson MD, Oshlack A: **A scaling normalization method for differential expression analysis of RNA-seq data.** *Genome Biol* 2010, **11:**R25.

14.   Li HS, Ou-Yang L, Zhu Y, Yan H, Zhang XF: **scDEA: differential expression analysis in single-cell RNA-sequencing data via ensemble learning.** *Brief Bioinform* 2021.

460 15.   Jaakkola MK, Seyednasrollah F, Mehmood A, Elo LL: **Comparison of methods to detect differentially expressed genes between single-cell populations.** *Brief Bioinform* 2017, **18:**735-743.

16.   Soneson C, Robinson MD: **Bias, robustness and scalability in single-cell differential expression analysis.** *Nat Methods* 2018, **15:**255-261.

465 17.   Zhu A, Ibrahim JG, Love MI: **Heavy-tailed prior distributions for sequence count data: removing the noise and preserving large differences.** *Bioinformatics* 2019, **35:**2084-2092.

18.   Hao J, Cao W, Huang J, Zou X, Han ZG: **Optimal Gene Filtering for Single-Cell data (OGFSC)-a gene filtering algorithm for single-cell RNA-seq data.** *Bioinformatics* 2019,
470   **35:**2602-2609.

19. Martinez-Jimenez CP, Eling N, Chen HC, Vallejos CA, Kolodziejczyk AA, Connor F, Stojic L, Rayner TF, Stubbington MJT, Teichmann SA, et al: **Aging increases cell-to-cell transcriptional variability upon immune stimulation.** *Science* 2017, **355:**1433-+.

20. Kiselev VY, Yiu A, Hemberg M: **scmap: projection of single-cell RNA-seq data across data sets.** *Nat Methods* 2018, **15:**359-362.

21. Mathys H, Davila-Velderrain J, Peng Z, Gao F, Mohammadi S, Young JZ, Menon M, He L, Abdurrob F, Jiang X, et al: **Single-cell transcriptomic analysis of Alzheimer's disease.** *Nature* 2019, **570:**332-337.

22. Kim C, Gao R, Sei E, Brandt R, Hartman J, Hatschek T, Crosetto N, Foukakis T, Navin NE: **Chemoresistance Evolution in Triple-Negative Breast Cancer Delineated by Single-Cell Sequencing.** *Cell* 2018, **173:**879-893 e813.

23. Huang S, Ziegler CGK, Austin J, Mannoun N, Vukovic M, Ordovas-Montanes J, Shalek AK, von Andrian UH: **Lymph nodes are innervated by a unique population of sensory neurons with immunomodulatory potential.** *Cell* 2021, **184:**441-459 e425.

24. Law CW, Chen Y, Shi W, Smyth GK: **voom: Precision weights unlock linear model analysis tools for RNA-seq read counts.** *Genome Biol* 2014, **15:**R29.

25. McCarthy DJ, Chen Y, Smyth GK: **Differential expression analysis of multifactor RNA-Seq experiments with respect to biological variation.** *Nucleic Acids Res* 2012, **40:**4288-4297.

26. Li J, Tibshirani R: **Finding consistent patterns: a nonparametric approach for identifying differential expression in RNA-Seq data.** *Stat Methods Med Res* 2013, **22:**519-536.

27. Miao Z, Deng K, Wang X, Zhang X, Berger B: **DEsingle for detecting three types of differential expression in single-cell RNA-seq data.** *Bioinformatics* 2018, **34:**3223-3224.

28. Zappia L, Phipson B, Oshlack A: **Splatter: simulation of single-cell RNA sequencing data.** *Genome Biol* 2017, **18:**174.

29. Buttner M, Miao Z, Wolf FA, Teichmann SA, Theis FJ: **A test metric for assessing single-cell RNA-seq batch correction.** *Nat Methods* 2019, **16:**43-49.

500   30.   Vallejos CA, Risso D, Scialdone A, Dudoit S, Marioni JC: **Normalizing single-cell RNA sequencing data: challenges and opportunities.** *Nat Methods* 2017, **14**:565-571.

Figure. 1 The Evaluation of DE gene detection performance and the schematic of scCODE. (a) The evaluation of the performance of different analysis strategies. (b) The schematic of scCODE.

505   Figure. 2 The violin plot of FPR results of different analysis strategies. The results are exhibited separately by diverse DE methods, and 4 different colors of violins represents 4 gene filtering methods. Each violin contains the FPR values resulted from every null data instance, central line, median value; dotted line, first and third quantile. Outliers (larger than third quantile + 2*IQR or smaller than first quantile – 2*IQR) are removed.

510   Figure. 3 The SNR of DE gene detection on simulated datasets with true DE genes, exhibited by 4 characteristics: (a) SNR of log2(average+1). (b) SNR of log2(variance+1). (c) SNR of cv. (d) SNR of fraction of zeros. The color of box represents different DE methods; the colored rectangle represents different filtering methods. Dotted line, SNR = 0; n, number of simulated datasets. Outliers (larger than third quantile + 2*IQR or smaller than first quantile – 2*IQR) are excluded.

515   Figure. 4 The Evaluation of FDP, TPR and AUROC on multiple simulated datasets. (a) Observed FDP with an adjusted p-value cutoff of 0.05, separated by high FDP (average > 0.2) and low FDP (average FDP < 0.1). (b) Observed TPR with an adjusted p-value cutoff of 0.05. (c) Observed

AUROC with an adjusted p-value cutoff of 0.05. The colored boxes represent different DE methods; the lower horizontal-colored rectangles represent different filtering methods. n, number of simulated datasets. Outliers (larger than third quantile + 2*IQR or smaller than first quantile – 2*IQR) are excluded.

Figure. 5 The reference AUROC of two simulated data with known true DE genes, and the spearman correlation between CDO, AUCC and AUROC. Left, Simulated-GSE48968-GPL13112 dataset; right, Simulated-GSE62270-GPL17021 dataset. (a) The heatmap of observed AUROC resulted by 40 analysis strategies on the two simulated data. (b) The correlation between CDO and AUROC, each point represents the result of one analysis strategy. (c) The correlation between AUCC and AUROC, each point represents the result of one analysis strategy.

Figure. 6 scCODE benchmarked on simulated datasets. (a) scCODE compared with the analysis strategies filtering by OGFSC, the TPR was calculated on the top n DE gene results detected by each analysis strategy, where n = the total number of DE genes set in each simulated dataset. (b) scCODE compared with analysis strategies filtering by conquer. (c) scCODE compared with analysis strategies filtering by scmap. (d) scCODE compared with analysis strategies without filtering genes.

Figure. 7 scCODE validated on a real scRNA-seq data comparing activated CD4+ T cells with naïve ones. (a) Heatmap of the average results of CDO and AUCC. White rectangle, 5 suitable

analysis strategies suggested by scCODE; black rectangle, top 5 unsuitable analysis strategies. (b) Bar plot of the DE gene information of the 5 suitable and unsuitable analysis strategies. Dotted line, consensus selection line, the genes above the line were selected. (c, d) Volcano map and consensus information of the top 500 DE genes collected from each of the analysis strategy; (e) suitable ((f) unsuitable) analysis strategies. Dotted line, p-adjusted = 0.05, logFC = $\pm$ 1. Detected times, gene detected times within the 5 analysis strategies selected. (e, f) GOBP and KEGG analyze of the up-regulated genes in activated CD4+ T cells from consensus DE genes of (e) suitable and (f) unsuitable analysis strategies. Top 5 results of GOBP and KEGG are exhibited.

Additional files

1. Supplementary table. 1 A excel lists the detail information of the experimental datasets and the simulated datasets, which were used for evaluation of DE gene detection performance.

2. Supplementary table. 2. A excel lists the demo results of scCODE applied on the immune-senescence dataset in this article. It contains 4 sheets, the results of Z-Score of AUCC, the results of Z-Score of CDO, the strategy information selected (default as top 5) and the information of the final DE genes results.

3. Supplementary figures. A PPT file includes all the supplementary figures in sequence. The figure legend and description are list in PPT comment of each page.

Evaluation of DE gene detection

Annotated scRNA-seq database

OGFSC
conquer
scmap
No filter

BPSC
DESeq2
edgeR
⋮
MAST
scDE
ttest

Metrics:
FPR
FDP
SNR
TPR
AUROC

CDO AUCC

scCODE

| Data preparation | Filter methods | DE methods | Evaluation | 'scCODE' R package |

scCODE

AUCC+CDO

OGFSC
conquer
scmap
no filter

edgeR DESingle limma DESeq2 sarr wilcoxtest BPSC ttest MAST scDD

Suitable / Unsuitable

scCODE

Top several analysis strategies: e.g.

(1) OGFSC & Limma
(2) OGFSC & ttest
(3) OGFSC & wilcoxtest
(4) conquer & ttest
(5) conquer & limma

Consensus selection

Times detected

N gene
1000
500
0

Selected

Suitable — Analysis strategies

Select DE genes by consensus concept

Sp100
Map2k3
Fos
Slamf1
Sell
Tcf7
⋮

DE genes

| scRNA-seq data | 'scCODE' R package | | | Output |

a log2(average+1)

b log2(variance+1)

c CV

d Fraction of zeros

SNR

DE methods

- edgeR
- DEsingle
- limma
- DESeq2
- samr
- wilcox-test
- BPSC
- t-test
- MAST
- scDD

Filtering methods: OGFSC | conquer | scmap | No filter

a

scCODE compared with filtering by OGFSC

b

scCODE compared with filtering by conquer

c

scCODE compared with filtering by scmap

d

scCODE compared with filtering by no filter