Olorofim and the azoles are antagonistic in *A. fumigatus* and functional genomic screens

reveal mechanisms of cross resistance.

Authors: Norman van Rhijn[1,2], Sam Hemmings[1], Clara Valero[1,3], Jorge Amich[1], Michael J

Bromley[1,2]

[1] Manchester Fungal Infection Group, Division of Evolution, Infection, and Genomics, Faculty of Biology,

Medicine and Health, University of Manchester, CTF Building, 46 Grafton Street, Manchester, M13 9NT, UK.

[2] Antimicrobial Resistance Network, University of Manchester, Manchester M13 9PT, UK

8
[3] Faculdade de Ciências Farmacêuticas de Ribeirão Preto, Departamento de Ciências Farmacêuticas, Universidade

de São Paulo, Avenida do Café S/N, Ribeirão Preto 14040-903, Brazil.

*To whom correspondence should be addressed:

Michael   J   Bromley,   Phone: (+44)   (0)161   275   1703,   email:

mike.bromley@manchester.ac.uk

Keywords: Aspergillus fumigatus, Olorofim, Orotomide, Antimicrobial resistance,

Antifungal, Transcription factor, Aspergillosis, Nitrogen metabolism, Antagonism

Abstract:

Aspergillosis, in its various manifestations, is a major cause of morbidity and mortality. Very

few classes of antifungal have been approved for clinical use to treat these diseases and

resistance to the first line therapeutics is increasing. A new class of antifungals, the orotomides,

are currently in development with the first compound in this class olorofim in late-stage clinical

trials. In this study, we characterise a network of genes that govern olorofim response in *A.*

fumigatus. We reveal that the number of transcription factors that regulate olorofim

susceptibility are far fewer than we have previously observed for the azoles and the change in

sensitivity observed in these isolates is less extreme. Intriguingly, loss of function in two higher

order transcriptional regulators, HapB a member of the heterotrimeric HapB/C/E (CBC)

complex or the regulator of nitrogen metabolic genes AreA, leads to cross resistance to both

the azoles and olorofim. However, a clinical azole resistant isolate with a point mutation in

HapE (*hapE^{P88L}*) retains sensitivity to olorofim. Our transcriptomic analysis suggests that

altered sensitivity to olorofim may emerge via modification of genes involved in the production

of pyrimidine biosynthetic precursors. Finally, we also show that the action of the azoles are

antagonistic to olorofim *in vitro*.

Introduction:

Invasive and chronic forms of aspergillosis affect over 3 million people resulting in excess of 300 thousand deaths per year [1]. Only three classes of antifungals are currently clinically available to treat aspergillosis, with the triazoles used as first-line therapy in most centres [2]. Resistance to the azoles is rising, which is linked to usage of triazole compounds in agri- and horticulture [3]. It is expected more resistant *A. fumigatus* will be seen due to increased azole used in farming linked to climate change [4]. In the last decade, isavuconazole is the only antifungal to have been approved by the Food and Drug Administration (FDA) for treatment of invasive aspergillosis [5], however many of the mechanisms that contribute to resistance to the other azoles also result in isavuconazole resistance. The development of novel classes of antifungals will be a key component to addressing the emerging resistance problem. Fortunately, there are a number of novel classes of antifungal currently in development for treatment of invasive aspergillosis (IA) including ibrexafungerp, which has recently (2021) gained FDA approval for treatment of vulvovaginal candidiasis and olorofim (phase 3) [6]. Olorofim (formerly known as F901318 and under development by F2G, Ltd.) is of particular interest as it is not only a member of a novel class of antifungal (the orotomides) but unlike ibrexafungerp has a novel mechanism of action that has not been exploited clinically [7]. As olorofim is orally bioavailable it presents a realistic alternative to the azoles for long-term treatment of chronic and allergic infections and especially resistant infections [8] and could potentially be used in combination therapy strategies to supress the emergence of resistance.

Olorofim acts by inhibiting the enzyme dihydroorotate dehydrogenase (DHODH), which is a crucial enzyme within the pyrimidine biosynthesis pathway and is required for both DNA and RNA synthesis [7]. Structural and biochemical analysis of DHODH suggests olorofim competes with CoQ to bind to DHODH, preventing the oxidation of dihydroorotate to orotate.

DHODHs are grouped into 2 classes according to sequence similarity and subcellular

localisation. Both mammals and most fungi have class 2 DHODH, which is bound to the inner

mitochondrial membrane [9]. The human DHODH only shares 30% protein sequence identity

with the *A. fumigatus* DHODH and olorofim has also been demonstrated to be >2,200-fold

more potent against the *A. fumigatus* enzyme [7]. Inhibition of the pyrimidine biosynthesis

pathway by olorofim prevents the germination of *A. fumigatus* conidia and causes hyphae to

undergo morphological changes [10]. Prolonged exposure of germlings and vegetative hyphae

to olorofim also causes extensive isotropic expansion that is then followed by cell lysis [11].

Olorofim has an effect on a wide range of fungi and has been shown to be effective against

Coccidioides immitis, *Scedosporium* spp., *Madurella mycetomatis*, *Lomentospora prolificans*

and several *Aspergillus* species [12-18]. Olorofim is also effective against triazole resistant *A.*

fumigatus isolates and cryptic *Aspergillus* species [19, 20]. In several murine models of

aspergillosis, olorofim treatment significantly reduced fungal burden and mortality [15].

However, olorofim has a reduced activity against *Fusarium solani* species complex and

Fusarium dimerum and is inactive against Mucorales [21]. To date, there is no reported

incidence of acquired olorofim resistance in clinical isolates of normally susceptible fungal

species.

In this study, we explore potential routes to olorofim resistance for *A. fumigatus* through

screening a 484-member *A. fumigatus* transcription factor null mutant library. We have

uncovered a total of four transcription factors, which regulate susceptibility to olorofim.

Existing published literature, and our phenotypic and transcriptomic data revealed these

transcription factors regulate genes involved in upstream processes of the pyrimidine

biosynthesis pathway. Notably two of these transcription factor null mutants, Δ*hapB* and

Δ*areA,* had elevated MICs to olorofim and are resistant to the azole class of antifungal

highlighting potential routes to cross resistance. As HapB is a member of the heterotrimeric

CCAAT-binding complex (which comprises HapB/C and E) we assessed the susceptibility of

a azole resistant clinical isolate with a mutation in *hapE* (*HapE^{P88L}*). Intriguingly this mutation

does not change susceptibility towards olorofim. Furthermore, we demonstrate that there is an

antagonism between the azoles and olorofim in three wild-type strains of *A. fumigatus* and the

olorofim resistant mutants. Despite our discovery that there are mechanisms that can reduce

susceptibility to both azoles and orotomides we feel these results do not preclude the

development of combination therapeutics with these two agents. Our study also provides data

that can be used to develop molecular diagnostic tests to identify olorofim resistant isolates in

clinic.

Materials and Methods:

Fungal strains

Conidia of *Aspergillus fumigatus* MFIG001 (a derivative of CEA10) and transcription factor

null mutants [22, 23] were prepared by inoculating strains in vented 25cm^2 tissue culture flasks

with Sabouraud Dextrose agar (Oxoid, Hampshire, England) and incubating at 37°C for 48

hours. Spores were harvested in PBS + 0.01% Tween-20 by filtration through Miracloth.

Spores were counted using a haemocytometer (Marienfeld Superior, Baden-Württemberg,

Germany). The clinical isolate VP067-36, containing the P88L allele of HapE, was a kind gift

from Willem Melchers and Paul Verweij [24]. To construct the HapE^{P88L} allele in MFIG001,

the *hapE* gene was amplified from VP067-36 using primers hapE_gene_Fw and

hapE_gene_Rv (**Supplementary Table 1**). This PCR product was transformed into *A.*

fumigatus MFIG001 using CRISPR-Cas9 mediated transformation, using hapE_392_revcom

as crRNA [25]. Transformants were selected using voriconazole (1 mg/L) containing YPS

plates. Transformation was confirmed by PCR amplification of the complete *hapE* gene

(hapE_Fw and hapE_Rv), followed by Sanger sequencing using hapE_Seq.

Olorofim MIC screening

Olorofim was a kind gift of F2G Ltd. The Minimum Inhibitory Concentration (MIC) of

olorofim against *A. fumigatus* was assessed using the European Committee for Antimicrobial

Susceptibility Testing (EUCAST) methodology [21, 26]. Briefly, 2x10^4 spores/mL (in 100 µl)

were added to a CytoOne® 96-well plate (StarLab, Brussels, Belgium) containing 1xRPMI-

1640 medium (Sigma Aldrich, St. Louis, MO), 165 mM MOPS buffer (pH 7.0), 2% glucose,

with olorofim 2-fold dilution series ranging from 0.1 µg/L to 0.25 mg/L and a drug free control

(n = 4). Additionally, a serial dilution of olorofim containing 10 mM uracil and uridine was

performed. 96-well plates were incubated at 37°C for 48 hours. The MIC was determined as

the minimum drug concentration at which no germination was observed. Optical density was

measured at 600 nm using a Synergy™ HTX Multi-Mode Microplate Reader (BioTek,

Winooski, VT). In keeping with research laboratory based definitions, but in contract to

definitions used clinically, we define *in vitro* resistance as a strain that is less susceptible to

drug than the parental isolate [27].

Olorofim sensitivity screening of the A. fumigatus transcription factor null mutant library

$2x10^4$ spores/mL from each of the 484 members of the transcription knockout library were

added to 1x RPMI-1640 medium, 165 mM MOPS buffer (pH 7.0), 2% glucose in each well of

a CytoOne® 96-well plate with 0.002 mg/L olorofim (n = 4). Plates were incubated at 37°C for

48 hours. Fitness was calculated by dividing the optical density of respective null mutants to

the MFIG001 control. Relative fitness in olorofim was calculated by dividing fitness in

olorofim with general growth fitness of the transcription factor null mutants using the same

microculture conditions in 1x RPMI-1640 medium, 165 mM MOPS buffer (pH 7.0), 2%

glucose without olorofim (n = 4). Optical density was measured at 600 nm on a Synergy™

HTX Multi-Mode Microplate Reader (BioTek, Winooski, VT).

RNA-extraction

$1x10^6$ spores/mL of *A. fumigatus* MFIG001, Δ*AFUB_056620* and Δ*AFUB_030440* were

inoculated into 50 mL of *Aspergillus* complete media (ACM) [28] and incubated for 18 hours

at 37°C in a rotary shaker (180 rpm). Mycelia were harvested using filtration through Miracloth

(Merck Millipore) and washed in 1x RPMI-1640 medium. Approximately 1g of mycelia was

added to shake flasks containing 50 mL RPMI-1640 medium, 165 mM MOPS buffer (pH 7.0),

2% glucose and incubated for 1 hour at 37°C in a rotary shaker (180rpm) in the presence or

absence of 0.062 mg/L olorofim (n = 3). Mycelia was filtered through Miracloth and

snapfrozen using liquid nitrogen and kept at -80°C until required.

To extract RNA, 1 mL of TRIzol reagent (Sigma Aldrich) and 710-1180 μm acid washed glass

beads (Sigma Aldrich) were added to frozen mycelia and placed in a TissueLyser II® (Qiagen,

Hilden, Germany) for 3 minutes at 30 Hz. The solution was centrifuged (12,000 rpm) for 1

minute at 4°C. The aqueous phase was added to 200 μL of chloroform and centrifuged (12,000

rpm) for 10 minutes at room temperature. The supernatant was added to 0.2 M sodium citrate,

0.3 M sodium chloride and 25% (v/v) isopropanol and left at room temperatures for 10 minutes.

This solution was centrifuged (12,000 rpm) for 15 minutes at 4°C. The supernatant was

removed; the pellet was washed in 70% (v/v) ethanol and resuspended in RNase free water

(Thermo Fisher Scientific, Waltham, MA). RNA samples were treated with RQ1 RNase-Free

DNase (Promega, Madison, WI) and purified using a RNeasy Mini Kit (Qiagen). RNA quality

and quantity were assessed using gel electrophoresis and using a NanoDrop^{TM} 2000/2000c

Spectrophotometer (Thermo Fisher Scientific). All RNA extractions were carried out in

triplicate.

Transcriptomic Analysis

RNA sequencing was carried out by the Genomic Technologies Core Facility (GTCF) at The

University of Manchester. Sequencing libraries were prepared from mRNA using TruSeq$^{®}$

Stranded mRNA assay (Illumina, San Diego, CA). Samples were sequenced on a single lane

on an Illumina HiSeq2500 (Illumina). Low-quality reads of resulting fastq files were removed

using FastQC and trimmed using Trimmomatic (Quality >20, Sliding window average of 4

bases) [29]. Bowtie was used to align libraries to the *A. fumigatus* A1163 genome assembly

GCA_000150145.1 with gene annotation from CADRE/Ensembl Fungi v24 [30]. Differential

expression analysis of was performed using DESeq2 [31].

Functional category and gene ontology enrichment analysis was carried out using FungiFun2

2.2.8, converting genes to Af293 gene names to allow using the KEGG option [32]. Genes that

showed over 2-fold in differential expression and Benjamin-Hochberg FDR <0.01 underwent

enrichment analysis. StringsDB analysis was performed by only including genes with at least

two connections.

Phenotypic analysis

For colony images, 500 spores per isolate were spotted onto solid ACM or Aspergillus Minimal

Media (AMM) and left to dry. Plates were incubated at 37°C for 72 hours and imaged. Growth

on solid AMM supplemented with different nitrogen sources (50 mM ammonium tartrate, 10

mM sodium nitrate, 10 mM L-glutamine, 10 mM urea or 10 mM L-proline) were assessed by

spotting 500 spores from each isolate (n = 3). Plates were incubated at 37°C for 72 hours. MICs

were determined using the same supplementation as the phenotypic test with a serial dilution

of olorofim (ranging from 0.1 μg/L to 0.25 mg/L). 96-well plates were incubated for 48 hours

at 37°C and growth was determined by microscopic evaluation.

Checkerboard assays

For assessing drug combination efficacies of itraconazole and olorofim against *A. fumigatus*,

we used a checkerboard assay similar to EUCAST MIC testing described above. Twofold serial

dilutions of itraconazole were prepared across the X-axis and olorofim serial dilutions across

the Y-axis. The MIC was determined by microscopy by visually assessing the well containing

the lowest drug concentration with non-germinated spores. The fractional inhibitory

concentration index (FICI) was calculated as the MIC in combination divided by the MIC of

individual drugs [33].

Disk assays

4×10^4 conidia of the relevant *A. fumigatus* strain were evenly distributed on solidified 1xRPMI

1640 (Sigma), 165 mM MOPS buffer (pH 7.0), 2% glucose. One 6 mm antibiotic assay disk

(Whatman) was placed on the middle of the plate or two disks at a fixed distance, and 10 μL

of voriconazole (0.8 mg/mL), olorofim (0.005 mg/L) or H_2O_2 (30%) were added to each of

them. The plates were incubated at 37°C for 48 hours and imaged.

Data availability

RNA-seq data is available from ArrayExpress as experiment: E-MTAB-10590. The differential

expression output from DESeq2 is included as Supplementary Data 1. (reviewer access:

Reviewer_E-MTAB-10590 Password: pptwwqmj)

Acknowledgements

The authors would like to thank F2G for supplying the drug. We would also like to thank the

Genomic Technology Core Facility, the Bioinformatics Core Facility in the University of

Manchester for their technical support. This work was supported by the Wellcome Trust grant

number 219551/Z/19/Z and 208396/Z/17/Z to M.B. CV is funded by postdoctoral fellowship

from Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP-BEPE 2020/01131-

5).

Competing interests

Michael Bromley is a former employee of F2G Ltd. F2G currently funds a PhD position in the

laboratory. F2G was not involved in any of the experimentation or analysis of data in this study.

Author contributions

214 N.v.R designed and performed the experiments, analysis, wrote and edited the manuscript.
215 S.H. designed and performed experiments and analysis. C.V. designed and performed
216 experiments and analysis. J.A. designed and performed experiments and analysis. M.B.
217 provided the funding, designed the experiments, wrote and edited the manuscript
218
219
Results:

The effect of olorofim on A. fumigatus MFIG001 results in minor transcriptional changes

in the pyrimidine biosynthetic pathway.

In order to standardise assays throughout, the Minimum Inhibitory Concentration (MIC) of

olorofim against *Aspergillus fumigatus* MFIG001 was determined. The MIC was defined as

the minimum concentration of olorofim at which no germination from *Aspergillus* spores was

observed by microscopic evaluation (**Supplemental Figure 1**). Microscopic evaluation

revealed the MIC of olorofim to be 0.06 mg/L for *A. fumigatus* MFIG001, consistent with

previous findings of other *A. fumigatus* isolates [19]. The effect of olorofim on growth of *A.*

fumigatus was further evaluated by measuring optical density of the plates used to determine

the MIC (**Figure 1a**). The maximal growth observed (OD_{600} = 0.39) and minimal growth

observed (OD_{600} = 0.04) was separated by a 64-fold difference in drug concentration, which is

in stark contrast to the inhibitory effects of the azoles on *A. fumigatus* where the difference

between maximal and minimal growth typically occurs over a drug concentration not

exceeding 8-fold (**Supplemental Figure 2**). As this range is so broad, we consider it useful to

measure the concentration at which growth is inhibited by 50% (herein referred to as IC50 [34]

to distinguish from MIC50 which is an MIC determination made of populations). For

MFIG001, the IC50 for olorofim is 0.012 mg/L whereas for itraconazole its 0.65 mg/L [22].

As olorofim inhibits pyrimidine biosynthesis, it would be expected that the action of the drug

would be fully reversed by supplementing the media with an excess of exogenous pyrimidines

[7]. To confirm growth inhibition was due to directly targeting the pyrimidine biosynthesis

pathway, the MIC was determined with the addition of 10 mM uridine and 10 mM uracil

(**Figure 1b**). Under these conditions there was no observed reduction in *A. fumigatus* growth,

and at all olorofim concentrations the median OD_{600} did not fall below control levels indicating

that there are no significant off target effects of this drug.

To facilitate our understanding of how olorofim impacts the transcriptome of *A. fumigatus*,

RNA-seq analysis was performed. Upon olorofim exposure (1x MIC), a modest 185 genes and

41 genes respectivley were up- and downregulated (**Figure 1c**). Our expectation was that

several genes in the immediate pyrimidine biosynthesis pathway would be upregulated

however   only   the   multifunctional   carbamoyl-phosphate   synthase/aspartate

carbamoyltransferase (PyrABCN, AFUB_077330) enzyme, which is upstream of DHODH and

converts carbamoyl-P to N-carbamoyl-L-aspartate, was upregulated by Log2 fold >1. Instead,

genes associated with pathways that synthesise precursors of the pyrimidine biosynthetic

pathway were identified including oxaloacetate metabolism and glutamate biosynthesis

(**Figure 1d,e**). Genes associated with tyrosine metabolism; secondary metabolite biosynthesis,

glycolysis/gluconeogenesis and valine, leucine and isoleucine degradation were enriched

among downregulated genes (**Figure 1d,e**). Interestingly, several proteins involved in

ergosterol   biosynthesis   (HmgA,   Erg13a   and   Erg26)   were   differentially   expressed

(**Supplemental Data 1**).

Deletion of HapB, AreA, DevR and AcdX changes olorofim susceptibility.

In order to identify novel transcriptional regulators associated with differential olorofim

susceptibility, the COFUN transcription factor knockout (TFKO) library was screened against

olorofim at a concentration that reduces growth of the isogenic wildtype isolate (MFIG001) by

c 20% (0.002 mg/L). At this concentration we were able to identify strains that have the

potential to be resistant or hypersensitive (**Figure 2a**).

Three transcription factor null mutants (Δ*areA*, Δ*hapB* and Δ*devR*) showed reproducible

increased relative fitness in the presence of olorofim and elevated MICs compared to MFIG001

(**Figure 2c, d and e**). Remarkably, two of these mutants (Δ*areA* and Δ*hapB*) are also resistant

to the azole class of antifungals. Loss of AreA, a transcription factor that has a global role in

activating expression of genes involved in nitrogen acquisition and processing [35] or loss of

HapB, which along with HapC and HapE, comprise the CCAAT Binding Complex (CBC) [36]

resulted in a 2-fold increase in MIC to olorofim when compared to the isotype control

MFIG001; IC50 values for these strains were 0.04 mg/L (4-fold increase) and 0.07 mg/L (8

fold-increase), respectively (**Figures 2c and 2e**). The Δ*devR* mutant showed a significant

reduction in susceptibility to olorofim at concentrations ranging from 0.008 mg/L to 0.06 mg/L

(MIC) and had an IC50 of 0.025 mg/L (**Figure 2d**). Although the MIC for this strain increased

to >0.125 mg/L most spores did not germinate at this concentration making this phenotype

reminiscent of tolerance **(Supplemental Figure 1)**.

One isolate (ΔAFUB_056620, Δ*acdX*) showed a significant increase in sensitivity to olorofim

and had an MIC of 0.03 mg/L and a IC50 of 0.006 mg/L, 2-fold lower than *A. fumigatus*

MFIG001 (**Figure 2b**). The *acdX* gene encodes a 612 amino acid transcription factor which, a

SMART domain search showed, contains six WD40 repeat units but no other functional

domains. A reciprocal BLAST of the AFUB_056620 protein sequence found a match to the

Saccharomyces cerevisiae transcription factor Spt8. However, the proteins only share 44%

identity of the entire protein sequence. In *S. cerevisiae*, Spt8 forms part of the SAGA (Spt-

Ada-Gcn5-acetyltransferase) complex [37] which is known to act as a transcriptional activator

under several stress conditions. While the orthologue of AcdX in other fungi generally contains

six WD40 domains, in species such as *N. crassa* and *A. terreus* only five domains are present,

however the significance of this is unclear. In *A. nidulans* AcdX has been described to be

functional in the SAGA complex and is involved in repressing genes in in acetate metabolism

and has a regulatory role in the proline metabolic pathway [38].

Transcription factor mutants with altered suseptibility to olorofim have defects in nitrogen

assimilation.

Further phenotypic analysis of the null mutants with differential susceptibility to olorofim

revealed that all had growth defects on *Aspergillus* Complete Medium (ACM) (**Figure 3a and**

b) and *Aspergillus* Minimal Medium (AMM) (**Figure 3a and c**). The *hapB, devR, areA* and

acdX null mutants showed a reduction of radial growth on ACM of 28%, 22%, 12% and 24%

respectively when compared to the isotype control. On AMM, the *hapB* mutant showed

increase radial growth (58%) however, colony growth was more diffuse than the isotype strain

(**Figure 3a and c**). As olorofim inhibits DHODH, which acts within the pyrimidine

biosynthetic pathway, and we observed that olorofim results in transcriptional repositioning of

several pathways linked to nitrogen processing we hypothesised that these growth defects could

be nitrogen source dependent. Indeed, the radial growth defect of strains were reversed by

differential supplementation (Δ*devR* by proline*;* Δ*acdX* by glutamine or proline and Δ*areA* by

glutamine). Furthermore, we observed an interesting correlation between radial growth on the

different nitrogen containing media and olorofim sensitivity. While the radial growth of the

isotype isolate differed little between nitrogen sources (**Figure 3d and Supplemental Figure**

3) the strains lacking the ability to effectively utilise nitrate as a sole nitrogen source were all

less susceptible to olorofim. Taken together, this suggests that defects in nitrogen utilisation in

these transcription factor null mutants could be linked to their olorofim susceptibility.

Changes in susceptibility to olorofim in Δ*devR* **and** Δ*acdX* **mutants** ***are caused by opposing***

regulation of pathways preceding pyrimidine biosynthesis

In order to characterise the basis of differential olorofim susceptibility in the Δ*devR* and Δ*acdX*

mutants the transcriptomes of these two mutants were compared to the wild-type

(**Supplemental Data 2, Supplemental Figure 4**). In the absence of olorofim 510 and 137

genes were respectively downregulated and upregulated in the Δ*devR* isolate while 212 were

downregulated and 194 upregulated upon olorofim exposure. In the absence of olorofim,

notable enriched functional categories included downregulation of genes involved in tyrosine

metabolism and an upregulation of genes involved in the biosynthesis of branched chain amino

acids and metabolism of arginine and proline, the latter of which was also seen under olorofim

exposure (**Figure 4a**). A detailed pathway analysis of genes involved in the conversion of

metabolites towards L-glutamate and through to orotate revealed that proline uptake and

degradation were upregulated in the *devR* null mutant (**Figure 4c**). Other pathways that

contribute to orotate precursors were also significantly upregulated, notably the nitrate

assimilation pathway (NAP [*crnA, niaD, niiA*]), and glutamate, glutamine and carbomyl-P

synthesis. Pathways that compete with orotidine biosynthesis for L-glutamate were not

differentially regulated in any of the assessed mutants (**Supplemental Figure 5).** Our

transcriptional data therefore suggests that nitrogen metabolism maybe altered in this strain in

ways that favor the generation of precursors for orotate biosynthesis and hence could explain

the reduced sensitivity of *devR* null mutant to olorofim.

The transcriptome of the olorofim hypersensitive Δ*acdX* mutant also revealed that proline and

arginine metabolism were upregulated compared to the wild-type however genes involved in

the NAP and glutamate, glutamine and carbomyl-P synthesis pathways were downregulated

suggesting that AcdX and DevR have directly opposing functions on these linked pathways

(**Figure 4d**) and providing further evidence to suggest that regulation of these pathways is

important for olorofim sensitivity.

Our transcriptomic data led us to assess the effect of pyrimidine pathway precursors on

olorofim susceptibility in the transcripton factor null mutants (**Supplemental Figure 6)**. Using

nitrate as the most abundant, but not sole nitrogen source increases susceptibility of *A.*

fumigatus MFIG001 to olorofim. Interestingly this increase in susceptibility is lost in the

Δ*devR*, perhaps as a consequence of the upregulation of the NAP in this mutant. In keeping

with this, the resistance phenoptype observed with the loss of AreA, a regulator that activates

the NAP, is lost. The hypersensitive phenotype in the *acdX* null mutant was exaggerated when

arginine and proline were used as sole nitrogen sources. These data, combined with results

from our transcriptomic analysis suggest that modification of environmental nitrogen sources

and or dysregulation of nitrogen metabolism is directly linked to changes in olorofim

sensitivity.

A clinical isolate with an hapE allele known to contribute to azole resistance does not affect

olorofim sensitivity.

HapB is a member of the heterotrimeric CBC complex along with HapE and HapC. We have

previously shown that this complex negatively regulates sterol biosynthesis and loss of any

member of this complex leads to azole resistance in *A. fumigatus* [22]. As the Δ*hapB* mutant

exhibited an olorofim resistance phenotype, we assessed the olorofim susceptibility for a

clinical isolate (V067-36) which has non-synonomous SNP in the *hapE* gene (resulting in a

P88L substitution) and which has been demonstrated to affect azole susceptibility. The MIC of

this isolate and an isogenic isolate with a wild-type *hapE* allele from the same patient were

identical for olorofim. To further confirm that the *hapE*^{P88L} allele did not impact olorofim

susceptibility, we reconstructed this point mutation in the MFIG001 background. While a

Δ*hapB* mutant showed decreased susceptibility to both olorofim and itraconazole, the

MFIG001 *hapE^{P88L}* strain only showed decreased susceptibility to itraconazole (**Figure 5**).

The action of the azoles and olorofim are antagonistic.

As we have identified two genes whose loss of function can result in cross resistance, we

assessed if both drugs when used in combination would be synergistic against the respective

mutants. We first evaluated the potential synergism of both compounds against the isotype

control MFIG001. To our surprise given the distinct mechanisms of action of the orotomides

and the azoles, we observed a clear antagonism between itraconazole and olorofim. The

antagonism also persisted in the Δ*hapB,* the V067-36 isolate and the *hapE^{P88L}* isolate (**Figure**

5). We evaluated our previously published RNA-seq data to assess if itraconazole increased

expression of genes involved in olorofim biosynthesis however contrary to our expectations,

many of the genes in the pyrimidine biosynthetic pathway were highly downregulated, even

when itraconazole was used at low levels **(Supplemental Data 1)**.

We assessed if the antagonism persisted in multiple genetic backgrounds and to an alternative

azole (voriconazole). Consistent with our previous observations the wild-type isolate CEA10,

which is the parental strain from which MFIG001 derived, the ATCC46645 isolate and an azole

resistant isolate with the TR_{34} L98H allele isolated from a patient, all showed an antagonistic

relationship between olorofim and voriconazole (**Figure 6**). Interestingly our data suggests that

the antagonism is predominantly unidirectional, with the azoles inhibiting the action of

olorofim in both solid and liquid media for all strains.

Discussion

Olorofim is a novel antifungal, currently in phase 3 clinical trials. It has a broad spectrum of activity against most moulds and acts by inhibiting the pyrimidine biosynthetic pathway through disruption of DHODH activity [7]. Our preliminary analysis of the inhibitory effects of olorofim revealed that the MIC and the IC50 were separated over a relatively large concentration range (5-fold). This contrasts with what is seen with itraconazole and other azoles where this concentration spread is typically 2-fold. The clinical implication of this finding remains unclear, however it is likely that olorofim will support clearance of an infection at doses well below the MIC. At these lower concentrations however, exposure to drug will be imparting selective pressure and has the potential to induce the production of mutagenic precursors that may drive the emergence of resistance as has been shown for several antibiotics [39]. As with other anti-infectives that act by inhibiting a single biological target there is clear potential for emergence of resistance. Understanding these mechanisms will provide a framework for development of diagnostics to detect resistance rapidly in the clinic.

Our previous survey of azole sensitivity in the *A. fumigatus* COFUN transcription factor knockout library [22] revealed 6 null mutants that had decreased sensitivity (ranging from 4 to 6-fold increase in MIC compared to the isogenic control) and 6 had increased sensitivity (4 to 8-fold decrease in MIC) to itraconazole. Here our screen revealed that only 1 mutant (Δ*acdX*) showed increased sensitivity while 3 showed decreased sensitivity (Δ*hapB*, Δ*devR*, Δ*areA*) to olorofim and the changes in sensitivity in these isolates were less extreme than seen for the azoles. It is unsurprising, given the mechanism of action of olorofim, that the transcription factors that we have identified in this screen either have well defined roles in regulating nitrogen utilisation or have been linked to this function in our study.

What is remarkable however given the distinct mechanisms of actions of the two compound classes, loss of function of either of AreA and HapB results in cross-class resistance to both

the azoles and orotomides. HapB is a member of the heterotrimeric CCAAT-binding complex

(CBC) and alongside HapC and HapE regulates the expression of over a third of the genome

[40] including several genes involved in ergosterol biosynthesis. The *hapB* null displayed the

highest levels of resistance to olorofim and was able to germinate at 0.12 mg/L, which is 8-

fold higher than the parental isolate but within the concentration range needed for clinical

utility. In *A. nidulans* AreA is a positive regulator of many genes that are required for utilisation

of nitrogen sources other than glutamate or ammonia [41] with loss of function resulting in an

inability to utilise amongst other nitrogen sources, nitrate, nitrite, uric acid and many amino

acids [42]. Reassuringly however, drug concentrations in animal models are tolerated well

above the increased MIC levels of the null mutants identified in this screen. Dosing 8 mg/kg at

8 hour intervals in mice results in peak serum levels of 2.5-3 mg/L [43]. Olorofim can be

tolerated at doses as high as 30 mg/kg intravenously, giving scope for higher drug levels *in*

vivo if required. In cynomolgus monkeys a single oral dose of olorofim resulted in peak levels

of 0.605-0.914 mg/L in serum for female and male animals, respectively [44].

Recently a clinical isolate, V067-36, has been described that harbours a mutant HapE allele

(HapE[P88L]) which directly contributes to azole resistance [24]. Even though this mutation

appears to affect the way the CBC interacts with its gDNA target, independent of other co-

factors [40, 45, 46] our data shows that this allele does not affect susceptibility of *A. fumigatus*

to olorofim and is therefore less important at regulating a response to the orotomides. It is also

noteworthy that to date, no further documented cases of azole resistance have been linked to

CBC function and even though AreA is partially dispensable for virulence in *A. fumigatus* [41],

no cases of clinical resistance have been linked to AreA function. Therefore, although

concerning, the cross-resistance phenotypes we have observed may not have major clinical

significance. If olorofim is ultimately licenced for use, monitoring for co-resistance before

switching therapy would be a sensible precaution.

Our studies have shown there is a clear antagonism of the azoles and olorofim. Whilst

concerning the antagonism is only evident when relatively low levels of both drugs are used.

It is interesting to note that the TR$_{34}$ L98H isolate used in this study has reduced susceptibility

to olorofim when compared to the CEA10 isolate and the antagonism drives the MIC above

0.5 mg/L. If these drugs are to be used in combination in a clinical setting, careful evaluation

of respective drug levels at the site of infection to ensure sufficient concentration of drug to

avoid antagonism would be sensible. The consequences of using azoles and olorofim in

combination for treatment of strains harbouring the TR$_{34}$ L98H allele also needs further

evaluation.

In summary, we have explored the mechanism behind olorofim susceptibility through

a systematic analysis of the COFUN transcription factor null library. All the mutants we

identified that had altered sensitivity to olorofim have associated defects in nitrogen

metabolism and two of these mutants Δ*devR* and Δ*acdX* regulate genes involved in metabolic

pathways immediately upstream of the pyrimidine pathway potentially leading to a differential

flux of metabolites into this pathway. Although we have identified two transcription factors,

the CBC and AreA, that regulate cross resistance to both the azoles and olorofim we are

reassured that the only recognised incidence of clinical azole resistance linked to CBC function

(HapE P88L) does not lead to olorofim resistance.

References

1.   Bongomin, F., et al., *Global and Multi-National Prevalence of Fungal Diseases-*
Estimate Precision. J Fungi (Basel), 2017. **3**(4).

2.   Patterson, T.F., et al., *Practice Guidelines for the Diagnosis and Management of*
Aspergillosis: 2016 Update by the Infectious Diseases Society of America. Clin Infect
Dis, 2016. **63**(4): p. e1-e60.

3.   Snelders, E., et al., *Possible environmental origin of resistance of Aspergillus fumigatus*
to medical triazoles. Appl Environ Microbiol, 2009. **75**(12): p. 4053-7.

4.   van Rhijn, N. and M. Bromley, *The Consequences of Our Changing Environment on*
Life Threatening and Debilitating Fungal Diseases in Humans. Journal of Fungi, 2021.
7(5): p. 367.

5.   Maertens, J.A., et al., *Isavuconazole versus voriconazole for primary treatment of*
invasive mould disease caused by Aspergillus and other filamentous fungi (SECURE):
a phase 3, randomised-controlled, non-inferiority trial. Lancet, 2016. **387**(10020): p.
760-9.

6.   Rauseo, A.M., et al., *Hope on the Horizon: Novel Fungal Treatments in Development.*
Open Forum Infect Dis, 2020. **7**(2): p. ofaa016.

7.   Oliver, J.D., et al., *F901318 represents a novel class of antifungal drug that inhibits*
dihydroorotate dehydrogenase. Proc Natl Acad Sci U S A, 2016. **113**(45): p. 12809-
12814.

8.   Hope, W.W., et al., *Pharmacodynamics of the Orotomides against Aspergillus*
fumigatus: New Opportunities for Treatment of Multidrug-Resistant Fungal Disease.
mBio, 2017. **8**(4).

9.   Boschi, D., et al., *Dihydroorotate dehydrogenase inhibitors in anti-infective drug*
research. Eur J Med Chem, 2019. **183**: p. 111681.

10. du Pre, S., et al., *The Dynamic Influence of Olorofim (F901318) on the Cell*
Morphology and Organization of Living Cells of Aspergillus fumigatus. J Fungi
(Basel), 2020. **6**(2).

11. du Pre, S., et al., *Effect of the Novel Antifungal Drug F901318 (Olorofim) on Growth*
and Viability of Aspergillus fumigatus. Antimicrob Agents Chemother, 2018. **62**(8).

12. Biswas, C., et al., *In vitro activity of the novel antifungal compound F901318 against*
Australian Scedosporium and Lomentospora fungi. Med Mycol, 2018. **56**(8): p. 1050-
1054.

13. Kirchhoff, L., et al., *Antibiofilm activity of antifungal drugs, including the novel drug*
olorofim, against Lomentospora prolificans. J Antimicrob Chemother, 2020.

14. Lim, W., et al., *Madurella mycetomatis, the main causative agent of eumycetoma, is*
highly susceptible to olorofim. J Antimicrob Chemother, 2020. **75**(4): p. 936-941.

15. Seyedmousavi, S., et al., *Efficacy of Olorofim (F901318) against Aspergillus*
fumigatus, A. nidulans, and A. tanneri in Murine Models of Profound Neutropenia and
Chronic Granulomatous Disease. Antimicrob Agents Chemother, 2019. **63**(6).

16. Wiederhold, N.P., D. Law, and M. Birch, *Dihydroorotate dehydrogenase inhibitor*
F901318 has potent in vitro activity against Scedosporium species and Lomentospora
prolificans. J Antimicrob Chemother, 2017. **72**(7): p. 1977-1980.

17. Wiederhold, N.P., et al., *The Orotomide Olorofim Is Efficacious in an Experimental*
Model of Central Nervous System Coccidioidomycosis. Antimicrob Agents Chemother,
2018. **62**(9).

18. Lackner, M., et al., *Dihydroorotate dehydrogenase inhibitor olorofim exhibits*
promising activity against all clinically relevant species within Aspergillus section
Terrei. J Antimicrob Chemother, 2018. **73**(11): p. 3068-3073.

19.   Buil, J.B., et al., *In vitro activity of the novel antifungal compound F901318 against*
difficult-to-treat Aspergillus isolates. J Antimicrob Chemother, 2017. **72**(9): p. 2548-
2552.

20.   Rivero-Menendez, O., M. Cuenca-Estrella, and A. Alastruey-Izquierdo, *In vitro activity*
of olorofim (F901318) against clinical isolates of cryptic species of Aspergillus by
EUCAST and CLSI methodologies. J Antimicrob Chemother, 2019. **74**(6): p. 1586-
1590.

21.   Jorgensen, K.M., et al., *EUCAST Determination of Olorofim (F901318) Susceptibility*
of Mold Species, Method Validation, and MICs. Antimicrob Agents Chemother, 2018.
62(8).

22.   Furukawa, T., et al., *The negative cofactor 2 complex is a key regulator of drug*
resistance in Aspergillus fumigatus. Nat Commun, 2020. **11**(1): p. 427.

23.   Bertuzzi, M., et al., *On the lineage of Aspergillus fumigatus isolates in common*
laboratory use. Med Mycol, 2020.

24.   Camps, S.M., et al., *Discovery of a HapE mutation that causes azole resistance in*
Aspergillus fumigatus through whole genome sequencing and sexual crossing. PLoS
One, 2012. **7**(11): p. e50034.

25.   van Rhijn, N., et al., *Development of a marker-free mutagenesis system using CRISPR-*
Cas9 in the pathogenic mould Aspergillus fumigatus. Fungal Genetics and Biology,
2020. **145**: p. 103479.

26.   Rodriguez-Tudela, J.L., et al., *Epidemiological cutoffs and cross-resistance to azole*
drugs in Aspergillus fumigatus. Antimicrob Agents Chemother, 2008. **52**(7): p. 2468-
72.

27.   Berman, J. and D.J. Krysan, *Drug resistance and tolerance in fungi.* Nat Rev Microbiol,
2020. **18**(6): p. 319-331.

28.   Barratt, R.W., G.B. Johnson, and W.N. Ogata, *Wild-type and mutant stocks of*

526   *Aspergillus nidulans.* Genetics, 1965. **52**(1): p. 233-46.

527   29.   Bolger, A.M., M. Lohse, and B. Usadel, *Trimmomatic: a flexible trimmer for Illumina*

528   *sequence data.* Bioinformatics, 2014. **30**(15): p. 2114-20.

529   30.   Howe, K.L., et al., *Ensembl Genomes 2020-enabling non-vertebrate genomic research.*

530   Nucleic Acids Res, 2020. **48**(D1): p. D689-D695.

531   31.   Love, M.I., W. Huber, and S. Anders, *Moderated estimation of fold change and*

532   *dispersion for RNA-seq data with DESeq2.* Genome Biol, 2014. **15**(12): p. 550.

533   32.   Priebe, S., et al., *FungiFun2: a comprehensive online resource for systematic analysis*

534   *of gene lists from fungal species.* Bioinformatics, 2015. **31**(3): p. 445-6.

535   33.   Odds, F.C., *Synergy, antagonism, and what the chequerboard puts between them.*

536   Journal of Antimicrobial Chemotherapy, 2003. **52**(1): p. 1-1.

537   34.   Soothill, J.S., R. Ward, and A.J. Girling, *The IC50: an exactly defined measure of*

538   *antibiotic sensitivity.* J Antimicrob Chemother, 1992. **29**(2): p. 137-9.

539   35.   Amaar, Y.G. and M.M. Moore, *Mapping of the nitrate-assimilation gene cluster (crnA-*

540   *niiA-niaD) and characterization of the nitrite reductase gene (niiA) in the opportunistic*

541   *fungal pathogen Aspergillus fumigatus.* Curr Genet, 1998. **33**(3): p. 206-15.

542   36.   Hortschansky, P., et al., *Deciphering the combinatorial DNA-binding code of the*

543   *CCAAT-binding complex and the iron-regulatory basic region leucine zipper (bZIP)*

544   *transcription factor HapX.* J Biol Chem, 2015. **290**(10): p. 6058-70.

545   37.   Sermwittayawong, D. and S. Tan, *SAGA binds TBP via its Spt8 subunit in competition*

546   *with DNA: implications for TBP recruitment.* EMBO J, 2006. **25**(16): p. 3791-800.

547   38.   Georgakopoulos, P., R.A. Lockington, and J.M. Kelly, *SAGA complex components and*

548   *acetate repression in Aspergillus nidulans.* G3 (Bethesda), 2012. **2**(11): p. 1357-67.

39.   Kohanski, M.A., M.A. DePristo, and J.J. Collins, *Sublethal antibiotic treatment leads*
to multidrug resistance via radical-induced mutagenesis. Mol Cell, 2010. **37**(3): p. 311-
20.

40.   Gsaller, F., et al., *Sterol Biosynthesis and Azole Tolerance Is Governed by the Opposing*
Actions of SrbA and the CCAAT Binding Complex. PLoS Pathog, 2016. **12**(7): p.
e1005775.

41.   Hensel, M., et al., *The role of the Aspergillus fumigatus areA gene in invasive*
pulmonary aspergillosis. Mol Gen Genet, 1998. **258**(5): p. 553-7.

42.   Arst, H.N. and D.J. Cove, *Nitrogen metabolite repression in Aspergillus nidulans.*
Molecular and General Genetics MGG, 1973. **126**(2): p. 111-141.

43.   Hope, W.W., et al., *Pharmacodynamics of the orotomides against Aspergillus*
fumigatus: new opportunities for treatment of multidrug-resistant fungal disease.
MBio, 2017. **8**(4).

44.   Law, D., et al. *Pharmacokinetics of the Novel Antifungal Agent F901318 in Mice, Rats*
and Cynomolgus monkey. in *Fifty-fifth Interscience Conference on Antimicrobial*
Agents and Chemotherapy. San Diego, California. 2015.

45.   Furukawa, T., et al., *The fungal CCAAT-binding complex and HapX display highly*
variable but evolutionary conserved synergetic promoter-specific DNA recognition.
Nucleic acids research, 2020. **48**(7): p. 3567-3590.

46.   Hortschansky, P., et al., *The CCAAT-binding complex (CBC) in Aspergillus species.*
Biochim Biophys Acta, 2017. **1860**(5): p. 560-570.

Figure 1: The response of *A. fumigatus* to olorofim. (a) MIC of *A. fumigatus* MFIG001 to

olorofim as measured by OD_{600} by EUCAST methodology. (b) Addition of 10mM uracil and

10mM uridine reverses the action of olorofim to *A. fumigatus* MFIG001 (n=3). (c) Volcano

plot of RNA-seq of *A. fumigatus* A1160p+ exposed to olorofim. 185 genes (blue dots) and 41

genes (red dots) were considered downregulated and upregulated, respectively (>2-fold

differentially regulated, p<0.05). (d) KEGG pathways that are enriched within differentially

regulated genes, blue categories are associated with downregulated genes, red with upregulated

genes. (e) Interactions of proteins involved in response to olorofim as determined by

StringsDB. Proteins derived from upregulated transcripts are in red, downregulated in blue.

Figure 2: Olorofim susceptibility screening of the COFUN transcription factor knockout

library. (a) Relative fitness of each individual strain was assessed by normalising to fitness in

non-drug condition (n=3). TF null mutants that are of particular interest are highlighted. (b-e)

MIC to olorofim of the TF null mutants, (b) for Δ*acdX*, (c) for Δ*hapB*, (d) for Δ*devR*, (e) for

Δ*areA*, as determined by OD_{600} (n=3). Statistical difference was assessed by Two-way

ANOVA with Sidaks multiple comparison test (* p<0.05, ** p<0.01, *** p<0.001, ***

p<0.001).

Figure 3: Phenotypic evaluation of TF null mutants. (a) 500 spores of TF null mutants and

MFIG001+ were spotted on Aspergillus Complete Medium and Aspergillus Minimal Medium

and incubated for 48 hours at 37º Celsius. (b-c) Radial growth of TF null mutants and A1160p+

on ACM (b) and AMM (c), after 72 hours at 37º Celsius (n=3) (d) TF null mutants spotted on

AMM supplemented with 50 mM ammonium tartrate, 10 mM sodium nitrate, 10 mM L-

glutamine, 10 mM urea or 10 mM L-proline (n=3) Statistical difference was assessed by two-

way ANOVA with Dunn's correction (p-values < 0.05 are shown).

Figure 4: RNA-seq analysis of Δ*devR* and Δ*acdX* exposed to olorofim. (a) KEGG pathways

enriched for down- (blue) or upregulated (red) genes in RPMI-1640 or upon olorofim exposure

for ∆*devR* compared to *A. fumigatus* A1160p+. (b) KEGG pathways enriched for down- (blue)

or upregulated (red) genes in RPMI-1640 or upon olorofim exposure for ∆*acdX* compared to

A. fumigatus A1160p+. (c) Detailed analysis of genes involved in pathways upstream of and

including the pyrimidine pathway. The target of olorofim, DHODH, is highlighted. Blue is

more than 1-fold downregulated, yellow more than 1-fold upregulated; red is more than 5-fold

upregulated. The right of each box is associated with ∆*acdX*, left with ∆*devR*. (d) Heatmap of

genes involved in the pyrimidine pathway and component upstream of this pathway.

Figure 5: Checkerboard assays for CBC complex associated strains. Checkerboard assays

were performed (n=3) by using a double dilution series of itraconazole (16-0.06 mg/L) along

the X-axis and olorofim (0.5-0.003 mg/L) along the Y-axis. The ∆*hapB* and hapE^{P88L} isolates

are in the MFIG001 background, the V067-36 isolate is a clinical isolate containing the P88L

point mutation within HapE. The FICI is shown in the right top corner of each individual plot

to show the antagonistic effect. Values greater than 4 indicate antagonism.

Figure 6: The antagonistic effect between voriconazole and olorofim. (a) Checkerboard

assays were performed (n=2) by using a double dilution series of voriconazole (16-0.06 mg/L)

along the X-axis and olorofim (0.500-0.007 mg/L) along the Y-axis. Two wild-type isolates

(CEA10 and ATCC46645) were assessed and an azole resistant TR$_{34}$ L98H isolate. The FICI

is shown in the right top corner of each individual plot to show the antagonistic effect. (b)

Representative images of disk assays of the CEA10, ATCC46645 and TR$_{34}$ L98H isolates to

voriconazole and olorofim. A clear disruption of the olorofim halo can be observed proximal

to the voriconazole halo.

Supplemental Figure 1: Microscopical evaluation of *A. fumigatus* MFIG001 and

transcription factor null mutants upon olorofim exposure. Representative images were

taken from the MIC 96-well plates by stereomicroscope. Maximum concentration at which

growth is observed is highlighted in black for each strain.

Supplemental Figure 2: Determination of IC50 for itraconazole. MIC to olorofim in RPMI-

1640 was determined according to EUCAST methodology for *A. fumigatus* MFIG001. OD600

was measured after 48 hours to determine growth quantitatively.

Supplemental Figure 3: Images of nitrogen spot tests of TF null mutants. Growth of TF

null mutants and wild-type was assessed on AMM supplemented with 50 mM ammonium

tartrate, 10 mM sodium nitrate, 10 mM L-glutamine, 10 mM urea or 10 mM L-proline (n=3).

Images were taken after 72 hours at 37 Celsius.

Supplemental Figure 4: Volcano plots of Δ*acdX* and Δ*devR* RNA-seq data. Volcano plots

showing $\log_2$(Fold-Change) compared to A1160p+ on the X-axis and $-\log_{10}$(False Discovery

Rate) from DESeq2 on the Y-axis. Genes considered differentially regulated are coloured in

red, and number of genes differentially regulated are noted in each respective corner of volcano

plots.

Supplemental Figure 5: Pathways utilising L-glutamate. Additional pathways utilising L-

glutamate. Genes involved in these reactions are shown, none of these were differentially

regulated in Δ*acdX* or Δ*devR*.

Supplemental Figure 6: Olorofim MICs in supplemental RPMI-1640. MICs according to

EUCAST methodology in RPMI-1640 supplemented with 20 mM arginine, 10 mM nitrate, 20

mM proline or 50 mM glutamine. Addition of nitrate changed Olorofim susceptibility by 2-

fold for all strains except Δ*devR*.

Supplemental Table 1: **Oligos and crRNA used in this study.**

(a)

| | MFIG001 | Δ*hapB* | Δ*devR* | Δ*acdX* | Δ*areA* |

ACM

AMM

(b) ACM

Radial growth (mm)

MFIG001 Δ*hapB* Δ*devR* Δ*acdX* Δ*areA*

(c) AMM

Radial growth (mm)

MFIG001 Δ*hapB* Δ*devR* Δ*acdX* Δ*areA*

(d)

Radial Growth (mm)

0.0001
<0.0001
0.0001
<0.0001
0.0137
0.0001
0.0004
0.0302
<0.0001
<0.0001
<0.0001

Ammonium Nitrate Glutamine Urea Proline

MFIG001 Δ*hapB* Δ*devR* Δ*acdX* Δ*areA*

(a)

(b)

Oligo name	Oligo sequence (5 to 3')
hapE_gene_Fw	AGCTCTATCCAGCTGAAGTG
hapE_gene_Rv	AGATTTCTCTCGCTGACTGA
hapE_Fw	TCAAGGGCTACAAGGCACAG
hapE_Rv	AGATATCGCAGCCTTTCGCA
hapE_Seq	CCGCGACATACTAACGACCT
hapE_392_revcom	ATAGGTCGTTAGTATGTCGC

655