Interrogating Genomic Data in the Phylogenetic Placement of Treeshrews Reveals

Potential Sources of Conflict.

Authors

Alexander Knyshov[1,*], Yana Hrytsenko[1,2], Robert Literman[1,3], and Rachel S. Schwartz[1]

1. Department of Biological Sciences, University of Rhode Island, South Kingstown,

RI, USA

2. Department of Computer Science and Statistics, University of Rhode Island, South

Kingstown, RI, USA

3. Current address: Center for Food Safety and Applied Nutrition, Office of Regulatory

Science, U.S. Food and Drug Administration, College Park, MD, USA

* Corresponding author, e-mail: aknyshov@uri.edu

Abstract

The position of some taxa on the Tree of Life remains controversial despite the increase

in genomic data used to infer phylogenies. While analyzing large datasets alleviates stochastic

errors, it does not prevent systematic errors in inference, caused by both biological (e.g.,

incomplete lineage sorting, hybridization) and methodological (e.g., incorrect modeling,

erroneous orthology assessments) factors. In this study, we systematically investigated factors

that could result in these controversies, using the treeshrew (Scandentia, Mammalia) as a study

case. Recent studies have narrowed the phylogenetic position of treeshrews to three competing

hypotheses: sister to primates and flying lemurs (Primatomorpha), sister to rodents and

lagomorphs (Glires), or sister to a clade comprising all of these. We sampled 50 mammal species

including three treeshrews, a selection of taxa from the potential sister groups, and outgroups.

Using a large diverse set of loci, we assessed support for the alternative phylogenetic position of

treeshrews. The results suggest that the data has statistical support for two hypotheses for the

placements of treeshrews, sister to Primatomorpha and to Primatomorpha+Glires. While we

observe differences in properties of loci of different types (e.g., CDS, intron, etc.) with respect to

the strength of the signal, the support for any particular topology is not dependent on the

properties of the data. Rather, we show that the method of phylogenetic signal assessment, as

well as whether the signal is measured using the full dataset or only loci with the strongest

signal, impact the results much more.

Key words: phylogenomics, contentious relationship, incomplete lineage sorting,

likelihood, site concordance factor, sisrs

37   In the past two decades, DNA sequence data has become the main source of information

38 for reconstructing phylogenetic relationships. When only a few short gene regions were available

39 for tree inference, stochastic error was often large, resulting in some areas of phylogenies with

40 poor resolution and low branch support. However, the size of phylogenetic datasets has increased

41 dramatically with the advent of High Throughput Sequencing, making it possible to overcome

42 stochastic errors in small datasets and establish some evolutionary relationships with greater

43 certainty (e.g., (Dunn et al. 2008). At the same time, large datasets on their own may still not be

44 able to resolve all phylogenetic relationships due to systematic error (Rodríguez-Ezpeleta et al.

45 2007). Reanalyses of the same genome-scale datasets with different tools or different data

46 subsets, have resulted in strong support for alternate hypotheses (Chakrabarty et al. 2017;

47 Esselstyn et al. 2017; Reddy et al. 2017). Some relationships remain controversial despite

48 extensive work (e.g. early-diverging animals (Jékely and Budd 2021)], birds (Reddy et al. 2017;

49 Braun et al. 2019)], and placental mammals [Smith et al. 2015, (Murphy et al. 2021)]).

50 Moreover, phylogenetic conflict may be obscured by the inability of traditional support metrics

51 to capture variation in very large datasets, as both non-parametric bootstrap and posterior

52 probability metrics tend to provide strong support for conflicting topologies (Kumar et al. 2012;

53 Salichos and Rokas 2013).

54   Apparent systematic discordance may be caused by an array of factors. Biological

55 processes such as incomplete lineage sorting (ILS) or hybridization are predicted to be most

56 impactful for areas of the tree of life with rapid diversification (i.e., short internodes) (Rannala et

57 al. 2020). Systematic issues in data processing (such as extensive contamination or homology

58 inference errors) can bias the phylogenetic results of any dataset and are predicted to have a

59 stronger impact on less informative datasets applied to controversial relationships (Simion et al.

2020). Addressing these issues can explain conflicts and resolve phylogenies, as seen in taxa

ranging from tunicates to turtles (Brown and Thomson 2017; Simion et al. 2020). Finally, a lack

of information in the data may cause essentially arbitrary topological resolution depending on

minor factors influencing particular analyses (e.g., (Simion et al. 2020).

In order to disentangle these factors, the array of phylogenetic support methods have

evolved to elucidate the presence, strength, and possible cause of phylogenetic conflicts (Simon

2020). For example, the SH-aLRT test (Guindon et al. 2010) assesses likelihood differences

between the maximum likelihood (ML) tree and the nearest neighbor interchange (NNI)

alternatives for each node, making it a good generic test for informativeness of data. To

interrogate locus-specific phylogenetic support, likelihoods (Lee and Hugall 2003; Shen et al.

2017) or marginal likelihoods (Brown and Thomson 2017) can be compared across genes or

partitions of interest. This approach provides a gauge of the variation in support among loci and

has proven effective in determining the homogeneity of the conflict, and is especially useful at

identifying overly impactful outliers (Brown and Thomson 2017; Springer and Gatesy 2018).

The site concordance factor (sCF) (Minh et al. 2020a) provides a quartet-based support measure

in a parsimony framework, and can be used both on the entire dataset and on a locus-by-locus

basis. However, despite the availability of the aforementioned techniques, for many empirical

datasets and contentious phylogenetic relationships the extent and potential causes of the

conflicts remain elusive.

The position of the treeshrews (Scandentia) is an example of one of the glaring

uncertainties in the mammal tree of life (Hallström and Janke 2010; Murphy et al. 2021).

Treeshrews are part of the Euarchontoglires, a lineage that also includes rodents, lagomorphs,

primates, and colugos. Historically, morphological analyses supported grouping treeshrews with

colugos (Dermoptera) (O'Leary et al. 2013); however, genomic data now overwhelmingly

support the sister-group relationship between Dermoptera and Primates (Primatomorpha)

(Meredith et al. 2011; Esselstyn et al. 2017). One of the currently prevailing hypotheses is that of

treeshrews as the sister to the Primatomorpha (Fig. 1) (Hallström and Janke 2008; McCormack et

al. 2012; Zoonomia Consortium 2020). This placement, forming the clade Euarchonta, has been

recovered by numerous analyses, although typically with relatively low confidence.

Alternatively, several phylogenomic analyses have supported the sister-group relationship of

treeshrews with rodents and lagomorphs (Glires; Fig. 1) (Tarver et al. 2016; Liu et al. 2017; Du

et al. 2019). Finally, one of the most recent molecular analyses (Fig. 1) found that

Primatomorpha and Glires themselves form a monophyletic clade, with treeshrews equidistantly

related to both (Esselstyn et al. 2017). Analyses recovering each of the topologies have varied in

their sampling of species and higher order taxa, the number of sites analyzed, and methodology

(Song et al. 2012; Esselstyn et al. 2017; Vachaspati and Warnow 2018), meaning that both

biological and methodological factors may be at play when considering the conflicting

taxonomic groupings. Thus, with copious genomic data across mammals and a limited set of

hypotheses, the placement of treeshrews presents an ideal case study for phylogenetic conflict

interrogation (Zoonomia Consortium 2020).

Fig. 1. The taxon set and the phylogeny used for evaluations. Different resolutions of node 1 are

shown on the right. Node 2 indicates the most recent common ancestor of Primates, this node was used as

a control node and was compared to the two alternative nearest neighbor interchanges around the

subtending branch.

Here we examine the support for the alternative hypotheses of the placement of

treeshrews on the mammal tree of life as a case study to understand why we continue to struggle

to understand some evolutionary relationships despite genome-scale data. Using millions of

variable orthologous sites spread across mammal genomes, we assessed the support for each

hypothesis in both parsimony and maximum likelihood frameworks. We observe that the

majority of loci are only weakly informative with respect to the phylogenetic placement of

treeshrews. The hypothesis receiving the most support depends on both the method of

phylogenetic signal assessment and which loci are selected. These results first suggest that

treeshrews, Glires, and Primatomorpha might have diversified in rapid succession, and it is

challenging to unambiguously distinguish the order of splitting. However, more importantly, the

impact of locus selection on phylogenetic results even with very large datasets presents some

concern. We suggest that the approach we take here to examine support for alternative

hypotheses and gauge the amount of phylogenetic and non-phylogenetic signal, and its potential

sources, should be extended to examine other contentious relationships.

Material and Methods

Taxon Selection

In order to comprehensively interrogate the conflict in the phylogenetic placement of the

treeshrews, we sampled taxa from this group and all major lineages surrounding its most likely

position (Primates, Dermoptera, Rodentia, Lagomorpha), as well as five outgroup taxa from

Laurasiatheria (Fig. 1). Sampling as many diverse representatives of each group as the publicly

available data allow should help to more accurately detect saturating substitutions by breaking

long branches and thus improve the signal-to-noise ratio (Baurain et al. 2007; Philippe et al.

2011). Thus we attempted to sample every family with publicly available short read whole

genome sequencing data. A total of 50 taxa were included in the dataset (Fig. 1, Table S1).

Raw Reads

Raw Illumina reads were downloaded from the European Nucleotide Archive FTP server

(accession numbers are listed in Table S1). The mean number of paired-end (PE) reads per

sample was 955,859,108 (for details see the info in Table S1). Read quality was assessed with

FastQC (Andrews 2010).

Obtaining Orthologous Loci

136   Putatively orthologous loci were obtained through the SISRS v2.1 pipeline (Schwartz et

137 al. 2015; Harkins et al. 2016; Literman and Schwartz 2021; Literman et al. 2022). Briefly, reads

138 were trimmed using BBDuk (Bushnell 2021) and subsampled using BBMap v38.81 suite scripts

139 (Bushnell 2021) to 10X (0.2X each) assuming an average taxon genome size of 3.5 Gbps.

140 Subsampled reads were assembled in Ray v2.3.1 (Boisvert et al. 2010) to produce a composite

141 genome as the basis of identifying orthologs. The composite assembly was used as a reference

142 for mapping all taxon reads with Bowtie 2 v2.3.5.1 (Langmead and Salzberg 2012; Langmead et

143 al. 2019).

144   A new step was implemented in SISRS to produce a consensus sequence for each locus

145 (based on the composite genome assembly) for each taxon using the Bowtie 2 aligned data. Sites

146 were called conservatively; sites with depth lower than 3 for the most frequent allele were called

147 as N. Additionally, heterozygous sites were kept track of and consensus sequences with

148 heterozygosity over 1% were discarded, as in the initial tests we empirically determined this to

149 be indicative of improper read alignment. The procedure to produce alignment-based consensus

150 outlined above was chosen over existing solutions (i.e., bcftools consensus), as the latter

151 increases the chances of reference sequence sites propagating into final taxon sequences when

152 coverage or mapping quality are low. This default behavior is undesirable for the SISRS pipeline

153 when analyzing very divergent species, as the reference composite genome sequence is likely to

154 differ significantly from any particular taxon sequence.

155   Loci were filtered to those containing at least 50% of the taxa, at least one taxon from

156 each of the five major lineages and the outgroup, and to have no more than two-thirds of the

157 sequence missing for each taxon. After filtering, the loci were aligned using MAFFT v7.475

158 (Katoh and Standley 2013) with the --auto flag.

We filtered potentially misaligned or paralogous data using the Branch Length

Correlation (BLC) metric (Simion et al. 2017), which identifies loci whose gene trees contain tip

branch lengths least correlated with the concatenation-based tree. Excessively long branches in

gene trees may be indicative of alignment or paralogy problems, potentially leading to

systematically biased results (Simion et al. 2017). Because the dataset was too large to

concatenate all loci, we estimated the concatenation-based tree by analyzing concatenated

alignments of 1000 arbitrary loci each in IQ-TREE v2.1.2 with all default parameters except -m

GTR+G -bb 1000 -alrt 1000. We then averaged the branch lengths of the obtained trees to

produce a set of concatenation-based branch lengths. We reconstructed a gene tree for each locus

using IQ-TREE v2.1.2 with all default parameters except -alrt 1000 -m GTR+G (Minh et al.

2020b). We then used the lm function in R to fit the linear model and compute the correlation

coefficient, and only retained loci with R-squared more than 0.25.

Annotation

We hypothesized that locus type (e.g. whether sites fall within coding sequences, introns,

etc.) may be associated with the phylogenetic signal for each of the alternative hypotheses, as a

similar association has been shown for the overall strength of the signal (Literman and Schwartz

2021). Thus, we annotated the obtained SISRS loci using the human genome as a reference.

Human sequences were extracted from the above generated SISRS loci and BLASTN 2.9.0+

(Altschul et al. 1990; Camacho et al. 2009) was used to align these sequences to the human

genome assembly GRCh38.p13 (Schneider et al. 2017). A custom script was used to filter the

BLASTN output table and convert it into a BED file. Coordinates of the resulting alignments on

the human genome chromosomes were intersected (Quinlan and Hall 2010) with the GFF file

containing annotations. Another custom script was used to process the annotated BED file and

produce a table with a proportion of each type of genomic feature for each locus. We used the

following feature types: CDS (coding sequence), UTR (untranslated region, defined as a

combination of gene, mRNA, and exon, but no CDS), intron (defined as a combination of gene

and mRNA, but no exon), lncRNA (long non-coding RNA), pseudogene, other (other features),

and unannotated sequence. For the purposes of subsequent analyses, each SISRS locus was

assigned to only one of the types of features based on which type covered the highest proportion

of the locus.

Assessment of Locus Properties

We hypothesized that in addition to the annotation type, other properties of loci may be

correlated with the support for particular phylogenetic hypotheses (Mongiardino Koch 2021).

QUAST v5.0.2 (Gurevich et al. 2013) was used to assess the basic properties of the assembly, so

that their covariation with phylogenetic signal can be investigated. For each locus, we scored the

number of taxa with data, locus length, proportion of missing data, the proportion of variable and

parsimony informative sites, and GC content using AMAS (Borowiec 2016). The number of taxa

and proportion of missing data quantify different levels of missing data, shown to be impactful

when non-randomly distributed within phylogenetic datasets (Xi et al. 2016). The locus length

and proportions of variable and parsimony informative sites are associated with the number of

sites available for model estimates and for resolving relationships (Shen et al. 2016; Steenwyk et

al. 2020). Quantifying locus GC content is helpful in determining skewed base composition,

which might be indicative of GC-biased gene conversion (Duret and Galtier 2009), which has

been shown to impact phylogenetic signal (Romiguier et al. 2013).

Good fit of a substitution model allows more accurate discrimination between

synapomorphic and convergent substitutions (Philippe et al. 2011), thus we assessed the absolute

fit of a GTR+G model to each locus using PhyloMAd (Duchêne et al. 2017, 2018a, 2018b). The

GTR+G model was chosen for all loci as it is one of the most generic substitution models and

could be fit to a wide diversity of loci (Sumner et al. 2012; Abadi et al. 2019). As a metric of the

adequacy of the model, we used the Mahalanobis distance (Mahalanobis 1936; Drummond and

Suchard 2008), which serves as a summary statistic in PhyloMAd. Specifically, we recorded the

number of standard deviations of the predictive distribution (SDPD) between the mean of the

distribution and the test statistic for the empirical data (Duchêne et al. 2017). The SDPD is

expected to be closer to 0 for loci with a good model fit and be further away for loci with a poor

model fit. Because minimization of the substitution saturation levels has been shown to improve

the phylogenetic signal (Philippe et al. 2011), we assessed saturation in our dataset. Following

(Borowiec et al. 2015), we compared corrected (TN93) and uncorrected (raw) pairwise distances

in R 4.1.0 (R Core Team 2021) with the package ape 5.5 (Paradis and Schliep 2019). A linear

model was fit to the data to estimate the slope of the regression line, with higher values

corresponding to lower relative saturation. We performed a principal component analysis on the

locus features using R with the ade4 v1.7-17 package (Thioulouse et al. 1997). The results of the

analysis were used to assess the correlation between different locus properties and ensure the

independence of the chosen metrics.

222   *Phylogenetic Signal Assessment*

223   To assess overall phylogenetic signal, we first obtained an estimate of the phylogeny

based on the complete dataset. We performed a coalescent analysis in SVDQuartets (Chifman

and Kubatko 2014) as implemented in PAUP* Version 4.0a, build 168 (Swofford 2003). Only

parsimony-informative sites were included and we ran 1000 bootstrap replicates to assess the

support for each of the treeshrew placements.

To investigate the phylogenetic signal on a locus-by-locus basis, we used two metrics as

proxies for the signal: a log-likelihood difference (Lee and Hugall 2003; Shen et al. 2017)

between competing hypotheses (delta log-likelihood, ΔlnL), and a difference in sCF (Minh et al.

2020a) between competing hypotheses. In addition to measuring the phylogenetic signal for the

competing hypotheses of the treeshrew placement, we also assessed the signal for the monophyly

of primates in order to compare the support for a contentious node with the support for a well-

supported split of similar age.

Because assessment of both ΔlnL and sCF required predetermined topologies, we used

the mammalian phylogenetic tree of (Armstrong et al. 2020), which was based on the taxa with

genomic data available through the Zoonomia project (Zoonomia Consortium 2020) and

included 49 of the 50 species we sampled. We pruned the tree to retain only the taxa that we

sampled and, since *Tupaia belangeri* was not included in the original tree, we grafted this species

as the sister to *Tupaia chinensis* based on (Fan et al. 2013). This topology is well-supported and

uncontroversial with regard to the major lineages (Kumar et al. 2017). Because the resulting tree

represented only one possible treeshrew placement (Scandentia+Primatomorpha), we obtained

the two alternative placements using the phangorn package v2.7.0 (Schliep 2011) in R. These

three topologies were used for subsequent evaluations (Fig. 1). To examine the phylogenetic

signal for primates, we used the original tree (Fig. 1) with monophyletic primates and two trees

with the alternative topologies around the primates branch (i.e. Haplorhini or Strepsirrhini sister

to Dermoptera).

Each locus was fit to each phylogenetic hypothesis in IQ-TREE v2.1.2 (Minh et al.

2020b) in two different ways (individually and as part of a larger concatenated alignment) and

log-likelihoods were recorded. The approach of fitting loci individually was used to minimize the

probability of a single underlying tree assumption violation, which is very likely in the

concatenation framework (Rannala et al. 2020). On the other hand, because the loci were on

average very short, we hypothesized that the estimates of model parameters may be too

inaccurate compared to model estimates for longer alignments (Xia 2020). Thus we also assessed

the likelihood-based signal by fitting concatenated partitioned alignments to the alternative

topologies, which is a more common approach for conflict interrogation (Shen et al. 2017).

Because the total amount of sequence data exceeded the capabilities of the compute cluster

system (total length over 15Mbp, see the results), we made the concatenated alignment

computationally tractable: loci were arbitrarily binned into groups of 1000, and loci of each

group were concatenated. For concatenated alignments we exported per site log-likelihoods and

summed them for each included locus according to its coordinates in the alignment (Lee and

Hugall 2003; Shen et al. 2017). For both types of likelihood signal assessment we scaled locus

log-likelihood by locus length to estimate a mean per site log-likelihood and to avoid bias

associated with locus length (Walker et al. 2020).

To assess the overall support across the entire dataset we summed log-likelihoods per

locus to produce total log-likelihood for each topology. On a locus by locus basis, the strength of

the likelihood support is typically calculated between two competing hypotheses as the

difference in log-likelihood (ΔlnL), with positive and negative values representing the support

for one or another topology (Shen et al. 2017). Because we considered three competing

hypotheses, per site log-likelihood values for each topology were standardized to be at 0 for a

hypothesis with the lowest likelihood. Positive values for a given hypothesis thus indicate the

ΔlnL-based support for it compared to the least likely hypothesis. We classified loci into three

groups, based on topology, based on which loci had the highest log-likelihood. Loci having two

best resolutions or equal likelihood between the alternative resolutions were excluded from this

assessment.

In addition to assessing the overall distribution of ΔlnL, we separately analyzed the 1000

loci with the largest differences in fit between the alternative topologies, and by extension with a

strongest phylogenetic signal (the "outliers" dataset), as well as the full dataset with these loci

excluded (the "non-outliers" dataset).

We used IQ-TREE v2.1.2 to calculate sCF (Minh et al. 2020a). For the

Scandentia+Primatomorpha and Scandentia+Glires hypotheses we recorded the sCF of the

branch subtending the corresponding clades, while for the Scandentia+(Primatomorpha+Glires)

hypothesis we recorded the sCF for the branch subtending the Primatomorpha+Glires clade. In

addition, we assessed the correlation between sCF and ΔlnL on a locus-by-locus basis to assess

similarities between parsimony-based and model-based phylogenetic signals.

We assessed the significance of the difference in total log-likelihood across the dataset

for all competing hypotheses using the approximately unbiased (AU) test (Shimodaira 2002)

implemented in the R package scaleboot (Shimodaira 2008). The significance of the differences

in the numbers of loci favoring each topological resolution was assessed using Pearson's chi-

squared test (Pearson 1900; Agresti 2018), while Kruskal-Wallis One-Way ANOVA (Kruskal

and Wallis 1952) followed by pairwise Wilcoxon Rank-Sum Test (Wilcoxon 1945) with

Bonferroni correction were used to compare per locus ΔlnL and sCF distributions between the

competing topologies.

Interrelationship Between Phylogenetic Signal and Loci Properties

We tested for a correlation between values of each locus property, and the phylogenetic

signal towards particular topological resolutions of the treeshrews. For this question, we

analyzed the distribution of ΔlnL in groups of loci favoring each topology, as well as the

distribution of previously assessed locus properties in each group. We used Kruskal-Wallis

(KW) One-Way ANOVA (Kruskal and Wallis 1952) followed by pairwise Wilcoxon Rank-Sum

Test (Wilcoxon 1945) with Bonferroni correction to compare the distributions of the properties

of the loci favoring competing hypotheses, except for the locus type proportions, differences in

which were assessed using Pearson's chi-squared test.

All scripts for data analysis are available at:

https://github.com/AlexKnyshov/TreeshrewProject

Results

Orthologous Loci Generation

On average, 573,949,175 reads per sample (min 90,024,816, max 1,868,772,590, median

576,806,927) passed the trimming and were used for the SISRS pipeline. A total of 11,997,130

SISRS contigs containing 2,208,378,641bp were identified. The largest contig spanned

11,534bp, N50 was 189, L50 was 4,693,728. After filtering based on missing data criteria and

the BLC metric, a total of 83,055 loci containing 15,680,010bp were retained for subsequent

analyses.

Properties of Selected Loci

Results of the evaluations are listed in Table S2 and summarized in Table S3 and Figure

S1. The number of taxa varied from 25 (minimum passing filter) to 50, with a median of 36. The

average locus length was 189bp. Proportions of variable and parsimony informative sites as well

as percent missing were nearly normally distributed with means of 0.58, 0.33, and 22%

respectively. Mean GC content was 40.3%, although significant differences in GC content were

observed between some locus types (Fig. S2). On average, the estimated saturation of the SISRS

loci was low, with the mean slope of saturation curve equal to 0.78. The model fit SDPD formed

a bimodal distribution, and most loci had a relatively poor model fit (SDPD peak around 9.54),

while a smaller group of loci had better model fit (SDPD peak around 5.84). Since the behavior

of different test statistics implemented in PhyloMAd is an active area of research (Duchêne et al.

2018a), it is unclear how a poor fit with respect to the Mahalanobis distance-based SPDP relates

to the phylogenetic noise. Most features were orthogonal to each other in the PCA,

demonstrating a relatively high level of independence, while proportions of variable and

parsimony informative characters strongly covaried (Figure S3). Among the loci successfully

mapped to the reference and passing the filtering, types of genomic features were annotated in

the following proportions: CDS 14.49%, intron 32.84%, lncRNA 11.00%, pseudogene 0.23%,

UTR 4.92%, other 0.44%, and unannotated 26.04%.

Coalescence analysis results

The SVDQuartets inference on all (parsimony-informative) sites yielded a 100% support

for the Primatomorpha-sister hypothesis of treeshrew placement and a 100% support for

monophyletic primates. Nevertheless, the tree had a Robinson-Foulds distance of 6 (normalized

distance of 6.4%) to the reference tree. The differences were restricted to relationships within the

outgroup, the placement of Sciuromorpha within rodents, as well as relationships within

Lemuriformes in primates. Most of the tree branches had a 100% bootstrap support with the

exception of two branches within the outgroup, which received slightly lower support (99.2-

99.9%).

Phylogenetic Signal in the Treeshrew case

Likelihood-based signal for individually fit loci.— Summed log-likelihood of all loci

individually fit to each treeshrew placement topology was as follows: -117,594,213 for the

Primatomorpha sister, -117,594,082 for the Glires sister, and -117,590,171 for the

Primatomorpha+Glires sister (Fig. 2a). AU test results indicate that the Primatomorpha+Glires

sister hypothesis was significantly more likely (AU test p-value = 0). The number of loci

favoring the Primatomorpha+Glires sister hypothesis (N = 19622) was significantly higher (X-

squared = 1161.765, df = 2, p-value = 5.32e-253) than the number of loci favoring the other two

topologies (N = 14653 and 14047) (Fig. 2c). However, the likelihood-based signal of loci

favoring the Primatomorpha sister (median ΔlnL = 0.0039) was significantly higher than that of

the Glires sister (median ΔlnL = 0.0034, W = 107379002, p-value = 5.97e-10) or the

Primatomorpha+Glires sister (median ΔlnL = 0.0035, W = 151974436, p-value = 3.78e-19) (Fig.

2e). The signal distribution of the top 1000 loci ranked by their ΔlnL values was different, with

significantly more loci (X-squared = 98.942, df = 2, p-value = 3.27e-22) supporting the

Primatomorpha sister (N = 452) than the Primatomorpha+Glires sister (N = 351) (Figs 2c, 3a).

The likelihood-based signal of outlier loci was not significantly different between hypotheses

(KW ANOVA, X-squared = 0.12891, p-value = 0.9376) (Figs 2e, 3a). The ΔlnL distributions

and counts of favoring loci in the dataset with top 1000 outliers removed were similar to that of

the overall dataset, with the Primatomorpha+Glires favoring loci being prevalent (X-squared =

1166.801, df = 2, p-value = 4.29e-254) (Fig. 2c), but the Primatomorpha favoring loci having

stronger signal (median ΔlnL = 0.0035, W = 101073540, p-value = 1.68e-04) (Fig. 2e).

Fig. 2. Likelihood-based signal distribution shows ambiguous support for alternative hypotheses of

treeshrew placement. The left panel (a, c, and e) shows the results based on loci fit to the trees individually, while

the right panel (b, d, and f) shows the same based on loci fit in concatenation, with considerable differences in

phylogenetic signal between the two methods of assessment. Each panel features three groups of loci (all, with

outliers excluded, and just the outliers), with signal of outliers being different from that of the rest of the dataset.

Top row (a and b) shows summed likelihood across all loci for each hypothesis. Middle row (c and d) shows counts

of loci that favor (have the highest likelihood) for each of the hypotheses. Bottom row shows the average per-locus

likelihood difference between the most and the least likely hypotheses, horizontal lines and points show medians and

means respectively.

Likelihood-based signal for loci fit in concatenation.— Summed log-likelihood of loci fit

to each topology in concatenated alignments was -121,169,491 for Primatomorpha sister, -

121,170,580 for Glires sister, and -121,171,209 for Primatomorpha+Glires sister (Fig. 2b). Thus

in concatenation fit the Primatomorpha sister hypothesis was the most likely (AU test p-value =

0.003717603). This was also reflected in the numbers of loci favoring each of the hypotheses:

the number of loci favoring the Primatomorpha sister (N = 26261) was significantly higher (X-

squared = 7.865172, df = 2, p-value = 0.0196) than the number of loci favoring the alternatives

(N = 25808 and 25645) (Fig. 2d). However, the likelihood-based signal strength of loci favoring

the Primatomorpha+Glires sister (median ΔlnL = 0.0009) was higher than the signal strength for

the alternatives (median ΔlnL = 0.0007 and 0.0006, W = 304381022 and 299618847, and p-

values = 1.88e- 55 and 2.35e-104 respectively) (Fig. 2f). The signal distribution of the top 1000

loci ranked by their ΔlnL values resembled the overall dataset signal, with significantly more loci

(X-squared = 503.34, df = 2, p-value = 5.02e-110) favoring the Primatomorpha sister (N = 647)

than the Primatomorpha+Glires sister (N = 76) or the Glires sister (N = 277) (Figs 2d, 3b). The

likelihood-based signal strength of outlier loci was not significantly different between the two

hypotheses with the highest median signal (W = 97952, p-value = 0.074) (Figs 2f, 3b). The ΔlnL

distributions in the dataset with top 1000 outliers removed were similar to that of the overall

dataset, with the Primatomorpha+Glires sister loci having stronger signal (median ΔlnL =

0.0009, W = 297286484, p-value = 8.37e-68) (Fig. 2f). However, unlike the results of the dataset

with per locus likelihood-based signal, in concatenation there were no significant differences in

counts between groups of loci favoring alternative hypotheses when the top 1000 outliers were

removed (X-squared = 0.13502, df = 2, p-value = 0.935) (Fig. 2d).

Fig. 3. Likelihood support among loci with strong signal (referred to as outliers) for the three competing

hypotheses of treeshrew placement, with loci fit individually (a) or in concatenation (b). The figures demonstrate

that in both cases more outliers favor the Primatomorpha sister hypothesis, although a noticeable number of loci

favor the alternatives. Each panel shows the difference in likelihood (ΔlnL), obtained via fitting each locus to the

three competing hypotheses, and rescaled by the locus length. If a point is at the center or close to it, there is no or

little difference in likelihood between the topologies for a given locus thus low-likelihood data was omitted. Points

plotted further away from the center correspond to loci having higher likelihood under one or two topologies. Each

locus point is color coded by density value in that area of the plot.

sCF-based signal.— Differences in the distribution of per locus sCF in groups of loci

favoring alternative treeshrew topologies were significant (KW ANOVA, X-squared = 28.533, p-

value = 6.372e-07), however the highest median value was the same for two hypotheses, the

Glires sister and the Primatomorpha+Glires sister (median sCF = 40.04 in both groups, W =

347550702, p-value = 4.38e-1) (Fig. 4). Counts of loci favoring each of the alternative treeshrew

410   topologies were also significantly different (X-squared = 328.1939, df = 2, p-value = 5.41e-72),

411   but with more loci (N = 30082) favoring the Primatomorpha sister (Fig. 4).

413   Fig. 4. Site concordance factor (sCF) distribution shows weak support for the alternative hypotheses of

414   treeshrew placement, with the Primatomorpha sister having slightly higher counts of loci and all hypotheses having

415   nearly identical median per locus sCF. The left panel shows the distribution of per-locus sCF values, horizontal bars

416   and points show medians and means respectively, and the dashed line shows the majority threshold of sCF (Minh et

417   al 2020). The right panel shows the numbers of loci favoring each topology based on sCF value.

419   The relationship between ΔlnL and sCF for particular loci was positive but very weak

420   (Fig. S6) both in the per locus fit approach and in the concatenation approach (R-squared 0.07-

421   0.08, p-value < 2.2e-308) (Fig. S4). The same relationship was even more weak for the outlier

422   loci (R-squared 0.008 - 0.03, p-value < 2.2e-308) (Fig. S4).

424   *Phylogenetic signal in the Primates case*

425   *Likelihood-based signal for individually fit loci.—* Summed log-likelihood of all loci fit

individually was -117,594,244 for the control case of monophyletic primates, compared to -

117,695,132 and -117,697,088 for the two alternative NNIs (Fig. 5a). AU test results indicate

that primate monophyly was a significantly more likely hypothesis (AU test p-value = 0). The

number of loci favoring monophyletic primates (N = 40812) was significantly different (X-

squared = 28431.62, df = 2, p-value < 2.2e-308) from the numbers of loci favoring the two

alternative topologies (N = 11527 and 10400) (Fig. 5c). The likelihood-based signal of loci

favoring monophyletic primates (median ΔlnL = 0.019) was also significantly stronger (W =

316670653, p-value < 2.2e-308) than that of the loci favoring the alternatives (median ΔlnL =

0.006 and 0.007) (Fig. 5e). The signal distribution of the top 1000 loci ranked by their ΔlnL

values was different (X-squared = 1934.546, df = 2, p-value < 2.2e-308), with vast majority of

loci (N = 989) favoring the monophyletic primates over the two alternatives (N = 6 and 5) (Figs

5c, 6a). The ΔlnL distributions and counts of favoring loci in the dataset with top 1000 outliers

removed were similar to that of the overall dataset, favoring the monophyletic primates (ΔlnL: W

= 306389180, p-value < 2.2e-308; counts: X-squared = 27021.47, df = 2, p-value < 2.2e-308)

(Figs 5d, e). However, the likelihood-based signal of outlier loci was significantly stronger for an

alternative topology compared to the monophyletic primates (W = 818, p-value = 0.029),

highlighting a possibility for spurious results obtained from likelihood outliers (Figs 5e, 6a).

Fig. 5. Likelihood-based signal distribution mostly shows strong support for monophyletic primates. The

left panel (a, c, and e) shows the results based on loci fit to the trees individually, while the right panel (b, d, and f)

shows the same based on loci fit in concatenation, with minimal differences in phylogenetic signal between the two

methods of assessment. Each panel features three groups of loci (all, with outliers excluded, and just the outliers),

with almost all outliers favoring monophyletic primates. Top row (a and b) shows summed likelihood across all loci

for each hypothesis. Middle row (c and d) shows counts of loci that favor (have the highest likelihood) for each of

the hypotheses. Bottom row shows the average per-locus likelihood difference between the most and the least likely

hypotheses, horizontal lines and points show medians and means respectively.

Likelihood-based signal for loci fit in concatenation.— Summed log-likelihood of all loci

in concatenation was -121,169,473 for the monophyletic primates, compared to -121,258,158

and -121,258,201 for the two alternative NNIs (Fig. 5b). AU test results indicate that primates

monophyly was a significantly more likely hypothesis (AU p-value = 0). The number of loci

favoring monophyletic primates (N = 48412) was significantly higher (X-squared = 28892.28, df

= 2, p-value < 2.2e-308) than the numbers of loci favoring the two alternative topologies (N =

13531 and 16242) (Fig. 5d). The strength of the likelihood-based signal of loci favoring

monophyletic primates (median ΔlnL = 0.0095) was also significantly higher (W = 495031550,

p-value < 2.2e-308) than that of the loci favoring the alternatives (median ΔlnL = 0.0032 and

0.0030) (Fig. 5f). The signal distribution of the top 1000 loci ranked by their ΔlnL values was

different (X-squared = 1994.006, df = 2, p-value < 2.2e-308), with all except one locus favoring

the monophyletic primates over the two alternatives (Figs 5d, 6b). The likelihood-based signal of

outlier loci was not significantly different between hypotheses (KW ANOVA, X-squared =

0.096123, p-value = 0.7565) (Figs 5f, 6b). The ΔlnL distributions and counts of favoring loci in

the dataset with top 1000 outliers removed were similar to that of the overall dataset, favoring

the monophyletic primates (ΔlnL: W = 481514670, p-value < 2.2e-308; counts: X-squared =

27557.74, df = 2, p-value < 2.2e-308) (Figs 5d, f).

Fig. 6. Likelihood support among outliers for monophyletic primates, with loci fit individually (a) or in

concatenation (b). In both cases the majority of outliers favor the monophyly as opposed to the alternative

hypotheses. Each panel shows the difference in likelihood (ΔlnL), obtained via fitting each locus to the three

competing hypotheses, and rescaled by the locus length. If a point is at the center or close to it, there is no or little

difference in likelihood between the topologies for a given locus thus low-likelihood data was omitted. Points

plotted further away from the center correspond to loci having higher likelihood under one or two topologies. Each

locus point is color coded by density value in that area of the plot. The subtree in the center shows the location of the

node for which the alternative hypotheses were considered.

sCF-based signal.— Differences in the distribution of per locus sCF between the

monophyletic primates and two alternative NNIs were significant (W = 540094683, p-value <

2.2e-308), with the sCF-based signal of the loci favoring monophyletic primates (median sCF =

49.78) being stronger than that of loci favoring two alternative NNIs (median sCF = 41.22 and

40.69) (Fig. 7). The median sCF support for the monophyletic primates was higher than that for

any of the treeshrew topologies. Counts of loci favoring the monophyletic primates and

alternative topologies were also significantly different (X-squared = 15976.57, df = 2, p-value <

2.2e-308), with the majority of loci (N = 44549) supporting primate monophyly (Fig. 7). Thus,

sCF based phylogenetic signal supported primates monophyly much more strongly than any of

the competing treeshrew topologies.

Fig. 7. Site concordance factor distribution shows strong support for monophyletic primates. The left panel

shows the distribution of per-locus sCF values, horizontal bars and points show medians and means respectively,

and the dashed line shows the majority threshold of sCF (Minh et al 2020). The right panel shows the numbers of

loci favoring each topology based on sCF value.

The relationship between ΔlnL and sCF across all loci for the primates case (standardized

slope 0.43-0.47, R-squared 0.19-0.22, p-value < 2.2e-308) was stronger than in the case of

competing treeshrew topologies (Fig. S5). The same relationship for the outliers was weak (R-

squared 0.008-0.03, p-value < 2.2e-308) (Fig. S5).

Support for Treeshrew Relationships Using Loci Supporting Monophyletic Primates

Only 0.6 to 0.9% of top 1000 outlier loci overlapped between the Treeshrew and the

Primates analyses, suggesting that the signal in loci varies among nodes. When fit individually,

both loci that supported monophyly of primates and those that contradicted statistically favored

the Primatomorpha+Glires placement of treeshrew (X-squared = 802.4895 and 141.5346, p-

value = 5.52e-175 and 1.85e-31 respectively). Specifically, 42% of loci that favor primates'

monophyly favored the dominant Primatomorpha+Glires hypothesis, while only 38% of loci that

did not favor primates monophyly supported that treeshrew placement.

When fit in concatenation, the non-outlier loci that favor monophyletic primates loci

favored three competing hypotheses in roughly the same proportions (~33%, X-squared =

3.904185, df = 2, p-value = 0.142), similarly to the overall signal of non-outliers in

concatenation (see above). However, loci that rejected monophyly of primates had a slightly

higher but nevertheless significant proportion of support for the Primatomorpha sister (34%, X-

squared = 9.281649, df = 2, p-value = 0.00965).

Interrelationship Between Phylogenetic Signal and Loci Properties

We hypothesized that annotation and other properties of loci may be associated with the

signal for each of the competing topologies. When fit individually, loci favoring each of the

treeshrew placements had different distributions of most of the properties we assessed,

specifically between the loci favoring the Primatomorpha+Glires-sister and the other alternatives

(with the exception no differences in alignment length and model fit) (Fig. 8, Table S3). When fit

in concatenation, similarly, most properties had significant differences between treeshrew

placement topologies, although there was a less clear pattern of which properties were

statistically different between which hypotheses (Fig. 9, Table S3).

Fig. 8. Properties of the loci favoring each of the treeshrew phylogenetic hypotheses when loci fit to

alignments individually. Most properties have distributions that look very similar between competing treeshrew

topologies. Nevertheless distributions of all properties except for alignment length and model fit SDPD have

statistically significant differences. The Primatomorpha+Glires-sister hypothesis is the most frequently supported

across all types of loci.

The proportions of locus types in loci favoring all competing treeshrew topologies were

significant when loci were fit individually, while only the proportions between the

Primatomorpha-sister and Primatomorpha+Glires-sister were significantly different when loci

were fit in concatenation. The counts of loci favoring each treeshrew topology in each locus type

overall reflected the total dataset counts. When fit individually, most locus types favored the

Primatomorpha+Glires sister (with exception of "other" where the difference was not significant,

primarily due to small sample size) (Fig. 8). When fit in concatenation, only the "unannotated"

group had significant differences between the competing treeshrew topologies, with other locus

types having statistically indistinguishable counts of loci supporting the alternatives (Fig. 9).

Fig. 9. Properties of the loci favoring each of the treeshrew phylogenetic hypotheses when loci fit to

alignments in concatenation. Most properties have distributions that look very similar between competing treeshrew

topologies. Nevertheless distributions of all properties except for the number of taxa have statistically significant

differences between at least two hypotheses. Only among the unannotated loci there were statistically significant

prevalence of the Primatomorpha-sister favoring loci.

Discussion

Overall, the SISRS loci contained strong phylogenetic signal, as measured on an

uncontroversial split (Primates). Both metrics used as proxies for phylogenetic signal - ΔlnL and

sCF - showed that the overall dataset clearly favored the topology with monophyletic primates

over the two alternative NNIs. With respect to phylogenetic placement of treeshrews, despite

observing a strong support for the Primatomorpha-sister hypothesis in the coalescent analysis of

all parsimony informative sites, we found disagreements among different proxies for per-locus

phylogenetic signal. Additionally, as opposed to the Primates split case, the overall modest per-

site ΔlnL-based signal suggests that the majority of genomic markers contain only weak

phylogenetic signal for short branch splits, and some contain non-phylogenetic signal.

Phylogenetic Signal, the Competing Treeshrew Hypotheses, and Primate Monophyly

Overall, two topologies (Primatomorpha-sister and Primatomorpha+Glires-sister) were

supported across different analyses. Differences in which particular hypothesis was supported

were observed between likelihood-based and site concordance-based analyses, between two

types of likelihood-based signal assessment, whether selection was based on average per-locus

signal of loci or numbers of loci, as well as whether the entire dataset, only likelihood-based

outliers, or the dataset without outliers were considered (Fig. 10). The sCF based phylogenetic

signal was also ambiguous with respect to placement of treeshrews.

Fig. 10. Phylogenetic signal for competing hypotheses summarized over all metrics, highlighting

conflicting phylogenetic signal with respect to a contentious node (treeshrew placement) while showing a very

strong support for an uncontroversial split (primates monophyly). Color encodes the hypothesis supported by each

metric in each subset, gray represents no significant difference between the top two or all three hypotheses, values in

the cells are p-values.

When the likelihood-based signal was assessed individually per locus, the total signal

(summed lnL across all loci) as well as counts of supporting loci favored the

Primatomorpha+Glires-sister. However, per locus signal strength was higher in the group of loci

favoring the Primatomorpha-sister. Additionally, the total signal and counts in the strong signal

outliers subset also supported the Primatomorpha-sister. When the likelihood-based signal was

assessed in concatenation, the signal distribution changed: the total signal and counts were

unequivocal except for when the strong signal outliers were included, while per locus signal

strength was higher in the group of loci favoring the Primatomorpha+Glires-sister. In both types

of likelihood assessment we observed a discordance between loci that have stronger signal and

loci that have weaker signal (compare per locus signal strength and locus counts on the figure).

With respect to the sCF-based signal, the majority of the loci favored the Primatomorpha-sister,

although sCF-based signal strength for that hypothesis was not significantly higher than for the

others. Thus, despite that the results of coalescent analysis convey superficial uniformity of the

support for the Primatomorpha-sister hypothesis, there is a significant conflict between

individual loci in the dataset. Given that, we conclude that based on our data it is impossible to

unequivocally resolve treeshrew position.

The overall ambiguity of the signal with respect to the treeshrew placement was

considerably larger than for the case of primate monophyly, which is consistent with a notion of

rapid radiation between the major Euarchontoglires lineages (Zhou et al. 2015; Esselstyn et al.

2017). However, even for primate monophyly the support was never maximal (all loci), thus

showing detectable conflict even in the uncontroversial split. Similar results were found by

(Esselstyn et al. 2017) when examining the likelihood support for a well-established monophyly

of Glires. We thus argue that non-phylogenetic signal is noticeable when examining both the

well supported and the poorly supported nodes, however in the case of the poorly supported one

the relatively small amount of phylogenetic signal prevents us from unambiguously resolving the

topology.

Locus support (or lack of) for monophyletic primates did not impact the signal

distribution with respect to competing treeshrew topologies. Loci that favor monophyly of

primates constitute about two thirds of the overall dataset depending on the method of signal

assessment, and the distribution of the signal for treeshrew placement in this subset replicated

that in the overall (unfiltered) dataset. Thus based on our results it can be concluded that filtering

loci based on the support for an uncontroversial relationship would not necessarily improve the

phylogenetic signal to help resolve a contentious node.

Impact of Individual vs in Concatenation Assessment on Phylogenetic Signal

Our results show stark contrast between the two types of assessment of phylogenetic

signal, by fitting locus alignments to topologies individually, or as a part of concatenated

alignments. We hypothesize that poor model fit for short single-locus alignments may be one

reason why the results of likelihood assessments between the two methods differ (Xia 2020). The

SISRS loci in our dataset had a mean length of about 200bp, which for conserved loci is likely

insufficient to adequately estimate substitution model parameters. Thus the parameter estimates

were likely to have random noise, which might increase the likelihood for the alternative

topologies and decrease the likelihood for the true tree. The use of short loci has been criticized

previously on the grounds of poor model fit and resulting poor gene tree estimates (Bryant and

Hahn 2020; Rannala et al. 2020). While normally model choice and fit does not impact the

resulting topology as much as it impacts branch length estimates, model accuracy could be

critical for nodes that are difficult to resolve (Kelchner and Thomas 2007). This is well

illustrated in the case of the order of divergence of Porifera and Ctenophora, where using

different types of models results in reconstruction of two alternative topological hypotheses (Li

et al. 2021).

However, it is also possible that model parameters may be estimated incorrectly due to

concatenating loci. Model parameter estimation for concatenated datasets is commonly done

with the help of the tree reconstructed on the entire matrix (in the case of IQ-TREE, it is a

parsimony-based tree (Minh et al. 2020b)]). Such a procedure would violate the assumption of

multispecies coalescence (MSC) and potentially bias individual loci to support the predominant

topology, which they would otherwise reject (Walker et al. 2020). Longer concatenated loci have

been noted as violating the assumptions of MSC and loci across our large dataset are practically

guaranteed to violate the assumption of a single generating tree (Rannala et al. 2020). Recent

work has noted discordance between reconstructed gene trees for individual loci and the tree

supported by the same loci when part of a larger supermatrix (Walker et al. 2020).

Overall, it is worth keeping in mind both the issue of model fit in short loci leading to

erroneous support, as well as the issues of both model fit and lost signal in concatenated datasets,

as it is unclear which approach leads to a more accurate phylogenetic signal inference. We thus

conclude that both approaches should be used when interrogating phylogenetic conflicts.

Heterogeneity of Signal within the Dataset and Impacts of Strong Signal Outliers

Our results suggest that even a large dataset with loci relatively uniformly sampled across

gene types can have some heterogeneity of phylogenetic signal, with a group of loci with a

strong phylogenetic signal favoring a different topology compared to a group of loci with a weak

signal. The selected phylogenetic hypothesis would depend on the proportions of these two

groups of loci in a dataset. In this context, we suggest exercising caution with picking loci with

the top most strong phylogenetic signal, since the phylogenetic error might be different for such

subset and mislead the analysis. For many phylogenomic subsampling methods, loci are selected

based on alignments between a handful of reference genomes (Faircloth et al. 2012; Lemmon et

al. 2012), and are typically checked to reproduce a well-supported phylogeny as a control

measure in the design phase (Faircloth 2017). Our results show that such (non-random) filtering

could bias results away from the signal across the genome.

Signal Metrics Discordance on a Locus-by-locus Basis

Despite the congruence between mean ΔlnL and mean sCF in the entire dataset, there was

no strong correlation between the two metrics on a locus-by-locus basis. This is not surprising

given that sCF is computed in a parsimony framework and is not impacted by model parameter

misspecification. On the other hand, sCF is a quartet-based metric and can be stochastic when

the conservative nature of SISRS loci results in a small number of informative quartets (Minh et

al. 2020a). Interestingly, there was a relatively strong positive correlation between ΔlnL and sCF

for the loci supporting the monophyletic primates. We hypothesize that these loci, even despite

their short size, have enough signal to overcome stochasticity in model fit and/or quartet

sampling, and thus bring the two metrics to congruence. Consequently, that would mean that in

the cases with a strong ΔlnL-sCF signal discordance loci might have elevated levels of noise

compared to the phylogenetic signal. We hypothesize that model fit accuracy might explain some

of the differences between the two types of ΔlnL-based signal and the sCF-based signal

assessments.

Loci Properties and Signal for Treeshrew Placement

It has been shown that locus type (Literman and Schwartz 2021) and locus properties

(Mongiardino Koch 2021) are related to the overall amount of phylogenetic signal. However, we

did not observe any large differences in locus properties with respect to a particular conflicting

topology (see Figures). While minor but significant differences of some properties could be

predictive of the amount of signal and noise in the data, more sensitive methods would be needed

to tease this out. It is also possible that the signal for each of the topologies is randomly

distributed throughout the genomes independent of the locus properties, and any difference

found between groups of loci favoring competing hypotheses would bear no implication across

different datasets and phylogenetic problems.

Taxon Sampling and Signal for Treeshrew Placement

Taxon sampling has been shown previously to affect phylogenetic estimates (Rosenberg

and Kumar 2003; Nabhan and Sarkar 2012). In this study, Dermoptera, Lagomorpha, and

treeshrews were represented by only a few species each and were constrained to have at least one

species present in each locus. Thus the sampling of the diverse Primates and rodents primarily

determined the overall variance in taxon sampling, as well as the fractions of Primatomorpha and

Glires. While our taxon set was relatively skewed towards a high fraction of Glires (0.55),

similar fractions of Glires were included in prior analyses supporting all alternative hypotheses

of treeshrew placement. In the datasets recovering the Glires sister, the fraction of Glires ranged

between 0.39 and 0.44 (Song et al. 2012; Romiguier et al. 2013; Tarver et al. 2016; Liu et al.

2017), while in the datasets finding the Primatomorpha sister this fraction ranged between 0.31

and 0.5 (Hallström and Janke 2008; McCormack et al. 2012; Kumar et al. 2013; Esselstyn et al.

2017). Thus, although the high Glires sampling might be seen as a concern, our results are

consistent with results from lower Glires sampled datasets. However, it is possible that more

intricate properties of taxon sampling (such as presence or absence of particular species) are at

play here.

SISRS Data and Conflict Interrogation

Our results also contain several practical outcomes with regard to SISRS-based data and

the newly introduced orthologous locus extraction pipeline. The recovered SISRS loci in our

case are relatively conserved, which is unsurprising given the methodology of the pipeline.

Although SISRS alignments were considerably shorter (~200bp vs ~700bp (Esselstyn et al.

2017)]), the proportion of parsimony informative sites in our dataset was in the same range as in

some of the mammal UCE studies (McCormack et al. 2012; Esselstyn et al. 2017), amounting on

average to about 70 parsimony informative sites per alignment. Small locus size, however,

should not impact the results much since we did not use individual loci to produce and analyze

gene trees (apart from the BLC filtering). It is conceivable that the model parameter estimation

could be compromised when small loci fit to topologies individually. On the other hand, longer

loci are likely to violate the single underlying tree assumption. While it is unclear which option

balances the phylogenetic signal and noise better, we explored both individual fit and

concatenation fit approaches in detail. Additionally, we also used other metrics to estimate

phylogenetic signal, including the total likelihood summed across all sites of all loci, thus we

argue that short loci produced by SISRS do not pose any significant obstacle to the result

interpretation.

Given that SISRS loci contain considerable proportions of both coding and regulatory

elements, as well as a large amount of normally not used unannotated (intergenic) loci (Literman

and Schwartz 2021), SISRS datasets offer a good diversity of locus types. In contrast, most other

methods focus on subsets of particular locus types (i.e., UCE, protein-coding sequences, introns)

or small subsets of randomly targeted markers (i.e., RADSeq methods), which have the potential

to bias results (Literman and Schwartz 2021).

Conclusion

Our work highlights the importance of per-locus phylogenetic signal interrogation (Smith

et al. 2015; Brown and Thomson 2017; Shen et al. 2017) as a valuable tool to examine support

for alternative phylogenetic relationships. While our initial phylogeny estimation supported the

treeshrew as the sister to Primatomorpha, this interrogation process revealed a quite pronounced

support for the Primatomorpha+Glires sister hypothesis as well. The methodology for conflict

interrogation utilized in this study allowed us to ascertain the overall magnitude of the conflict

and the amount of signal supporting each conflicting topology in a locus-by-locus framework.

The techniques allowed us to disentangle the phylogenetic signal of strong signal outliers from

that of the remainder of the dataset and discover that these can contain support for different

topologies. A non-random subsetting of strong signal outliers could thus bias results away from

the signal across the genome and mislead the phylogenetic inference. This suggests that filtering

loci (or selecting particular loci to sequence) should be done with caution due to the potential to

estimate a phylogeny that is not concordant with an estimate based on a more complete dataset.

Using locus-by-locus interrogation we also found discordance between the model-based and

parsimony-based methods, which we in part attribute to the adequacy of model fit to the data.

Overall, with a wealth of phylogenetic information provided by SISRS, we support two

alternate phylogenetic placements for treeshrews, even though our initially estimated phylogeny

strongly supported a single hypothesis. Two hypotheses, Primatomorpha-sister or

Primatomorpha+Glires-sister receive statistical support, depending on the method of

phylogenetic signal assessment, as well as whether the signal outlier loci were included or

excluded from the analysis. This observation, even with an extremely large DNA sequence

dataset, suggests that treeshrews, Glires, and Primatomorpha might have diversified in rapid

succession.

Acknowledgments

This research was funded by a grant to R. Schwartz from the National Science

Foundation (DBI-1942273). We thank members of the Schwartz lab, as well as three reviewers,

for valuable comments and suggestions. The High Performance Research Computing facility at

URI is acknowledged for providing the computational resources for the analyses in this

manuscript.

References

Abadi S., Azouri D., Pupko T., Mayrose I. 2019. Model selection may not be a mandatory step
for phylogeny reconstruction. Nat. Commun. 10:934.

Agresti A. 2018. An Introduction to Categorical Data Analysis. John Wiley & Sons.

Altschul S.F., Gish W., Miller W., Myers E.W., Lipman D.J. 1990. Basic local alignment search
tool. J. Mol. Biol. 215:403–410.

Andrews S. 2010. FastQC: A Quality Control Tool for High Throughput Sequence Data.
Available from http://www.bioinformatics.babraham.ac.uk/projects/fastqc/.

Armstrong J., Hickey G., Diekhans M., Fiddes I.T., Novak A.M., Deran A., Fang Q., Xie D.,
Feng S., Stiller J., Genereux D., Johnson J., Marinescu V.D., Alföldi J., Harris R.S.,
Lindblad-Toh K., Haussler D., Karlsson E., Jarvis E.D., Zhang G., Paten B. 2020.
Progressive Cactus is a multiple-genome aligner for the thousand-genome era. Nature.
587:246–251.

Baurain D., Brinkmann H., Philippe H. 2007. Lack of resolution in the animal phylogeny:
closely spaced cladogeneses or undetected systematic errors? Mol. Biol. Evol. 24:6–9.

Boisvert S., Laviolette F., Corbeil J. 2010. Ray: simultaneous assembly of reads from a mix of
high-throughput sequencing technologies. J. Comput. Biol. 17:1519–1533.

Borowiec M.L. 2016. AMAS: a fast tool for alignment manipulation and computing of summary
statistics. PeerJ. 4:e1660.

Borowiec M.L., Lee E.K., Chiu J.C., Plachetzki D.C. 2015. Extracting phylogenetic signal and
accounting for bias in whole-genome data sets supports the Ctenophora as sister to
remaining Metazoa. BMC Genomics. 16:987.

Braun E.L., Cracraft J., Houde P. 2019. Resolving the Avian Tree of Life from Top to Bottom:
The Promise and Potential Boundaries of the Phylogenomic Era. In: Kraus R.H.S., editor.
Avian Genomics in Ecology and Evolution: From the Lab into the Wild. Cham: Springer
International Publishing. p. 151–210.

Brown J.M., Thomson R.C. 2017. Bayes Factors Unmask Highly Variable Information Content,
Bias, and Extreme Influence in Phylogenomic Analyses. Syst. Biol. 66:517–530.

Bryant D., Hahn M.W. 2020. The Concatenation Question. In: Scornavacca, C., Delsuc, F., and

Galtier, N., editor. Phylogenetics in the Genomic Era. No commercial publisher | Authors
open access book. p. 3.4:1–3.4:23.

Bushnell B. 2021. BBTools. BBMap https://sourceforge. net/projects/bbmap/.

Camacho C., Coulouris G., Avagyan V., Ma N., Papadopoulos J., Bealer K., Madden T.L. 2009.
BLAST+: architecture and applications. BMC Bioinformatics. 10:421.

Chakrabarty P., Faircloth B.C., Alda F., Ludt W.B., Mcmahan C.D., Near T.J., Dornburg A.,
Albert J.S., Arroyave J., Stiassny M.L.J., Sorenson L., Alfaro M.E. 2017. Phylogenomic
Systematics of Ostariophysan Fishes: Ultraconserved Elements Support the Surprising Non-
Monophyly of Characiformes. Syst. Biol. 66:881–895.

Chifman J., Kubatko L. 2014. Quartet inference from SNP data under the coalescent model.
Bioinformatics. 30:3317–3324.

Drummond A.J., Suchard M.A. 2008. Fully Bayesian tests of neutrality using genealogical
summary statistics. BMC Genet. 9:68.

Duchêne D.A., Duchêne S., Ho S.Y.W. 2017. New Statistical Criteria Detect Phylogenetic Bias
Caused by Compositional Heterogeneity. Mol. Biol. Evol. 34:1529–1534.

Duchêne D.A., Duchêne S., Ho S.Y.W. 2018a. PhyloMAd: efficient assessment of
phylogenomic model adequacy. Bioinformatics. 34:2300–2301.

Duchêne D.A., Duchêne S., Ho S.Y.W. 2018b. Differences in Performance among Test Statistics
for Assessing Phylogenomic Model Adequacy. Genome Biol. Evol. 10:1375–1388.

Dunn C.W., Hejnol A., Matus D.Q., Pang K., Browne W.E., Smith S.A., Seaver E., Rouse G.W.,
Obst M., Edgecombe G.D., Sørensen M.V., Haddock S.H.D., Schmidt-Rhaesa A., Okusu
A., Kristensen R.M., Wheeler W.C., Martindale M.Q., Giribet G. 2008. Broad
phylogenomic sampling improves resolution of the animal tree of life. Nature. 452:745–
749.

Duret L., Galtier N. 2009. Biased gene conversion and the evolution of mammalian genomic
landscapes. Annu. Rev. Genomics Hum. Genet. 10:285–311.

Du Y., Wu S., Edwards S.V., Liu L. 2019. The effect of alignment uncertainty, substitution
models and priors in building and dating the mammal tree of life. BMC Evol. Biol. 19:203.

Esselstyn J.A., Oliveros C.H., Swanson M.T., Faircloth B.C. 2017. Investigating Difficult Nodes
in the Placental Mammal Tree with Expanded Taxon Sampling and Thousands of
Ultraconserved Elements. Genome Biol. Evol. 9:2308–2321.

Faircloth B.C. 2017. Identifying conserved genomic elements and designing universal bait sets to
enrich them. Methods Ecol. Evol. 8:1103–1112.

Faircloth B.C., McCormack J.E., Crawford N.G., Harvey M.G., Brumfield R.T., Glenn T.C.

2012. Ultraconserved elements anchor thousands of genetic markers spanning multiple
evolutionary timescales. Syst. Biol. 61:717–726.

Fan Y., Huang Z.-Y., Cao C.-C., Chen C.-S., Chen Y.-X., Fan D.-D., He J., Hou H.-L., Hu L.,
Hu X.-T., Jiang X.-T., Lai R., Lang Y.-S., Liang B., Liao S.-G., Mu D., Ma Y.-Y., Niu Y.-
Y., Sun X.-Q., Xia J.-Q., Xiao J., Xiong Z.-Q., Xu L., Yang L., Zhang Y., Zhao W., Zhao
X.-D., Zheng Y.-T., Zhou J.-M., Zhu Y.-B., Zhang G.-J., Wang J., Yao Y.-G. 2013.
Genome of the Chinese tree shrew. Nat. Commun. 4:1426.

Guindon S., Dufayard J.-F., Lefort V., Anisimova M., Hordijk W., Gascuel O. 2010. New
algorithms and methods to estimate maximum-likelihood phylogenies: assessing the
performance of PhyML 3.0. Syst. Biol. 59:307–321.

Gurevich A., Saveliev V., Vyahhi N., Tesler G. 2013. QUAST: quality assessment tool for
genome assemblies. Bioinformatics. 29:1072–1075.

Hallström B.M., Janke A. 2008. Resolution among major placental mammal interordinal
relationships with genome data imply that speciation influenced their earliest radiations.
BMC Evol. Biol. 8:162.

Hallström B.M., Janke A. 2010. Mammalian evolution may not be strictly bifurcating. Mol. Biol.
Evol. 27:2804–2816.

Harkins K.M., Schwartz R.S., Cartwright R.A., Stone A.C. 2016. Phylogenomic reconstruction
supports supercontinent origins for Leishmania. Infect. Genet. Evol. 38:101–109.

Jékely G., Budd G.E. 2021. Animal Phylogeny: Resolving the Slugfest of Ctenophores, Sponges
and Acoels? Curr. Biol. 31:R202–R204.

Katoh K., Standley D.M. 2013. MAFFT multiple sequence alignment software version 7:
improvements in performance and usability. Mol. Biol. Evol. 30:772–780.

Kelchner S.A., Thomas M.A. 2007. Model use in phylogenetics: nine key questions. Trends
Ecol. Evol. 22:87–94.

Kruskal, Wallis. 1952. Use of ranks in one-criterion variance analysis. J. Am. Stat. Assoc.
47:583–621.

Kumar S., Filipski A.J., Battistuzzi F.U., Kosakovsky Pond S.L., Tamura K. 2012. Statistics and
truth in phylogenomics. Mol. Biol. Evol. 29:457–472.

Kumar S., Stecher G., Suleski M., Hedges S.B. 2017. TimeTree: A Resource for Timelines,
Timetrees, and Divergence Times. Mol. Biol. Evol. 34:1812–1819.

Kumar V., Hallström B.M., Janke A. 2013. Coalescent-based genome analyses resolve the early
branches of the euarchontoglires. PLoS One. 8:e60019.

Langmead B., Salzberg S.L. 2012. Fast gapped-read alignment with Bowtie 2. Nat. Methods.

9:357–359.

Langmead B., Wilks C., Antonescu V., Charles R. 2019. Scaling read aligners to hundreds of
threads on general-purpose processors. Bioinformatics. 35:421–432.

Lee M.S.Y., Hugall A.F. 2003. Partitioned likelihood support and the evaluation of data set
conflict. Syst. Biol. 52:15–22.

Lemmon A.R., Emme S.A., Lemmon E.M. 2012. Anchored hybrid enrichment for massively
high-throughput phylogenomics. Syst. Biol. 61:727–744.

Literman R.A., Ott B.M., Wen J., Grauke L.J., Schwartz R.S., Handy S.M. 2022. Reference-free
discovery of nuclear SNPs permits accurate, sensitive identification of Carya (hickory)
species and hybrids. Appl. Plant Sci. 10:e11455.

Literman R., Schwartz R. 2021. Genome-scale profiling reveals noncoding loci carry higher
proportions of concordant data. Mol. Biol. Evol. 38:2306–2318.

Liu L., Zhang J., Rheindt F.E., Lei F., Qu Y., Wang Y., Zhang Y., Sullivan C., Nie W., Wang J.,
Yang F., Chen J., Edwards S.V., Meng J., Wu S. 2017. Genomic evidence reveals a
radiation of placental mammals uninterrupted by the KPg boundary. Proc. Natl. Acad. Sci.
U. S. A. 114:E7282–E7290.

Li Y., Shen X.-X., Evans B., Dunn C.W., Rokas A. 2021. Rooting the Animal Tree of Life.
Evolution. 38:4322–4333.

Mahalanobis P.C. 1936. On the generalized distance in statistics. Proceedings of the National
Institute of Science of India.:49–55.

McCormack J.E., Faircloth B.C., Crawford N.G., Gowaty P.A., Brumfield R.T., Glenn T.C.
2012. Ultraconserved elements are novel phylogenomic markers that resolve placental
mammal phylogeny when combined with species-tree analysis. Genome Res. 22:746–754.

Meredith R.W., Janečka J.E., Gatesy J., Ryder O.A., Fisher C.A., Teeling E.C., Goodbla A.,
Eizirik E., Simão T.L.L., Stadler T., Rabosky D.L., Honeycutt R.L., Flynn J.J., Ingram
C.M., Steiner C., Williams T.L., Robinson T.J., Burk-Herrick A., Westerman M., Ayoub
N.A., Springer M.S., Murphy W.J. 2011. Impacts of the Cretaceous Terrestrial Revolution
and KPg extinction on mammal diversification. Science. 334:521–524.

Minh B.Q., Hahn M.W., Lanfear R. 2020a. New Methods to Calculate Concordance Factors for
Phylogenomic Datasets. Mol. Biol. Evol. 37:2727–2733.

Minh B.Q., Schmidt H.A., Chernomor O., Schrempf D., Woodhams M.D., von Haeseler A.,
Lanfear R. 2020b. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic
Inference in the Genomic Era. Mol. Biol. Evol. 37:1530–1534.

Mongiardino Koch N. 2021. Phylogenomic subsampling and the search for phylogenetically
reliable loci. Mol. Biol. Evol. 38:4025–4038.

Murphy W.J., Foley N.M., Bredemeyer K.R., Gatesy J., Springer M.S. 2021. Phylogenomics and
the Genetic Architecture of the Placental Mammal Radiation. Annu Rev Anim Biosci. 9:29–
53.

Nabhan A.R., Sarkar I.N. 2012. The impact of taxon sampling on phylogenetic inference: a
review of two decades of controversy. Brief. Bioinform. 13:122–134.

O'Leary M.A., Bloch J.I., Flynn J.J., Gaudin T.J., Giallombardo A., Giannini N.P., Goldberg
S.L., Kraatz B.P., Luo Z.-X., Meng J., Ni X., Novacek M.J., Perini F.A., Randall Z.S.,
Rougier G.W., Sargis E.J., Silcox M.T., Simmons N.B., Spaulding M., Velazco P.M.,
Weksler M., Wible J.R., Cirranello A.L. 2013. The placental mammal ancestor and the
post-K-Pg radiation of placentals. Science. 339:662–667.

Paradis E., Schliep K. 2019. ape 5.0: an environment for modern phylogenetics and evolutionary
analyses in R. Bioinformatics. 35:526–528.

Pearson K. 1900. X. On the criterion that a given system of deviations from the probable in the
case of a correlated system of variables is such that it can be reasonably supposed to have
arisen from random sampling. Lond. Edinb. Dublin Philos. Mag. J. Sci. 50:157–175.

Philippe H., Brinkmann H., Lavrov D.V., Littlewood D.T.J., Manuel M., Wörheide G., Baurain
D. 2011. Resolving difficult phylogenetic questions: why more sequences are not enough.
PLoS Biol. 9:e1000602.

Quinlan A.R., Hall I.M. 2010. BEDTools: a flexible suite of utilities for comparing genomic
features. Bioinformatics. 26:841–842.

Rannala B., Edwards S.V., Leaché A., Yang Z. 2020. The Multi-species Coalescent Model and
Species Tree Inference. In: Scornavacca, C., Delsuc, F., and Galtier, N., editor.
Phylogenetics in the Genomic Era. No commercial publisher | Authors open access book. p.
3.3:1–3.3:21.

R Core Team. 2021. R: A Language and Environment for Statistical Computing. .

Reddy S., Kimball R.T., Pandey A., Hosner P.A., Braun M.J., Hackett S.J., Han K.-L.,
Harshman J., Huddleston C.J., Kingston S., Marks B.D., Miglia K.J., Moore W.S., Sheldon
F.H., Witt C.C., Yuri T., Braun E.L. 2017. Why Do Phylogenomic Data Sets Yield
Conflicting Trees? Data Type Influences the Avian Tree of Life more than Taxon Sampling.
Syst. Biol. 66:857–879.

Rodríguez-Ezpeleta N., Brinkmann H., Roure B., Lartillot N., Lang B.F., Philippe H. 2007.
Detecting and overcoming systematic errors in genome-scale phylogenies. Syst. Biol.
56:389–399.

Romiguier J., Ranwez V., Delsuc F., Galtier N., Douzery E.J.P. 2013. Less is more in
mammalian phylogenomics: AT-rich genes minimize tree conflicts and unravel the root of
placental mammals. Mol. Biol. Evol. 30:2134–2144.

Rosenberg M.S., Kumar S. 2003. Taxon sampling, bioinformatics, and phylogenomics. Syst.
Biol. 52:119–124.

Salichos L., Rokas A. 2013. Inferring ancient divergences requires genes with strong
phylogenetic signals. Nature. 497:327–331.

Schliep K.P. 2011. phangorn: phylogenetic analysis in R. Bioinformatics. 27:592–593.

Schneider V.A., Graves-Lindsay T., Howe K., Bouk N., Chen H.-C., Kitts P.A., Murphy T.D.,
Pruitt K.D., Thibaud-Nissen F., Albracht D., Fulton R.S., Kremitzki M., Magrini V.,
Markovic C., McGrath S., Steinberg K.M., Auger K., Chow W., Collins J., Harden G.,
Hubbard T., Pelan S., Simpson J.T., Threadgold G., Torrance J., Wood J.M., Clarke L.,
Koren S., Boitano M., Peluso P., Li H., Chin C.-S., Phillippy A.M., Durbin R., Wilson
R.K., Flicek P., Eichler E.E., Church D.M. 2017. Evaluation of GRCh38 and de novo
haploid genome assemblies demonstrates the enduring quality of the reference assembly.
Genome Res. 27:849–864.

Schwartz R.S., Harkins K.M., Stone A.C., Cartwright R.A. 2015. A composite genome approach
to identify phylogenetically informative data from next-generation sequencing. BMC
Bioinformatics. 16:193.

Shen X.-X., Hittinger C.T., Rokas A. 2017. Contentious relationships in phylogenomic studies
can be driven by a handful of genes. Nat Ecol Evol. 1:126.

Shen X.-X., Salichos L., Rokas A. 2016. A Genome-Scale Investigation of How Sequence,
Function, and Tree-Based Gene Properties Influence Phylogenetic Inference. Genome Biol.
Evol. 8:2565–2580.

Shimodaira H. 2002. An approximately unbiased test of phylogenetic tree selection. Syst. Biol.
51:492–508.

Shimodaira H. 2008. Testing regions with nonsmooth boundaries via multiscale bootstrap. J.
Stat. Plan. Inference. 138:1227–1241.

Simion P., Delsuc F., Philippe H. 2020. To What Extent Current Limits of Phylogenomics Can
Be Overcome? In: Scornavacca, C., Delsuc, F., and Galtier, N., editor. Phylogenetics in the
Genomic Era. No commercial publisher | Authors open access book. p. 2.1:1–2.1:34.

Simion P., Philippe H., Baurain D., Jager M., Richter D.J., Di Franco A., Roure B., Satoh N.,
Quéinnec É., Ereskovsky A., Lapébie P., Corre E., Delsuc F., King N., Wörheide G.,
Manuel M. 2017. A Large and Consistent Phylogenomic Dataset Supports Sponges as the
Sister Group to All Other Animals. Curr. Biol. 27:958–967.

Simon C. 2020. An Evolving View of Phylogenetic Support. Syst. Biol. 0:1–8.

Smith S.A., Moore M.J., Brown J.W., Yang Y. 2015. Analysis of phylogenomic datasets reveals
conflict, concordance, and gene duplications with examples from animals and plants. BMC
Evol. Biol. 15:150.

Song S., Liu L., Edwards S.V., Wu S. 2012. Resolving conflict in eutherian mammal phylogeny
using phylogenomics and the multispecies coalescent model. Proceedings of the National
Academy of Sciences. 109:14942–14947.

Springer M.S., Gatesy J. 2018. Pinniped Diphyly and Bat Triphyly: More Homology Errors
Drive Conflicts in the Mammalian Tree. J. Hered. 109:297–307.

Steenwyk J.L., Buida T.J. 3rd, Li Y., Shen X.-X., Rokas A. 2020. ClipKIT: A multiple sequence
alignment trimming software for accurate phylogenomic inference. PLoS Biol.
18:e3001007.

Sumner J.G., Jarvis P.D., Fernández-Sánchez J., Kaine B.T., Woodhams M.D., Holland B.R.
2012. Is the general time-reversible model bad for molecular phylogenetics? Syst. Biol.
61:1069–1074.

Swofford D.L. 2003. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods).
Version 4. .

Tarver J.E., Dos Reis M., Mirarab S., Moran R.J., Parker S., O'Reilly J.E., King B.L., O'Connell
M.J., Asher R.J., Warnow T., Peterson K.J., Donoghue P.C.J., Pisani D. 2016. The
Interrelationships of Placental Mammals and the Limits of Phylogenetic Inference. Genome
Biol. Evol. 8:330–344.

Thioulouse J., Chessel D., Doledec S., Olivier J.-M. 1997. ADE-4: a multivariate analysis and
graphical display software. Stat. Comput. 7:75–83.

Vachaspati P., Warnow T. 2018. SVDquest: Improving SVDquartets species tree estimation
using exact optimization within a constrained search space. Mol. Phylogenet. Evol.
124:122–136.

Walker J.F., Shen X.-X., Rokas A., Smith S.A., Moyroud E. 2020. Disentangling biological and
analytical factors that give rise to outlier genes in phylogenomic matrices.
bioRxiv.:2020.04.20.049999.

Wilcoxon F. 1945. Individual Comparisons by Ranking Methods. Biometrics Bulletin. 1:80–83.

Xia X. 2020. A Mathematical Primer of Molecular Phylogenetics. CRC Press.

Xi Z., Liu L., Davis C.C. 2016. The Impact of Missing Data on Species Tree Estimation. Mol.
Biol. Evol. 33:838–860.

Zhou X., Sun F., Xu S., Yang G., Li M. 2015. The position of tree shrews in the mammalian
tree: Comparing multi-gene analyses with phylogenomic results leaves monophyly of
Euarchonta doubtful. Integr. Zool. 10:186–198.

Zoonomia Consortium. 2020. A comparative genomics multitool for scientific discovery and
conservation. Nature. 587:240–245.

Supplementary information

Data available from the Dryad Digital Repository:

http://dx.doi.org/10.5061/dryad.[NNNN]

Supplementary Figures

Fig. S1. Properties of the evaluated SISRS loci. A set of properties on the left (blue)

contains size and information related features, a set of properties on the right (pink) contains

compositional features, the bottom center section shows counts of different locus types.

Fig. S2. GC content of different locus types in the SISRS dataset.

Fig. S3. Interrelationship between evaluated loci properties, assessed via a principal

component analysis. Only the first two PCA components are shown.

Fig. S4. Interrelationship between ΔlnL and sCF with respect to competing treeshrew

placements. Top panels correspond to individual locus fit, bottom to fit in concatenation.

Fig. S5. Interrelationship between ΔlnL and sCF with respect to monophyly of primates.

Top panels correspond to individual locus fit, bottom to fit in concatenation.

1002

1003

Supplementary Tables

Tab. S1. Sample information

Tab. S2. Loci properties and signal

Tab. S3. Results of the statistical analyses

Miscellaneous

File. S1. Compressed sequences of SISRS loci

1010