Study on the effect of a triple cancer treatment of propolis, thermal cycling-hyperthermia, and low-intensity ultrasound on PANC-1 cells

Yu-Yi Kuo[1,2], Wei-Ting Chen[1,2], Guan-Bo Lin[1,2], Chueh-Hsuan Lu[1,2], and Chih-Yu Chao[1,2,3,*]

[1] Department of Physics, Lab for Medical Physics & Biomedical Engineering, National Taiwan University, Taipei 10617, Taiwan

[2] Biomedical & Molecular Imaging Center, National Taiwan University College of Medicine, Taipei 10051, Taiwan

[3] Graduate Institute of Applied Physics, Biophysics Division, National Taiwan University, Taipei 10617, Taiwan

* Corresponding author

E-mail: cychao@phys.ntu.edu.tw (CYC)

Abstract

To reduce side effects and enhance treatment efficacy, study on combination therapy for pancreatic cancer, a deadly cancer, has gained much attraction in recent years. In this study, we propose a novel triple treatment combining propolis and two physical stimuli—thermal cycling-hyperthermia (TC-HT) and low-intensity ultrasound (US). The study found that, after the triple treatment, the cell viability of a human cancer cell line PANC-1 decreased to a level 80% less than the control, without affecting the normal pancreatic cells. Another result was excessive accumulation of reactive oxygen species (ROS) after the triple treatment, leading to the amplification of apoptotic pathway through the MAPK family and mitochondrial dysfunction, while activating autophagy simultaneously. This study, to the best of our knowledge, is the first attempt to combine TC-HT, US, and a natural compound in cancer treatment. The combination of TC-HT and US also promotes the anticancer effect of the heat-sensitive chemotherapy drug cisplatin on PANC-1 cells. It is expected that optimized parameters for different agents and different types of cancer will expand the methodology on oncological therapy in a safe manner.

Introduction

Cancer is one of the most dreadful diseases and the second leading cause of death around the world. Among all kinds of cancers, pancreatic cancer is the most threatening one, due to its high death rate and low five-year survival rate [1]. Existing therapies for pancreatic cancer, including surgery, radiation, and chemotherapy, all involve major risks, such as tumor recurrence, refractory, and serious side effects [2], as a result of which development of new therapies is of the utmost importance. A popular option is combination therapy, administering two or more anticancer agents to attain a synergistic effect [3]. However, the interaction between drugs may lead to unexpected competition and even harmful side effects [4, 5], jeopardizing patients' health, let alone improving therapeutic efficacy.

An emerging option in combination therapy is physical stimulus, whose effects on cellular physiology have been reported in several studies [6-8]. Our team has also looked into the feasibility of combing drug therapy and physical stimuli, such as heat [9], electric field [10], and magnetic field [11]. Ultrasound (US) is also a therapeutic tool with extensive application, such as developing internal images [12, 13], transporting liposomes to increase agent-delivery rate [14, 15], and ablating tumors from normal tissues [16, 17]. The previous study also found US as a helpful method in inhibiting significantly the viability of PANC-1

cells [18]. However, US may also entail a number of risks, such as harm to normal tissues

around tumors due to overheating caused by high-intensity US [19, 20]. In addition,

liposomes may induce myocardial injury during transport by US [21, 22]. In view of this,

integrating low-intensity US [23, 24] with a non-hazardous agent is important for expanding

the use of US in therapy.

Accordingly, the study employed natural compounds from herbal medicines as

anticancer agent. Among natural compounds with therapeutic potential, propolis has been

found to be effective in inhibiting several cancer cell lines [25-27]. In our previous study,

propolis was applied, along with thermal cycling-hyperthermia (TC-HT) as a physical

stimulus [28], in cancer treatment, but the effect still lags behind the in vitro efficacy of

chemotherapy drugs.

In this paper, the study introduced low-intensity US as a secondary helper, on top of

TC-HT, in order to further augment the anticancer effect of propolis. The novel triple

treatment turned out to inhibit the viability of PANC-1 cells significantly, approaching the in

vitro efficacy of chemotherapy drugs, without damaging the normal human pancreatic duct

cells and skin cells. The study found that the triple treatment manipulated the phosphorylation

levels of MAPK family, thereby activating the intracellular apoptotic signalling. Besides, the

autophagy signalling was noticed being greatly activated by the triple treatment. Moreover, it

was found in the upstream that intracellular reactive oxygen species (ROS) also increased

greatly, thereby boosting the death rate of PANC-1 cells.

Materials and Methods

Cell culture and propolis treatment

The human pancreatic cancer cell line PANC-1 and the normal human embryonic skin cell

line Detroit 551 were obtained from Bioresource Collection and Research Center (Hsinchu,

Taiwan). Normal human pancreatic duct H6c7 cells were purchased from Kerafast, Inc.

(Boston, MA, USA). PANC-1 and Detroit 551 cells were cultured respectively in DMEM and

EMEM (both from Hyclone, South Logan, UT, USA) supplemented with 10% fetal bovine

serum (FBS) (Hyclone) and 1% penicillin-streptomycin (Gibco Life Technologies, Grand

Island, NY, USA). H6c7 cells were maintained in keratinocyte-serum free medium

(Invitrogen, Life Technologies, Grand Island, NY, USA) supplemented with human

recombinant epidermal growth factor, bovine pituitary extract (Invitrogen), and 1%

penicillin-streptomycin (Gibco Life Technologies). All cells were maintained in a humidified

5% CO_2 incubator at 37 ☐ and subcultured by 0.05% trypsin–0.5 mM EDTA solution (Gibco

Life Technologies). Once the confluences reached suitable percentages, cells were plated in

96-well or 35-mm-diameter culture dishes (Thermo Fisher Scientific, Inc., Waltham, MA,

USA) for in vitro experiments after 24 h. Propolis was purchased from Grandhealth™ (Grand

Health Inc, Richmond, BC, Canada), and cisplatin was obtained from Sigma-Aldrich (St.

Louis, MO, USA). All agents were mixed with culture medium to the desired concentration

and were incubated with cells for 1h before treating physical stimuli.

Ultrasound exposure

The US exposure system consisted of a function generator (SG382; Stanford Research

Systems, Sunnyvale, CA, USA), a power amplifier (25a250a; amplifier research, Souderton,

PA, USA), and a planar transducer (A104S-RM; Olympus NDT Inc., Waltham, MA, USA).

Continuous pulses were produced using the function generator with the following parameters:

-10 dBm amplitude, 1 ms pulse period, and 0.5 ms pulse width. The cell culture plate or dish

was placed on the ceramic transducer (resonance frequency 2.25 MHz), which converted

electrical signals into acoustic power (**Fig 1A**). To avoid undesirable thermal effects induced

by US, the output power of the spatial average intensity of the US exposure was adjusted to

be 0.3 W/cm^2 according to the previous studies [29, 30].

Fig 1. Experimental setups for the triple treatment.

35-mm culture dishes were placed on (A) a ceramic transducer and (B) a modified PCR

machine for the exposures of US and TC-HT, respectively. (C) The schematic representation

of the TC-HT temperature settings. (D) Cell temperature detected by a needle thermocouple

when TC-HT was implemented. (E) Experimental schedule of the triple treatment with

different exposing order of US and TC-HT.

Thermal cycling-hyperthermia (TC-HT) treatment

A modified polymerase chain reaction (PCR) system was used to perform TC-HT (**Fig 1B**).

Thermal cycler (model 2720) was purchased from Applied Biosystems (Thermo Fisher

Scientific). The system was repeatedly brought to the desired high temperature state and

followed by a cooling stage to achieve a series of short period of heat exposure within the

desired time (**Fig 1C**). The experimental setup and administration of TC-HT (10-cycles) have

been previously described [28] with optimum results. The actual temperatures the cancer cells

sensed were measured by a needle thermocouple, ranging in 45~40.5 □ (**Fig 1D**). Ultrasound

exposure was applied either before or after the TC-HT treatment, for different combination

tests. During the TC-HT treatment (~ 45 min), the control and propolis-treated groups were

under room temperature (RT) without a 5% CO_2 environment. After the treatments, cells were

maintained in the cell culture incubator for the following experiments.

MTT assay

3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide (MTT) (Sigma-Aldrich) was

dissolved in distilled water to prepare a 5 mg/ml stock solution. The treated PANC-1 cells

were incubated for 4 h at 37 □ with a final MTT concentration 0.5 mg/ml in DMEM culture

medium to assess the cell viabilities. The formazan crystals were dissolved by equal volume

of the solubilizing buffer of 10% sodium dodecyl sulfate (SDS) (Bioshop Canada Inc.,

Burlington, ON, Canada) solution in 0.01 N hydrochloric acid (HCl) (Echo Chemical Co. Ltd.,

Taiwan) at 37 □ overnight. The absorbance of each well was detected by Multiskan GO

microplate Spectrophotometer (Thermo Fisher Scientific), and the quantity of formazan was

determined by the absorbance at 570 nm, with a background subtraction at 690 nm. The cell

viabilities were expressed in percentage and the untreated control was set at 100%.

Treatment with ROS scavenger

PANC-1 cells were seeded into 96-well or 35-mm-diameter culture dishes overnight. For

ROS inhibition analysis, cells were pretreated with 5 mM N-acetyl-cysteine (NAC)

(Sigma-Aldrich) in culture medium for 1 h and subsequently treated with propolis and/or

physical stimuli.

Apoptotic analysis by flow cytometry

PANC-1 cells were collected 24 h after treatments and then rinsed twice with ice-cold

phosphate buffered saline (PBS) (Hyclone). The apoptotic rates were analyzed by the

Annexin V-FITC and propidium iodide (PI) double detection kit (BD Biosciences, San Jose,

CA, USA), and the rinsed cells were resuspended in binding buffer containing Annexin

V-FITC and PI and then incubated at RT for 15 min in the dark. Apoptotic signals were

detected by FACSCanto II system (BD Biosciences).

ROS and mitochondrial membrane potential (MMP) analyses by flow cytometry

ROS was detected using the fluorescent dye dihydroethidium (DHE) (Sigma-Aldrich), and

the loss of MMP was determined using the lipophilic cationic fluorescent dye

3,3'-dihexyloxacarbocyanine iodide ($DiOC_6(3)$) (Enzo Life Sciences, Inc., Plymouth Meeting,

PA, USA). PANC-1 cells were harvested 24 h after treatments and rinsed with PBS before

staining. Rinsed cells were resuspended and then incubated with 5 μM DHE or 20 nM

$DiOC_6(3)$ in PBS at 37 □ for 30 min in the dark. The fluorescence signals were measured by

FACSCanto II system (BD Biosciences) with the PE channel (for DHE staining) or FL1

channel (for $DiOC_6(3)$ staining).

Western blot analysis

Protein expression levels of PANC-1 cells were quantified by western blot analysis. Cells

were rinsed with PBS and then lysed in the lysis buffer (50 mM Tris-HCL, pH 7.4, 0.15 M

NaCl, 0.25% deoxycholic acid, 1% NP-40, 1% Triton X-100, 0.1 % SDS, 1 mM EDTA)

(Millipore, Billerica, MA, USA), supplemented with active protease (Millipore) and

phosphatase inhibitor cocktail (Cell signaling Technology, Danvers, MA, USA). After

centrifugation, the supernatants were collected and the protein concentrations were quantified

by Bradford protein assay (Bioshop, Inc.). Equal amount of proteins (20 μg) were resolved by

10% SDS-polyacrylamide gel electrophoresis (SDS-PAGE) and then transferred onto

polyvinylidene fluoride (PVDF) membranes (Millipore). 5% skim milk powder or 5% bovine

serum albumin in TBST (20 mM Tris-base, pH 7.6, 0.15 M NaCl, 0.1% Tween 20) was used

to block nonspecific antibody binding sites for 1 h at RT. Afterwards, the blocked membranes

were probed with specific primary antibodies against p-ERK, p-JNK, PARP (Cell signaling),

LC3 and GAPDH (Gentex, Irvine, CA, USA) at 4 □ overnight. The membranes were rinsed

with TBST buffer three times and then incubated with horseradish peroxidase-conjugated

goat anti-rabbit secondary antibodies (Jackson ImmunoResearch Laboratories, West Grove,

PA, USA) in a blocking solution at RT for 1 h. Immunoreactivity signal was amplified by an

enhanced chemiluminescence (ECL) substrate (Advansta, San Jose, CA, USA) and detected

by an imaging system Amersham Imager 600 (GE Healthcare Life Sciences). GAPDH was

used as the loading control to normalize the relative folds of targeting proteins.

Statistical analysis

Experiments were repeated three times for validation, and statistical analyses were performed

using one-way analysis of variance (ANOVA) by OriginPro 2015 software (OriginLab).

Results were expressed as the mean ± standard deviation, and were considered to be

statistically significant when *p*-values were less than 0.05.

Results

Triple treatment greatly inhibits the viability of PANC-1 cells

The cell viability of PANC-1 cells versus the propolis concentration was performed in a

gradient manner. When the propolis was less than 0.5%, as shown in **Fig 2A**, there was no

notable inhibition effect on MTT results. However, when it exceeded 0.5%, the cell viability

dropped significantly. Therefore, a moderate propolis concentration 0.3% was chosen for the

following experiments. Next, the physical stimuli of TC-HT and low-intensity US were

introduced to affect the viability of PANC-1 cells. In our study, 10-cycles TC-HT and 2.25

MHz US with intensity $0.3W/m^2$ and duration 30 minutes were chosen to avoid the

thermotoxicity on PANC-1 cells. As shown in **Fig 2B**, we found that the combination of

TC-HT and US was also innocent to PANC-1 cells, but when 0.3% propolis was involved in

the triple treatment, the viability of PANC-1 cells was greatly inhibited. It was also noted that

the implementation order of TC-HT and US in triple treatment was influential. When TC-HT

was performed prior to US (TC-HT + US) in the presence of 0.3% propolis, the cell viability

of PANC-1 cells was significantly suppressed down to 17.1%, cutting more than 80% of the

viability of the untreated control and thus approaching the in vitro efficacy of chemotherapy

drugs. In comparison, the treatment that US was performed prior to TC-HT (US + TC-HT)

showed a less inhibition effect (43.1% viability), and hence we adopted the implementation

order TC-HT + US as the protocol of the triple treatment in the subsequent experiments.

Furthermore, in all double treatments, only 0.3% propolis + TC-HT showed notable inhibition

effect (48.9% viability) on PANC-1 cells, which was consistent with our previous results [28].

However, 0.3% propolis + US performed a relatively poor inhibition effect (65.4% viability)

on PANC-1 cells, and as a result it was also not included in the following experiments. **Fig**

2C showed the light microscope images of PANC-1 cells 24 h after each treatment, and the

cell morphologies demonstrated an evident inhibition effect on PANC-1 cells after the triple

treatment. Moreover, normal cells such as the human skin cells Detroit 551 (**Fig 2D**) and

human pancreatic duct cells H6c7 (**Fig 2E**) were not significantly affected by the triple

treatment as well as all the other treatments. The result indicates that the triple treatment

could have a good selective effect on carcinoma cells and normal cells, which makes it safer

and more feasible in anticancer treatment.

Fig 2. Viability inhibition effects of propolis on PANC-1, Detroit 551, and H6c7 cells.

MTT assay was conducted to determine the viabilities of PANC-1 cells after the treatment of

(A) different propolis concentrations and (B) different combinations of physical stimulations.

(C) Representative light microscope images of PANC-1 cells after each treatment, scale bar =

100 μm. The viabilities of (D) normal human skin cells Detroit 551 and (E) normal human

pancreatic duct cells H6c7 were measured 24 h after each treatment. Data were presented as

the mean ± standard deviation in triplicate. (***P<0.001).

Triple treatment increases intracellular ROS levels synergistically

Intracellular ROS is an important regulator of cell death. It has been reported that heat and

low-intensity US could elevate the intracellular ROS level [31, 32]. We further investigated

whether ROS was increased in response to the triple treatment, so the DHE was used in this

experiment to determine the level of superoxide radical anion ($O_2^{\cdot-}$) in PANC-1 cells after

each treatment. As shown in **Fig 3A** and **3B**, it was found that propolis hardly changed the

fluorescence signals. Although propolis + TC-HT significantly deformed the fluorescence

intensity distribution in an enhanced manner (1.6-fold to control), it did not significantly

differ from the enhancement induced by TC-HT alone. In addition, US elevated ROS levels

as well, though not as many as TC-HT. Noticeably, the triple treatment showed a significant

accumulation of the intracellular ROS (up to 2.1-fold of the control group), which was also

significantly higher than the TC-HT + 0.3% propolis treatment. The result suggested that the

triple treatment boosted up the generation of ROS in PANC-1 cells, and could result in

enhanced cell death rate after the treatment.

Fig 3. The triple treatment raised the generation of intracellular ROS.

(A) Intracellular superoxide radical anion ($O_2^{\cdot-}$) levels of PANC-1 cells were determined by

flow cytometry with the fluorescent dye DHE. (B) Quantification of the mean DHE

fluorescence levels after each treatment. Data were presented as the mean $\pm$ standard

deviation in triplicate. (***$P<0.001$).

Triple treatment increases mitochondrial apoptosis and autophagy in PANC-1

cells

It has been known that the enhanced intracellular ROS levels were positively correlated to

mitochondrial apoptosis [33]. In our work, the apoptotic rates of PANC-1 cells after various

treatments were analyzed by the flow cytometry with the fluorescence dye Annexin V and PI

(**Fig 4A & 4B**). The triple treatment caused 55.3% apoptotic rate, which was significantly

higher than the 23.5% of propolis + TC-HT. The cell apoptosis results observed here were

highly consistent with the results of the accumulated ROS levels in PANC-1 cells after the

same treatment, as described in **Fig 3**. Furthermore, the mitochondrial membrane potential

(MMP) was assessed using $DiOC_6(3)$ fluorescence staining by flow cytometric analysis. As

shown in **Fig 4C** and **4D**, the ratio of the cells exhibiting MMP loss was significantly

promoted to 23.3% after the double treatment of propolis + TC-HT, and it was further

elevated significantly to 34.7% by adopting the triple treatment. These results showed that the

triple treatment could decrease MMP level, and hence caused mitochondrial dysfunction. The

decreased MMP level was an indicator of mitochondrial apoptosis, and since the results of

apoptosis assay (**Fig 4A & 4B**) and MMP assay (**Fig 4C & 4D**) were quite similar, we

believe that the mitochondrial dysfunction was implicated in the apoptosis of PANC-1 cells

via the triple treatment.

Fig 4. Triple treatment increased apoptosis on PANC-1 cells via mitochondrial

dysfunction.

(A) Apoptosis after treatment was analyzed via flow cytometry with Annexin V-FITC/PI

double staining, and (B) the apoptotic percentage (Q2+Q4) were calculated. (C) MMP level

after treatment was analyzed via flow cytometry with $DiOC_6(3)$ staining, and (D) the

percentage of cells with the loss of MMP (M1) was calculated. Data were presented as the

mean ± standard deviation in triplicate. (***P<0.001, **P<0.01).

In addition to apoptosis of PANC-1 cells, our previous study has also shown that US alone or

accompanied with other treatments could promote autophagy in PANC-1 cells [18]. In this

experiment, autophagy was also examined after each treatment. The conversion of precursor

LC3-I to the product LC3-II (ratio of LC3-II/LC3-I) quantified the autophagy activity, and the

western blot result showed that both US and TC-HT could slightly elevate the autophagy

activity in PANC-1 cells (**Fig 5**). Moreover, propolis + TC-HT treatment further enhanced

this activity, but it was significantly less than the autophagy activity produced by the triple

treatment, as shown in **Fig 5**. Hence, the triple treatment in this study was also an effective

promoter of autophagy in PANC-1 cells.

Fig 5. Autophagy activation after the triple treatment in PANC-1 cells.

(A) Representative western blots of autophagy related proteins LC3-I and LC3-II and (B) the

quantification of LC3-II/LC3-I ratio. GAPDH was used as loading control. Data were

presented as the mean ± standard deviation in triplicate. (***P<0.001, **P<0.01).

The apoptosis induced by triple treatment is regulated through MAPK pathway

The activation of apoptotic signalling was examined by western blot analysis. As shown in

Fig 6A, we found that the PARP cleavage was significantly increased (2.9-fold of control)

after propolis + TC-HT treatment on PANC-1 cells. Noticeably, the PARP cleavage was

further promoted significantly to 6.2-fold of control by US in the triple treatment (**Fig 6A**).

Together with the previous flow cytometry results of apoptosis and MMP, it was pointed out

that propolis + TC-HT could activate the mitochondrial apoptosis signalling in PANC-1 cells,

and US in the triple treatment could further help this cascade to realize a near-chemotherapy

level treatment in vitro.

Moreover, it was known that the PARP cleavage could be modulated by MMP level,

and mitochondrial dysfunction could also be regulated by the excessive intracellular ROS via

MAPK family [34]. In MAPK family, the phosphorylated ERK (p-ERK) level represented the

activation of cell survival [35], while the phosphorylated JNK (p-JNK) and p38 (p-p38) levels

were the indicator of cell death [36, 37]. In this study, it was found that the p-ERK level was

suppressed by propolis + TC-HT treatment (0.30-fold), and was further down-regulated when

US was introduced in the triple treatment (0.15-fold) (**Fig 6B**). In addition, the p-JNK and

p-p38 levels both exhibited a reverse performance, which were promoted the most in the

triple treatment (8.7-fold & 9.2-fold, respectively) (**Fig 6C & 6D**). These results were

consistent with the results of ROS and MMP assessments by flow cytometry. Therefore, we

speculated that the excess intracellular ROS induced by the triple treatment regulated the

activation of the MAPK family and thus caused mitochondrial dysfunction and the cascade of

apoptosis.

Fig 6. Triple treatment modulated apoptosis via regulating the MAPK family.

Representative western blots of the apoptosis-related proteins and the quantification of (A)

the PARP cleavage (ratio of cleaved PARP/full length PARP), the phosphorylation level of

(B) ERK, (C) JNK, and (D) p38. GAPDH was used as loading control. Data were presented

as the mean ± standard deviation in triplicate. (***$P<0.001$, **$P<0.01$, *$P<0.05$).

ROS scavenger attenuates the apoptosis induced by triple treatment

To further confirm that the cell death after the triple treatment was regulated by the

generation of intracellular ROS, the ROS scavenger NAC was applied in the experiment [38].

5 mM NAC was incubated with PANC-1 cells 1 h prior to the triple treatment. As shown in

Fig 7A, the inhibitory effect of the triple treatment was restored by NAC, and NAC itself did

not affect the viability of PANC-1 cells. Similar results were also observed in the activation

of apoptotic pathway, as shown in **Fig 7B**. NAC alone did not affect PARP cleavage, but it

significantly down-regulated the triple treatment-promoted PARP cleavage (**Fig 7B**).

Therefore, the results supported our speculation that the triple treatment could induce

mitochondrial apoptosis of PANC-1 cells via the excessive increment of intracellular ROS.

Fig 7. ROS inhibition blocked the cell death and the activation of apoptosis pathways.

(A) Cell viabilities of PANC-1 cells and (B) the quantification of PARP cleavage (ratio of

cleaved PARP/full length PARP) after the triple treatment with or without 1h NAC

pretreatment. GAPDH was used as loading control. Data were presented as the mean ±

standard deviation in triplicate. (***P<0.001, **P<0.01).

Triple treatment can be applied with chemotherapy drug as a novel anticancer

treatment

In this study, we have shown that the method TC-HT followed by mild US exposure could

further amplify the anticancer effect of propolis. But, the question is whether the TC-HT +

US method can be expanded to the existing chemotherapy drugs. Cisplatin, for instance, was

a commonly used clinical chemotherapy drug for several kinds of cancers such as lung,

ovarian, breast, and brain cancer [39]. Besides, it has also been reported that cisplatin was

sensitive to heat [40]. However, the conventional therapeutic dosage of cisplatin could cause

severe side effects to the patients [41, 42], and therefore it was important to develop a new

method to reduce the effective dose of cisplatin. As a result, we applied the method of TC-HT

+ US with cisplatin on the PANC-1 cells to investigate the potential of this triple treatment

method. A relatively low dose of 1 μM cisplatin and short incubation time 24 h was chosen

[43, 44] for the MTT assay independently or in combination with the physical stimulations.

After the treatment, the viability of PANC-1 cells was just slightly inhibited by 7.5% by

individual cisplatin, and TC-HT + cisplatin treatment also did not alter the viability of

PANC-1 cells. The triple treatment, however, promoted the inhibitory effect significantly up

to 48.2% (**Fig 8**). Compared to the conventional results of cisplatin, our method could not

only reduce the effective dose but also boost up the anticancer effect of cisplatin. Therefore,

the method of physical stimuli in the study should hold great potential to apply onto other

heat sensitive chemotherapy drugs or anticancer agents to achieve a better anticancer effect.

Fig 8. The method of triple treatment promoted the inhibitory effect of the heat sensitive

chemotherapy drug cisplatin.

Cell viabilities of PANC-1 cells treated with cisplatin, TC-HT + cisplatin, and the triple

treatment of cisplatin. Cisplatin concentrations in all treatments were 1 μM. Data were

presented as the mean ± standard deviation in triplicate. (***P<0.001).

Discussion

Combination treatment augments the anticancer effect of individual agents by activating

multiple pathways, thereby lowering the necessary dosage of the agents to a level harmless to

normal cells and human health. However, research on combination anticancer agents is costly

and time-consuming [45]. Moreover, unpredictable molecular interactions may be detrimental

to patients' health [5]. Therefore, the application of physical stimuli has been considered as a

potential candidate for combinative anticancer treatments in combating cancer. Following

several combination treatments of physical stimuli and herbal compounds proposed by our

team previously [46], the study put forth the combination treatment of propolis, TC-HT, and

low-intensity US, proving that it can inhibit pancreatic cancer cell line PANC-1 at a level

close to the in vitro efficacy of chemotherapy drugs.

In cancer cells, intracellular ROS levels have been known to be the main source of the

oxidative stress [47], a key factor for cell viability [48]. Elevated ROS level has been shown

to be able to activate some signalling pathways associated with cell proliferation, apoptosis,

autophagy, and cell cycle progression [49]. Heat treatment and low-intensity US both can

increase the intracellular ROS levels [31, 32]. In this work, the study shows that TC-HT

significantly augments the intracellular ROS levels of PANC-1 cells (**Fig 3A** and **Fig 3B**), at

an extent higher than the individual effects of propolis and low-intensity US. It was found that

the combination of propolis and TC-HT did not further elevate the levels in PANC-1 cells.

Nevertheless, the triple treatment showed a great improvement effect, doubling the

intracellular ROS levels of PANC-1 cells. It has been known that excessive intracellular ROS

level could activate the apoptotic pathway cascade, increasing the apoptosis in the carcinoma

cells [33]. Our study demonstrated that the apoptotic rate of PANC-1 cells was elevated along

with the increase of the intracellular ROS levels. The result showed that the triple treatment

induced the highest apoptotic rate, compared with other approaches, suggesting its ability to

regulate the death of PANC-1 cells via excessive intracellular ROS accumulation. To

demonstrate the crucial role of the intracellular ROS levels, NAC was employed in the

following experiments on the triple treatment. While independent NAC pretreatment did not

affect the viabilities of PANC-1 cells (**Fig 7A**), it protected the cells from cytotoxicity in the

triple treatment. Hence, ROS elevation played a key role in the anticancer effect in the triple

treatment.

The initiation of apoptosis, a common cell death mechanism, is closely related to the

function of mitochondria, which is the chemical-energy source of cells and critical for the

viability of cells [50]. Its dysfunction could induce a series of biochemical cascade of

apoptosis, thereby blocking cell proliferation [51]. Meanwhile, the activated members of the

MAPK family, such as ERK, JNK, and p38 were deemed to be capable of regulating the

dysfunction of mitochondria, and the elevated ROS level was shown to be conducive to the

activation of p38 and JNK but down-regulate the activation of ERK [52]. The results suggest

that excessive ROS could cause the mitochondrial apoptotic rate via activating the MAPK

family members. In this work, our study showed that the triple treatment raised significantly

the apoptotic rate of PANC-1 cells (**Fig 4A** and **4B**), while decreasing the MMP level, which

led to the dysfunction of mitochondria (**Fig 5A** and **5B**). Furthermore, the triple treatment

increased the phosphorylated levels of p38 and JNK significantly, while inhibiting the

phosphorylation of ERK (**Fig 6**), underscoring its ability to manipulate the function of

mitochondria via the ROS-activated MAPK family proteins.

The injured mitochondria would release cytochrome-c into the cytoplasm [53], cleaving

caspase 9 and thus activating caspase 3 in the downstream [54], which entered further the

nucleus and cleaved PARP. Then PARP would lose its enzyme activity, initiating apoptosis

irreversibly [55]. The study demonstrated that the triple treatment could induce mitochondrial

dysfunction via regulating the phosphorylation level of the members of MAPK family. In

addition, the triple treatment also increased PARP cleavage significantly, compared with the

combination treatment of propolis and TC-HT (**Fig 6A**). Furthermore, as shown in **Fig 7B**, it

was found that the triple treatment-promoted PARP cleavage was significantly suppressed by

NAC, which indicates that increased ROS level was the key regulator for the apoptotic effect

of the triple treatment.

The triple treatment in the study included two physical stimuli, TC-HT and US, which

could be integrated for simultaneous implementation via the high-intensity focused ultrasound

(HIFU). Being able to raise temperature in the exposed region for energy transfer [19, 20],

HIFU has been applied for tumor ablation for over a decade [16, 56]. The temperature

increase could be electrically controlled and directed to the targeted region [57]. Therefore,

with the help of HIFU, the triple treatment could be applied clinically. To augment the effect,

cisplatin, a common thermal sensitive chemotherapy drug, was incorporated as a substitute

for propolis into the triple treatment. It was shown that the inhibited viability of PANC-1 cells

by cisplatin was further suppressed with a large extent when both of TC-HT and US were

introduced into the treatment (**Fig 8**). The dosage of cisplatin can be reduced to a lower

concentration, without compromising the anticancer effect. As a result, the triple treatment

has the potential supplementing the administration of drugs, not only augmenting the effect

but also reducing the dosage of the latter.

In summary, this study proposed for the first time an effective triple cancer treatment

combining propolis, TC-HT, and low-intensity US, which could significantly suppress the

growth of PANC-1 cells via an ROS-modulated mitochondrial apoptosis, with a performance

comparable to chemotherapy. The study also showed that the triple treatment could induce

mitochondrial dysfunction via the regulation of MAPK family, resulting in apoptosis via the

up-regulated PARP cleavage. It also demonstrated that the ROS level plays a key role in the

performance of the triple treatment. In addition, chemotherapy drugs, such as cisplatin, can be

incorporated into the treatment as substitute for propolis. The triple treatment incorporating

cisplatin also exhibited a much higher effect in inhibiting cancer cell growth than the cisplatin

alone, promising to increase the performance and safety of the existing cancer therapy.

Overall, the study proposed employment of physical stimuli, as a promising option in cancer

therapy.

Funding

This work was supported by grants from Ministry of Science and Technology

(MOST 110-2112-M-002-004, MOST 109-2112-M-002-004, MOST

108-2112-M-002-016, and MOST 105-2112-M-002-006-MY3 to CYC) and Ministry

of Education (MOE 106R880708 to CYC) of the Republic of China. The funders had

no role in study design, data collection and analysis, decision to publish, or

preparation of the manuscript.

Competing Interests

455   The authors have declared that no competing interests exist.

Acknowledgments

The authors would like to acknowledge the service provided by Technology Commons in College of Life Science, National Taiwan University for use of flow cytometry system.

Author Contributions

Conceptualization: Chih-Yu Chao.

Data Curation: Yu-Yi Kuo, Chih-Yu Chao.

Formal analysis: Yu-Yi Kuo, Wei-Ting Chen, Guan-Bo Lin, Chueh-Hsuan Lu, Chih-Yu Chao.

Funding acquisition: Chih-Yu Chao.

Investigation: Yu-Yi Kuo, Wei-Ting Chen, Chih-Yu Chao.

Project Administration: Chih-Yu Chao.

Supervision: Chih-Yu Chao.

Validation: Yu-Yi Kuo, Wei-Ting Chen.

Writing – original draft: Yu-Yi Kuo, Chih-Yu Chao.

Writing – review & editing: Chih-Yu Chao.

References

1. Siegel RL, Miller KD, Jemal A. Cancer statistics, 2019. CA Cancer J Clin. 2019; 69(1): 7-34.

2. Lee JC, Ahn S, Cho IK, Lee J, Kim J, Hwang JH. Management of recurrent pancreatic cancer after surgical resection: a protocol for systematic review, evidence mapping and meta-analysis. BMJ Open. 2018; 8(4):e017249.

3. Yap TA, Omlin A, de Bono JS. Development of therapeutic combinations targeting major cancer signaling pathways. J Clin Oncol. 2013; 31(12):1592-1605.

4. Terrault NA. Benefits and risks of combination therapy for hepatitis B. Hepatology. 2009; 49(5 Suppl):S122-8.

5. Lee JJ, Kong M, Ayers GD, Lotan R. Interaction index and different methods for determining drug interaction in combination therapy. J Biopharm Stat. 2007; 17(3):461-80.

6. Soares PI, Ferreira IM, Igreja RA, Novo CM, Borges JP. Application of hyperthermia for cancer treatment: recent patents review. Recent Pat Anticancer Drug Discov. 2012; 7(1):64-73.

7. Davies AM, Weinberg U, Palti Y. Tumor treating fields: a new frontier in cancer therapy. Ann N Y Acad Sci. 2013; 1291(1):86-95.

8. Wood AKW, Sehgal CM. A review of low-intensity ultrasound for cancer therapy. Ultrasound Med Biol. 2015; 41(4):905-928.

9. Lu CH, Chen WT, Hsieh CH, Kuo YY, Chao CY. Thermal cycling-hyperthermia in combination with polyphenols, epigallocatechin gallate and chlorogenic acid, exerts synergistic anticancer effect against human pancreatic cancer PANC-1 cells. PLoS One. 2019; 14(5):e0217676.

10. Lu CH, Lin SH, Hsieh CH, Chen WT, Chao CY. Enhanced anticancer effects of low-dose curcumin with non-invasive pulsed electric field on PANC-1 cells. Onco Targets Ther. 2018; 11:4723-32.

503   11. Chen WT, Lin GB, Lin SH, Lu CH, Hsieh CH, Ma BL, et al. Static magnetic field
504   enhances the anticancer efficacy of capsaicin on HepG2 cells via capsaicin receptor
505   TRPV1. PLoS One. 2018; 13(1):e0191078.

506   12. O'Keeffe D, Mamtora H. Ultrasound in clinical orthopaedics. J Bone Joint Surg. 1992;
507   74(4):488-94.

508   13. Bendick PJ, Brown OW, Hernandez D, Glover JL, Bove PG. Three-dimensional
509   vascular imaging using Doppler ultrasound. Am J Surg. 1998; 176(2):183-7.

510   14. Ferrara K, Pollard R, Borden M. Ultrasound microbubble contrast agents:
511   fundamentals and application to gene and drug delivery. Annu Rev Biomed Eng.
512   2007; 9:415-47.

513   15. Wang TY, Wilson KE, Machtaler S, Willmann JK. Ultrasound and microbubble
514   guided drug delivery: mechanistic understanding and clinical implications. Curr
515   Pharm Biotechnol. 2013; 14(8):743-52.

516   16. Zhou YF. High intensity focused ultrasound in clinical tumor ablation. World J Clin
517   Oncol. 2011; 2(1):44070.

518   17. Lang BH, Wu ALH. High intensity focused ultrasound (HIFU) ablation of benign
519   thyroid nodules—a systematic review. J Ther Ultrasound. 2017; 5:11.

520   18. Hsieh CH, Lu CH, Kuo YY, Chen WT, Chao CY. Studies on the non-invasive
521   anticancer remedy of the triple combination of epigallocatechin gallate, pulsed
522   electric field, and ultrasound. PLoS One. 2018; 13(8):e0201920.

523   19. van den Bijgaart RJ, Eikelenboom DC, Hoogenboom M, Fütterer JJ, den Brok MH,
524   Adema GJ. Thermal and mechanical high-intensity focused ultrasound: perspectives
525   on tumor ablation, immune effects and combination strategies. Cancer Immunol
526   Immunother. 2017; 66(2):247-58.

527   20. Dubinsky TJ, Cuevas C, Dighe MK, Kolokythas O, Hwang JH. High-intensity
528   focused ultrasound: current potential and oncologic applications. AJR Am J
529   Roentgenol. 2008; 190(1):191-9.

530   21. Zhang B, Zhou H, Cheng Q, Lei L, Hu B. Low-frequency low energy ultrasound
531   combined with microbubbles induces distinct apoptosis of A7r5 cells. Mol Med Rep.
532   2014; 10(6):3282-8.

533   22. Vancraeynest D, Havaux X, Pasquet A, Gerber B, Beauloye C, Rafter P, et al.
534   Myocardial injury induced by ultrasound-targeted microbubble destruction: evidence
535   for the contribution of myocardial ischemia. Ultrasound Med Biol. 2009;
536   35(4):672-9.

23. Johns LD. Nonthermal effects of therapeutic ultrasound: the frequency resonance hypothesis. J Athl Train. 2002; 37(3):293-9.

24. Bazou D, Maimon N, Munn LL, Gonzalez I. Effects of low intensity continuous ultrasound (LICU) on mouse pancreatic tumor explants. Appl Sci. 2017; 7(12):1275.

25. Taira N, Nguyen BC, Be Tu PT, Tawata S. Effect of Okinawa propolis on PAK1 activity, caenorhabditis elegans longevity, melanogenesis, and growth of cancer cells. J Agric Food Chem. 2016; 64(27):5484-9.

26. Li H, Kapur A, Yang JX, Srivastava S, McLeod DG, Paredes-Guzman JF, et al. Antiproliferation of human prostate cancer cells by ethanolic extracts of Brazilian propolis and its botanical origin. Int J Oncol. 2007; 31(3):601-6.

27. Xuan H, Li Zhen, Yan H, Sang Q, Wang K, He Q, et al. Antitumor activity of Chinese propolis in human breast cancer MCF-7 and MDA-MB-231 cells. Evid Based Complement Alternat Med. 2014; 2014:280120.

28. Chen WT, Sun YK, Lu CH, Chao CY. Thermal cycling as a novel thermal therapy to synergistically enhance the anticancer effect of propolis on PANC□1 cells. Int J Oncol. 2019; 55(3):617-28.

29. Love L, Kremkau F. Intracellular temperature distribution produced by ultrasound. J Acoust Soc Am.1980; 67(3):1045-50.

30. Nahirnyak V, Mast TD, Holland CK. Ultrasound-induced thermal elevation in clotted blood and cranial bone. Ultrasound Med Biol. 2007; 33(8):1285-95.

31. Hou CH, Lin FL, Hou SM, Liu JF. Hyperthermia induces apoptosis through endoplasmic reticulum and reactive oxygen species in human osteosarcoma cells. Int J Mol Sci. 2014; 15(10):17380-95.

32. Xia C, Zeng H, Zheng Y. Low-intensity ultrasound enhances the antitumor effects of doxorubicin on hepatocellular carcinoma cells through the ROS-miR-21-PTEN axis. Mol Med Rep. 2020; 21(3):989-98.

33. Redza-Dutordoir M, Averill-Bates DA. Activation of apoptosis signalling pathways by reactive oxygen species. Biochim Biophys Acta. 2016; 1863(12):2977-92.

34. Zang YQ, Feng YY, Luo YH, Zhai YQ, Ju XY, Feng YC, et al. Quinalizarin induces ROS-mediated apoptosis via the MAPK, STAT3 and NF-κB signaling pathways in human breast cancer cells. Mol Med Rep. 2019; 20(5):4576-86.

35. Meng J, Fang B, Liao Y, Chresta CM, Smith PD, Roth JA. Apoptosis induction by MEK inhibition in human lung cancer cells is mediated by Bim. PLoS One. 2010; 5(9):e13026.

36. Chiyo T, Fujita K, Iwama H, Fujihara S, Tadokoro T, Ohura K, et al. Galectin-9
induces mitochondria-mediated apoptosis of esophageal cancer in vitro and in vivo in
a xenograft mouse model. Int J Mol Sci. 2019; 20(11):2634.

37. Wu YJ, Su TR, Dai GF, Su JH, Liu CI. Flaccidoxide-13-acetate-induced apoptosis in
human bladder cancer cells is through activation of p38/JNK, mitochondrial
dysfunction, and endoplasmic reticulum stress regulated pathway. Mar Drugs. 2019;
17(5):287.

38. Aruoma OI, Halliwell B, Hoey BM, Butler J. The antioxidant action of
N-acetylcysteine: its reaction with hydrogen peroxide, hydroxyl radical, superoxide,
and hypochlorous acid. Free Radic Biol Med. 1989; 6(6):593-7.

39. Dasari S, Tchounwou PB. Cisplatin in cancer therapy: molecular mechanisms of
action. Eur J Pharmacol. 2014; 740:364-78.

40. Behrouzkia Z, Joveini Z, Keshavarzi B, Eyvazzadeh N, Aghdam RZ. Hyperthermia:
How Can It Be Used? Oman Med J. 2016; 31(2):89-97.

41. Florea AM, Büsselberg D. Cisplatin as an anti-tumor drug: cellular mechanisms of
activity, drug resistance and induced side effects. Cancers. 2011; 3(1):1351-1371.

42. Ghosh S. Cisplatin: the first metal based anticancer drug. Bioorg Chem. 2019;
88:102925.

43. Cregan IL, Dharmarajan AM, Fox SA. Mechanisms of cisplatin-induced cell death in
malignant mesothelioma cells: Role of inhibitor of apoptosis proteins (IAPs) and
caspases. Int J Oncol. 2013; 42(2):444-52.

44. Beer M, Kuppalu N, Stefanini M, Becker H, Schulz I, Manoli S, et al. A novel
microfluidic 3D platform for culturing pancreatic ductal adenocarcinoma cells:
comparison with in vitro cultures and in vivo xenografts. Sci Rep. 2017; 7(1):1325.

45. Huang H, Zhang P, Qu XA, Sanseau P, Yang L. Systematic prediction of drug
combinations based on clinical side-effects. Sci Rep. 2014; 4:7160.

46. Lu CH, Kuo YY, Lin GB, Chen WT, Chao CY. Application of non-invasive
low-intensity pulsed electric field with thermal cycling-hyperthermia for
synergistically enhanced anticancer effect of chlorogenic acid on PANC-1 cells.
PLoS ONE. 2020; 15(1):e0222126.

47. Liou GY, Storz P. Reactive oxygen species in cancer. Free Radic Res. 2010;
44(5):479-96.

48. Kannan K, Jain SK. Oxidative stress and apoptosis. Pathophysiology. 2000;
7(3):153-63.

49. Aggarwal V, Tuli HS, Varol A, Thakral F, Yerer MB, Sak K. Role of reactive oxygen
species in cancer progression: molecular mechanisms and recent advancements.
Biomolecules. 2019; 9(11):735.

50. Osellame LD, Blacker TS, Duchen MR. Cellular and molecular mechanisms of
mitochondrial function. Best Pract Res Clin Endocrinol Metab. 2012; 26(6):711-23.

51. Elmore S. Apoptosis: a review of programmed cell death. Toxicol Pathol. 2007;
35(4):495-516.

52. Yuan L, Wang J, Xiao H, Wu W, Wang Y, Liu X. MAPK signaling pathways
regulate mitochondrial-mediated apoptosis induced by isoorientin in human
hepatoblastoma cancer cells. Food Chem Toxicol. 2013; 53:62-8

53. Garrido C, Galluzzi L, Brunet M, Puig PE, Didelot C, Kroemer G. Mechanisms of
cytochrome c release from mitochondria. Cell Death Differ. 2006; 13(9):1423-33.

54. Brentnall M, Rodriguez-Menocal L, De Guevara RL, Cepero E, Boise LH. Caspase-9,
caspase-3 and caspase-7 have distinct roles during intrinsic apoptosis. BMC Cell Biol.
2013; 14:32.

55. Jimenez F, Aiba☐Masago S, Al Hashimi I, Vela☐Roch N, Fernandes G, Yeh CK, et
al. Activated caspase 3 and cleaved poly(ADP-ribose)polymerase in salivary
epithelium suggest a pathogenetic mechanism for Sjögren's syndrome. Rheumatology.
2002; 41(3):338-42.

56. Duc NM, Keserci B. Emerging clinical applications of high-intensity focused
ultrasound. Diagn Interv Radiol. 2019; 25(5):398-409.

57. Park S, Pham NT, Huynh HT, Kang HW. Development of temperature
controller-integrated portable HIFU driver for thermal coagulation. Biomed Eng
Online. 2019; 18(1):77.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8