1 **A Comprehensive Roadmap Towards Generation of Influenza B Reporter Assay Using a**
2 **Single DNA Polymerase Based Cloning of Reporter RNA Construct.**

3 Nandita Kedia[1], Saptarshi Banerjee[1], Arindam Mondal[1*]

4 [1]School of Bioscience, Indian Institute of Technology Kharagpur, Kharagpur -721302, India
5
6 *Corresponding author
7

Abstract:

Mini-genome reporter assay is a key tool for conducting RNA virus research. But, procedural
complications and lack of adequate literature pose major challenge towards developing these
assay systems. Here we present a novel yet generic and simple cloning strategy for construction
of influenza B virus reporter RNA template and describe extensive standardization of the reporter
RNP/ polymerase activity assay for monitoring viral RNA synthesis in infection free setting. Using
this assay system, we, for the first time showed the effect of viral protein NS1 and host protein
PKC-Delta upon influenza B virus RNA synthesis. Additionally, the assay system showed
promising results in evaluating efficacy of antiviral drugs targeting viral RNA synthesis and virus
propagation. Together, this work offers a detailed protocol for standardization of influenza virus
mini-genome assay and an excellent tool for screening of host factors and antivirals in a fast, user
friendly and high throughput manner.

Keywords:

21 Influenza B virus, engineering reporter construct, ribonucleoprotein particles, reporter based
22 activity assay, screening of host factors and antivirals.

Introduction:

First discovered in 1940[1], the influenza B virus has since been causing significant morbidity and
mortality in the global population[2]. As per the recent surveillance (seasons 2010–2018)
influenza B viruses are responsible for 15-30% of the total influenza like illness, with a number of
complications like fevers, body ache, fatigue and even life threatening acute respiratory distress
syndrome for patients having pre-existing lung diseases[3, 4]. There are two different lineages of
influenza B virus, Victoria and Yamagata, which circulate in the human population with various
degrees of predominance in different influenza seasons[5, 6]. Due to constant increase in the
influenza B virus related infections and limited cross protection offered by influenza B vaccine
against both of these lineages, there is a gradual transition from trivalent (against two subtypes
of FluA and one lineage of FluB) to quadrivalent (against two subtypes of FluA and two lineages
of FluB) flu shots offered across the world[7]. In spite of its immense importance in the context of
global healthcare ecosystem, influenza B virus research has drawn significantly lesser attention
in comparison to the closely related influenza A viruses, largely due to the scarcity of the tools
required to study virus replication cycle. This also severely restricts antiviral drug discovery
directed towards influenza B virus therapy[8].

Influenza viruses are segmented negative sense RNA viruses of the *Orthomyxoviridae* family.
Amongst the four types A, B, C and D, only influenza A and B cause human epidemics[9]. Viral

genome consists of eight different segments each of which remains enwrapped with multiple
copies of nucleoprotein (NP) in their oligomeric form and associates with a single copy of RNA
dependent RNA polymerase (RdRp) to form the ribonucleoprotein complexs or RNPs[10]. RNPs
are the self-sufficient machinery for driving different modes of RNA dependent RNA synthesis
events including viral gene expression and genome replication, hence reside at the center of virus
replication cycle[11]. This is why reporter RNP based assay systems remain one of the invaluable
tools for studying virus replication, host-pathogen interaction and high-throughput screening of
antivirals without handling infectious virus particles and hence avoiding biosafety associated
procedural complications[12, 13].
50
Influenza virus genomic segments are single stranded RNA that are devoid of the 5'-Cap and 3'-
Poly(A) tail structures[14]. Different segments harbor conserved untranslated regions of variable
lengths both at the 5' and 3' ends, which bracket single or multiple open reading frames (in the
antisense orientation) encoding viral proteins[14, 15]. Terminal regions of the 5' and 3' UTRs
contain complementarity resulting in a partial duplex structure (also known as "panhandle RNA"
or "cork-screw RNA") that serves as the promoter for RdRp[16]. Additionally, the UTRs contain
cis-acting elements, necessary and sufficient for transcription and replication of viral[17–19]and
non-viral reporter genes[12, 16, 20, 21]. Several groups have established reporter RNA based
assay systems where viral open reading frames have been replaced with reporter genes of
fluorescence or chemiluminescent proteins[12, 13, 16, 22–25]. These reporter RNA templates,
when expressed inside the cells in combination with NP and RdRp proteins, reconstitute reporter
RNPs. RNA synthesis activity of these RNPs could be measured by quantifying the extent of
reporter gene expression. Although appears to be straightforward, successful establishment of
reporter assay system requires (i) complicated cloning strategies to synthesize the reporter RNA
construct, (ii) construction of plasmids for expression of viral NP and RdRp subunits (PB1, PB2
and PA), and (iii) optimized expression of the reporter RNA and viral proteins in required
stoichiometric amounts that leads to the reconstruction of reporter RNPs with maximum efficiency.
68
So far, different strategies have been used to construct plasmids expressing influenza A and B
virus reporter RNA template (reporter plasmids)[12, 13, 24–26]. In few studies, reporter luciferase
gene was amplified using primers containing long overhangs corresponding to 3' and 5' UTR
regions of influenza A or B viruses; resulting PCR fragment harboring luciferase ORF flanked by
the viral UTRs were then inserted into the target vector for RNA polymerase-I driven expression
of the same, using conventional restriction digestion and subsequent ligation method[12, 21, 27].
Alternatively, viral 5' and 3'-UTR containing vectors (amplified using inverse PCR from the cDNA
clone of the corresponding segment) were ligated with reporter gene insert predigested with
compatible restriction enzyme sites[28]. In an independent cloning strategy, a double-stranded
DNA linker encompassing 5' and 3'-UTRs was inserted into the vector in-between the Pol-I
promoter and terminator sequences with the help of compatible restriction enzyme sites. The
reporter gene was then inserted between the UTRs using a second restriction enzyme site[24].
These restriction enzyme based cloning strategies are laborious and often introduce additional
nucleotides between the UTRs and the reporter gene which may interfere with the activity of the
cis/trans acting elements[24]. In order to avoid these constraints, restriction enzyme free cloning
methods utilizing vectors and inserts containing overlapping sequences have also been

implemented. For example, inserts containing reporter genes flanked by the 5' and 3'-UTRs were
created using long overhang primers (containing the UTR regions) which was then stitched to the
vector through the use of specialized proprietary enzymes/kits[13, 29]. With the inherent
limitations of the aforesaid cloning techniques, scarcity of information about the extensive
experimental protocol makes it difficult to establish and standardize the reporter based RNP
activity assay for influenza viruses. This situation gets further complicated for influenza B viruses
due to the larger size of the UTRs compared to the same for influenza A viruses.
92
Here we present a novel yet fairly simple cloning strategy, independent of any restriction enzyme
or specialized reagents or kits, to construct a firefly luciferase based reporter plasmid capable of
generating reporter genome template for influenza B/Brisbane/60/2008 virus. Additionally, we
present extensive standardization of this reporter plasmid based RNP activity assay through
optimization of various parameters regulating viral RNA synthesis. Using the reporter assay
system, we showed for the first time the effect of viral non-structural protein-1 (NS1) and host
protein kinase C delta (PKCD) upon influenza B virus RNA synthesis. We also demonstrated the
ability of this assay system to be used as a high throughput screening platform for the identification
of antiviral drugs specifically inhibiting RNA polymerase activity of the virus. Together this work
presents a great resource for cloning, standardization and implementation of reporter based RNP
activity assay for influenza and other related viruses.
104

Materials and methods:

Cell lines and Viruses
Human embryonic kidney 293T (HEK 293T) cells were maintained in Dulbecco's modified Eagle's
medium (DMEM; Invitrogen, Cat no. #12800017) supplemented with 10% (v/v) fetal bovine serum
(FBS; Invitrogen, Cat no. #10082147), 2mM GlutaMAX™ (Invitrogen, Cat no. #35050061), 1%
penicillin-streptomycin (Invitrogen, Cat no. #1514122) incubated at 37°C in a humidified 5% CO_2
incubator. Madin-Darby Canine Kidney (MDCK) cells were maintained in same conditions with
10% FBS (FBS; Invitrogen Cat no. #10270106). Influenza B/Brisbane/60/2008 virus was used in
this study.
114

Virus amplification and RNA Extraction
3×10^6 MDCK cells were seeded in 10 cm dishes, 24 hours before the infection. Prior to infection,
cell monolayer was washed with PBS twice and subsequently infected at an M.O.I. of 0.001. For
each 10cm plate, 1 mL virus inoculum was prepared in virus growth media (VGM; containing
DMEM, 0.2% bovine serum albumin (Sigma; Cat no. #A8412), 25 mM N-(2-hydroxyethyl)
piperazine-N′-ethanesulfonic acid (HEPES; Invitrogen, Cat no. #15630080) buffer, 2mM
GlutaMAX™, 1% penicillin-streptomycin and 0.5 µg/mL TPCK-trypsin (Thermo Scientific™, Cat
no. #20233)). Virus attachment was performed with 1ml of inoculum for 1 hour at 37°C in
humidified 5% CO_2 incubator with intermittent shaking at every 10 minutes to prevent drying of
cell monolayer and homogenous distribution of the inoculum. Post attachment, each 10 cm plate
was supplemented with 6 mL of VGM and incubated either at 37°C for Influenza A virus and 33°C
for Influenza B virus in humidified 5% CO_2 incubator. At 72 hours post infection, supernatant was
collected and centrifuged at 3200 g for 10 minutes at 4°C to remove the cell debris. The

supernatants were collected and aliquots were stored at 80°C refrigerator for further
applications[30].
130
Reverse transcription (RT)-PCR
Viral RNA was extracted from amplified virus stock using Trizol reagent (Invitrogen; Cat no.
#15596018). Reverse transcription was carried out using primer 'Uni9', which is complementary
to all of the individual genomic RNA segments of Influenza B/Brisbane/60/2008 virus[31] with
Verso cDNA Synthesis Kit (Thermo Scientific™; #AB1453A) according to manufacturer's
instructions. After the RT reaction, segment 6 (NA-NB) of the viral genome was amplified using
the 'NA-NB_F' and 'NA-NB_R' primers with the help of Phusion High-Fidelity DNA Polymerase
(Thermo Scientific™, Cat no. #F530S). The PCR product was purified using PCR Purification Kit
(Invitrogen; Cat No. # K310001) and the yield and quality of the purified product was checked
measuring the absorbance at 260 and 280nm and subsequently by running it on agarose gel.
141
142 **PCR and Cloning**
143
i)　　**Amplification of 5' and 3' UTR:** The RT-PCR amplified DNA corresponding to
segment 6 of viral genome was used as a template for the amplification of 5'UTR and
3'UTR using Phusion High-Fidelity DNA Polymerase using 5' UTR_F, 5' UTR_R and 3'
UTR_F, 3' UTR_R primers pairs and the PCR product were purified using Quick gel
extraction kit (Invitrogen: Cat no. # K210012). The purified double-stranded 5'UTR
fragment and 3'UTR fragment were then used as primers for the amplification of
luciferase gene described in the next section.
151
ii)　　**Preparation of Insert:** The luciferase ORF was amplified using the pHH21-vNA-Luc
as a template, kindly provided by Dr. Andrew Mehle. The double-stranded 5'UTR and
3'UTR fragments, synthesized in the previous step, were used as primers (5uM final
concentration) for the PCR amplification of Luciferase ORF using Phusion high fidelity
DNA polymerase, following manufacturer's protocol. The PCR product was analyzed
on 0.8% agarose gel and purified using PCR purification kit.
158
159　　**iii)**　**Preparation of Vector:** The vector was amplified using the pHH21-vNA-Luc as a
160　　　　　　template. PHH21_F' & 'PHH21_R' primers were used to amplify and linearize the
161　　　　　　vector using Phusion high fidelity DNA polymerase using 5x Phusion GC rich buffer
162　　　　　　following manufacturers' protocol. The PCR product was analyzed on 0.8% agarose
163　　　　　　gel and purified using PCR purification kit.
164
iv)　**Circular Polymerase Extension Cloning:** In the final CPEC assembly reaction, the
purified linearized vector and inserts are mixed together maintaining a vector to insert
molar ratio (V:I) = 1:1, 1:2 or 1:3, with the PCR reaction components, that consists of
0.4ul of 40mM dNTPs, 4ul of 5x Phusion HF Buffer, 0.25 µl of Phusion High-Fidelity
DNA Polymerase, 100ng of Vector and required amount of Insert to make the final
volume of 20ul. The composition of the CPEC reaction is identical to that of a standard
Phusion PCR mix, except that, there is a high concentration of dNTPs and no primers
are added. The thermal cycling conditions were as follows: 98 ∘C for 30 s, 25 cycles
of 98 ∘C for 10 s, 72 ∘C for 1 min 30 s, followed by 72 ∘C for 10 min. To assess if a
CPEC reaction is successful or not, 5ul of the product was analyzed by agarose gel

175	electrophoresis. The V:I=1:3 reaction showing highest intensity of the high molecular
176	weight band corresponding to the total length of vector and Insert was selected for
177	transformation.
178	

Transformation
E. Coli. DH5 alpha competent cells were prepared by modified rubidium chloride method as
described by Glover et al[32]. For transformation, 10ul of the CPEC reaction mixture were added
to the competent cells. Followed by an incubation of 30 minutes, cells were subjected to a brief
heat shock at 42^0C for 35 seconds followed by 5 minutes incubation in ice. Cells were then grown
in 400ul of Luria bertani broth (Himedia, Cat no. # M1245; 2.5% in double distilled water) at 37^0C
incubator for 1.5 hours with shaking at 220rpm and the entire culture volume was spread upon
LB-agar plates containing 100ug/ml of Ampicillin. Colony PCR was performed to identify positive
clones which were then verified through Sanger sequencing.
188

Generation of polymerase protein expressing plasmids
The PB2, PB1 and PA ORFs were cloned into the pCDNA-3X-FLAG vector (generously provided
by Dr. Andrew Mehle) which is a modified version of pcDNA3.1 (addgene) vector expressing
proteins under CMV promoter. This vector contains three FLAG epitopes joined in tandem (3X-
FLAG) after the NotI site at its MCS followed by a Cytosine. This results in the expression of a
protein having tri-alanine linker in between the individual ORFs and the C terminal 3X-FLAG tag.
For the expression of untagged version of each RdRp subunit, the stop codon has been kept
intact at the end of ORF. For the expression of V5 tagged NP, pcDNA3 vector has been modified
in order to have glycine-glycine-serine-glycine linker in between the ORF and the C terminal V5
epitope tag. Briefly, two primers of 58 nucleotide length were annealed to create double-stranded
piece of DNA having sticky ends on both sides (BamHI restriction site at the beginning of the
sequence and the NotI site at the end of the sequence). The thermal protocol for ramp down
annealing was as follows: 95 ∘C for 5 minutes, 70 cycles of 95^0 C (-1^0C/cycle) each for 1 minute,
followed by hold at 4∘C. The V5 linker was phosphorylated at 5' end by treatment with T4
Polynucleotide Kinase (PNK, Cat no. # M0201S). The pcDNA3.1 (addgene) vector was digested
with BamHI and NotI, treated with Calf alkaline phosphatase (CIP; NEB, Cat no. #M0290S) and
ligated with the V5 linker in V:I = 1:20 ratio.

Each individual insert fragment have been amplified using the cDNA template with the primers
containing restriction enzyme overhangs. The PB2, PB1 and PA have been amplified with EcoRI
and NotI overhang in two different PCR sets, one omitting the stop codon and the other including
the stop codon in the reverse primer. The NP have been amplified using primers with KpnI and
BamHI overhangs. For each amplification, 50 µl of PCR reaction consisted of 10 µl of 5x Phusion
HF buffer, 5 µl each of the forward and reverse 5 µM primers, 5 µl of 2mM dNTPs, 5 µl of cDNA
template, 19.5 µl sterile nuclease free water and 0.5 µl of Phusion High-Fidelity DNA
Polymerase. The modified pcDNA3-3X-FLAG have been digested with EcoRI & NotI (NEB) and
the pcDNA3-V5 vector has been digested with KpnI & BamHI, followed by treatment with CIP.
The digested vectors as well as insert fragments were gel excised and ligated in a vector to insert
ratio of 1:3 using T4 DNA ligase (Thermo Scientific™, Cat no. # EL0011) as per manufacturer's
protocol. 10 ul of ligation mixture was transformed into DH5 alpha competent cells. Sequences of
all the clones have been verified by sequencing.

Transfection
For examining protein expression level of each plasmid, HEK 293T cells were transfected using
lipofectamine3000 (Invitrogen: Cat no. # L3000015) as per manufacturer's instructions. All the
plasmids were prepared using plasmid DNA isolation kit (Promega, Cat No. # A1222). The

pcDNA3.1 blank vector was used in the control cells. The media was changed 12 hours post
transfection and incubated for 36 (or stated otherwise) hours following transfection.
225
Western Blot
Protein levels for transiently transfected cells were assessed by Western blotting. Transfected
cells were lysed for 20 minutes in pre-chilled Co-Immunoprecipitation buffer (50mM Tris-HCl pH
7.4, 150mM NaCl, 0.53% NP-40) containing protease inhibitors (Roche-Sigma, Cat no. #
11873580001) and phosphatase inhibitor and subsequently total protein samples were separated
via 8% SDS-PAGE and transferred to polyvinylidene difluoride (PVDF, Bio-Rad, Cat no.
#1620177) membrane using transfer buffer (25 mM Tris, 191 mM glycine, 0.025% SDS & 10%
methanol (vol/vol) in Trans-Blot Turbo Transfer System (Bio-Rad). After incubation with 5% nonfat
milk in TBST (10 mM Tris, pH 8.0, 150 mM NaCl, 0.5% Tween 20) for 60 min, the membrane was
washed once with TBST and incubated with antibodies against 3xFLAG (1:5000, Sigma, Cat no.#
F3165), V5 (1:5000, CST, Cat no. # D3H8Q), HA (1:5000, Cat no. # C29F4), BNP (1:5000,
generated in collaboration with BioBharati LifeScience Pvt. Ltd, India), at 4 °C for 12 h.
Membranes were washed with TBST three times for 5 minutes and incubated with a 1:25,000
dilution of horseradish peroxidase-conjugated anti-mouse (Sigma, Cat no. # A9044-2ML6) or anti-
rabbit (Sigma, Cat no. # A0545-1ML) antibodies for 1 h. Blots were washed with TBST three times
for 5 minutes and developed with the ECL system (ThermoFisher, Cat No. #34095) according to
the manufacturer's protocols.
243
Polymerase activity assay
HEK293T cells were co-transfected with 94.11 ng of each of the pcDNA3-PB2-FLAG, pcDNA-
PB1, 11.76 ng of pcDNA-PA, 100 ng of pcDNA3-BNP and 100 ng of pHH21-vNA-Luc plasmids.
The additional plasmids were used in various amount upto 150 ng and is topped up by blank
vector in order to keep the amount of DNA same in all the sets. In absence of additional plasmid,
the total amount of RNP reconstitution plasmids were increased upto 500ng keeping the ratio of
each RNP component same (1/4th BNP, 1/4th pHH21-vRNA-Luc, and the half of the total amount
of DNA will be divided as 8:8:1 ratio for PB2, PB1 and PA). The transfection mix were prepared
with Lipofectamine 3000 as per manufacturer's protocol using optiMEM (Thermo Scientific™, Cat
no. #31985-070). At 12 hours, the media was changed very carefully without dislodging any cell
that may result in error in further signal. The cells were harvested at 36 hours (or as mentioned)
post transfection and luciferase activity assay was performed using Promega Luciferase Assay
System (Promega, #E1500 & #E1910), as per manufacturer's protocol using luminometer
(Promega Glomax 20/20).
258
Ribavirin and Favipiravir dose-response assays in HEK 293T cells
Ribavirin (Sigma-Aldrich, Cat no. # R9644) was dissolved in water to prepare 80 mM stock and
Favipiravir (MedChemExpress, Cat no. #HY-14768) was dissolved in DMSO to prepare a 200
mM stock which were aliquoted and stored at – 80^{0}C until used. 0.2 x 10^{6} HEK 293T cells were
seeded in 24-well plates and post 24 hours were treated with specified concentrations of ribavirin
or favipiravir for 2.5 h at 37^{0} C with 5% CO2. Subsequently, cells were transfected with
lipofectamine 3000 as per manufacturers protocol and incubated in fresh media containing
specified concentrations of drugs were for 36 hours. Polymerase activity was then assayed as
described above. The IC50 value was calculate by fitting the data to four parameter nonlinear
equation.
269
3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium Bromide (MTT) Assay
30,000 HEK293T cells were seeded in 96-well plate. After 24 hours of seeding, the cells were
treated with different concentrations of drugs in triplicates for 36 hours. Post-treatment, 100 µL of
MTT reagent (5 mg/mL, SRL) dissolved in phosphate-buffered saline (PBS) was added to the

cells and incubated for 3 h at 37 °C. Subsequently, the reagent was removed and the formazan
crystals were dissolved by adding 100 µL of dimethyl sulfoxide (DMSO) (Sigma) in each well. The
absorbance of the suspension was measured at 595 nm using an Epoch 2 microplate reader
(BioTek Instruments). The percentages of metabolically active cells were compared with the
percentage of control cells treated with vehicle control.
279
Transfection-Infection assay
0.1×10^6 HEK 293T cells were seeded in 48-well plates. After 20 hours of seeding, the cells were
transfected with 0.25 ug of reporter plasmid using lipofectamine 3000 and 22 hours post
transfection, the cells were pre-treated with half-maximal inhibitory concentrations (IC50) of the
drugs (Ribavirin:18.54 uM; Favipiravir:25.46 uM) for 2.5 h at 37^0 C with 5% CO2. Following 2.5
hours of treatment, the cells were infected with Influenza B virus at an MOI of 0.1 in presence of
drugs and the polymerase activity was assayed at 16 hours of post infection.
287
Statistical analysis
The arithmetic mean and standard deviation of the firefly luciferase signal were calculated from
three biological replicates for each experiment. The data were plotted in bar diagram with
standard deviations as error bars. For dual luciferase assay, the firefly luciferase signal was
normalized with the *Renilla* luciferase signal for each replicate. In firefly luciferase assays
involving the host factors, viral factors, and antiviral molecules and dual luciferase assays;
normalized mean and standard deviation were calculated against the control. The normalized
mean was calculated by dividing the arithmetic mean of the experimental sets by the mean of the
control set and converting it to a percentage value. Normalized standard deviation was calculated
by normalizing the coefficient of variations against the control and augmenting it with the normal
mean. A two-tailed Student's T-test was performed for comparison of the data sets.
299
Results:

Generation of influenza B virus reporter plasmid for expression of viral reporter RNA in
mammalian cells

In order to establish a reporter based RNP activity assay, a template RNA harboring the reporter
gene flanked by the viral UTRs needs to be expressed under the control of RNA polymerase I
promoter. This ensures that the reporter RNAs are devoid of any 5'- or 3'- terminal modifications,
hence mimicking authentic viral genomic RNA. To achieve this, firstly we have constituted the
"insert" harboring the firefly luciferase gene in reverse orientation flanked by viral 5'- and 3'- UTRs.
Subsequently, this cassette was introduced into the pHH21 vector in between the RNA
polymerase I promoter and terminator. The entire process of constituting authentic viral UTRs,
assembling them with the reporter gene and introducing this cassette into the pHH21 vector
utilized a single DNA polymerase enzyme without the need of any restriction enzyme or
specialized kits (Figure 1).

Viral genomic RNA, purified from influenza B/Brisbane/60/2008 virus particles (Figure 1A), was
used as a source for the amplification of the 144nt 5'- and 95nt 3'-UTRs, using sequence specific
primers with 15-20 nucleotide overhanging sequences corresponding to the vector and the
reporter gene (Figure 1B, 2A). The resulting PCR products thus contain (i) viral 5'-UTR flanked
by the overlapping sequences with the Pol-I promoter and 3'-termini of the reporter gene and (ii)
viral 3' UTR region flanked by the overlapping sequence with 5'-termini of reporter gene and Pol-
I terminator. These double-stranded PCR products were then used as primers to amplify the

firefly luciferase gene from the pHH21-vNA-Luc plasmid, kindly provided by Prof. Andrew Mehle,
University of Wisconsin Madison (Figure1C, 2B). The final PCR product, constitutes reporter gene
flanked by viral 5'- and 3'- UTR regions along with partial sequences from the Pol-I promoter and
terminator regions at the extreme 5'- and 3'- termini respectively. In order to synthesize the final
reporter plasmid construct, named as pHH21-BNA-Luc, this cassette was inserted into the pHH21
vector (amplified in a separate PCR reaction; Figure 2C) using the Circular Polymerase Extension
Cloning (CPEC), as originally described by Quan et al[33, 34] (outlined in the Figure 1D). A vector
to insert molar ratio of 1:3 generated maximum amount of assembled product (Figure 2D).
Reaction product was transformed in chemically competent *E. coli* and successful incorporation
of the insert was confirmed by the colony PCR screening method. All the PCR amplifications were
performed using a single Phusion High-Fidelity DNA polymerase as described in further detail in
the methods section.

To reconstitute functional reporter RNPs inside the cells, reporter RNA template needs to be co-
expressed with NP and the RdRp subunits, PB1, PB2 and PA (Figure 1E, F). The RdRp subunits
were cloned into the pCDNA-3X-FLAG vector (generously provided by Dr. Andrew Mehle) under
the control of CMV promoter with the help of EcoRI and NotI restriction enzymes, which results in
incorporation of a tri-alanine linker in between the individual ORFs and the three FLAG epitopes
joined in tandem (3X-FLAG). For expression of untagged proteins, ORFs with the stop codon
were cloned using the same strategy. The NP gene was cloned into a modified pcDNA3 vector
harboring V5 epitope tag (mentioned in the methods section) with the help of the KpnI and BamHI
sites, with a glycine-glycine-serine-glycine linker in between the ORF and the V5 epitope tag.

Standardization of influenza B virus RNP activity assay

Influenza B virus reporter RNPs were reconstituted in HEK293T cells through transient
transfection of the reporter plasmid (pHH21-BNA-Luc) either in the absence (negative control) or
the presence of the plasmids expressing PB1, PB2, PA and NP proteins (Figure 3A). Cells were
harvested at 24 hours of post-transfection and luciferase activity was measured to quantitate the
Influenza B RNP activity. To our surprise, the positive control set showed only 10^4 signal
(luciferase light unit or RLU) which is only two log higher than the negative control set (10^2 RLU),
suggesting suboptimal activity of the reconstituted RNPs. This could be due to the poor
expression levels of the RdRp subunits, PB1 and PB2, in comparison to the PA and NP proteins,
as observed from the western blot analysis using specific antibodies, hence prohibiting the
successful assembly of reporter RNPs inside the cell. To investigate this further, we examined
the sequence of the constructs carefully and noticed that all of the protein expression plasmids
lack the Kozak sequence which may result in their suboptimal translation. The PA gene have a
Guanosine after the ATG sequence which is a part of Kozak sequence. PB1 and PB2 do not have
this Guanosine after the start codon ATG. Thus half Kozak sequence is conserved in PA gene
which makes its expression better than PB2 and PB1.

Figure 1: An illustrative overview of cloning strategy of influenza B reporter plasmid and reporter assay (A) Total RNA
was isolated from amplified stocks of Influenza B/Brisbane/60/2008 virus (B) total RNA was converted to cDNA by
performing RT-PCR and 5' & 3' UTRs were amplified using specific primers containing overhangs. (C) The double-
stranded 5' and 3' UTRs containing overlapping regions were used as primers to amplify the luciferase ORF. (D) The
resulting PCR product was used as an insert for CPEC assembly with the PCR amplified vector fragment. (E-F) The
generated reporter and other protein expressing plasmids upon co-transfection in HEK293T cells reconstitute the
luciferase RNP's that express luciferase enzyme under the control of the viral promoter. The quantification of the
luciferase signal gives the measure of viral polymerase activity.

To address this, we performed site-directed mutagenesis to introduce partial Kozak sequences in
each of these plasmids without any alteration in the ORF and repeated the polymerase activity
assay with them. As evident from figure 3A, the introduction of the Kozak sequence significantly
boosted the expression of all of the RNP proteins which together resulted in reporter activity of
10^6 RLU, four logs higher than the negative control set. Interestingly, the expression levels of
the PA subunit still remained severalfold higher than the other two subunits of RdRp, PB1 and
PB2 (Figure 3A).

373
374 **Figure 2:** PCR amplification and CPEC reaction for construction of reporter construct: Agarose gel electrophoresis
375 images of (A) PCR amplification products corresponding to the 5'UTR and 3'UTR of NA-NB segment. (B) PCR
376 amplification product of luciferase ORF using double-stranded PCR products corresponding to 5' and 3' UTRs as
377 primers. (C) PCR amplification of pHH21 vector. (D) CPEC products with different ratio of vector to insert.
378

Precise abundance of the PB1, PB2 and PA subunits in equimolar amounts is a pre-requisite for
the successful assembly of the heterotrimeric RdRp complex and hence, reconstitution of reporter
RNPs to optimum levels. Therefore, we tried to optimize the amount of the plasmids to be
transfected in order to have a comparable expression of the RdRp subunits. Reporter RNPs were

reconstituted using different ratios of RdRp subunit plasmids, while keeping the amount of the
reporter RNA and NP plasmid constant. As shown in Figure 3B, increasing the amount of PB1
and PB2 expressing plasmids compared to the PA led to a gradual increase in reporter activity
and a ratio of 8:8:1 for PB1: PB2: PA resulted in comparable expression of all three polymerase
subunits and maximum reporter activity. Subsequently, keeping the ratio of the polymerase
subunit plasmids constant, we increased the amount of the NP expressing plasmid, which
resulted in increase in the reporter activity, hence stretching the sensitivity of this reporter assay
to the maximum level (Figure 3C). The NP to polymerase proportion up to 1:2 results in increase
in polymerase activity. Further increase in the amount of NP does not result in substantial increase
in polymerase activity. Hence for our further experiments, we have used this ratio of RNP
reconstituting plasmids. Once we optimize the amounts of various plasmids reconstituting reporter
RNPs, we have performed a time kinetics experiment in order to assess the optimum time
required to obtain a signal of 10^6 RLU or more (Figure 3D).

Figure 3: Optimization of the reporter system: (A) Effect of Kozak sequence on expression of viral polymerase proteins
and influenza B RNP activity assay (B) Reporter RNP activity assay with different ratio of PA protein expression plasmid
with respect to PB1 and PB2. (C) Reporter RNP activity assay with various amount of NP expression plasmid. (D)
Optimization of time for reporter activity assay (E) Optimization of incubation temperature for influenza B RNP activity

assay (n=3 ± standard deviation, *p<0.05 one-way ANOVA with post hoc Student's t-test when compared to the
preceding set, for Figure E, comparison was performed in between two RNP positive sets, ns = not significant).

A time dependent increase in the reporter activity was observed which reaches a plateau by 42
hours of post transfection. Additionally, influenza B polymerase activity was assessed at different
temperatures (33^0C and 37^0C) by reconstituting the polymerase through transient transfection at
37^0C for 12 hours followed by an additional incubation of 30 hours at respective temperatures
(Figure 3E). As observed at 37^0C, reporter activity was almost two fold higher than the activity at
33^0C, a data corroborated perfectly with the previous results obtained by Santos et al[35].
Together, we present a fast sensitive and high throughput reporter assay for monitoring influenza
B virus RNA synthesis in an infection free setting.

All of the assays were performed in 24, 48 and 96 well plates, in triplicates for each of the
biological sets (data presented in this manuscript is from 24 well plates), hence confirming that
this assay system is high-throughput compatible. Additionally, five log difference between the
signal and background readouts provides a wide dynamic range for this reporter based assay
system. Together, we have been able to establish a fast, reliable and high-throughput compatible
assay system for monitoring influenza B virus RNP/ polymerase activity, which is suitable for
assessing the effect of various viral or cellular factors in modulating RNP activity and hence viral
RNA synthesis.

The influenza B RNP activity assay is suitable for evaluating the efficacy of viral or host
factors in regulating viral RNA synthesis.

To this end, we set out to evaluate the efficacy of the newly developed polymerase activity assay
in identifying novel viral and host factors that may regulate viral RNA synthesis. Influenza virus
Nonstructural Protein 1 (NS1) is a multifunctional protein participating mainly in the suppression
of antiviral defense mechanisms exerted by a wide variety of host factors [36, 37]. Additionally,
influenza A virus NS1 protein has been shown to boost viral RNA synthesis[38–40], possibly
through interfering with antiviral activity DDX21 and RAP55 [41, 42]. While the immune
suppression activity of influenza B NS1 was well studied[37, 43, 44], little is known about the role
of NS1 in regulating influenza B virus RNA synthesis. Hence, we evaluated the ability of influenza
B virus NS1 protein to promote viral RNA synthesis with the newly developed reporter RNP activity
assay. Influenza B virus NS1 ORF was cloned into the pCDNA-3X-FLAG vector that resulted in
the expression of the C-terminally FLAG tagged NS1 protein. Influenza B reporter RNPs were
reconstituted in HEK293T cells either in the absence or presence of increasing amounts of NS1
protein and reporter activity was monitored to assess the extent of viral RNA synthesis. Increasing
amount of NS1 resulted in 1.5 to 2 folds increase in reporter activity (Figure 4A) establishing it as
a positive regulator of viral RNA synthesis. Furthermore, reconstituting reporter RNPs in the
presence of NS1 presents an assay system that closely resembles RNA synthesis, which occurs
during the course of infection.

Subsequently, we tested the ability of a host factor to regulate influenza B virus RNA synthesis
using our reporter RNP activity assay. Host Protein Kinase C, specifically the delta isoform, has
been shown to positively influence influenza A virus RNA synthesis by regulating the
phosphorylation and subsequent assembly of viral nucleoproteins into RNPs. Interestingly, the
constitutively active catalytic domain of PKC delta (PKCD), when overexpressed, negatively

regulates influenza A virus RNA synthesis [45]. To determine the role of PKCD in regulating
influenza B virus RNA synthesis we employed the newly developed reporter RNP activity assay.
As evident from figure 4B, increasing amounts of PKCD resulted in a gradual decrease in RNP
activity and hence viral RNA synthesis without any severe impact upon the translation of viral
proteins. These data not only substantiates the role of PKCD in regulating influenza B virus RNP
activity, but also validates the efficacy of our assay system in studying the effect of pro- or antiviral
factors regulating viral RNA synthesis.

Figure 4: Effect of host & viral factors upon viral RNA synthesis: (A) Effect of an increasing amount of viral NS1 protein
on Influenza B RNP activity assay (B) Effect of an increasing amount of constitutively active host protein kinase c delta
(PKCD) protein on B RNP activity assay. (n=3± standard deviation. *p<0.05 one-way ANOVA with *post hoc* student's
t-test when compared to the preceding set.

Reporter based RNP activity assay as high throughput screening platform of antiviral drugs.

Finally, we intend to establish the suitability of the RNP activity assay as a high throughput
screening platform of antiviral drugs that can inhibit viral RNA synthesis and hence virus
replication. Ribavirin and Favipiravir are nucleoside (purine) analogues, which inhibit the
replication of a wide variety of RNA viruses by acting as an alternative substrate for viral RNA
polymerase[46, 47]. Additionally, Ribavirin also inhibits inosine monophosphate

Figure 5: Effect of antiviral drugs upon viral RNA synthesis in infection-free and infection setting: (A, B) MTT assay to
determine the cytotoxicity of Ribavirin and Favipiravir on HEK293T cells. (C, D & E) Effect of Ribavirin and Favipiravir
on influenza B virus RNP activity. Viral polymerase proteins in HEK293T cells are expressed either by transient
transfection (C, D) or by infecting the cells with Influenza B virus (E). (n=3 ± standard deviation, *p<0.05 one-way
ANOVA with post hoc Student's t-test when compared to the preceding set, for Figure E, comparison was performed
with control set, ns = not significant).

dehydrogenase thereby depleting the GTP and creating an imbalance in the nucleotide pool
inside the cell[48]. Both Ribavirin and Favipiravir has been approved as chemoprophylaxis as well
as therapy against influenza A and B viruses[49–51]. Hence, we used these two drugs as positive
controls to test the efficacy of our assay system for antiviral screening. MTT assay was performed
in HEK293T cells (Figure 4A, B), where neither of the drugs show any cytotoxicity. HEK293T cells
were pretreated with different concentrations of the drugs followed by forward transfection to
reconstitute the influenza B reporter RNPs and subsequent incubation with the drugs for 36 hours.
Reporter activities were measured and expressed as relative percentages with respect to the
vehicle control. Data presented in Figure 5 C, D, shows a dose-dependent decrease in the
reporter activity and hence viral RNA synthesis with increasing amounts of the drugs with IC50
values of 18.54 uM and 25.46 uM for Ribavirin and Favipiravir respectively.

To further extend the scope of the assay system, we sought to check if this system is capable of
assessing the effect of host factors or antivirals upon the overall progress of infection. For this
purpose, we transfected HEK293T cells with the reporter construct and subsequently infected

them with influenza B virus at 20 hours post transfection. It is expected that in infected cells
reporter RNA template will get transcribed with the help of RdRp and NP proteins expressed from
viral genomic RNA segments. As evident from Figure 5E, infected cells supported successful
generation of reporter RNPs and hence showed high reporter activity, while the uninfected cells
showed no such effect. Interestingly, when parallel set of cells were treated with Ribavirin and
Favipiravir prior to infection with Influenza B virus, significant reduction in reporter activity were
observed, hence suggesting an overall reduction in viral gene expression and hence virus
replication in presence of the drugs. Together our data reconfirms the activity of the two well
established antiviral drugs against influenza B virus RNA synthesis machinery and also
establishes the newly developed reporter based influenza B virus RNP activity assay as a high
throughput screening platform of antivirals specifically inhibiting viral RNA synthesis.

494
Discussion:

Luciferase based reporter assay systems remain a key tool for analyzing gene expression in a
wide variety of organisms; viruses are not exceptions. While, for positive sense RNA viruses,
introduction of the single sub-genomic reporter RNA template in cells is sufficient for expression
of reporter genes; for negative sense RNA viruses, RNP associated viral proteins needs to be
synthesized along with the reporter RNA in order to reconstitute complete RNPs, which then leads
to the expression of reporter enzyme as a proxy of viral genes[12, 52, 53]. This is why, successful
reconstruction of reporter viral RNPs require extensive cloning of multiple RNA and protein
expressing constructs, standardization of their expression in right stoichiometric ratios and
optimization of other crucial parameters like time, temperatures etc. Although, several groups
have reported reporter assay systems for monitoring influenza A and B virus RNA synthesis, non-
availability of detailed methodical description makes the process of establishing the assay system
non-trivial[12, 13, 24–26]. In this work, we have established a firefly luciferase-based influenza B
virus RNP activity assay and presented the detailed methodology of the entire procedure which
could be easily followed for the development of such viral and non-viral reporter assay systems.

We have introduced a unique cloning strategy for the construction of the influenza B virus reporter
RNA construct that is devoid of restriction enzymes or any other specialized enzymes. This
cloning strategy utilizes a single DNA polymerase, which is widely used for regular molecular
biology work and hence easily available. Using this polymerase, two consecutive PCR
amplification reactions led to the generation of the reporter RNA cassette encompassing the
reporter ORF flanked by viral 5'- and 3'-UTR regions which were then inserted into the vector
using CPEC cloning method. While the vector and the insert used for CPEC, are also compatible
for Gibson assembly based cloning method, we intentionally avoided use of any specialized
enzymes to make the overall procedure simple and user-friendly that could be adapted for cloning
of any other reporter RNA constructs. In addition to reporter RNA construct, we also cloned ORFs
corresponding to viral PB1, PB2, PA and NP proteins and optimized their expression to
reconstitute reporter RNPs at maximum levels. The robustness of this assay system was
substantiated by testing the efficacy of antiviral drugs, Ribavirin and Favipiravir, to inhibit influenza
B virus RNA synthesis either in the context of reconstituted RNPs (through transfection) or during
the course of infection. The fact that the reporter RNA template can be preferentially recognized

by viral NP and RdRp subunits to reconstitute reporter RNPs during the course of infection,
confirms that the reporter RNA mimics viral genomic RNA segments and hence validates its
suitability to be used for the study of viral RNA synthesis and effect of various viral and host
factors upon the same. In fact, using the newly developed reporter RNP system, we for the first
time showed that viral NS1 protein can boost influenza B virus RNA synthesis and constitutively
active form of host PKCD can downregulate the same. While effect of NS1 and PKCD proteins
has been previously characterized in case of influenza A virus[38–40, 45], our results
substantiates that these proteins participate similarly regulate influenza B virus RNA synthesis as
well.

Altogether, we present a comprehensive roadmap for development, characterization and
validation of a reporter-based Influenza B virus polymerase/ RNP activity assay and made it
generic enough to be followed by others who intend to develop similar assay systems for influenza
and other negative sense RNA viruses. We also made all the resources publicly available (upon
request) to enrich the armoury for combating influenza B viruses and hope that it will be widely
utilized to identify new therapeutic strategies against this deadly human pathogen.

Table 1. Primers used in this study.

Name of the primer	Sequence (5'->3')
Uni 9	AGCAGAAGC
NA-NB_F	AGTAGTAACAAGAGCATTTTTCAG
NA-NB_R	AGCAGAAGCAGAGCATC
5' UTR_F	CATTTTGGGCCGCCGGGTTATTAGTAGTAACAAGAGCATTTTTCAG
5' UTR_R	CGGAAAGATCGCCGTGTAATGGAGGAATGGTTGAGTC
3' UTR_F	CTTTATGTTTTTGGCGTCTTCCATTGTTCATTTTTGGCCTATTTG
3' UTR_R	CCTCCGAAGTTGGGGGGGGAGCAGAAGCAGAGCATCTTC
PHH21_F	CCCCCCCAACTTCGGAGG
PHH21_R	AATAACCCGGCGGCCCAAAATG
PHH21 SEQ2	AAAACGCTGGGCGTTAATCAAAGAGGCG
PHH21 SEQ1	GGGGGACACTTTCGGACATCTGGTC
pcDNA3-V5 For	GATCCGGAGGTAAGCCTATCCCTAACCCTCTCCTCGGTCTCGATTCTACGTAGTAAGC
pcDNA3-V5 Rev	GGCCGCTTACTACGTAGAATCGAGACCGAGGAGAGGGTTAGGGATAGGCTTACCTCCG
NP_For	ATTCGGGGTACCGCCACCATGTCCAACATGGATATTGACG
NP_Rev_V5	ATTCGCGGATCCACCATAATCGAGGTCATCATAATCCTC
NP_Rev_Stop	ATTCGCGGATCCTTAATAATCGAGGTCATCATAATCCTC
PB1_For	TAAGCGGAATTCACCATGAATATAAATCCTTATTTTCTCTTC
PB1_Rev_FLAG	ATTGAGGCGGCCGCTATGTACCCAATCTCACCAAG
PB1_Rev_Stop	ATTGAGGCGGCCGCTTATATGTACCCAATCTCACC
PB2_For	TAAGCGGAATTCACCATGACATTGGCCAAAATTGAATTG
PB2_Rev_FLAG	ATTGAGGCGGCCGCGCTCAAGGCCCACCCC
PB2_Rev_Stop	ATTGAGGCGGCCGCTTAGCTCAAGGCCCACCC
PA_For	TAAGCGGAATTCACCATGGATACTTTTATTACAAGAAACT
PA_Rev_FLAG	ATTGAGGCGGCCGCTTCGTCCATAATCTCGTC
PA_Rev_Stop	ATTGAGGCGGCCGCTTATTCGTCCATAATCTCGTC

Table 2: PCR conditions used in the study for amplification of individual inserts.

PCR amplification	Forward Primer	Reverse Primer	Denaturation (temp./duration)	Cyclic denaturation, annealing and elongation (temp./duration)	No. of cycles	Final elongation
NA-NB fragment	NA-NB_F	NA-NB_R	98^0C/30 seconds	98^0C/10 seconds 60^0C/30 seconds 72^0C/95 seconds	35	72^0C/5 minutes
5' UTR	5' UTR_F	5' UTR_R	98^0C/30 seconds	98^0C/10 seconds 60^0C/25 seconds 72^0C/5 seconds	35	72^0C/5 minutes
3' UTR	3' UTR_F	3' UTR_R	98^0C/30 seconds	98^0C/10 seconds 59^0C/25 seconds 72^0C/5 seconds	35	72^0C/5 minutes
Insert for Reporter plasmid	5' UTR double-stranded PCR product	3' UTR double-stranded PCR product	98^0C/30 seconds	98^0C/10 seconds 72^0C/80 seconds	35	72^0C/5 minutes
Vector for Reporter plasmid	PHH21_F	PHH21_R	98^0C/30 seconds	98^0C/10 seconds 67^0C/30 seconds 72^0C/90 seconds	35	72^0C/5 minutes
PB1-FLAG	PB1_For	PB1_Rev_FLAG	98^0C/30 seconds	98^0C/10 seconds 60^0C/30 seconds 72^0C/90 seconds	35	72^0C/10 minutes
PB1-STOP	PB1_For	PB1_Rev_Stop	98^0C/30 seconds	98^0C/10 seconds 58^0C/30 seconds 72^0C/90 seconds	35	72^0C/10 minutes
PB2-FLAG	PB2_For	PB2_Rev_FLAG	98^0C/30 seconds	98^0C/10 seconds 69^0C/30 seconds 72^0C/90 seconds	35	72^0C/10 minutes
PB2-STOP	PB2_For	PB2_Rev_Stop	98^0C/30 seconds	98^0C/10 seconds 69^0C/30 seconds 72^0C/90 seconds	35	72^0C/10 minutes
PA-FLAG	PA_For	PA_Rev_FLAG	98^0C/30 seconds	98^0C/10 seconds 60^0C/30 seconds 72^0C/90 seconds	35	72^0C/10 minutes
PA-STOP	PA_For	PA_Rev_Stop	98^0C/30 seconds	98^0C/10 seconds 58^0C/30 seconds 72^0C/90 seconds	35	72^0C/10 minutes
NP-FLAG	NP_For	NP_Rev_V5	98^0C/30 seconds	98^0C/10 seconds 64^0C/30 seconds 72^0C/90 seconds	35	72^0C/10 minutes
NP-STOP	NP_For	NP_Rev_Stop	98^0C/30 seconds	98^0C/10 seconds 66^0C/30 seconds 72^0C/90 seconds	35	72^0C/10 minutes

Declaration of competing interest:

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgement:

We sincerely acknowledge Prof. Andrew Mehle (University of Wisconsin Madison) for proving us various valuable plasmids. We acknowledge Indian Institute of Technology Kharagpur for providing us infrastructural support. A.M. acknowledges DBT, Ramalingaswami re-entry

fellowship (BT/RLF/Re-entry/02/2015), DST-SERB, Early Career Research Award (ECR/2017/001896) and MHRD, "Scheme for Transformational and Advanced Research in Science" {STARS/APR2019/BS/369/FS (Project ID: 369)} for financial support. S.B. (File number 09/081(1301)/2017-EMR-I) and N.K. (File number 09/081(1316)/2017-EMR-I) acknowledge the Council of Scientific and Industrial Research for their fellowship.

Address of correspondence:

Dr. Arindam Mondal, Assistant Professor, School of Bioscience, Indian Institute of Technology Kharagpur, Kharagpur-721302, Dist- Paschim Medinipur, West Bengal, India. E-mail address: arindam.mondal@iitkgp.ac.in.

References:

1. Francis T (1940) A new type of virus from epidemic influenza. Science (80-) 92:405–408 . doi: 10.1126/SCIENCE.92.2392.405

2. Sharma L, Rebaza A, Cruz CS Dela (2019) When "B" becomes "A": the emerging threat of influenza B virus. Eur Respir J 54: . doi: 10.1183/13993003.01325-2019

3. U.S. Influenza Surveillance: Purpose and Methods | CDC. https://www.cdc.gov/flu/weekly/overview.htm. Accessed 2 Nov 2021

4. Flu Symptoms & Complications | CDC. https://www.cdc.gov/flu/symptoms/symptoms.htm. Accessed 2 Nov 2021

5. JA M, T S, AR I (2004) Multiple genotypes of influenza B virus circulated between 1979 and 2003. J Virol 78:12817–12828 . doi: 10.1128/JVI.78.23.12817-12828.2004

6. Xu C, Chan K-H, Tsang TK, et al (2015) Comparative Epidemiology of Influenza B Yamagata- and Victoria-Lineage Viruses in Households. Am J Epidemiol 182:705–713 . doi: 10.1093/AJE/KWV110

7. Ambrose CS, Levin MJ (2012) The rationale for quadrivalent influenza vaccines. Hum Vaccin Immunother 8:81 . doi: 10.4161/HV.8.1.17623

8. Dumm RE, Heaton NS The Development and Use of Reporter Influenza B Viruses. doi: 10.3390/v11080736

9. Krammer F, Smith GJD, Fouchier RAM, et al (2018) Influenza. Nat. Rev. Dis. Prim. 4:1–21

10. Bouvier NM, Palese P (2008) THE BIOLOGY OF INFLUENZA VIRUSES. Vaccine 26:D49

11. H F, AP W, L C, et al (2019) Structures of influenza A virus RNA polymerase offer insight into viral genome replication. Nature 573:287–290 . doi: 10.1038/S41586-019-1530-7

12. Lutz A, Dyall J, Olivo PD, Pekosz A (2005) Virus-inducible reporter genes as a tool for detecting and quantifying influenza A virus replication. J Virol Methods 126:13–20 . doi: 10.1016/j.jviromet.2005.01.016

13. Zhu W, Zhou J, Qin K, et al (2011) A reporter system for assaying influenza virus RNP functionality based on secreted Gaussia luciferase activity. Virol J 8:29 . doi: 10.1186/1743-422X-8-29

14. Dadonaite B, Gilbertson B, Knight ML, et al (2019) The structure of the influenza A virus genome.
Nat Microbiol 2019 411 4:1781–1789 . doi: 10.1038/s41564-019-0513-7

15. Szymkowiak C, Kwan WS, Su Q, et al (2003) Rapid method for the characterization of 3′ and 5′
UTRs of influenza viruses. J Virol Methods 107:15–20 . doi: 10.1016/S0166-0934(02)00184-2

16. Flick R, Hobom G (1999) Interaction of influenza virus polymerase with viral RNA in the
'corkscrew' conformation. J Gen Virol 80:2565–2572 . doi: 10.1099/0022-1317-80-10-2565

17. Ferhadian D, Contrant M, Printz-Schweigert A, et al (2018) Structural and Functional Motifs in
Influenza Virus RNAs. Front Microbiol 0:559 . doi: 10.3389/FMICB.2018.00559

18. Park YW, Katze MG (1995) Translational Control by Influenza Virus IDENTIFICATION OF CIS-
ACTING SEQUENCES AND TRANS-ACTING FACTORS WHICH MAY REGULATE SELECTIVE
VIRAL mRNA TRANSLATION*. doi: 10.1074/jbc.270.47.28433

19. Fodor E, Palese † Peter, Brownlee GG, et al (1998) Attenuation of Influenza A Virus mRNA Levels
by Promoter Mutations. J Virol 72:6283 . doi: 10.1128/jvi.72.8.6283-6290.1998

20. Neumann G, Zobel A, Hobom G (1994) RNA Polymerase I-Mediated Expression of Influenza Viral
RNA Molecules. Virology 202:477–479 . doi: 10.1006/VIRO.1994.1365

21. Sherry L, Smith M, Davidson S, Jackson D (2014) The N Terminus of the Influenza B Virus
Nucleoprotein Is Essential for Virus Viability, Nuclear Localization, and Optimal Transcription and
Replication of the Viral Genome. doi: 10.1128/JVI.01542-14

22. Luytjes W, Krystai M, Enami M, et al (1989) Amplification, Expression, and Packaging of a Foreign
Gene by Influenza Virus. Cell 59:1107–1113

23. Parvin JD, Palese P, Honda A, et al (1989) Promoter analysis of influenza virus RNA polymerase.
J Virol 63:5142–5152 . doi: 10.1128/JVI.63.12.5142-5152.1989

24. Li Y, Larrimer A, Curtiss T, et al Influenza virus assays based on virus-inducible reporter cell lines.
doi: 10.1111/j.1750-2659.2009.00095.x

25. Wang Z, Duke GM (2007) Virology Journal Cloning of the canine RNA polymerase I promoter and
establishment of reverse genetics for influenza A and B in MDCK cells. doi: 10.1186/1743-422X-4-
102

26. Moncorge O, Long JS, Cauldwell A V., et al (2013) Investigation of Influenza Virus Polymerase
Activity in Pig Cells. J Virol 87:384–394 . doi: 10.1128/JVI.01633-12

27. Wakai C, Iwama M, Mizumoto K, Nagata K (2011) Recognition of Cap Structure by Influenza B
Virus RNA Polymerase Is Less Dependent on the Methyl Residue than Recognition by Influenza A
Virus Polymerase. J Virol 85:7504 . doi: 10.1128/JVI.02375-10

28. Muramoto Y, Takada A, Fujii K, et al (2006) Hierarchy among Viral RNA (vRNA) Segments in
Their Role in vRNA Incorporation into Influenza A Virions. J Virol 80:2318–2325 . doi:
10.1128/JVI.80.5.2318-2325.2006

29. Zhang Z, Zhang H, Xu L, et al (2020) Selective usage of ANP32 proteins by influenza B virus
polymerase: Implications in determination of host range. doi: 10.1371/journal.ppat.1008989

30. Eisfeld AJ, Neumann G, Kawaoka Y (2014) Influenza A virus isolation, culture and identification.
Nat Protoc 9:2663–2681 . doi: 10.1038/NPROT.2014.180

31.   Ivshina A V., Vodeiko GM, Kuznetsov VA, et al (2004) Mapping of Genomic Segments of Influenza
B Virus Strains by an Oligonucleotide Microarray Method. J Clin Microbiol 42:5793 . doi:
10.1128/JCM.42.12.5793-5801.2004

32.   Glover DM (1985) DNA cloning: a practical approach. Volume 1

33.   Quan J, Tian J (2009) Circular Polymerase Extension Cloning of Complex Gene Libraries and
Pathways. PLoS One 4:6441 . doi: 10.1371/journal.pone.0006441

34.   Quan J, Tian J (2011) Circular polymerase extension cloning for high-throughput cloning of
complex and combinatorial DNA libraries. Nat Protoc 6:242–251 . doi: 10.1038/nprot.2010.181

35.   Santos JJS, Finch C, Sutton T, et al (2017) Development of an Alternative Modified Live Influenza
B Virus Vaccine. J Virol 91: . doi: 10.1128/JVI.00056-17/FORMAT/EPUB

36.   Tisoncik JR, Billharz R, Burmakina S, et al (2011) The NS1 protein of influenza A virus suppresses
interferon-regulated activation of antigen-presentation and immune-proteasome pathways. J Gen
Virol 92:2093 . doi: 10.1099/VIR.0.032060-0

37.   Nogales A, Aydillo T, Ávila-Pérez G, et al (2019) Functional Characterization and Direct
Comparison of Influenza A, B, C, and D NS1 Proteins in vitro and in vivo. Front Microbiol 10:2862 .
doi: 10.3389/FMICB.2019.02862/BIBTEX

38.   Mahmoudian S, Auerochs S, Gröne M, Marschall M (2009) Influenza A virus proteins PB1 and
NS1 are subject to functionally important phosphorylation by protein kinase C. J Gen Virol
90:1392–1397 . doi: 10.1099/VIR.0.009050-0

39.   Huang X, Zheng M, Wang P, et al (2017) ARTICLE An NS-segment exonic splicing enhancer
regulates influenza A virus replication in mammalian cells. doi: 10.1038/ncomms14751

40.   Salvatore M, Basler CF, Parisien J-P, et al (2002) Effects of Influenza A Virus NS1 Protein on
Protein Expression: the NS1 Protein Enhances Translation and Is Not Required for Shutoff of Host
Protein Synthesis. J Virol 76:1206–1212 . doi: 10.1128/JVI.76.3.1206-1212.2002

41.   Chen G, Liu CH, Zhou L, Krug RM (2014) Cellular DDX21 RNA helicase inhibits influenza A virus
replication but is counteracted by the viral NS1 protein. Cell Host Microbe 15:484–493 . doi:
10.1016/j.chom.2014.03.002

42.   Mok BW-Y, Song W, Wang P, et al (2012) The NS1 Protein of Influenza A Virus Interacts with
Cellular Processing Bodies and Stress Granules through RNA-Associated Protein 55 (RAP55)
during Virus Infection. J Virol 86:12695 . doi: 10.1128/JVI.00647-12

43.   Dauber B, Heins G, Wolff T (2004) The Influenza B Virus Nonstructural NS1 Protein Is Essential
for Efficient Viral Growth and Antagonizes Beta Interferon Induction. J Virol 78:1865 . doi:
10.1128/JVI.78.4.1865-1872.2004

44.   Yuan W, Krug RM (2001) Influenza B virus NS1 protein inhibits conjugation of the interferon (IFN)-
induced ubiquitin-like ISG15 protein. EMBO J 20:362 . doi: 10.1093/EMBOJ/20.3.362

45.   Mondal A, Dawson AR, Potts GK, et al (2017) Influenza virus recruits host protein kinase C to
control assembly and activity of its replicat ion machinery. Elife 6: . doi: 10.7554/eLife.26910

46.   Graci JD, Cameron CE (2006) Mechanisms of action of ribavirin against distinct viruses. Rev Med
Virol 16:37–48 . doi: 10.1002/RMV.483

47.   Furuta Y, Komeno T, Nakamura T (2017) Favipiravir (T-705), a broad spectrum inhibitor of viral
666   RNA polymerase. Proc Jpn Acad Ser B Phys Biol Sci 93:449–463 . doi: 10.2183/PJAB.93.027

667   48.   Gish RG (2006) Treating HCV with ribavirin analogues and ribavirin-like molecules. J Antimicrob
668   Chemother 57:8–13 . doi: 10.1093/JAC/DKI405

669   49.   Oxford JS (1975) Inhibition of the replication of influenza A and B viruses by a nucleoside
670   analogue (ribavirin). J Gen Virol 28:409–414 . doi: 10.1099/0022-1317-28-3-409

671   50.   Cheung PPH, Watson SJ, Choy KT, et al (2014) Generation and characterization of influenza A
672   viruses with altered polymerase fidelity. Nat Commun 2014 51 5:1–13 . doi: 10.1038/ncomms5794

673   51.   McCracken EA, McCracken EA (1976) Double-Blind Clinical Assessment of Ribavirin (Virazole) in
674   the Prevention of Induced Infection with Type B Influenza Virus. J Infect Dis 133:A109–A113 . doi:
675   10.1093/INFDIS/133.SUPPLEMENT_2.A109

676   52.   Ren L, Peng Z, Chen X, Ouyang H (2016) Live Cell Reporter Systems for Positive-Sense Single
677   Strand RNA Viruses. Appl Biochem Biotechnol 178:1567 . doi: 10.1007/S12010-015-1968-5

678   53.   Su J, Dou Y, You Y, Cai X (2015) Application of minigenome technology in virology research of
679   the Paramyxoviridae family. J Microbiol Immunol Infect 48:123–129 . doi:
680   10.1016/J.JMII.2014.02.008