Loss of self-tolerance leads to altered gene expression and IMD pathway activation in

Drosophila melanogaster

Authors: Pooja Kr[1], Ashley L. Waring-Sparks[1], Nicholas M. Bretz[1,2], and Nathan T. Mortimer[1,2]

Affiliation: [1]School of Biological Sciences, Illinois State University, Normal, IL, 61790, USA; [2] Department of Biochemistry and Biophysics, Oregon State University, Corvallis, OR, 97331, USA

*Correspondence: Nathan T. Mortimer, Department of Biochemistry and Biophysics, Oregon State University, Corvallis, OR, 97331, USA, nathan.mortimer@oregonstate.edu

ABSTRACT

Immune self-tolerance is the ability of a host's immune system to recognize and avoid triggering immune responses against self-tissue. This allows the host to avoid self-directed immune damage while still responding appropriately to pathogen infection. A breakdown of self-tolerance can lead to an autoimmune state in which immune cells target healthy self-tissue, leading to inflammation and tissue damage. In order to better understand the basic biology of autoimmunity and the role of the innate immune system in maintaining self-tolerance, we have recently characterized the *Drosophila melanogaster tuSz* autoimmune mutant. This mutant strain can serve as a model of innate immune mediated self-tolerance, and here we identify transcripts that are deregulated in flies experiencing a loss of self-tolerance. We find that these changes include the activation of the Relish/NFκB transcription factor, alterations in transcripts encoding proteins predicted to mediate organismal metabolism, and a downregulation of transcripts linked to developmental processes. We further find that NFκB signaling plays a protective role against loss of self-tolerance in *Drosophila.* Our findings provide insight into the transcriptional and physiological changes underlying self-tolerance and autoimmunity.

1. INTRODUCTION

The maintenance of self-tolerance, the process in which self-tissues are identified and self-directed immune mechanisms are repressed, is an important characteristic of the immune system (Medzhitov and Janeway, 2002; Romagnani, 2006). The breakdown of the immune system's ability to discriminate healthy self-tissue from non-self pathogens can lead to damage of a host's own tissues by the immune system. Multiple studies have revealed the importance of self-tolerance, and the failure of this system has been implicated in a variety of autoimmune disorders and other immune mediated diseases. Although much of the research uncovering mechanisms of immune recognition and self-tolerance has focused on adaptive immune

mechanisms, a growing body of work has demonstrated the importance of innate immune

mechanisms in the suppression of autoimmunity (Toubi and Vadasz, 2019; Waldner, 2009).

The genetic model organism *Drosophila melanogaster* is a valuable model for

understanding conserved innate immune mechanisms (Lemaitre and Hoffmann, 2007). Flies are

host to a wide range of pathogens including bacteria, fungi, viruses, and parasites and use both

humoral and cell mediated immune responses to combat these pathogens (Brennan and

Anderson, 2004). In *Drosophila*, hemocytes (immune cells) known as plasmatocytes are found

in circulation, and following stimulation, surveil the hemolymph (body cavity) for the presence of

invading pathogens and tissue damage. These macrophage-like plasmatocytes are responsible

for the phagocytosis of microbial pathogens and are involved in the encapsulation response

against macroparasites including parasitoid wasp eggs (Honti et al., 2014; Mortimer et al., 2012;

Parsons and Foley, 2016). Another immune cell type, the lamellocytes, are generally not

present in a healthy fly larva, but are induced following parasitoid wasp infection and are

required for melanotic encapsulation of parasitoid eggs and other foreign objects (Anderl et al.,

2016; Honti et al., 2014; Mortimer et al., 2012).

Drosophila hemocytes also play a role in self-recognition and maintaining self-tolerance.

We have recently characterized the *Drosophila tuSz^1* temperature sensitive mutant strain which

mounts a self-directed encapsulation response at the restrictive temperature (Mortimer et al.,

2021). We found that the loss of self-tolerance in *tuSz^1* is due to two genetic changes. The first

is a gain of function mutation in *hopscotch* (*hop*), the *Drosophila* homolog of Janus Kinase

(JAK). The JAK-STAT signaling pathway plays a key role in immune priming, and the *hop^{Sz}*

allele leads to hemocyte activation and the ectopic production of lamellocytes in naïve larvae

(Mortimer et al., 2021). These hemocyte phenotypes are also seen in larvae with the *hop^{Tum}*

gain of function allele (Hanratty and Dearolf, 1993; Luo et al., 1995), but interestingly, even

though *hop^{Tum}* and *tuSz^1* mutants both have hemocyte activation and excess lamellocyte

production, hop^{Tum} mutants do not display an autoimmune phenotype (Mortimer et al., 2021).

This would suggest that JAK-STAT mediated immune activation alone is insufficient for the loss

of self-tolerance, and that loss of self-tolerance and immune priming are genetically separable.

The second mutation in the $tuSz^1$ strain is a conditional loss of function mutation in the GCS1

gene. At the restrictive temperature, the $GCS1^{Sz}$ allele leads to disruption of protein N-

glycosylation of extracellular matrix proteins (Mortimer et al., 2021). Together these findings

define a system in which Drosophila self-tolerance is maintained by the ability of primed

hemocytes to sense a self-signal that is encoded in the N-glycosylation of extracellular matrix

proteins, and the loss of this self-recognition leads to the melanotic encapsulation of self-

tissues. Thus, hemocytes can mediate both the activation of an immune response and the

maintenance of self-tolerance thereby regulating immune homeostasis.

Here, we utilize the Drosophila $tuSz^1$ and hop^{Tum} mutant strains to identify key

transcriptional changes underlying immune priming, self-tolerance, and autoimmunity.

Differentially expressed transcripts often reveal disease-induced changes (Porcu et al., 2021),

so our analysis of transcript abundance changes in these mutants may allow us to identify

processes that occur in autoimmune reactions following the loss of self-tolerance. Our results

further suggest that the pro-inflammatory Rel/NFκB signaling pathway is dysregulated in the

$tuSz^1$ loss of self-tolerance phenotype, and plays a protective role in this response.

2. MATERIALS AND METHODS

2.1. Insect Stocks

The D. melanogaster strains w^{1118} (BDSC: 5905), $tuSz^1$ (BDSC: 5834),

PBac{GFP.FPTB-Rel}VK00037 (Rel-GFP) (BDSC: 81268), hop^{Tum} (BDSC: 8492), P{VALIUM20-

GAL4.1}attP2 (UAS-GAL4-RNAi) (BDSC: 35784), P{TRiP.HM05154}attP2 (UAS-Rel-RNAi-1)

(BDSC: 28943), P{TRiP.HMS00070}attP2 (UAS-Rel-RNAi-2) (BDSC: 33661),

P{TRiP.HMS00063}attP2 (*UAS-Dredd-RNAi*) (BDSC: 34070), *P{TRiP.HMC04952}attP40* (*UAS-*

key-RNAi) (BDSC: 57759), and   *PGRP-LB^{Delta}* (BDSC: 55715) were obtained from the

Bloomington Drosophila Stock Center (BDSC). *msn-GAL4* was obtained from R. Schulz

(Tokusumi et al., 2009). All fly stocks were maintained on standard *Drosophila* medium on a 12-

hour light-dark cycle. The study also uses the parasitoid wasp *Leptopilina boulardi* (strain Lb17)

which is maintained on the *Canton S D. melanogaster* strain (BDSC: 64349).

2.2 RNA Isolation

Total RNA was extracted from late third instar w^{1118}, $tuSz^1$ and hop^{Tum} larvae raised at

28ºC. In total, three biological replicates were prepared with each biological replicate consisting

of 20 pooled larvae. RNA was extracted from whole larvae using Trizol (Thermo Fisher

Scientific) followed by QIAGEN RNeasy Micro clean-up (QIAGEN; both according to

manufacturer's instructions). RNA concentration was measured using a nanodrop and the

integrity was verified using agarose gel electrophoresis. The RNA samples were sent to

Novogene (Sacramento, CA) for RNA sequencing (150bp paired end reads, Illumina). RNAseq

data are available through the NCBI Gene Expression Omnibus, study GSE189383.

2.3 Differential Gene Expression Analysis

Sequence data were obtained in the form of FASTQ reads and analyzed using the

Galaxy web platform (https://usegalaxy.org/) (Afgan et al., 2018). Initial quality control was

performed using FastQC (version 0.11.8) (Andrews, 2010) to trim reads below a quality score

threshold of 30. Reads were then mapped to the *D. melanogaster* reference genome dm6 using

HISAT2 (version 2.1.0) (Kim et al., 2015). The resulting BAM format files were analyzed

using featureCounts (version 1.6.3) (Liao et al., 2014) to generate read counts for each gene in

the reference genome annotation. The counts generated by featureCounts were subsequently

used to determine the differentially expressed genes using DESeq2 (version 2.11.40.6) (Love et

al., 2014). The results were filtered and significant differentially expressed genes (DEGs) were

identified after applying significance cut-offs of a false discovery rate (FDR) < 0.05 and $\log_2$ fold

change ≥ 1 (for upregulated transcripts) or ≤ -1 (for downregulated transcripts).

2.4 Functional Annotation

The Gene Ontology (GO) Resource (http://geneontology.org/) and PANTHER

classification system (Ashburner et al., 2000; Gene Ontology Consortium, 2021; Mi et al., 2013)

were used for GO term analysis of each list of DEGs. The FDR threshold to determine term

enrichment was 0.05.

2.5 Rel Localization

To determine Rel subcellular localization in hemocytes in control and mutant larvae,

w^{1118}, $tuSz^1$, and hop^{Tum} unmated females were crossed to *Rel-GFP* males and the crosses

were raised at 28°C. From these crosses we selected w^{1118} /Y; Rel-GFP/+ control larvae and

hop^{Tum}/Y; Rel-GFP/+ and *$tuSz^1$/Y; Rel-GFP/+* mutant larvae. Five male third instar larvae from

each genotype were bled in PBS and hemocytes were stained using DAPI (1:500; Invitrogen) to

mark nuclei and mounted in Vectashield mounting medium. To determine Rel localization in

hemocytes of parasitoid infected larvae, late second instar w^{1118} /Y; Rel-GFP/+ larvae were

placed on 35mm Petri dishes filled with Drosophila medium either alone (naïve control) or

together with *L. boulardi* wasps (infected) at 25°C. Larvae were dissected at 72-hours post

infection (hpi; and age-matched controls) in PBS and hemocytes were stained using DAPI

mounted in Vectashield for imaging. Cells were imaged using a Leica SP8 confocal microscope

in the ISU Confocal Microscopy Facility to assay Rel-GFP localization.

The relative proportion of Rel-GFP signal coming from the nucleus vs cytoplasm (N/C

ratio) was calculated using Fiji (Schindelin et al., 2012). Image analysis was performed as

described in (Kelley and Paschal, 2019; Noursadeghi et al., 2008). Specifically, the DAPI stain

was used to mask the nucleus allowing for mean cytoplasmic and nuclear fluorescence intensity

values to be independently determined. The N/C ratio was then calculated based on the ratio of

mean nuclear intensity to mean cytoplasmic intensity. Cells with an N/C ratio greater than 1.5

were considered to have nuclear Rel-GFP localization and cells with an N/C ratio less than 1.5

were considered to have diffuse Rel-GFP (Rel-GFP exclusion from the nucleus was not

observed). This relative localization measure is a more reliable measure than absolute

fluorescence intensity, being robust to stochastic cell-to-cell differences in Rel expression

(Noursadeghi et al., 2008). The proportion of cells with nuclear Rel-GFP localization in the

mutant or infected backgrounds was compared to w^{1118} using Fisher's exact test.

2.6 Statistical Analysis

All statistical analyses were done in the R statistical computing environment (R Core

Team, 2021). The plotMDS function from the limma R package (Ritchie et al., 2015) was used

to plot the first two principal components to illustrate the relationships in the expression profiles

of the studied genotypes. For RNAi experiments, phenotypic penetrance data were analyzed

using Fisher's exact test, and p values were adjusted using Holm's method to correct for

multiple comparisons. Genetic interactions between *tuSz* and *PGRP-LB* were tested using

ANOVA of aligned rank transformed data using the ARTool R package (Wobbrock et al., 2011)

and pairwise comparisons were made using Tukey's HSD from the multcomp R package

(Hothorn et al., 2008). Graphs were produced using the ggplot2 and ggvenn R packages

(Wickham, 2009; Yan, 2021) and heatmaps were produced using the gplots R package

(Warnes et al., 2022).

3. RESULTS AND DISCUSSION

3.1 Identification of differentially expressed genes in *tuSz^1* and *hop^{Tum}*

To gain a better understanding of the changes involved in JAK-STAT mediated immune

priming and the consequences of the loss of self-tolerance, we performed RNA sequencing on

w^{1118}, $tuSz^1$ and hop^{Tum} larvae raised at 28ºC to assess differential transcript abundance. The

resulting sequence reads were mapped to the *D. melanogaster* genome (Table S1) to quantify

transcript levels. We find that each genotype shows a distinct expression profile (Figure 1A),

and that the genotypes are separable by plotting the first two principal components of all

transcript quantification data (Figure 1B). To identify genes of interest that may be differentially

expressed, we separately compared transcript levels in each mutant genotype to the w^{1118}

control genotype (Tables S2-S5). Both $tuSz^1$ and hop^{Tum} have gain of function mutations in *hop*,

so any genes that are differentially expressed in both genotypes are likely to be regulated by the

JAK-STAT pathway, or to be altered because of JAK-STAT pathway-mediated changes in cell

physiology. We found 832 shared genes that were differentially expressed in both $tuSz^1$ and

hop^{Tum} mutants compared to wildtype, with 384 genes that were upregulated, and 438 genes

that were downregulated (Figure 1C-D). Because the JAK-STAT pathway plays a key role in

immune activation (Agaisse and Perrimon, 2004), these identified genes might be relevant to

the priming of the immune response.

To begin characterizing these putative immune activation genes, we conducted Gene

Ontology (GO) analysis to identify the biological processes and molecular functions that are

enriched among the common set of differentially expressed genes (Table 1). Analysis of genes

that were upregulated in both $tuSz^1$ and hop^{Tum} identified GO terms that have been linked to

host immunity including defense response to other organism (GO:0098542) and response to

external stimulus (GO:0009605). General immune response genes identified in these categories

(Figure 2A) include five members of the antimicrobial peptide (AMP)-like *Bomanin* gene family

(*BomBc1*, *BomS1*, *BomS2*, *BomS4*, *BomT1*) (Clemmons et al., 2015) and the Cecropin family

AMP gene *CecC* (Tryselius et al., 1992), along with several immune receptors (*NimB2*, *NimC1*,

NimC4) (Kurucz et al., 2007; Somogyi et al., 2010), the prophenoloxidase family members

PPO1 and *PPO3* (Dudzic et al., 2015; Nam et al., 2008), and the *induced by infection* (*IBIN*)

gene (Valanne et al., 2019). We also found differential expression of important immune related

signaling pathways such as the JAK-STAT signaling pathway components *dome*, *et* and *upd3*

(Myllymäki and Rämet, 2014), the Toll pathway genes *nec* and *SPE* (Kambris et al., 2006), and

Rac2 and *Rap1* which are linked to hemocyte function (Huelsmann et al., 2006; Williams et al.,

2005). The *tuSz^1* and *hop^{Tum}* mutations both lead to the ectopic production of lamellocytes, and

accordingly, we see lamellocyte related genes such as *ItgaPS4*, *ItgaPS5*, and *drip* upregulated

in these genotypes (Stofanko et al., 2010; Tattikota et al., 2020). The proteolysis (GO:0006508)

GO term is also enriched among genes upregulated in both *tuSz^1* and *hop^{Tum}*. Notably, 29

members of the S1A serine protease family are significantly upregulated in both *tuSz^1* and

hop^{Tum}, including *CG10764* and *CG4793* which have recently been shown to be involved in the

cellular immune response to parasitoid infection (Kr et al., 2021).

The induction of host immunity is associated with life history tradeoffs, primarily in the

redistribution of energy resources from development or reproduction to the production of an

immune response (Zuk and Stoehr, 2002). We found that genes involved with cellular lipid

metabolism (GO:0044255) are enriched among the common transcripts upregulated in *tuSz^1*

and *hop^{Tum}*, and conversely that genes linked to development, including regulation of

multicellular organismal development (GO:2000026), tissue morphogenesis (GO:0048729), and

cell fate commitment (GO:0045165) are among those transcripts downregulated in both *tuSz^1*

and *hop^{Tum}* (Table 1). The upregulation of transcripts involved in metabolism and

downregulation of transcripts linked to developmental processes possibly reflects the immune

response-development tradeoff, and suggests that this tradeoff may be regulated in part by

JAK-STAT signaling or downstream processes. Further study of these genes may help to

provide an understanding of the energy trade-offs and other physiological changes that occur

during immune responses.

3.2 Loss of self-tolerance in *tuSz^1* mutants results in differential gene expression

Despite the similarity in ectopic JAK-STAT pathway activation between the *tuSz^1* and

hop^{Tum} mutants, *tuSz^1* specifically displays a loss of self-tolerance phenotype (Mortimer et al.,

2021). We investigated transcripts that are specifically differentially expressed in *tuSz^1* to

identify genes that might be involved in innate immune self-tolerance and autoimmune

mechanisms. Out of 1174 transcripts that were differentially expressed in *tuSz^1* but not *hop^{Tum}*

(referred to as *tuSz^1* specific DEGs), we find 683 transcripts that are significantly upregulated

and 491 transcripts that are significantly downregulated (Figure 1C-D). It is likely that this class

of *tuSz^1* specific DEGs represents genes that are altered due to the physiological changes seen

in larvae during the autoimmune response to loss of self-tolerance rather than direct targets of

JAK-STAT signaling.

In contrast to the list of upregulated transcripts in common to *tuSz^1* and *hop^{Tum}*, GO Term

analysis of these *tuSz^1* specific DEGs (Table 2) failed to find enrichment of any GO categories

associated with positive regulation of host immunity. Indeed, one of the most highly enriched

terms was the negative regulation of antimicrobial peptide biosynthetic process (GO:0002806).

This term was linked to peptidoglycan recognition protein (PGRP) encoding genes, and we find

that six PGRP gene transcripts are altered in *tuSz^1*, but not *hop^{Tum}*, mutants (Figure 2A). There

are 13 members of this family in the *Drosophila melanogaster* genome (Table 3), which

primarily play roles in the regulation of the Toll and Imd immune signaling pathways (Kurata,

2014). There are five PGRP genes that encode negative regulators of Imd signaling, and we

find that all five of these are specifically upregulated in *tuSz^1* mutants, with the sixth altered

PGRP playing an unknown role (Table 3). The finding that whereas positive regulators of

immunity are identified as upregulated genes in both *tuSz^1* and *hop^{Tum}*, the enrichment of genes

involved in negative regulation of the Imd pathway specifically in *tuSz^1* mutants suggests a

differential role of immune mechanisms in JAK-STAT mediated immune priming and in the

autoimmune response to a loss of self-tolerance. More specifically, the regulatory PGRP genes

we find elevated specifically in *tuSz^1* larvae are induced downstream of Imd signaling in a

negative feedback regulatory loop (Zaidman-Rémy et al., 2006). This finding suggests that Imd

pathway activity may be altered in the autoimmune environment resulting from the loss of self-

tolerance in *tuSz^1* larvae.

We additionally find 26 genes that map to the chitin-based cuticle development (GO:

0040003) term among the *tuSz^1* specific DEGs, including members of the *Lcp*, *Acp*, *Twdl*, *Cpr*,

and *Ccp* gene families (Table 2). Expression of cuticle genes has been linked to infection in *D.*

melanogaster and other insects, although a role in host immunity has yet to be identified (Hou et

al., 2021; Pantha et al., 2021; Yadav et al., 2017). Many of the additional enriched GO

categories among the *tuSz^1* specific DEGs are associated with stress responses and

metabolism, perhaps providing insight into physiological changes in *tuSz^1* mutant larvae due to

the occurrence of an autoimmune response (Table 2). We find an enrichment of oxidoreductase

activity (GO: 0016491) genes, including the glutathione transferase (GST) encoding genes

GstD2, GstD6, GstD7, GstD8, GstE1, GstE3, GstE5, and *GstE7* (Low et al., 2007). We also find

that 18 cytochrome P450 (CYP450) encoding genes are upregulated in *tuSz^1* mutants.

CYP450s are involved in a wide range of cellular activities including roles in hormone signaling

and detoxification of xenobiotics (Bergé et al., 1998; Chung et al., 2009). The specific

upregulation of these stress response genes suggests that loss of self-tolerance may lead to the

production of an internal oxidizing environment.

As mentioned previously, metabolic changes often accompany immune responses, and

we find that two general regulators of host metabolism, the *Adipokinetic hormone receptor*

(*AkhR*) (Bharucha et al., 2008) and *Linking immunity and metabolism* (*Lime*) (Figure 2)

(Mihajlovic et al., 2019), are upregulated in *tuSz[1]* mutants. *Lime* encodes a transcription factor

that plays a key role in metabolic changes following infection. Accordingly, we find that genes

associated with a wide range of metabolic activities including generation of precursor

metabolites and energy (GO:0006091), carbohydrate derivative metabolic process

(GO:1901135), alpha-amino acid metabolic process (GO:1901605), and lipid metabolic process

(GO:0006629) were enriched among those upregulated in *tuSz[1]* larvae (Table 2). Further

analysis of these genes and the mechanisms they mediate will help to identify the metabolic

pathways involved in the response to a loss of self-tolerance.

Along with upregulated transcripts, transcripts that are downregulated in these

genotypes may also provide insight into the cellular activities that are altered. We find that

genes associated with tissue morphogenesis (GO:0048729) and the establishment of tissue

polarity (GO:0007164) are enriched among the downregulated *tuSz[1]* specific DEGs (Table 2).

The loss of cellular polarity and disrupted tissue organization has also been observed in both

autoimmune mouse models and patients with autoimmune conditions such as inflammatory

bowel disease (Ahmad et al., 2017; Guo and Shen, 2021; Klunder et al., 2017). This may

suggest that tissue structural maintenance is a key self-tolerance checkpoint, in agreement with

previous findings in *Drosophila* (Kim and Choe, 2014; Rizki and Rizki, 1974).

3.3 Relish/NFκB is activated in immune tissues in *tuSz[1]* mutants

Based on the specific upregulation of Imd-responsive PGRP genes (Table 3), we

hypothesized that Imd signaling may be dysregulated in the *tuSz[1]* loss of self-tolerance

phenotype. While the role of the Imd pathway in mediating the *Drosophila* humoral immune

response to Gram-negative bacteria has been well established (Lemaitre et al., 1995), the

mechanism by which it may participate in the response to loss of self-tolerance is unknown. The

tuSz^1 autoimmune mutant may therefore provide an effective model to begin exploring the role

of the Imd pathway in self-tolerance and autoimmune reactions.

The Imd pathway mainly responds to infection via differential expression of target genes

(Kleino and Silverman, 2014). A previous study has identified genes whose expression is

modified by constitutive Imd pathway activation (Davoodi et al., 2019). We find that 58 of the

324 genes identified in this study also showed differential expression in *tuSz^1* larvae (Figure 2B)

(Table S6), including three of the identified PGRP genes (*PGRP-LB*, *PGRP-LF*, *PGRP-SC2*),

and other immune genes such as *AttA, Rala, Gbp1* and *GILT2*. The altered expression of Imd

pathway genes is not seen in *hop^{Tum}* mutants (Figure 2B), and the *tuSz^1* expression profile is

clearly distinct from both *w^{1118}* and *hop^{Tum}* by principal components analysis of Imd pathway

transcript levels (Figure 2C). Notably, *w^{1118}* and *hop^{Tum}* samples are indistinguishable in this

analysis, suggesting minimal alteration of Imd signaling in *hop^{Tum}*. These findings provide

support for the hypothesis that Imd signaling is dysregulated in *tuSz^1* mutants in response to the

loss of self-tolerance rather than as a direct result of JAK-STAT pathway activation. The overlap

of gene expression changes between *tuSz^1* mutants and constitutive Imd activation suggests

that the Imd pathway is activated in *tuSz^1* mutant larvae.

Gene expression downstream of Imd is largely mediated by the activation of Relish

(Rel), a *D. melanogaster* member of the NF-κB family of transcriptional activators (Hetru and

Hoffmann, 2009). Rel activation is regulated by signal-dependent translocation into the nucleus.

Whereas full-length Rel protein is autoinhibited and remains in the cytoplasm, Imd activation

results in cleavage of Rel and the production of an active N-terminal fragment, which

translocates to the nucleus to drive gene expression (Stöven et al., 2000). To determine

whether Rel is activated in *tuSz^1* mutants, we used a *Rel-GFP* fusion protein encoding

transgenic line to track the subcellular localization within the main immune tissues, the

hemocytes and fat body. This reporter has GFP fused to the N-terminal domain of Relish, and

GFP localization in the nucleus versus cytoplasm gives a relative readout of pathway activity.

Rel-GFP expression was observed in plasmatocytes in *tuSz^1/Y; Rel-GFP/+* and *w^{1118}/Y;*

Rel-GFP/+ larvae (Figure 3A-B). We found that *Rel-GFP* was significantly more likely to be

concentrated in the nucleus in plasmatocytes from *tuSz^1/Y; Rel-GFP/+* larvae compared to *w^{1118}*

/Y; Rel-GFP/+ control larvae (Fisher's exact test: p = 1.2x10^{-5}; odds ratio = 9.02) or compared to

hop^{Tum} mutant larvae (Fisher's exact test: p = 9.7x10^{-4}; odds ratio = 5.87) (Figure 3D),

suggesting that *Rel* activation is induced specifically in *tuSz^1* mutants. In addition, *Rel-GFP*

expression and nuclear localization were observed in lamellocytes from *tuSz^1/Y; Rel-GFP/+*

larvae (Figure 3C), suggesting that the Imd pathway is also active in this hemocyte type in the

tuSz^1 mutant background.

To determine if this Rel activation is common to cellular immune responses in

Drosophila or if it may be specific to loss of self-tolerance reactions, we assayed *Rel-GFP*

localization in the immune cells of *w^{1118}/Y; Rel-GFP/+* larvae during the response to parasitoid

wasp infection. *w^{1118}/Y; Rel-GFP/+* larvae were infected by *Leptopilina boulardi* and hemocytes

were isolated and imaged 72hpi. We observed minimal *Rel-GFP* expression in both

plasmatocytes (Figure 4A-B) and lamellocytes (Figure 4C) in these infected larvae. Circulating

non-hemocyte cells (marked by an asterisk* in Figure 4B-C) showed strong fluorescence and

acted as an imaging control. These results suggest that Rel and the Imd pathway are not likely

to be involved in the cellular immune response to parasitoid infection. In agreement with this

finding, a previous study has shown that *Rel* mutants have a normal encapsulation response

against parasitoid wasp infection (Hedengren et al., 1999).

To test whether loss of *Rel* modifies the *tuSz^1* loss of self-tolerance phenotype, we used

RNA interference (RNAi) to knock-down *Rel* transcripts in immune cells in the *tuSz^1* mutant

background. In comparison with a genetic control (*GAL4-RNAi*), the expression of either of two

distinct *Rel* knockdown constructs (*Rel-RNAi-1* and *Rel-RNAi-2*) in immune cells leads to an

increased penetrance of the autoimmune phenotype (*Rel-RNAi-1*: Fisher's exact test: p =

4.3×10^{-3}; *Rel-RNAi-2*: Fisher's exact test: p = 3.7×10^{-5}) (Figure 5A). To further test the role of *Rel*

activation, we used knockdown constructs targeting two distinct positive regulators of *Rel*: *key,* a

component of the IκB Kinase complex which is required for *Rel* activation (Ertürk-Hasdemir et

al., 2009) (*key-RNAi*) and *Dredd,* the caspase that directly cleaves and activates *Rel* (Stöven et

al., 2003) (*Dredd-RNAi*). Similar to *Rel-RNAi-1* and *-2, Dredd-RNAi* mediated knockdown

enhances the penetrance of the $tuSz^1$ phenotype (Fisher's exact test: p = 1.0×10^{-6}) (Figure 5A).

Additionally, *key-RNAi* mediated knockdown shows considerable lethality in the $tuSz^1$

background, which we have previously observed with mutations that strongly enhance the $tuSz^1$

phenotype (Mortimer et al., 2021). These results suggest that *Rel*/NFκB transcriptional activity is

protective against loss of self-tolerance.

Rel/NFκB transcriptional activity is associated with pro-inflammatory signaling in

Drosophila and other species (Ganesan et al., 2011; Lawrence, 2009; Libert et al., 2006;

Myllymäki et al., 2014). The observed Rel activation suggests that inflammation is a feature of

loss of self-tolerance responses in *Drosophila*, a mechanism that is conserved in human

autoimmune disease (Furman et al., 2019; Zhang et al., 2017). The finding that loss of Rel

activation enhances the $tuSz^1$ phenotype is surprising given that the constitutive activation of

NF-κB is implicated in the pathogenesis of multiple forms of autoimmune disease (Ilchovska and

Barrow, 2021; Miraghazadeh and Cook, 2018; Sun, 2017; Zhang et al., 2017). However, loss of

mutations in NF-κB family genes including the *Rel* homolog RELA have also been linked to the

development of autoimmune disorders (Barnabei et al., 2021; Comrie et al., 2018). This

suggests that, in some situations, NF-κB signaling can be protective against disruptions in self-

tolerance and progression of autoimmune disorders. Our findings of the protective effect of *Rel*

in *Drosophila* may provide a model to further explore this underappreciated role of NFκB

signaling.

To better understand the role of Rel activation in the response to loss of self-tolerance in

Drosophila, we tested the effect of a loss of function mutation in the negative regulatory PRGP

family protein PGRP-LB. We find that in the *tuSz^1* genetic background, a heterozygous deletion

mutation of the *PGRP-LB* gene dominantly and significantly suppresses the *tuSz^1* mutant

phenotype (Full genotype: *tuSz^1/Y*; *PGRP-LB^{Delta}/+* . Tukey's HSD: t = -5.196; p = 3.7x10^{-3})

(Figure 5B). Interestingly, the loss of one copy of the *PGRP-LB* gene in an otherwise wildtype

background does not lead to a visible phenotype (Full genotype: *w^{1118}/Y*; *PGRP- LB^{Delta}/+*.

Tukey's HSD: t = 0.0; p = 1.0) (Figure 5B).

The negative regulatory PGRPs are amidase or amidase-like proteins that suppress the

Imd/*Rel* pathway by either degrading (amidase PGRPs) or scavenging (amidase-like PGRPs)

peptidoglycans (PGNs), the bacterial cell well components that trigger pathway activation

(Bischoff et al., 2006; Maillet et al., 2008; Zaidman-Rémy et al., 2006). PGRP-LB is an active

amidase that is specific for *meso*-diaminopimelic acid-containing peptidoglycans (DAP-PGN),

which are found predominantly in gram negative bacteria (Zaidman-Rémy et al., 2006). Based

on this mode of action, we would not predict that loss of PGRP-LB would have an impact in the

sterile inflammation environment of *tuSz^1* mutants. However, structural analysis of PGRPs

suggests a mechanism through which these proteins may have target-binding specificity beyond

the PGN sub-type (Kim et al., 2003; Royet et al., 2011). Multiple examples of PGN-independent

activity have also been observed in diverse species (Lu et al., 2006; Sharma et al., 2011; Tydell

et al., 2006). The loss of self phenotype in *tuSz^1* mutants arises due to defects in the N-

glycosylation of extracellular matrix proteins (Mortimer et al., 2021). Perhaps these mis-

glycosylated proteins serve as an altered-self signal (Maverakis et al., 2015), which is

recognized by PGRPs as an additional, and thus far unidentified, target. If PGRP-LB recognizes

mis-glycolysated proteins to trigger an immune response, it could explain the strong

suppression of the tuSz^1 mutants by heterozygosity for the *PGRP-LB* gene. Conversely, PGRPs

have also been linked to modulating host immunity and tolerance to the gut microbiota (Bosco-

Drayon et al., 2012). It may be that the loss of *PGRP-LB* leads to alterations in the gut

microbiome, which may have direct or indirect effects on the loss of self-tolerance immune

response that characterizes *tuSz^1* mutants.

4. CONCLUSION

Immune self-tolerance is essential for maintaining immune homeostasis, and breakdown

of self-tolerance mechanisms results in autoimmunity and other immune related diseases.

Although great progress has been made in understanding self-tolerance in the context of

adaptive immunity, innate immune mechanisms regulating self-tolerance and its role in

autoimmunity remain less well characterized. In this study we have identified genes that are

differentially expressed following loss of self-tolerance in the *Drosophila melanogaster tuSz^1*

mutant. These results provide insight into the biological processes that take place in *tuSz^1*

mutant larvae and may contribute toward a better understanding of the physiological changes in

loss of self-tolerance autoimmune reactions.

Our findings also serve to highlight the complex relationship between the regulation of

immune signaling mechanisms and the development of autoimmune phenotypes. We find that,

similar to human autoimmune disease, the loss of self-tolerance in *Drosophila* leads to elevated

Rel/NFκB signaling. Dysregulation of NFκB through both ectopic activation or loss of function is

associated with the pathogenesis of autoimmunity, and we find that the pathway plays a

protective role in *Drosophila*. This will provide a tractable genetic model for further studies into

the complex roles of NFκB in autoimmune disorders. Additionally, we have identified a role for a

peptidoglycan recognition protein in regulating the response to a loss of self-tolerance in the

absence of bacterial peptidoglycans. These PGRPs are widely conserved from insects to
vertebrates, but their possible roles outside of bacterial pattern recognition have yet to be
explored. We have identified the intriguing possibility that certain PGRP proteins may recognize
self or lack-of-self epitopes and contribute to loss of self-tolerance reactions. Again, the loss of
self-tolerance *Drosophila* mutant lines may provide a model to better understand these
functions.

ACKNOWLEDGEMENTS

405 We thank Joshua Hill and Elise N. Le for their assistance on this project. The ISU Confocal
406 Microscopy Facility was funded by NSF grant DBI-1828136. Stocks obtained from the
407 Bloomington Drosophila Stock Center (NIH P40OD018537) were used in this study.

FUNDING

409 Research reported in this publication was supported by the National Institute of General Medical
410 Sciences of the National Institutes of Health under Award Number R35GM133760 to NTM, a
411 Grant-in-Aid of Research from the National Academy of Sciences administered by Sigma Xi,
412 The Scientific Research Society to PK, and the Illinois State University College of Arts and
413 Sciences University Research Grants program.

TABLES

415 **Table 1.** Enriched Gene Ontology (GO) Terms among DEG in common to both *tuSz[1]* and *hop[Tum]*

GO Term	Fold enrichment	FDR
Upregulated genes		
Defense response to other organism (GO:0098542)	2.58	0.014
Response to external stimulus (GO:0009605)	2.11	2.6×10^{-4}

Proteolysis (GO:0006508)	2.87	1.2×10^{-9}
Cellular lipid metabolic process (GO:0044255)	2.43	0.022
Downregulated genes		
Developmental process (GO:0032502)	1.59	4.7×10^{-4}
Regulation of multicellular organismal development (GO:2000026)	2.24	0.015
Tissue morphogenesis (GO:0048729)	2.15	0.029
Cell fate commitment (GO:0045165)	2.99	0.005

417 **Table 2.** Enriched GO Terms among *tuSz^1* specific DEGs

GO Term	Fold enrichment	FDR
Upregulated genes		
Negative regulation of antimicrobial peptide biosynthetic process (GO:0002806)	12.42	0.031
Chitin-based cuticle development (GO: 0040003)	2.89	4.9×10^{-6}
Oxidoreductase activity (GO: 0016491)	2.80	5.4×10^{-13}
Generation of precursor metabolites and energy (GO:0006091)	2.32	0.023
Carbohydrate derivative metabolic process (GO:1901135)	2.14	5.0×10^{-5}
Alpha-amino acid metabolic process (GO:1901605)	3.29	0.006
Lipid metabolic process (GO:0006629)	1.98	0.003
Downregulated genes		
Tissue morphogenesis (GO:0048729)	2.43	2.3×10^{-4}

Establishment of tissue polarity (GO:0007164)	4.17	0.049
Appendage development (GO:0048736)	2.63	0.004
Central nervous system development (GO:0007417)	2.53	0.042
Cell differentiation (GO:0030154)	1.53	0.043
Pattern specification process (GO:0007389)	2.09	0.035
Regulation of cell communication (GO:0010646)	1.83	0.025
Positive regulation of stem cell proliferation (GO:2000648)	9.6	0.003
Metamorphosis (GO:0007552)	2.42	0.002

419 **Table 3.** Expression of PGRP family genes in $tuSz^1$ and hop^{Tum} compared to w^{1118} control
420 larvae.

Gene	Pathway	Function	$tuSz^1$ log$_2$FC	hop^{Tum} log$_2$FC
PGRP-LA	IMD	Positive Regulator	-0.38	-0.30
PGRP-LB	IMD	Negative Regulator	**1.83***	-0.18
PGRP-LC	IMD	Positive Regulator	-1.12	-0.63
PGRP-LD	Unknown	Unknown	**-2.77***	-0.81
PGRP-LE	IMD	Positive Regulator	-.018	-0.75
PGRP-LF	IMD	Negative Regulator	**1.37***	0.32
PGRP-SA	Toll	Positive Regulator	1.67	0.99
PGRP-SB1	IMD	Immune Effector	-0.92	-1.95
PGRP-SB2	IMD	Immune Effector	Not detected	Not detected
PGRP-SC1a	IMD	Negative Regulator	**3.46***	1.70
PGRP-SC1b	IMD	Negative Regulator	**3.38***	1.51
PGRP-SC2	IMD	Negative Regulator	**1.23***	-0.19
PGRP-SD	Toll	Positive Regulator	0.66	-0.83

* indicates FDR < 0.05 compared to w^{1118} control; $\log_2 FC = \log_2$ Fold Change

FIGURE LEGENDS

Figure 1. Gene expression profiles from $tuSz^1$, hop^{Tum} and w^{1118}. (A) Heatmap of all genes

showing altered expression between w^{1118} and at least one mutant genotype. See Table S1 for

a description of the samples. (B) Multidimensional scaling plot of the first two principal

components of gene expression across the samples. (C-D) Venn diagrams illustrating the

numbers of transcripts that are (C) upregulated and (D) downregulated in the indicated

genotypes in comparison with w^{1118} control larvae.

Figure 2. Expression of immune genes in in $tuSz^1$, hop^{Tum} and w^{1118}. (A) Heatmap showing

expression of representative immune genes identified by Gene Ontology analysis in $tuSz^1$,

hop^{Tum} and w^{1118}. See Table S1 for a description of the samples. (B) Heatmap showing

expression of Imd pathway genes identified by Davoodi *et al.* (2019) in the indicated genotypes.

See Table S1 for a description of the samples. (C) Multidimensional scaling plot of the first two

principal components of differentially expressed Imd pathway genes between the samples.

Figure 3. Localization of the Rel-GFP fusion protein in hemocytes. Hemocytes were dissected

from w^{1118}/Y; *Rel-GFP/+* and $tuSz^1/Y$; *Rel-GFP/+* larvae raised at 28°C, fixed, and stained with

propidium iodide (PI). Shown are representative images of hemocytes: brightfield (A-C), GFP to

visualize Rel-GFP localization (A'-C') and PI to mark nuclei (A''-C''). Plasmatocytes are shown

in A and B, lamellocytes are shown in C. (D) Bar graph showing the proportion of hemocytes

with a predominantly nuclear Rel-GFP localization in the indicated genotypes (see Materials and

Methods for analysis details). Complete genotypes $tuSz^1/Y$; *Rel-GFP/+, hop^{Tum}/Y; Rel-GFP/+*

and w^{1118}/Y; *Rel-GFP/+* are simplified to $tuSz^1$, hop^{Tum}, and w^{1118} for clarity. Distinct letter labels

denote differences with $p < 0.05$.

Figure 4. Expression of the Rel-GFP fusion protein in hemocytes from parasitoid infected w^{1118}/Y; Rel-GFP/+ larvae. Brightfield (A-C) and GFP (A'-C') images. Rel-GFP expression in an unknown non-hemocyte cell type is indicated by * (B-C).

Figure 5. Rel/NFκB signaling modifies the $tuSz^1$ phenotype. (A) Penetrance of the $tuSz^1$ phenotype in adult flies of the indicated genotypes. Data are plotted as the average penetrance ± standard error across performed replicates. The mean penetrance of each replicate is visualized with open circles. * p < 0.05 compared to *GAL4-RNAi*. (B) Penetrance of the $tuSz^1$ phenotype in adult flies of the indicated genotypes. Data are plotted as the average penetrance ± standard error across performed replicates. The mean penetrance of each replicate is visualized with open circles. Distinct letter labels denote differences with p < 0.05 by Tukey's HSD.

REFERENCES

Afgan, E., Baker, D., Batut, B., van den Beek, M., Bouvier, D., Čech, M., Chilton, J., Clements, D., Coraor, N., Grüning, B.A., Guerler, A., Hillman-Jackson, J., Hiltemann, S., Jalili, V., Rasche, H., Soranzo, N., Goecks, J., Taylor, J., Nekrutenko, A., Blankenberg, D., 2018. The Galaxy platform for accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucleic Acids Research 46, W537–W544. https://doi.org/10.1093/nar/gky379

Agaisse, H., Perrimon, N., 2004. The roles of JAK/STAT signaling in *Drosophila* immune responses. Immunol. Rev. 198, 72–82.

Ahmad, R., Sorrell, M.F., Batra, S.K., Dhawan, P., Singh, A.B., 2017. Gut permeability and mucosal inflammation: bad, good or context dependent. Mucosal Immunol 10, 307–317. https://doi.org/10.1038/mi.2016.128

Anderl, I., Vesala, L., Ihalainen, T.O., Vanha-aho, L.-M., Andó, I., Rämet, M., Hultmark, D., 2016. Transdifferentiation and proliferation in two distinct hemocyte lineages in *Drosophila melanogaster* larvae after wasp infection. PLOS Pathogens 12, e1005746. https://doi.org/10.1371/journal.ppat.1005746

Andrews, S., 2010. FastQC: A quality control tool for high throughput sequence data [WWW Document]. URL http://www.bioinformatics.babraham.ac.uk/projects/fastqc/ \ (accessed 8.30.19).

Ashburner, M., Ball, C.A., Blake, J.A., Botstein, D., Butler, H., Cherry, J.M., Davis, A.P., Dolinski, K., Dwight, S.S., Eppig, J.T., Harris, M.A., Hill, D.P., Issel-Tarver, L., Kasarskis, A., Lewis, S., Matese, J.C., Richardson, J.E., Ringwald, M., Rubin, G.M., Sherlock, G., 2000. Gene ontology: tool for the unification of biology. The Gene Ontology Consortium. Nat Genet 25, 25–29. https://doi.org/10.1038/75556

479   Barnabei, L., Laplantine, E., Mbongo, W., Rieux-Laucat, F., Weil, R., 2021. NF-κB: At the
480   borders of autoimmunity and inflammation. Front Immunol 12, 716469.
481   https://doi.org/10.3389/fimmu.2021.716469
482   Bergé, J.B., Feyereisen, R., Amichot, M., 1998. Cytochrome P450 monooxygenases and
483   insecticide resistance in insects. Philos Trans R Soc Lond B Biol Sci 353, 1701–1705.
484   https://doi.org/10.1098/rstb.1998.0321
485   Bharucha, K.N., Tarr, P., Zipursky, S.L., 2008. A glucagon-like endocrine pathway in *Drosophila*
486   modulates both lipid and carbohydrate homeostasis. J Exp Biol 211, 3103–3110.
487   https://doi.org/10.1242/jeb.016451
488   Bischoff, V., Vignal, C., Duvic, B., Boneca, I.G., Hoffmann, J.A., Royet, J., 2006.
489   Downregulation of the *Drosophila* immune response by peptidoglycan-recognition
490   proteins SC1 and SC2. PLoS Pathog 2, e14.
491   https://doi.org/10.1371/journal.ppat.0020014
492   Bosco-Drayon, V., Poidevin, M., Boneca, I.G., Narbonne-Reveau, K., Royet, J., Charroux, B.,
493   2012. Peptidoglycan sensing by the receptor PGRP-LE in the *Drosophila* gut induces
494   immune responses to infectious bacteria and tolerance to microbiota. Cell Host Microbe
495   12, 153–165. https://doi.org/10.1016/j.chom.2012.06.002
496   Brennan, C.A., Anderson, K.V., 2004. *Drosophila*: The genetics of innate immune recognition
497   and response. Annual Review of Immunology 22, 457–483.
498   https://doi.org/10.1146/annurev.immunol.22.012703.104626
499   Chung, H., Sztal, T., Pasricha, S., Sridhar, M., Batterham, P., Daborn, P.J., 2009.
500   Characterization of *Drosophila melanogaster* cytochrome P450 genes. Proc Natl Acad
501   Sci U S A 106, 5731–5736. https://doi.org/10.1073/pnas.0812141106
502   Clemmons, A.W., Lindsay, S.A., Wasserman, S.A., 2015. An effector peptide family required for
503   *Drosophila* Toll-mediated immunity. PLOS Pathogens 11, e1004876.
504   https://doi.org/10.1371/journal.ppat.1004876
505   Comrie, W.A., Faruqi, A.J., Price, S., Zhang, Y., Rao, V.K., Su, H.C., Lenardo, M.J., 2018.
506   RELA haploinsufficiency in CD4 lymphoproliferative disease with autoimmune
507   cytopenias. J Allergy Clin Immunol 141, 1507-1510.e8.
508   https://doi.org/10.1016/j.jaci.2017.11.036
509   Davoodi, S., Galenza, A., Panteluk, A., Deshpande, R., Ferguson, M., Grewal, S., Foley, E.,
510   2019. The immune deficiency pathway regulates metabolic homeostasis in *Drosophila*. J
511   Immunol 202, 2747–2759. https://doi.org/10.4049/jimmunol.1801632
512   Dudzic, J.P., Kondo, S., Ueda, R., Bergman, C.M., Lemaitre, B., 2015. *Drosophila* innate
513   immunity: regional and functional specialization of prophenoloxidases. BMC Biol 13, 81.
514   https://doi.org/10.1186/s12915-015-0193-6
515   Ertürk-Hasdemir, D., Broemer, M., Leulier, F., Lane, W.S., Paquette, N., Hwang, D., Kim, C.-H.,
516   Stöven, S., Meier, P., Silverman, N., 2009. Two roles for the *Drosophila* IKK complex in
517   the activation of Relish and the induction of antimicrobial peptide genes. Proceedings of
518   the National Academy of Sciences 106, 9779–9784.
519   https://doi.org/10.1073/pnas.0812022106
520   Furman, D., Campisi, J., Verdin, E., Carrera-Bastos, P., Targ, S., Franceschi, C., Ferrucci, L.,
521   Gilroy, D.W., Fasano, A., Miller, G.W., Miller, A.H., Mantovani, A., Weyand, C.M.,
522   Barzilai, N., Goronzy, J.J., Rando, T.A., Effros, R.B., Lucia, A., Kleinstreuer, N., Slavich,
523   G.M., 2019. Chronic inflammation in the etiology of disease across the life span. Nat
524   Med 25, 1822–1832. https://doi.org/10.1038/s41591-019-0675-0
525   Ganesan, S., Aggarwal, K., Paquette, N., Silverman, N., 2011. NF-κB/Rel proteins and the
526   humoral immune responses of *Drosophila melanogaster*. Curr Top Microbiol Immunol
527   349, 25–60. https://doi.org/10.1007/82_2010_107
528   Gene Ontology Consortium, 2021. The Gene Ontology resource: enriching a GOld mine.
529   Nucleic Acids Res 49, D325–D334. https://doi.org/10.1093/nar/gkaa1113

Guo, C., Shen, J., 2021. Cytoskeletal organization and cell polarity in the pathogenesis of Crohn's disease. Clinic Rev Allerg Immunol 60, 164–174. https://doi.org/10.1007/s12016-020-08795-5

Hanratty, W.P., Dearolf, C.R., 1993. The *Drosophila Tumorous-lethal* hematopoietic oncogene is a dominant mutation in the *hopscotch* locus. Mol. Gen. Genet. 238, 33–37.

Hedengren, M., BengtÅsling, Dushay, M.S., Ando, I., Ekengren, S., Wihlborg, M., Hultmark, D., 1999. Relish, a central factor in the control of humoral but not cellular immunity in *Drosophila*. Molecular Cell 4, 827–837. https://doi.org/10.1016/S1097-2765(00)80392-5

Hetru, C., Hoffmann, J.A., 2009. NF-κB in the Immune Response of Drosophila. Cold Spring Harb Perspect Biol 1, a000232. https://doi.org/10.1101/cshperspect.a000232

Honti, V., Csordás, G., Kurucz, É., Márkus, R., Andó, I., 2014. The cell-mediated immunity of *Drosophila melanogaster*: Hemocyte lineages, immune compartments, microanatomy and regulation. Developmental & Comparative Immunology, Drosophila Immunity 42, 47–56. https://doi.org/10.1016/j.dci.2013.06.005

Hothorn, T., Bretz, F., Westfall, P., 2008. Simultaneous inference in general parametric models. Biom J 50, 346–363. https://doi.org/10.1002/bimj.200810425

Hou, Z., Shi, F., Ge, S., Tao, J., Ren, L., Wu, H., Zong, S., 2021. Comparative transcriptome analysis of the newly discovered insect vector of the pine wood nematode in China, revealing putative genes related to host plant adaptation. BMC Genomics 22, 189. https://doi.org/10.1186/s12864-021-07498-1

Huelsmann, S., Hepper, C., Marchese, D., Knöll, C., Reuter, R., 2006. The PDZ-GEF Dizzy regulates cell shape of migrating macrophages via Rap1 and integrins in the *Drosophila* embryo. Development 133, 2915–2924. https://doi.org/10.1242/dev.02449

Ilchovska, D.D., Barrow, D.M., 2021. An Overview of the NF-kB mechanism of pathophysiology in rheumatoid arthritis, investigation of the NF-kB ligand RANKL and related nutritional interventions. Autoimmun Rev 20, 102741. https://doi.org/10.1016/j.autrev.2020.102741

Kambris, Z., Brun, S., Jang, I.-H., Nam, H.-J., Romeo, Y., Takahashi, K., Lee, W.-J., Ueda, R., Lemaitre, B., 2006. *Drosophila* immunity: A large-scale *in vivo* RNAi screen identifies five serine proteases required for Toll activation. Current Biology 16, 808–813. https://doi.org/10.1016/j.cub.2006.03.020

Kelley, J.B., Paschal, B.M., 2019. Fluorescence-based quantification of nucleocytoplasmic transport. Methods 157, 106–114. https://doi.org/10.1016/j.ymeth.2018.11.002

Kim, D., Langmead, B., Salzberg, S.L., 2015. HISAT: a fast spliced aligner with low memory requirements. Nat Methods 12, 357–360. https://doi.org/10.1038/nmeth.3317

Kim, M.J., Choe, K.-M., 2014. Basement membrane and cell integrity of self-tissues in maintaining *Drosophila* immunological tolerance. PLoS Genet 10. https://doi.org/10.1371/journal.pgen.1004683

Kim, M.-S., Byun, M., Oh, B.-H., 2003. Crystal structure of peptidoglycan recognition protein LB from *Drosophila melanogaster*. Nat Immunol 4, 787–793. https://doi.org/10.1038/ni952

Kleino, A., Silverman, N., 2014. The *Drosophila* IMD pathway in the activation of the humoral immune response. Dev Comp Immunol 42, 25–35. https://doi.org/10.1016/j.dci.2013.05.014

Klunder, L.J., Faber, K.N., Dijkstra, G., van IJzendoorn, S.C.D., 2017. Mechanisms of cell polarity-controlled epithelial homeostasis and immunity in the intestine. Cold Spring Harb Perspect Biol 9, a027888. https://doi.org/10.1101/cshperspect.a027888

Kr, P., Lee, J., Mortimer, N.T., 2021. The S1A protease family members *CG10764* and *CG4793* regulate cellular immunity in *Drosophila*. microPublication Biology 2021. https://doi.org/10.17912/micropub.biology.000370

Kurata, S., 2014. Peptidoglycan recognition proteins in *Drosophila* immunity. Dev. Comp. Immunol. 42, 36–41. https://doi.org/10.1016/j.dci.2013.06.006

Kurucz, É., Márkus, R., Zsámboki, J., Folkl-Medzihradszky, K., Darula, Z., Vilmos, P., Udvardy, A., Krausz, I., Lukacsovich, T., Gateff, E., Zettervall, C.-J., Hultmark, D., Andó, I., 2007. Nimrod, a putative phagocytosis receptor with EGF repeats in *Drosophila* plasmatocytes. Current Biology 17, 649–654. https://doi.org/10.1016/j.cub.2007.02.041

Lawrence, T., 2009. The nuclear factor NF-kappaB pathway in inflammation. Cold Spring Harb Perspect Biol 1, a001651. https://doi.org/10.1101/cshperspect.a001651

Lemaitre, B., Hoffmann, J., 2007. The host defense of *Drosophila melanogaster*. Annual Review of Immunology 25, 697–743. https://doi.org/10.1146/annurev.immunol.25.022106.141615

Lemaitre, B., Kromer-Metzger, E., Michaut, L., Nicolas, E., Meister, M., Georgel, P., Reichhart, J.M., Hoffmann, J.A., 1995. A recessive mutation, *immune deficiency (imd)*, defines two distinct control pathways in the *Drosophila* host defense. Proc. Natl. Acad. Sci. U.S.A. 92, 9465–9469.

Liao, Y., Smyth, G.K., Shi, W., 2014. featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics 30, 923–930. https://doi.org/10.1093/bioinformatics/btt656

Libert, S., Chao, Y., Chu, X., Pletcher, S.D., 2006. Trade-offs between longevity and pathogen resistance in *Drosophila melanogaster* are mediated by NFkappaB signaling. Aging Cell 5, 533–543. https://doi.org/10.1111/j.1474-9726.2006.00251.x

Love, M.I., Huber, W., Anders, S., 2014. Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology 15, 550. https://doi.org/10.1186/s13059-014-0550-8

Low, W.Y., Ng, H.L., Morton, C.J., Parker, M.W., Batterham, P., Robin, C., 2007. Molecular evolution of glutathione S-transferases in the genus *Drosophila*. Genetics 177, 1363–1375. https://doi.org/10.1534/genetics.107.075838

Lu, X., Wang, M., Qi, J., Wang, H., Li, X., Gupta, D., Dziarski, R., 2006. Peptidoglycan recognition proteins are a new class of human bactericidal proteins. Journal of Biological Chemistry 281, 5895–5907. https://doi.org/10.1074/jbc.M511631200

Luo, H., Hanratty, W.P., Dearolf, C.R., 1995. An amino acid substitution in the *Drosophila hopTum-l* Jak kinase causes leukemia-like hematopoietic defects. EMBO J 14, 1412–1420.

Maillet, F., Bischoff, V., Vignal, C., Hoffmann, J., Royet, J., 2008. The *Drosophila* peptidoglycan recognition protein PGRP-LF blocks PGRP-LC and IMD/JNK pathway activation. Cell Host Microbe 3, 293–303. https://doi.org/10.1016/j.chom.2008.04.002

Maverakis, E., Kim, K., Shimoda, M., Gershwin, M.E., Patel, F., Wilken, R., Raychaudhuri, S., Ruhaak, L.R., Lebrilla, C.B., 2015. Glycans in the immune system and the altered glycan theory of autoimmunity: A critical review. J Autoimmun 0, 1–13. https://doi.org/10.1016/j.jaut.2014.12.002

Medzhitov, R., Janeway, C.A., 2002. Decoding the patterns of self and nonself by the innate immune system. Science 296, 298–300. https://doi.org/10.1126/science.1068883

Mi, H., Muruganujan, A., Casagrande, J.T., Thomas, P.D., 2013. Large-scale gene function analysis with the PANTHER classification system. Nat Protoc 8, 1551–1566. https://doi.org/10.1038/nprot.2013.092

Mihajlovic, Z., Tanasic, D., Bajgar, A., Perez-Gomez, R., Steffal, P., Krejci, A., 2019. Lime is a new protein linking immunity and metabolism in *Drosophila*. Developmental Biology 452, 83–94. https://doi.org/10.1016/j.ydbio.2019.05.005

Miraghazadeh, B., Cook, M.C., 2018. Nuclear Factor-kappaB in autoimmunity: man and mouse. Frontiers in Immunology 9, 613. https://doi.org/10.3389/fimmu.2018.00613

Mortimer, N.T., Fischer, M.L., Waring, A.L., Kr, P., Kacsoh, B.Z., Brantley, S.E., Keebaugh, E.S., Hill, J., Lark, C., Martin, J., Bains, P., Lee, J., Vrailas-Mortimer, A.D., Schlenke,

T.A., 2021. Extracellular matrix protein N-glycosylation mediates immune self-tolerance in *Drosophila melanogaster*. PNAS 118. https://doi.org/10.1073/pnas.2017460118

Mortimer, N.T., Kacsoh, B.Z., Keebaugh, E.S., Schlenke, T.A., 2012. Mgat1-dependent N-glycosylation of membrane components primes *Drosophila melanogaster* blood cells for the cellular encapsulation response. PLoS Pathog 8, e1002819. https://doi.org/10.1371/journal.ppat.1002819

Myllymäki, H., Rämet, M., 2014. JAK/STAT pathway in *Drosophila* immunity. Scand J Immunol 79, 377–385. https://doi.org/10.1111/sji.12170

Myllymäki, H., Valanne, S., Rämet, M., 2014. The *Drosophila* Imd signaling pathway. The Journal of Immunology 192, 3455–3462. https://doi.org/10.4049/jimmunol.1303309

Nam, H.-J., Jang, I.-H., Asano, T., Lee, W.-J., 2008. Involvement of pro-phenoloxidase 3 in lamellocyte-mediated spontaneous melanization in *Drosophila*. Mol Cells 26, 606–610.

Noursadeghi, M., Tsang, J., Haustein, T., Miller, R.F., Chain, B.M., Katz, D.R., 2008. Quantitative imaging assay for NF-κB nuclear translocation in primary human macrophages. J Immunol Methods 329, 194–200. https://doi.org/10.1016/j.jim.2007.10.015

Pantha, P., Chalivendra, S., Oh, D.-H., Elderd, B.D., Dassanayake, M., 2021. A tale of two transcriptomic responses in agricultural pests via host defenses and viral replication. Int J Mol Sci 22, 3568. https://doi.org/10.3390/ijms22073568

Parsons, B., Foley, E., 2016. Cellular immune defenses of *Drosophila melanogaster*. Dev. Comp. Immunol. 58, 95–101. https://doi.org/10.1016/j.dci.2015.12.019

Porcu, E., Sadler, M.C., Lepik, K., Auwerx, C., Wood, A.R., Weihs, A., Sleiman, M.S.B., Ribeiro, D.M., Bandinelli, S., Tanaka, T., Nauck, M., Völker, U., Delaneau, O., Metspalu, A., Teumer, A., Frayling, T., Santoni, F.A., Reymond, A., Kutalik, Z., 2021. Differentially expressed genes reflect disease-induced rather than disease-causing changes in the transcriptome. Nat Commun 12, 5647. https://doi.org/10.1038/s41467-021-25805-y

R Core Team, 2021. R: A language and environment for statistical computing.

Ritchie, M.E., Phipson, B., Wu, D., Hu, Y., Law, C.W., Shi, W., Smyth, G.K., 2015. limma powers differential expression analyses for RNA-sequencing and microarray studies. Nucleic Acids Research 43, e47–e47. https://doi.org/10.1093/nar/gkv007

Rizki, T.M., Rizki, R.M., 1974. Topology of the caudal fat body of the *tumor-w* mutant of *Drosophila melanogaster*. J. Invertebr. Pathol. 24, 37–40.

Romagnani, S., 2006. Immunological tolerance and autoimmunity. Intern Emerg Med 1, 187–196. https://doi.org/10.1007/bf02934736

Royet, J., Gupta, D., Dziarski, R., 2011. Peptidoglycan recognition proteins: modulators of the microbiome and inflammation. Nat Rev Immunol 11, 837–851. https://doi.org/10.1038/nri3089

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., Tinevez, J.-Y., White, D.J., Hartenstein, V., Eliceiri, K., Tomancak, P., Cardona, A., 2012. Fiji: an open-source platform for biological-image analysis. Nature Methods 9, 676–682. https://doi.org/10.1038/nmeth.2019

Sharma, P., Dube, D., Singh, A., Mishra, B., Singh, N., Sinha, M., Dey, S., Kaur, P., Mitra, D.K., Sharma, S., Singh, T.P., 2011. Structural basis of recognition of pathogen-associated molecular patterns and inhibition of proinflammatory cytokines by camel peptidoglycan recognition protein. Journal of Biological Chemistry 286, 16208–16217. https://doi.org/10.1074/jbc.M111.228163

Somogyi, K., Sipos, B., Pénzes, Z., Andó, I., 2010. A conserved gene cluster as a putative functional unit in insect innate immunity. FEBS Lett 584, 4375–4378. https://doi.org/10.1016/j.febslet.2010.10.014

Stofanko, M., Kwon, S.Y., Badenhorst, P., 2010. Lineage tracing of lamellocytes demonstrates
Drosophila macrophage plasticity. PLoS ONE 5, e14051.
https://doi.org/10.1371/journal.pone.0014051
Stöven, S., Ando, I., Kadalayil, L., Engström, Y., Hultmark, D., 2000. Activation of the
Drosophila NF-κB factor Relish by rapid endoproteolytic cleavage. EMBO Rep 1, 347–
352. https://doi.org/10.1093/embo-reports/kvd072
Stöven, S., Silverman, N., Junell, A., Hedengren-Olcott, M., Erturk, D., Engström, Y., Maniatis,
T., Hultmark, D., 2003. Caspase-mediated processing of the *Drosophila* NF-κB factor
Relish. Proceedings of the National Academy of Sciences 100, 5991–5996.
https://doi.org/10.1073/pnas.1035902100
Sun, S.-C., 2017. The non-canonical NF-κB pathway in immunity and inflammation. Nat Rev
Immunol 17, 545–558. https://doi.org/10.1038/nri.2017.52
Tattikota, S.G., Cho, B., Liu, Y., Hu, Y., Barrera, V., Steinbaugh, M.J., Yoon, S.-H., Comjean, A.,
Li, F., Dervis, F., Hung, R.-J., Nam, J.-W., Ho Sui, S., Shim, J., Perrimon, N., 2020. A
single-cell survey of *Drosophila* blood. eLife 9, e54818.
https://doi.org/10.7554/eLife.54818
Tokusumi, T., Sorrentino, R.P., Russell, M., Ferrarese, R., Govind, S., Schulz, R.A., 2009.
Characterization of a lamellocyte transcriptional enhancer located within the *misshapen*
gene of *Drosophila melanogaster*. PLOS ONE 4, e6429.
https://doi.org/10.1371/journal.pone.0006429
Toubi, E., Vadasz, Z., 2019. Innate immune-responses and their role in driving autoimmunity.
Autoimmun Rev 18, 306–311. https://doi.org/10.1016/j.autrev.2018.10.005
Tryselius, Y., Samakovlis, C., Kimbrell, D.A., Hultmark, D., 1992. *CecC*, a cecropin gene
expressed during metamorphosis in *Drosophila* pupae. Eur J Biochem 204, 395–399.
https://doi.org/10.1111/j.1432-1033.1992.tb16648.x
Tydell, C.C., Yuan, J., Tran, P., Selsted, M.E., 2006. Bovine Peptidoglycan Recognition Protein-
S: antimicrobial activity, localization, secretion, and binding properties. The Journal of
Immunology 176, 1154–1162. https://doi.org/10.4049/jimmunol.176.2.1154
Valanne, S., Salminen, T.S., Järvelä-Stölting, M., Vesala, L., Rämet, M., 2019. Immune-
inducible *IBIN* connects immunity and metabolism in *Drosophila melanogaster*. PLOS
Pathogens 15, e1007504. https://doi.org/10.1371/journal.ppat.1007504
Waldner, H., 2009. The role of innate immune responses in autoimmune disease development.
Autoimmun Rev 8, 400–404. https://doi.org/10.1016/j.autrev.2008.12.019
Warnes, G.R., Bolker, B., Bonebakker, L., Gentleman, R., Huber, W., Liaw, A., Lumley, T.,
Maechler, M., Magnusson, A., Moeller, S., Schwartz, M., Venables, B., Galili, T., 2022.
gplots: Various R Programming Tools for Plotting Data.
Wickham, H., 2009. ggplot2: Elegant graphics for data analysis, Use R! Springer-Verlag, New
York. https://doi.org/10.1007/978-0-387-98141-3
Williams, M.J., Ando, I., Hultmark, D., 2005. *Drosophila melanogaster* Rac2 is necessary for a
proper cellular immune response. Genes to Cells 10, 813–823.
https://doi.org/10.1111/j.1365-2443.2005.00883.x
Wobbrock, J.O., Findlater, L., Gergle, D., Higgins, J.J., 2011. The aligned rank transform for
nonparametric factorial analyses using only anova procedures, in: Proceedings of the
SIGCHI Conference on Human Factors in Computing Systems, CHI '11. Association for
Computing Machinery, Vancouver, BC, Canada, pp. 143–146.
https://doi.org/10.1145/1978942.1978963
Yadav, S., Daugherty, S., Shetty, A.C., Eleftherianos, I., 2017. RNAseq analysis of the
Drosophila response to the entomopathogenic nematode *Steinernema*. G3 (Bethesda)
7, 1955–1967. https://doi.org/10.1534/g3.117.041004
Yan, L., 2021. ggvenn: Draw Venn Diagram by "ggplot2."

Zaidman-Rémy, A., Hervé, M., Poidevin, M., Pili-Floury, S., Kim, M.-S., Blanot, D., Oh, B.-H.,
Ueda, R., Mengin-Lecreulx, D., Lemaitre, B., 2006. The *Drosophila* amidase PGRP-LB
modulates the immune response to bacterial infection. Immunity 24, 463–473.
https://doi.org/10.1016/j.immuni.2006.02.012
Zhang, Q., Lenardo, M.J., Baltimore, D., 2017. 30 years of NF-κB: a blossoming of relevance to
human pathobiology. Cell 168, 37–57. https://doi.org/10.1016/j.cell.2016.12.012
Zuk, M., Stoehr, A.M., 2002. Immune defense and host life history. Am. Nat. 160 Suppl 4, S9–
S22. https://doi.org/10.1086/342131
738

A

Phenotypic Penetrance

GAL4-RNAi   Rel-RNAi-1   Rel-RNAi-2   Dredd-RNAi

B

Phenotypic Penetrance

w^{1118} /Y w^{1118} /Y; PGRP-LB^{Delta} /+ $tuSz^1$ /Y $tuSz^1$ /Y; PGRP-LB^{Delta} /+