Loss of self-tolerance leads to altered gene expression and IMD pathway activation in

Drosophila melanogaster

Authors: Pooja Kr and Nathan T. Mortimer

Affiliation: School of Biological Sciences, Illinois State University, Normal, IL, 61790, USA

*Correspondence: Nathan T. Mortimer, School of Biological Sciences, Campus Box 4120,

Illinois State University, Normal, IL 61790, USA, ntmorti@ilstu.edu

ABSTRACT

Immune self-tolerance is the ability of a host's immune system to recognize and avoid triggering immune responses against self-tissue. This allows the host to avoid self-directed immune damage while still responding appropriately to pathogen infection. A breakdown of self-tolerance can lead to an autoimmune state in which immune cells target healthy self-tissue, leading to inflammation and tissue damage. In order to better understand the basic biology of autoimmunity and the role of the innate immune system in maintaining self-tolerance, we have recently characterized the *Drosophila melanogaster tuSz* autoimmune mutant. This mutant strain can serve as a model of innate immune mediated self-tolerance, and here we identify transcripts that are deregulated in flies experiencing a loss of self-tolerance. We found that these changes include the ectopic activation of pro-inflammatory signaling through the Relish/NFκB transcription factor, alterations in transcripts encoding proteins predicted to mediate organismal metabolism, and a downregulation of transcripts linked to developmental processes. This study can provide insight into the transcriptional and physiological changes underlying self-tolerance and autoimmunity.

INTRODUCTION

The maintenance of self-tolerance, the process in which self-tissues are identified and self-directed immune mechanisms are repressed, is an important characteristic of the immune system that protects an organism from pathogen infection (Medzhitov and Janeway 2002; Romagnani 2006). The breakdown of the immune system's ability to discriminate healthy self-tissue from non-self pathogens can lead to damage of a host's own tissues by the immune system. Multiple studies have revealed the importance of self-tolerance and failure of this system has been implicated in a variety of autoimmune disorders and other immune mediated diseases. Although much of the research uncovering mechanisms of immune recognition and

self-tolerance has focused on adaptive immune mechanisms, a growing body of work has

demonstrated the importance of innate immune mechanisms in the suppression of

autoimmunity (Waldner 2009; Toubi and Vadasz 2019).

Innate immune responses are evolutionarily conserved and serve as a first line of

defense against infection (Hoffmann *et al.* 1999). The innate immune system relies upon germ-

line encoded receptors, signaling components and other molecules for the recognition and

destruction of pathogens. When activated, these receptors induce intracellular signaling that

results in the activation of genes involved in host defense. The inappropriate activation and

targeting of these innate immune effectors against self-tissue can contribute to autoimmune

pathology (Toubi and Vadasz 2019). However, mechanisms underlying innate immune

mediated self-tolerance and the genes associated with the maintenance of this system remain

largely unexplored.

The genetic model organism *Drosophila melanogaster* is a valuable model for

understanding conserved innate immune mechanisms (Lemaitre and Hoffmann 2007). Flies are

host to a wide range of pathogens including bacteria, fungi, viruses, and parasites and use both

humoral and cell mediated immune responses to combat these pathogens (Brennan and

Anderson 2004). In *Drosophila*, hemocytes (immune cells) known as plasmatocytes are found in

circulation, and following stimulation, surveil the hemolymph (body cavity) for the presence of

invading pathogens and tissue damage. These macrophage-like plasmatocytes are responsible

for the phagocytosis of microbial pathogens and are involved in the encapsulation response

against macroparasites including parasitoid wasp eggs (Mortimer *et al.* 2012; Honti *et al.* 2014;

Parsons and Foley 2016). Another immune cell type, the lamellocytes, are generally not present

in a healthy fly larva, but are induced following parasitoid wasp infection and are required for

melanotic encapsulation of parasitoid eggs and other foreign objects (Mortimer *et al.* 2012;

Honti *et al.* 2014; Anderl *et al.* 2016).

Drosophila hemocytes also play a role in self-recognition and maintaining self-tolerance.

We have recently characterized the *Drosophila tuSz[1]* temperature sensitive mutant strain which

mounts a self-directed encapsulation response at the restrictive temperature (Mortimer *et al.*

2021). We found that the loss of self-tolerance in *tuSz[1]* is due to two genetic changes. The first

is a gain of function mutation in *hopscotch* (*hop*), the *Drosophila* orthologue of Janus Kinase

(JAK). The JAK-STAT signaling pathway plays a key role in immune priming, and the *hop[Sz]*

allele leads to hemocyte activation and the ectopic production of lamellocytes in naïve larvae

(Mortimer *et al.* 2021). These hemocyte phenotypes are also seen in the *hop[Tum]* gain of function

allele (Hanratty and Dearolf 1993; Luo *et al.* 1995), but interestingly, even though *hop[Tum]* and

tuSz[1] mutants both have hemocyte activation and excess lamellocyte production, *hop[Tum]*

mutants do not display an autoimmune phenotype (Mortimer *et al.* 2021). This would suggest

that JAK-STAT mediated immune activation alone is insufficient for the loss of self-tolerance,

and that loss of self-tolerance and immune priming are genetically separable. The second

mutation in the *tuSz[1]* strain is a conditional loss of function mutation in the *GCS1* gene. At the

restrictive temperature, the *GCS1[Sz]* allele leads to disruption of protein N-glycosylation of

extracellular matrix proteins (Mortimer *et al.* 2021). Together these findings define a system in

which *Drosophila* self-tolerance is maintained by the ability of primed hemocytes to sense a self-

signal that is encoded in the N-glycosylation of extracellular matrix proteins, and the loss of this

self-recognition leads to the melanotic encapsulation of self-tissues. Thus, hemocytes can

mediate both the activation of an immune response and the maintenance of self-tolerance

thereby regulating immune homeostasis.

Here, we utilize the *Drosophila tuSz[1]* and *hop[Tum]* mutant strains to identify key

transcriptional changes underlying immune priming, self-tolerance, and autoimmunity.

Differentially expressed transcripts often reveal disease-induced changes (Porcu *et al.* 2021), so

our analysis of transcript abundance changes in these mutants may allow us to identify

processes that occur in autoimmune reactions following the loss of self-tolerance. Our results

further suggest that the pro-inflammatory Immune deficiency (Imd) signaling pathway may be

dysregulated in the $tuSz^1$ loss of self-tolerance phenotype.

MATERIALS AND METHODS

Insect Stocks

87   The *D. melanogaster* strains w^{1118} (BDSC: 5905), $tuSz^1$ (BDSC: 5834),

PBac{GFP.FPTB-Rel}VK00037 (*Rel-GFP*) (BDSC: 81268), hop^{Tum} (BDSC: 8492) were obtained

from the Bloomington Drosophila Stock Center (BDSC). All fly stocks were maintained on

standard *Drosophila* medium on a 12-hour light-dark cycle. The study also uses the parasitoid

wasp *Leptopilina boulardi* (strain Lb17) which is maintained on the *Canton S D. melanogaster*

strain.

RNA Isolation

94   Total RNA was extracted from late third instar w^{1118}, $tuSz^1$ and hop^{Tum} larvae raised at

28º. In total, three biological replicates were prepared with each biological replicate consisting of

20 pooled larvae. RNA was extracted from whole larvae using Trizol (Thermo Fisher Scientific)

followed by QIAGEN RNeasy Micro clean-up (QIAGEN; both according to manufacturer's

instructions). RNA concentration was measured using a nanodrop and the integrity was verified

using agarose gel electrophoresis. The RNA samples were sent to Novogene (Sacramento, CA)

for RNA sequencing (150bp paired end reads, Illumina).

Differential Gene Expression Analysis

102   Sequence data were obtained in the form of FASTQ reads and analyzed using the

Galaxy web platform (https://usegalaxy.org/) (Afgan *et al.* 2018). Initial quality control was

performed using FastQC (version 0.11.8) (Andrews 2010) to trim reads below a quality score

threshold of 30. Reads were then mapped to the *D. melanogaster* reference genome dm6 using

HISAT2 (version 2.1.0) (Kim *et al.* 2015). The resulting BAM format files were analyzed

using featureCounts (version 1.6.3) (Liao *et al.* 2014) to generate read counts for each gene in

the reference genome annotation. The counts generated by featureCounts were subsequently

used to determine the differentially expressed genes using DESeq2 (version 2.11.40.6) (Love *et*

al. 2014). The results were filtered and significant differentially expressed genes (DEGs) were

identified after applying significance cut-offs of a false discovery rate (FDR) < 0.05 and $\log_2$ fold

change ≥ 1 (for upregulated transcripts) or ≤ -1 (for downregulated transcripts).

Functional Annotation

114   The Gene Ontology (GO) Resource (http://geneontology.org/) and PANTHER

115   classification system (Ashburner *et al.* 2000; Mi *et al.* 2013; Gene Ontology Consortium 2021)

116   were used for GO term analysis of each list of DEGs. The FDR threshold to determine term

117   enrichment was 0.05.

Statistical Analysis

119   All statistical analyses described in the following sections were done in the R statistical

120   computing environment (R Core Team 2021) as detailed. Graphs were produced using the

121   ggplot2 and ggvenn R packages (Wickham 2009; Yan 2021).

Quantitative PCR (qPCR)

qPCR was used to confirm the differential expression of selected genes of interest.

Primers for individual genes were designed according to the FlyPrimerBank (Hu *et al.* 2013) and

obtained from Integrated DNA Technologies (Table S1). Ten larvae from each genotype raised

at 28⁰ were collected and total RNA was extracted using Trizol (Thermo Fisher Scientific). RNA

concentration was measured using a Nanodrop and samples were normalized to 2.5 μg for

reverse transcription into cDNA using SuperScript IV (Thermo Fisher Scientific). cDNA samples

were diluted 1:100 in nuclease-free water and 1 μl was used as a template for qPCR

experiments. qPCR was performed using the PowerUp SYBR Green Supermix (Thermo Fisher

Scientific). Samples were run as technical triplicates and a total of three biological replicates

were performed for each primer in each genotype.

For comparison across genotypes, transcript levels for the genes of interest were

normalized to the *Arp1* housekeeping control from the same sample. Within each biological

replicate, the ΔCt value was calculated by subtracting the Ct value of *Arp1* from the Ct value of

each experimental primer set in each genotype. Differences in transcript ΔCt values were tested

across biological replicates using mixed linear models, with genotype as a fixed effect and

biological replicate as a random effect using the lmerTest package (Kuznetsova *et al.* 2017) in

R. Relative quantification was calculated as $2^{-\Delta\Delta Ct}$ (where ΔΔCt is the difference between the

tuSz^1 ΔCt values and *w^{1118}* control ΔCt values for each gene of interest).

Rel Localization

To determine Rel subcellular localization in hemocytes in control and mutant larvae,

w^{1118} and *tuSz^1* unmated females were crossed to *Rel-GFP* males and the crosses were raised

at 28ºC. From these crosses we selected *w^{1118}/Y; Rel-GFP/+* control larvae and *tuSz^1/Y; Rel-*

GFP/+ mutant larvae. Five male third instar larvae from each genotype were bled in PBS and

hemocytes were stained using DAPI (1:500; Invitrogen) to mark nuclei and mounted in

Vectashield mounting medium. To determine Rel localization in hemocytes of parasitoid infected

larvae, late second instar *w^{1118}/Y; Rel-GFP/+* larvae were placed on 35mm Petri dishes filled

with Drosophila medium only (uninfected control) or together (infected) with *L. boulardi* wasps at

25°C. Larvae were dissected at 72-hours post infection (hpi) in PBS and hemocytes were

stained using DAPI mounted in Vectashield for imaging. To determine the *Rel-GFP* expression

in fat body tissue, anterior and posterior regions of fat bodies from w^{1118}/Y; *Rel-GFP/+* and

tuSz^1/Y; *Rel-GFP/+* larvae raised at 28° were dissected in PBS and mounted in Vectashield

mounting medium. Cells and tissues were imaged using a Leica SP8 confocal microscope in the

ISU Confocal Microscopy Facility to assay Rel localization.

The relative proportion of Rel-GFP signal coming from the nucleus vs cytoplasm (N/C

ratio) was calculated using Fiji (Schindelin *et al.* 2012). Image analysis was performed as

described in (Noursadeghi *et al.* 2008). Specifically, the DAPI stain was used to mask the

nucleus allowing for mean cytoplasmic and nuclear fluorescence intensity values to be

independently determined. The N/C ratio was then calculated based on the ratio of mean

nuclear intensity to mean cytoplasmic intensity. Cells with an N/C ratio greater than 1.5 were

considered to have nuclear Rel-GFP localization and cells with an N/C ratio less than 1.5 were

considered to have diffuse Rel-GFP (Rel-GFP exclusion from the nucleus was not observed).

The proportion of cells with nuclear Rel-GFP localization in the *tuSz^1* genetic background was

compared to w^{1118} using Fisher's exact test in R.

RESULTS AND DISCUSSION

Identification of differentially expressed genes in *tuSz^1* and *hop^{Tum}*

To gain a better understanding of the changes involved in JAK-STAT mediated immune

priming and the consequences of the loss of self-tolerance, we performed RNA sequencing on

w^{1118}, *tuSz^1* and *hop^{Tum}* larvae raised at 28⁰ to assess differential transcript abundance. The

resulting sequence reads were mapped to the *D. melanogaster* genome (Table S2) to quantify

transcript levels. We separately compared transcript levels in each mutant genotype to the w^{1118}

control genotype (Tables S3-S6). Both *tuSz^1* and *hop^{Tum}* have gain of function mutations in *hop*,

so any genes that are differentially expressed in both genotypes are likely to be regulated by the

JAK-STAT pathway. We found 832 shared genes that were differentially expressed in both

$tuSz^1$ and hop^{Tum} mutants compared to wildtype, with 384 genes that were upregulated, and 438

genes that were downregulated (Figure 1). Because the JAK-STAT pathway plays a key role in

immune activation (Agaisse and Perrimon 2004), these identified genes might be relevant to the

priming of the immune response.

To begin characterizing these putative immune activation genes, we conducted Gene

Ontology (GO) analysis to identify the biological processes and molecular functions that are

enriched among the common set of differentially expressed genes (Table 1). Analysis of genes

that were upregulated in both $tuSz^1$ and hop^{Tum} identified GO terms that have been linked to

host immunity including defense response to other organism (GO:0098542) and response to

external stimulus (GO:0009605). General immune response genes identified in these categories

include five members of the antimicrobial peptide (AMP)-like *Bomanin* gene family (*BomBc1*,

BomS1, *BomS2*, *BomS4*, *BomT1*) (Clemmons *et al.* 2015) and the Cecropin family AMP gene

CecC (Tryselius *et al.* 1992), along with several immune receptors (*NimB2*, *NimC1*, *NimC4*)

(Kurucz *et al.* 2007; Somogyi *et al.* 2010), the prophenoloxidase family members *PPO1* and

PPO3 (Nam *et al.* 2008; Dudzic *et al.* 2015), and the *induced by infection* (*IBIN*) gene (Valanne

et al. 2019). We also found differential expression of important immune related signaling

pathways such as the JAK-STAT signaling pathway components *dome*, *et* and *upd3* (Myllymäki

and Rämet 2014), the Toll pathway genes *nec* and *SPE* (Kambris *et al.* 2006), and *Rac2* and

Rap1 which are linked to hemocyte function (Williams *et al.* 2005; Huelsmann *et al.* 2006). The

$tuSz^1$ and hop^{Tum} mutations both lead to the ectopic production of lamellocytes, and accordingly,

we see lamellocyte related genes such as *ItgaPS4*, *ItgaPS5*, and *drip* upregulated in these

genotypes (Stofanko *et al.* 2010; Tattikota *et al.* 2020). The proteolysis (GO:0006508) GO term

is also enriched among genes upregulated in both $tuSz^1$ and hop^{Tum}. Notably, 29 members of

the S1A serine protease family are significantly upregulated in both $tuSz^1$ and hop^{Tum}, including

CG10764 and *CG4793* which have recently been shown to be involved in the cellular immune

response to parasitoid infection (Kr *et al.* 2021).

The induction of host immunity is associated with life history tradeoffs, primarily in the

redistribution of energy resources from development or reproduction to the production of an

immune response (Zuk and Stoehr 2002). We found that genes involved with cellular lipid

metabolism (GO:0044255) are enriched among the transcripts upregulated in *tuSz¹* and *hop^{Tum}*,

and conversely that genes linked to development, including regulation of multicellular

organismal development (GO:2000026), tissue morphogenesis (GO:0048729), and cell fate

commitment (GO:0045165) are among those transcripts downregulated in both *tuSz¹* and

hop^{Tum} (Table 1). The upregulation of transcripts involved in metabolism and downregulation of

transcripts linked to developmental processes possibly reflects the immune response-

development tradeoff, and suggests that this tradeoff may be regulated in part by JAK-STAT

signaling or downstream processes. Further study of these genes may help to provide an

understanding of the energy trade-offs and other physiological changes that occur during

immune responses.

Loss of self-tolerance in *tuSz¹* mutants results in differential gene expression

Despite the similarities between the *tuSz¹* and *hop^{Tum}* mutants, *tuSz¹* specifically

displays a loss of self-tolerance phenotype (Mortimer *et al.* 2021), so we wanted to investigate

transcripts that were specifically differentially expressed in *tuSz¹* to identify genes that might be

involved in innate immune self-tolerance and autoimmune mechanisms. Out of 1154 transcripts

that were differentially expressed in *tuSz¹* but not *hop^{Tum}* (referred to as *tuSz¹* specific DEGs),

we found 670 transcripts that were significantly upregulated and 484 transcripts that were

significantly downregulated (Figure 1).

In contrast to the list of upregulated transcripts in common to $tuSz^1$ and hop^{Tum}, GO Term

analysis of these $tuSz^1$ specific DEGs (Table 2) failed to find enrichment of any GO categories

associated with positive regulation of host immunity. Indeed, one of the most highly enriched

terms was the negative regulation of antimicrobial peptide biosynthetic process (GO:0002806).

This term was linked to peptidoglycan recognition protein (PGRP) encoding genes, and we

found that six PGRP gene transcripts are upregulated in $tuSz^1$ mutants. There are 13 members

of this family in the *Drosophila* genome (Table 3), which primarily play roles in the regulation of

the Toll and Imd immune signaling pathways (Kurata 2014). There are five PGRP genes that

encode negative regulators of Imd signaling, and we found that all five of these are specifically

upregulated in $tuSz^1$ mutants, with the sixth altered PGRP playing an unknown role (Table 3).

The finding that whereas positive regulators of immunity were identified as upregulated genes in

both $tuSz^1$ and hop^{Tum}, the enrichment of genes involved in negative regulation of the Imd

pathway specifically in $tuSz^1$ mutants suggests a differential role of immune mechanisms in

immune priming and the autoimmune response to a loss of self-tolerance. We additionally found

26 genes that map to the chitin-based cuticle development (GO: 0040003) term among the

$tuSz^1$ specific DEGs, including members of the *Lcp*, *Acp*, *Twdl*, *Cpr*, and *Ccp* gene families.

Expression of cuticle genes has been linked to infection in *D. melanogaster* and other insects,

although a role in host immunity has yet to be identified (Yadav *et al.* 2017; Hou *et al.* 2021;

Pantha *et al.* 2021).

Many of the additional enriched GO categories among the $tuSz^1$ specific DEGs are

associated with stress responses and metabolism, perhaps providing insight into physiological

changes in $tuSz^1$ mutant larvae due to the occurrence of an autoimmune response (Table 2).

We found an enrichment of oxidoreductase activity (GO: 0016491) genes, including the

glutathione transferase (GST) encoding genes *GstD2, GstD6, GstD7, GstD8, GstE1, GstE3,*

GstE5, and *GstE7* (Low *et al.* 2007). We also found that 18 cytochrome P450 (CYP450)

encoding genes are upregulated in *tuSz[1]* mutants. CYP450s are involved in a wide range of

cellular activities including roles in hormone signaling and detoxification of xenobiotics (Bergé *et*

al. 1998; Chung *et al.* 2009). The specific upregulation of these stress response genes suggests

that loss of self-tolerance may lead to the production of an internal oxidizing environment.

As mentioned previously, metabolic changes often accompany immune responses, and

we found that two general regulators of host metabolism, the *Adipokinetic hormone receptor*

(*AkhR*) (Bharucha *et al.* 2008) and *Linking immunity and metabolism* (*Lime*) (Mihajlovic *et al.*

2019), are upregulated in *tuSz[1]* mutants. *Lime* encodes a transcription factor that plays a key

role in metabolic changes following infection. Accordingly, we found that genes associated with

a wide range of metabolic activities including generation of precursor metabolites and energy

(GO:0006091), carbohydrate derivative metabolic process (GO:1901135), alpha-amino acid

metabolic process (GO:1901605) and lipid metabolic process (GO:0006629) were enriched

among those upregulated in *tuSz[1]* larvae (Table 2). Further analysis of these genes and the

mechanisms they mediate will help to identify the metabolic pathways involved in the response

to a loss of self-tolerance.

Along with upregulated transcripts, transcripts that are downregulated in these

genotypes may also provide insight into the cellular activities that are altered. We found that

genes associated with tissue morphogenesis (GO:0048729) and the establishment of tissue

polarity (GO:0007164) are enriched among the downregulated *tuSz[1]* specific DEGs (Table 2).

The loss of cellular polarity and disrupted tissue organization has also been observed in both

autoimmune mouse models and patients with autoimmune conditions such as inflammatory

bowel disease (Ahmad *et al.* 2017; Klunder *et al.* 2017; Guo and Shen 2021). This may suggest

that tissue structural maintenance is a key self-tolerance checkpoint, in agreement with previous

findings in *Drosophila* (Rizki and Rizki 1974; Kim and Choe 2014; Mortimer *et al.* 2021).

Relish/NFκB is activated in immune tissues in *tuSz[1]* mutants

Based on the specific upregulation of PGRP genes that act as negative regulators of the

Imd pathway (Table 3) (Bischoff *et al.* 2006; Zaidman-Rémy *et al.* 2006; Maillet *et al.* 2008), we

hypothesized that Imd signaling may be dysregulated in the *tuSz[1]* loss of self-tolerance

phenotype. While the role of the Imd pathway in mediating the *Drosophila* humoral immune

response to Gram-negative bacteria has been well established (Lemaitre *et al.* 1995), the

mechanism by which it may participate in the response to loss of self-tolerance is unknown. The

tuSz[1] autoimmune mutant may therefore provide an effective model to begin exploring the role

of the Imd pathway in self-tolerance and autoimmune reactions.

To verify the elevated expression of PGRP genes in *tuSz[1]* larvae, we assayed the

expression of *PGRP-SC1a* and *PGRP-SC2* in *tuSz[1]* and *w[1118]* by qPCR. As expected, both

PGRP-SC1a and *PGRP-SC2* were significantly upregulated in *tuSz[1]* compared to *w[1118]* (*PGRP-*

SC1a: 11.3-fold upregulated, $p < 2.2e-16$; *PGRP-SC2*: 1.3-fold upregulated, $p = 0.0094$) (Figure

2A-B). To further evaluate the ability of qPCR to detect altered expression in *tuSz[1]* mutants, we

also assayed transcript levels of *scrib* (a cell polarity gene) and observed the expected

downregulation in *tuSz[1]* larvae compared to *w[1118]* (1.7-fold downregulated, $p = 1.4 \times 10^{-8}$)

(Figure 2C).

The Imd pathway mainly responds to infection via differential expression of target genes

(Kleino and Silverman 2014). A previous study has identified genes whose expression is

modified by constitutive Imd pathway activation (Davoodi *et al.* 2019). We found that 58 of the

324 genes identified in this study also showed differential expression in *tuSz[1]* larvae (Table S7),

including three of the identified PGRP genes (*PGRP-LB*, *PGRP-LF*, *PGRP-SC2*), and other

immune genes such as *AttA, Rala, Gbp1* and *GILT2*. These findings provide support for the

hypothesis that Imd signaling is dysregulated in *tuSz[1]* mutants.

Gene expression downstream of Imd is largely mediated by the activation of *Relish*

(*Rel*), a *D. melanogaster* member of the NF-κB family of transcriptional activators (Hetru and

Hoffmann 2009). *Rel* activation is regulated by signal-dependent translocation into the nucleus.

Whereas full-length Rel protein is autoinhibited and remains in the cytoplasm, Imd activation

results in cleavage of Rel and the production of an active N-terminal fragment, which

translocates to the nucleus to drive gene expression (Stöven *et al.* 2000). To determine whether

Rel is activated in *tuSz^1* mutants, we used a *Rel-GFP* fusion protein to track the subcellular

localization within the main immune tissues, the hemocytes and fat body. This reporter has GFP

fused to the N-terminal domain of *Relish*, and GFP localization in the nucleus versus cytoplasm

gives a relative readout of pathway activity.

Rel-GFP expression was observed in plasmatocytes in *tuSz^1/Y; Rel-GFP/+* and *w^{1118}/Y;*

Rel-GFP/+ larvae (Figure 3A-B). We found that *Rel-GFP* was significantly more likely to be

concentrated in the nucleus in plasmatocytes from *tuSz^1/Y; Rel-GFP/+* larvae compared to *w^{1118}*

/Y; Rel-GFP/+ control larvae (Fisher's exact test: p = 1.23x10^{-5}; odds ratio = 9.02) (Figure 3D),

suggesting that Imd/*Rel* activation is induced in *tuSz^1* mutants. In addition, *Rel-GFP* expression

and nuclear localization were observed in lamellocytes from *tuSz^1/Y; Rel-GFP/+* larvae (Figure

3C), suggesting that the Imd pathway is also active in this hemocyte type in the *tuSz^1* mutant

background.

To determine if this *Rel* activation is common to cellular immune responses in

Drosophila or if it may be specific to loss of self-tolerance reactions, we assayed *Rel-GFP*

expression and localization in the immune cells of *w^{1118}/Y; Rel-GFP/+* larvae during the

response to parasitoid wasp infection. *w^{1118}/Y; Rel-GFP/+* larvae were infected by *Leptopilina*

boulardi and hemocytes were isolated and imaged 72hpi. We observed minimal *Rel-GFP*

expression in both plasmatocytes (Figure 4A-B) and lamellocytes (Figure 4C) in these infected

larvae. Circulating non-hemocyte cells (marked by an asterisk* in Figure 4B-C) showed strong

Rel-GFP expression and acted as an imaging control. These results suggest that *Rel* and the

Imd pathway are not likely to be involved in the cellular immune response to parasitoid infection.

In agreement with this finding, a previous study has shown that *Rel* mutants have a normal

encapsulation response against parasitoid wasp infection (Hedengren *et al.* 1999).

To test whether *Rel* activation is specific to hemocytes or seen in other immune tissues

in *tuSz^1* mutants, we imaged fat body tissue dissected from *tuSz^1/Y; Rel-GFP/+* and *w^1118 /Y;*

Rel-GFP/+ larvae raised at 28°. We observed minimal *Rel-GFP* expression throughout the fat

body of control larvae (Figure 5A). The *tuSz^1* self-encapsulation phenotype is specific to the

posterior fat body tissue in mutant larvae, so we imaged both anterior and posterior fat body

regions. We found that similar to control, *Rel-GFP* showed minimal expression in anterior fat

body tissue from *tuSz^1/Y; Rel-GFP/+* larvae (Figure 5B). However, *Rel-GFP* expression was

elevated in posterior fat body tissue, including at the sites of self-encapsulation, in *tuSz^1/Y; Rel-*

GFP/+ larvae (Figure 5C).

Rel/NFκB transcriptional activity is associated with pro-inflammatory signaling in

Drosophila and other species (Libert *et al.* 2006; Lawrence 2009; Ganesan *et al.* 2011;

Myllymäki *et al.* 2014). The observed *Rel* activation suggests that inflammation is a feature of

loss of self-tolerance responses in *Drosophila*, a mechanism that is conserved in human

autoimmune disease (Zhang *et al.* 2017; Furman *et al.* 2019). Interestingly, *tuSz^1* mutants

appear to attempt to block pro-inflammatory *Rel* activity by inducing transcription of negative

regulators of the Imd pathway. A similar mechanism is seen in the establishment of immune

tolerance; immune tolerance can be defined as an attempt by the host to mitigate the effects of

infection rather than eliminating the infecting pathogen (Ayres and Schneider 2012), and is often

accompanied by a downregulation of pro-inflammatory signaling (Sears *et al.* 2011; Vale *et al.*

2014). Indeed, a recent study has revealed that the downregulation of Imd signaling plays an

important role in establishing immune tolerance in *Drosophila* (Prakash *et al.* 2021), highlighting

this possible relationship.

The constitutive activation of NF-κB is associated with various chronic inflammatory

conditions, autoimmune diseases, and cancer (Baker *et al.* 2011; Zhang *et al.* 2017; Sun 2017).

Studies using *in vivo* mouse models and human patients with autoimmune disorders have

demonstrated that NF-κB signaling is implicated in the pathogenesis of a number of

autoimmune diseases, such as rheumatoid arthritis, inflammatory bowel disease, multiple

sclerosis, and systemic lupus erythematosus (Atreya *et al.* 2008; Zubair and Frieri 2013;

Miraghazadeh and Cook 2018; Zhou *et al.* 2020; Ilchovska and Barrow 2021). The involvement

of aberrant NF-κB activity in disease has attracted focus in the therapeutic targeting of this

signaling pathway (Viatour *et al.* 2005; Lin *et al.* 2010). Thus, our model may serve as a suitable

system to further understand the role of NF-κB in autoimmunity and allow for investigation into

its full potential as a therapeutic target.

noncoding RNAs are differentially expressed in *hop^{Tum}* and *tuSz^1* mutants

Similar to a recent study looking at the transcriptional response to aging and viral

infection in *Drosophila* (Sheffield *et al.* 2021), we found widespread changes in the expression

of predicted noncoding RNAs (ncRNAs) in *tuSz^1* and *hop^{Tum}* larvae (Figure 6A-B). ncRNAs are a

class of RNA molecules that are transcribed from the genome but are not translated into

proteins. Major classes of ncRNAs include ribosomal RNAs (rRNAs), transfer RNAs (tRNAs),

small ncRNAs, and long ncRNAs (Cech and Steitz 2014). While the roles of rRNAs and tRNAs

are well characterized, multiple studies have found that small and long ncRNAs participate in a

range of biological processes such as genome imprinting, organismal development, and

disease pathogenesis (Wang and Chang 2011; Anastasiadou *et al.* 2018; Fasolo *et al.* 2019),

as well as immune regulation (Atianand *et al.* 2017; Yuan *et al.* 2018).

ncRNAs are categorized according to their transcript size, with small ncRNAs being

<200 nucleotides (nt). These small ncRNA species include microRNAs (miRNAs), small nuclear

RNAs (snRNAs), and small nucleolar RNAs (snoRNAs), and only a limited number of small

RNAs are differentially expressed in $tuSz^1$ and hop^{Tum} larvae (Table 4). On the other hand, a

large number of long ncRNAs (lncRNAs; >200 nt) show differential expression in these mutants

(Figure 6C-D). Amongst the lncRNAs upregulated in both $tuSz^1$ and hop^{Tum}, *Induced by infection*

(*IBIN*) is the only lncRNA that has been studied in the context of *Drosophila* immunity, although

it has been proposed that *IBIN* encodes a short peptide that confers its function (Valanne *et al.*

2019; Ebrahim *et al.* 2021). The significant upregulation of lncRNAs in both genotypes highlights

the potential importance of lncRNAs in mediating immune priming and autoimmune responses.

CONCLUSION

Immune self-tolerance is essential for maintaining immune homeostasis, and breakdown

of self-tolerance mechanisms results in autoimmunity and other immune related diseases.

Although great progress has been made in understanding self-tolerance in the context of

adaptive immunity, innate immune mechanisms regulating self-tolerance and its role in

autoimmunity remain less well characterized. In this study we have identified genes that are

differentially expressed following loss of self-tolerance in the *Drosophila melanogaster tuSz^1*

mutant. These results provide insight into the biological processes that take place in $tuSz^1$

mutant larvae and may contribute toward a better understanding of the events that take place in

loss of self-tolerance autoimmune reactions.

DATA AVAILABILITY

RNAseq data will be made available through the NCBI Gene Expression Omnibus (submission

pending). The authors are happy to make these data available during the review process, and

we will update with the NCBI GEO accession number when available.

ACKNOWLEDGEMENTS

We thank Joshua Hill and Elise N. Le for their assistance on this project. The ISU Confocal

Microscopy Facility was funded by NSF grant DBI-1828136. Stocks obtained from the

Bloomington Drosophila Stock Center (NIH P40OD018537) were used in this study.

FUNDING

Research reported in this publication was supported by the National Institute of General Medical

Sciences of the National Institutes of Health under Award Number R35GM133760 to NTM, a

Grant-in-Aid of Research from the National Academy of Sciences administered by Sigma Xi,

The Scientific Research Society to PK, and the Illinois State University College of Arts and

Sciences University Research Grants program.

FIGURE LEGENDS

Figure 1. Venn diagrams illustrating the numbers of transcripts that are (A) upregulated and (B)

downregulated in the indicated genotypes in comparison with w^{1118} control larvae.

Figure 2. Mean relative quantification ± standard error of *PGRP-SC1a* (A), *PGRP-SC2* (B) and

scrib (C) transcript levels in *tuSz^1* mutant larvae compared to w^{1118} controls. Transcripts were

quantified by qPCR, * indicates $p < 0.05$ relative to w^{1118} control.

Figure 3. Localization of the Rel-GFP fusion protein in hemocytes. Hemocytes were dissected

from w^{1118}/Y; *Rel-GFP/+* and *tuSz^1*/Y; *Rel-GFP/+* larvae raised at 28°C, and stained with

propidium iodide (PI). Shown are representative images of hemocytes: brightfield (A-C), GFP to

visualize Rel-GFP localization (A'-C') and PI to mark nuclei (A''-C''). Rel-GFP shows enriched

nuclear localization in plasmatocytes (B-B'') and lamellocytes (C-C'') in *tuSz^1*/Y; *Rel-GFP/+*, but

not in plasmatocytes (A-A'') in w^{1118}/Y; *Rel-GFP/+*. Note that w^{1118}/Y; *Rel-GFP/+* larvae do not

produce lamellocytes. (D) Bar graph showing the proportion of hemocytes with a predominantly

nuclear Rel-GFP localization (see Materials and Methods for analysis details). Complete

genotypes *tuSz^1/Y*; *Rel-GFP/+* and *w^{1118}/Y*; *Rel-GFP/+* are simplified to *tuSz^1* and *w^{1118}* for

clarity. * indicates p < 0.05 relative to *w^{1118}/Y*; *Rel-GFP/+* control.

Figure 4. Expression of the Rel-GFP fusion protein in hemocytes from parasitoid infected

w^{1118}/Y; *Rel-GFP/+* larvae. Brightfield (A-C) and GFP (A'-C') images. The Rel-GFP fusion

protein is barely detectable in plasmatocytes (A, B) and lamellocytes (C). Strong Rel-GFP

expression is seen in an unknown non-hemocyte cell type (indicated by * in B and C).

Figure 5. Expression of the Rel-GFP fusion protein in the fat body dissected from *w^{1118}/Y*; *Rel-*

GFP/+ and *tuSz^1/Y*; *Rel-GFP/+* larvae raised at 28°C. Minimal Rel-GFP expression is observed

in fat body tissue dissected from *w^{1118}/Y*; *Rel-GFP/+* (A-A') or anterior fat body tissue dissected

from *tuSz^1/Y*; *Rel-GFP/+* (B-B') larvae. Increased Rel-GFP expression is observed in posterior

fat body tissue dissected from *tuSz^1/Y*; *Rel-GFP/+* (C-C') larvae. The posterior fat body is the

site of the loss of self-tolerance in *tuSz^1* mutant larvae, and the characteristic self-encapsulation

phenotype is evident in (C).

Figure 6. Venn diagrams illustrating the numbers of differentially expressed ncRNA transcripts

that are (A) upregulated and (B) downregulated, and the numbers of differentially expressed

lncRNA transcripts that are (C) upregulated and (D) downregulated in the indicated genotypes

in comparison with *w^{1118}* control larvae.

REFERENCES

Afgan, E., D. Baker, B. Batut, M. van den Beek, D. Bouvier *et al.*, 2018 The Galaxy platform for

accessible, reproducible and collaborative biomedical analyses: 2018 update. Nucleic

Acids Research 46: W537–W544.

Agaisse, H., and N. Perrimon, 2004 The roles of JAK/STAT signaling in *Drosophila* immune

responses. Immunol. Rev. 198: 72–82.

Ahmad, R., M. F. Sorrell, S. K. Batra, P. Dhawan, and A. B. Singh, 2017 Gut permeability and

mucosal inflammation: bad, good or context dependent. Mucosal Immunol 10: 307–317.

Anastasiadou, E., L. S. Jacob, and F. J. Slack, 2018 Non-coding RNA networks in cancer. Nat

Rev Cancer 18: 5–18.

Anderl, I., L. Vesala, T. O. Ihalainen, L.-M. Vanha-aho, I. Andó *et al.*, 2016 Transdifferentiation

and proliferation in two distinct hemocyte lineages in *Drosophila melanogaster* larvae

after wasp infection. PLOS Pathogens 12: e1005746.

Andrews, S., 2010 FastQC: A quality control tool for high throughput sequence data.

Ashburner, M., C. A. Ball, J. A. Blake, D. Botstein, H. Butler *et al.*, 2000 Gene ontology: tool for

the unification of biology. The Gene Ontology Consortium. Nat Genet 25: 25–29.

Atianand, M. K., D. R. Caffrey, and K. A. Fitzgerald, 2017 Immunobiology of Long Noncoding

RNAs. Annu Rev Immunol 35: 177–198.

Atreya, I., R. Atreya, and M. F. Neurath, 2008 NF-kappaB in inflammatory bowel disease. J

Intern Med 263: 591–596.

Ayres, J. S., and D. S. Schneider, 2012 Tolerance of infections. Annual Review of Immunology

30: 271–294.

Baker, R. G., M. S. Hayden, and S. Ghosh, 2011 NF-κB, inflammation, and metabolic disease.

Cell Metab 13: 11–22.

Bergé, J. B., R. Feyereisen, and M. Amichot, 1998 Cytochrome P450 monooxygenases and

insecticide resistance in insects. Philos Trans R Soc Lond B Biol Sci 353: 1701–1705.

Bharucha, K. N., P. Tarr, and S. L. Zipursky, 2008 A glucagon-like endocrine pathway in

Drosophila modulates both lipid and carbohydrate homeostasis. J Exp Biol 211: 3103–

3110.

Bischoff, V., C. Vignal, B. Duvic, I. G. Boneca, J. A. Hoffmann *et al.*, 2006 Downregulation of the

Drosophila immune response by peptidoglycan-recognition proteins SC1 and SC2. PLoS

Pathog 2: e14.

Brennan, C. A., and K. V. Anderson, 2004 *Drosophila*: The genetics of innate immune

recognition and response. Annual Review of Immunology 22: 457–483.

Cech, T. R., and J. A. Steitz, 2014 The noncoding RNA revolution-trashing old rules to forge

new ones. Cell 157: 77–94.

Chung, H., T. Sztal, S. Pasricha, M. Sridhar, P. Batterham *et al.*, 2009 Characterization of

Drosophila melanogaster cytochrome P450 genes. Proc Natl Acad Sci U S A 106: 5731–

5736.

Clemmons, A. W., S. A. Lindsay, and S. A. Wasserman, 2015 An effector peptide family

required for *Drosophila* Toll-mediated immunity. PLOS Pathogens 11: e1004876.

Davoodi, S., A. Galenza, A. Panteluk, R. Deshpande, M. Ferguson *et al.*, 2019 The immune

deficiency pathway regulates metabolic homeostasis in *Drosophila*. J Immunol 202:

2747–2759.

Dudzic, J. P., S. Kondo, R. Ueda, C. M. Bergman, and B. Lemaitre, 2015 *Drosophila* innate

immunity: regional and functional specialization of prophenoloxidases. BMC Biol 13: 81.

Ebrahim, S. A. M., G. J. S. Talross, and J. R. Carlson, 2021 Sight of parasitoid wasps

accelerates sexual behavior and upregulates a micropeptide gene in *Drosophila*. Nat

Commun 12: 2453.

Fasolo, F., K. Di Gregoli, L. Maegdefessel, and J. L. Johnson, 2019 Non-coding RNAs in

cardiovascular cell biology and atherosclerosis. Cardiovasc Res 115: 1732–1756.

Furman, D., J. Campisi, E. Verdin, P. Carrera-Bastos, S. Targ *et al.*, 2019 Chronic inflammation

in the etiology of disease across the life span. Nat Med 25: 1822–1832.

Ganesan, S., K. Aggarwal, N. Paquette, and N. Silverman, 2011 NF-κB/Rel proteins and the

humoral immune responses of *Drosophila melanogaster*. Curr Top Microbiol Immunol

349: 25–60.

Gendron, C. M., and S. D. Pletcher, 2017 MicroRNAs *mir-184* and *let-7* alter *Drosophila*

metabolism and longevity. Aging Cell 16: 1434–1438.

Gene Ontology Consortium, 2021 The Gene Ontology resource: enriching a GOld mine. Nucleic

Acids Res 49: D325–D334.

Guo, C., and J. Shen, 2021 Cytoskeletal organization and cell polarity in the pathogenesis of

Crohn's disease. Clinic Rev Allerg Immunol 60: 164–174.

Hanratty, W. P., and C. R. Dearolf, 1993 The *Drosophila Tumorous-lethal* hematopoietic

oncogene is a dominant mutation in the *hopscotch* locus. Mol. Gen. Genet. 238: 33–37.

Hedengren, M., BengtÅsling, M. S. Dushay, I. Ando, S. Ekengren *et al.*, 1999 Relish, a central

factor in the control of humoral but not cellular immunity in *Drosophila*. Molecular Cell 4:

827–837.

Hetru, C., and J. A. Hoffmann, 2009 NF-κB in the immune response of *Drosophila.* Cold Spring

Harb Perspect Biol 1: a000232.

Hoffmann, J. A., F. C. Kafatos, C. A. Janeway, and R. A. Ezekowitz, 1999 Phylogenetic

perspectives in innate immunity. Science 284: 1313–1318.

Honti, V., G. Csordás, É. Kurucz, R. Márkus, and I. Andó, 2014 The cell-mediated immunity of

Drosophila melanogaster: Hemocyte lineages, immune compartments, microanatomy

and regulation. Developmental & Comparative Immunology 42: 47–56.

Hou, Z., F. Shi, S. Ge, J. Tao, L. Ren *et al.*, 2021 Comparative transcriptome analysis of the

newly discovered insect vector of the pine wood nematode in China, revealing putative

genes related to host plant adaptation. BMC Genomics 22: 189.

Hu, Y., R. Sopko, M. Foos, C. Kelley, I. Flockhart *et al.*, 2013 FlyPrimerBank: an online

database for Drosophila melanogaster gene expression analysis and knockdown

evaluation of RNAi reagents. G3 (Bethesda) 3: 1607–1616.

Huelsmann, S., C. Hepper, D. Marchese, C. Knöll, and R. Reuter, 2006 The PDZ-GEF Dizzy

regulates cell shape of migrating macrophages via Rap1 and integrins in the *Drosophila*

embryo. Development 133: 2915–2924.

Ilchovska, D. D., and D. M. Barrow, 2021 An Overview of the NF-kB mechanism of

pathophysiology in rheumatoid arthritis, investigation of the NF-kB ligand RANKL and

related nutritional interventions. Autoimmun Rev 20: 102741.

Kambris, Z., S. Brun, I.-H. Jang, H.-J. Nam, Y. Romeo *et al.*, 2006 *Drosophila* immunity: A

large-scale *in vivo* RNAi screen identifies five serine proteases required for Toll

activation. Current Biology 16: 808–813.

Kim, M. J., and K.-M. Choe, 2014 Basement membrane and cell integrity of self-tissues in

maintaining *Drosophila* immunological tolerance. PLoS Genet 10:.

Kim, D., B. Langmead, and S. L. Salzberg, 2015 HISAT: a fast spliced aligner with low memory

requirements. Nat Methods 12: 357–360.

Kleino, A., and N. Silverman, 2014 The *Drosophila* IMD pathway in the activation of the humoral

immune response. Dev Comp Immunol 42: 25–35.

Klunder, L. J., K. N. Faber, G. Dijkstra, and S. C. D. van IJzendoorn, 2017 Mechanisms of cell

polarity-controlled epithelial homeostasis and immunity in the intestine. Cold Spring Harb

Perspect Biol 9: a027888.

Kr, P., J. Lee, and N. T. Mortimer, 2021 The S1A protease family members *CG10764* and

CG4793 regulate cellular immunity in *Drosophila*. microPublication Biology 2021:.

Kurata, S., 2014 Peptidoglycan recognition proteins in *Drosophila* immunity. Dev. Comp.

Immunol. 42: 36–41.

Kurucz, É., R. Márkus, J. Zsámboki, K. Folkl-Medzihradszky, Z. Darula *et al.*, 2007 Nimrod, a putative phagocytosis receptor with EGF repeats in *Drosophila* plasmatocytes. Current Biology 17: 649–654.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen, 2017 lmerTest package: Tests in linear mixed effects models. Journal of Statistical Software 82: 1–26.

Lawrence, T., 2009 The nuclear factor NF-kappaB pathway in inflammation. Cold Spring Harb Perspect Biol 1: a001651.

Lemaitre, B., and J. Hoffmann, 2007 The host defense of *Drosophila melanogaster*. Annual Review of Immunology 25: 697–743.

Lemaitre, B., E. Kromer-Metzger, L. Michaut, E. Nicolas, M. Meister *et al.*, 1995 A recessive mutation, *immune deficiency (imd)*, defines two distinct control pathways in the *Drosophila* host defense. Proc. Natl. Acad. Sci. U.S.A. 92: 9465–9469.

Liao, Y., G. K. Smyth, and W. Shi, 2014 featureCounts: an efficient general purpose program for assigning sequence reads to genomic features. Bioinformatics 30: 923–930.

Libert, S., Y. Chao, X. Chu, and S. D. Pletcher, 2006 Trade-offs between longevity and pathogen resistance in *Drosophila melanogaster* are mediated by NFkappaB signaling. Aging Cell 5: 533–543.

Lin, Y., L. Bai, W. Chen, and S. Xu, 2010 The NF-kappaB activation pathways, emerging molecular targets for cancer prevention and therapy. Expert Opin Ther Targets 14: 45–55.

Love, M. I., W. Huber, and S. Anders, 2014 Moderated estimation of fold change and dispersion for RNA-seq data with DESeq2. Genome Biology 15: 550.

Low, W. Y., H. L. Ng, C. J. Morton, M. W. Parker, P. Batterham *et al.*, 2007 Molecular evolution of glutathione S-transferases in the genus *Drosophila*. Genetics 177: 1363–1375.

Luo, H., W. P. Hanratty, and C. R. Dearolf, 1995 An amino acid substitution in the *Drosophila hopTum-l* Jak kinase causes leukemia-like hematopoietic defects. EMBO J 14: 1412–1420.

Maillet, F., V. Bischoff, C. Vignal, J. Hoffmann, and J. Royet, 2008 The *Drosophila* peptidoglycan recognition protein PGRP-LF blocks PGRP-LC and IMD/JNK pathway activation. Cell Host Microbe 3: 293–303.

Medzhitov, R., and C. A. Janeway, 2002 Decoding the patterns of self and nonself by the innate immune system. Science 296: 298–300.

Mi, H., A. Muruganujan, J. T. Casagrande, and P. D. Thomas, 2013 Large-scale gene function analysis with the PANTHER classification system. Nat Protoc 8: 1551–1566.

Mihajlovic, Z., D. Tanasic, A. Bajgar, R. Perez-Gomez, P. Steffal *et al.*, 2019 Lime is a new protein linking immunity and metabolism in *Drosophila*. Developmental Biology 452: 83–94.

Miraghazadeh, B., and M. C. Cook, 2018 Nuclear Factor-kappaB in autoimmunity: man and mouse. Frontiers in Immunology 9: 613.

Mortimer, N. T., M. L. Fischer, A. L. Waring, P. Kr, B. Z. Kacsoh *et al.*, 2021 Extracellular matrix protein N-glycosylation mediates immune self-tolerance in *Drosophila melanogaster*. PNAS 118:.

Mortimer, N. T., B. Z. Kacsoh, E. S. Keebaugh, and T. A. Schlenke, 2012 Mgat1-dependent N-glycosylation of membrane components primes *Drosophila melanogaster* blood cells for the cellular encapsulation response. PLoS Pathog 8: e1002819.

Myllymäki, H., and M. Rämet, 2014 JAK/STAT pathway in *Drosophila* immunity. Scand J Immunol 79: 377–385.

Myllymäki, H., S. Valanne, and M. Rämet, 2014 The *Drosophila* Imd signaling pathway. The Journal of Immunology 192: 3455–3462.

Nam, H.-J., I.-H. Jang, T. Asano, and W.-J. Lee, 2008 Involvement of pro-phenoloxidase 3 in

lamellocyte-mediated spontaneous melanization in *Drosophila*. Mol Cells 26: 606–610.

Noursadeghi, M., J. Tsang, T. Haustein, R. F. Miller, B. M. Chain *et al.*, 2008 Quantitative

imaging assay for NF-κB nuclear translocation in primary human macrophages. J

Immunol Methods 329: 194–200.

Pantha, P., S. Chalivendra, D.-H. Oh, B. D. Elderd, and M. Dassanayake, 2021 A tale of two

transcriptomic responses in agricultural pests via host defenses and viral replication. Int

J Mol Sci 22: 3568.

Paredes, J. C., D. P. Welchman, M. Poidevin, and B. Lemaitre, 2011 Negative regulation by

amidase PGRPs shapes the *Drosophila* antibacterial response and protects the fly from

innocuous infection. Immunity 35: 770–779.

Parsons, B., and E. Foley, 2016 Cellular immune defenses of *Drosophila melanogaster*. Dev.

Comp. Immunol. 58: 95–101.

Porcu, E., M. C. Sadler, K. Lepik, C. Auwerx, A. R. Wood *et al.*, 2021 Differentially expressed

genes reflect disease-induced rather than disease-causing changes in the

transcriptome. Nat Commun 12: 5647.

Prakash, A., K. M. Monteith, and P. F. Vale, 2021 Negative regulation of IMD contributes to

disease tolerance during systemic bacterial infection in *Drosophila*:, 2021.09.23.461574

p.

R Core Team, 2021 R: A language and environment for statistical computing.

Rizki, T. M., and R. M. Rizki, 1974 Topology of the caudal fat body of the *tumor-w* mutant of

Drosophila melanogaster. J. Invertebr. Pathol. 24: 37–40.

Romagnani, S., 2006 Immunological tolerance and autoimmunity. Intern Emerg Med 1: 187–

196.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair *et al.*, 2012 Fiji: an open-

source platform for biological-image analysis. Nature Methods 9: 676–682.

Sears, B. F., J. R. Rohr, J. E. Allen, and L. B. Martin, 2011 The economy of inflammation: when

is less more? Trends in Parasitology 27: 382–387.

Sheffield, L., N. Sciambra, A. Evans, E. Hagedorn, C. Goltz *et al.*, 2021 Age-dependent

impairment of disease tolerance is associated with a robust transcriptional response

following RNA virus infection in *Drosophila*. G3 Genes|Genomes|Genetics 11:.

Sokol, N. S., P. Xu, Y.-N. Jan, and V. Ambros, 2008 *Drosophila let-7* microRNA is required for

remodeling of the neuromusculature during metamorphosis. Genes Dev 22: 1591–1596.

Somogyi, K., B. Sipos, Z. Pénzes, and I. Andó, 2010 A conserved gene cluster as a putative

functional unit in insect innate immunity. FEBS Lett 584: 4375–4378.

Stofanko, M., S. Y. Kwon, and P. Badenhorst, 2010 Lineage tracing of lamellocytes

demonstrates *Drosophila* macrophage plasticity. PLoS ONE 5: e14051.

Stöven, S., I. Ando, L. Kadalayil, Y. Engström, and D. Hultmark, 2000 Activation of the

Drosophila NF-κB factor Relish by rapid endoproteolytic cleavage. EMBO Rep 1: 347–

352.

Sun, S.-C., 2017 The non-canonical NF-κB pathway in immunity and inflammation. Nat Rev

Immunol 17: 545–558.

Tattikota, S. G., B. Cho, Y. Liu, Y. Hu, V. Barrera *et al.*, 2020 A single-cell survey of *Drosophila*

blood (B. Lemaître, A. Akhmanova, & B. Lemaître, Eds.). eLife 9: e54818.

Toubi, E., and Z. Vadasz, 2019 Innate immune-responses and their role in driving autoimmunity.

Autoimmun Rev 18: 306–311.

Tryselius, Y., C. Samakovlis, D. A. Kimbrell, and D. Hultmark, 1992 *CecC*, a cecropin gene

expressed during metamorphosis in *Drosophila* pupae. Eur J Biochem 204: 395–399.

Valanne, S., T. S. Salminen, M. Järvelä-Stölting, L. Vesala, and M. Rämet, 2019 Immune-

inducible *IBIN* connects immunity and metabolism in *Drosophila melanogaster*. PLOS

Pathogens 15: e1007504.

Vale, P. F., A. Fenton, and S. P. Brown, 2014 Limiting Damage during infection: lessons from

infection tolerance for novel therapeutics. PLOS Biology 12: e1001769.

Viatour, P., M.-P. Merville, V. Bours, and A. Chariot, 2005 Phosphorylation of NF-kappaB and

IkappaB proteins: implications in cancer and inflammation. Trends Biochem Sci 30: 43–

52.

Waldner, H., 2009 The role of innate immune responses in autoimmune disease development.

Autoimmun Rev 8: 400–404.

Wang, K. C., and H. Y. Chang, 2011 Molecular mechanisms of long noncoding RNAs. Mol Cell

43: 904–914.

Wickham, H., 2009 *ggplot2: Elegant graphics for data analysis.* Springer-Verlag, New York.

Williams, M. J., I. Ando, and D. Hultmark, 2005 *Drosophila melanogaster* Rac2 is necessary for

a proper cellular immune response. Genes to Cells 10: 813–823.

Yadav, S., S. Daugherty, A. C. Shetty, and I. Eleftherianos, 2017 RNAseq analysis of the

Drosophila response to the entomopathogenic nematode *Steinernema*. G3 (Bethesda)

7: 1955–1967.

Yan, L., 2021 *ggvenn: Draw Venn Diagram by "ggplot2."*

Yuan, X., N. Berg, J. W. Lee, T.-T. Le, V. Neudecker *et al.*, 2018 MicroRNA miR-223 as

regulator of innate immunity. J Leukoc Biol 104: 515–524.

Zaidman-Rémy, A., M. Hervé, M. Poidevin, S. Pili-Floury, M.-S. Kim *et al.*, 2006 The *Drosophila*

amidase PGRP-LB modulates the immune response to bacterial infection. Immunity 24:

463–473.

Zhang, Q., M. J. Lenardo, and D. Baltimore, 2017 30 years of NF-κB: a blossoming of relevance

to human pathobiology. Cell 168: 37–57.

Zhou, Y., C. Cui, X. Ma, W. Luo, S. G. Zheng *et al.*, 2020 Nuclear Factor κB (NF-κB)-Mediated

Inflammation in Multiple Sclerosis. Front Immunol 11: 391.

Zubair, A., and M. Frieri, 2013 NF-κB and systemic lupus erythematosus: examining the link. J.

Nephrol. 26: 953–959.

Zuk, M., and A. M. Stoehr, 2002 Immune defense and host life history. Am. Nat. 160 Suppl 4:

S9–S22.

A $w^{1118}/Y; Rel\text{-}GFP/+$

A' Rel-GFP

B $tuSz^1/Y; Rel\text{-}GFP/+$

B'

C $tuSz^1/Y; Rel\text{-}GFP/+$

C'

Table 1. Enriched Gene Ontology (GO) Terms among DEG in common to both *tuSz[1]* and *hop[Tum]*

GO Term	Fold enrichment	FDR
Upregulated genes		
Defense response to other organism (GO:0098542)	2.58	0.014
Response to external stimulus (GO:0009605)	2.11	2.6×10^{-4}
Proteolysis (GO:0006508)	2.87	1.2×10^{-9}
Cellular lipid metabolic process (GO:0044255)	2.43	0.022
Downregulated genes		
Developmental process (GO:0032502)	1.59	4.7×10^{-4}
Regulation of multicellular organismal development (GO:2000026)	2.24	0.015
Tissue morphogenesis (GO:0048729)	2.15	0.029
Cell fate commitment (GO:0045165)	2.99	0.005

Table 2. Enriched GO Terms among *tuSz[1]* specific DEGs

GO Term	Fold enrichment	FDR
Upregulated genes		
Negative regulation of antimicrobial peptide biosynthetic process (GO:0002806)	12.42	0.031
Chitin-based cuticle development (GO: 0040003)	2.89	4.9×10^{-6}
Oxidoreductase activity (GO: 0016491)	2.80	5.4×10^{-13}
Generation of precursor metabolites and energy (GO:0006091)	2.32	0.023
Carbohydrate derivative metabolic process (GO:1901135)	2.14	5.0×10^{-5}
Alpha-amino acid metabolic process (GO:1901605)	3.29	0.006
Lipid metabolic process (GO:0006629)	1.98	0.003
Downregulated genes		

Tissue morphogenesis (GO:0048729)	2.43	2.3×10^{-4}
Establishment of tissue polarity (GO:0007164)	4.17	0.049
Appendage development (GO:0048736)	2.63	0.004
Central nervous system development (GO:0007417)	2.53	0.042
Cell differentiation (GO:0030154)	1.53	0.043
Pattern specification process (GO:0007389)	2.09	0.035
Regulation of cell communication (GO:0010646)	1.83	0.025
Positive regulation of stem cell proliferation (GO:2000648)	9.6	0.003
Metamorphosis (GO:0007552)	2.42	0.002

Table 3. Expression of PGRP family genes in $tuSz^1$ and hop^{Tum} compared to w^{1118} control larvae.

Gene	Pathway	Function	$tuSz^1$ log$_2$FC	hop^{Tum} log$_2$FC
PGRP-LA	IMD	Positive Regulator	-0.38	-0.30
PGRP-LB	IMD	Negative Regulator	**1.83***	-0.18
PGRP-LC	IMD	Positive Regulator	-1.12	-0.63
PGRP-LD	Unknown	Unknown	**-2.77***	-0.81
PGRP-LE	IMD	Positive Regulator	-.018	-0.75
PGRP-LF	IMD	Negative Regulator	**1.37***	0.32
PGRP-SA	Toll	Positive Regulator	1.67	0.99
PGRP-SB1	IMD	Immune Effector	-0.92	-1.95
PGRP-SB2	IMD	Immune Effector	Not detected	Not detected
PGRP-SC1a	IMD	Negative Regulator	**3.46***	1.70
PGRP-SC1b	IMD	Negative Regulator	**3.38***	1.51
PGRP-SC2	IMD	Negative Regulator	**1.23***	-0.19
PGRP-SD	Toll	Positive Regulator	0.66	-0.83

* indicates FDR < 0.05 compared to w^{1118} control; log$_2$FC = log$_2$ Fold Change

Table 4. Number of differentially expressed ncRNAs of each class that are upregulated and downregulated in the indicated genotype in comparison with w^{1118} control larvae.

ncRNA Class	*tuSz*[1]		*hop*[Tum]		Both mutants	
	Up	Down	Up	Down	Up	Down
lncRNAs	55	136	53	130	14	68
miRNAs	0	5	4	1	0	1
snRNAs	1	2	3	1	1	1
snoRNAs	2	2	4	1	2	1