The global spread and invasion capacities of alien ants

Mark K. L. Wong[1,2], Evan P. Economo[3], and Benoit Guénard[2]*

[1] Department of Zoology, University of Oxford, Oxford, United Kingdom

[2] School of Biological Sciences, The University of Hong Kong, Hong Kong SAR, China

[3] Biodiversity and Biocomplexity Unit, Okinawa Institute of Science and Technology Graduate University, Okinawa, Japan

*Corresponding author: bguenard@hku.hk

Abstract

The ecological and economic impacts of biological invasions are usually highly conspicuous, but these are the

outcome of a global, multistage process that is obscured from view. For most taxa, we lack a large-scale picture

of the movements of alien species, the biases and filters that promote or inhibit their spread at each stage, and

blind spots in our ability to detect species during their spread. For instance, countries rely heavily on customs

interceptions to prevent new species introductions, but their efficacy for detecting invaders remains unclear. To

address these gaps, we synthesize and provide data at unprecedented geographic resolution on the global

diversity and distribution of alien ants, a pervasive group strongly impacting humans and ecosystems.

From >145,000 records spanning 602 regions, we identify 522 ant species exhibiting human-mediated

spread, doubling recent estimates of their diversity. We show that movements of alien ant species across regions

globally are non-random and, moreover, that these flows differ by the extents to which species invade—ranging

from arrival to indoor establishment, naturalization, and harmful status. Importantly, we find that almost two-

thirds of the 309 species that naturalize globally—most of which are ground- and litter-dwelling—are absent

from customs interceptions, which record disproportionately high numbers of arboreal species. Our results

illustrate the vast, yet uneven extent of ant invasions globally, and suggest that most alien species bypass

biosecurity controls while spreading successfully worldwide. This raises doubts on the efficacy of current

customs interceptions procedures and highlights a need for radically new approaches.

Keywords

Biodiversity, Formicidae, invasive, species, insects

INTRODUCTION

Incessant movements of humans and goods are driving an unprecedented exchange of organisms worldwide, and the ensuing biological invasions cost the global economy US$26.8 billion annually (Elton, 1958; Seebens et al., 2017; Diagne et al., 2021). The spread and impacts of alien species introduced deliberately for economic or cultural purposes, such as various ornamental plants and mammalian predators, are relatively well studied (Doherty et al., 2016; Van Kleunen et al., 2020). However, barring a few exceptions (Kraemer et al., 2015), there is little information about the global spread of the great majority of alien species that stowaway on transport and that are introduced unintentionally (Turbelin et al., 2017). Furthermore, the extent to which such species, especially numerous invertebrates, plants, and fungi, are detected in customs interceptions—which provide vital gatekeeping functions for the biosecurity of most countries—remains unclear.

As not all alien species pose equal threats, effective management of invasions requires accurate prediction of species' capacities to naturalize and cause harm. From an ecological perspective, this "invasion capacity" of a species is determined by the extent to which its individuals or propagules can overcome geographical barriers to their transport, and—upon subsequent arrival—overcome intrinsic demographic- and extrinsic environmental barriers to their establishment (survival and reproduction) and spread of viable populations (Blackburn et al., 2011). Although geographic patterns of species exchange have been documented at the global scale for several groups (Van Kleunen et al., 2015; Dawson et al., 2017), differences among species' invasion capacities are seldom addressed. Most studies focus only on "naturalized" or "invasive" species—subsets of alien species that can overcome all barriers discussed above (Van Kleunen et al., 2015; Dawson et al., 2017). Other studies have treated records of alien species that are unique to customs interceptions or indoor settings similarly to records of those with established populations in the wild (Miravete et al., 2014; Bertelsmeier et al., 2017, Bertelsmeier & Ollier, 2021). Such approaches overlook fundamental differences in invasion outcomes and may mischaracterize global patterns of invasions. Moreover, they miss key opportunities to understand the mechanisms responsible for species' differing invasion capacities and to develop targeted biosecurity measures in response.

Ants are archetypal stowaway organisms (Weber, 1939) and excellent models for understanding the global spread of unintentionally introduced species. Many alien ants naturalize, and multiple cause severe ecological, economic, and social damages globally (Holway et al., 2002; Diagne et al., 2021). Records of ant species' occurrences across regions worldwide have also been consolidated recently (Guénard et al., 2017). Nonetheless, there have been no attempts to investigate the global spread of alien ant species while explicitly accounting for species' invasion capacities. For instance, in previous studies summarizing the diversity and distribution of alien ants (McGlynn, 1999; Suarez et al., 2010; Miravete et al., 2014; Bertelsmeier, 2021), records of species that were only transported to but not established within a particular region were aggregated with those of other species that established non-native populations within that region. Neglecting such differences in invasion capacities can bias perceptions of biosecurity threats as well as species' viable geographic ranges, which are used widely in ecological niche modelling and climate-change predictions. Similar issues may arise from the aggregation of records from distinct climatic regions or biogeographic realms within coarse country-level boundaries (e.g., Australia, Brazil, China, and the USA in Bertelsmeier, 2021). A more detailed assessment— one addressing current gaps in information on species' invasion capacities, barriers to invasion, and geographic

scale—is therefore needed to clarify the global spread of alien ants as well as the biosecurity threats originating

from and arriving at different parts of the world.

Here, we address key gaps in the knowledge of biological invasions by unintentionally introduced species. We

(i) consolidate the total diversity of alien ant species and map their global distribution at unprecedented

geographic resolution; (ii) clarify species' capacities to overcome different barriers to invasion at regional and

global scales; (iii) contrast the diversity, geographic spread, and habitat use among alien ants with different

invasion capacities, as well as between alien ant species and all ant species; and (iv) test whether species that are

spreading around the world are being detected at customs interceptions, and if there is any ecological bias to

their detectability. Our test of habitat use as a basic correlate of invasion capacity is premised on evidence that

ant assemblages using different vertical habitat strata are distinct in composition and ecological interactions

(Ryder Wilkie et al., 2010; Wong & Guénard, 2017). Alien species occupying different strata may also use

different introduction pathways (e.g., arboreal species in imports of fruits and wood; litter-dwelling species in

imports of substrates) and therefore not be similarly detected in customs interceptions. Investigating alien

species' habitat may thus help identify major gaps in biosecurity measures.

Steps to accomplish the above aims are briefly described as follows (see Methods for additional details). We

assembled data on ant species' occurrences within their native and non-native ranges globally using the Global

Ant Biodiversity Informatics (GABI) database (Guénard et al., 2017). The resultant dataset included 146,917

records corresponding to 17,948 occurrences of alien ant species across 602 non-overlapping regions (all areas

where ants occur on Earth). We then organized the data according to the framework of Blackburn et al. (2011),

which clarifies different invasion barriers. First, we summarized the native occurrences of all alien ant species

across the 602 regions globally. Second, we used locality information to distinguish all non-native occurrences

of alien ant species by invasion extent. We distinguished occurrences of species in customs interceptions (i.e.,

species overcoming geographical barriers to transport); occurrences of species established in indoor settings

such as buildings and greenhouses (i.e., species overcoming demographic barriers and some environmental

barriers to establishment); and occurrences of species established in outdoor settings including semi-natural and

natural habitats (i.e., species overcoming most barriers to establishment). We also distinguished occurrences of

species known to impact the ecosystems they invade as these are of utmost priority to invasive species

management. Third, we identified the vertical habitat strata (i.e., the arboreal, ground-surface, or litter-and-soil

strata; after Lucky et al., 2013) occupied by each alien ant species. The procedures allowed for comparing the

diversity, geographic spread, and habitat use of alien ants with different invasion capacities globally. We also

compared these to the native ranges and habitats of 15,618 ant species and subspecies (i.e., 99% of all extant

taxa, Bolton, 2021; henceforth "all ant species") to identify significant deviations from null expectations.

RESULTS AND DISCUSSION

We identified a total of 522 alien ant species, which have been recorded as native in 522 regions and non-native in 489 regions, out of a total of 602 regions globally (suggesting that 113 regions are currently devoid of any alien ants based on current knowledge). Our data—available in Supporting information[1]—doubles the most recent estimate of alien ant diversity and improves the geographic resolution for mapping their global distributions (cf. 252 species distributed across 195 country-level units in Bertelsmeier, 2021). We placed alien ant species into four groups (Box 1) based on their maximum advancement—at the global scale—along a gradient of invasion extent and biosecurity threat (Fig. 1).

We found that the 522 alien ant species differed significantly (Chi-square tests, $P<0.05$) from the pool of all ant species globally in terms of their taxonomic composition, geographic origins, and use of vertical habitat strata (Fig. 1; Supporting information). This shows that alien ant species comprise a non-random subset of total ant diversity, with disproportionate contributions of species from distinct geographic source pools and vertical habitat strata. Moreover, we found that the taxonomic composition, geographic origins, and use of vertical habitat strata often varied significantly (Chi-square tests, $P<0.05$) among alien species with different invasion capacities at the global scale (Fig. 1; Supporting information). Different sets of factors may thus determine alien species' capacities to overcome the geographical, demographic, and environmental barriers encountered during the successive invasion stages of transport, establishment, and spread (Blackburn et al., 2011). Alternatively, the patterns may also reflect differences in the likelihood of detecting different alien species in different settings and invasion stages (see "Blind spots in customs interceptions" below).

---

[1] Supporting information will be made available upon publication and is not included in this preprint.

Box 1. Invasion capacities of alien ants at global scale
Not all alien species possess equal capacities to invade. At the global scale, the invasion capacities of the 522 known alien ant species can be classified under four levels based on species' maximum success in overcoming different barriers to invasion across 602 regions. The species in each level of invasion capacity are listed in Supporting information.

Level I: Transported Only
162 alien ant species have non-native records that only include individuals in customs interceptions at ports of entry. While such species can overcome barriers to their uptake and transport as stowaways on vehicles and cargo, successfully entering new destinations, they fail to overcome demographic and environmental barriers to their establishment of populations at these destinations. These ants therefore do not represent immediate threats to biosecurity, but their arrival should continue to be monitored.

Level II: Established Indoors Only
51 alien ant species can establish non-native populations within indoor settings only. Like Transported Only species, Established Indoors Only species can overcome barriers to their uptake and transport but upon arrival fail to overcome environmental barriers to establish non-native populations in the wild. These species can, however, overcome demographic barriers to establish (i.e., survive and reproduce) non-native populations in locations buffered from external environments, such as in buildings and greenhouses, where they may impact human activity as indoor pests. Given evidence that these alien species retain their demographic capacities under specific environmental conditions (i.e., indoor settings), they are ideal models for understanding environmental barriers to invasions.

Levels III and IV: Naturalized and Harmful
Of most concern to biosecurity, no less than 309 species—the majority of alien ant species globally—can establish non-native populations in outdoor settings in at least one region. Such ants possess substantially greater invasion capacities than the species described above. Not only can Naturalized and Harmful species overcome geographical, demographic, and environmental barriers to their establishment of non-native populations—even reaching high densities (Sanders et al., 2003), but doing so critically raises their chances for subsequent uptake and transport to expand their non-native ranges (Bertelsmeier & Ollier., 2021). Seventeen well-studied species (the Argentine Ant, Red Imported Fire Ant, African Big-Headed Ant, and others) are distinguished as Harmful (Level IV) based on extensive evidence of their direct and severe impacts on biodiversity and ecosystems, which often swiftly follow naturalization (Supporting information). However, this by no means implies that the remaining 292 Naturalized species (Level III) are of low concern; only that the impacts of these other species represent a critical knowledge gap, with some considered potential pests (e.g., the Ghost Ant, *Tapinoma melanocephalum*). It is unclear whether discrete ecological barriers regulate a transition from naturalized species to "harmful" or "invasive" species; the existence of a barrier also hinges on the precise definition and context of impact—a complex and disputed topic (Colautti & MacIsaac, 2004). Nonetheless, this barrier is tentatively termed "regulation" in Fig. 1.

Examples of alien ant species with different invasion capacities at the global scale. Clockwise from top-left: *Meranoplus bicolor*, a Level I: Transported Only species recorded from customs interceptions in New Zealand and Louisiana; *Carebara diversa*, a Level II: Established Indoors Only species recorded from populations in a storage hangar in Italy; *Tetramorium bicarinatum*, a Level III: Naturalized species with non-native wild populations in multiple regions worldwide; *Solenopsis geminata*, a Level IV: Harmful species with non-native populations that have demonstrated harmful effects on native biodiversity in multiple regions worldwide. Images courtesy of Francois Brassard.

NB: The global classification of alien ants' invasion capacities presented here is based on species' *maximum* invasion extents across multiple regions globally. That is, the invasion capacities of individual species can vary at smaller scales, such as among different zoogeographic realms or individual regions. For instance, a species with an invasion capacity of "Level II: Established Indoors Only" globally may fail to establish indoors in or even enter some regions but not others; a species with an invasion capacity of "Level III: Naturalized" globally may fail to naturalize in or even enter some regions but not others. In addition to global-scale invasion capacities, realm- and region-scale invasion capacities were determined for alien species and used in analyses (Figs. 2,3;5).

Figure 1. Differences among ant species' invasion capacities, source realms, and vertical habitat strata globally. Top row (Invasion Capacity) summarizes the diversity of species in the global pool of ants [All Ants (Native)] as well as for 522 alien ant species in four levels of invasion capacity and biosecurity threat (Level I: Transported Only, Level II: Established Indoors Only, Level III: Naturalized, and Level IV: Harmful) that increase from left to right. Invasion capacities reflect species' relative capacities to overcome successive barriers to invasion (vertical dotted lines) at the global scale. For species in each group, the second row (Source Realms) shows the distribution of their native regions across eleven major zoogeographic realms; the third row (Habitat Strata) shows their distribution across three major vertical habitat strata in terrestrial ecosystems; and the fourth row (Management) summarizes the most relevant actions for biosecurity management, which vary in complexity. Images courtesy of Alex Wild.

158 Global maps of alien species diversity and flows in the numbers of species exchanged between geographic
159 realms are key information for understanding region-specific biosecurity risks and macroscale drivers of
160 invasion (Van Kleunen et al., 2015). We found, however, that both the observed hotspots of alien ant species
161 diversity and flows among geographic realms changed markedly when species of different invasion capacities
162 were considered (Fig. 2–3).
163

165 **Figure 2.** Global distributions (across 602 regions) of the origins of all alien ant species and the destinations of alien ant
166 species of different invasion capacities (at region-scale). Region-scale invasion capacities are based on species' maximum
167 invasion extents within individual regions.
168

170 **Figure 3.** Flows in the diversity of alien ant species of different invasion capacities (at realm-scale) among eleven major
171 zoogeographic realms. Realm-scale invasion capacities are based on species' maximum invasion extents within individual
172 zoogeographic realms. Flows with <10 species are not shown.
173

Invasion hotspots in individual regions and major zoogeographic realms
Across 602 potential destination regions globally, the hotspots where most alien ant species had established
non-native populations within indoor settings were poorly correlated (r=-0.03) with those where most had
overcome environmental barriers to establish non-native populations in the wild (Fig. 2). The highest diversity

of naturalized and harmful alien ant species was found in the southern United States and various oceanic islands
(Florida, Hawaii, and Mascarene Islands record 72, 58, and 55 species respectively). In contrast, regions in
Europe showed the highest diversity of alien species with non-native populations that were restricted to indoor
settings (United Kingdom, France, and Norway record 58, 26 and 23 species, respectively).

Distinctions in invasion capacities were likewise crucial when defining hotspots in alien ant diversity among
Earth's eleven major zoogeographic realms (after Holt et al., 2013). Although the Palearctic realm recorded the
highest diversity of alien ant species overall (156 species), most records comprised species that were only
transported to or which had established non-native populations only in indoor settings within this realm (Fig. 4).
Fewer than half of all alien ant species in the Palearctic realm had overcome environmental barriers and
naturalized—64 species—a number lower than that of substantially smaller areas such as the Saharo-Arabian
and Madagascan realms (Fig. 4). In contrast, most alien ant species had naturalized in the Nearctic (108 species)
and Oceanian (99 species) realms (Fig. 4).

These global patterns align with those observed for naturalized plant and reptile species, which also peak in
diversity in North America (Van Kleunen et al., 2015; Dawson et al., 2017). As postulated for many groups, an
extensive history and high volumes of goods imported to the Nearctic from multiple regions globally could have
substantially lowered geographical barriers to the transport of alien ant species to this realm (Hulme, 2009).
Nonetheless, the large number of studies on biological invasions in the Nearctic (Pyšek et al., 2008) may also
explain the many naturalized species detected. The high diversity of naturalized alien ant species in the
Oceanian realm, which comprises numerous island systems, aligns with broad patterns observed in many groups
(Moser et al., 2018), and may result from various ecological mechanisms that weaken barriers to the
establishment of alien populations in islands, such as reduced biotic resistance and enemy release (Elton, 1958;
Moser et al., 2018).

The relatively low diversity of naturalized alien ant species in the Palearctic realm is striking, given that
geographical barriers to the transport of alien species to this realm would presumably have been extremely low
for an extensive period throughout historic European colonization and maritime trade (Bertelsmeier et al.,
2017). It also contrasts the high diversity of naturalized plant species reported in Europe (Van Kleunen et al.,
2015). Environmental barriers to establishment, such as climatic or biotic resistance associated with the
pervasive temperate climate across the Palearctic realm, may strongly mitigate ant invasions (Lach, 2021).

Two lines of evidence support the hypothesis that the Palearctic realm is distinctly characterized by a
combination of low geographical barriers to transport but high environmental barriers to establishment. First, it
supports the world's highest diversity of alien ant species that establish non-native populations only in indoor
settings (Fig. 4). Second, in comparison with the Palearctic realm, there is a disproportionately high diversity of
species that have successfully naturalized in subtropical regions in the Nearctic realm such as Florida,
Louisiana, and Mississippi (Fig. 4), which have likewise seen extensive historic maritime activity (Bertelsmeier
et al., 2017), but which are subject to warmer climates. Alternatively, it is possible that some ant species thought
to be native to the Palearctic realm may represent old introductions—in particular, from Mediterranean regions

(Seifert et al., 2017)—that naturalized and spread very early throughout the realm. Additional information on

population genetics would help discriminate such "cryptic" alien species.

Figure 4. Numbers of alien ant species exported and imported among eleven major zoogeographic realms. Imported alien ant

species are classified under different invasion capacities (at realm-scale), based on species' maximum invasion extents within

individual zoogeographic realms.

Flow patterns among major zoogeographic realms

Flow patterns of alien ant species exchanged between the eleven major zoogeographic realms changed

substantially depending on the invasion capacities of the species considered (Fig. 3). Given their potential

threats to biosecurity, we tested whether flows of naturalized ant species (including harmful ones) between

individual realms were significantly above or below those expected from random (Fig. 5). We identified the

Oriental realm as a significant exporter of naturalized and harmful ant species to six of ten other realms

globally, exporting no less than 19 and as many as 52 species to these realms. (Fig. 5). Similarly, Turbelin et al.

(2017) identified the Asia-Pacific region as the largest exporter of invasive alien species across multiple plant

and animal groups, with most invasive alien species being native to China and India. While such patterns are

attributed to complex and interacting biogeographic and economic factors, the patterns for ants nonetheless

highlight a need for emphasized biosecurity measures to mitigate the spread of unintentionally introduced alien

species from Asia.

Figure 5. Flows in the diversity of alien ant species with an invasion capacity (at realm-scale) of Level III: Naturalized or higher exchanged among eleven major zoogeographic realms. First diagram shows flows for all realms; remaining diagrams show flows for individual realms. Diagrams illustrate results of resampling tests comparing the observed flows to 999 randomly drawn samples, where flows with values significantly higher than those drawn under random expectations (i.e., in the upper 2.5 percentile) are shown in red, and flows with values significantly lower than those drawn under random expectations (i.e., in the lower 2.5 percentile) are shown in blue; in diagrams for individual realms, flows with values which do not differ significantly from those drawn under random expectations are also shown in grey. Flows with <10 species are not shown.

Blind spots in customs interceptions

The early detection and interception of alien species largely determines the success of biosecurity measures overall (Lodge et al., 2006). Our data showed that 291 alien ant species were recorded in customs interceptions globally (Supporting information). Remarkably, absent from this list were as many as two-thirds of all naturalized (198 species) species and three harmful species at the global scale (Supporting information). Many of these species likely evaded detection in repeated instances; as for numerous alien organisms, naturalized ant populations are unlikely to establish unless multiple colonization events have already occurred (Bertelsmeier & Ollier, 2021). While the general shortage of resources for screening at ports of entry surely accounts for this key blind spot in invasion management, biases in detection may also play a role.

Ecological studies show that systematic biases in detection can produce misestimates in the occurrences of species with (in)conspicuous traits or residing in (un)commonly sampled habitats (Gu & Swihart, 2004). Such biases in the detection of alien organisms are a less explored but nonetheless crucial aspect of biosecurity. We found a possible detection bias in customs interceptions toward arboreal ant species, which predominantly nest and forage in vegetation. Among the 291 alien ant species recorded in customs interceptions globally, the proportion of arboreal species was double and the proportion of litter-dwelling species less than half that among the 309 species capable of naturalizing (including harmful ones) at the global scale (Chi-square test, P<0.001; Supporting information). Furthermore, as many as 87% of the 198 species with naturalization capacity at global scale that were never recorded in customs interceptions were ground- and litter-dwelling (Supporting information). A bias toward arboreal species could occur if inspections of imports focused mainly on plant materials such as fruits and wood, as is often the case when screening for agricultural and domestic pests such as leaf-chewing or wood-boring insects (Kenis et al., 2007). Such measures place less emphasis on imported substrates and may therefore overlook myriad soil fauna that can disrupt native ecosystems (Ferlian et al., 2018).

Aside from potential detection biases, the high proportions of arboreal species among alien ants that are transported only (30%) as compared those that are naturalized or harmful species (≤16%) (Fig. 1) may suggest that arboreal species have low success in establishing alien populations owing to environmental barriers. Arboreal alien ants may face direct biotic resistance from interspecific competition, which is often especially intense in vegetation, owing to the patchy distribution of limiting resources such as nest space and food (Mottl et al., 2021). Arboreal alien ants may also depend on mutualistic or facilitative interactions with plants or herbivorous insects that fail to establish in new regions owing to climatic factors.

Conclusion

The spread of alien species globally is an ongoing ecological disaster. However, as shown here in the case of ants, not all alien species possess equal capacities to overcome geographical and ecological barriers and achieve successful invasions. Alien species with different invasion capacities originate from and accumulate at different regions globally, due to the influence of environmental barriers. Given the limited resources available for invasion management, recipient regions should prioritize biosecurity measures towards corresponding donor regions of species with high invasion capacities. Because the invasion capacity of species is driven in part by

288 ecological attributes, such as affinity for different vertical habitat strata—biosecurity measures that do not take

289 these differences into account (e.g., customs inspections) may achieve limited efficacy. Indeed, the fact that the

290 majority of alien ant species being spread by humans have not been recorded by customs interceptions indicates

291 that existing screening methods are woefully inadequate and need to be modified and/or supplemented by new

292 approaches (e.g., Yasashimoto et al., 2021).

METHODS

Data compilation and organization

Data were compiled as part of the Global Ant Biodiversity Informatics (GABI) project; the details of data compilation are described fully in Guénard et al. (2017), and an update is presented here. In total, nearly two million records for nominal species were compiled from 10,342 scientific publications, as well as 82 and 16 public and private databases, respectively. Dubious and erroneous records were identified based on their mentions in the literature, cross-referencing when updates to checklists of species were available, and through direct communication with numerous experts since 2012.

Native and non-native ranges were identified for each species based on their mentions in the literature and the number of records present for the species (or in a few cases genera) within a particular biogeographic realm or part of it. Information about the locality and habitat of collection were also used to determine the native or non-native status of a species within a particular region. For instance, species encountered only within highly disturbed habitats within a particular region, one of the characteristics of tramp species (Passera, 1994), and with an uncertain native range were considered non-native within the region. For several species that displayed extensive and continuous distributions, some uncertainty was inevitable when demarcating the precise limits of their native and non-native ranges at the scale of individual regions; nonetheless these uncertainties would have limited influence on patterns of species distributions at the larger scale of zoogeographic realms. The resultant dataset, which included all ant species that had at least one non-native record globally, consisted of 146,917 records (including native and non-native records); 4,127 of these were subsequently identified as dubious records. We further summarized the data by geographic regions (Guénard et al., 2017) to yield 17,948 occurrences of alien ant species across 602 non-overlapping regions (i.e., all areas where ants occur on Earth).

Details about the collection locality, sampling method, or any other relevant information associated with each non-native record of a species were used to identify the geographical, demographic, and environmental barriers to invasion that the species had overcome (or failed to overcome) and therefore its invasion extent within that particular region (as described in Introduction). We compared the invasion extents of all non-native records for individual species to determine their invasion capacities (Level I: Transported Only, Level II: Established Indoors Only, or Level III: Naturalized) at the scale of individual regions and zoogeographic realms, as well as at the global scale. In the common case where a species had multiple non-native records with differing invasion extents at a given scale, we used a hierarchical approach and inferred the species' invasion capacity from its maximum invasion extent at that scale.

To provide additional information to guide biosecurity measures, we scrutinized the pool of species with an invasion capacity of "Level III: Naturalized" and from these distinguished 17 "harmful" species for which there was evidence of their impacts on native biota in any of their invaded regions globally (Supporting information). We assigned the 17 species an invasion capacity of "Level IV: Harmful" in all non-native regions where they were previously recorded as "Level III: Naturalized." We used this conservative approach because in the interests of biosecurity it would be safer to assume that a naturalized ant species could impact native biota if it

368 had demonstrable impacts in other regions than to assume otherwise. For instance, naturalized ant species have
369 shown to drive less obvious but nonetheless serious losses in functional diversity before more conspicuous
370 declines in species numbers are observed (Wong et al., 2020).
371
It should be noted that while some studies of biological invasions have considered species to be "invasive" or
"harmful" solely based on their geographic spread (Colautti & MacIsaac, 2004), such parameters are far more
difficult to establish for ants because most parts of the world have received very limited sampling efforts
(Guénard et al., 2012); even in frequently sampled areas, important habitat strata are often overlooked (e.g.,
Wong & Guénard 2017). Furthermore, species of some genera are extremely hard to identify in the absence of
taxonomic expertise (e.g., *Cardiocondyla*, *Nylanderia*, *Pheidole*). Thus, we emphasized alien ant species'
impacts on native biota (as compared to their geographic spread) when flagging species that would be of the
greatest priority for biosecurity. In this regard, our list of 17 harmful alien ant species differs from the IUCN
invasive species list (IUCN, 2021), which includes 19 species but for which, to the best of our knowledge, seven
are lacking in evidence of impacts on native biota but may act as pests that are restricted to indoor environments
or highly modified systems (see Supporting information for a comparison of species in both lists and the
evidence for their impacts). In addition, the IUCN invasive species list lacks some species, such as *Nylanderia*
fulva, for which impacts on native biota have already been demonstrated (LeBrun et al., 2013).
385
We identified the vertical habitat strata used by each alien ant species based on a literature search as well as the
genus-level classification of vertical habitat strata-use as reported in Lucky et al. (2013) with necessary updates
to adhere to recent taxonomic changes. Three habitat strata were considered: the arboreal ("arboreal"), ground-
surface ("ground"), and the litter-and-soil ("litter") strata (Lucky et al., 2013). Ant species' affinity for each of
these strata was determined based on their foraging and nesting behaviors (see Lucky et al., 2013 for details). In
coding the habitat strata of alien ant species in the present study, any species belonging to a genus that
exclusively used a single stratum as reported in Lucky et al. (2013) was coded as that stratum; for instance, all
species of *Hypoponera*, which nest and forage in leaf litter or soil, were coded as "litter." For all other species, a
literature search—typically using the web resources AntWiki (AntWiki, 2021) and AntWeb (AntWeb, 2021) to
identify primary literature—was conducted to determine the stratum or strata used. If we found clear evidence
for a species' use of more than one stratum, all relevant strata (i.e., up to a maximum of three) were coded for
that species. For instance, the Ghost Ant, *Tapinoma melanocephalum*, a species displaying extremely high
environmental plasticity, was coded as "arboreal" and "ground" and "litter" as it has been observed foraging and
nesting in all three strata (e.g., Sharaf et al., 2017). The vertical habitat strata of alien ant species in our study are
provided in Supporting information.
401
Data analysis
We used Chi-square tests to compare the frequency distributions of the native zoogeographic realms and vertical
habitat strata between alien ant species in the four different invasion capacities (defined at the global scale). We
also compared each of these to the frequency distributions of the native zoogeographic realms of all ant species
(using data from GABI, Guénard et al., 2017) and the vertical habitat strata of all ant genera (using updated data
from Lucky et al., 2013). We characterized flows of naturalized ant species (including harmful ones) within and

between different zoogeographic realms, and used resampling tests (after Van Kleunen et al., 2020) to assess whether the observed flows were statistically larger or smaller than expected. The resampling tests involved comparing the observed flows to flows based on 9,999 random draws from the full list of naturalized ant species. If the observed number was in the upper or lower 2.5% quantiles of the resampled values, the flows were considered to be significantly higher or lower, respectively, than expected.

REFERENCES

AntWeb (2021). AntWeb. Version 8.66. California Academy of Science. https://www.antweb.org. Accessed 17 November 2021.

AntWiki (2021). AntWiki. https://www.antwiki.org. Accessed 17 November 2021.

Bertelsmeier, C. (2021). Globalization and the anthropogenic spread of invasive social insects. Current Opinion in Insect Science.

Bertelsmeier, C., & Ollier, S. (2021). Bridgehead effects distort global flows of alien species. Diversity and Distributions, 27(11), 2180-2189.

Bertelsmeier, C., Ollier, S., Liebhold, A., & Keller, L. (2017). Recent human history governs global ant invasion dynamics. Nature ecology & evolution, 1(7), 1-8.

Blackburn, T. M., Pyšek, P., Bacher, S., Carlton, J. T., Duncan, R. P., Jarošík, V., ... & Richardson, D. M. (2011). A proposed unified framework for biological invasions. Trends in ecology & evolution, 26(7), 333-339.

Bolton, B. 2021. An online catalog of the ants of the world. Available from https://antcat.org

Colautti, R. I., & MacIsaac, H. J. (2004). A neutral terminology to define 'invasive' species. Diversity and distributions, 10(2), 135-141.

Dawson, W., Moser, D., Van Kleunen, M., Kreft, H., Pergl, J., Pyšek, P., ... & Essl, F. (2017). Global hotspots and correlates of alien species richness across taxonomic groups. Nature Ecology & Evolution, 1(7), 1-7.

Diagne, C., Leroy, B., Vaissière, A. C., Gozlan, R. E., Roiz, D., Jarić, I., ... & Courchamp, F. (2021). High and rising economic costs of biological invasions worldwide. Nature, 592(7855), 571-576.

Doherty, T. S., Glen, A. S., Nimmo, D. G., Ritchie, E. G., & Dickman, C. R. (2016). Invasive predators and global biodiversity loss. Proceedings of the National Academy of Sciences, 113(40), 11261-11265.

Elton, C. S. (1958). The ecology of invasions by plants and animals. Methuen.

Ferlian, O., Eisenhauer, N., Aguirrebengoa, M., Camara, M., Ramirez-Rojas, I., Santos, F., ... & Thakur, M. P. (2018). Invasive earthworms erode soil biodiversity: A meta-analysis. Journal of Animal Ecology, 87(1), 162-172.

Fournier, A., Penone, C., Pennino, M. G., & Courchamp, F. (2019). Predicting future invaders and future invasions. Proceedings of the National Academy of Sciences, 116(16), 7905-7910.

Gu, W., & Swihart, R. K. (2004). Absent or undetected? Effects of non-detection of species occurrence on wildlife–habitat models. Biological Conservation, 116(2), 195-203.

Guénard, B., Weiser, M. D., & Dunn, R. R. (2012). Global models of ant diversity suggest regions where new discoveries are most likely are under disproportionate deforestation threat. Proceedings of the National Academy of Sciences, 109(19), 7368-7373.

Guenard, B., Weiser, M. D., Gomez, K., Narula, N., & Economo, E. P. (2017). The Global Ant Biodiversity Informatics (GABI) database: synthesizing data on the geographic distribution of ant species (Hymenoptera: Formicidae). Myrmecological News, 24, 83-89.

Holt, B. G., Lessard, J. P., Borregaard, M. K., Fritz, S. A., Araújo, M. B., Dimitrov, D., ... & Rahbek, C. (2013). An update of Wallace's zoogeographic regions of the world. Science, 339(6115), 74-78.

Holway, D. A., Lach, L., Suarez, A. V., Tsutsui, N. D., & Case, T. J. (2002). The causes and consequences of ant invasions. Annual review of ecology and systematics, 33(1), 181-233.

Hulme, P. E. (2009). Trade, transport and trouble: managing invasive species pathways in an era of
globalization. Journal of applied ecology, 46(1), 10-18.

IUCN (2021). SSC Invasive Species Specialist Group: Global Invasive Species Database Available from
http://www.iucngisd.org/gisd/

Kenis, M., Rabitsch, W., Auger-Rozenberg, M. A., & Roques, A. (2007). How can alien species inventories and
interception data help us prevent insect invasions?. Bulletin of entomological research, 97(5), 489-502.

Kraemer, M. U., Sinka, M. E., Duda, K. A., Mylne, A. Q., Shearer, F. M., Barker, C. M., ... & Hay, S. I. (2015).
The global distribution of the arbovirus vectors Aedes aegypti and Ae. albopictus. elife, 4, e08347.

Lach, L. (2021). Invasive ant establishment, spread, and management with changing climate. Current Opinion in
Insect Science.

LeBrun, E. G., Abbott, J., & Gilbert, L. E. (2013). Imported crazy ant displaces imported fire ant, reduces and
homogenizes grassland ant and arthropod assemblages. Biological Invasions, 15(11), 2429-2442.

Lodge, D. M., Williams, S., MacIsaac, H. J., Hayes, K. R., Leung, B., Reichard, S., ... & McMichael, A. (2006).
Biological invasions: recommendations for US policy and management. Ecological applications, 16(6),
2035-2054.

Lucky, A., Trautwein, M. D., Guenard, B. S., Weiser, M. D., & Dunn, R. R. (2013). Tracing the rise of ants-out
of the ground. PLOS one, 8(12), e84012.

McGlynn, T. P. (1999). The worldwide transfer of ants: geographical distribution and ecological invasions.
Journal of biogeography, 26(3), 535-548.

Miravete, V., Roura-Pascual, N., Dunn, R. R., & Gómez, C. (2014). How many and which ant species are being
accidentally moved around the world?. Biology letters, 10(8), 20140518.

Moser, D., Lenzner, B., Weigelt, P., Dawson, W., Kreft, H., Pergl, J., ... & Essl, F. (2018). Remoteness
promotes biological invasions on islands worldwide. Proceedings of the National Academy of Sciences,
115(37), 9270-9275.

Mottl, O., Yombai, J., Novotný, V., Leponce, M., Weiblen, G. D., & Klimeš, P. (2021). Inter-specific
aggression generates ant mosaics in canopies of primary tropical rainforest. Oikos.

Passera, L. (1994) Characteristics of tramp species. In Williams, D.F. (Ed.) Exotic ants: Biology, impact and
control of introduced species. Westview Press, Boulder, Colorado USA.

Pyšek, P., Richardson, D. M., Pergl, J., Jarošík, V., Sixtová, Z., & Weber, E. (2008). Geographical and
taxonomic biases in invasion ecology. Trends in ecology & evolution, 23(5), 237-244.

Ryder Wilkie, K. T., Mertl, A. L., & Traniello, J. F. (2010). Species diversity and distribution patterns of the
ants of Amazonian Ecuador. Plos one, 5(10), e13146.

Sanders, N. J., Gotelli, N. J., Heller, N. E., & Gordon, D. M. (2003). Community disassembly by an invasive
species. Proceedings of the National Academy of Sciences, 100(5), 2474-2477.

Seebens, H., Blackburn, T. M., Dyer, E. E., Genovesi, P., Hulme, P. E., Jeschke, J. M., ... & Essl, F. (2017). No
saturation in the accumulation of alien species worldwide. Nature communications, 8(1), 1-9.

Seifert, B., d'Eustacchio, D., Kaufmann, B., Centorame, M., Lorite, P., & Modica, M. (2017). Four species
within the supercolonial ants of the Tapinoma nigerrimum complex revealed by integrative taxonomy
(Hymenoptera: Formicidae). Myrmecological News, 24, 123-144.

Sharaf, M. R., Fisher, B. L., Collingwood, C. A., & Aldawood, A. S. (2017). Ant fauna (Hymenoptera:
Formicidae) of the Socotra Archipelago (Yemen): zoogeography, distribution and description of a new
species. Journal of Natural History, 51(5-6), 317-378.

Suarez, A. V., McGlynn, T. P., & Tsutsui, N. D. (2010). Biogeographic and taxonomic patterns of introduced
ants. Ant ecology, 233-244.

Turbelin, A. J., Malamud, B. D., & Francis, R. A. (2017). Mapping the global state of invasive alien species:
patterns of invasion and policy responses. Global Ecology and Biogeography, 26(1), 78-92.

Van Kleunen, M., Dawson, W., Essl, F., Pergl, J., Winter, M., Weber, E., ... & Pyšek, P. (2015). Global
exchange and accumulation of non-native plants. Nature, 525(7567), 100-103.

Van Kleunen, M., Xu, X., Yang, Q., Maurel, N., Zhang, Z., Dawson, W., ... & Fristoe, T. S. (2020). Economic
use of plants is key to their naturalization success. Nature communications, 11(1), 1-12.

Weber, N. A. (1939). Tourist ants. Ecology, 20(3), 442-446.

Wong, M. K. L, & Guenard, B. (2017). Subterranean ants: summary and perspectives on field sampling
methods, with notes on diversity and ecology (Hymenoptera: Formicidae). Myrmecological News, 25, 1-
16.

Wong, M. K. L., Guénard, B., & Lewis, O. T. (2020). The cryptic impacts of invasion: functional
homogenization of tropical ant communities by invasive fire ants. Oikos, 129(4), 585-597.

Yasashimoto, T., Sakata, M. K., Sakita, T., Nakajima, S., Ozaki, M., & Minamoto, T. (2021). Environmental
DNA detection of an invasive ant species (Linepithema humile) from soil samples. Scientific reports,
11(1), 1-12.