Microbiota-instructed regulatory T cell differentiation is mediated by a distinct RORγt[+] antigen presenting cell subset

Ranit Kedmi[1], Tariq Najar[1], Kailin R. Mesa[1], Allyssa Grayson[1,2], Lina Kroehling[1], Yuhan Hao[3,4], Stephanie Hao[5], Maria Pokrovskii[1,13], Mo Xu[1,14], Jhimmy Talbot[1,15], Jiaxi Wang[6], Joe Germino[6], Caleb A. Lareau[7], Ansuman T. Satpathy[7], Mark S. Anderson[6], Terri M. Laufer[8], Iannis Aifantis[9], Juliet M. Bartleson[10,16], Paul M. Allen[10], Helena Paidassi[11], James M. Gardner[6,12], Marlon Stoeckius[5,17], Dan R. Littman[1,2†]

[1] Molecular Pathogenesis Program, The Kimmel Center for Biology and Medicine of the Skirball Institute, New York University School of Medicine, New York, NY, USA

[2] Howard Hughes Medical Institute, New York, NY, USA

[3] Center for Genomics and Systems Biology, New York University, New York, NY, USA

[4] New York Genome Center, New York, NY, USA

[5] Technology Innovation Lab, New York Genome Center, New York, NY, USA

[6] Diabetes Center, University of California, San Francisco, San Francisco, CA, USA

[7] Department of Pathology, Stanford University, Stanford, CA 94305, USA. Parker Institute for Cancer Immunotherapy, Stanford University, Stanford, CA 94305, USA, Gladstone-UCSF Institute of Genomic Immunology, San Francisco, CA 94158, USA

[8] Department of Medicine, Perelman School of Medicine, University of Pennsylvania and Department of Medicine, C. Michael Crescenz Veterans Administration Medical Center, Philadelphia, PA, USA

[9] Department of Pathology, New York University School of Medicine, New York, NY, USA

[10]   Department of Pathology and Immunology, Washington University School of Medicine, St.

Louis, MO, USA

[11]   CIRI, Centre International de Recherche en Infectiologie "Team Normal and Pathogenic B

cells" Univ Lyon Inserm U1111 Université Claude Bernard Lyon 1 CNRS UMR5308 ENS de

Lyon, F-69007 Lyon France.

[12]   Department of Surgery, University of California, San Francisco, San Francisco, CA, USA

Current addresses: [13]Calico Life Sciences, LLC, South San Francisco, CA, USA; [14]National

Institute for Biological Sciences, Beijing, China; [15]Fred Hutchinson Cancer Center, Seattle, WA

USA; [16]Buck Institute for Research on Aging, Novato, CA, USA;[17]10X Genomics, Stockholm,

Sweden

[†]   E-mail: dan.littman@med.nyu.edu

Abstract

The mutualistic relationship of gut-resident microbiota and cells of the host immune system promotes homeostasis that ensures maintenance of the microbial community and of a poised, but largely non-aggressive, immune cell compartment[1,2]. Consequences of disturbing this balance, by environmental or genetic factors, include proximal inflammatory conditions, like Crohn's disease, and systemic illnesses, both metabolic and autoimmune. One of the means by which this equilibrium is achieved is through induction of both effector and suppressor or regulatory arms of the adaptive immune system. In mice, *Helicobacter* species induce regulatory (iTreg) and follicular helper (Tfh) T cells in the colon-draining mesenteric lymph nodes under homeostatic conditions, but can instead induce inflammatory Th17 cells when iTreg cells are compromised[3,4]. How *Helicobacter hepaticus* and other gut bacteria direct T cells to adopt distinct functions remains poorly understood. Here, we investigated which cells and molecular components are required to convey the microbial instruction for the iTreg differentiation program. We found that antigen presentation by cells expressing RORγt, rather than by classical dendritic cells, was both required and sufficient for iTreg induction. These RORγt$^+$ cells, likely to be type 3 innate lymphoid cells (ILC3) and/or a recently-described population of Aire$^+$ cells termed Janus cells[5], require the MHC class II antigen presentation machinery, the chemokine receptor CCR7, and α_v integrin, which activates TGF-β, for iTreg cell differentiation. In the absence of any of these, instead of iTreg cells there was expansion of microbiota-specific pathogenic Th17 cells, which were induced by other antigen presenting cells (APCs) that did not require CCR7. Thus, intestinal commensal microbes and their products target multiple APCs with pre-determined features suited to directing appropriate T cell differentiation programs, rather than a common APC that they endow with appropriate functions. Our results illustrate the ability of microbiota to exploit specialized functions

of distinct innate immune system cells, targeting them to achieve the desired composition of

equipoised T cells, thus maintaining tolerance.

Main

A subset of bacterial species among the hundreds that comprise the gut microbiota elicit

stereotypic antigen-specific T cell differentiation programs through mechanisms yet to be

elucidated. It is generally accepted that conventional (or classical) dendritic cells (cDC) that

migrate from tissue to the inductive sites in lymph nodes present microbial antigens to activate

and promote differentiation of naive antigen-specific T cells[4,6-10]. However, the antigen presenting

cells (APCs) that execute these functions according to instructions of distinct intestinal bacterial

species have not been clearly defined. We have chosen to study the APC requirements for T cell

responses to *H. hepaticus* (Hh), which elicits iTreg, Tfh, or pathogenic Th17 cells under different

conditions[3].

Selective requirement of APC subsets for gut microbiota-specific iTreg and inflammatory

Th17 cell differentiation

To study the properties of antigen-presenting cells that direct the differentiation of microbiota-

specific iTreg cells, we transferred into Hh-colonized mice naive Hh-specific CD4$^+$ T cells from

HH7-2 TCR transgenic mice[3]. CFSE-labeled transferred T cells exhibited robust proliferation by

day 3 in colon-draining C1 mesenteric lymph nodes (MLN) of wild type (WT) mice, with up-

regulation of Foxp3 and RORγt, characteristic of colonic iTreg cells[11,12]. In contrast, in mice

deficient for antigen presentation by DC (and, potentially, other cells: *CD11c-Cre;I-Ab$^{f/f}$*,

designated as *MHCII^{ΔCD11c}*), there was no expression of Foxp3 by the Hh-specific T cells, but,

surprisingly, there was substantial proliferation of these cells, with up-regulation of RORγt (Fig.

1a). At 2-3 weeks after transfer, there was expansion in the colonic lamina propria of RORγt- and

T-bet-expressing Hh-specific T cells in mutant mice, characteristic of a pro-inflammatory program

(Extended Data Fig. 1a-c). Endogenous T cells also displayed this phenotype, with fewer Foxp3$^+$

RORγt$^+$ iTreg and expansion of Th17 cells (Extended Data Fig. 1b). This result suggested that

antigen presentation by *Cd11c*-lineage cells is required for iTreg cell differentiation, but that it is

dispensable for the differentiation of pro-inflammatory Th17 cells.

The chemokine receptor CCR7 mediates migration of DC and T cells into lymph nodes, where

adaptive immune responses are initiated, and is critical for tolerogenic responses to food

antigens[13]. In CCR7-deficient mice, iTreg induction in response to Hh colonization failed

(Extended Data Fig. 1d), in agreement with the recent demonstration of a CCR7 requirement for

the differentiation of iTreg cells specific for other *Helicobacter* species[6]. However, we observed

robust priming and proliferation of RORγt$^+$ Hh-specific T cells in CCR7-deficient mice (Extended

Data Fig. 1d). In *CD11c-Cre;Ccr7$^{f/f}$* conditional mutant mice[14] (designated *Ccr7^{ΔCD11c}*), transferred

Hh-specific T cells failed to differentiate into iTreg cells, despite exhibiting robust proliferation, as

in CCR7-deficient mice (Fig. 1b). In the colon of these mice, *Hh*-specific iTreg cells were rare, but

there was accumulation of inflammatory Th17 cells along with elongation of the crypts (Fig. 1c,

Extended Data Fig. 1e,f). Together, our results indicate that, unlike iTreg, microbiota-specific

inflammatory Th17 cell differentiation does not depend on either CCR7 or antigen presentation

by CD11c-expressing cells.

Antigen presentation by *RORγt*-lineage cells is required for microbiota-induced iTreg cell

differentiation

Classical DC, which have been broadly divided into cDC1 and cDC2, comprise multiple cell

subsets that differ in their ontogeny, location, and transcription factor dependency[15,16]. Both cDC1

and cDC2 have been proposed to initiate iTreg responses, largely based on their ability to induce

Treg cell differentiation *in vitro*[9,17]. However, *in vivo* depletion of cDC1 or cDC2 failed to

phenocopy iTreg loss (Extended Data Fig. 2a-c), in agreement with previous reports[6,18-20]. These

findings suggested that antigen presentation by a rare uncharacterized *Cd11c*-lineage myeloid or

non-myeloid cell subset is required for iTreg cell differentiation. To identify putative antigen

presenting cell populations targeted by *CD11c-cre*, we performed CITE-seq analysis of cells

isolated from MLN of Hh-colonized *CD11c-Cre;ROSA26LSLtdTomato* (designated *tdTomato-*

ON$^{\Delta CD11c}$) fate-map mice (Fig. 2a and Extended Data Fig. 3a,b). In addition to expected myeloid

cell subsets, we identified both ILC3 and a recently described Aire⁺ RORγt⁺ population, named

Janus cells (JC)[5,21], among the tdTomato⁺ cells, and these also expressed MHCII (Fig. 2b and

Extended Data Fig. 3b). Using a gating strategy for JC (Extended Data Fig. 4a-d), we confirmed

that they, like ILC3, express *Rorc* (Extended Data Fig. 3a). Consistent with this, GFP⁺ ILC3 and

JC from the C1 MLN of *RORγt-eGFP* mice expressed both CD11c and CD11b or mostly CD11c,

respectively and were both targeted by *CD11c-Cre* (Fig. 2c, Extended Data Fig. 3b and 4e).

Accordingly, many fewer RORγt⁺ cells from MLN of *MHCII*$^{\Delta CD11c}$ mice expressed MHCII, as

compared to WT littermates (Extended Data Fig. 5a).

Expression of MHCII by ILC3 was reported to prevent microbiota-dependent expansion of

inflammatory T cells in the intestine, and was proposed to mediate negative selection of those

cells[22]. Our results suggested that *CD11c-Cre*-expressing RORγt⁺ cells may, instead, be required

for the differentiation of microbiota-specific iTreg cells. This was confirmed by examining iTreg

cell differentiation in mice whose MHCII was inactivated in presumptive ILC3 and JC (*RORγt-*

Cre;I-AB$^{fl/fl}$*,* designated as *MHCII*$^{\Delta RORγt}$). Despite expression of MHCII in cDC2 of these mutant

mice (Fig. 2d), there was complete loss of Hh-specific iTreg cell differentiation in colon-draining

MLN, but intact priming and subsequent expansion of pathogenic Th17 cells in the large intestine

lamina propria (Fig. 2e,f, Extended Data Fig. 5b). Both the donor Hh-specific and host T cells

exhibited loss of iTreg and increase in IFNγ- and IL-17A-producing CD4$^+$ T cells in the large

intestine lamina propria (Fig. 2f, Extended Data Fig. 5c-e). Similar results were observed with Hh-

specific T cells in another conditional mutant strain, *RORγt-Cre; H2-DMa$^{f/f}$* mice (*H2-DMa^{ΔRORγt}*)

deficient in H2-DMa, the mouse equivalent of HLA-DM, required for displacement of invariant

chain peptide and loading of processed peptide on MHCII molecules[23]. This result confirms that

the antigen processing machinery is required in RORγt$^+$ APCs for induction of microbiota-specific

iTregs (Extended Data Fig. 5f).

CCR7-dependent migration by *RORγt*-lineage cells is required for microbiota-induced

iTreg cell differentiation

Intestinal ILC3 have been reported to also employ CCR7 for migration to draining MLN[24,25]. We

found expression of *Ccr7* in both ILC3 and JC (Extended Data Fig. 3a and 6a). We therefore

asked whether CCR7 is required for migration of RORγt$^+$ APCs rather than classical DC for iTreg

induction. Indeed, Hh-specific iTreg cell differentiation was abrogated in the MLN of *Ccr7^{ΔRORγt}*

mice (Fig. 3a). In the colon of *Ccr7^{ΔRORγt}* mutant mice, there was skewing of Hh-specific T cells

towards a Th1-Th17 inflammatory program after 2-3 weeks of adoptive T cell transfer, although

this was less marked than in *Ccr7^{ΔCD11c}* mice, and elongation of crypts was not observed (Fig. 3b

and data not shown). To determine whether loss of CCR7 affects migration of RORγt$^+$ cells to the

MLN, we reconstituted irradiated mice with equal numbers of WT and *RORγt-Cre;Ccr7$^{f/f}$*

(*Ccr7^{ΔRORγt}*) bone marrow cells. Although there was no effect on the ratio of WT and mutant

migratory DC, there was a substantial reduction in the proportion of CCR7-deficient lymphoid

tissue inducer (LTi)-like ILC3, but no significant difference in the proportions of other RORγt[+] cells

in the MLN (Fig 3c and Extended Data Fig. 6b-d). Similarly, in $Ccr7^{\Delta CD11c}$ mice, LTi-like ILC3 lost

CCR7 expression and were reduced in number in the MLN (Extended Data Fig. 6e,f). To further

study a potential role for JC, we examined the T cell response to Hh in *RORγt-cre;Aire^{f/f}* mice and

in Aire-DTR bone marrow-reconstituted mice treated with diphtheria toxin. In neither case was

there an effect on Hh-specific iTreg cell differentiation (Extended Data Fig. 7a,b). We cannot,

however, rule out a role for JC, as Aire may not be required for the microbiota-dependent iTreg

inductive function of these cells and residual cells in DT-treated animals may be sufficient to

support iTreg cell differentiation. Together, these results indicate that RORγt[+] APC, either LTi-like

ILC3 or JC, migrate to the MLN, where they present microbial antigen to naïve T cells to induce

iTreg cell differentiation, and that their failure to migrate results, instead, in inflammatory Th17 cell

differentiation.

Integrin α_v expression by *RORγt*-lineage antigen presenting cells is required for iTreg

differentiation

Differentiation of iTreg cells requires TGF-β signaling in CD4[+] T cells, and defects in this pathway

result in spontaneous colitis[3,11]. TGF-β is released from its latent form on cell surfaces or

extracellular matrix following physical interaction with integrins $\alpha_v\beta_6$ or $\alpha_v\beta_8$[26-28]. Loss of integrins

β_8 or α_v in hematopoietic cells, including in *CD11c-Cre:Itgb8^{f/f}* mice, resulted in reduced colonic

Tregs and in multiorgan inflammation[29-31]. Consistent with those observations, differentiation of

adoptively transferred Hh-specific T cells into iTreg cells was abrogated both following treatment

of mice with anti-β8 antibody and in *CD11c-Cre;Itgav^{f/f}* (Itgav$^{\Delta CD11c}$) recipient mice (Fig. 4a,b). We

examined iTreg cell differentiation in mice with conditional inactivation of *Itgav* in RORγt[+] APCs

and in T cells (*Itgav^{\Delta RORγt}*) (Fig. 4c,d). In the colon-draining MLN of *Itgav^{\Delta RORγ}* mice, there was loss

of integrin α_v (CD51) expression on both ILC3 and JC, and adoptively transferred Hh-specific T

cells failed to express Foxp3 (Fig. 4c and Extended Data Fig. 8a). The MLN were increased in

size (Extended Data Fig. 8b) and the Hh-specific T cells expressed RORγt, but, unlike in control

littermates, they also had elevated T-bet, along with a substantial decrease in CCR6, consistent

with reduced TGF-β signaling (Extended Data Fig. 8c). Notably, β_8 antibody blockade resulted in

the same phenotype (Extended Data Fig. 8d). In the colonic lamina propria of *Itgav$^{\Delta ROR\gamma t}$* mice,

there was loss of both Hh-specific and host-derived iTreg cells, with skewing of CD4$^+$ T cells

towards IFNγ^+ Th1 and pathogenic Th17 programs (Fig. 4d, Extended Data Fig. 8e), suggesting

that expression of integrin α_v on RORγt+ APCs, likely in partnership with integrin β_8, is a general

requirement for intestinal iTreg cell differentiation. Although ILC3 express higher levels of CD51

(Extended Data Fig. 8a), single-cell RNA sequencing analysis of GFP$^+$ cells from pooled lymph

nodes of *Aire* reporter mice[32] showed clustering of JCs into three discrete subpopulations, with

JC2 and JC3 expressing high levels of *Itgav* and *Itgb8* (Extended Data Fig.9a-e). JC and a fraction

of ILC3, obtained from *Itgb8-tdTomato* reporter mice[33], were found to express tdTomato

(Extended Data Fig 9f), and we therefore cannot exclude the requirement of either cell type for

iTreg differentiation. Differentiation of iTreg cells was normal in *CD4-Cre;Itgav$^{f/f}$* mice (Extended

Data Fig. 8f), consistent with a role of the integrin in RORγt$^+$ APCs rather than in TCR$\alpha\beta$ T cells.

Furthermore, we reconstituted mice after irradiation with a mix of MHCII$^{\Delta CD11c}$ and *Itgav$^{\Delta ROR\gamma t}$* bone

marrow cells, resulting in binary expression of MHCII or integrin α_v. In these mice, iTreg cell

differentiation was abolished, consistent with a requirement for both antigen presentation and

activation of TGF-β by the same APC, coupling T cell activation with differentiation cues (Fig. 4e-

f, Extended Data Fig. 8g).

Antigen presentation by RORγt$^+$ APCs is sufficient to promote iTreg cell differentiation

Our results support a role for *RORγt⁺* APCs in microbiota-specific iTreg differentiation, but do not

rule out a requirement for additional conventional antigen presenting cells. Moreover, a population

of DC (T-bet-negative cDC2) was recently reported to express RORγt[34], raising the possibility that

neither ILC3 nor JC is relevant in iTreg cell differentiation. To examine a potential role for a rare

DC subset, we employed *zbtb46-Cre*, considered to specifically target cDC, and *zbtb46* reporter

mice. Because *zbtb46-Cre;Ccr7^{f/-}* mice (CCR7^{Δzbtb46}) were unable to support microbiota-

dependent iTreg cell differentiation (Extended Data Fig. 10a), we profiled by CITE-seq sorted

cells from C1 mLN of *zbtb46-eGFP ; RORγt-Cre;ROSA26LSLtdTomato* (*tdTomato-ON^{RORγt}*) mice,

expressing one or both fluorescent reporters, gated to exclude B and T cells (Extended Data Fig.

10b). Surprisingly, GFP expression was identified on all ILC3 and fate-mapped JC. Zbtb46

expression on ILC3 was confirmed using *mKate2-ON^{zbtb46}*; *RORγt-eGFP* mice (Extended Data

Fig. 10c). We identified a few migratory cDC2 among tdTomato⁺ GFP⁺ cells in *zbtb46-eGFP;*

tdTomato-ON^{RORγt} mice, but these did not form a subcluster to suggest a unique gene signature

and did not exhibit active *Rorc* or integrin α_v mRNA and protein expression (Extended Data Fig.

10d-f), suggesting that they are unlikely to be RORγt⁺ APCs required to direct iTreg cell

differentiation. We performed three-dimensional intravital imaging studies to visualize interactions

of newly-primed Hh-specific T cells with DC and *RORγt⁺* APCs populations in C1 MLN. We utilized

mKate2-ON^{zbtb46}; *RORγt-eGFP* mice to visualize cDC (mKate2⁺) and *RORγt⁺* APCs (eGFP⁺ and

mKate2). However, given that the efficiency of *zbtb46-Cre*-mediated activation of the mKate2

reporter was low (~20% for *RORγt+* APCs and cDC populations) (Extended Data Fig. 11a), we

additionally utilized cell morphology and size analysis to distinguish the few host-derived GFP⁺ T

cells from GFP⁺ *RORγt⁺* APCs. We next transferred dye-labeled *Nur77-eGFP* Hh-specific T cells

into these fluorescent reporter host mice to measure direct interactions of primed T cells (which

up-regulate *Nur77*) with DC and *RORγt⁺* APCs at 15 h after adoptive transfer. Approximately 81%

of GFP⁺ primed Hh-specific T cells were found in contact with at least one *RORγt⁺* APC

with/without DC, as opposed to only ~31% of the non-primed T cells (Extended Data Fig. 11b,c).

Our imaging study and results with conditional mutant mice did not rule out a contribution by DC

towards Hh-specific T cell activation and iTreg differentiation. We therefore wished to determine

whether antigen presentation limited to only *RORγt-Cre*-expressing cells was sufficient to allow

for iTreg cell differentiation. For this purpose, we used mice that express MHCII only in *RORγt⁺*

APCs, and not in DC or other APCs (*RORγt-Cre;I-AB⁻/lsl* , designated *MHCII-ON^{RORγt}*). We

reconstituted irradiated congenic mice with bone marrow from *MHCII^{ΔCD11c}* mice with or without

bone marrow from WT or *MHCII-ON^{RORγt}* mice (Fig. 5a). Flow cytometry analysis confirmed MHCII

expression by *RORγt⁺* cells, but not DC from the C1 MLN and the colon lamina propria of

MHCII^{ΔCD11c};MHCII-ON^{RORγt} bone marrow-reconstituted mice (Fig. 5b, Extended Data Fig.

12a). As expected, in control mice reconstituted with only *MHCII^{ΔCD11c}* bone marrow cells,

in which no MHCII expression was detected in either *RORγt⁺* APC or DC, there was no

differentiation of adoptively transferred Hh-specific iTreg cells, but there was, instead,

differentiation of RORγt⁺Tbet⁺ inflammatory Th17 cells (Fig. 5c,d). In contrast, antigen

presentation by RORγt⁺ cells alone was sufficient to rescue iTreg cell differentiation and

suppression of inflammatory T cells in response to Hh colonization, as seen in mice reconstituted

with *MHCII^{ΔCD11c}* plus *MHCII-ON^{RORγt}* bone marrow cells (Fig. 5c,d), or with only *MHCII-ON^{RORγt}*

cells (Extended Data Fig. 12c). There was similar rescue of endogenous iTreg cell differentiation

in mice having the gain-of-function MHCII in RORγt⁺ cells (Extended Data Fig. 12b). It should be

noted that whereas there was rescue of Hh-directed iTreg cell differentiation in mice reconstituted

with *MHCII-ON^{RORγt}* bone marrow, there was a marked absence of Bcl6-expressing Tfh cells,

which are also induced by Hh (Fig. 5c,d; Extended Data Fig. 12c). Interestingly, Tfh cells were

present in mice reconstituted with only *MHCII-ON^{CD11c}* bone marrow, suggesting that their

differentiation requires antigen presentation by DC and/or B cells, as proposed previously[35]

(Extended Data Fig. 12c). We conclude that *RORγt$^+$* APC, ILC3 and/or JC, are specialized to

prime naive microbiota-specific T cells and guide their differentiation into iTregs, but other APCs

are required to guide the differentiation of microbiota-specific pathogenic Th17 cells and Tfh cells.

Discussion

The composition of the intestinal microbiota influences host immune functions that contribute to

anti-microbial host defense, inflammatory disease, and anti-tumor immunity. Transmission of

information from gut microbes to immune system cells remains poorly understood. The current

results indicate that RORγt$^+$ cells, either JC, whose transcriptional profile suggests a role in

promoting immunological tolerance[5], or type 3 innate lymphoid cells previously implicated in

restraining microbiota-dependent Th1/Th17 inflammatory responses in the gut[36,37], do so in large

part by conveying signals from the microbiota to naive bacteria-specific T cells, activating them

and guiding their differentiation towards a unique iTreg cell program. RORγt$^+$ cells defective for

CCR7-mediated migration (either to or within the MLN), MHCII antigen presentation, or $\alpha v\beta 8$

function (presumably through activation of TGF-β), failed to induce the iTreg program. Intriguingly,

under such circumstances there was expansion of pathogenic Th17 cells that promoted intestinal

inflammation. We previously demonstrated that a T cell-intrinsic c-Maf deficiency prevented iTreg

cell differentiation, and similarly allowed for microbiota-dependent differentiation of pathogenic

Th17 cells[3]. Together, these findings suggest that iTreg cells restrain the priming, proliferation,

and differentiation of Th17 cells in the MLN. The APC(s) that directs pathogenic Th17 cell

differentiation does not require CCR7, and it is not targeted in CD11c-Cre mice, and is, therefore,

most likely not a cDC, but its identity is currently not known. One hypothesis is that iTreg cells

might not only inhibit Th17 cell differentiation and function but also may inhibit the function of the

CCR7-independent Th17 inducer APC (Extended Data Fig. 13).

It has been proposed that intrinsically different APC subsets direct distinct T cell responses[38], but

such processes have been difficult to demonstrate in the setting of immune responses *in vivo*.

Our study shows that a unique RORγt$^+$ cell type instructs naïve microbiota-specific CD4$^+$ T cells

to become iTreg cells, but does not support the differentiation of other T cell programs, including

Tfh cells, that are normally also induced by Hh intestinal colonization. Although our results are

most compatible with the RORγt$^+$ APC being an ILC3 or Aire$^+$ JC subset, we cannot rule out that

it may represent a novel RORγt$^+$ cell type that cannot yet be categorized as either lymphoid or

DC-like. Definitive identification and characterization of this cell awaits more specific genetic tools

than those currently available. Nevertheless, our results clearly demonstrate the existence of

multiple APCs that are targeted by a specific commensal microbe to instruct diverse effector T

cell functions (Extended Data Fig. 13). The APCs may act hierarchically, as exemplified here by

RORγt$^+$ cells that supersede the function of Th17-inducing APC. The existence and identity of

distinct cellular circuits responsible for the induction of iTregs and other T cell functional subsets

offers the opportunity to investigate the corresponding cells in humans and, potentially, to

modulate them therapeutically.

Acknowledgements

We thank members of the Littman lab, Juan J. Lafaille, and Susan Schwab for valuable discussion

and critical reading of the manuscript and Gabriela Romero-Meza for assistance with

experiments. We thank Dean Sheppard for advice and providing ADWA11 blocking Ab, S.Y. Kim

and the NYU Rodent Genetic Engineering Laboratory (RGEL) for rederivation of mutant mice,

and Cindy Loomis and the Experimental Pathology Research Laboratory of NYULMC for histology

of intestine samples. The Microscopy Core and the Genome Technology Core are partially

supported by NYU Cancer Center Support Grant NIH/NCI P30CA016087 at the Laura and Isaac

Perlmutter Cancer Center, S10 RR023704-01A1 and NIH S10 ODO019974-01A1. The

Experimental Pathology Research Laboratory is supported by National Institutes of Health Shared

Instrumentation grants S10OD010584-01A1 and S10OD018338-01. Caleb A. Lareau, Ansuman

T. Satpathy and James M. Gardner are recipients of the IGVF award UM1HG012076. This work

was supported by an Irvington Institute fellowship from the Cancer Research Institute (R.K.) and

a Jane Coffin Childs Fund fellowship (K.R.M.), the Helen and Martin Kimmel Center for Biology

and Medicine (D.R.L.); National Institutes of Health grants R01AI139540 (P.M.A.) and

R01AI158687 (D.R.L.), and the Howard Hughes Medical Institute (D.R.L.).

Author Contributions

R.K., T.N., K.R.M. and D.R.L. designed the study and analyzed the data; R.K. and T.N. performed

mouse genetic experiments with assistance from A.G; M.P., M.X., and J.T. performed early

experiments to launch the study. Intravital multiphoton microscopy (K.R.M. and R.K.), CITE-seq

studies (R.K, S.H. and M.S.), scRNA-seq (A.T.S., C.A.L), bioinformatics analyses (R.K, L.K., Y.H.,

J.G.). J.W., M.S.A., and J.M.G. provided biological samples, genomics data, and experimental

support. H.P., T.M.L., I.A., J.M.B., and P.M.A. contributed mouse strains and phenotypic analysis

(H.P.). R.K. and D.R.L. wrote the manuscript, with input from the other authors. D.R.L.

supervised the research.

References

References

1   Hooper, L. V., Littman, D. R. & Macpherson, A. J. Interactions between the microbiota

and the immune system. *Science* **336**, 1268-1273, doi:10.1126/science.1223490 (2012).

2   Ansaldo, E., Farley, T. K. & Belkaid, Y. Control of Immunity by the Microbiota. *Annu Rev*

Immunol **39**, 449-479, doi:10.1146/annurev-immunol-093019-112348 (2021).

3   Xu, M. *et al.* c-MAF-dependent regulatory T cells mediate immunological tolerance to a

gut pathobiont. *Nature* **554**, 373-377, doi:10.1038/nature25500 (2018).

4   Chai, J. N. *et al.* Helicobacter species are potent drivers of colonic T cell responses in

homeostasis and inflammation. *Science immunology* **2**, doi:10.1126/sciimmunol.aal5068

(2017).

5   Wang, J., Lareau, C.A., Bautista, J.L., Gupta, A.R., Sandor, K., Germino, J., Yin, Y.,

Arvedson, M.P., Reeder, G.C., Cramer, N.T., Xie, F., Ntranos,V., Satpathy, A.T., Anderson,

M.S., Gardner, J.M. Single-cell multiomics defines tolerogenic extrathymic Aire-

expressing populations with unique homology to thymic epithelium. *Sci. Immunol.* **6**

(2021).

6   Russler-Germain, E. V. *et al.* Gut Helicobacter presentation by multiple dendritic cell

subsets enables context-specific regulatory T cell generation. *eLife* **10**,

doi:10.7554/eLife.54792 (2021).

7   Darrasse-Jèze, G. *et al.* Feedback control of regulatory T cell homeostasis by dendritic

cells in vivo. *The Journal of experimental medicine* **206**, 1853-1862,

doi:10.1084/jem.20090746 (2009).

8   Durai, V. & Murphy, K. M. Functions of Murine Dendritic Cells. *Immunity* **45**, 719-736,

doi:10.1016/j.immuni.2016.10.010 (2016).

9   Esterházy, D. *et al.* Classical dendritic cells are required for dietary antigen-mediated

induction of peripheral T(reg) cells and tolerance. *Nature immunology* **17**, 545-555,

doi:10.1038/ni.3408 (2016).

10   Nussenzweig, M. C., Steinman, R. M., Gutchinov, B. & Cohn, Z. A. Dendritic cells are

accessory cells for the development of anti-trinitrophenyl cytotoxic T lymphocytes. *The*

Journal of experimental medicine **152**, 1070-1084, doi:10.1084/jem.152.4.1070 (1980).

11   Nutsch, K. *et al.* Rapid and Efficient Generation of Regulatory T Cells to Commensal

Antigens in the Periphery. *Cell Rep* **17**, 206-220, doi:10.1016/j.celrep.2016.08.092

(2016).

12   Esterhazy, D. *et al.* Compartmentalized gut lymph node drainage dictates adaptive

immune responses. *Nature* **569**, 126-130, doi:10.1038/s41586-019-1125-3 (2019).

13   Worbs, T. *et al.* Oral tolerance originates in the intestinal immune system and relies on

antigen carriage by dendritic cells. *The Journal of experimental medicine* **203**, 519-527,

doi:10.1084/jem.20052016 (2006).

14   Koscso, B. *et al.* Gut-resident CX3CR1(hi) macrophages induce tertiary lymphoid

structures and IgA response in situ. *Science immunology* **5**,

doi:10.1126/sciimmunol.aax0062 (2020).

15   Mildner, A. & Jung, S. Development and function of dendritic cell subsets. *Immunity* **40**,

642-656, doi:10.1016/j.immuni.2014.04.016 (2014).

16   Anderson, D. A., 3rd, Dutertre, C. A., Ginhoux, F. & Murphy, K. M. Genetic models of

human and mouse dendritic cell development and function. *Nat Rev Immunol* **21**, 101-

115, doi:10.1038/s41577-020-00413-x (2021).

17   Coombes, J. L. *et al.* A functionally specialized population of mucosal CD103+ DCs

induces Foxp3+ regulatory T cells via a TGF-beta and retinoic acid-dependent

mechanism. *The Journal of experimental medicine* **204**, 1757-1764,

doi:10.1084/jem.20070590 (2007).

18   Persson, E. K. *et al.* IRF4 transcription-factor-dependent CD103(+)CD11b(+) dendritic

cells drive mucosal T helper 17 cell differentiation. *Immunity* **38**, 958-969,

doi:10.1016/j.immuni.2013.03.009 (2013).

19   Pool, L., Rivollier, A. & Agace, W. W. Deletion of IRF4 in Dendritic Cells Leads to Delayed

Onset of T Cell-Dependent Colitis. *J Immunol* **204**, 1047-1055,

doi:10.4049/jimmunol.1900775 (2020).

20   Wohn, C. *et al.* Absence of MHC class II on cDC1 dendritic cells triggers fatal

autoimmunity to a cross-presented self-antigen. *Science immunology* **5**,

doi:10.1126/sciimmunol.aba1896 (2020).

21   Yamano, T. *et al.* Aire-expressing ILC3-like cells in the lymph node display potent APC

features. *The Journal of experimental medicine* **216**, 1027-1037,

doi:10.1084/jem.20181430 (2019).

22   Hepworth, M. R. *et al.* Immune tolerance. Group 3 innate lymphoid cells mediate

intestinal selection of commensal bacteria-specific CD4(+) T cells. *Science* **348**, 1031-

1035, doi:10.1126/science.aaa4812 (2015).

23   Bartleson, J. M. *et al.* Strength of tonic T cell receptor signaling instructs T follicular

helper cell-fate decisions. *Nature immunology* **21**, 1384-1396, doi:10.1038/s41590-020-

0781-7 (2020).

24   Mackley, E. C. *et al.* CCR7-dependent trafficking of RORgamma(+) ILCs creates a unique

microenvironment within mucosal draining lymph nodes. *Nat Commun* **6**, 5862,

doi:10.1038/ncomms6862 (2015).

25   Kim, M. H., Taparowsky, E. J. & Kim, C. H. Retinoic Acid Differentially Regulates the

Migration of Innate Lymphoid Cell Subsets to the Gut. *Immunity* **43**, 107-119,

doi:10.1016/j.immuni.2015.06.009 (2015).

26   Wang, R. *et al.* GARP regulates the bioavailability and activation of TGFbeta. *Mol Biol*

Cell **23**, 1129-1139, doi:10.1091/mbc.E11-12-1018 (2012).

27   Lienart, S. *et al.* Structural basis of latent TGF-beta1 presentation and activation by

GARP on human regulatory T cells. *Science* **362**, 952-956, doi:10.1126/science.aau2909

(2018).

28   Qin, Y. *et al.* A Milieu Molecule for TGF-beta Required for Microglia Function in the

Nervous System. *Cell* **174**, 156-171 e116, doi:10.1016/j.cell.2018.05.027 (2018).

29   Lacy-Hulbert, A. *et al.* Ulcerative colitis and autoimmunity induced by loss of myeloid

alphav integrins. *Proc Natl Acad Sci U S A* **104**, 15823-15828,

doi:10.1073/pnas.0707421104 (2007).

30   Paidassi, H. *et al.* Preferential expression of integrin alphavbeta8 promotes generation

of regulatory T cells by mouse CD103+ dendritic cells. *Gastroenterology* **141**, 1813-1820,

doi:10.1053/j.gastro.2011.06.076 (2011).

31   Travis, M. A. *et al.* Loss of integrin alpha(v)beta8 on dendritic cells causes autoimmunity

and colitis in mice. *Nature* **449**, 361-365, doi:10.1038/nature06110 (2007).

32   Gardner, J. M. *et al.* Deletional tolerance mediated by extrathymic Aire-expressing cells.

Science **321**, 843-847, doi:10.1126/science.1159407 (2008).

33   Nakawesi, J. *et al.* alphavbeta8 integrin-expression by BATF3-dependent dendritic cells

facilitates early IgA responses to Rotavirus. *Mucosal Immunol* **14**, 53-67,

doi:10.1038/s41385-020-0276-8 (2021).

34   Brown, C. C. *et al.* Transcriptional Basis of Mouse and Human Dendritic Cell

Heterogeneity. *Cell* **179**, 846-863 e824, doi:10.1016/j.cell.2019.09.035 (2019).

35   Barnett, L. G. *et al.* B cell antigen presentation in the initiation of follicular helper T cell

and germinal center differentiation. *J Immunol* **192**, 3607-3617,

doi:10.4049/jimmunol.1301284 (2014).

36   Hepworth, M. R. *et al.* Innate lymphoid cells regulate CD4+ T-cell responses to intestinal

commensal bacteria. *Nature* **498**, 113-117, doi:10.1038/nature12240 (2013).

37   Goto, Y. *et al.* Segmented filamentous bacteria antigens presented by intestinal

dendritic cells drive mucosal Th17 cell differentiation. *Immunity* **40**, 594-607,

doi:10.1016/j.immuni.2014.03.005 (2014).

38   Yin, X., Chen, S. & Eisenbarth, S. C. Dendritic Cell Regulation of T Helper Cells. *Annu Rev*

Immunol **39**, 759-790, doi:10.1146/annurev-immunol-101819-025146 (2021).

Figure 1. Distinct requirements for antigen presentation and CCR7 expression in differentiation of iTreg versus pathogenic Th17 cells.

a-b, Hh-specific T cell proliferation and differentiation in Hh-colonized *CD11c-Cre;I-Ab^{f/f}* (*MHCII^{ΔCD11c}*) (n=3) and *I-Ab^{f/f}* or *CD11c-Cre;I-Ab^{f/+}* littermate control mice (n=6) (a) and in *Ccr7^{ΔCD11c}* (n = 7) and littermate control mice (n = 9) (b). CFSE-labeled naïve TCR transgenic Hh7-2 T cells from the C1 MLN were assessed for cell proliferation and expression of FoxP3 and RORγt at 3 days after adoptive transfer. Representative flow cytometry (left) and aggregate results (right). Data summarize two (a) and three (b) independent experiments. **c,** Representative H&E histology in large intestine of mice with indicated genotypes. Scale bars are 1 mm and 50μm, for 1.25X and 20X. respectively. All statistics were calculated by unpaired two-sided Welch's t-test. Error bars denote mean ± s.d. *p*-values are indicated in the figure.

Figure 2. Antigen presentation by RORγt⁺ cells is required for microbiota-induced iTreg cell differentiation.

a-b, UMAP visualization of the *tdTomato-ON^CD11c* fate-map cell CITE-seq dataset, analyzed by the WNN method (a), and Violin plot showing MHCII protein levels in the different cell clusters (b). MLN cells from Hh-colonized *tdTomato-ON^CD11c* fate-map mice were gated for TCRβ⁻, TCRγδ⁻, B220⁻, and tdTomato⁺ cells were sorted for CITE-seq analysis. Cells were sorted from two mice and labeled by hashing antibodies (n=2). **c,** CD11c and CD11b staining of ILC3 and JC from MLN of Hh-colonized *RORγt-eGFP* mice, gated as indicated. **d,** MHCII expression in RORγt⁺ cells (top) and DCs (bottom) from the MLN of Hh-colonized mice of the indicated genotypes. RORγt⁺ cells were gated as TCRβ⁻, TCRγδ⁻, B220⁻, RORγt⁺; DC were gated as TCRβ⁻, TCRγδ⁻, B220⁻, CD90⁻, CD11c⁺, CD11b⁺ SIRPα⁺. **e,** Hh7-2 cell proliferation and differentiation in *MHCII^ΔRORγt* (n = 6) and *I-Ab^f/f* littermate control mice (n = 6) at 3 days after adoptive transfer into Hh-colonized mice. **f,** Hh7-2 T cell differentiation profiles (upper) and cytokine production (lower) in the large intestine lamina propria at 22 days after transfer into *MHCII^ΔRORγt* (n=11) and littermate controls (n=9). Differentiation was assessed by expression of

473   Foxp3, RORγt with or without T-bet, and T-bet. Data summarize two independent experiments.

474   All statistics were calculated by unpaired two-sided Welch's t-test. Error bars denote mean ± s.d.

475   *p*-values are indicated on the figure.

Figure 3. RORγt⁺ cells require CCR7 to promote iTreg cell differentiation. a-b,

Representative flow cytometry profiles (left) and aggregate data (right) for Hh7-2 T cell

proliferation and differentiation in MLN at 3 days (a) and their phenotype in large intestine at 14

days (b) following adoptive transfer into *Ccr7^{ΔRORγt}* and littermate control mice. MLN: Control

mice n=10, *Ccr7^{ΔRORγt}* mice n=20; LI: Control mice n=10, *Ccr7^{ΔRORγt}* mice n=9. Data summarize

three independent experiments. Statistics were calculated by unpaired two-sided Welch's t-test.

c, Analysis of CCR7 requirement for RORγt⁺ cell accumulation in the MLN. Irradiated CD45.1

mice were reconstituted with equal number of bone marrow cells from CD45.2 *Ccr7^{ΔRORγt}* and

CD45.1/CD45.2 WT mice or with CD45.2 WT and CD45.1/CD45.2 WT mice as controls.

Scheme is shown at the top (created with BioRender.com). Aggregate data shows the

frequency in MLN and colon lamina propria of CD45.2 WT (n=4) or CD45.2 *Ccr7^{ΔRORγt}* (n=9)

cells within each subset, as indicated. Data summarize three independent experiments. All

491   statistics were calculated by unpaired two-sided Welch's t-test. Error bars denote mean ± s.d. *p*-

492   values are indicated in the figure.

Figure 4. Role of integrin $\alpha_v\beta_8$ in RORγt$^+$ antigen presenting cell-dependent iTreg cell

differentiation. a, Frequency of iTreg cells among proliferating donor-derived Hh-specific cells

in the MLN at 3 days after transfer of naïve CFSE-labeled Hh7-2 T cells into mice treated with

200μg of ADWA11 blocking antibody (n=8) or left untreated (n=8), on the day of adoptive

transfer. Data summarize three independent experiments. **b,** Hh7-2 T cell proliferation and

differentiation in the MLN of *Itgav$^{\Delta CD11c}$* (n=5) and littermate controls (n=15) at 3 days after

adoptive transfer. Data summarize three independent experiments. **c,** Proliferation and

differentiation of Hh-specific iTreg cells in the MLN of *Itgav$^{\Delta ROR\gamma t}$* (n=6) and littermate control

mice (n=8). CFSE-labeled Hh7-2 T cells were analyzed at 3 days following their adoptive

transfer into Hh-colonized mice. Representative flow cytometry profiles (left) and aggregate data

(right). Data summarize three independent experiments. **d,** Transcription factor expression in

Hh7-2 T cells (left panels) and in endogenous CD4$^+$ T cells (right panels) from colon lamina

propria (LP) at 10 days after adoptive transfer into *Itgav$^{\Delta ROR\gamma t}$* mice (n=3) and control littermates

(n=6). Data summarize two independent experiments. Representative dot plots and aggregate

data are shown (right panels). **e,** Scheme for mixed bone marrow chimeric mouse experiment,

with control, *Itgav*$^{\Delta ROR\gamma t}$ or *MHCII*$^{\Delta CD11c}$ cells administered to irradiated host mice (created with

BioRender.com). **f,** Bar graphs showing iTreg frequency among Hh7-2 T cells in the colon LP at

10 days after their transfer into the bone marrow chimeric mice, reconstituted with different

combinations of donor cells as indicated. Control mice (n=3), *MHCII*$^{\Delta CD11c}$ (n=3), *MHCII*$^{\Delta CD11c}$

and WT (n=4), *Itgav*$^{\Delta ROR\gamma t}$ and WT (n=4) and *MHCII*$^{\Delta CD11c}$ and *Itgav*$^{\Delta ROR\gamma t}$ (n=4). All statistics

were calculated by unpaired two-sided Welch's t-test. Error bars denote mean ± s.d. *p*-values

are indicated in the figure.

Figure 5. Antigen presentation by RORγt⁺ cells is sufficient to promote iTreg cell

differentiation. a, Experimental design (created with BioRender.com). **b,** MHCII frequency in

donor bone marrow-derived cDC2 (gated TCRβ⁻, TCRγδ, B220⁻, CD45.2⁺, CD11c⁺, CD11b⁺

Sirpa⁺) and RORγt⁺ cells (gated as TCRβ⁻, TCRγδ⁻, B220⁻, RORγt⁺, CD45.2⁺) in MLN from

chimeric mice reconstituted with combinations of donor BM cells as indicated. **c,** Representative

flow cytometry of Hh7-2 T cell differentiation in colon lamina propria of Hh-colonized bone

marrow chimeric mice, 12 days after transfer of naive TCR transgenic T cells. **d,** Aggregate data

for differentiation of Hh7-2 T cells in bone marrow chimeric mice reconstituted with cells of

indicated genotypes. WT (n=10), MHCII^{ΔCD11c} (n=9), MHCII^{ΔCD11c} and WT (n=8), MHCII^{ΔCD11c} and

MHCII-ON^{ΔRORγt} (n=9). Data summarize three independent experiments.   All statistics were

calculated by unpaired two-sided Welch's t-test. Error bars denote mean ± s.d. *p*-values are

indicated in the figure.

Extended Data Fig. 1. Cells targeted by CD11c-Cre and consequences for Hh-specific T cell differentiation. a, Phenotype of Hh7-2 TCR transgenic T cells in the colon lamina propria at 10 days after transfer into Hh-colonized $MHCII^{\Delta CD11c}$ (n=3) and control mice (n=7), as indicated. **b,** Phenotype of host CD4$^+$ T cells from mice in (a); $MHCII^{\Delta CD11c}$ (n=3) and control mice (n=10), as indicated. **c,** Cytokine profile of Hh7-2 T cells shown in (a); $MHCII^{\Delta CD11c}$ (n=4) and control mice (n=3). **d,** Proliferation and differentiation of Hh-specific iTreg and Th17 cells in the MLN of $Ccr7^{-/-}$ (n=5) and littermate control mice (n=5). CFSE-labeled Hh7-2 T cells were analyzed at 3 days following their adoptive transfer into Hh-colonized mice. Data summarize two independent experiments. **e-f,** Transcription factor (e) and intracellular cytokine (f) profiles of Hh7-2 T cells in the large intestine of $Ccr7^{\Delta CD11c}$ (n=7 or 5, for transcription factors and cytokines, respectively) and littermate control (n=5) mice, at 10 days after adoptive transfer. Data summarize two independent experiments. Representative flow panels and aggregate data are shown for each analysis. All statistics were calculated by unpaired two-sided Welch's t-test. Error bars denote mean ± s.d. *p*-values are indicated in the figure.

Extended Data Fig. 2. cDC1 or cDC2 are not required for iTreg differentiation. a-c,

Proportion in MLN of Hh7-2 with the iTreg phenotype at 3 days after transfer into $BATF3^{-/-}$ (a)

(n=7), $IRF4^{\Delta CD11c}$ (b) (n=6), and *huLangerin (CD207)-DTA* (c) (n=12) mice (red) and indicated

littermate controls (black). Data summarize at least two independent experiments. All statistics

were calculated by unpaired two-sided Welch's t-test. Error bars denote mean ± s.d. *p*-values

are indicated in the figure.

Extended Data Fig. 3. Identification of CITE-seq-assigned clusters of sorted *tdTomato-*

***ON*$^{\Delta CD11c}$ fate-mapped cells**. **a,** Stacked violin plots for selected (curated) and top DEG (data-

driven) of tdTomato$^+$ cells sorted from MLN of Hh-colonized mice. **b,** Ridge Plot of cell surface

markers for each cluster.

Extended Data Fig. 4. Phenotypic discrimination of ILC3 and JC. **a,** Dot plots for selected

(curated) DEG and cell surface markers for the indicated clusters, obtained from CITE-seq

analysis of *tdTomato-ON$^{\Delta CD11c}$* fate-mapped cells. **b,** Flow cytometry profiling of CXCR6,

CD127(IL-7R), CCR6 CD25 and CD40 on ILC3 (red) and JC (blue), pre-gated on TCRβ⁻,

TCRγδ⁻, B220⁻, RORγt⁺, MHCII⁺. **c,** gating strategy for JC using cell surface staining as

indicated. **d,** Flow cytometry profiling of JC and DC markers, showing that migratory cDC are

excluded from CD11c^{low} CCR6⁺ gating. **e,** TdTomato levels in ILC3 (TCRβ⁻, TCRγδ⁻, B220⁻,

MHCII⁺ CCR6⁺, Il7R⁺) and JC (TCRβ⁻, TCRγδ⁻, B220⁻, MHCII⁺ CCR6⁺, Il7R⁻) from the MLN of

Hh-colonized *tdTomato-ON^{CD11c}* fate-map mice.

Extended Data Fig. 5. Antigen presentation by RORγt⁺ cells is required for microbiota-induced iTreg cell differentiation. a, MHCII expression in RORγt⁺ cells (top) and DC (bottom) from the MLN of Hh-colonized *MHCII^ΔCD11c* mice (n=6 and 5) and littermate controls (n =10 and 8). RORγt⁺ cells were gated as TCRβ⁻, TCRγδ⁻, B220⁻, RORγt⁺; DC were gated as TCRβ⁻, TCRγδ⁻, B220⁻, CD90⁻, CD11c^high. **b,** Bar graph showing frequency of iTreg among Hh7-2 T cells, measured as in Figure 2e. **c-d,** Representative dot plots showing Hh7-2 T cell differentiation (c) and cytokine (d) profiles in colon lamina propria at 22 days after adoptive transfer into Hh-colonized *MHCII^ΔRORγt* and littermate controls. **e,** Representative and aggregate data of transcription factor profiles of host CD4⁺ T cells in colon lamina propria of mice shown in (c) and (d). **f,** Hh7-2 cell proliferation and differentiation in the MLN of *H2-DMa^ΔRORγt* (*RORγt-Cre; H2-Dma^f/f*) (n=11) and littermate controls (*RORγt-Cre; H2-DMa^+/f*) (n=5) at 3 days after transfer of CFSE labeled naïve Hh7-2, cell proliferation and FoxP3 were assessed in cells isolated from C1 MLN. Representative flow cytometry (left) and aggregate data from multiple animals (right).

Data summarize two independent experiments. All statistics were calculated by unpaired two-

sided Welch's t-test. Error bars denote mean ± s.d. p-values are indicated in the figure.

Extended Data Fig. 6. Differential requirements for CCR7 in iTreg and effector Th17 cell

differentiation. a, Cell surface expression of CCR7 on *CD11c-Cre* fate-mapped ILC3 (TCRβ⁻,

TCRγδ⁻, B220⁻, MHCII⁺, CCR6⁺, IL-7R⁺) and JC (TCRβ⁻, TCRγδ⁻, B220⁻, MHCII+, CCR6+, IL-7R⁻)

in the MLN. **b-c,** Analysis of DC counts in MLN (b) and large intestine (c) of WT and *Ccr7^{ΔRORγt}*

mixed bone marrow chimeric mice described in Figure 3c. Counts in the MLN of DC subsets

derived from bone marrow (b); frequencies of CCR7⁺ among total colonic DCs (c) (n=4).

Statistics were calculated using paired two-sided t-test. **d,** Analysis of CD45.2 frequencies within

donor cells is presented for ILC3 (TCRβ⁻, TCRγδ⁻, B220⁻, MHCII⁺, RORγt ⁺, IL-7R⁺) and JC

(TCRβ⁻, TCRγδ⁻, B220⁻, MHCII⁺, RORγt ⁺, IL-7R⁻) in the MLN of WT and *Ccr7^{ΔRORγt}* mixed bone

marrow chimeric mice described in Figure 3c. **e,** Cell surface expression of CCR7 in colonic

ILC3 (TCRβ⁻, TCRγδ⁻, B220⁻, CD90⁺, RORγt⁺, CD25⁺, CD4⁺) from *Ccr7^{ΔRORγt}* (n=3), *Ccr7^{ΔCD11c}*

(n=2) and control Hh-colonized mice (n=2). **f,** Cell surface expression of CCR7 and CD11b in

ILC3-gated MLN cells (TCRβ⁻, TCRγδ⁻, B220⁻, IL-7R⁺, CCR6⁺, CD25⁺) from *Ccr7^{ΔCD11c}* (n=4) and

control Hh-colonized mice (n=4).

Extended Data Fig. 7. Analysis of Aire$^+$ JC function in differentiation of Hh-specific iTreg

cells. a, Lethally irradiated mice were reconstituted with BM cells from CD45.2 *Aire-DTR* or

CD45.2 WT mice. One month after reconstitution, mice were colonized with Hh, and one week

later were treated with Diphtheria toxin (DT, Sigma-Aldrich) for 3 sequential days (at a dose of

25 ng/g mice). CD45.1/CD45.2 CFSE-labeled Hh7-2 T cells (1 x 10^5) were transferred

intravenously into the mice on the first day of DT treatment. Bar graph of proportion of

proliferating Foxp3$^+$ Hh7-2 T cells in the MLN of mice reconstituted with Aire-DTR BM (n=5) or

with WT BM (n=4) (left); *Aire* mRNA in the spleen of the treated mice was blindly scored using

RNAscope analysis. **b,** Proliferation and differentiation of CFSE-labeled Hh7-2 T cells in the

MLN of *RORγt-Cre;Aire$^{f/f}$* (n=5) and control *Aire$^{+/f}$* littermates (n=7) at 3 days after adoptive

transfer. Data summarize three independent experiments. All statistics, except for b and c, were

calculated by unpaired two-sided Welch's t-test. Error bars denote mean ± s.d. *p*-values are

indicated in the figure.

Extended Data Fig. 8. Effect of integrin $\alpha_v\beta_8$ blockade or α_v inactivation on microbiota-

dependent T cell differentiation. a, Expression of integrin α_v (CD51) in fate-mapped $ROR\gamma t^+$

cell subsets from MLN of wild type and *Itgav$^{\Delta ROR\gamma t}$* mice. **b,** C1 MLN from *Itgav$^{\Delta CD11c}$* (n=3) and

littermate controls (n=4), 10 days after Hh colonization. **c-d,** Flow cytometry profiling of

transcription factors and CCR6 in proliferating CFSE-labeled Hh7-2 in the MLN at 3 days after

adoptive transfer into *Itgav$^{\Delta ROR\gamma t}$* (n=3) and littermate control mice (n=3) (b) or into mice treated

with ADWA11(n=4) (as in Fig. 4a) or untreated control littermates (n=4) (c). Summary data of

results in (b) and (c) are shown below. **e,** Intracellular IFNγ and T-bet expression in

PMA/Ionomycin-stimulated Hh7-2 T cells isolated from colon lamina propria of *Itgav$^{\Delta ROR\gamma t}$*(n=3)

and control littermates (n=5), 10 days after adoptive transfer. **f,** Frequency of iTreg cells among

proliferating Hh7-2 in the MLN at 3 days after adoptive transfer into *Itgav$^{\Delta CD4}$* (n=4) and control

littermates (n=7). Data summarize two independent experiments. **g,** Integrin α_v and MHCII cell

surface expression in ILC3 (gated as $TCR\beta^-$, $TCR\gamma\delta^-$, $B220^-$, $ROR\gamma t^+$, $CD90^+$, $CD25^+$ $CD45.2^+$)

isolated from MLN of bone marrow chimeric mice, reconstituted with different combinations of

donor cells as indicated and colonized with Hh for 10 days. Data summarized below for control

(n=3), $MHCII^{\Delta CD11c}$ (n=3), $MHCII^{\Delta CD11c}$ and WT (n=4), and $MHCII^{\Delta CD11c}$ and $Itgav^{\Delta ROR\gamma t}$ (n=4)

reconstituted mice. All Statistics were calculated by unpaired two-sided Welch's t-test. Error

bars denote mean ± s.d. p-values are indicated in the figure.

Extended Data Fig. 9. *Itgav and Itgb8* **expression in ILC3 and JC. a**, tSNE plot with Leiden

clustering of scRNAseq of pooled GFP[+] sorted and unsorted cells, as indicated, from pooled

lymph nodes of Adig mice[32]. **b,** tSNE feature plots showing *Aire, Itgav, and Itgb8* levels in the

cell clusters. **c,** UMAP plot of Aire+ JC and ILC3 populations from pooled datasets as indicated

with associated feature plots. **d,** top differentially expressed genes per pseudobulk cluster in (c),

shown by heatmap. **e,** dot plot of selected genes in JC and ILC3 clusters. **f,** Representative flow

cytometry (left) and aggregate results (right) of tdTomato[+] JC and ILC3, gated as indicated, in

C1 MLN of *tdTomato^ΔItgb8* mice[33] (n=4) and littermate controls (n=4). Aggregate data (right) show

percent tdTomato+ cells among total ILC3 and JC and number of reporter-positive cells in the

C1 MLN of each mouse. All statistics were calculated by unpaired two-sided Welch's t-test.

Error bars denote mean ± s.d. *p*-values are indicated in the figure.

Extended Data Fig. 10. Analysis of RORγt-expressing cells in the MLN. a, Hh7-2

proliferation and differentiation in MLN of *Ccr7^Δzbtb46^* (n= 4) and control littermates (n=5), at 3

days after transfer of the naïve cells. Data in the right panel summarize three independent

experiments. **b,** UMAP visualization of CITE-seq datasets obtained from 3 distinct sorted

populations (GFP⁺, GFP⁺ tdTomato⁺ and tdTomato⁺) isolated from C1 MLN of *Zbtb46-eGFP ;*

*tdTomato-ON^ΔRORγt^* mice (n=2), analyzed by the WNN method. **c,** Flow cytometry analysis of

fate-mapped C1 MLN cells from *RORγt-eGFP;mKate2-ON^Δzbtb46^* mice, gated for the indicated

cell subsets. **d-e,** Feature plot showing *Rorc* (d) and integrin α_v (e) levels in the cell clusters

identified in the CITE-seq analysis shown in (b); Positive cells are layered in front. All statistics

were calculated by unpaired two-sided Welch's t-test. Error bars denote mean ± s.d. *p*-values

are indicated in the figure.

Extended Data Fig. 11. Intravital tracking of RORγt-expressing cells and DC interactions

with Hh-specific T cells during priming in the MLN. a, Flow cytometry analysis of fate-

mapped C1 MLN cells from *RORγt-eGFP;mKate2-ON$^{\Delta zbtb46}$* mice, gated for the indicated cell

subsets (ILC3 were gated as TCRβ⁻, TCRγδ⁻, B220⁻, MHCII⁺, RORγt-eGFP⁺, CCR6⁺, CD25⁺ and

JC as TCRβ⁻, TCRγδ⁻, B220⁻, MHCII⁺, RORγt-eGFP⁺, CD25⁻). Note that there is incomplete

excision of the transcriptional stop signal by *zbtb46-Cre*. **b,** Representative image of cell-cell

interactions of recently primed Hh-specific T cells with DC and RORγt-expressing cells. *Nur77-*

eGFP tracer-labeled Hh7-2 T cells were transferred into of *RORγt-eGFP;mKate2-ON^{zbtb46}* Hh-

colonized mice. Cell colocalization of primed Hh7-2 (tracer dye⁺, GFP⁺) or naïve Hh7-2 (tracer

dye⁺, GFP⁻) T cells with cDC (mKate2⁺ with dendritic morphology), RORγt-expressing cells

(eGFP⁺, mKate2⁺ or eGFP⁺ alone with amoeboid morphology), or both were visualized using

intravital multiphoton microscopy of the C1 MLN at 15 h after transfer. Note that Cell Tracer

fluorescent labeling provides clear spatial discrimination of *RORγt-eGFP* and *Nur77-eGFP*

expressing cells. **c,** Quantification and graphical representation of the total and individual rates

of interaction of RORγt-expressing cells or cDC populations with primed or naïve Hh7-2 T

cells. Data summarize cell-cell interactions from six 0.25mm³ three-dimensional regions of C1

MLN, (n=72 total Hh7-2 T cells), (n=49 primed and 23 naïve Hh7-2 T cells). All statistics were

683   calculated by unpaired two-sided Welch's t-test. Error bars denote mean ± s.d. *p*-values are

684   indicated in the figure.

Extended Data Fig. 12. Gain-of-function expression of MHCII in RORγt⁺ cells rescues

bone marrow-derived iTreg cell differentiation. a, Aggregate data showing MHCII frequency

on donor-derived DC and RORγt⁺ cells in MLN and colon lamina propria from chimeric mice

reconstituted with combinations of donor BM cells as indicated, with representative flow

cytometry panel in Fig. 5b. MLN: WT (n=5), *MHCII^ΔCD11c* (n=5), *MHCII^ΔCD11c* and WT (n=8),

MHCII^ΔCD11c and *MHCII-ON^ΔRORγt* (n=7). Colon: WT (n=4), *MHCII^ΔCD11c* (n=6), *MHCII^ΔCD11c* and

WT (n=8), *MHCII^ΔCD11c* and *MHCII-ON^ΔRORγt* (n=6). **b,** Donor bone marrow-derived CD4⁺ T cell

differentiation in colon lamina propria from chimeric mice reconstituted with combinations of BM

cells as indicated. Representative flow panels (left) and aggregate data (right). WT (n=10),

MHCII^ΔCD11c (n=11), *MHCII^ΔCD11c* and WT (n=8), *MHCII^ΔCD11c* and *MHCII-ON^ΔRORγt* (n=9). Colon:

WT (n=4), *MHCII^ΔCD11c* (n=6), *MHCII^ΔCD11c* and WT (n=8), *MHCII^ΔCD11c* and *MHCII-ON^ΔRORγt* (n=7).

c, Representative flow cytometry (left) and aggregate data (right) of Hh7-2 T cell differentiation

in colon lamina propria of Hh-colonized bone marrow chimeric mice reconstituted with cells of

indicated genotypes, 12 days after transfer of naive TCR transgenic T cells. WT (n=8),

701   *MHCII$^{\Delta CD11c}$ (n=7), MHCII-ON$^{\Delta CD11c}$ (n=5), and MHCII-ON$^{\Delta ROR\gamma t}$ (n=4)*. Data summarize two or

702   three independent experiments. All statistics were calculated by unpaired two-sided Welch's t-

703   test. Error bars denote mean ± s.d. *p*-values are indicated in the figure.

Extended Data Fig. 13. Schematic of the requirement of distinct APC subsets for T cell

differentiation. CCR7 and integrin $\alpha_v\beta_8$ are required in RORγt$^+$ APCs for iTreg cell

differentiation. Note that other APCs, with differential requirements for CCR7 expression, are

involved in the priming and differentiation of pathogenic Th17 and Tfh cells.

Methods

Mice

Mice were bred and maintained in the Alexandria Center for the Life Sciences animal facility of the New York University School of Medicine, in specific pathogen-free conditions. C57BL/6 mice (Jax# 000664), *Batf3$^{-/-}$ (B6.129S(C)-Batf3tm1Kmm/J* #Jax 013755), *Itgav$^{f/f}$* (*B6.129P2(Cg)-Itgav^{tm2Hyn}/J* Jax# 032297, CD45.1 mice (*B6.SJL-Ptprca Pepcb/BoyJ*, Jax# 002014), *CD4-Cre* (*Tg(Cd4-cre)1Cwi/BfluJ,* Jax# 017336), *CD11c-Cre* (*B6.Cg-Tg(Itgax-cre)1-1Reiz/J* #Jax 008068), *Ccr7$^{-/-}$* (*B6.129P2(C)-Ccr7tm1Rfor/J*, Jax# 006621), *I-AB$^{f/f}$* (*B6.129X1-H2-Ab1tm1Koni/J* #Jax 013181), *Zbtb46-Cre* (*B6.Cg-Zbtb46tm3.1(cre)Mnz/J* #Jax 028538), *Zbtb46-eGFP* (*B6.129S6(C)-Zbtb46tm1.1Kmm/J* #Jax 027618), tdTomato^{LSL} (*B6;129S6-Gt(ROSA)26Sortm14(CAG-tdTomato)Hze/J* #Jax 007908), *Aire$^{f/f}$* (*B6.Cg-Airetm1Dfil/J* #Jax 031409), *Nur77-eGFP* (*C57BL/6-Tg(Nr4a1-EGFP/cre)820Khog/J* #Jax 016617), CD90.1 (*B6.PL-Thy1a/CyJ* #Jax 000406) mice were purchased from Jackson Laboratories. ROR*γ*t-Cre and Hh7-2tg were generated in our laboratory and previously described[3,39]. BAC-transgenic Rorc(t)-GfpTG were generated by G. Eberl's lab[40]. *huLangerin (CD207)-DTA* mice were kindky provided by Daniel H. Kaplan[41]. Ccr7$^{f/f}$ mice were previously described[14]. *mKate2^{LSL}* mice[42] were provided by Scott Lowe. I-AB lox-STOP-lox[35], Aire-DTR[43], H2-DMa1$^{f/f}$ [23], Itgb8-tdTomato[33] Adig (Aire-Driven Igrp-Gfp)[32] mice have been described. Littermates with matched sex (both males and females) were used. Mice in all the experiments were 6–12 weeks old at the starting point of treatment. Animal sample size estimates were determined using power analysis (power=90% and alpha=0.05) based on the mean and standard deviation from our previous studies and/or pilot studies using 4–5 animals per group. All animal procedures were performed in accordance with protocols approved by the Institutional Animal Care and Usage Committee of New York University School of Medicine.

Antibodies, intracellular staining and flow cytometry

The following monoclonal antibodies were purchased from eBiosciences, BD Pharmingen or BioLegend: CD3 (145-2C11), CD4 (RM4-5), CD25 (PC61), CD44 (IM7), CD45.1 (A20), CD45.2 (104), CD90.1 (HIS51), CD90.2 (53-2.1), CD19 (1D3), CD45R (RA3-6B2), CD127 (A7R34), CD51 (RMV-7), IA/IE (56-5321-82), CCR6 (3D6), Ncr1 (29A1.4), NK1.1 (PK136), CD62L (MEL-14), CXCR5 (L138D7), TCRβ (H57-597), TCR Vβ6 (RR4-7), Bcl-6 (K112-91), Foxp3 (FJK-16s), RORγt (B2D or Q31-378), T-bet (eBio4B10), IL-17A (eBio17B7) and IFN-γ (XM61.2), CD11c (N418), CD11b (M1/70), CX3CR1 (SA011F11), Ly6c (HK1.4), SIRPa (P84), Ly6G (1A8), CD273 (TY25), Clec12a (5D3), CD103 (M290), XCR1 (ZET), F4/80 (BM8), CCR7 (4B12), CXCR6 (SA051D1), CD40 (HM40-3), 4′,6-diamidino-2-phenylindole (DAPI) or Live/dead fixable blue (ThermoFisher) was used to exclude dead cells.

For transcription factor staining, cells were stained for surface markers, followed by fixation and permeabilization before nuclear factor staining according to the manufacturer's protocol (Foxp3 staining buffer set from eBioscience). For cytokine analysis, cells were incubated for 5 h in RPMI with 10% FBS, phorbol 12-myristate 13-acetate (PMA) (50 ng/ml; Sigma), ionomycin (500 ng/ml; Sigma) and GolgiStop (BD). Cells were stained for surface markers before fixation and permeabilization, and then subjected to intracellular cytokine staining according to the manufacturer's protocol (Cytofix/Cytoperm buffer set from BD Biosciences). Flow cytometric analysis was performed on an LSR II (BD Biosciences) or Cytek Aurora (Cytek) or an Aria II (BD Biosciences) and analyzed using FlowJo software (Tree Star).

Flow cytometry gating strategy

Hh7-2 gating: FSC, SSC; Live Dead$^-$,singlets, Dump$^-$ (B220,TCRgd, Ly6G), MHCII$^-$, CD4$^+$, TCRβ$^+$, VB6$^+$, CD45.1$^+$ or CD90.1$^+$; **cDC gating**: FSC, SSC; Live Dead$^-$,singlets, Dump$^-$ (B220, TCRβ, TCRgd, Ly6G), CD11c$^+$ and CD11b$^+$, SIRPa$^{low-moderate}$ (remove CD11c$^-$, SIRPA^{high}); **cDC2**

gating (unless mentioned otherwise): FSC, SSC; Live Dead$^-$,singlets, Dump$^-$ (B220,TCRgd,

Ly6G), CD11c$^+$ and CD11b$^+$, SIRPa$^{low-moderate}$ (remove CD11c$^-$, SIRPA^{high}), Clec12a$^-$ SIRPa$^+$;

migratory cDC2 gating: FSC, SSC; Live Dead$^-$,singlets, Dump$^-$ (B220,TCRgd, Ly6G), CD11c$^+$

and CD11b$^+$, SIRPa$^{low-moderate}$ (remove CD11c$^-$, SIRPA^{high}), Clec12a$^-$ SIRPa$^+$, PDL2$^+$; **NCR1$^+$ ILC3**

gating: FSC, SSC; Live Dead$^-$,singlets, Dump$^-$ (B220,TCRgd, Ly6G), TCRβ^- , CD90+, Il7R$^+$,

RORγt $^+$, CCR6$^-$, NCR1$^+$; **JC** (using internal staining): FSC, SSC; Live Dead$^-$,singlets, Dump$^-$

(B220,TCRgd, Ly6G), TCRβ^- , RORγt $^+$, MHCII+ CXCR6$^-$, Il7R$^-$; **JC** (excluding internal staining):

FSC, SSC; Live Dead$^-$,singlets, Dump$^-$ (B220,TCRgd, Ly6G), TCRβ^- , CD11c$^{low-negative}$, MHCII$^+$,

CCR6$^+$,CXCR6$^-$, Il7R$^-$; **ILC3**: FSC, SSC; Live Dead$^-$,singlets, Dump$^-$ (B220,TCRgd, Ly6G),

TCRβ^- , CXCR6$^+$, Il7R$^+$, RORγt $^+$; **LTi-Like ILC3** (using internal staining): FSC, SSC; Live Dead$^-$

,singlets, Dump$^-$ (B220,TCRgd, Ly6G), TCRβ^- , MHCII+, CXCR6$^+$, Il7R$^+$, RORγt $^+$, CCR6$^+$, CD25$^+$;

LTi-Like ILC3 (excluding internal staining): FSC, SSC; Live Dead$^-$,singlets, Dump$^-$ (B220,TCRgd,

Ly6G), TCRβ^- , CD11c$^{low-negative}$, MHCII$^+$, CCR6$^+$,CXCR6$^+$, Il7R$^+$

Isolation of lymphocytes and APCs

After removal of caecal patches, large intestine tissues were sequentially treated with PBS

containing 1 mM DTT at room temperature for 10 min, twice with 5 mM EDTA at 37 °C for 10 min

to remove epithelial cells, and then minced and dissociated in digestion buffer (RPMI containing

collagenase (1 mg ml−1 collagenase D; Roche), DNase I (100 µg ml−1; Sigma), dispase (0.1 U

ml−1; Worthington) and 10% FBS) with constant stirring at 37 °C 55 min. Leukocytes were

collected at the interface of a 40%/80% Percoll gradient (GE Healthcare). Lymph nodes were

mechanically disrupted for lymphocyte isolation. For isolation of myeloid cells and ILC, lymph

nodes were mechanically disrupted with digestion buffer with constant stirring at 37 °C 30 min.

H. hepaticus **culture and oral infection**

H. hepaticus was kindly provided by Dr. James Fox (MIT). Hh was cultured and administrated as

was previously described[3]. Frozen stock aliquots of *H. hepaticus* were stored in Brucella broth

with 20% glycerol and frozen at −80°C. The bacteria were grown on blood agar plates (TSA with

5% sheep blood, Thermo Fisher). Inoculated plates were placed into a hypoxia chamber (Billups-

Rothenberg), and anaerobic gas mixture consisting of 80% nitrogen, 10% hydrogen, and 10%

carbon dioxide (Airgas) was added to create a micro-aerobic atmosphere, in which the oxygen

concentration was 3~5%. The micro-aerobic jars containing bacterial plates were left at 37°C for

4 days before animal inoculation. For oral infection, *H. hepaticus* was resuspended in Brucella

broth by application of a pre-moistened sterile cotton swab applicator tip to the colony surface.

0.3 mL bacterial suspension was administered to each mouse by oral gavage. Mice were

inoculated for a second dose after 3 days.

Adoptive transfer of Hh7-2 TCR transgenic cells

Adoptive transfer of Hh7-2 was done as was previously described[3], with minor modifications.

Recipient mice were colonized with *H. hepaticus* by oral gavage seven days before adoptive

transfer.   Spleens and lymph nodes from donor Hh7-2 TCRtg mice were collected and

mechanically disassociated. Red blood cells were lysed using ACK lysis buffer (Lonza). Naive

Hh7-2 T cells were sorted as $CD4^+TCR\beta^+CD44^{lo}CD62L^{hi}CD25^-V\beta6^+$ (HH7-2tg), on the Aria II (BD

Biosciences). For analysis of early differentiation, cells were additionally labeled with CFSE

(ThermoFisher). Cells were resuspended in PBS on ice and 100K were transferred into congenic

isotype-labelled recipient mice by retro-orbital injection. Cells from MLN were analyzed 3 days

after transfer and cells from colon LP were analyzed 10-14 days after transfer.

CITE-seq

CITE-seq and cell hashing were performed as described[44,45] with minor modifications. Single-cell

suspensions were obtained from digests of C1 MLN of t*dTomato-ON*[CD11c] or *tdTomato-ON*[RORγt] *;*

zbtb46-eGFP mice that had been colonized with *Helicobacter* for 7 days. Cells were sorted on a

BD FACSAriaII using a 100-μm nozzle. Dead cells as well as T cells and B cells were gated out

using DAPI, TCRβ, TCRγδ and B220 antibodies. From t*dTomato-ON*[CD11c] mice, the tdTomato+

population was collected separately from two mice. From *tdTomato-ON*[RORγt] *; zbtb46-eGFP* mice,

we collected 3 populations from two separate mice: GFP⁺, GFP⁺ tdTomato⁺ and tdTomato⁺.

Sorted cells were stained separately with hashing antibodies (Biolegend)[45]. After removal of

excess hashing antibodies, we combined the samples and stained them with CITE-seq

antibodies, conjugated using iEDDA click chemistry to barcode oligos as described before[46]. In

addition, we included some commercially available Totalseq-A antibodies for CD11c, CD90.2,

CD185, CD51 and CD127 (Biolegend). Post-sorting and staining, cells were run through the

standard 10x Chromium (v3) protocol up until cDNA amplification, with the following modification:

For the cDNA PCR step, 0.2uM of ADT additive primer (5'CCTTGGCACCCGAGAATTCC) and

0.2uM of HTO additive primer (5'GTGACTGGAGTTCAGACGTGTGCTC) were added to the

cDNA amplification master mix. Post cDNA amplification, a 0.6X SPRI cleanup was performed to

separate the cDNA fraction (on beads) from the smaller ADT and HTO fractions (in supernatant).

The cDNA fraction was converted into a 3' tag gene expression library according to the 10x

Genomics Single Cell Genomics Protocol (v3). Supernatant from cleanup was kept for ADT and

HTO preparation.

To the supernatant, another 1.4X SPRI was added to bring the total SPRI concentration to 2X.

After washing the beads in 80% ethanol and eluting in water, a second round of 2X SPRI cleanup

was performed to remove any residual primer carryover from the cDNA PCR. Post cleanup, eluate

was taken into ADT PCR amplification (using TruSeq Small RNA RPIx primer

(5'CAAGCAGAAGACGGCATACGAGXXXXXXXXGTGACTGGAGTTCCTTGGCACCCGAGAAT

TCCA) and SI PCR primer

(5'AATGATACGGCGACCACCGAGATCTACACTCTTTCCCTACACGACGCTC)), and into

hashtag amplification (using TruSeq D7xx

(5'CAAGCAGAAGACGGCATACGAGATXXXXXXXXGTGACTGGAGTTCAGACGTGTGC) and

SI PCR primers). Libraries were pooled and sequenced on a 100 cycle Novaseq S1 flowcell, with

the configuration of 30 base pairs for R1, and 92 base pairs for R2. Additional protocol details can

be found for CITE-seq and cell hashing at www.cite-seq.com.

Post sequencing, gene expression count matrices were generated using cellranger version 5.0

using the refdata-gex-mm10-2020-A reference library provided by 10x Genomics, with the

additional sequences of cre, eGFP, and tdTomato. Counts matrices for hashtags and antibodies

were generated using CITE-seq-Count version 1.4.4. Downstream analysis was then performed

in R using Seurat.

Quality control and doublet removal: We initially selected all cells that were detected in our RNA-

seq, cell hashing, and ADT libraries. We removed cells with < 700 detected genes, but also

removed cells which had an aberrantly high number of genes (more than 5,000 genes) and a high

percentage of mitochondria genes (more than 6%). Additionally, we removed the cells which

were attached by the clumps of antibodies and had a too high number of ADT or HTO UMIs (more

than 5,000 ADT UMI and 4,000 HTO UMI). We used our previously described hashing-based

doublet detection strategy[45], implemented in HTODemux, to identify doublets that represent two

or more cells representing different samples. Only Singlet cells were used for the downstream

analysis.

Multimodal analysis: We normalized RNA data using SCTransform[47] and applied the centered-

log ratio (CLR) transformation to normalize ADT data within each cell. We used principal

component analysis (PCA) to reduce dimensionality of both datasets. Then we took both top 20

RNA and protein PCA dimensions as the input of weighted nearest neighbors (WNN) method[48]

to construct the multimodal weighted KNN graph. To cluster our multimodal dataset, we first used

the weighted KNN graph to generate a shared nearest neighbor graph (SNN) and then apply the

graph-based smart local moving (SLM) algorithm (https://doi.org/10.1140/epjb/e2013-40829-0)

on this SNN graph to find clusters with 0.8 resolution. We performed differential expression on all

pairs of clusters for both RNA and protein markers, and merged clusters that did not exhibit clear

evidence of separation. All samples were clustered together and separated later for further

analysis as indicated.

CITE-seq data projection in Flowjo

For gating analysis, scaled and normalized ADT counts together with cluster identity, UMAP and

UWnn coordinates were exported into a csv format that was uploaded in FlowJo.

Single cell RNAseq of Aire+ cells

Four week-old Adig mouse lymph nodes (cervical, brachial, axillary, inguinal, and mesenteric)

(n=3) were pooled in digestion medium consisting of RPMI 1640 with 2% fetal bovine serum (FBS)

(Sigma-Aldrich), deoxyribonuclease (DNase) (100 µg/ml; Roche) and Liberase (50 µg/ml; Roche),

minced and agitated at 37°C for 30 min, and passed through a 70-µm filter. Cells were

resuspended for magnetic column enrichment (Miltenyi LD column depletion with streptavidin

microbeads and biotinylated antibodies against B220, Ter119, TCRbeta, CD3e). Cells were then

either processed directly for 10x single-cell analysis as reference sample or sorted by flow

cytometry for all live GFP+ cells. Cells were sorted into PBS with 0.04% bovine serum albumin

(BSA). Cell viability and counts were evaluated with Vi-CELL XR (Beckman Coulter), and samples

with viability >85% were used for sequencing.

For analysis, sequencing files were aligned to the mm10 mouse reference genome with the 10x

Genomics Cell Ranger (v3.1.0) count method using the default parameters. Raw count files were

then processed in Python (v3.9.7), removing doublets from each sample using Scrublet[49] with a

doublet score threshold of 0.3. Samples were merged and filtered, removing genes with no

counts, and retaining cells with 600 to 5,000 genes and 1,000 to 30,000 counts, leaving a total of

35,797 cells and 22,979 genes in the dataset. The raw counts were used to train a model of gene

expression using scvi-tools (v0.14.6) with each sample as a batch key. This model was used to

generate normalized expression values for all genes scaled to a library size of 100,000 and create

a tSNE representation of the data using Scanpy[50] (v1.8.2) with the default parameters. Leiden

clustering with a resolution of 0.8 gave 29 clusters which were assigned cell types after removing

three low quality clusters. For targeted analysis of JC, the data was subset on the three JC

clusters and a UMAP was created using Scanpy with a minimum distance of 0.5 and spread of 1.

Differential gene expression of JC clusters was performed using scvi-tools[51], filtering on genes

with a bayes factor greater than 3, mean log fold change greater than 0, and proportion of cells

with non-zero expression greater than 0.1. The z-score of the average expression of the top 10

DE genes from each cluster was used for visualization. For low-dimensional embeddings of

feature plots, scVI normalized expression was used, clipping the top and bottom 1% of expression

values in the full dataset to the maximum and minimum of the color scale to prevent outliers from

skewing the visualization. To display expression of key features in dotplots, log-normalized and

scaled counts for each gene were averaged and standardized across cell types to have a value

ranging from 0 to 1.

RNAscope

Fresh-frozen spleen and lymph nodes were sectioned at 8um and then fixed overnight at 4°C in

10% neutral buffered formalin (Thermo Fisher Scientific, Waltham, MA) before proceeding with

an RNAscope RED 2.5 HD Chromogenic Assay kit (Advanced Cell Diagnostics, Newark, CA) for

detection of Aire mRNA. DapB probe was used as negative control and Polr2a probe was used

as positive control. Semi-quantitative scores were determined in a blinded fashion based on the

number of Aire$^+$ cells per section.

Integrin β_8 (ADWA-11) blocking Ab

We injected IP 200ug of ADWA-11 into mice colonized with Hh for 7 days. On the same day we

adoptively transferred 100K CFSE-labeled naive Hh7-2 cells and tested their proliferation and

differentiation in the C1 MLN, 3 days after the transfer.

Generation of bone marrow (BM) chimeric reconstituted mice

To generate chimeric mice, 4-5 week old CD45.1 mice were irradiated twice with 500 rads/mouse

at an interval of 2-5 h (X-RAD 320 X-Ray Irradiator). A day after, bone marrow (BM) mononuclear

cells were isolated from donor mice, as indicated in each experiment, by flushing the femur bones.

Red blood cells were lysed with ACK Lysing Buffer, and lymphocytes were depleted for Thy1.2

using magnetic microbeads (Miltenyi). BM cells were resuspended in PBS and a total 3-4 x 10^6

BM cells were injected intravenously into the irradiated mice. In case of mixed BMC reconstitution,

a ratio of 1:1 was used. Mice were kept for a week on broad spectrum antibiotics (1 mg/mL

sulfamethoxazole and 0.2 mg/ mL trimethoprim), followed by microbiome reconstitution by fecal

gavage. Mice were reconstituted for 1-2 months before Hh colonization. After 7 weeks, peripheral

blood samples were collected and analyzed by FACS 7 to check for reconstitution.

Intravital multiphoton microscopy

Naive Hh7-2 T cells were isolated from *Nur77-eGFP* Hh7-2tg mice, labeled with Cell tracker dye

(eBioscience Cell Proliferation Dye eFluor 450), and transferred into *mKate2-ON[zbtb46];RORγt-*

eGFP mice that had been colonized with Hh for 6 days. Fifteen hours following adoptive transfer

of Hh T cells, mice were euthanized and C1 MLN were immediately isolated and mounted in cold

RPMI with 10% FCS for intravital multiphoton microscopy. Image stacks were acquired with an

Olympus multiphoton FVMPE-RS system equipped with both InSight X3 and Mai Tai Deepsee

(Spectra-Physics) tunable Ti:Sapphire lasers. To acquire serial optical sections, a laser beam

(780 nm for eFluor™ 450 and 940 nm for simultaneous excitation of eGFP and mKate2) was

focused through a water immersion lens (N.A. 1.05; Olympus) and scanned with a field of view of

0.5 mm^2, at 600 Hz. Z-stacks were acquired in 2 mm steps to image a total depth of 150-200 mm

of tissue.

Image analysis

Raw image stacks were imported into Fiji (NIH) for T cell colocalization analysis. Provided images

are presented as a maximal projection of 3–6 mm optical sections. For visualizing individual

labelled cells expressing both the mKate2 and eGFP, the brightness and contrast were adjusted

accordingly to single positive green (eGFP) and red (mKate2) cells. Adoptively transferred Hh7-

2tg T cells were identified via positive labeling with cell proliferation dye eFluor 450. Primed Hh7-

2tg T cells were identified via expression of the *Nur77-eGFP* reporter. Cell identity was scored by

a combination of both fluorescent reporter expression as well as cell morphology. Specifically,

cells expressing mKate2 with a dendritic cell shape were scored as cDC, while cells expressing

both *RORγt-eGFP* and mKate2 (or *eGFP* alone) with an amoeboid (non-spherical) cell shape

were scored as ILC3. T cell interactions with cDC or ILC3 was strictly measured as direct (<1

micron) colocalization of cells with respective fluorescent and cell morphology combinations.

Statistical analysis

For animal studies, mutant and control groups did not always have similar standard deviations and therefore an unpaired two-sided Welch's t-test was used. Error bars represent ± s.d. Animal sample size estimates were determined using power analysis (power = 90% and α = 0.05) based on the mean and s.d. from our previous studies and/or pilot studies using 4–5 mice. No samples were excluded from analysis. For analysis of ILC3 counts in MLN and LP of chimeric mice, a paired two-sided t-test was used.

Data availability

Data generated for this project are available at the Gene Expression Omnibus with the accession code GSE190372 and XXX. Published data GSE176282 was used for analysis.

Code availability

All code used for analysis in this manuscript is available at https://github.com/nygctech/Kedmi-CITEseq.

Additional References

39   Eberl, G. & Littman, D. R. Thymic origin of intestinal alphabeta T cells revealed by fate mapping of RORgammat+ cells. *Science* **305**, 248-251, doi:10.1126/science.1096472 (2004).

40   Lochner, M. *et al.* In vivo equilibrium of proinflammatory IL-17+ and regulatory IL-10+ Foxp3+ RORgamma t+ T cells. *The Journal of experimental medicine* **205**, 1381-1393, doi:10.1084/jem.20080034 (2008).

41   Kaplan, D. H., Jenison, M. C., Saeland, S., Shlomchik, W. D. & Shlomchik, M. J. Epidermal

langerhans cell-deficient mice develop enhanced contact hypersensitivity. *Immunity* **23**,

611-620, doi:10.1016/j.immuni.2005.10.008 (2005).

42   Dow, L. E. *et al.* Conditional reverse tet-transactivator mouse strains for the efficient

induction of TRE-regulated transgenes in mice. *PLoS One* **9**, e95236,

doi:10.1371/journal.pone.0095236 (2014).

43   Metzger, T. C. *et al.* Lineage tracing and cell ablation identify a post-Aire-expressing

thymic epithelial cell population. *Cell Rep* **5**, 166-179, doi:10.1016/j.celrep.2013.08.038

(2013).

44   Stoeckius, M. *et al.* Simultaneous epitope and transcriptome measurement in single

cells. *Nat Methods* **14**, 865-868, doi:10.1038/nmeth.4380 (2017).

45   Stoeckius, M. *et al.* Cell Hashing with barcoded antibodies enables multiplexing and

doublet detection for single cell genomics. *Genome Biol* **19**, 224, doi:10.1186/s13059-

018-1603-1 (2018).

46   van Buggenum, J. A. *et al.* A covalent and cleavable antibody-DNA conjugation strategy

for sensitive protein detection via immuno-PCR. *Sci Rep* **6**, 22675,

doi:10.1038/srep22675 (2016).

47   Hafemeister, C. & Satija, R. Normalization and variance stabilization of single-cell RNA-

seq data using regularized negative binomial regression. *Genome Biol* **20**, 296,

doi:10.1186/s13059-019-1874-1 (2019).

48   Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587

e3529, doi:10.1016/j.cell.2021.04.048 (2021).

49   Wolock, S. L., Lopez, R. & Klein, A. M. Scrublet: Computational Identification of Cell

Doublets in Single-Cell Transcriptomic Data. *Cell Syst* **8**, 281-291 e289,

doi:10.1016/j.cels.2018.11.005 (2019).

50   Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data

analysis. *Genome Biol* **19**, 15, doi:10.1186/s13059-017-1382-0 (2018).

51   Lopez, R., Regier, J., Cole, M. B., Jordan, M. I. & Yosef, N. Deep generative modeling for

single-cell transcriptomics. *Nat Methods* **15**, 1053-1058, doi:10.1038/s41592-018-0229-

2 (2018).

1020