1 Characterization of Reactive Oxygen Species Signaling changes in a cell culture model of
2 skeletal muscle ageing, and its application to screening pharmacokinetically-relevant
3 exposures of dietary polyphenols for bioactivity.

4   N. Hayes[1], M. Fogarty[2], L. Sadofsky[3], and H.S. Jones[4]*

[1]Department of Biological and Marine Sciences, University of Hull, Hull, HU6 7RX.
[2]Leeds Trinity University, Leeds, LS18 5HD.
[3]Centre for Atherothrombotic and Metabolic Research, Hull York Medical School, Hull, HU6
7RX.
[4]Institute of Cancer Therapeutics, University of Bradford, Bradford, BD7 1DP.
*Corresponding author (h.s.jones@bradford.ac.uk)

11
Data availability statement
The data that support the findings of this study are available from the corresponding author
upon reasonable request.

Funding statement
NH is supported by a University of Hull-funded PhD studentship. No other funding sources
supported this research.

19
Conflict of interest disclosure
There are no conflicts of interest for any of the authors of the manuscript

Abstract
Age-related frailty is a significant health and social care burden, however treatment options
are limited. There is currently a lack of suitable cell culture model for screening large
numbers of test compounds to identifying those which can potentially promote healthy
skeletal muscle function. This paper describes the characterization of reactive oxygen and
nitrogen species (RONS) signalling changes in young and aged myoblasts and myotubes
using the C_2C_{12} cell line, and the application of aged myoblast and myotube cultures to
assess the effect of dietary polyphenols on RONS signalling. Aged myoblasts and myotubes
were observed to have significantly increased reactive oxygen species levels ($p<0.01$ and
$p<0.001$ respectively), increases in nitric oxide levels ($p<0.05$ for myoblasts and myotubes),
and lipid peroxidation markers ($p<0.05$ for myoblasts and myotubes). A panel of nine
polyphenols were assessed in aged myoblasts and myotubes using concentrations and
incubation times consistent with known pharmacokinetic parameters for these compounds.
Of these, although several polyphenols were seen to reduce single markers of RONS
signalling, only kaempferol and resveratrol consistently reduced multiple markers of RONS
signalling with statistical significance in both cell models. Modulation of cellular enzymatic
antioxidant activities (superoxide and catalase) was assessed as a possible mechanism of
action for these polyphenols, and although both superoxide and catalase activities were
significantly reduced in aged (versus young) myotubes ($p<0.01$ and $p<0.05$ respectively), no
effect of polyphenol treatment on these enzyme activities were observed. Overall, this
research has shown the utility of the C_2C_{12} model, as both myoblasts and myotubes, as a
suitable cell model for screening compounds for modulating RONS signalling in aged
muscle, and that resveratrol and kaempferol (using pharmacokinetically-informed exposures)
can modulate RONS signalling in skeletal muscle cells after an acute exposure.

47
48
49
50

Introduction

Age-related muscle dysfunction is a major healthcare issue in elderly patient, and is associated with significant reductions in functional capacity, independence, and quality of life (Int. J. Endocrinol., 2021, 5563960). It is also associated with increased risk of chronic debilitating disorders (such as obesity), falls, hospitalization and mortality (Int. J. Endocrinol., 2021, 5563960). Currently, the management of age-related muscle dysfunction consists of diet and lifestyle changes (e.g., increased physical exercise, intake of protein, vitamin D and calcium, weight management), and symptomatic treatments such as pain relief (Ther. Adv. Muscluloskelet Dis., 2021, 13, 1759720X211009018). These approaches have had some success in improving patient quality of life, however this outcome is not consistent for all patients (Ther. Adv. Muscluloskelet Dis., 2021, 13, 1759720X211009018). Due to the increasing life expectancy, the number of elderly patients suffering with age-related loss of muscle function is predicted to also increase (Int. J. Endocrinol., 2021, 5563960). Thus, identifying new treatments for age-related muscle dysfunction is an important clinical need.

Reactive oxygen and nitrogen species (RONS) signaling is well-accepted to have important physiological signaling functions in healthy skeletal muscle, and its dysregulation is also strongly implicated in age-related skeletal muscle dysfunction (J. Physiol., 2016, 594, 5081; Biogerontology, 2020, 21, 475). This makes measurements of RONS signaling important biomarkers, and therapeutic targets of skeletal muscle dysfunction. The current state of play in this area has recently been reviewed by Thoma et al (Biogerontology, 2020, 21, 475), highlighting current attempts to modulate RONS signaling have produced mixed results, dependent on the model systems and dosing regimes used (Biogerontology, 2020, 21, 475).

The dietary polyphenol resveratrol has shown encouraging results in terms of reducing RONS signaling dysfunction and improvement in aged muscle function in both cell culture and mouse models (Sci. Rep., 2015, 28, 8093; J. Nutr. Biochem., 2017, 50, 103; J. Physiol. Sci., 2018, 68, 681). There are some limitations to these studies, particularly the use of supra-physiological concentrations of polyphenols for supra-physiological exposure times, this lack of reflection of the pharmacokinetics of polyphenols is becoming a more recognized issue within dietary polyphenol research (J. Agric. Food Chem., 66, 7857; J. Agric. Food Chem., 66, 8221). Taken together, this suggests that dietary polyphenols such as resveratrol could have therapeutic application to age-related muscle dysfunction, however a more thorough and physiologically-relevant assessment of these compounds needs to be done.

This study, using the C_2C_{12} murine myotube model described by Sharples et al (J. Cell. Physiol., 2010, 225, 240), characterizes changes in RONS signaling between young and aged myoblasts and myotubes, which has not been previously reported for this model system. A range of dietary polyphenols, using pharmacokinetics-informed concentrations and exposure durations, were assessed for effects on RONS signaling in aged cells, to identify those which ameliorate age-related changes in these markers. Those polyphenols which alter ROS signaling in the screening assays were then further characterized for effects on ROS signaling in aged myoblasts and myotubes.

Materials and Methods

Materials

All materials were purchased from Sigma-Aldrich (Poole, UK) unless otherwise stated.

Cell culture

Murine myoblasts (C_2C_{12}) were purchased from Public Health England at Passage 13 and
seeded into either T-75 flasks (Biolite, Thermofisher Scientific) , 6 well plates (Biolite,
Thermofisher) or 96 well plates (Biolite, Thermofisher Scientific) at a density of $2.75x10^4$
cells/cm^2. Myoblasts were cultured in high glucose Dulbecco's Modified Eagle Medium
(DMEM) containing L-glutamine, 10% v/v FBS, 100 U penicillin, 100 µg/mL streptomycin,
and 12.5µg/mL amphotericin B, to a confluency of 80%, and reseeded using a 0.5% trypsin
solution. To differentiate myoblasts into myotubes, cells were serum-starved (DMEM media
containing L- Glutamine, 2% v/v horse serum, 100 U penicillin, 100 µg/mL streptomycin, and
12.5µg/mL amphotericin B) for 5-7 days, with media refreshed every 2 days.
110
111   To age myoblast and myotube cultures, myoblasts were continuously cultured for 50
112   population doublings as outlined by Sharples et al 2010. Myoblasts less than passage 20
113   were defined as "young", with those above passage 30 as "aged". For aged myotubes,
114   myoblasts were cultured to the appropriate passage number, and were then differentiated as
115   described above.
116
Assessment of myotube length and width
C_2C_{12} myotubes were cultured as described above and imaged by Olympus IX71 Inverted
Fluorescence Microscope (10x UPlanXApo Objective), and the thickness of myotubes was
determined using the ImageJ software (Version 1.53m). Five 6-well plates were used for
each sample group 3 images per well, 5 fibres per image, a mean of three measurements of
each myotube diameter was taken and one centermost measurement was taken for length.
123
124
FOX assay
Lysates of myoblasts and myotubes grown in T-75 flasks were prepared after washing
cultures with phosphate buffered saline (PBS). Lysates were stored at -20°C until analysed.
Cell lysate or hydrogen peroxide standards (80 µL, 0-5 µM) were added into wells of a 96-
well plate, with FOX reagent (7.6 mg of xylenol orange, 9.8 mg of ammonium ferrous
sulphate and 1.8217 g of sorbitol together in 100 mL of 25 mM sulphuric acid, 120 µL per
well) added to each well, and incubated for 30 min in the dark. Absorbance was measured
at λ=560 nm using a Tecan infinite 200 mx-pro plate reader. Lipohydroperoxide
concentrations were calculated for each sample using the hydrogen peroxide standard
curve, and then normalized to the protein content of each lysate using Bradford assay.
135
TBARS assay
Lysates of myoblasts and myotubes grown in T-75 flasks were prepared after washing
cultures with PBS. Lysate was collected using a cell scraper and suspended in 1.5 mL PBS
and stored at -20°C until analysed. The following steps were performed on ice and light
shielded wherever possible.1 mL of cell lysate suspension was added to 100 µL of 6%
Perchloric acid (PCA) and incubated for 10 mins, Supernatant was spun off on max speed,
4°C for 5 mins. Supernatant was then neutralised with 10 µL of 6 M Potassium Hydroxide
Supernatant was spun off on max speed, 4°C for 5 mins and filtered through a 0.22 µm filter.
40 µL of sample or 1,1,3,3-Tetraethoxypropane (TEP)/ 40% ethanol standard (0-10 µM)
were combined with 200 µL 438 mM Phosphoric acid, 200 µL 41.6 mM Thiobarituric acid
(TBA) and 350 µL 18.2 MΩ Water. Samples and standards were then heated at 90°C for 60
min in the dark. Samples were then transferred to amber HPLC vials and sealed with a crimp
top lid.
Chromatographic separation was achieved using an Agilent G1311A quaternary pump,
Agilent G1367E1260 Infinity High Performance Autosampler and Agilent 1100 Series
G1316A COLCOM Column Compartment connected to an Agilent 1260 Infinity

Fluorescence Detector. An Agilent ZORBAX Eclipse Plus C18 column (5 µm pore size, 4.6 x
250 mm, Agilent Technologies, Cheshire, UK) was used for separating the analytes as
detailed below: Solvent A consisted of Phosphate buffer (70% 25 mM KH_2PO_4 : 30% 25 mM
Na_2HPO_4) and Solvent B was Acetonitrile. The Column was maintained at 25°C, Flow rate
was 1 mL/min, injection volume was 25 µL, starting conditions were equilibrated over 10
min. The method began at 25% Solvent B from 0-1.5 min, rising to 50% over 1.51-3 min,
holding at 50% until 4 min then dropping at 4.01 min, finally holding at 25% until 6 min. Using
FLD λex = 532 nm and λem = 553 nm, TBARS were detected at RT = 3.5 min.

160
161
DHE assay
Myoblasts and myotubes were cultured in a 96-well plate as described above, leaving the
edge wells of the plate free of cells. Once cells had reached the desired confluence or
differentiated into myotubes, cell culture media was removed from the wells and replaced
with PBS, containing 10 µM Dihydroethidium (DHE, Abcam, UK, 100 µL per well). The edge
wells were filled with 100 µL PBS. The well plates also contained cell only control wells (no
DHE added), DHE only control wells (no cells), and positive control wells (10 µM DHE, 1 mU
xanthine oxidase, 1 mM xanthine, in PBS). The plate was incubated at 37°C in a BMG
Labtech fluorostar omega plate reader, with fluorescence intensity of each well measured
each minute (using orbital averaging) for 30 min at λ_{ex} 544 nm and λ_{em} 590 nm. At the end
of the assay, the wells were emptied and replaced with 100 µL of PBS, and frozen before
quantification of protein content for each well using the Bradford assay. The data was
analysed by calculating the linear rate of DHE fluorescence increase for each well, which
was then corrected for protein content and the rate of the positive control samples. This was
done to ensure reproducibility between different plates and experimental days.

177
178
DCF assay
Myoblasts and myotubes were cultured in a 96-well plate as described above, leaving the
edge wells of the plate free of cells. Once cells had reached the desired confluence or
differentiated into myotubes, cell culture media was removed from the wells and replaced
with PBS, containing 10 µM Dichlorofluorescein (DCF, Abcam, UK, 100 µL per well). The
edge wells were filled with 100 µL PBS. The well plates also contained cell only control
wells (no DHE added), DHE only control wells (no cells), and positive control wells (10 µM
DCF, 1 mU xanthine oxidase, 1 mM xanthine, in PBS). The plate was incubated at 37°C in
a BMG Labtech fluorostar omega plate reader, with fluorescence intensity of each well
measured each minute (using orbital averaging) for 30 min at λ_{ex} 485 nm and λ_{em} 520 nm.
At the end of the assay, the wells were emptied and replaced with 100 µL of PBS, and
frozen before quantification of protein content for each well using the Bradford assay. The
data was analysed by calculating the linear rate of DCF fluorescence increase for each well,
which was then corrected for protein content and the rate of the positive control samples.
This was done to ensure reproducibility between different plates and experimental days.

194
DAF assay
Myoblasts and myotubes were cultured in a 96-well plate as described above, leaving the
edge wells of the plate free of cells. Once cells had reached the desired confluence or
differentiated into myotubes, cell culture media was removed from the wells and replaced
with PBS, containing 2 µM 4,5-Diaminofluorescein diacetate (DAF, Abcam, UK, 100 µL per
well). The edge wells were filled with 100 µL PBS. The well plates also contained cell only
control wells (no DAF added), DAF only control wells (no cells), and positive control wells (2
µM DAF, 10 µM PAPA NONOate, in PBS). The plate was incubated at 37°C in a TECAN

Infinite 200 PRO plate reader, with fluorescence intensity of each well measured each
minute (using orbital averaging) for 30 min at λ_{ex} 495 nm and λ_{em} 515 nm. At the end of the
assay, the wells were emptied and replaced with 100 μL of PBS, and frozen before
quantification of protein content for each well using the Bradford assay. The data was
analysed by calculating the linear rate of DAF fluorescence increase for each well, which
was then corrected for protein content and the rate of the positive control samples. This was
done to ensure reproducibility between different plates and experimental days.
210
PC assay
Dinitrophenylhydrazine (10 mM in 0.5 M phosphoric acid, 100 μL per well) was added to
wells containing 50 μL of cell lysate (prepared in PBS) and 50 μL of 18.2 MΩ water and
incubated for 10 min at room temperature in the dark. To each well, 50 μL of 6 M sodium
hydroxide was added, and the plate was incubated for 10 min at room temperature, followed
by measurement of absorbance at 450 nm using a Tecan infinite 200 mx-pro plate reader.
Protein carbonyl concentrations were determined for blank corrected samples using the
Beer-Lambert law (molar extinction coefficient = 33131 $M^{-1}cm^{-1}$), followed by normalization to
the protein content of each lysate.
220
Enzymatic antioxidant activity assays
Superoxide dismutase enzyme activity was quantified using the Abcam SOD activity assay
kit (Ab65354, Abcam, UK) as directed by the manufacturer. Catalase activity was
determined using the Abcam catalase activity assay kit (Ab83464, Abcam, UK) as directed
by the manufacturer. Enzyme activity (measured in units) was corrected for the protein
concentration of samples (determined by Bradford assay as described below) for both
enzyme activities.
228
Bradford assay
Cell lysates (5 μL per well) were added to a 96 well plate, along with BSA standards (
0-1.4 mg/mL in PBS, 5 μL per well) in triplicate. Bradford reagent (200 μL per well) was
added to these samples and standards and the plate was incubated for 45 min in the dark at
room temperature. Absorbance at 595 nm was measured using a Tecan infinite200 mx-pro
plate reader. Protein concentration per μL of cell lysate was determined for each sample
using the calibration curve of BSA standards.
236
237
238 *Statistical analysis*
239 All data was checked for normality of distribution using a Shapiro-Wilks test. Data which
240 was normal was analysed using an independent t-test or one-way ANOVA as indicated in
241 the figure legends. Non-parametric data was analysed using either a Mann-Whitney U or
242 Kruskal-Wallis test as indicated in the figure legends.
243
244
245
246 **Results**
247
Aged myotubes show morphological changes compared to young myotubes
Myotube width and length were compared for young and aged cultures (Figure 1). Aged
myoblasts were significantly thinner (Figure 1A, p<0.001, unpaired t-test) and significantly
shorter (Figure 1B, p<0.001, Mann-Whitney U test) than young myotubes. This confirmed
that, under the culture conditions used in this study, the aged myotubes showed a
dysfunctional phenotype compared to young cells.

254
255

256
Figure 1: Effect of ageing protocol on myotube width and length. Representative
images of young (A) and aged (B) myotubes used in analysis of myotube length and width.
Scale bars equate to 100 µm. Quantification of myotube width (C) and length (D) were
significantly different between young and aged. Individual replicate experiments are
represented by each data-point on the graph, with the median value indicated by the
horizontal line. Young and aged datasets were compared using unpaired t-test or Mann-
Whitney U test, with **** p<0.001). Panel A shows myotube width (n=15 per group) and
panel B shows myotube length (n=15 per group).

*Aged myoblasts show increased markers of RONS and oxidative stress compared to young
myoblasts*
Both young and aged myoblasts were assessed for markers of ROS production, lipid
oxidation and protein oxidation (Figure 2). ROS was significantly increased in aged
myoblasts compared with young myoblasts by DHE assay (Figure 2A and 1B, p<0.01 and
p<0.05 respectively, unpaired t-test). Two markers of lipid oxidation were assessed (FOX
assay and TBARS assay), with both markers showing a median increase in aged (compared
with young) myoblasts (Figure 2C and D, p>0.05 for FOX assay, p<0.05 for TBARS assay).
Protein carbonyls (measured using dinitrophenylhydrazine), as a marker of protein oxidation,
showed no differences between young and aged myoblasts (Figure 2E, p>0.05, unpaired t-
test). Nitric oxide levels, measured by the DAF assay, were also increased in aged
myoblasts compared to young myoblast cultures (Figure 2F, p<0.05, unpaired t-test).

Aged myotubes show increased markers of RONS and oxidative stress compared to young
myotubes
Both young and aged myotubes were assessed for markers of ROS production, lipid
oxidation and protein oxidation (Figure 3). ROS was significantly increased in aged
myoblasts compared with young myotubes by DHE assay (Figure 3A, $p<0.001$, unpaired t-
test), however this difference was not observed when ROS levels were assessed using the
DCF assay (Figure 3B, $p>0.05$, unpaired t-test). Two markers of lipid oxidation were
assessed (FOX assay and TBARS assay), with FOX assay showing a significant increase in
aged compared to young, and TBARS assay showing no change in aged (compared with
young) myotubes (Figure 3C and D, $p>0.05$ for FOX assay, $p<0.05$ for TBARS assay).
Protein carbonyls, as a marker of protein oxidation, showed no differences between young
and aged myotubes (Figure 3E, $p>0.05$, unpaired t-test). Nitric oxide levels were also
increased in aged myotubes compared to young myotube cultures (Figure 3F, $p<0.05$,
unpaired t-test).
294
295
Screening of polyphenols at physiologically-relevant exposures in aged myoblasts shows
differential effects on RONS signalling
The dietary polyphenols quercetin, chrysin, curcumin, ellagic acid, kaempferol, resveratrol,
epigallocatechin gallate, 3-hydroxy-tyrosol and gallic acid we assessed for effects on ROS
levels in aged myoblasts (Figure 4) using the DHE assay. In aged myoblasts, curcumin and
kaempferol (at 1 µM, 4 h) significantly reduced ROS production (Figure 4A, curcumin 56% ±
4%, kaempferol 54% ± 6%, $p<0.05$, Kruskal-Wallis test with Dunn's multiple comparison
versus solvent control). ROS production was also reduced by chrysin, ellagic acid and
resveratrol, albeit not with statistical significance (Figure 4A).
305
Chrysin, curcumin, kaempferol, resveratrol and epigallocatechin gallate (EGCG) were
selected for further characterisation, where the effect of exposure to these compounds (at 1
µM and 100 nM, 4 h) upon nitric oxide production (Figure 4B) and lipid peroxidation (Figure
4C and D) was determined. Nitric oxide production was significantly reduced by both
curcumin (1 µM: $p<0.0001$; 100 nM: $p<0.01$) and resveratrol (1 µM: $p<0.05$; 100 nM: $p<0.01$,
2-way ANOVA with Tukey's multiple comparisons test). Chrysin (100 nM, 4 h) also
approached statistical significance ($p=0.052$, 2-way ANOVA with Tukey's multiple
comparisons test).
314
Kaempferol (at 1 µM and 100 nM, 4 h) and EGCG (100 nM, 4 h) reduced lipohydroperoxide
concentrations in aged myoblasts (Figure 4C), however only kaempferol (1 µM, 4 h) reached
statistical significance ($p<0.05$, 2-way ANOVA with Tukey's multiple comparisons test),
whereas 100 nM kaempferol and 100 nM EGCG reached near-significance ($p=0.058$ and
$p=0.068$ respectively).   In the TBARS assay (Figure 4D), significant reductions in lipid
peroxidation were observed for chrysin (1 µM: $p<0.01$; 100 nM: $p<0.05$), curcumin (100 nM:
$p<0.05$), kaempferol (1 µM: $p<0.01$; 100 nM: $p<0.01$), and resveratrol (1 µM: $p<0.01$; 100
nM: $p<0.05$; all the above significances tested using a 2-way ANOVA with Tukey's multiple
comparisons test).
324
325

326
Figure 2: Characterisation of RONS signaling in young and aged myoblasts. Individual
replicate experiments are represented by each data-point on the graph, with the median
value indicated by the horizontal line. Young and aged datasets were compared using
unpaired t-tests, with * p<0.05 and **p<0.01) Panel A: ROS levels were assessed in young
and aged myoblast cultures by DHE fluorescence. Panel B: ROS levels were assessed in
young and aged myoblast cultures by DCF fluorescence. Panel C: Lipohydroperoxides were
measured by FOX assay. Panel D: Thiobuturic acid reactive substances were measured by
HPLC TBARS assay. Panel E: Protein carbonyls measured using dinitrophenylhydrazine.
Panel F: Nitric oxide levels measured using the DAF.

336
Figure 3: Characterisation of RONS signaling in young and aged myotubes. Individual
replicate experiments are represented by each data-point on the graph, with the median
value indicated by the horizontal line. Young and aged datasets were compared using
unpaired t-tests, with ns p>0.05, * p<0.05 and **p<0.01)   Panel A: ROS levels were
assessed in young and aged myotube cultures by DHE fluorescence. Panel B: ROS levels
were assessed in young and aged myotube cultures by DCF fluorescence. Panel C:
Lipohydroperoxides were measured by FOX assay. Panel D: Thiobuturic acid reactive
substances were measured by HPLC TBARS assay. Panel E: Protein carbonyls measured
using dinitrophenylhydrazine. Panel F: Nitric oxide levels measured using DAF.

347
348
Figure 4: Screening of dietary polyphenols at physiologically-relevant concentrations
for effects on ROS production in aged myoblasts, and characterization of hits for
effects on lipid peroxidation and nitric oxide levels.   Individual replicate experiments are
represented by each data-point on the graph, with the median value indicated by the
horizontal line. The dashed line on the graphs indicates the solvent control level. Panel A:
The effect of a range of polyphenols (1 µM, 4 h) on ROS levels was determined using the
DHE assay. Although several polyphenols reduced ROS levels in aged myoblasts, only

curcumin and kaempferol reached statistical significance (p<0.05, Kruskal-Wallis test with
Dunn's multiple comparison versus solvent control). Panel B-D: The effects of selected
polyphenols (1 µM (circles) or 100 nM (triangles), 4 h) on nitric oxide levels (DAF assay,
Panel B), lipohydroperoxide concentration (Fox assay, Panel C), and TBARS (Panel D) were
determined.   Statistical analysis of these data used two-way ANOVA with Tukey's multiple
comparisons test.
362
363 *Screening of polyphenols at physiologically-relevant exposures in aged myotubes shows*
364 *differential effects on RONS signalling and myotube morphology.*
365
The dietary polyphenols quercetin, chrysin, curcumin, ellagic acid, kaempferol, resveratrol,
epigallocatechin gallate, 3-hydroxy-tyrosol and gallic acid were assessed for effects on ROS
levels in aged myotubes (Figure 5) using the DHE assay. In aged myotubes, curcumin and
kaempferol (at 1 µM, 4 h) significantly reduced ROS production (Figure 5A, curcumin 50% ±
9%, kaempferol 49% ± 2%, p<0.05, Kruskal-Wallis test with Dunn's multiple comparison
versus solvent control). ROS production was also reduced by chrysin, ellagic acid and
resveratrol, albeit not with statistical significance (Figure 5A).
373
Chrysin, curcumin, kaempferol, resveratrol and epigallocatechin gallate (EGCG) were
selected for further characterisation, where the effect exposure to these compounds (at 1 µM
and 100 nM, 4 h) upon nitric oxide production (Figure 5B) and lipid peroxidation (Figure 5C
and D) was determined. Nitric oxide production was significantly reduced by chrysin (100
nM: p<0.01), curcumin (1 µM: p<0.0001; 100 nM: p<0.0001) and resveratrol (1 µM: p<0.001;
100 nM: p<0.0001, 2-way ANOVA with Tukey's multiple comparisons test).
Kaempferol (at 1 µM and 100 nM, 4 h) and resveratrol (100 nM, 4 h) reduced
lipohydroperoxide concentrations in aged myotubes (Figure 5C) with statistical significance
(kaempferol 1 µM: p<0.05; 100 nM: p<0.01; resveratrol 100 nM: p<0.05; 2-way ANOVA with
Tukey's multiple comparisons test).   In the TBARS assay (Figure 5D), no significant
reductions in lipid peroxidation were observed.
385
386

387
388
Figure 5: Screening of polyphenols in aged myotubes for effects on ROS levels, and
characterization of hits. Individual replicate experiments are represented by each data-
point on the graph, with the median value indicated by the horizontal line. The dashed line
on the graphs indicates the solvent control level. Panel A: The effect of a range of
polyphenols (1 µM, 4 h) on ROS levels was determined using the DHE assay. Although
several polyphenols reduced ROS levels in aged myoblasts, only curcumin and kaempferol
reached statistical significance (p<0.05, Kruskal-Wallis test with Dunn's multiple comparison
versus solvent control). Panel B-D: The effects of selected polyphenols (1 µM (circles) or
100 nM (triangles), 4 h) on nitric oxide levels (DAF assay, Panel B), lipohydroperoxide
concentration (Fox assay, Panel C), and TBARS (Panel D) were determined. Statistical
analysis of these data used two-way ANOVA with Tukey's multiple comparisons test.
Statistical significance is indicated in the main text.
401

402 *The modulation of ROS and macromolecule oxidation in aged skeletal myotubes by dietary*
403 *polyphenols is not due to upregulation of enzymatic antioxidant capacity.*
404
One potential mechanism of action for the observed effects of acute polyphenol treatment is
via the upregulation of cellular antioxidant defences. To test this hypothesis, superoxide
dismutase (SOD) and catalase (CAT) activities were assessed in myotube cultures (Figure
6). Statistically significant decreases in SOD (Figure 6A, $p<0.01$, unpaired t-test) and CAT
(Figure 6B, $p<0.05$, unpaired t-test) activities were observed when comparing young and
aged myotubes, suggesting that cellular antioxidant defence may be compromised in aged
myotubes. Dietary polyphenol treatment of aged myotubes did not alter either SOD (Figure
6C) or CAT (Figure 6D) activities, suggesting upregulation of cellular enzymatic antioxidant
capacity is not part of the mechanism of action for the observed reduction of ROS and lipid
oxidation in aged myotubes.
415
416

417
418
Figure 6: Superoxide dismutase and catalase activities in aged myotubes are not
modulated by polyphenol exposure. Panel A and C show catalase activity assays, panel
B and D show superoxide dismutase (SOD) assays. Both catalase (A) and SOD (B)
activities are significantly reduced in aged myotubes compared to young myotubes (*
$p<0.05$, **$p<0.01$, unpaired t-test). Polyphenol exposure (100 nM, 4 h) does not significantly
alter Catalase (C) or SOD (D) activities in aged myotubes ($p>0.05$, one-way ANOVA and
Kruskal-Wallis test respectively). Individual replicate experiments are represented by each
data-point on the graph, with the median value indicated by the horizontal line.
427

428   *Acute kaempferol exposure of aged myoblasts partially induces a young phenotype when*
429   *differentiated into myotubes.*
430
To understand if acute exposure to these polyphenols impacted on myotube morphology,
myoblasts were treated with 100 nM for 4 h of Quercetin, Kaempferol, EGCG or Resveratrol
prior to differentiation into myotubes (Figure 7). Kaempferol exposure resulted in a
significant ($p<0.05$) increase in myotube length (Figure 7C), but no change in myotube width
(Figure 7D). When aged myotubes were incubated with these polyphenols after
differentiation, no changes in in myotube length or width were observed (Figure 8C and 8D).
437
438
439

440
441
Figure 7: The effect of acute polyphenol exposure on myotube length and width.
Panels A and B show representative images of myotubes resulting from the differentiation of
solvent control (A) and kaempferol-treated (100 nM, 4 h, Panels B) aged myoblasts treated
before differentiation. Scale bars indicate 100 µm. Panels C (myotube length) and D
(myotube width) for solvent only or polyphenol-treated (100 nM, 4 h) myoblasts prior to
differentiation. Individual replicate experiments are represented by each data-point on the
graph, with the median value indicated by the horizontal line. Datasets were compared

using a Kruskal Wallis test (p<0.01), with Dunn's multiple comparisons test versus the
solvent control; * p<0.05.
451
452

453
454
Figure 8: The effect of acute polyphenol exposure post differentation on myotube
length and width. Panels A and B show representative images of aged myotubes treated
with solvent control (A) and kaempferol-treated (100 nM, 4 h, Panels B) after differentiation.
Panels C (myotube length) and D (myotube width) for solvent only or polyphenol-treated
(100 nM, 4 h) myoblasts post differentiation. Individual replicate experiments are
represented by each data-point on the graph, with the median value indicated by the
horizontal line. Datasets were compared using a Kruskal Wallis test (p>0.05).
462
463
464
Discussion
Although oxidative stress endpoints have been previously assessed in C_2C_{12} cells, this study
is the first (to authors knowledge) to undertake a thorough assessment of RONS signaling
changes between young and aged (by multiple population doublings) myoblast and myotube
cell cultures, and use this model to screen for potential modifiers of RONS signaling
dysfunction in aged skeletal muscle. Previous studies of oxidative stress in C_2C_{12} have used
a range of stressors such as TNF-α (J Pharm Biomed Anal, 2021, 204, 114271), LPS (Poult

Sci. 2019 Jul 1;98(7):2756-2764), and hydrogen peroxide (J Biosci Bioeng. 2021
May;131(5):572-578; Molecules 2021 Jan 4;26(1):215; J Pharmacol Exp Ther. 2021
Mar;376(3):385-396; Front Cell Dev Biol. 2020 Sep 11;8:541260), whereas this study uses
repeated population doublings to mimic the ageing process. The C_2C_{12} model used in this
study has been well-characterized in a physiological context (such as alterations to myotube
structure, time to differentiation from myoblasts to myotubes, maintenance of a skeletal
muscle phenotype, e.g., J. Cell. Physiol., 2010, 225, 240) to reflect an aged skeletal muscle
phenotype, an observation our study agrees with (based on a reduction of myotube length
and width compared with young cells, Figure 1), and this study has now demonstrated that it
also reflects biochemical alterations in RONS signaling, such as those reported from aged
muscle biopsies. Increases in ROS levels and markers of lipid oxidation have been shown
for both myoblasts and myotubes in aged cultures compared with young cultures, whereas
protein carbonyl levels were not different in young and aged cultures (Figures 2 and 3).
These observations show good agreement with human biopsy data for studies comparing
young and aged volunteers, showing consistent increases in lipid oxidation markers (biopsy
data is quite variable for protein carbonyl level, with some studies showing differences and
others no changes) supporting the validity of the myoblast and myotube culture models
(Free Rad. Biol. Med., 1999, 26, 303; Redox Biol., 2015, 5, 267; Free Rad. Biol. Med., 2006,
41, 797). Increases in nitric oxide levels have also been reported in aged skeletal muscle
samples from rodent models (e.g., Free Radic Biol Med. 2015 Jan;78:82-8) and 3-
nitrotyrosine (a marker of protein nitrosylation and surrogate marker for nitric oxide) has
been reported to increase in human biopsy samples from aged compared with young
participants (Cells 2019, 8(12), 1525). Taken together, these data suggest that the C_2C_{12}
model used in this study is representative of aged skeletal muscle biopsy samples in both
phenotypic and RONS signalling contexts.
497
A major limitation of skeletal muscle ageing and sarcopenia research is a lack of a tractable
model to screen compounds for potential beneficial effects on muscle ageing and
dysfunction. Although there are a range of *in vivo* model systems (e.g., murine and other
rodent models), these lack the throughput and capacity to screen a large number of
compounds. This study demonstrates that the C_2C_{12} ageing model can address this issue.
Both aged myoblast and myotube cultures were used as a screening tool to identify dietary
polyphenols with bioactivity on RONS signaling from a wider panel of polyphenols. Chrysin,
curcumin, kaempferol and resveratrol were identified to reduce elevated markers of RONS
levels and lipid peroxidation at physiologically-relevant exposures. Kaempferol also had a
significant effect on myotube length, when incubated with aged myoblasts prior to
differentiation (Figure 7), which is representative of a younger phenotype in this cell model of
ageing. This screening approach, combined with the use of compound exposure
concentrations and durations guided by known pharmacokinetic parameters for these
compounds, are two areas in which this work has advanced the field.
512
Resveratrol, curcumin, epigallocatechin gallate and quercetin have all been previously
assessed for effects on skeletal muscle aging in cell culture and rodent models (Int J Mol
Sci, 2019, 20, 1178; J Agric Food Chem, 2021, 69, 6214; FASEB, 2014, 28, 1159.4;
Antioxidants, 2021, 10, 476; Protein and Cell, 2019, 10, 770; Braz. Pharm Sci, 2020, 56; J
Frail Ageing, 2015, 4 209; Exp Gerontol, 2008, 43, 176; Arch Biochem Biophys, 2020, 692,
108511; Biochem and Biophys Res Comm, 2017, 489, 142). These studies have shown that
these polyphenols alter ROS signaling in skeletal muscle cells and lysate, and some, but not
all, rodent studies report improvements in muscle function with chronic exposures. A major
limitation of these studies is that the *in vitro* experiments use exposures which do not reflect
human pharmacokinetic data (for example, Chang et al exposed C_2C_{12} cells to 50 µM

epigallocatechin gallate for 24 h, whereas the reported Cmax and half-life for
epigallocatechin galate in humans is 447 ± 300 nM and 3.6 ± 1.4 h (Arch Biochem Biophys,
2020, 692, 108511, Cancer Epidemiol Biomarkers Prev. 2002, 1025-32.). Additionally, even
the rodent model studies highlighted above use impractical doses of polyphenols when
scaled to human studies (for example, both Braz. Pharm Sci, 2020, 56, and J Frail Ageing,
2015, 4 209-215 dosed rats with 200 mg/kg body weight per day, which when scaled to a 75
kg human (as detailed in J Basic Clin Pharm, 2016, 7, 27-31, taking body surface area into
consideration) equates to a bolus of 2.43 g of pure quercetin or epigallocatechin gallate per
day respectively. Human clinical trials using quercetin report minor side effects at long term
doses over 1g, (J Am Heart Assoc, 2016, 5, e002713), suggesting that the exposures used
in the rodent models are not reflective of what would be used in humans. This lack of
relevance to human pharmacokinetics in both in vitro and animal model experiments means
that it is difficult to extrapolate these findings to humans. This study has, for in vitro models,
used more pharmacokinetically-relevant exposures, using a 4 h exposure at 1 μM for initial
screening, and a lower concentration (100 nM, 4 h) which is also consistent with reported
pharmacokinetic data for the test compounds (Cancer Epidemiol Biomarkers Prev. 2008
Jun;17(6):1411-7; Cancer Epidemiol Biomarkers Prev. 2002, 1025-32; Br. J. Clin. Pharm.,
51 (2) (2001), pp. 143-146; Ann N Y Acad Sci. 2011 Jan;1215:9-15; Nutrients 2019, 11(10),
2288).
542
This study has also undertaken some limited mechanistic investigations, to try to understand
how dietary polyphenols such kaempferol modulate RONS signalling, via the assessment of
enzymatic antioxidant activities (SOD and catalase) in aged myotube cultures (Figure 6).
Kaempferol has been reported in a variety of studies to modulate SOD and catalase activity
in a variety of health and disease states (Drug and chemical toxicology, 39(3), 239–247;
Oxidative Medicine and Cellular Longevity, vol. 2018, Article ID 1610751; Redox Report,
20:5, 198-209). It should be noted, however, that the application of physiologically-relevant
kaempferol exposures in these studies is variable. This study reports that aged myotubes
have reduced SOD and catalase activities compared to young myotubes, and that none of
the polyphenols tested induced either SOD or catalase activities in aged myotube cultures.
This suggests that modulation of enzymatic antioxidant capacity is not the mechanism of
action of the polyphenols tested which underpins the reduction in ROS and lipid peroxidation
markers reported above (Figures 4 and 5).
556
It should be noted that there are some potentially important limitations with the experimental
design in this study. Dietary polyphenols undergo significant conjugative metabolism (e.g.,
glucuronidation, sulfation, methylation), however due to a lack of availability of human
polyphenol metabolites, they were not used in the experiments described in this study.
Additionally, the experiments described above assess the effects of acute exposures to
dietary polyphenols, whereas a chronic dosing is more likely to be experienced by humans.
It should also be acknowledged that this study only assesses the direct impact of polyphenol
exposure on skeletal muscle cells, and does not integrate effects on other cell types or
systems (e.g., neuronal cells, blood supply).
566
567
Conclusions
Overall, this study has characterized a myoblast and myotube cell culture model of skeletal
muscle ageing for changes in RONS signaling. These cultures show similar changes to
those reported for human skeletal muscle biopsies, and the modulation of RONS signaling
by physiologically-relevant dietary polyphenol exposures in aged myoblasts and myotubes

has been demonstrated. This model system can be used to identify bioactive compounds
which may be beneficial for restoring normal RONS signaling in aged skeletal muscle.
575
576
577
578

579 **References**

580 Ali AM, Kunugi H. Screening for Sarcopenia (Physical Frailty) in the COVID-19 Era. Int J
581 Endocrinol. 2021 May 21;2021:5563960.
582
583 Papageorgiou M, Biver E. Interactions of the microbiome with pharmacological and non-
584 pharmacological approaches for the management of ageing-related musculoskeletal
585 diseases. Ther Adv Musculoskelet Dis. 2021 Apr 20;13:1759720X211009018.
586
587 Powers SK, Radak Z, Ji LL. Exercise-induced oxidative stress: past, present and future. J
588 Physiol. 2016 Sep 15;594(18):5081-92.
589
590 Thoma A, Akter-Miah T, Reade RL, Lightfoot AP. Targeting reactive oxygen species (ROS)
591 to combat the age-related loss of muscle mass and function. Biogerontology. 2020
592 Aug;21(4):475-484.
593
594 Bosutti A, Degens H. The impact of resveratrol and hydrogen peroxide on muscle cell
595 plasticity shows a dose-dependent interaction. Sci Rep. 2015 Jan 28;5:8093.
596
597 Haramizu S, Asano S, Butler DC, Stanton DA, Hajira A, Mohamed JS, Alway SE. Dietary
598 resveratrol confers apoptotic resistance to oxidative stress in myoblasts. J Nutr Biochem.
599 2017 Dec;50:103-115
600
601 Muhammad MH, Allam MM. Resveratrol and/or exercise training counteract aging-
602 associated decline of physical endurance in aged mice; targeting mitochondrial biogenesis
603 and function. J Physiol Sci. 2018 Sep;68(5):681-688.
604
605 Ávila-Gálvez MÁ, González-Sarrías A, Espín JC. In Vitro Research on Dietary Polyphenols
606 and Health: A Call of Caution and a Guide on How To Proceed. J Agric Food Chem. 2018
607 Aug 1;66(30):7857-7858.
608
609 Mena P, Del Rio D. Gold Standards for Realistic (Poly)phenol Research. J Agric Food
610 Chem. 2018 Aug 8;66(31):8221-8223.
611
612 Sharples AP, Al-Shanti N, Stewart CE. C2 and C2C12 murine skeletal myoblast models of
613 atrophic and hypertrophic potential: relevance to disease and ageing? J Cell Physiol. 2010
614 Oct;225(1):240-50.
615
616 Ju L, Zhang J, Wang F, Zhu D, Pei T, He Z, Han Z, Wang M, Ma Y, Xiao W. Chemical
617 profiling of Houttuynia cordata Thunb. by UPLC-Q-TOF-MS and analysis of its antioxidant
618 activity in C2C12 cells. J Pharm Biomed Anal. 2021 Sep 10;204:114271.
619
620 Liu S, Adewole D, Yu L, Sid V, Wang B, O K, Yang C. Rutin attenuates inflammatory
621 responses induced by lipopolysaccharide in an in vitro mouse muscle cell (C2C12) model.
622 Poult Sci. 2019 Jul 1;98(7):2756-2764.
623

624 Fujita H, Mae K, Nagatani H, Horie M, Nagamori E. Effect of hydrogen peroxide
625 concentration on the maintenance and differentiation of cultured skeletal muscle cells. J
626 Biosci Bioeng. 2021 May;131(5):572-578.
627
628 Kim YS, Yuk HJ, Kim DS. Effect of Jakyakgamcho-Tang Extracts on H_2O_2-Induced C2C12
629 Myoblasts. Molecules. 2021 Jan 4;26(1):215.
630
631 Hosoda R, Hamada H, Uesugi D, Iwahara N, Nojima I, Horio Y, Kuno A. Different
632 Antioxidative and Antiapoptotic Effects of Piceatannol and Resveratrol. J Pharmacol Exp
633 Ther. 2021 Mar;376(3):385-396.
634
Li J, Yang Q, Han L, Pan C, Lei C, Chen H, Lan X. C2C12 Mouse Myoblasts Damage
Induced by Oxidative Stress Is Alleviated by the Antioxidant Capacity of the Active
Substance Phloretin. Front Cell Dev Biol. 2020 Sep 11;8:541260.
Mecocci P, Fanó G, Fulle S, MacGarvey U, Shinobu L, Polidori MC, Cherubini A, Vecchiet J,
Senin U, Beal MF. Age-dependent increases in oxidative damage to DNA, lipids, and
proteins in human skeletal muscle. Free Radic Biol Med. 1999 Feb;26(3-4):303-8.
641
642 Lourenço Dos Santos S, Baraibar MA, Lundberg S, Eeg-Olofsson O, Larsson L, Friguet B.
643 Oxidative proteome alterations during skeletal muscle ageing. Redox Biol. 2015 Aug;5:267-
644 274.
645
646 Barreiro E, Coronell C, Laviña B, Ramírez-Sarmiento A, Orozco-Levi M, Gea J; PENAM
647 Project. Aging, sex differences, and oxidative stress in human respiratory and limb muscles.
648 Free Radic Biol Med. 2006 Sep 1;41(5):797-809.
649
650 Pearson T, McArdle A, Jackson MJ. Nitric oxide availability is increased in contracting
651 skeletal muscle from aged mice, but does not differentially decrease muscle superoxide.
652 Free Radic Biol Med. 2015 Jan;78:82-8.
653
654 Picca A, Mankowski RT, Kamenov G, Anton SD, Manini TM, Buford TW, Saini SK, Calvani
655 R, Landi F, Bernabei R, Marzetti E, Leeuwenburgh C. Advanced Age Is Associated with Iron
656 Dyshomeostasis and Mitochondrial DNA Damage in Human Skeletal Muscle. Cells. 2019
657 Nov 27;8(12):1525.
658
659 Receno CN, Liang C, Korol DL, Atalay M, Heffernan KS, Brutsaert TD, DeRuisseau KC.
660 Effects of Prolonged Dietary Curcumin Exposure on Skeletal Muscle Biochemical and
661 Functional Responses of Aged Male Rats. Int J Mol Sci. 2019 Mar 7;20(5):1178.
662
663 ee DY, Chun YS, Kim JK, Lee JO, Ku SK, Shim SM. Curcumin Attenuates Sarcopenia in
664 Chronic Forced Exercise Executed Aged Mice by Regulating Muscle Degradation and
665 Protein Synthesis with Antioxidant and Anti-inflammatory Effects. J Agric Food Chem. 2021
666 Jun 9;69(22):6214-6228.
667
668 Wawrzyniak, N., Duarte, A., Nguyen, L., Joseph, A.-M., Layne, A., Criswell, D. and
669 Adhihetty, P. (2014), Effect of short-term dietary curcumin supplementation on mitochondrial
670 regulatory proteins in muscle and brown adipose tissue of aged mice (1159.4). The FASEB
671 Journal, 28: 1159.4.
672

673 Lee DY, Chun YS, Kim JK, Lee JO, Lee YJ, Ku SK, Shim SM. Curcumin Ameliorated
674 Oxidative Stress and Inflammation-Related Muscle Disorders in C2C12 Myoblast Cells.
675 Antioxidants (Basel). 2021 Mar 17;10(3):476.
676
677 Geng L, Liu Z, Wang S, Sun S, Ma S, Liu X, Chan P, Sun L, Song M, Zhang W, Liu GH, Qu
678 J. Low-dose quercetin positively regulates mouse healthspan. Protein Cell. 2019
679 Oct;10(10):770-775.
680
681 lm-Eldeen, Abeer et al. Quercetin modulates age-induced changes in the transcript levels of
682 some apoptosis related genes in the skeletal muscles of male rats. Brazilian Journal of
683 Pharmaceutical Sciences [online]. 2020, v. 56 [Accessed 7 December 2021] , e18861.
684 https://doi.org/10.1590/s2175-979020200003180861
685
686 M. Meador ; K.A. Mirza ; M. Tian ; M.B. Skelding ; L.A. Reaves ; N.K. Edens ; M.J. Tisdale ;
687 S.L. Pereira (2015): The green tea polyphenol Epigallocatechin-3-gallate (EGCg) attenuates
688 skeletal muscle atrophy in a rat model of sarcopenia. The Journal of Frailty and Aging (JFA).
689 http://dx.doi.org/10.14283/jfa.2015.58
690
691 Senthil Kumaran V, Arulmathi K, Srividhya R, Kalaiselvi P. Repletion of antioxidant status by
692 EGCG and retardation of oxidative damage induced macromolecular anomalies in aged rats.
693 Exp Gerontol. 2008 Mar;43(3):176-83.
694
695 Chang YC, Liu HW, Chan YC, Hu SH, Liu MY, Chang SJ. The green tea polyphenol
696 epigallocatechin-3-gallate attenuates age-associated muscle loss via regulation of miR-486-
697 5p and myostatin. Arch Biochem Biophys. 2020 Oct 15;692:108511.
698
699 Kim AR, Kim KM, Byun MR, Hwang JH, Park JI, Oh HT, Kim HK, Jeong MG, Hwang ES,
700 Hong JH. Catechins activate muscle stem cells by Myf5 induction and stimulate muscle
701 regeneration. Biochem Biophys Res Commun. 2017 Jul 22;489(2):142-148.
702
703 Lee MJ, Maliakal P, Chen L, Meng X, Bondoc FY, Prabhu S, Lambert G, Mohr S, Yang CS.
704 Pharmacokinetics of tea catechins after ingestion of green tea and (-)-epigallocatechin-3-
705 gallate by humans: formation of different metabolites and individual variability. Cancer
706 Epidemiol Biomarkers Prev. 2002 Oct;11(10 Pt 1):1025-32.
707
708 Nair AB, Jacob S. A simple practice guide for dose conversion between animals and human.
709 J Basic Clin Pharm. 2016 Mar;7(2):27-31.
710
711 Serban MC, Sahebkar A, Zanchetti A, Mikhailidis DP, Howard G, Antal D, Andrica F, Ahmed
712 A, Aronow WS, Muntner P, Lip GY, Graham I, Wong N, Rysz J, Banach M; Lipid and Blood
713 Pressure Meta-analysis Collaboration (LBPMC) Group. Effects of Quercetin on Blood
714 Pressure: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. J Am
715 Heart Assoc. 2016 Jul 12;5(7):e002713.
716
717 Vareed SK, Kakarala M, Ruffin MT, Crowell JA, Normolle DP, Djuric Z, Brenner DE.
718 Pharmacokinetics of curcumin conjugate metabolites in healthy human subjects. Cancer
719 Epidemiol Biomarkers Prev. 2008 Jun;17(6):1411-7.
720
721 Lee MJ, Maliakal P, Chen L, Meng X, Bondoc FY, Prabhu S, Lambert G, Mohr S, Yang CS.
722 Pharmacokinetics of tea catechins after ingestion of green tea and (-)-epigallocatechin-3-

723 gallate by humans: formation of different metabolites and individual variability. Cancer
724 Epidemiol Biomarkers Prev. 2002 Oct;11(10 Pt 1):1025-32.
725
726 Walle T, Otake Y, Brubaker JA, Walle UK, Halushka PV. Disposition and metabolism of the
727 flavonoid chrysin in normal volunteers. Br J Clin Pharmacol. 2001 Feb;51(2):143-6.
728
729 Walle T. Bioavailability of resveratrol. Ann N Y Acad Sci. 2011 Jan;1215:915.
730
731 Dabeek WM, Marra MV. Dietary Quercetin and Kaempferol: Bioavailability and Potential
732 Cardiovascular-Related Bioactivity in Humans. Nutrients. 2019 Sep 25;11(10):2288.
733
734 Boadi, W. Y., Amartey, P. K., & Lo, A. (2016). Effect of quercetin, genistein and kaempferol
735 on glutathione and glutathione-redox cycle enzymes in 3T3-L1 preadipocytes. Drug and
736 chemical toxicology, 39(3), 239–247.
737
738 Weiwei Du, Yuanlong An, Xiangdong He, Donglei Zhang, Wei He, "Protection of Kaempferol
739 on Oxidative Stress-Induced Retinal Pigment Epithelial Cell Damage", Oxidative Medicine
740 and Cellular Longevity, vol. 2018, Article ID 1610751.
741
742 Khalid S. Al-Numair, Govindasamy Chandramohan, Chinnadurai Veeramani & Mohammed
743 A. Alsaif (2015) Ameliorative effect of kaempferol, a flavonoid, on oxidative stress in
744 streptozotocin-induced diabetic rats, Redox Report, 20:5, 198-209.
745
746 N. Hayes, M. Fogarty, L. Sadofsky, H.S. Jones. Characterization of Reactive Oxygen
747 Species Signaling changes in a cell culture model of skeletal muscle ageing, and its
748 application to screening pharmacokinetically-relevant exposures of dietary polyphenols for
749 bioactivity. bioRxiv 2021.11.20.469396; doi: https://doi.org/10.1101/2021.11.20.469396
750
751
752

Acknowledgements

NH was funded by a University of Hull PhD studentship. We thank Dr Roger Sturmey and
Mr Andrew Gordon for their assistance with the TBARS assay. A preprint of this work has
previously been published at bioRxiv (bioRxiv 2021.11.20.469396)

Author statements

All the authors declare that there are no conflicts of interest with this research.
760

761 HSJ: Designed the study, planned experiments, analysed data, wrote the manuscript.
762 NH: Designed the study, planned experiments, analysed data, wrote the manuscript.
763 MF: Designed the study, wrote the manuscript
764 LS: Designed the study, wrote the manuscript.
765