Evolution of immune genes in island birds: reduction in population sizes can explain island syndrome

Mathilde BARTHE[1*], Claire DOUTRELANT[2], Rita COVAS[3-5], Martim MELO[3-5], Juan Carlos ILLERA[6], Marie-Ka TILAK[1], Constance COLOMBIER[1], Thibault LEROY[1,7], Claire LOISEAU[2,3a], Benoit NABHOLZ[1,8a]

[1] ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France

[2] Centre d'Ecologie Fonctionnelle et Evolutive, CNRS, Univ Montpellier, EPHE, IRD, Montpellier, France

[3] CIBIO, Research Centre in Biodiversity and Genetic Resources, InBIO Associated Laboratory, Campus Agrário de Vairão, Vairão, Portugal

[4] MHNC-UP, Natural History and Science Museum of the University of Porto, Porto, Portugal

[5] DST/NRF Centre of Excellence, FitzPatrick Institute, University of Cape Town, Rondebosch, South Africa

[6] Biodiversity Research Institute (CSIC-Oviedo University-Principality of Asturias), Oviedo University, Mieres, Spain

[7] IRHS-UMR1345, Université d'Angers, INRAE, Institut Agro, SFR 4207 QuaSaV, 49071, Beaucouzé, France

[8] Institut universitaire de France, Paris

[*] Corresponding author : mathilde.barthe.pro@gmail.com

[a] Last co-authors

Abstract

Shared ecological conditions encountered by species that colonize islands often lead to the evolution of convergent phenotypes, commonly referred to as the "island syndrome". Reduced immune functions have been previously proposed to be part of this syndrome, as a consequence of the reduced diversity of pathogens on island ecosystems. According to this hypothesis, immune genes are expected to exhibit genomic signatures of relaxed selection pressure in island species. In this study, we used comparative genomic methods to study immune genes in island species (N = 20) and their mainland relatives (N = 14). We gathered public data as well as generated new data on innate (TLR: Toll-Like Receptors, BD: Beta Defensins) and acquired immune genes (MHC: Major Histocompatibility Complex classes I and II), but also on hundreds of genes with various immune fonctions. As a control, we used a set of 97 genes, not known as involved in immune functions based on the literature, to account the increased drift effects for the lower effective population sizes in island species. We used synonymous and non-synonymous variations to estimate the selection pressure acting on immune genes. BDs and TLRs have higher ratios of non-synonymous over synonymous polymorphisms (Pn/Ps) than randomly selected control genes suggesting that they evolve under a different selection regime than non-immune related genes. However, simulations analyses show that this is unlikely to be explained by ongoing positive selection or balancing selection. For the MHC evolving under balancing selection, we used simulation to estimate the impact of population size variation. We found a significant effect of drift on immune genes of island species leading to a reduction in genetic diversity and efficacy of selection. However, the intensity of relaxed selection was not significantly different from control genes, except for MHC class II genes. These genes exhibit a significantly higher level of non-synonymous loss of polymorphism than expected assuming only drift and an evolution under frequency dependent selection, possibly due to a reduction of extracellular parasite communities on islands. Overall, our results showed that demographic effects lead to a decrease in the immune functions of island species, but the relaxed selection caused by a reduced parasite pressure may only occur in some immune genes categories.

Keywords: genetic drift, island evolution, immunity, Toll-Like Receptors, Beta-Defensins, major histocompatibility complex, molecular evolution, population genomics

Introduction

Island colonizers face new communities of competitors, predators and parasites in a small area with limited resources, which generally result in high extinction rates of colonizers (Losos and Ricklefs, 2009). Oceanic island faunas are characterized by a low species richness, coupled with high population densities for each species (MacArthur and Wilson, 1967; Warren et al., 2015) - which translates in communities with, on average, low levels of inter-specific interaction and high levels of intra-specific competition (but see Rando et al., 2010 for an example of character displacement due to competition among island finch species). These shared island characteristics are thought to underlie the evolution of convergent phenotypes, in what is called the 'island syndrome' (Baeckens and Van Damme, 2020). Convergence has been documented in multiple traits, such as size modification (dwarfism or gigantism; Lomolino, 2005), reduction of dispersal (Baeckens and Van Damme, 2020) shift towards K life history strategies (Boyce, 1984; Covas, 2012; MacArthur and Wilson, 1967), evolution of generalist traits (Blondel, 2000; Warren et al., 2015), or changes in colour and acoustic signals (Doutrelant et al., 2016; Grant, 1965).

Reduced immune function has also been hypothesized as an island syndrome trait, directly linked to reduced parasite pressure on islands (Lobato et al., 2017; Matson and Beadell, 2010; Wikelski et al., 2004). Island parasite communities are i) less diverse (Beadell et al., 2006; Illera et al., 2015; Loiseau et al., 2017; Maria et al., 2009; Pérez-Rodríguez et al., 2013), and ii) could be less virulent due to the expansion of the ecological niche expected by the theory of island biogeography. In fact, island parasites are generally more generalists than their mainland counterparts, which could lead to a reduced virulence due to the trade-off between replication capacity and resistance against host immune defenses (Garamszegi, 2006; Hochberg and Møller, 2001; Pérez-Rodríguez et al., 2013). Overall, a reduction of parasitic pressure should lead to a weakening of the immune system due to the costs of maintaining efficient immune functions (Lindström et al., 2004; Matson and Beadell, 2010; Wikelski et al., 2004). Such reduction may have important implications for the ability of these populations to resist or tolerate novel pathogens. The introduction of avian malaria in the Hawaiian archipelago, and the subsequent extinctions and population declines of many endemic species is the most emblematic example (Van Riper III et al., 1986; Wikelski et al., 2004).

Immunological parameters, such as blood leukocyte concentration, antibodies or other immune proteins (e.g. haptoglobin), hemolysis, and hemagglutination (Lee et al., 2006; Matson and

Beadell, 2010) may serve as proxies to determine population immune functions. To date, the

majority of studies that focused on island avifauna have found ambiguous results, with either

no support for a reduced immune response on island species (Beadell et al., 2007; Matson,

2006), or contrasted results, such as a lower humoral component (total immunoglobulins) on

islands, but a similar innate component (haptoglobin levels) between island and mainland

species (Lobato et al., 2017). The use of immune parameters as proxies of immune function is

fraught with difficulties (Lobato et al., 2017). The study of molecular evolution of immune

genes therefore represents an alternative strategy to tackle this question. However, it is

necessary to distinguish neutral effects, the demographic effects resulting from island

colonization, from selective ones, the potential relaxation of selection pressures due to the

changes in the pathogen community.

First, the bottleneck experienced by species during island colonization leads to a decrease in

genetic variability (Frankham, 1997). A reduced genetic diversity at loci involved in immunity

should have a direct implication on immune functions (Hale and Briskie, 2007 but see ; Hawley

et al., 2005; Spurgin et al., 2011). Second, small population sizes increase genetic drift, which

may counteract the effect of natural selection on weakly deleterious mutations (Ohta, 1992).

Several recent studies found a greater load of deleterious mutations in island species (Kutschera

et al., 2020; Leroy et al., 2021b; Loire et al., 2013; Robinson et al., 2016; Rogers and Slatkin,

2017). Finally, it is necessary to differentiate genes involved in the innate versus the acquired

immune response. The innate immune response is the first line of defense and is composed of

phagocytes, macrophages and dendritic cells. These cells allow non-specific recognition of

pathogens (Akira, 2003; Alberts et al., 2002). For example, Toll-Like Receptors (TLR;

transmembrane proteins) trigger a chain reaction leading to the production of various

substances, including antimicrobial peptides such as beta-defensins (BD) that have active

properties in pathogen cell lysis (Velová et al., 2018). On the other hand, the acquired immune

system allows a specific response, characterized by immune memory. Major

Histocompatibility Complex (MHC) genes code for surface glycoproteins that bind to antigenic

peptides, and present them to the cells of the immune system; class I and II genes ensure the

presentation of a broad spectrum of intra- and extracellular-derived peptides, respectively

(Klein, 1986). Although all these genes are directly involved in the identification and

neutralization of pathogens, previous studies found that they evolve under different selection

regimes: TLRs and BDs are under purifying selection which usually results in the selective

removal of deleterious alleles and stabilizing selection (Grueber et al., 2014; van Dijk et al.,
2008), whereas MHC genes are under balancing selection (Bernatchez and Landry, 2003).

Recent studies on birds (Gonzalez-Quevedo et al., 2015a, 2015b), amphibians (Belasen et al.,
2019), and lizards (Santonastaso et al., 2017) found that the demographic history of island
populations led to the loss of genetic variation at immune genes involved in pathogen
recognition, such as TLRs and MHC. For example, (Santonastaso et al., 2017) demonstrated
that the polymorphism pattern in MHC genes and microsatellites covary positively with island
area in *Podarcis* lizards, suggesting a dominant role for genetic drift in driving the evolution
of the MHC. Gonzalez-Quevedo, et al. (2015a) found a similar pattern comparing TLR and
microsatellite polymorphism in the Berthelot pipit, *Anthus berthelotii*, an endemic species from
Macaronesia, supporting a predominant role of genetic drift in TLR evolution. However, these
studies did not explicitly test the hypothesis of a relaxed selection pressure on islands imposed
by an impoverished parasite community. All other things being equal, it is expected that the
polymorphism pattern of a coding sequence decreases with population size (Buffalo, 2021;
Leroy et al., 2021b). Therefore, a decrease in polymorphism with population size could not be
taken as a proof of a relaxation in the selection pressure.

Here, we study a dataset of 34 bird species (20 insular and 14 mainland species; Figure 1)
combining the 24 species of Leroy et al. (2021b) and 10 newly generated by targeted-capture
sequencing (Table 1). To be able to demonstrate a change in natural selection, a traditional
approach is to contrast polymorphism of synonymous sites (Ps) with polymorphism of non-
synonymous sites (Pn). Synonymous mutations refer to mutations that do not alter amino acid
sequences, whereas non-synonymous mutations do.

Following population genetic theory, in a diploid population, $Ps = 4\,Ne\,\mu$ and $Pn = 4\,Ne\,\mu\,f$,
where Ne is the effective population size, μ is the mutation rate and f is a function that integrates
the probability of an allele to segregate at a given frequency. f depends on the distribution of
the fitness effect (DFE) of mutations (Eyre-Walker and Keightley, 2007). This distribution
scales with Ne as the fitness effect is dependent on Ne multiplied by the coefficient of selection
s (Kimura, 1962). The nearly-neutral theory predicts that the DFE includes a large proportion
of mutations with a $Ne*s$ close to 0 (Ohta, 1992). As a consequence, an increase of Ne will lead
to an increase of the fitness effect of weakly deleterious mutations, in such a way that these
mutations will be more easily removed from the population by natural selection, therefore
reducing Pn relative to Ps leading to a negative correlation between Pn/Ps and Ps (through Ne;

Welch et al., 2008). The presence of linked positively selected mutations does not change
qualitatively this relationship (Castellano et al., 2018; Chen et al., 2020 and our simulations
below).

Shifts in the parasitic community on islands are expected to have an impact on the ratio Pn/Ps
of immune genes. However, the fixation probability depends on the product $Ne\ s$, and a
variation in Ne is also expected to impact the efficacy of selection and thus the ratio Pn/Ps
across the entire transcriptome, particularly in the presence of slightly deleterious mutations
(Charlesworth and Eyre-Walker, 2008; Leroy et al., 2021b; Loire et al., 2013; Ohta, 1992). In
addition, due to their lower population sizes, island birds compared to continental species
exhibit a genome-wide reduction in genetic diversity and efficacy of selection (Kutschera et
al., 2020; Leroy et al., 2021b). Therefore, we expect a similar reduction in immune genes
diversity even without any change in the parasite pressure.

To disentangle the effect of population size from a change in parasite pressure and estimate the
impact of demography on the efficacy of selection, we randomly selected protein-coding genes
(i.e., control genes) involved in various biological functions (Fijarczyk et al., 2016; Leroy et
al., 2021b). The selection pressure acting on the randomly selected control genes is expected
to be similar between island and mainland bird species. Therefore, the variation of Pn/Ps of the
control genes is only dependent on the variation of Ne. In contrast, if a reduced parasite pressure
on islands directly impacts the evolution of immune genes, the Pn/Ps of immune genes is
expected to show a larger variation between island and continental species than the control
genes. More specifically, for genes under purifying selection, non-synonymous weakly
deleterious mutations, normally eliminated under strong selection, would be maintained,
leading to an increase of Pn/Ps. By contrast, for genes under balancing selection, non-
synonymous advantageous mutations, normally maintained in the polymorphism under strong
selection, would be fixed or eliminated leading to a decrease of Pn/Ps (Figure 2).

Figure 1: Phylogeny based on mitochondrial genes of species from the dataset reconstructed by maximum likelihood method (IQTREE model GTR+Gamma). Species names in yellow indicate island species, and in green, mainland species. Ultrafast bootstrap values are provided in the supplementary methods. Some relationships are poorly supported. Bird representations are not to scale. Photos from top to bottom : *P. major, C. caeruleus, P. trochilus, Z. borbonicus, T. pelios, F. albicollis, C. olivacea, P. acuticauda, P. grandis, F. coelebs, C. fusca, G. conirostris*. Photo credits: A. Chudý, F. Desmoulins, E. Giacone, G. Lasley, Lianaj, Y. Lyubchenko, B. Nabholz, J.D. Reynolds, K. Samodurov, A. Sarkisyan, Wimvz, Birdpics, T. Aronson, G. Lasley*,* P. Vos (iNaturalist.org); M. Gabrielli (*Zosterops borbonicus*).

Figure 2: Conceptual diagram showing the expected results under the hypothesis of a relaxation in the selection pressure of the immune genes in island species due to a change in the parasitic community. A) Genes evolving under purifying selection where control genes are randomly selected protein-coding genes. B) Genes evolving under balancing selection where controls are obtained from SLiM simulations of genes evolving under the same balancing selection but different population size. Under the hypothesis of a relaxed selection as a consequence of the reduced diversity of pathogens on island ecosystems, the difference in Pn/Ps between categories (ΔPn/Ps) is expected to be different between species' origin, leading to a statistical interaction between gene categories and origin.

Methods

Dataset

Alignments of Coding DNA Sequences (CDS) of individuals from 24 species were obtained from Leroy et al. (2021b). In addition, data for ten other species (six and four from islands and mainland, respectively) were newly generated for this study by targeted-capture sequencing. Blood samples and subsequent DNA extractions were performed by different research teams. The complete dataset consisted of 34 bird species (20 and 14 insular and mainland species respectively; Table 1; Figure 1). We filtered alignments in order to retain only files containing a minimum of five diploid individuals per site (Table 1).

Sequence enrichment was performed using MYBaits Custom Target Capture Kit targeting 21 immune genes: 10 Toll-Like receptors (TLR), 9 Beta Defensins (BD), 2 Major Histocompatibility Complex (MHC) and 97 control genes (see below). We followed the manufacturer's protocol (Rohland and Reich, 2012). Illumina high-throughput sequencing using a paired-end 150 bp strategy was performed by Novogene (Cambridge, UK).

Table 1: List of species and sampling localities, along with the type of data obtained and the number of individuals (N).

Species	Origin	Island/Country	N	Reference genome	Reference for population genomics data	Type of data
Cyanistes teneriffae palmae	Island	La Palma	15	*Cyanistes caeruleus* (This study)	(Mueller et al., 2016)	Capture
Cyanistes teneriffae teneriffae	Island	Tenerife	14			
Cyanistes caeruleus	Mainland	France	15			
Parus major	Mainland	Europe	10	*Parus major* (Laine et al., 2016)	(Corcoran et al., 2017)	Whole genome
Phylloscopus trochilus	Mainland	Europe	9	*Phylloscopus trochilus* (Lundberg et al., 2017)	(Lundberg et al., 2017)	Whole genome
Zosterops virens	Mainland	South Africa	7	*Zosterops borbonicus* (Leroy et al., 2021a)	(Leroy et al., 2021b)	Whole genome
Zosterops olivaceus	Island	Réunion	15			
Zosterops mauritianus	Island	Mauritius	9			
Zosterops borbonicus	Island	Réunion	25			
Ficedula semitorquata	Mainland	Europe	20	*Ficedula albicollis* (Ellegren et al., 2012)	(Ellegren et al., 2012)	Whole genome
Ficedula albicollis	Mainland	Europe	20			
Ficedula speculigera	Mainland	Nord Africa	20			
Ficedula hypoleuca	Mainland	Europe	20			
Turdus olivaceofuscus	Island	São Tomé	15	*Turdus pelios* (This study)	This study	Capture
Turdus pelios	Mainland	Gabon	15			
Cyanomitra olivacea	Island	Príncipe	15	*Cyanomitra olivacea* (This study)	This study	Capture
Cyanomitra olivacea	Mainland	Gabon	15			
Ploceus grandis	Island	São Tomé	13	*Ploceus cucullatus* (This study)	This study	Capture
Ploceus princeps	Island	Príncipe	13			
Ploceus nigerrimus	Mainland	Cameroon Gabon	14			
Poephila acuticauda acuticauda	Mainland	Australia	10	*Taeniopygia guttata* (Warren et al., 2010)	(Singhal et al., 2015)	Whole genome
Taeniopygia guttata castanotis	Mainland	Australia	19			
Fringilla teydea	Island	Tenerife	10	*Fringilla coelebs* (Recuerda et al., 2021)	(Leroy et al., 2021b)	Whole genome
Fringilla canariensis palmae	Island	La Palma	15			
Fringilla coelebs	Mainland	Spain	9			
Certhidea olivacea	Island	Santiago (Galápagos)	5	*Geospiza fortis* (Zhang et al., 2012)	(Lamichhaney et al., 2015)	Whole genome
Certhidea fusca	Island	San Cristobal (Galápagos)	10			
Certhidea fusca	Island	Española (Galápagos)	10			
Geospiza difficilis	Island	Pinta(Galápagos)	10			
Platyspiza crassirostris	Island	Santa Cruz (Galápagos)	5			
Pinaroloxias inornata	Island	Coco (Galápagos)	8			
Camarhynchus pallidus	Island	Santa Cruz (Galápagos)	5			
Geospiza difficilis	Island	Wolf (Galápagos)	8			
Geospiza conirostris	Island	Española (Galápagos)	10			

Newly generated draft genome sequence

We generated whole genome sequences at moderate coverage (~40X) for *Turdus pelios*, *Ploceus cucullatus* and *Cyanomitra olivacea* (from Gabon). Library preparation from blood

DNA samples and Illumina high-throughput sequencing using a paired-end 150 bp strategy were performed at Novogene (Cambridge, UK). Raw reads were cleaned using FastP (vers. 0.20.0; Chen et al., 2018). Genomes assemblies were performed using SOAPdenovo (vers. 2.04) and Gapcloser (v1.10) (Luo et al., 2012) with parameters "-d 1 -D 2" and a kmers size of 33. Protein annotation was performed by homology detection using genBlastG (She et al., 2011; http://genome.sfu.ca/genblast/download.html) and the transcriptome of the collared flycatcher (*Ficedula albicollis*; assembly FicAlb1.5; Ellegren et al., 2012) as reference.

Capture data processing

Reads from targeted-capture sequencing were cleaned with FastP (vers. 0.20.0; Chen et al., 2018). Reads of each individual were mapped respectively to the nearest available reference genomes using bwa mem (vers. 0.7.17; Li, 2013; Table 1), with default parameters. Samtools (vers. 1.3.1; Li et al., 2009) and Picard (vers. 1.4.2; Picard Toolkit 2019) were used to convert the mapping files, order and index reads according to their position on the chromosomes (or scaffolds) of the reference genomes or on the draft genomes generated in this study for *Ploceus*, *Cyanomitra* and *Turdus*. Duplicate reads were marked using MarkDuplicates (vers. 1.140; Picard Toolkit 2019). SNP calling was performed with Freebayes (vers. 1.3.1; Garrison and Marth, 2012). Freebayes output file (VCF file) was converted to a fasta file by filtering out sites with a minimum quality of 40 and a sequencing depth between 10 and 1000X (sites outside these thresholds were treated as missing data, i.e., 'N'). CDS were then extracted from the alignments using the coordinates of the annotations (gff files). CDS were aligned using MACSE (vers. 2.03; Ranwez et al., 2011) to prevent frameshift mutation errors and GNU-parallel (Tange, 2018) was used to parallelise the computation.

Selection and identification of immune and control genes

We defined several groups of immune genes to compare with the control genes. The control group consisted of 97 protein-coding genes randomly selected in the genome of *Zosterops borbonicus* (Leroy et al., 2021a). These control genes allowed the estimation of the average selection pressure that a gene, not involved in the immune response, undergoes in the genome. These genes are single copy (absence of paralogue) and have a variable GC content representative of the whole transcriptome.

For the immune genes, we selected three sets of genes from i) a limited set of genes (Core
Group) where functions are unambiguously related to immunity, and ii) two larger sets of genes
(Database-group & Sma3s-group), obtained through an automatic annotation pipeline.

The Core Group included MHC class I and class II genes, 10 Toll-Like Receptors (TLRs;
Velová et al., 2018) and 9 Beta Defensins (BD; Chapman et al., 2016). The Database group
included genes identified by Immunome Knowledge Base (Ortutay and Vihinen, 2009,
http://structure.bmc.lu.se/idbase/IKB/; last access 04/02/2020) and InnateDB (Breuer et al.,
2013, http://www.innatedb.com ; last access 04/02/2020). We also added a set of genes for
which the genetic ontology indicated a role in immune functions. To do so, we used the chicken
(*Gallus gallus*) annotation (assembly GRCg6a downloaded from Ensembl database in March
2020; https://www.ensembl.org/). We identified genes with the terms "immun*" or
"pathogen*" in their Gene Ontology identifiers description (directory obtained from
http://geneontology.org/). This set included 2605 genes considered to be involved in immunity,
although some may be only indirectly involved in immunity or have a small impact on immune
functions. Finally, the third set of genes (Sma3s-group) has been built up through the Sma3s-
group program (vers. 2; Munoz-Mérida et al., 2014). This program annotated sequences in
order to be associated with biological functions through gene ontology identifiers. The
annotation of the genome of *F. albicollis* allowed us to identify 3136 genes associated with the
genetic ontology "immune system processes". Like for the Database group, this set may include
genes with various functions in the immune response. It should be noted that Sma3s-group and
Database-group are not mutually exclusive, and some genes are present in both groups. An
analysis was performed to identify and exclude genes under balancing selection from Database-
group and Sma3s-group sets using BetaScan (vers. 2; Siewert and Voight, 2020), due to the
potentially antagonistic responses of these genes. Very few genes (only 2 and 3 genes from
Database-group and Sma3s-group sets) were identified and removed from the analysis (see
Detection of genes under balancing selection in Supplementary Methods).

Test for contamination and population structure

We use the program CroCo (vers. 1.1; Simion et al., 2018) to identify candidates for cross-
species contamination (see supplementary materials for details). Overall, we did not detect a
clear case of cross-species contamination in our dataset (Figure S1). Contigs identified as
potential contamination always involve a pair of species belonging to the same genus. In this

case, contamination could be difficult to identify due to the low genetic divergence between
species.

For the newly sequences species, we also performed PCA analyses on using allele frequencies
of control genes. We use the function dudi.pca of adegenet R packages (Jombart and Ahmed,
2011). This analysis aims to check for population structure and to detect potentially
problematic individuals (i.e., contaminated individuals). This analysis led to the exclusion of 4
individuals (*Ploceus princeps* P6-174; *P. grandis* ST10_094; *P. nigerrimus* G3_016; *C.*
teneriffae TF57) for which we suspected contamination. Otherwise, no extra population
structure was detected (Figure S2-S4).

Hidden paralogy

We compute the statistic Fis = 1-H0/He where H0 is the average number of heterozygous
individual observed (H0 = #heterozygous / n ; where n is the sample size) and He is the
expected number of heterozygous individuals at Hardy-Weinberg (HW) equilibrium (He =
(n/(n-1) 2 * p * (1-p))*n where n is the sample size and p the allele frequency of a randomly
chosen allele). Fis varies between -1 and 1 with positive value representing excess of
homozygous individuals and negative value representing excess of heterozygous individuals
compared to the HW proportions. Gene with high value of nucleotide diversity (Pi) and
negative value of Fis could represent potential cases where hidden paralogous sequences have
not been separated and where all the individuals present heterozygous sites in the positions
where a substitution occurred between the paralogous copies. Five TLR21 genes appear
problematic (Pi > 0.01 and Fis < -0.5; Figure S5) and were excluded from further analyses.

The MHC genes are more difficult to analyse. Indeed, heterozygosity could be comparable to
divergence under balancing selection. This makes the identification of orthologs very difficult.
We identify a variable number of genes among species (from 1 to 10 genes for MHC class I
and MHC class II). We checked the sequence similarity for the 10 copies of the MHC class II
in *F. albicollis* and the 7 copies of the MHC classI genes in *C. caeruleus* using cd-hit (Fu et
al., 2012). For MHC class II, sequence divergences are always higher than 15% indicating that
reads will likely be correctly assigned to their corresponding gene copy. For MHC class I,
sequence identities could be as high as 95%. In this case, we rely on the fact that the reads from
very similar paralogous copies will not be confidently assigned to a gene copy sequence by the
mapping software. This will lead to a low mapping score quality and are likely to be discarded

during the genotype calling procedure. For example, 3 out of 7 genes of the *Cyanistes* MHC
class I genes could not have been correctly genotyped and are missing from our final dataset.

Data Analysis

SLiM simulations

We use SLiM (vers. 3.3.2; Haller and Messer, 2017) to estimate the impact of demographic
changes on polymorphism patterns under various selection regimes. The following parameters
were used in all simulations. Sequences of 30kb with a mutation of $4.6e^{-9}$
substitutions/site/generation were simulated (Smeds et al., 2016). Recombination was set to be
equal to mutation rate. Introns/exons pattern was reproduced by simulating fragments of 3kb
separated by one bp with a very high recombination rate of 0.1 rec./site/generation. Five types
of mutations were possible: i) neutral synonymous mutations, ii) non-synonymous mutations
with a Distribution of Fitness Effect (DFE) following a gamma law of mean = -0.025 and shape
= 0.3, which corresponds to the DFE estimated in Passerines by Rousselle et al. (2020), iii)
codominant non-synonymous mutations positively selected with s = 0.1, iv) non-synonymous
mutations under balancing selection with an effect on fitness initially set at 0.01 but re-
estimated by the program at each generation according to the mutation frequency in the
population, thus including a frequency-dependent effect and v) non-synonymous mutations
under overdominance using with a dominance coefficient of 1.2.

We simulate a coding sequence organization where positions one and two of the codons were
considered as non-degenerated sites, with the non-synonymous types of mutations previously
described are possible in various proportions. The third position was considered as completely
neutral where only synonymous mutations could appear.

In the absence of control genes evolving under balancing selection, we use SLiM to generate a
set of control genes for this category. We simulate two populations of 270,000 and 110,000
individuals, representing mainland and island effective population size respectively.

We also explore the effect of positive and balancing selection on the pattern of Ps and Pn/Ps in
a population of size 50,000, 110,000, 270,000 and 500,000. In order to speed up the
computational time, we reduce the population size by a factor 100 and rescale mutation rate,
recombination rate and selection coefficient accordingly.

All the details of the simulation parameters, calculations of non-synonymous polymorphism

rate (Pn) and synonymous polymorphism rate (Ps) of simulated sequences, as well as SLiM

command lines are provided in Supplementary Methods and Supplementary Materials.

Polymorphism analyses

Synonymous (Ps) and non-synonymous (Pn) nucleotide diversity were estimated from

seq_stat_coding written from the Bio++ library (Available as Supplementary data; Guéguen et

al., 2013). The mean Pn/Ps was computed as the sum of Pn over the sum of Ps (Wolf et al.,

2009). Ps of concatenated sequences of control genes were estimated for each species of our

dataset. For the whole-genome sequence species, we compared the Pn/Ps and Ps estimated

obtained using the 97 control genes with the values from Leroy et al., (2021b; ~5000 genes

used in their study). Pn/Ps and Ps correlations showed a R^2 of 0.6 and 0.95 respectively (Figure

S6). Thus, the 97 control genes used in our study were representative of a larger set. This

allowed us to identify *Phylloscopus trochilus* as an outlier. Unlike for all other species (e.g.

Fringilla coelebs, Figure S7), synonymous polymorphism level was correlated to the amount

of missing data in *P. trochilus* alignments (Figure S7). As such, we excluded *P. trochilus* from

further analysis.

The mean Pn/Ps, calculated from the concatenated sequences of genes from the same gene

class (control genes; BD; TLR; MHC I; MHC II; Database-group; Sma3s-group), was

estimated for each bird species. Alternative transcripts were identified based on the genomic

position in the GFF file. If several transcripts were available, one transcript was randomly

selected. Pn/Ps estimates based on less than four polymorphic sites were excluded from the

analysis, as were those with no polymorphic non-synonymous sites.

Statistical analyses

To estimate the impact of demographic history on genome-wide polymorphism of island

species and the potentially reduced constraints on their immune genes, we computed the ratio

of non-synonymous nucleotide diversity over synonymous nucleotide diversity (Pn/Ps). A

linear mixed model was performed, using the Pn/Ps ratio as dependent variable and, as

explanatory variables, the mainland or insular origin of species as well as the category of genes

(packages lme4 and lmerTest (Bates et al., 2012; Kuznetsova et al., 2017)). In order to take the

phylogenetic effect into account, the taxonomic rank "family" was included as a random effect

in the model. We also used a generalized linear mixed model (using the function glmer of the

package lme4) with the family "Gamma(link="log")" which leads to the same results (Figure S15 to S24). Five linear mixed models were defined i) null model, ii) model with only the origin parameter, iii) model with only the gene category parameter, iv) model using both origin and gene category parameters, and finally v) model including those two parameters and the interaction effect. In some cases, the phylogenetic effect was difficult to estimate because the number of species per family was reduced to one. In that case, we chose to reduce the number of families by grouping Turdidae with Muscicapidae, Nectariniidae, and Estrildidae with Ploceidae and Fringillidae within Thraupidae. The results obtained with these family groupings were similar to the original model (Table S1), except when stated. The categories Database-group and Sma3s-group were tested separately from the Core group because they contained hundreds of genes annotated using the automatic pipeline that were only available for species with genome wide data. Database-group and Sma3s-group were not analysed simultaneously because they contained a partially overlapping set of genes. Finally, genes evolving under purifying selection and genes evolving under balancing selection were also analysed separately. Model selection was based on two methods. First, we use the difference in corrected Akaike Information Criterion (ΔAICc) calculated using the qpcR package (Spiess and Spiess, 2018). Second, a model simplification using an ANOVA between models was also performed.

We also tested an alternative model using the difference between Pn/Ps of immune genes and control genes (ΔPn/Ps) as dependent variable, and species origin as explanatory variable. Under the hypothesis of a relaxation in selection pressure on islands due to a change in the parasite community, we expect the ΔPn/Ps to be higher on island species compared to the mainland ones and, therefore, the species origin (i.e., mainland or island) to be significant. In this model, we used the Phylogenetic Generalized Least Squares model (PGLS; implemented in the "nlme" packages; Pinheiro et al., 2017). This model assumes that the covariance between species follows a Brownian motion evolution process along the phylogeny (implemented using the "corBrownian" function from the '"ape"' package; Paradis and Schliep, 2019). The species phylogeny was estimated using mitochondrial genes and a maximum likelihood inference implemented in IQTREE (model GTR+Gamma and ultrafast bootstrap; Nguyen et al., 2014; median of 11,134 bp analysed per species). The phylogeny with the bootstrap support is provided as supplementary material.

384   All the statistical analyses were performed using R (R Core Team, 2018), and dplyr package

385   (Wickham, 2016). Graphical representations were done using ggplot2, ggrepel, ggpubr and

386   ggpmisc (Aphalo, 2020; Kassambara, 2018; Slowikowski et al., 2018; Wickham, 2016).

Results

For the 150 individuals (10 species with 15 individuals each) for which we generated new data

by targeted capture sequencing, an average of 3.3 million paired-ends reads per individual was

generated (Table S1). After mapping, genotyping and cleaning, we analysed 86 control and 16

immune genes on average per species, out of the 141 targeted genes (120 control and 21

immune related genes; Table S4). For the species with whole-genome sequences, we analysed

106 control and 20 immune genes on average per species, out of the 141 targeted genes, and

875 and 688 genes on average in the Database-group and Sma3s-group respectively (Table S4).

For the species for which full genome sequences were available, the Ps and Pn/Ps estimated

using the control genes reflect the Ps and Pn/Ps of the whole transcriptome (Figure S6).

Population genetics of BD and TLR Immune genes

In order to characterize the selection regimes shaping the BD and TLR polymorphisms, we
first analyze the variation of Pn/Ps ratios among gene categories using a linear mixed model.

Figure 3: Pn/Ps according to species origin (mainland in green and insular in orange) for

different gene categories under purifying selection. The number of species (N), and the mean

Pn/Ps are shown for each modality.

Model selection based on AICc as well as model selection approach based on simplification

with ANOVA identified the model including the origin (i.e., mainland or island) and gene

category without interaction (Table 2). In this model, island origin of species is associated with

a greater Pn/Ps (0.14 vs. 0.10; Table 3; p < 0.01). Gene categories corresponding to TLRs and

BDs showed a significantly higher Pn/Ps than control genes (Table 3; p < 0.001). Our statistical

analysis confirms that island birds have a higher Pn/Ps ratio than mainland relatives, in

agreement with the nearly-neutral theory of evolution. It also reveals that immune genes have

a higher Pn/Ps than randomly selected control genes suggesting that BD and TLR evolve under

a different selection regime than non-immune related genes.

Next, we investigate the cause of the higher Pn/Ps of immune genes by testing three hypotheses.

First, we exclude a bias due to a lower number of genes in immune genes, and therefore higher

variance in the estimation of Pn/Ps, in immune genes. Immune genes still have significantly

higher Pn/Ps compared to a random subsample of control genes of comparable size (Figure S8

& S9). Second, the Pn/Ps of immune genes could be inflated by positive selection. It is well

known that immune genes are subject to frequent adaptation due to harm race evolution with

pathogens (Enard et al., 2016; Shultz and Sackton, 2019; Velová et al., 2018). We evaluate the

effect of positively selected genes on the Pn/Ps using SLiM simulations with both positively

and negatively selected mutations. The presence of recurrent positive selection could increase

the Pn/Ps leading to a higher Pn/Ps in immune genes if this category is more prone to adaptive

evolution (Figure 4A). However, positive selection always leads to a drastic decrease in Ps due

to genetic sweep effect at linked sites (Figure 4B). BDs and TLRs have a slightly higher or

similar Ps than control genes (Figure S9, mean Ps = 0.007, 0.004 and 0.003 for BDs, TLRs and

control genes respectively, effect of gene category p < 0.1) and, as a consequence, even if

positive selection is likely to have impacted the evolution of immune genes, it is not the cause

of the higher Pn/Ps observed here. Third, balancing selection could be present, at least

temporarily, in the evolution of BDs and TLRs genes (Kloch et al., 2018; Levy et al., 2020).

Simulation analyses confirm that balancing selection causes an increase of Ps and Pn/Ps (Figure

4C & 4D). However, a change in effective population size has an opposite effect on the Pn/Ps

according to the type of selection. In the presence of slightly deleterious mutations, Pn/Ps

decreases with *Ne* whereas it increases in the presence of balancing selection. Island birds have

higher Pn/Ps ratios than mainland birds for BDs and TLRs. Therefore, we can rule out

balancing selection as the main factor explaining the high Pn/Ps of immune genes because, in

this case, Pn/Ps of island birds should be lower. The last possible explanation we can think of

is a relaxed selection of immune genes. It is likely that immune genes are overall less

constrained than the control genes. It has been shown that evolutionary constraints are more

related to gene expression than to function (Drummond et al., 2005; Drummond and Wilke,

2008) and therefore, functionally important genes could still have a high Pn/Ps.

Overall, our analyses do not support a strong impact of ongoing adaptive mutation or balancing

selection on BDs and TLRs. However, these immune genes do not evolve as the random genes

not involved in immune functions and present a significantly higher Pn/Ps of 0.20 (p < 0.001 ;

Table 3).

No evidence of a reduced impact of the parasite communities on the polymorphism pattern

immunes genes in island birds

For BDs and TLRs, the best model selected includes the origin (i.e., mainland or island) and

gene category without interaction (see above and Table 2). This model has no interaction

between origin and gene categories invalidating the hypothesis of a reduced parasite

communities on island (Figure 2).

<u>Figure 4:</u> Neutral polymorphism (Ps) and ratio of selected over neutral polymorphism (Pn/Ps)

estimated from SLiM simulations. A) Pn/Ps as a function of population size, N and B) Ps as a

function of N. In both A and B, colour indicates the frequency of positively selected mutation compare to deleterious mutation. C) Pn/Ps as a function of N and D) Ps as a function of N. In both C and D, yellow indicates simulations with overdominance mutation (h = 1.2) and negatively selected mutations and green indicates simulations with only negatively selected mutations.

Table 2: Statistical model explaining Pn/Ps variation of Toll-Like Receptors, Beta-Defensins genes, and control genes. The p-values of ANOVA test between simpler models are not reported if a more complex model explains a larger proportion of the variance.

Model		Model selection by AIC			ANOVA test			
n°	Details	AICc	ΔAICc	Likelihood	n° 1	2	3	4
1	Pn/Ps~ 1+ category +origin+ category *origin	-5.39	8.83	0.01		0.63		
2	Pn/Ps~ 1+ category +origin	-14.22	0	1			0.002	3.71E-05
3	Pn/Ps~1+ category	-11.8	2.42	0.3				
4	Pn/Ps~1+ origin	-6.83	7.39	0.02				
5	Pn/Ps~1	-6.44	7.78	0.02				

Table 3: Summary of the best statistical model selected using AICc explaining variation in Pn/Ps in control genes, Toll-Like receptors and Beta-Defensins genes under purifying selection with origin, gene category parameters. * indicates significances : * < 0.05; ** < 0.01; *** < 0.001.

Model	Parameters		Estimate	P.value	
	Origin	Category			
Origin Intercept *mainland*		*Control genes*	0.10	2.65E-02	*
and Gene	*island*		0.14	4.56E-03	**
category		*Toll-Like Receptors*	0.20	7.43E-05	***
(n°2)		*Beta-Defensins genes*	0.20	3.16E-04	***

For larger sets of genes, identified using an automatic pipeline and gene annotation, model selection based on AICc and simplification with ANOVA (Table S5, S8) identified models that included origine parameters which associated a higher Pn/Ps of at least 0.07 for island species (p < 0.001; Table S6, S7, S9, S10, Figure 5). Selection model by simplification with

ANOVA identified models with interaction effect between origin and gene category associated
with a reduced Pn/Ps for immune genes of island species that invalidate our hypothesis (Table
S7, S10).

The alternative statistical approach using the difference between Pn/Ps of immune genes and
control genes (ΔPn/Ps) as dependent variable, and species origin as explanatory variable under
a PGLS framework lead to similar results. Island was never associated to a statistically higher
ΔPn/Ps (table S2) providing no support for an increased relaxed selection of immune genes in
island species.

Figure 5: Boxplot of Pn/Ps according to species origin (mainland in green and insular in orange)
for different gene categories under purifying selection. The number of individuals (N), and the
mean Pn/Ps are shown for each modality.

Genes under balancing selection

First, we estimated the effect of population size variation on the Pn/Ps of the genes evolving
under balancing selection by simulating sequences under frequency dependent or
overdominance selection using SLiM (see Methods and Supplementary Methods). The
simulation under frequency dependence selection revealed an average Pn/Ps equal to 0.8 for
island species and 1.2 for mainland species (Figure 6). Under overdominance, simulated
sequences from island and mainland populations respectively have an average Pn/Ps equal to
0.54 and 1.03 (Figure 6).

<u>Figure 6</u>: Boxplot of Pn/Ps according to species origin (mainland in green and insular in orange)
for different gene categories under balancing selection. The number of species (N), and the
mean Pn/Ps are shown for each modality. The control groups correspond to the results obtained
from simulated sequence via SLiM (see Methods and Supplementary Methods Simulation of
control genes under balancing selection).

Using simulations under frequency dependence selection as well as simulations under the
overdominance, model selection by AIC identifies the model with origin as the best, contrary
to the method by simplification with ANOVA which identified the full model therefore
including significant interaction between origin and genes category (Table 4). This interaction
effect is significant for the MHC II ($p < 0.05$, Table S12) but not for MHC I. As expected,
island species have a significantly lower Pn/Ps in MHC genes compared to mainland species
($p < 0.01$; except for the full model based on control genes evolving under overdominance
Table S12).

<u>Table 4</u>: Statistical model explaining Pn/Ps variation of genes under balancing selection (i.e
MHC class I and II), and simulated sequences under i) frequency dependence or ii)
overdominance. The p-values of ANOVA test between simpler models are not reported if a
more complex model explains a larger proportion of the variance.

Model			Model selection by AIC			ANOVA test			
Type of balancing selection	n°	Details	AICc	ΔAICc	Likelihood	n°1	2	3	4
Frequency	1	Pn/Ps~1+ category +origin+ category	157.17	5.62	0.06		0.019		

dependence		*origin				
	2	Pn/Ps~1+ category +origin	157.85	6.31	0.04	
	3	Pn/Ps~1+ category	187.58	36.04	0.00	
	4	Pn/Ps~1+ origin	151.54	0.00	1.00	
	5	Pn/Ps~1	180.52	28.97	0.00	
Overdominance	1	Pn/Ps~1+ category +origin+ category *origin	140,56	8,50	0,01	0.024
	2	Pn/Ps~1+ category +origin	140,56	8,50	0,01	
	3	Pn/Ps~1+ category	185,91	53,85	0,00	
	4	Pn/Ps~1+ origin	132,05	0,00	1,00	
	5	Pn/Ps~1	177,54	45,49	0,00	

Discussion

On oceanic islands, the depauperate parasite community is expected to lead to a relaxation of selection on the immune system. In this study, we found support for such an effect, but only on MHC class II genes and using simulated sequences under balancing selection as control. No effect was detected for MHC class I genes nor for innate immune genes (TLRs and BDs), evolving under purifying selection. On these gene sets, increased drift effects on island populations limit the efficacy of selection in accordance with the nearly-neutral theory (Ohta, 1992). The ability to distinguish between the selective and nearly-neutral processes (relaxed selection due to environmental change vs. drift) could only be achieved by our approach of using random genes (i.e., "control genes") to estimate the genome-wide effect of potential variation in effective population size between populations.

Effects of effective population size variation

Our results support the nearly-neutral theory of evolution for those genes under purifying
selection, whereby strong genetic drift acting on small island populations reduces the efficacy
of natural selection, leading to an increase in non-synonymous nucleotide diversity compared
to the mostly neutral, synonymous nucleotide diversity (i.e., Pn/Ps; Ohta, 1992). This is
materialized by a genome-wide increase in frequency of weakly deleterious mutations
(Kutschera et al., 2020; Leroy et al., 2021b; Loire et al., 2013; Robinson et al., 2016; Rogers
and Slatkin, 2017).

For genes evolving under balancing selection, we performed simulations under the hypotheses
of overdominance (heterozygote advantage) or frequency dependence (rare-allele advantage).
Our results showed reduced Pn/Ps for smaller population sizes (Figure 6, S10, S11). This
simulation confirmed our expectations (Figure 2) that a reduction in the efficacy of selection
results in a decrease in the frequency of non-synonymous polymorphism, as, under normal
circumstances, selection maintains those mutations at intermediate frequencies. It also matches
what we obtained in the empirical results, where both MHC classes I and II had a reduced Pn/Ps
in island birds. This result supports that the fitness effect of having non-synonymous
polymorphisms segregating at high frequencies is not strong enough to counteract entirely the
effect of genetic drift on islands, therefore extending the nearly-neutral theory to the
overdominance type of selection.

Effects of selection on immune genes

For immune genes, we try to characterize the nature of the selection acting on BDs and TLRs
genes. Comparing those genes with control genes and using simulations, we were able to rule
out that directional positive selection and balancing selection had a major impact shaping the
polymorphism of these immune genes. In contrast, the pattern of Pn/Ps between island and
mainland populations is in line with the effect of purifying selection in the presence of slightly
deleterious mutation. However, no effect was detected on insular species, beyond what could
be attributed to genetic drift. This is in line with the result of Gonzalez-Quevedo et al. (2015b)
and Grueber et al. (2013) who found that TLR genetic diversity was mostly influenced by
genetic drift. At first sight, this result seems not in line with the fact that island parasite
communities are less diverse (Beadell et al., 2006; Loiseau et al., 2017; Maria et al., 2009;
Pérez-Rodríguez et al., 2013; but see Illera et al., 2015). However, a reduced pathogens number
has also been found to be associated with a higher prevalence in birds and reptiles from the

Macaronesian archipelago (Illera and Perera, 2020). Therefore, these two patterns, i.e. a less
diverse pathogen's community on islands with a higher prevalence, could still imply a strong
selection pressure on immune genes.

In contrast, for MHC genes that unambiguously evolve under balancing selection, MHC class
II genes presented a reduction in non-synonymous polymorphism larger than the effects of drift
alone, when simulated sequences are used as control. This was the only case where a role for
relaxed selection pressures in the molecular evolution of immune genes could be invoked.

Our results are in accordance with the hypothesis of Lee (2006), which proposes that innate
and acquired immunity may exhibit distinct responses to changes in pressures due to different
costs and benefits. However, our result contrasts with the study of Santonastaso et al. (2017)
that identified no change in selection pressures on MHC II genes in a lizard species, concluding
that their evolution was mostly governed by drift. Similarly, Agudo et al. (2011) also found a
prominent role for genetic drift over selection in the evolution of MHC II genes in the Egyption
vulture (*Neophron percnopterus*).

Our results rely on simulations that may be affected by the choice of the parameter values.
First, we performed simulations using a fixed effective population size (*Ne*) estimated from the
polymorphism data. Using others values of *Ne* had a weak impact on the relative difference
between island and mainland species for the overdominance type of selection (Figure S10,
S11). Secondly, we simulated two types of selection, namely overdominance (Doherty and
Zinkernagel, 1975) and frequency dependence (Slade and McCallum, 1992), but it has been
argued that the maintenance of MHC polymorphism could be the result of fluctuating selection
(Hill, 1991). Additionally, recombination and gene conversion has also been put forward as a
mechanism responsible for generating diversity (Spurgin et al., 2011). Therefore, our results
for the MHC II, which is based on the relative difference between Pn/Ps of island and mainland
species comparing empirical and simulated data, should be taken cautiously as their
significance can be dependent on the specific parameters that we used, although we did our
best to select a realistic range of parameters.

The observed difference between MHC class I and II could be explained by their different
pathogen targets: MHC class I genes are primarily involved in the recognition of intracellular
pathogens (Kappes and Strominger, 1988), while MHC class II genes are directly involved in
the recognition of extracellular pathogens (Bjorkman and Parham, 1990). These differences

could lead to variable selection pressures depending on the extracellular versus intracellular

parasite communities present on islands. In addition, the relaxed selection pressures on MHC

II genes from adaptive immunity is in line with a reduction in acquired immunity parameters

observed by Lobato et al. (2017) that used partly the same sets of species.

As a perspective of our work, we should mention that there is an extensive variation in the

number of MHC gene copies across the avian phylogeny (Minias et al., 2019; O'Connor et al.,

2020). Particularly, it was recently discovered that Passerines have a very dynamic evolution

of duplication/loss events compared to other birds (Minias et al., 2019). Here, we used the two

copies of MHC gene I and II currently annotated in the collared flycatcher genome as target

sequences for our targeted-capture sequencing. The future improvement of genome assembly,

thanks to the development of long-reads technology (Peona et al., 2021, 2018), will certainly

help to precisely annotate all MHC copies and to study the whole repertoire of MHC genes.

Consequences of drift effect and selection on immunity

The potential relaxation of the natural selection acting on immune genes in island species is

expected to reduce immune functions and increase susceptibility of island populations to

pathogens. This is true even if this relaxation is only the consequence of a reduction in the

effective population size and not caused by a reduction of the pressure exerted by the parasitic

community. This is in line with the results of Hawley et al. (2005) and Belasen et al. (2019)

who showed that a decrease in diversity of immune loci (MHC II or through immune proxy)

was associated with a reduction in immune functions. It should be noted that even if migration

rate is reduced on islands, sedentary and endemic island species are not completely free from

the exposure of exogen pathogens through migratory birds (Levin et al., 2013).

As a final remark, we would like to stress that more research is still needed (i) to ascertain both

selection pressures on innate and adaptive immune responses and the load of deleterious

mutations due to drift, also identified by an increasing body of work (Loire et al., 2013;

Robinson et al., 2016; Rogers and Slatkin, 2017; Kutschera et al., 2020; Leroy et al., 2021b),

and (ii) to describe island parasite communities. To date, most of the studies investigated

intracellular parasite communities on islands, and more specifically haemosporidian parasites,

avian pox and coccidian parasites (Cornuault et al., 2012; Illera et al., 2015, 2008; Ishtiaq et

al., 2010; Loiseau et al., 2017; Martinez et al., 2015; Padilla et al., 2017; Pérez-Rodríguez et

al., 2013; Silva-Iturriza et al., 2012), whereas very few evaluated the extracellular parasite

diversity, such as helminths (Nieberding et al., 2006) but see the review of Illera and Perera

(2020) for reptiles. Metabarcoding of parasites is a new technique to evaluate at the same time

both communities of intracellular and extracellular parasites (Bourret et al., 2021) and might

therefore be a promising approach to evaluate their communities in island and mainland

populations.

Conclusion

Our comparative population genomics study has investigated the combined effects of drift and

selection on immune genes from island and mainland passerines. The study of synonymous

and non-synonymous polymorphism of these genes confirmed that island species, with smaller

population sizes than their mainland counterparts, were more impacted by drift, which induces

a load of weakly deleterious mutations in their genome. Indeed most of the genes studied here

involved in the immune response do not show a statistically different pattern from control

genes. Only MHC II genes, involved in the recognition of extracellular pathogens, showed a

reduction in their non-synonymous polymorphism in island species. This response, which may

be attributed to reduced selection pressures on these genes, could be associated with the

suspected reduced parasitic communities on islands. The increased load of deleterious

mutations as well as the potential relaxed selection pressures on MHC II support the reduced

immune functions of island species, which could be added to the list of other convergent

responses of the island syndrome.

Data availability

Datasets, scripts, supplementary figures and texts are available on figshare :

https://figshare.com/s/ab7004cc2f4415b4058f. The reads newly generated for this study have

been deposited in the NCBI Sequence Read Archive under the bioproject PRJNA724656.

Acknowledgments

In Gabon, we thank the Director and the guides of the Lekedi Park, Marie Charpentier for her

help in organizing the expedition, and Alexandre Vaz for field assistance and outreach work.

In São Tomé and Príncipe, we thank the Directorate of the Environment and the Department

for Nature Conservation, its directors—Arlindo Carvalho and Victor Bonfim—Guilhermino,

the Association Monte Pico, its president Luis Mário, and its members. Philippe Perret, Octávio

Veiga, Bikegila, and Yelli provided invaluable assistance in the field. Permissions for

fieldwork were given by the authorities of São Tomé and Príncipe and Gabon (CENAREST

authorization No. AR0053/12/MENESTFPRSCJS/ CENAREST/CG/CST/CSAR). Permits for

the Canary Islands were provided by the Regional Government (Ref.: 2012/0710), and the

Cabildo of La Palma and Tenerife. In Montpellier, we thank the blue tit team

(https://oreme.org/observation/ecopop/mesanges/) for the capture of the individuals used in

this study. The analyses benefited from the Montpellier Bioinformatics Biodiversity (MBB)

platform services. This research was conducted in the scope of the international twin-lab "LIA

– Biodiversity and Evolution" between CIBIO (Portugal) and ISEM and CEFE-CNRS

(France).

Funding information

This research was funded by the Labex CeMEB (project ISLAND IMMUNITY) for BN, the

ANR (BirdIslandGenomic project, ANR-14-CE02-0002) for MB, BN, CL and CD, the

National Geographic Society (Grant/Award Number:W251-12), the British Ecological Society

(Grant/Award Number: 369/4558), the portuguese Foundation for Science and Technology

under the PTDC/BIA-EVL/29390/2017 "DEEP" Research Project for MM, RC and CL and

the Spanish Ministry of Science, Innovation and Universities, and the European Regional

Development Fund (Ref.: PGC2018-097575-B-I00) for JCI.

References

Agudo, R., Alcaide, M., Rico, C., Lemus, J.A., Blanco, G., Hiraldo, F., Donázar, J.A., 2011.
Major histocompatibility complex variation in insular populations of the Egyptian
vulture: inferences about the roles of genetic drift and selection. Mol. Ecol. 20, 2329–
2340. https://doi.org/10.1111/j.1365-294X.2011.05107.x
Akira, S., 2003. Toll-like receptor signaling. J Biol Chem 278, 38105–38108.
Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K., Walter, P., 2002. Innate immunity.
Mol. Biol. Cell.
Aphalo, P.J., 2020. ggpmisc: Miscellaneous Extensions to "ggplot2"(R package version 0.3.
6.
Baeckens, S., Van Damme, R., 2020. The island syndrome. Curr. Biol. 30, R338–R339.
Bates, D.M., Maechler, M., Bolker, B., Walker, S., 2012. Package 'lme4.' CRAN R Found
Stat Comput.
Beadell, J.S., Atkins, C., Cashion, E., Jonker, M., Fleischer, R.C., 2007. Immunological
change in a parasite-impoverished environment: divergent signals from four island
taxa. PLoS One 2:e896.
Beadell, J.S., Ishtiaq, F., Covas, R., Melo, M., Warren, B.H., Atkinson, C.T., Bensch, S.,
Graves, G.R., Jhala, Y.V., Peirce, M.A., 2006. Global phylogeographic limits of
Hawaii's avian malaria. Proc R Soc Lond B Biol Sci 273, 2935–2944.
Belasen, A.M., Bletz, M.C., S, L.D., Toledo, L.F., James, T.Y., 2019. Long-term habitat
fragmentation is associated with reduced MHC IIB diversity and increased infections
in amphibian hosts. Front Ecol Evol 6.
Bernatchez, L., Landry, C., 2003. MHC studies in nonmodel vertebrates: what have we

learned about natural selection in 15 years? J Evol Biol 16, 363–377.

Bjorkman, P.J., Parham, P., 1990. Structure, function, and diversity of class I major
histocompatibility complex molecules. Annu Rev Biochem 59, 253–288.

Blondel, J., 2000. Evolution and ecology of birds on islands: trends and prospects. Vie
MilieuLife Environ. 205–220.

Bourret, V., Gutiérrez López, R., Melo, M., Loiseau, C., 2021. Metabarcoding options to
study eukaryotic endoparasites of birds. Ecol. Evol. 11, 10821–10833.

Boyce, M.S., 1984. Restitution of gamma-and k-selection as a model of density-dependent
natural selection. Annu Rev Ecol Syst 15, 427–447.

Breuer, K., Foroushani, A.K., Laird, M.R., Chen, C., Sribnaia, A., Lo, R., Winsor, G.L.,
Hancock, R.E., Brinkman, F.S., Lynn, D.J., 2013. InnateDB: systems biology of
innate immunity and beyond—recent updates and continuing curation. Nucleic Acids
Res 41:D1228–D1233.

Buffalo, V., 2021. Quantifying the relationship between genetic diversity and population size
suggests natural selection cannot explain Lewontin's paradox. eLife 10, e67509.
https://doi.org/10.7554/eLife.67509

Castellano, D., James, J., Eyre-Walker, A., 2018. Nearly neutral evolution across the
Drosophila melanogaster genome. Mol. Biol. Evol. 35, 2685–2694.

Chapman, H., JR, O, H., AS, K., RH, C., RL, W., J., 2016. The evolution of innate immune
genes: purifying and balancing selection on β-defensins in waterfowl. Mol Biol Evol
33, 3075–3087.

Charlesworth, J., Eyre-Walker, A., 2008. The McDonald–Kreitman test and slightly
deleterious mutations. Mol Biol Evol 25, 1007–1015.

Chen, J., Glémin, S., Lascoux, M., 2020. From drift to draft: how much do beneficial
mutations actually contribute to predictions of Ohta's slightly deleterious model of
molecular evolution? Genetics 214, 1005–1018.

Chen, S., Zhou, Y., Chen, Y., Gu, J., 2018. fastp: an ultra-fast all-in-one FASTQ
preprocessor. Bioinformatics 34:i884–i890.

Corcoran, P., Gossmann, T.I., Barton, H.J., Slate, J., Zeng, K., 2017. Determinants of the
Efficacy of Natural Selection on Coding and Noncoding Variability in Two Passerine
Species. Genome Biol. Evol. 9, 2987–3007. https://doi.org/10.1093/gbe/evx213

Cornuault, J., Bataillard, A., Warren, B.H., Lootvoet, A., Mirleau, P., Duval, T., Milá, B.,
Thébaud, C., Heeb, P., 2012. The role of immigration and in-situ radiation in
explaining blood parasite assemblages in an island bird clade. Mol. Ecol. 21, 1438–
1452.

Covas, R., 2012. Evolution of reproductive life histories in island birds worldwide. Proc R
Soc B Biol Sci 279, 1531–1537.

Doherty, P.C., Zinkernagel, R.M., 1975. Enhanced immunological surveillance in mice
heterozygous at the H-2 gene complex. Nature 256, 50–52.

Doutrelant, C., Paquet, M., Renoult, J.P., Grégoire, A., Crochet, P.-A., Covas, R., 2016.
Worldwide patterns of bird colouration on islands. Ecol Lett 19, 537–545.

Drummond, D.A., Bloom, J.D., Adami, C., Wilke, C.O., Arnold, F.H., 2005. Why highly
expressed proteins evolve slowly. Proc. Natl. Acad. Sci. 102, 14338–14343.

Drummond, D.A., Wilke, C.O., 2008. Mistranslation-induced protein misfolding as a
dominant constraint on coding-sequence evolution. Cell 134, 341–352.

Ellegren, H., Smeds, L., Burri, R., Olason, P.I., Backström, N., Kawakami, T., Künstner, A.,
Mäkinen, H., Nadachowska-Brzyska, K., Qvarnström, A., 2012. The genomic
landscape of species divergence in Ficedula flycatchers. Nature 491, 756–760.

Enard, D., Cai, L., Gwennap, C., Petrov, D.A., 2016. Viruses are a dominant driver of protein
adaptation in mammals. elife 5, e12469.

Eyre-Walker, A., Keightley, P.D., 2007. The distribution of fitness effects of new mutations. Nat. Rev. Genet. 8, 610–618.

Fijarczyk, A., Dudek, K., Babik, W., 2016. Selective Landscapes in newt Immune Genes Inferred from Patterns of Nucleotide Variation. Genome Biol Evol 8, 3417–3432.

Frankham, R., 1997. Do island populations have less genetic variation than mainland populations? Heredity 78, 311–327.

Fu, L., Niu, B., Zhu, Z., Wu, S., Li, W., 2012. CD-HIT: accelerated for clustering the next-generation sequencing data. Bioinformatics 28, 3150–3152.

Garamszegi, L.Z., 2006. The evolution of virulence and host specialization in malaria parasites of primates. Ecol Lett 9, 933–940.

Garrison, E., Marth, G., 2012. Haplotype-based variant detection from short-read sequencing.

Gonzalez-Quevedo, C., Phillips, K.P., Spurgin, L.G., Richardson, D.S., 2015a. 454 screening of individual MHC variation in an endemic island passerine. Immunogenetics 67, 149–162. https://doi.org/10.1007/s00251-014-0822-1

Gonzalez-Quevedo, C., Spurgin, L.G., Illera, J.C., Richardson, D.S., 2015b. Drift, not selection, shapes toll-like receptor variation among oceanic island populations. Mol. Ecol. 24, 5852–5863.

Grant, P.R., 1965. The adaptive significance of some size trends in island birds. 355–367, Evolution.

Grueber, C.E., Wallis, G.P., Jamieson, I.G., 2014. Episodic positive selection in the evolution of avian toll-like receptor innate immunity genes. PloS One 9, e89632.

Grueber, C.E., Wallis, G.P., Jamieson, I.G., 2013. Genetic drift outweighs natural selection at toll-like receptor (TLR) immunity loci in a re-introduced population of a threatened species. Mol. Ecol. 22, 4470–4482.

Guéguen, L., Gaillard, S., Boussau, B., Gouy, M., Groussin, M., Rochette, N.C., Bigot, T., Fournier, D., Pouyet, F., Cahais, V., Bernard, A., Scornavacca, C., Nabholz, B., Haudry, A., Dachary, L., Galtier, N., Belkhir, K., Dutheil, J.Y., 2013. Bio++: Efficient Extensible Libraries and Tools for Computational Molecular Evolution. Mol. Biol. Evol. 30, 1745–1750. https://doi.org/10.1093/molbev/mst097

Hale, K.A., Briskie, J.V., 2007. Decreased immunocompetence in a severely bottlenecked population of an endemic New Zealand bird. Anim. Conserv. 10, 2–10.

Haller, B.C., Messer, P.W., 2017. SLiM 2: Flexible, interactive forward genetic simulations. Mol Biol Evol 34, 230–240.

Hawley, D.M., Sydenstricker, K.V., Kollias, G.V., Dhondt, A.A., 2005. Genetic diversity predicts pathogen resistance and cell-mediated immunocompetence in house finches. Biol Lett 1, 326–329.

Hill, A.V., 1991. HLA associations with malaria in Africa: some implications for MHC evolution, in: Molecular Evolution of the Major Histocompatibility Complex. Springer, pp. 403–420.

Hochberg, M.E., Møller, A.P., 2001. Insularity and adaptation in coupled victim–enemy associations. J Evol Biol 14, 539–551.

Illera, J.C., Emerson, B.C., Richardson, D.S., 2008. Genetic characterization, distribution and prevalence of avian pox and avian malaria in the Berthelot's pipit (Anthus berthelotii) in Macaronesia. Parasitol Res 103, 1435–1443.

Illera, J.C., Fernández-Álvarez, Á., Hernández-Flores, C.N., Foronda, P., 2015. Unforeseen biogeographical patterns in a multiple parasite system in Macaronesia. J. Biogeogr. 42, 1858–1870.

Illera, J.C., Perera, A., 2020. Where are we in the host-parasite relationships of native land vertebrates in Macaronesia? Ecosistemas.

Institute, B., 2019. "Picard Toolkit", Broad institute, GitHub repository. Picard Toolkit.

Ishtiaq, F., Clegg, S.M., Phillimore, A.B., Black, R.A., Owens, I.P., Sheldon, B.C., 2010.
Biogeographical patterns of blood parasite lineage diversity in avian hosts from
southern Melanesian islands. J. Biogeogr. 37, 120–132.
Jombart, T., Ahmed, I., 2011. adegenet 1.3-1: new tools for the analysis of genome-wide SNP
data. Bioinformatics 27, 3070–3071.
Kappes, D., Strominger, J.L., 1988. Human class II major histocompatibility complex genes
and proteins. Annu Rev Biochem 57, 991–1028.
Kassambara, A., 2018. ggpubr:"ggplot2" based publication ready plots. R Package Version
01, 7.
Kimura, M., 1962. On the Probability of Fixation of Mutant Genes in a Population. Genetics
47, 713–719.
Klein, J., 1986. Natural history of the major histocompatibility complex. Wiley.
Kloch, A., Wenzel, M.A., Laetsch, D.R., Michalski, O., Bajer, A., Behnke, J.M., Welc-
Fałęciak, R., Piertney, S.B., 2018. Signatures of balancing selection in toll-like
receptor (TLRs) genes–novel insights from a free-living rodent. Sci. Rep. 8, 1–10.
Kutschera, V.E., Poelstra, J.W., Botero-Castro, F., Dussex, N., Gemmell, N., Hunt, G.R.,
Ritchie, M.G., Rutz, C., Wiberg, R.A.W., Wolf, J.B.W., 2020. Purifying Selection in
Corvids Is Less Efficient on Islands. Mol. Biol. Evol.
https://doi.org/10.1093/molbev/msz233
Kuznetsova, A., Brockhoff, P.B., Christensen, R.H., 2017. lmerTest package: tests in linear
mixed effects models. J Stat Softw 82, 1–26.
Laine, V.N., Gossmann, T.I., Schachtschneider, K.M., Garroway, C.J., Madsen, O.,
Verhoeven, K.J., De Jager, V., Megens, H.-J., Warren, W.C., Minx, P., 2016.
Evolutionary signals of selection on cognition from the great tit genome and
methylome. Nat. Commun. 7, 1–9.
Lamichhaney, S., Berglund, J., Almén, M.S., Maqbool, K., Grabherr, M., Martinez-Barrio,
A., Promerová, M., Rubin, C.-J., Wang, C., Zamani, N., 2015. Evolution of Darwin/'s
finches and their beaks revealed by genome sequencing. Nature 518, 371–375.
Lee, J.W., Beebe, K., Nangle, L.A., Jang, J., Longo-Guess, C.M., Cook, S.A., Davisson,
M.T., Sundberg, J.P., Schimmel, P., Ackerman, S.L., 2006. Editing-defective tRNA
synthetase causes protein misfolding and neurodegeneration. Nature 443, 50–55.
https://doi.org/10.1038/nature05096
Lee, K.A., 2006. Linking immune defenses and life history at the levels of the individual and
the species. Integr Comp Biol 46, 1000–1015.
Leroy, Anselmetti, Y., Tilak, M.-K., Bérard, S., Csukonyi, L., Gabrielli, M., Scornavacca, C.,
Milá, B., Thébaud, C., Nabholz, B., 2021a. A bird's white-eye view on avian sex
chromosome evolution. Peer Community J. 1.
Leroy, Rousselle, M., Tilak, M.-K., Caizergues, A.E., Scornavacca, C., Recuerda, M., Fuchs,
J., Illera, J.C., De Swardt, D.H., Blanco, G., 2021b. Island songbirds as windows into
evolution in small populations. Curr. Biol. 31, 1303-1310. e4.
Levin, I.I., Zwiers, P., Deem, S.L., Geest, E.A., Higashiguchi, J.M., Iezhova, T.A., Jiménez-
Uzcátegui, G., Kim, D.H., Morton, J.P., Perlut, N.G., Renfrew, R.B., Sari, E.H.R.,
Valkiunas, G., Parker, P.G., 2013. Multiple Lineages of Avian Malaria Parasites
(Plasmodium) in the Galapagos Islands and Evidence for Arrival via Migratory Birds.
Conserv. Biol. 27, 1366–1377. https://doi.org/10.1111/cobi.12127
Levy, H., Fiddaman, S.R., Vianna, J.A., Noll, D., Clucas, G.V., Sidhu, J.K., Polito, M.J.,
Bost, C.A., Phillips, R.A., Crofts, S., 2020. Evidence of pathogen-induced
immunogenetic selection across the large geographic range of a wild seabird. Mol.
Biol. Evol. 37, 1708–1726.
Li, H., 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-

MEM. ArXiv Prepr. ArXiv13033997.
Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,
G., Durbin, R., 2009. The Sequence Alignment/Map format and SAMtools.
Bioinforma Oxf Engl 25, 2078–2079.
Lindström, K.M., Foufopoulos, J., Pärn, H., Wikelski, M., 2004. Immunological investments
reflect parasite abundance in island populations of Darwin's finches. Proc R Soc Lond
B Biol Sci 271, 1513–1519.
Lobato, E., Doutrelant, C., Melo, M., Reis, S., Covas, R., 2017. Insularity effects on bird
immune parameters: A comparison between island and mainland populations in West
Africa. Ecol. Evol. 7, 3645–3656.
Loire, E., Chiari, Y., Bernard, A., Cahais, V., Romiguier, J., Nabholz, B., Lourenço, J.M.,
Galtier, N., 2013. Population genomics of the endangered giant Galapagos tortoise.
Genome Biol. 14, R136. https://doi.org/10.1186/gb-2013-14-12-r136
Loiseau, C., Melo, M., Lobato, E., Beadell, J.S., Fleischer, R.C., Reis, S., Doutrelant, C.,
Covas, R., 2017. Insularity effects on the assemblage of the blood parasite community
of the birds from the Gulf of Guinea. J. Biogeogr. 44, 2607–2617.
Lomolino, M.V., 2005. Body size evolution in insular vertebrates: generality of the island
rule. J. Biogeogr. 32, 1683–1699. https://doi.org/10.1111/j.1365-2699.2005.01314.x
Losos, J.B., Ricklefs, R.E., 2009. Adaptation and diversification on islands. Nature 457, 830–
836.
Lundberg, M., Liedvogel, M., Larson, K., Sigeman, H., Grahn, M., Wright, A., Åkesson, S.,
Bensch, S., 2017. Genetic differences between willow warbler migratory phenotypes
are few and cluster in large haplotype blocks. Evol Lett 1, 155–168.
Luo, R., Liu, B., Xie, Y., Li, Z., Huang, W., Yuan, J., He, G., Chen, Y., Pan, Q., Liu, Y.,
2012. SOAPdenovo2: an empirically improved memory-efficient short-read de novo
assembler. Gigascience 1.
MacArthur, R.H., Wilson, E.O., 1967. The theory of island biogeography, in: The Theory of
Island Biogeography. Princeton university press.
Maria, L., Svensson, E., Ricklefs, R.E., 2009. Low diversity and high intra-island variation in
prevalence of avian Haemoproteus parasites on Barbados, Lesser Antilles.
Parasitology 136, 1121–1131.
Martinez, J., Vasquez, R.A., Venegas, C., Merino, S., 2015. Molecular characterisation of
haemoparasites in forest birds from Robinson Crusoe Island: is the Austral Thrush a
potential threat to endemic birds? Bird Conserv. Int. 25, 139–152.
Matson, K.D., 2006. Are there differences in immune function between continental and
insular birds? Proc. Biol. Sci. 273, 2267–2274.
https://doi.org/10.1098/rspb.2006.3590
Matson, K.D., Beadell, J.S., 2010. Infection, immunity, and island adaptation in birds.
Minias, P., Pikus, E., Whittingham, L.A., Dunn, P.O., 2019. Evolution of copy number at the
MHC varies across the avian tree of life. Genome Biol Evol 11, 17–28.
Mueller, J.C., Kuhl, H., Timmermann, B., Kempenaers, B., 2016. Characterization of the
genome and transcriptome of the blue tit Cyanistes caeruleus: polymorphisms, sex-
biased expression and selection signals. Mol. Ecol. Resour. 16, 549–561.
https://doi.org/10.1111/1755-0998.12450
Munoz-Mérida, A., Viguera, E., Claros, M.G., Trelles, O., Pérez-Pulido, A.J., 2014. Sma3s: a
three-step modular annotator for large sequence datasets. DNA Res 21, 341–353.
Nguyen, L.-T., Schmidt, H.A., Haeseler, A., Minh, B.Q., 2014. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol
32, 268–274.
Nieberding, C., Morand, S., Libois, R., Michaux, J., 2006. Parasites and the island syndrome:

890    the colonization of the western Mediterranean islands by Heligmosomoides polygyrus
891    (Dujardin, 1845. J Biogeogr 33, 1212–1222.
892 O'Connor, E.A., Hasselquist, D., Nilsson, J.-Å., Westerdahl, H., Cornwallis, C.K., 2020.
893    Wetter climates select for higher immune gene diversity in resident, but not
894    migratory, songbirds. Proc. R. Soc. B Biol. Sci. 287, 20192675.
895    https://doi.org/10.1098/rspb.2019.2675
896 Ohta, T., 1992. The nearly neutral theory of molecular evolution. Annu Rev Ecol Syst 23,
897    263–286.
898 Ortutay, C., Vihinen, M., 2009. Identification of candidate disease genes by integrating Gene
899    Ontologies and protein-interaction networks: case study of primary
900    immunodeficiencies. Nucleic Acids Res 37, 622–628.
901 Padilla, D.P., Illera, J.C., Gonzalez-Quevedo, C., Villalba, M., Richardson, D.S., 2017.
902    Factors affecting the distribution of haemosporidian parasites within an oceanic
903    island. Int. J. Parasitol. 47, 225–235.
904 Paradis, E., Schliep, K., 2019. ape 5.0: an environment for modern phylogenetics and
905    evolutionary analyses in R. Bioinformatics 35, 526–528.
906 Peona, V., Blom, M.P.K., Xu, L., Burri, R., Sullivan, S., Bunikis, I., Liachko, I., Haryoko, T.,
907    Jønsson, K.A., Zhou, Q., 2021. Identifying the causes and consequences of assembly
908    gaps using a multiplatform genome assembly of a bird-of-paradise. Mol Ecol Resour
909    21, 263–286.
910 Peona, V., Weissensteiner, M.H., Suh, A., 2018. How complete are "complete" genome
911    assemblies?—An avian perspective. Mol Ecol Resour 18, 1188–1195.
912 Pérez-Rodríguez, A., Ramírez, Á., Richardson, D.S., Pérez-Tris, J., 2013. Evolution of
913    parasite island syndromes without long-term host population isolation: Parasite
914    dynamics in Macaronesian blackcaps Sylvia atricapilla. Glob Ecol Biogeogr 22,
915    1272–1281.
916 Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Heisterkamp, S., Willigen, B., Maintainer, R.,
917    2017. Package 'nlme. Linear Nonlinear Mix Eff Models Version 3.
918 R Core Team, 2018. R: A language and environment for statistical computing.
919 Rando, J.C., Alcover, J.A., Illera, J.C., 2010. Disentangling Ancient Interactions: A New
920    Extinct Passerine Provides Insights on Character Displacement among Extinct and
921    Extant Island Finches. PLOS ONE 5:e12956.
922 Ranwez, V., Harispe, S., Delsuc, F., Douzery, E.J., 2011. MACSE: Multiple Alignment of
923    Coding SEquences accounting for frameshifts and stop codons. PLoS One 6:e22594.
924 Recuerda, M., Vizueta, J., Cuevas-Caballé, C., Blanco, G., Rozas, J., Milá, B., 2021.
925    Chromosome-level genome assembly of the common chaffinch (Aves: Fringilla
926    coelebs): a valuable resource for evolutionary biology. Genome Biol. Evol. 13,
927    evab034.
928 Robinson, J.A., Ortega-Del Vecchyo, D., Fan, Z., Kim, B.Y., Marsden, C.D., Lohmueller,
929    K.E., Wayne, R.K., 2016. Genomic flatlining in the endangered island fox. Curr Biol
930    26, 1183–1189.
931 Rogers, R.L., Slatkin, M., 2017. Excess of genomic defects in a woolly mammoth on
932    Wrangel island. PLoS Genet 13:e1006601.
933 Rohland, N., Reich, D., 2012. Cost-effective, high-throughput DNA sequencing libraries for
934    multiplexed target capture. Genome Res. 22, 939–946.
935 Rousselle, M., Simion, P., Tilak, M.-K., Figuet, E., Nabholz, B., Galtier, N., 2020. Is
936    adaptation limited by mutation? A timescale-dependent effect of genetic diversity on
937    the adaptive substitution rate in animals. PLoS Genet. 16, e1008668.
938 Santonastaso, T., Lighten, J., Oosterhout, C., Jones, K.L., Foufopoulos, J., Anthony, N.M.,
939    2017. The effects of historical fragmentation on major histocompatibility complex

class II β and microsatellite variation in the Aegean island reptile, Podarcis erhardii. Ecol Evol 7, 4568–4581.

She, R., Chu, J.S.-C., Uyar, B., Wang, J., Wang, K., Chen, N., 2011. genBlastG: using BLAST searches to build homologous gene models. Bioinforma Oxf Engl 27, 2141–2143.

Shultz, A.J., Sackton, T.B., 2019. Immune genes are hotspots of shared positive selection across birds and mammals. Elife 8, e41815.

Siewert, K.M., Voight, B.F., 2020. BetaScan2: Standardized Statistics to Detect Balancing Selection Utilizing Substitution Data. Genome Biol. Evol. 12, 3873–3877. https://doi.org/10.1093/gbe/evaa013

Silva-Iturriza, A., Ketmaier, V., Tiedemann, R., 2012. Prevalence of avian haemosporidian parasites and their host fidelity in the central Philippine islands. Parasitol. Int. 61, 650–657.

Simion, P., Belkhir, K., François, C., Veyssier, J., Rink, J.C., Manuel, M., Philippe, H., Telford, M.J., 2018. A software tool 'CroCo' detects pervasive cross-species contamination in next generation sequencing data. BMC Biol. 16, 1–9.

Singhal, S., Leffler, E.M., Sannareddy, K., Turner, I., Venn, O., Hooper, D.M., Strand, A.I., Li, Q., Raney, B., Balakrishnan, C.N., 2015. Stable recombination hotspots in birds. Science 350, 928–932.

Slade, R.W., McCallum, H.I., 1992. Overdominant vs. frequency-dependent selection at MHC loci. Genetics 132.

Slowikowski, K., Schep, A., Hughes, S., Lukauskas, S., Irisson, J.-O., Kamvar, Z.N., Ryan, T., Christophe, D., Hiroaki, Y., Gramme, P., 2018. Package ggrepel. Autom Position Non-Overlapping Text Labels ggplot2.

Smeds, L., Qvarnstrom, A., Ellegren, H., 2016. Direct estimate of the rate of germline mutation in a bird. Genome Res. gr-204669.

Spiess, A.-N., Spiess, M.A.-N., 2018. Package 'qpcR, in: Model. Anal. Real-Time PCRdata Httpscran R-Proj. OrgwebpackagesqpcRqpcR Pdf.

Spurgin, L.G., Van Oosterhout, C., Illera, J.C., Bridgett, S., Gharbi, K., Emerson, B.C., Richardson, D.S., 2011. Gene conversion rapidly generates major histocompatibility complex diversity in recently founded bird populations. Mol. Ecol. 20, 5213–5225. https://doi.org/10.1111/j.1365-294X.2011.05367.x

Tange, O., 2018. GNU parallel 2018.

van Dijk, A., Veldhuizen, E.J., Haagsman, H.P., 2008. Avian defensins. Vet. Immunol. Immunopathol. 124, 1–18.

Van Riper III, C., Van Riper, S.G., Goff, M.L., Laird, M., 1986. The epizootiology and ecological significance of malaria in Hawaiian land birds. Ecol. Monogr. 56, 327–344.

Velová, H., Gutowska-Ding, M.W., Burt, D.W., Vinkler, M., 2018. Toll-like receptor evolution in birds: gene duplication, pseudogenization, and diversifying selection. Mol Biol Evol 35, 2170–2184.

Warren, B.H., Simberloff, D., Ricklefs, R.E., Aguilée, R., Condamine, F.L., Gravel, D., Morlon, H., Mouquet, N., Rosindell, J., Casquet, J., 2015. Islands as model systems in ecology and evolution: Prospects fifty years after MacArthur-Wilson. Ecol. Lett. 18, 200–217.

Warren, W.C., Clayton, D.F., Ellegren, H., Arnold, A.P., Hillier, L.W., Künstner, A., Searle, S., White, S., Vilella, A.J., Fairley, S., 2010. The genome of a songbird. Nature 464, 757–762.

Welch, J.J., Eyre-Walker, A., Waxman, D., 2008. Divergence and Polymorphism Under the Nearly Neutral Theory of Molecular Evolution. J. Mol. Evol. 67, 418–426.

https://doi.org/10.1007/s00239-008-9146-9

Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis.

Wikelski, M., Foufopoulos, J., Vargas, H., Snell, H., 2004. Galápagos birds and diseases:
invasive pathogens as threats for island species. Ecol. Soc. 9.

Wolf, J.B.W., Künstner, A., Nam, K., Jakobsson, M., Ellegren, H., 2009. Nonlinear
Dynamics of Nonsynonymous (dN) and Synonymous (dS) Substitution Rates Affects
Inference of Selection. Genome Biol Evol 1, 308–319.

Zhang, G., Parker, P., Li, B., Li, H., Wang, J., 2012. The genome of Darwin's Finch
(Geospiza fortis). https://doi.org/10.5524/100040