Evolution of immune genes in island birds: reduction in population sizes can explain island syndrome

Mathilde BARTHE[1*], Claire DOUTRELANT[2], Rita COVAS[3,5], Martim MELO[3-6], Juan Carlos ILLERA[7], Marie-Ka TILAK[1], Constance COLOMBIER[1], Thibault LEROY[1,8], Claire LOISEAU[2,3,4a] , Benoit NABHOLZ[1,9a]

[1] ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France

[2] CEFE, CNRS, Univ Montpellier, EPHE, IRD, Montpellier, France

[3] CIBIO-InBio, Research Center in Biodiversity and Genetic Resources, Associated Laboratory, Campus Agrário de Vairão, Vairão, Portugal

[4] BIOPOLIS Program in Genomics, Biodiversity and Land Planning, CIBIO, Campus de Vairão, 4485-661 Vairão, Portugal

[5] FitzPatrick Institute, University of Cape Town, Rondebosch, South Africa

[6] MHNC-UP, Natural History and Science Museum of the University of Porto, Porto, Portugal

[7] Biodiversity Research Institute (CSIC-Oviedo University-Principality of Asturias), Oviedo University, Mieres, Spain

[8] IRHS-UMR1345, Université d'Angers, INRAE, Institut Agro, SFR 4207 QuaSaV, 49071, Beaucouzé, France

[9] Institut universitaire de France, Paris

[*] Corresponding author : mathilde.barthe.pro@gmail.com

[a] Last co-authors

Abstract

Shared ecological conditions encountered by species that colonize islands often lead to the

evolution of convergent phenotypes, commonly referred to as "island syndrome". Reduced

immune functions have been previously proposed to be part of this syndrome, as a consequence

of the reduced diversity of pathogens on island ecosystems. According to this hypothesis,

immune genes are expected to exhibit genomic signatures of relaxed selection pressure in

island species. In this study, we used comparative genomic methods to study immune genes in

island species (N = 20) and their mainland relatives (N = 14). We gathered public data as well

as generated new data on innate (TLR: Toll-Like Receptors, BD: Beta Defensins) and acquired

immune genes (MHC: Major Histocompatibility Complex classes I and II), but also on

hundreds of genes with various immune functions. As a control, we used a set of 97 genes, not

known to be involved in immune functions based on the literature, to account for the increased

drift effects of the lower effective population sizes in island species. We used synonymous and

non-synonymous variants to estimate the selection pressure acting on immune genes. We found

that BDs and TLRs have higher ratios of non-synonymous over synonymous polymorphisms

(Pn/Ps) than randomly selected control genes, suggesting that they evolve under a different

selection regime. However, simulations show that this is unlikely to be explained by ongoing

positive selection or balancing selection. For the MHC genes, which evolve under balancing

selection, we used simulations to estimate the impact of population size variation. We found a

significant effect of drift on immune genes of island species leading to a reduction in genetic

diversity and efficacy of selection. However, the intensity of relaxed selection was not

significantly different from control genes, except for MHC class II genes. These genes exhibit

a significantly higher level of non-synonymous loss of polymorphism than expected assuming

only drift and evolution under frequency dependent selection, possibly due to a reduction of

extracellular parasite communities on islands. Overall, our results showed that demographic

effects lead to a decrease in the immune functions of island species, but the relaxed selection

that is expected to be caused by a reduced parasite pressure may only occur in some categories

of immune genes.

Keywords: genetic drift, island evolution, immunity, Toll-Like Receptors, Beta-Defensins,

major histocompatibility complex, molecular evolution, population genomics

Introduction

Island colonizers face new communities of competitors, predators and parasites in a small area with limited resources, which generally results in high extinction rates of colonizers (Losos and Ricklefs, 2009). Oceanic island faunas are characterized by a low species richness, coupled with high population densities for each species (MacArthur and Wilson, 1967; Warren et al., 2015) - which translates into communities with, on average, lower levels of inter-specific interactions and higher levels of intra-specific competition (but see Rando et al., 2010 for an example of character displacement due to competition among island finch species). These shared island characteristics are thought to underlie the evolution of convergent phenotypes, in what is called the 'island syndrome' (Baeckens and Van Damme, 2020). Convergence has been documented in multiple traits, such as size modification (dwarfism or gigantism; Lomolino, 2005), reduction of dispersal (Baeckens and Van Damme, 2020), shift towards K life-history strategies (Boyce, 1984; Covas, 2012; MacArthur and Wilson, 1967), evolution of generalist traits (Blondel, 2000; Warren et al., 2015), or changes in colour and acoustic signals (Doutrelant et al., 2016; Grant, 1965).

Reduced immune function has also been hypothesized to be an island syndrome trait, directly linked to reduced parasite pressure on islands (Lobato et al., 2017; Matson and Beadell, 2010; Wikelski et al., 2004). Island parasite communities are i) less diverse (Beadell et al., 2006; Illera et al., 2015; Loiseau et al., 2017; Maria et al., 2009; Pérez-Rodríguez et al., 2013), and ii) could be less virulent due to the expansion of the ecological niche expected by the theory of island biogeography. In fact, island parasites are often more generalist than their mainland counterparts, which could lead to a reduced virulence due to the trade-off between replication capacity and resistance against host immune defenses (Garamszegi, 2006; Hochberg and Møller, 2001; Pérez-Rodríguez et al., 2013). Overall, a reduction of parasitic pressure should lead to a weakening of the immune system due to the costs of maintaining efficient immune functions (Lindström et al., 2004; Matson and Beadell, 2010; Wikelski et al., 2004). Such reduction may have important implications for the ability of these populations to resist or tolerate novel pathogens. The introduction of avian malaria in the Hawaiian archipelago, and the subsequent extinctions and population declines of many endemic species is the most emblematic example (Van Riper III et al., 1986; Wikelski et al., 2004).

Immunological parameters, such as blood leukocyte concentration, antibodies or other immune proteins (e.g. haptoglobin), hemolysis, and hemagglutination (Lee et al., 2006; Matson and

Beadell, 2010) may serve as proxies to determine population immune functions. To date, the
majority of studies that focused on island avifauna have found ambiguous results, with either
no support for a reduced immune response on island species (Beadell et al., 2007; Matson,
2006), or contrasting results, such as a lower humoral component (total immunoglobulins) on
islands, but a similar innate component (haptoglobin levels) between island and mainland
species (Lobato et al., 2017). The use of immune parameters as proxies of immune function is
fraught with difficulties (Lobato et al., 2017). The study of molecular evolution of immune
genes therefore represents an alternative strategy to tackle this question. However, it is
necessary to distinguish neutral effects (i.e. the demographic effects resulting from island
colonization) from selective ones, the potential relaxation of selection pressures due to the
changes in the pathogen community.

The bottleneck experienced by species during island colonization leads to a decrease in genetic
variability (Frankham, 1997). A reduced genetic diversity at loci involved in immunity should
have a direct implication on immune functions (Hale and Briskie, 2007 but see ; Hawley et al.,
2005; Spurgin et al., 2011). Also, small population sizes increase genetic drift, which may
counteract the effect of natural selection on weakly deleterious mutations (Ohta, 1992). Several
recent studies found a greater load of deleterious mutations in island species (Kutschera et al.,
2020; Leroy et al., 2021b; Loire et al., 2013; Robinson et al., 2016; Rogers and Slatkin, 2017).
Finally, it is necessary to differentiate genes involved in the innate versus the acquired immune
response. The innate immune response is the first line of defense and is composed of
phagocytes, macrophages and dendritic cells. These cells allow non-specific recognition of
pathogens (Akira, 2003; Alberts et al., 2002). For example, Toll-Like Receptors (TLR;
transmembrane proteins) trigger a chain reaction leading to the production of various
substances, including antimicrobial peptides such as beta-defensins (BD) that have active
properties in pathogen cell lysis (Velová et al., 2018). On the other hand, the acquired immune
system allows a specific response, characterized by immune memory. Major
Histocompatibility Complex (MHC) genes code for surface glycoproteins that bind to antigenic
peptides, and present them to the cells of the immune system; class I and II genes ensure the
presentation of a broad spectrum of intra- and extracellular-derived peptides, respectively
(Klein, 1986). Although all these genes are directly involved in the identification and
neutralization of pathogens, previous studies found that they evolve under different selection
regimes: TLRs and BDs are under purifying selection which usually results in the selective

removal of deleterious alleles and stabilizing selection (Grueber et al., 2014; van Dijk et al.,

2008), whereas MHC genes are under balancing selection (Bernatchez and Landry, 2003).

Recent studies on birds (Gonzalez-Quevedo et al., 2015a, 2015b), amphibians (Belasen et al.,

2019), and lizards (Santonastaso et al., 2017) found that the demographic history of island

populations led to the loss of genetic variants at immune genes involved in pathogen

recognition, such as TLRs and MHC. For example, Santonastaso et al., (2017) revealed that

the polymorphism pattern in MHC genes and microsatellites covary positively with island area

in *Podarcis* lizards, suggesting a dominant role for genetic drift in driving the evolution of the

MHC. Gonzalez-Quevedo, et al. (2015a) found a similar pattern comparing TLR and

microsatellite polymorphism in the Berthelot pipit, *Anthus berthelotii*, an endemic species from

Macaronesia, supporting a predominant role of genetic drift in TLR evolution. However, these

studies did not explicitly test the hypothesis of a relaxed selection pressure on islands imposed

by an impoverished parasite community. All other things being equal, it is expected that the

polymorphism of a coding sequence decreases with population size (Buffalo, 2021; Leroy et

al., 2021b). Therefore, a decrease in polymorphism with population size could not be taken as

a proof of a relaxation in the selection pressure.

To be able to demonstrate a change in natural selection, a traditional approach is to contrast

polymorphism of synonymous sites (Ps) with polymorphism of non-synonymous sites (Pn).

Synonymous mutations do not change amino acid sequences, whereas non-synonymous

mutations do. Thus, synonymous mutations are expected to be neutral while non-synonymous

could be subject to selection.

Following population genetic theory, in a diploid population, $Ps = 4\,Ne\,\mu$ and $Pn = 4\,Ne\,\mu\,f$,

where Ne is the effective population size, μ is the mutation rate and f is a function that integrates

the probability of an allele to segregate at a given frequency. f depends on the distribution of

the fitness effect (DFE) of mutations (Eyre-Walker and Keightley, 2007). This distribution

scales with Ne as the fitness effect is dependent on Ne multiplied by the coefficient of selection

s (Kimura, 1962). The nearly-neutral theory predicts that the DFE includes a large proportion

of mutations with a $Ne*s$ close to 0 (Ohta, 1992). As a consequence, an increase of Ne will lead

to an increase of the fitness effect of weakly deleterious mutations, in such a way that these

mutations will be more easily removed from the population by natural selection, therefore

reducing Pn relative to Ps, leading to a negative correlation between Pn/Ps and Ps (through Ne;

Welch et al., 2008). The presence of linked mutations, that are positively selected, does not

change this relationship qualitatively (Castellano et al., 2018; Chen et al., 2020 and our

simulations below).

Shifts in the parasitic community on islands are expected to have an impact on the Pn/Ps ratio

of immune genes. However, the fixation probability depends on the product $Ne*s$, and variation

in Ne is also expected to impact the efficacy of selection and thus the Pn/Ps ratio across the

entire transcriptome, particularly in the presence of slightly deleterious mutations

(Charlesworth and Eyre-Walker, 2008; Leroy et al., 2021b; Loire et al., 2013; Ohta, 1992). In

addition, due to their lower population sizes, island birds compared to continental species

exhibit a genome-wide reduction in genetic diversity and efficacy of selection (Kutschera et

al., 2020; Leroy et al., 2021b). Therefore, we expect a similar reduction in immune genes'

diversity even without any change in the parasite pressure.

To disentangle the effect of population size from a change in parasite pressure and estimate the

impact of demography on the efficacy of selection, we studied a dataset of 34 bird species (20

insular and 14 mainland species; Figure 1) combining the 24 species of Leroy et al. (2021b)

and 10 newly generated by targeted-capture sequencing (Table 1). We randomly selected

protein-coding genes (i.e., control genes) involved in various biological functions (Fijarczyk et

al., 2016; Leroy et al., 2021b). The selection pressure acting on the randomly selected control

genes is expected to be similar between island and mainland bird species. Therefore, the

variation of Pn/Ps of the control genes is only dependent on the variation of Ne. In contrast, if

a reduced parasite pressure on islands directly impacts the evolution of immune genes, the

Pn/Ps of immune genes is expected to show a larger variation between island and continental

species than the control genes. More specifically, for genes under purifying selection, non-

synonymous weakly deleterious mutations, normally eliminated under strong selection, would

be maintained, leading to an increase of Pn/Ps. By contrast, for genes under balancing selection,

non-synonymous advantageous mutations, normally maintained in the polymorphism under

strong selection, would be fixed or eliminated leading to a decrease of Pn/Ps (Figure 2).

Lanius cristatus
Menura novaehollandiae
Parus major
Cyanistes palmae
Cyanistes caeruleus
Cyanistes teneriffae
Phylloscopus trochilus
Zosterops virens
Zosterops olivaceus
Zosterops borbonicus
Zosterops mauritianus
Turdus olivaceofuscus
Turdus pelios
Ficedula semitorquata
Ficedula albicollis
Ficedula hypoleuca
Ficedula speculigera
Cyanomitra olivacea mainland
Cyanomitra olivacea island
Poephila acuticauda
Taeniopygia guttata
Ploceus nigerrimus
Ploceus grandis
Ploceus princeps
Fringilla teydea
Fringilla coelebs
Fringilla canariensis palmae
Certhidea olivacea S
Certhidea fusca E
Certhidea fusca L
Geospiza difficilis P
Platyspiza crassirostris Z
Pinaroloxias inornata C
Camarhynchus pallidus Z
Geospiza conirostris E
Geospiza difficilis W

0.03

Figure 1: Phylogeny based on mitochondrial genes of species from the dataset reconstructed by
maximum likelihood method (IQTREE model GTR+Gamma). Species names in yellow indicate island
species, and in green, mainland species. Ultrafast bootstrap values are provided in the supplementary
methods. Some relationships are poorly supported. Bird representations are not to scale. Photos from
top to bottom : *P. major, C. caeruleus, P. trochilus, Z. borbonicus, T. pelios, F. albicollis, C. olivacea,*
P. acuticauda, P. grandis, F. coelebs, C. fusca, G. conirostris. Photo credits: A. Chudý, F. Desmoulins,
E. Giacone, G. Lasley, Lianaj, Y. Lyubchenko, B. Nabholz, J.D. Reynolds, K. Samodurov, A.
Sarkisyan, Wimvz, Birdpics, T. Aronson, G. Lasley, P. Vos (iNaturalist.org); M. Gabrielli (*Zosterops*
borbonicus).

164
Figure 2: Conceptual diagram showing the expected results under the hypothesis of a relaxation in the
selection pressure of the immune genes in island species due to a change in the parasitic community.
A) Genes evolving under purifying selection where control genes are randomly selected protein-coding
genes. B) Genes evolving under balancing selection where controls are obtained from SLiM simulations
of genes evolving under the same balancing selection but different population size. Under the hypothesis
of a relaxed selection as a consequence of the reduced diversity of pathogens on island ecosystems, the
difference in Pn/Ps between categories (ΔPn/Ps) is expected to be different between species' origin,
leading to a statistical interaction between gene categories and origin.

Methods
174

Dataset
175

Alignments of Coding DNA Sequences (CDS) of individuals from 24 species were obtained

from Leroy et al. (2021b). In addition, data for ten other species (six and four from islands and

mainland, respectively) were newly generated for this study by targeted-capture sequencing.

Blood samples and subsequent DNA extractions were performed by different research teams.

The complete dataset consisted of 34 bird species (20 and 14 insular and mainland species

respectively; Table 1; Figure 1). We filtered alignments in order to retain only files containing

a minimum of five diploid individuals per site (Table 1).

Sequence enrichment was performed using MYBaits Custom Target Capture Kit targeting 21

immune genes: 10 Toll-Like receptors (TLR), 9 Beta Defensins (BD), 2 Major

Histocompatibility Complex (MHC) and 97 control genes (see below). We followed the

manufacturer's protocol (Rohland and Reich, 2012). Illumina high-throughput sequencing,

using a paired-end 150 bp strategy, was performed by Novogene (Cambridge, UK).

Table 1: List of species and sampling localities, along with the type of data obtained and the number of individuals (N).

Species	Origin	Island/Country	N	Reference genome	Reference for population genomics data	Type of data
Cyanistes teneriffae palmae	Island	La Palma	15	*Cyanistes caeruleus* (This study)	(Mueller et al., 2016)	Capture
Cyanistes teneriffae teneriffae	Island	Tenerife	14			
Cyanistes caeruleus	Mainland	France	15			
Parus major	Mainland	Europe	10	*Parus major* (Laine et al., 2016)	(Corcoran et al., 2017)	Whole genome
Phylloscopus trochilus	Mainland	Europe	9	*Phylloscopus trochilus* (Lundberg et al., 2017)	(Lundberg et al., 2017)	Whole genome
Zosterops virens	Mainland	South Africa	7	*Zosterops borbonicus* (Leroy et al., 2021a)	(Leroy et al., 2021b)	Whole genome
Zosterops olivaceus	Island	Réunion	15			
Zosterops mauritianus	Island	Mauritius	9			
Zosterops borbonicus	Island	Réunion	25			
Ficedula semitorquata	Mainland	Europe	20	*Ficedula albicollis* (Ellegren et al., 2012)	(Ellegren et al., 2012)	Whole genome
Ficedula albicollis	Mainland	Europe	20			
Ficedula speculigera	Mainland	Nord Africa	20			
Ficedula hypoleuca	Mainland	Europe	20			
Turdus olivaceofuscus	Island	São Tomé	15	*Turdus pelios* (This study)	This study	Capture
Turdus pelios	Mainland	Gabon	15			
Cyanomitra olivacea	Island	Príncipe	15	*Cyanomitra olivacea* (This study)	This study	Capture
Cyanomitra olivacea	Mainland	Gabon	15			
Ploceus grandis	Island	São Tomé	13	*Ploceus cucullatus* (This study)	This study	Capture
Ploceus princeps	Island	Príncipe	13			
Ploceus nigerrimus	Mainland	Cameroon Gabon	14			
Poephila acuticauda acuticauda	Mainland	Australia	10	*Taeniopygia guttata* (Warren et al., 2010)	(Singhal et al., 2015)	Whole genome
Taeniopygia guttata castanotis	Mainland	Australia	19			
Fringilla teydea	Island	Tenerife	10	*Fringilla coelebs* (Recuerda et al., 2021)	(Leroy et al., 2021b)	Whole genome
Fringilla canariensis palmae	Island	La Palma	15			
Fringilla coelebs	Mainland	Spain	9			
Certhidea olivacea	Island	Santiago (Galápagos)	5	*Geospiza fortis* (Zhang et al., 2012)	(Lamichhaney et al., 2015)	Whole genome
Certhidea fusca	Island	San Cristobal (Galápagos)	10			
Certhidea fusca	Island	Española (Galápagos)	10			
Geospiza difficilis	Island	Pinta(Galápagos)	10			
Platyspiza crassirostris	Island	Santa Cruz (Galápagos)	5			
Pinaroloxias inornata	Island	Coco (Galápagos)	8			
Camarhynchus pallidus	Island	Santa Cruz (Galápagos)	5			
Geospiza difficilis	Island	Wolf (Galápagos)	8			
Geospiza conirostris	Island	Española (Galápagos)	10			

Newly generated draft genome sequence

We generated whole genome sequences at moderate coverage (~40X) for *Turdus pelios*, *Ploceus cucullatus* and *Cyanomitra olivacea* (from Gabon). Library preparation from blood DNA samples and Illumina high-throughput sequencing using a paired-end 150 bp strategy were performed at Novogene (Cambridge, UK). Raw reads were cleaned using FastP (vers. 0.20.0; Chen et al., 2018). Genomes assemblies were performed using SOAPdenovo (vers. 2.04) and Gapcloser (v1.10) (Luo et al., 2012) with parameters "-d 1 -D 2" and a kmers size of

33. Protein annotation was performed by homology detection using genBlastG (She et al., 2011; http://genome.sfu.ca/genblast/download.html) and the transcriptome of the collared flycatcher (*Ficedula albicollis*; assembly FicAlb1.5; Ellegren et al., 2012) as reference.

Capture data processing

Reads from targeted-capture sequencing were cleaned with FastP (vers. 0.20.0; Chen et al., 2018). Reads of each individual were mapped respectively to the nearest available reference genomes using bwa mem (vers. 0.7.17; Li, 2013; Table 1), with default parameters. Samtools (vers. 1.3.1; Li et al., 2009) and Picard (vers. 1.4.2; Picard Toolkit 2019) were used to convert the mapping files, order and index reads according to their position on the chromosomes (or scaffolds) of the reference genomes or on the draft genomes generated in this study for *Ploceus*, *Cyanomitra* and *Turdus*. Duplicate reads were marked using MarkDuplicates (vers. 1.140; Picard Toolkit 2019). SNP calling was performed with Freebayes (vers. 1.3.1; Garrison and Marth, 2012). Freebayes output file (VCF file) was converted to a fasta file by filtering out sites with a minimum quality of 40 and a sequencing depth between 10 and 1000X (sites outside these thresholds were treated as missing data, i.e., 'N'). CDS were then extracted from the alignments using the coordinates of the annotations (gff files). CDS were aligned using MACSE (vers. 2.03; Ranwez et al., 2011) to prevent frameshift mutation errors and GNU-parallel (Tange, 2018) was used to parallelise the computation.

Selection and identification of immune and control genes

We defined several groups of immune genes to compare with the control genes. The control group consisted of 97 protein-coding genes randomly selected in the genome of *Zosterops borbonicus* (Leroy et al., 2021a). These control genes allowed the estimation of the average selection pressure that a gene, not involved in the immune response, undergoes in the genome under a given effective population size. These genes were single copy (absence of paralogue) and had a variable GC content representative of the whole transcriptome.

For the immune genes, we selected three sets of genes from i) a limited set of genes (Core Group) where functions are unambiguously related to immunity, and ii) two larger sets of genes (Database-group & Sma3s-group), obtained through an automatic annotation pipeline.

The Core Group included MHC class I and class II genes, 10 Toll-Like Receptors (TLRs; Velová et al., 2018) and 9 Beta Defensins (BD; Chapman et al., 2016). The Database group

included genes identified by Immunome Knowledge Base (Ortutay and Vihinen, 2009,

http://structure.bmc.lu.se/idbase/IKB/; last access 04/02/2020) and InnateDB (Breuer et al.,

2013, http://www.innatedb.com ; last access 04/02/2020). We also added a set of genes for

which the genetic ontology indicated a role in immune functions. To do so, we used the chicken

(*Gallus gallus*) annotation (assembly GRCg6a downloaded from Ensembl database in March

2020; https://www.ensembl.org/). We identified genes with the terms "immun*" or

"pathogen*" in their Gene Ontology identifiers description (directory obtained from

http://geneontology.org/). This set included 2605 genes considered to be involved in immunity,

although some may be only indirectly involved in immunity or have a small impact on immune

functions. Finally, the third set of genes (Sma3s-group) has been built up through the Sma3s-

group program (vers. 2; Munoz-Mérida et al., 2014). This program annotated sequences in

order to be associated with biological functions through gene ontology identifiers. The

annotation of the genome of *F. albicollis* allowed us to identify 3136 genes associated with the

genetic ontology "immune system processes". Like for the Database group, this set may include

genes with various functions in the immune response. It should be noted that Sma3s-group and

Database-group were not mutually exclusive, and some genes were present in both groups. An

analysis was performed to identify and exclude genes under balancing selection from Database-

group and Sma3s-group sets using BetaScan (vers. 2; Siewert and Voight, 2020), due to the

potentially antagonistic responses of these genes. Very few genes (only 2 and 3 genes from

Database-group and Sma3s-group sets) were identified and removed from the analysis (see

Detection of genes under balancing selection in Supplementary Methods).

Test for contamination and population structure

We used the program CroCo (vers. 1.1; Simion et al., 2018) to identify candidates for cross-

species contamination (see supplementary materials for details). Overall, we did not detect a

clear case of cross-species contamination in our dataset (Figure S1). Contigs identified as

potential contamination always involve a pair of species belonging to the same genus. In this

case, contamination could be difficult to identify due to the low genetic divergence between

species.

For the newly sequenced species, we also performed PCA analyses using allele frequencies of

control genes. We used the function dudi.pca of adegenet R package (Jombart and Ahmed,

2011). This analysis aims to check for population structure and to detect potentially

problematic individuals (i.e., contaminated individuals). This analysis led to the exclusion of 4

individuals (*Ploceus princeps* P6-174; *P. grandis* ST10_094; *P. nigerrimus* G3_016; *C.*
teneriffae TF57) for which we suspected contamination. Otherwise, no extra population
structure was detected (Figure S2-S4).

Hidden paralogy

We computed the statistic $F_{IS} = 1-H_0/H_e$ where H_0 is the average number of heterozygous
individuals observed (H_0 = #heterozygous / n ; where n is the sample size) and H_e is the
expected number of heterozygous individuals at Hardy-Weinberg (HW) equilibrium (H_e =
(n/(n-1) 2 * p * (1-p))*n where n is the sample size and p the allele frequency of a randomly
chosen allele). F_{IS} varies between -1 and 1 with positive value representing excess of
homozygous individuals and negative value representing excess of heterozygous individuals
compared to the HW proportions. Gene with high value of nucleotide diversity (Pi) and
negative value of F_{IS} could represent a potential case where hidden paralogous sequences have
not been separated and where all the individuals present heterozygous sites in the positions
where a substitution occurred between the paralogous copies. Five sequences corresponding to
the TLR21 genes appeared problematic (Pi > 0.01 and F_{IS} < -0.5; Figure S5) and were excluded
from further analyses.

The MHC genes were more difficult to analyse. Indeed, heterozygosity could be comparable
to divergence under balancing selection. This made the identification of orthologs very
difficult. We identified a variable number of genes among species (from 1 to 10 genes for MHC
class I and MHC class II). We checked the sequence similarity for the 10 copies of the MHC
class II in *F. albicollis* and the 7 copies of the MHC class I genes in *C. caeruleus* using cd-hit
(Fu et al., 2012). For MHC class II, sequence divergences were always higher than 15%
indicating that reads were likely correctly assigned to their corresponding gene copy. For MHC
class I, sequence similarity could be as high as 95%. In this case, we relied on the fact that the
reads from very similar paralogous copies were not be confidently assigned to a gene copy
sequence by the mapping software. This should lead to a low mapping score quality and were
likely to be discarded during the genotype calling procedure. For example, 3 out of 7 of the
Cyanistes MHC class I genes were not correctly genotyped and were missing from our final
dataset.

Data Analysis

SLiM simulations

We used SLiM (vers. 3.3.2; Haller and Messer, 2017) to estimate the impact of demographic

changes on polymorphism patterns under various selection regimes. The following parameters

were used in all simulations. Sequences of 30kb with a mutation rate of $4.6e^{-9}$

substitutions/site/generation were simulated (Smeds et al., 2016). Recombination was set to be

equal to mutation rate. Introns/exons pattern was reproduced by simulating fragments of 3kb

separated by one bp with a very high recombination rate of 0.1 rec./site/generation. We chose

3kb because TLR CDS were typically single-exon sequences of 2-3kb (Velová et al., 2018).

Five types of mutations were possible: i) neutral synonymous mutations, ii) codominant non-

synonymous mutations with a Distribution of Fitness Effect (DFE) following a gamma law of

mean = -0.025 and shape = 0.3, which corresponds to the DFE estimated in Passerines by

Rousselle et al. (2020), iii) codominant non-synonymous mutations positively selected with s

= 0.1, iv) non-synonymous mutations under balancing selection with an effect on fitness

initially set at 0.01 but re-estimated by the program at each generation according to the mutation

frequency in the population, thus including a frequency-dependent effect and v) non-

synonymous mutations under overdominance with a dominance coefficient of 1.2.

We simulated a coding sequence organization where positions one and two of the codons were

considered as non-degenerated sites, with the non-synonymous types of mutations previously

described were possible in various proportions. The third position was considered as

completely neutral where only synonymous mutations could appear.

In the absence of control genes evolving under balancing selection, we used SLiM to generate

a set of control genes for this category. We simulated two populations of 270,000 and 110,000

individuals, representing mainland and island effective population size respectively.

We also explored the effect of positive and balancing selection on the pattern of Ps and Pn/Ps

in a population of size 50,000, 110,000, 270,000 and 500,000. In order to speed up the

computational time, we reduced the population size by a factor 100 and rescaled mutation rate,

recombination rate and selection coefficient accordingly running 10 replicates per simulation.

All the details of the simulation parameters, calculations of non-synonymous polymorphism

rate (Pn) and synonymous polymorphism rate (Ps) of simulated sequences, as well as SLiM

command lines are provided in Supplementary Methods and Materials.

Polymorphism analyses

Synonymous (Ps) and non-synonymous (Pn) nucleotide diversities were estimated from

seq_stat_coding written from the Bio++ library (Available as Supplementary data; Guéguen et

al., 2013). The mean Pn/Ps was computed as the sum of Pn over the sum of Ps (Wolf et al.,

2009). Ps of concatenated sequences of control genes were estimated for each species of our

dataset. For the whole-genome sequenced species, we compared the Pn/Ps and Ps estimated

from the 97 control genes with the values from Leroy et al., (2021b; ~5000 genes used in their

study). Pn/Ps and Ps correlations showed a R^2 of 0.6 and 0.95 respectively (Figure S6). Thus,

the 97 control genes used in our study were representative of the larger set of genes from Leroy

et al (2021b). This allowed us to identify *Phylloscopus trochilus* as an outlier. Unlike for all

other species (e.g. *Fringilla coelebs*, Figure S7), synonymous polymorphism level was

correlated to the amount of missing data in *P. trochilus* alignments (Figure S7). As such, we

excluded *P. trochilus* from further analysis.

The mean Pn/Ps, calculated from the concatenated sequences of genes from the same gene

class (control genes; BD; TLR; MHC I; MHC II; Database-group; Sma3s-group), was

estimated for each bird species. Alternative transcripts were identified based on the genomic

position in the GFF file. If several transcripts were available, one transcript was randomly

selected. Pn/Ps estimates based on less than four polymorphic sites were excluded from the

analysis, as were those with no polymorphic non-synonymous sites.

Statistical analyses

To estimate the impact of demographic history on genome-wide polymorphism of island

species and the potentially reduced constraints on their immune genes, we computed the ratio

of non-synonymous nucleotide diversity over synonymous nucleotide diversity (Pn/Ps). A

linear mixed model was performed, using the Pn/Ps ratio as dependent variable and, as

explanatory variables, the mainland or insular origin of species as well as the category of genes

(packages lme4 and lmerTest (Bates et al., 2012; Kuznetsova et al., 2017)). In order to take

into account the phylogenetic effect, the taxonomic rank "family" was included as a random

effect in the model. We also used a generalized linear mixed model (using the function glmer

of the package lme4) with the family "Gamma(link="log")" which led to the same results

(Figure S15 to S24). Five linear mixed models were defined i) model including origin and gene

category parameters and also the interaction effect ii) model using both origin and gene

category parameters, iii) model with only the gene category parameter, iv) model with only the

origin parameter, and finally v) null model. In some cases, the phylogenetic effect was difficult

to estimate because the number of species per family was reduced to one. In that case, we

choose to reduce the number of families by grouping Turdidae with Muscicapidae,

Nectariniidae, and Estrildidae with Ploceidae and Fringillidae within Thraupidae. The results

obtained with these family groupings were similar to the original model (Table S1), except

when stated. The categories Database-group and Sma3s-group were tested separately from the

Core group because they contained hundreds of genes annotated using the automatic pipeline

that were only available for species with genome wide data. Database-group and Sma3s-group

were not analysed simultaneously because they contained a partially overlapping set of genes.

Finally, genes evolving under purifying selection and genes evolving under balancing selection

were also analysed separately. Model selection was based on two methods. First, we used the

difference in corrected Akaike Information Criterion (ΔAICc) calculated using the qpcR

package (Spiess and Spiess, 2018). Second, a model simplification using an ANOVA between

models was also performed.

We also tested an alternative model using the difference between Pn/Ps of immune genes and

control genes (ΔPn/Ps) as dependent variable, and species origin as explanatory variable. Under

the hypothesis of a relaxation in selection pressure on islands due to a change in the parasite

community, we expected the ΔPn/Ps to be higher on island species compared to the mainland

ones and, therefore, the species origin (i.e., mainland or island) to be significant. In this model,

we used the Phylogenetic Generalized Least Squares model (PGLS; implemented in the "nlme"

packages; Pinheiro et al., 2017). This model assumed that the covariance between species

follows a Brownian motion evolution process along the phylogeny (implemented using the

"corBrownian" function from the "'ape'" package; Paradis and Schliep, 2019). The species

phylogeny was estimated using mitochondrial genes and a maximum likelihood inference

implemented in IQTREE (model GTR+Gamma and ultrafast bootstrap; Nguyen et al., 2014;

median of 11,134 bp analysed per species). The phylogeny with the bootstrap support is

provided as supplementary material.

All the statistical analyses were performed using R (R Core Team, 2018), and dplyr package

(Wickham, 2016). Graphical representations were done using ggplot2, ggrepel, ggpubr and

ggpmisc (Aphalo, 2020; Kassambara, 2018; Slowikowski et al., 2018; Wickham, 2016).

Results

For the 150 individuals (10 species with 15 individuals each) for which we generated new data

by targeted capture sequencing, an average of 3.3 million paired-ends reads per individual was

generated (Table S1). Additionally, we generated three new draft assemblies using 40x pair-

end illumina data for the species for which no closely related reference genomes were available.

N50 and total size were 1.11 Gb and 27.9 kb for *Cyanomitra olivaceus;* 1.10 Gb and 31.7 kb

for *Ploceus cucullatus* and 1.13 Gb and 14.3 kb for *Turdus pelios* respectively. After mapping,

genotyping and cleaning, we analysed 86 control and 16 immune genes on average per species,

out of the 141 targeted genes (120 control and 21 immune related genes; Table S4). For the

species with whole-genome sequences, we analysed 106 control and 20 immune genes on

average per species, out of the 141 targeted genes, and 875 and 688 genes on average in the

Database-group and Sma3s-group respectively (Table S4).

For the species for which full genome sequences were available, the Ps and Pn/Ps estimated

using the control genes reflect the Ps and Pn/Ps of the whole transcriptome (Figure S6).

Population genetics of BD and TLR immune genes

In order to characterize the selection regimes shaping the BD and TLR polymorphisms (Figure

3), we first analyzed the variation of Pn/Ps ratios among gene categories using a linear mixed

model.

Figure 3: Pn/Ps according to species origin (mainland in green and insular in orange) for different gene categories under purifying selection. The number of species (N), and the mean Pn/Ps are shown for each modality.

Model selection based on AICc as well as model selection approach based on simplification with ANOVA identified the model n° 2, including the origin (i.e., mainland or island) and gene category without interaction (Table 2). In this model, island origin of species is associated with a greater Pn/Ps (0.14 vs. 0.10; Table 3; $p < 0.01$). Gene categories corresponding to TLRs and BDs showed a significantly higher Pn/Ps than control genes (Table 3; $p < 0.001$). Our statistical analysis confirmed that island birds have a higher Pn/Ps ratio than mainland relatives, in agreement with the nearly-neutral theory of evolution. It also reveals that immune genes have a higher Pn/Ps than randomly selected control genes suggesting that BD and TLR evolve under a different selection regime than non-immune related genes.

Next, we investigated the cause of the higher Pn/Ps of immune genes by testing three hypotheses. First, we excluded a bias due to a lower number of immune genes, and therefore higher variance in the estimation of Pn/Ps in immune genes. Immune genes still had significantly higher Pn/Ps compared to a random subsample of control genes of comparable size (Figure S8 & S9). Second, the Pn/Ps of immune genes could be inflated by positive selection. It is well known that immune genes are subject to frequent adaptation due to arms race evolution with pathogens (Enard et al., 2016; Shultz and Sackton, 2019; Velová et al., 2018). We evaluated the effect of positively selected genes on the Pn/Ps using SLiM simulations with both positively and negatively selected mutations. The presence of recurrent

positive selection could increase the Pn/Ps leading to a higher Pn/Ps in immune genes if this

category was more prone to adaptive evolution (Figure 4A). However, positive selection

always led to a drastic decrease in Ps due to genetic sweep effect at linked sites (Figure 4B).

BDs and TLRs had a slightly higher or similar Ps than control genes (Figure S9, mean Ps =

0.007, 0.004 and 0.003 for BDs, TLRs and control genes respectively, effect of gene category

$p < 0.1$) and, as a consequence, even if positive selection is likely to have impacted the

evolution of immune genes, it is not the cause of the higher Pn/Ps observed here. Third,

balancing selection could be present, at least temporarily, in the evolution of BDs and TLRs

genes (Kloch et al., 2018; Levy et al., 2020). Simulation analyses confirmed that balancing

selection causes an increase of Ps and Pn/Ps (Figure 4C & 4D). However, a change in effective

population size had an opposite effect on the Pn/Ps according to whether selection was negative

or balancing. In the presence of slightly deleterious mutations, Pn/Ps decreases with Ne

whereas it increases in the presence of balancing selection. Island birds had higher Pn/Ps ratios

than mainland birds for BDs and TLRs. Therefore, we can rule out balancing selection as the

main factor explaining the high Pn/Ps of immune genes because, in this case, Pn/Ps of island

birds should be lower. Another possible explanation is a relaxed selection of immune genes. It

is likely that immune genes are overall less constrained than the control genes. It has been

shown that evolutionary constraints are more related to gene expression than to function

(Drummond et al., 2005; Drummond and Wilke, 2008) and therefore, functionally important

genes could still have a high Pn/Ps.

Overall, our analyses do not support a strong impact of ongoing adaptive mutation or balancing

selection on BDs and TLRs. However, these immune genes do not evolve as random genes

(not involved in immune functions) and present a significantly higher Pn/Ps of 0.20 ($p < 0.001$

; Table 3).

No evidence of a reduced impact of the parasite communities on the polymorphism pattern of

immunes genes in island birds

For BDs and TLRs, the best model selected includes the origin (i.e., mainland or island) and

gene category without interaction, corresponding to model n°2 (see above and Table 2). This

model has no interaction between origin and gene categories invalidating the hypothesis of a

reduced parasite communities on islands (Figure 2).

Figure 4: Neutral polymorphism (Ps) and ratio of selected over neutral polymorphism (Pn/Ps) estimated from SLiM simulations. A) Pn/Ps as a function of population size, N and B) Ps as a function of N. In both A and B, colour indicates the frequency of positively selected mutation compare to deleterious mutation. C) Pn/Ps as a function of N and D) Ps as a function of N. In both C and D, yellow indicates simulations with overdominance mutation (h = 1.2) and negatively selected mutations and green indicates simulations with only negatively selected mutations.

Table 2: Statistical model explaining Pn/Ps variation of Toll-Like Receptors, Beta-Defensins genes, and control genes. The p-values of ANOVA test between simpler models are not reported if a more complex model explains a larger proportion of the variance.

n°	Model Details	Model selection by AIC AICc	ΔAICc	Likelihood	ANOVA test n° 1	2	3	4
1	Pn/Ps~ 1+ category +origin+ category *origin	-5.39	8.83	0.01		0.63		
2	Pn/Ps~ 1+ category +origin	-14.22	0	1			0.002	3.71E-05
3	Pn/Ps~1+ category	-11.8	2.42	0.3				
4	Pn/Ps~1+ origin	-6.83	7.39	0.02				
5	Pn/Ps~1	-6.44	7.78	0.02				

465 <u>Table 3</u>: Summary of the model n°2, best statistical model selected using AICc explaining variation in
466 Pn/Ps in control genes, Toll-Like receptors and Beta-Defensins genes under purifying selection with
467 origin, gene category parameters. * indicates significant values : * < 0.05; ** < 0.01; *** < 0.001.

Model		Parameters		Estimate	P.value	
		Origin	*Category*			
Origin and	Intercept	*mainland*	*Control genes*	*0.10*	*2.65E-02*	*
Gene		*island*		*0.14*	*4.56E-03*	**
category			*Toll-Like Receptors*	*0.20*	*7.43E-05*	***
(n°2)			*Beta-Defensins genes*	*0.20*	*3.16E-04*	***

For larger sets of genes, identified using an automatic pipeline and gene annotation, model
selection based on AICc and simplification with ANOVA (Table S5, S8) identified models n°4
that included origine parameters which associated a higher Pn/Ps of at least 0.07 for island
species (p < 0.001; Table S6, S7, S9, S10, Figure 5). Model selection by simplification with
ANOVA identified models n°1 with interaction effect between origin and gene category
associated with a reduced Pn/Ps for TLR and BD genes of island species that invalidate our
hypothesis (Table S7, S10).

The alternative statistical approach using the difference between Pn/Ps of immune genes and
control genes (*Δ*Pn/Ps) as dependent variable, and species origin as explanatory variable under
a PGLS framework lead to similar results. Island was never associated to a statistically higher
*Δ*Pn/Ps (table S2) providing no support for an increased relaxed selection of immune genes in
island species.

480
481 <u>Figure 5</u>: Boxplot of Pn/Ps according to species origin (mainland in green and insular in orange) for

482   different gene categories under purifying selection. The number of individuals (N), and the mean Pn/Ps

483   are shown for each modality.

Genes under balancing selection

First, we estimated the effect of population size variation on the Pn/Ps of the genes evolving

under balancing selection by simulating sequences under frequency dependent or

overdominance selection using SLiM (see Methods and Supplementary Methods). The

simulation under frequency dependent selection revealed an average Pn/Ps equal to 0.8 for

island species and 1.2 for mainland species (Figure 6). Under overdominance, simulated

sequences from island and mainland populations respectively have an average Pn/Ps equal to

0.54 and 1.03 (Figure 6).

Figure 6: Boxplot of Pn/Ps according to species origin (mainland in green and insular in orange) for

different gene categories under balancing selection. The number of species (N), and the mean Pn/Ps are

shown for each modality. The control groups correspond to the results obtained from simulated

sequence via SLiM (see Methods and Supplementary Methods Simulation of control genes under

balancing selection).

Using simulations under frequency dependent selection as well as simulations under the

overdominance, model selection by AIC identifies the model n°4 with origin, contrary to the

method by simplification with ANOVA which identified the full model (model n°1) therefore

including significant interaction between origin and genes category (Table 4). This interaction

effect is significant for the MHC II ($p < 0.05$, Table S12) but not for MHC I. As expected,

island species have a significantly lower Pn/Ps in MHC genes compared to mainland species

(p < 0.01; except for the full model based on control genes evolving under overdominance

Table S12).

<u>Table 4</u>: Statistical model explaining Pn/Ps variation of genes under balancing selection (i.e MHC class

I and II), and simulated sequences under i) frequency dependent or ii) overdominance. The p-values of

ANOVA test between simpler models are not reported if a more complex model explains a larger

proportion of the variance.

Model			Model selection by AIC			ANOVA test			
Type of balancing selection	n°	Details	AICc	ΔAICc	Likelihood	n°1	2	3	4
Frequency dependent	1	Pn/Ps~1+ category +origin+ category *origin	157.17	5.62	0.06		0.019		
	2	Pn/Ps~1+ category +origin	157.85	6.31	0.04				
	3	Pn/Ps~1+ category	187.58	36.04	0.00				
	4	Pn/Ps~1+ origin	151.54	0.00	1.00				
	5	Pn/Ps~1	180.52	28.97	0.00				
Overdominance	1	Pn/Ps~1+ category +origin+ category *origin	140,56	8,50	0,01		0.024		
	2	Pn/Ps~1+ category +origin	140,56	8,50	0,01				
	3	Pn/Ps~1+ category	185,91	53,85	0,00				
	4	Pn/Ps~1+ origin	132,05	0,00	1,00				
	5	Pn/Ps~1	177,54	45,49	0,00				

Discussion

On oceanic islands, the depauperate parasite community is expected to lead to a relaxation of

selection on the immune system. In this study, we found support for such an effect, but only on

MHC class II genes and using simulated sequences under balancing selection as control. No

effect was detected for MHC class I genes nor for innate immune genes (TLRs and BDs),

evolving under purifying selection. On these sets of genes, increased drift effects on island

populations limit the efficacy of selection in accordance with the nearly-neutral theory (Ohta,

1992). The ability to distinguish between the selective and nearly-neutral processes (relaxed

selection due to environmental change vs. drift) could only be achieved by our approach of

using random genes (i.e., "control genes") to estimate the genome-wide effect of potential

variation in effective population size between populations.

Effects of effective population size variation

Our results support the nearly-neutral theory of evolution for those genes under purifying

selection, whereby strong genetic drift acting on small island populations reduces the efficacy

of natural selection, leading to an increase in non-synonymous nucleotide diversity compared

to the mostly neutral, synonymous nucleotide diversity (i.e., Pn/Ps; Ohta, 1992). This is

materialized by a genome-wide increase in frequency of weakly deleterious mutations

(Kutschera et al., 2020; Leroy et al., 2021b; Loire et al., 2013; Robinson et al., 2016; Rogers

and Slatkin, 2017).

For genes evolving under balancing selection, we performed simulations under the hypotheses

of overdominance (heterozygote advantage) or frequency-dependent (rare allele advantage).

Our results showed reduced Pn/Ps for smaller population sizes (Figure 6, S10, S11). This

simulation confirmed our expectations (Figure 2) that a reduction in the efficacy of selection

results in a decrease in the frequency of non-synonymous polymorphism, as, under normal

circumstances, selection maintains those mutations at intermediate frequencies. It also matches

what we obtained for the empirical results, where both MHC classes I and II had a reduced

Pn/Ps in island birds. This result supports that the fitness effect of having non-synonymous

polymorphisms segregating at high frequencies is not strong enough to counteract entirely the

effect of genetic drift on islands.

Effects of selection on immune genes

For immune genes, we tried to characterize the nature of the selection acting on BDs and TLRs

genes. Comparing those genes with control genes and using simulations, we were able to rule

out that directional positive selection and balancing selection had a major impact shaping the

polymorphism of these immune genes. In contrast, the pattern of Pn/Ps between island and

mainland populations is in line with the effect of purifying selection in the presence of slightly

deleterious mutations. However, no effect was detected on insular species, beyond what could

be attributed to genetic drift. This is in line with the result of Gonzalez-Quevedo et al. (2015b)

and Grueber et al. (2013) who found that TLR genetic diversity was mostly influenced by

genetic drift. At first sight, this result seems not in line with the fact that island parasite

communities are less diverse (Beadell et al., 2006; Loiseau et al., 2017; Maria et al., 2009;

Pérez-Rodríguez et al., 2013; but see Illera et al., 2015). However, a reduced number of

pathogens has also been found to be associated with a higher prevalence in birds and reptiles

from the Macaronesian archipelago (Illera and Perera, 2020). Therefore, these two patterns, i.e.

a less diverse pathogen's community on islands with a higher prevalence, could still imply a

strong selection pressure on immune genes.

In contrast, for MHC genes that unambiguously evolve under balancing selection, MHC class

II genes presented a reduction in non-synonymous polymorphism larger than the effects of drift

alone, when simulated sequences are used as control. This was the only case where a role for

relaxed selection pressures in the molecular evolution of immune genes could be invoked.

Our results are in accordance with the hypothesis of Lee (2006), which proposes that innate

and acquired immunity may exhibit distinct responses to changes in pressures due to different

costs and benefits. However, they contrast with the study of Santonastaso et al. (2017) that

identified no change in selection pressures on MHC II genes in a lizard species and concluded

that their evolution was mostly governed by drift. Similarly, Agudo et al. (2011) also found a

prominent role for genetic drift over selection in the evolution of MHC II genes in the Egyption

vulture (*Neophron percnopterus*).

Our results rely on simulations that may be affected by the choice of the parameter values.

First, we performed simulations using a fixed effective population size (*Ne*) estimated from the

polymorphism data. Using others values of *Ne* had a weak impact on the relative difference

between island and mainland species for the overdominance type of selection (Figure S10,

S11). Secondly, we simulated two types of selection, namely overdominance (Doherty and

Zinkernagel, 1975) and frequency-dependent (Slade and McCallum, 1992), but it has been

argued that the maintenance of MHC polymorphism could be the result of fluctuating selection

(Hill, 1991). Additionally, recombination has also been put forward as a mechanism

responsible for generating diversity (Spurgin et al., 2011). Therefore, our results for the MHC

II genes, which is based on the relative difference between Pn/Ps of island and mainland species

comparing empirical and simulated data, should be taken cautiously as their significance can

be dependent on the specific parameters that we used, although we did our best to select a
realistic range of parameters.

The observed difference between MHC class I and II could be explained by their different
pathogen targets: MHC class I genes are primarily involved in the recognition of intracellular
pathogens (Kappes and Strominger, 1988), while MHC class II genes are directly involved in
the recognition of extracellular pathogens (Bjorkman and Parham, 1990). These differences
could lead to variable selection pressures depending on the extracellular versus intracellular
parasite communities present on islands. In addition, the relaxed selection pressures on MHC
II genes from adaptive immunity is in line with a reduction in acquired immunity parameters
foundby Lobato et al. (2017).

Future work should take into account that there is an extensive variation in the number of MHC
gene copies across the avian phylogeny (Minias et al., 2019; O'Connor et al., 2020).
Particularly, it was recently discovered that Passerines have a very dynamic evolution of
duplication/loss events compared to other birds (Minias et al., 2019). Here, we used the two
copies of MHC gene I and II currently annotated in the collared flycatcher genome as target
sequences for our targeted-capture sequencing. The future improvement of genome assembly,
resulting from the development of long-reads technology (Peona et al., 2021, 2018), should
help to annotate with increased precision all MHC copies and to study the whole repertoire of
MHC genes.

Consequences of drift and selection on immunity

The potential relaxation of the natural selection acting on immune genes in island species is
expected to reduce immune functions and increase susceptibility of island populations to
pathogens. This is true even if this relaxation is only the consequence of a reduction in the
effective population size and not caused by a reduction of the pressure exerted by the parasitic
community. This is in line with the results of Hawley et al. (2005) and Belasen et al. (2019)
who showed that a decrease in diversity of immune loci (MHC II or through immune proxy)
was associated with a reduction in immune functions. It should be noted that even if migration
rate is reduced on islands, sedentary and endemic island species are not completely free from
the exposure of exogen pathogens through migratory birds (Levin et al., 2013).

As a final remark, we would like to stress that more research is needed (i) to ascertain both
selection pressures on innate and adaptive immune responses and the load of deleterious

mutations due to drift, also identified by an increasing body of work (Loire et al., 2013;
Robinson et al., 2016; Rogers and Slatkin, 2017; Kutschera et al., 2020; Leroy et al., 2021b),
and (ii) to better describe island parasite communities. To date, most of the studies investigated
intracellular parasite communities on islands, and more specifically haemosporidian parasites,
avian pox and coccidian parasites (Cornuault et al., 2012; Illera et al., 2015, 2008; Ishtiaq et
al., 2010; Loiseau et al., 2017; Martinez et al., 2015; Padilla et al., 2017; Pérez-Rodríguez et
al., 2013; Silva-Iturriza et al., 2012), whereas very few evaluated the extracellular parasite
diversity, such as helminths (Nieberding et al., 2006, but see the review of Illera and Perera
2020 for reptiles). Metabarcoding of parasites is a new technique to evaluate at the same time
both communities of intracellular and extracellular parasites (Bourret et al., 2021) and might
therefore be a promising approach to compare their communities in island and mainland
populations.

Conclusion

Our comparative population genomics study has investigated the combined effects of drift and
selection on immune genes from island and mainland passerines. The study of synonymous
and non-synonymous polymorphism of these genes confirmed that island species, with smaller
population sizes than their mainland counterparts, were more impacted by drift, which induces
a load of weakly deleterious mutations in their genome. Indeed most of the genes studied here
involved in the immune response do not show a statistically different pattern from control
genes. Only MHC II genes, involved in the recognition of extracellular pathogens, showed a
reduction in their non-synonymous polymorphism in island species. This response, which may
be attributed to reduced selection pressures on these genes, could be associated with the
suspected reduced parasitic communities on islands. The increased load of deleterious
mutations as well as the potential relaxed selection pressures on MHC II support the reduced
immune functions of island species, which could be added to the list of other convergent
responses of the island syndrome.

Data availability

Datasets, scripts, supplementary figures and texts are available on figshare :
https://figshare.com/s/ab7004cc2f4415b4058f. The reads newly generated for this study have
been deposited in the NCBI Sequence Read Archive under the bioproject PRJNA724656.

Acknowledgments

In Gabon, we thank the Director and the guides of the Lekedi Park, Marie Charpentier for her

help in organizing the expedition, and Elisa Lobato, Alexandre Vaz for field assistance and

outreach work. In São Tomé and Príncipe, we thank the Directorate of the Environment and

the Department for Nature Conservation, its directors—Arlindo Carvalho and Victor Bonfim—

Guilhermino, the Association Monte Pico, its president Luis Mário, and its members. Elisa

Lobato, Philippe Perret, Octávio Veiga, Bikegila, and Yelli provided invaluable assistance in

the field. Permissions for fieldwork were given by the authorities of São Tomé and Príncipe

and Gabon (CENAREST authorization No. AR0053/12/MENESTFPRSCJS/

CENAREST/CG/CST/CSAR). Permits for the Canary Islands were provided by the Regional

Government (Ref.: 2012/0710), and the Cabildo of La Palma and Tenerife. In Montpellier, we

thank the blue tit team (https://oreme.org/observation/ecopop/mesanges/) for the capture of the

individuals used in this study. The analyses benefited from the Montpellier Bioinformatics

Biodiversity (MBB) platform services. This research was conducted in the scope of the

international twin-lab "LIA – Biodiversity and Evolution" between CIBIO (Portugal) and

ISEM and CEFE-CNRS (France). This is ISEM publication n° ISEM 2022-223.

Funding information

This research was funded by the Labex CeMEB (project ISLAND IMMUNITY) for BN, CL

and CD, the ANR (BirdIslandGenomic project, ANR-14-CE02-0002) for MB and BN,, the

National Geographic Society (Grant/Award Number:W251-12), the British Ecological Society

(Grant/Award Number: 369/4558) to Elisa Lobato, RC and CD, the portuguese Foundation for

Science and Technology under the PTDC/BIA-EVL/29390/2017 "DEEP" Research Project for

MM, RC and CL, and the Spanish Ministry of Science, Innovation and Universities, and the

European Regional Development Fund (Ref.: PGC2018-097575-B-I00) for JCI.

References

Agudo, R., Alcaide, M., Rico, C., Lemus, J.A., Blanco, G., Hiraldo, F., Donázar, J.A., 2011.
Major histocompatibility complex variation in insular populations of the Egyptian
vulture: inferences about the roles of genetic drift and selection. Mol. Ecol. 20, 2329–
2340. https://doi.org/10.1111/j.1365-294X.2011.05107.x
Akira, S., 2003. Toll-like receptor signaling. J Biol Chem 278, 38105–38108.
Alberts, B., Johnson, A., Lewis, J., Raff, M., Roberts, K., Walter, P., 2002. Innate immunity.
Mol. Biol. Cell.
Aphalo, P.J., 2020. ggpmisc: Miscellaneous Extensions to "ggplot2"(R package version 0.3.

6.

Baeckens, S., Van Damme, R., 2020. The island syndrome. Curr. Biol. 30, R338–R339.

Bates, D.M., Maechler, M., Bolker, B., Walker, S., 2012. Package 'lme4.' CRAN R Found
Stat Comput.

Beadell, J.S., Atkins, C., Cashion, E., Jonker, M., Fleischer, R.C., 2007. Immunological
change in a parasite-impoverished environment: divergent signals from four island
taxa. PLoS One 2:e896.

Beadell, J.S., Ishtiaq, F., Covas, R., Melo, M., Warren, B.H., Atkinson, C.T., Bensch, S.,
Graves, G.R., Jhala, Y.V., Peirce, M.A., 2006. Global phylogeographic limits of
Hawaii's avian malaria. Proc R Soc Lond B Biol Sci 273, 2935–2944.

Belasen, A.M., Bletz, M.C., S, L.D., Toledo, L.F., James, T.Y., 2019. Long-term habitat
fragmentation is associated with reduced MHC IIB diversity and increased infections
in amphibian hosts. Front Ecol Evol 6.

Bernatchez, L., Landry, C., 2003. MHC studies in nonmodel vertebrates: what have we
learned about natural selection in 15 years? J Evol Biol 16, 363–377.

Bjorkman, P.J., Parham, P., 1990. Structure, function, and diversity of class I major
histocompatibility complex molecules. Annu Rev Biochem 59, 253–288.

Blondel, J., 2000. Evolution and ecology of birds on islands: trends and prospects. Vie
MilieuLife Environ. 205–220.

Bourret, V., Gutiérrez López, R., Melo, M., Loiseau, C., 2021. Metabarcoding options to
study eukaryotic endoparasites of birds. Ecol. Evol. 11, 10821–10833.

Boyce, M.S., 1984. Restitution of gamma-and k-selection as a model of density-dependent
natural selection. Annu Rev Ecol Syst 15, 427–447.

Breuer, K., Foroushani, A.K., Laird, M.R., Chen, C., Sribnaia, A., Lo, R., Winsor, G.L.,
Hancock, R.E., Brinkman, F.S., Lynn, D.J., 2013. InnateDB: systems biology of
innate immunity and beyond—recent updates and continuing curation. Nucleic Acids
Res 41:D1228–D1233.

Buffalo, V., 2021. Quantifying the relationship between genetic diversity and population size
suggests natural selection cannot explain Lewontin's paradox. eLife 10, e67509.
https://doi.org/10.7554/eLife.67509

Castellano, D., James, J., Eyre-Walker, A., 2018. Nearly neutral evolution across the
Drosophila melanogaster genome. Mol. Biol. Evol. 35, 2685–2694.

Chapman, H., JR, O, H., AS, K., RH, C., RL, W., J., 2016. The evolution of innate immune
genes: purifying and balancing selection on β-defensins in waterfowl. Mol Biol Evol
33, 3075–3087.

Charlesworth, J., Eyre-Walker, A., 2008. The McDonald–Kreitman test and slightly
deleterious mutations. Mol Biol Evol 25, 1007–1015.

Chen, J., Glémin, S., Lascoux, M., 2020. From drift to draft: how much do beneficial
mutations actually contribute to predictions of Ohta's slightly deleterious model of
molecular evolution? Genetics 214, 1005–1018.

Chen, S., Zhou, Y., Chen, Y., Gu, J., 2018. fastp: an ultra-fast all-in-one FASTQ
preprocessor. Bioinformatics 34:i884–i890.

Corcoran, P., Gossmann, T.I., Barton, H.J., Slate, J., Zeng, K., 2017. Determinants of the
Efficacy of Natural Selection on Coding and Noncoding Variability in Two Passerine
Species. Genome Biol. Evol. 9, 2987–3007. https://doi.org/10.1093/gbe/evx213

Cornuault, J., Bataillard, A., Warren, B.H., Lootvoet, A., Mirleau, P., Duval, T., Milá, B.,
Thébaud, C., Heeb, P., 2012. The role of immigration and in-situ radiation in
explaining blood parasite assemblages in an island bird clade. Mol. Ecol. 21, 1438–
1452.

Covas, R., 2012. Evolution of reproductive life histories in island birds worldwide. Proc R

Soc B Biol Sci 279, 1531–1537.

Doherty, P.C., Zinkernagel, R.M., 1975. Enhanced immunological surveillance in mice heterozygous at the H-2 gene complex. Nature 256, 50–52.

Doutrelant, C., Paquet, M., Renoult, J.P., Grégoire, A., Crochet, P.-A., Covas, R., 2016. Worldwide patterns of bird colouration on islands. Ecol Lett 19, 537–545.

Drummond, D.A., Bloom, J.D., Adami, C., Wilke, C.O., Arnold, F.H., 2005. Why highly expressed proteins evolve slowly. Proc. Natl. Acad. Sci. 102, 14338–14343.

Drummond, D.A., Wilke, C.O., 2008. Mistranslation-induced protein misfolding as a dominant constraint on coding-sequence evolution. Cell 134, 341–352.

Ellegren, H., Smeds, L., Burri, R., Olason, P.I., Backström, N., Kawakami, T., Künstner, A., Mäkinen, H., Nadachowska-Brzyska, K., Qvarnström, A., 2012. The genomic landscape of species divergence in Ficedula flycatchers. Nature 491, 756–760.

Enard, D., Cai, L., Gwennap, C., Petrov, D.A., 2016. Viruses are a dominant driver of protein adaptation in mammals. elife 5, e12469.

Eyre-Walker, A., Keightley, P.D., 2007. The distribution of fitness effects of new mutations. Nat. Rev. Genet. 8, 610–618.

Fijarczyk, A., Dudek, K., Babik, W., 2016. Selective Landscapes in newt Immune Genes Inferred from Patterns of Nucleotide Variation. Genome Biol Evol 8, 3417–3432.

Frankham, R., 1997. Do island populations have less genetic variation than mainland populations? Heredity 78, 311–327.

Fu, L., Niu, B., Zhu, Z., Wu, S., Li, W., 2012. CD-HIT: accelerated for clustering the next-generation sequencing data. Bioinformatics 28, 3150–3152.

Garamszegi, L.Z., 2006. The evolution of virulence and host specialization in malaria parasites of primates. Ecol Lett 9, 933–940.

Garrison, E., Marth, G., 2012. Haplotype-based variant detection from short-read sequencing.

Gonzalez-Quevedo, C., Phillips, K.P., Spurgin, L.G., Richardson, D.S., 2015a. 454 screening of individual MHC variation in an endemic island passerine. Immunogenetics 67, 149–162. https://doi.org/10.1007/s00251-014-0822-1

Gonzalez-Quevedo, C., Spurgin, L.G., Illera, J.C., Richardson, D.S., 2015b. Drift, not selection, shapes toll-like receptor variation among oceanic island populations. Mol. Ecol. 24, 5852–5863.

Grant, P.R., 1965. The adaptive significance of some size trends in island birds. 355–367, Evolution.

Grueber, C.E., Wallis, G.P., Jamieson, I.G., 2014. Episodic positive selection in the evolution of avian toll-like receptor innate immunity genes. PloS One 9, e89632.

Grueber, C.E., Wallis, G.P., Jamieson, I.G., 2013. Genetic drift outweighs natural selection at toll-like receptor (TLR) immunity loci in a re-introduced population of a threatened species. Mol. Ecol. 22, 4470–4482.

Guéguen, L., Gaillard, S., Boussau, B., Gouy, M., Groussin, M., Rochette, N.C., Bigot, T., Fournier, D., Pouyet, F., Cahais, V., Bernard, A., Scornavacca, C., Nabholz, B., Haudry, A., Dachary, L., Galtier, N., Belkhir, K., Dutheil, J.Y., 2013. Bio++: Efficient Extensible Libraries and Tools for Computational Molecular Evolution. Mol. Biol. Evol. 30, 1745–1750. https://doi.org/10.1093/molbev/mst097

Hale, K.A., Briskie, J.V., 2007. Decreased immunocompetence in a severely bottlenecked population of an endemic New Zealand bird. Anim. Conserv. 10, 2–10.

Haller, B.C., Messer, P.W., 2017. SLiM 2: Flexible, interactive forward genetic simulations. Mol Biol Evol 34, 230–240.

Hawley, D.M., Sydenstricker, K.V., Kollias, G.V., Dhondt, A.A., 2005. Genetic diversity predicts pathogen resistance and cell-mediated immunocompetence in house finches. Biol Lett 1, 326–329.

Hill, A.V., 1991. HLA associations with malaria in Africa: some implications for MHC evolution, in: Molecular Evolution of the Major Histocompatibility Complex. Springer, pp. 403–420.

Hochberg, M.E., Møller, A.P., 2001. Insularity and adaptation in coupled victim–enemy associations. J Evol Biol 14, 539–551.

Illera, J.C., Emerson, B.C., Richardson, D.S., 2008. Genetic characterization, distribution and prevalence of avian pox and avian malaria in the Berthelot's pipit (Anthus berthelotii) in Macaronesia. Parasitol Res 103, 1435–1443.

Illera, J.C., Fernández-Álvarez, Á., Hernández-Flores, C.N., Foronda, P., 2015. Unforeseen biogeographical patterns in a multiple parasite system in Macaronesia. J. Biogeogr. 42, 1858–1870.

Illera, J.C., Perera, A., 2020. Where are we in the host-parasite relationships of native land vertebrates in Macaronesia? Ecosistemas.

Institute, B., 2019. "Picard Toolkit", Broad institute, GitHub repository. Picard Toolkit.

Ishtiaq, F., Clegg, S.M., Phillimore, A.B., Black, R.A., Owens, I.P., Sheldon, B.C., 2010. Biogeographical patterns of blood parasite lineage diversity in avian hosts from southern Melanesian islands. J. Biogeogr. 37, 120–132.

Jombart, T., Ahmed, I., 2011. adegenet 1.3-1: new tools for the analysis of genome-wide SNP data. Bioinformatics 27, 3070–3071.

Kappes, D., Strominger, J.L., 1988. Human class II major histocompatibility complex genes and proteins. Annu Rev Biochem 57, 991–1028.

Kassambara, A., 2018. ggpubr:"ggplot2" based publication ready plots. R Package Version 01, 7.

Kimura, M., 1962. On the Probability of Fixation of Mutant Genes in a Population. Genetics 47, 713–719.

Klein, J., 1986. Natural history of the major histocompatibility complex. Wiley.

Kloch, A., Wenzel, M.A., Laetsch, D.R., Michalski, O., Bajer, A., Behnke, J.M., Welc-Falęciak, R., Piertney, S.B., 2018. Signatures of balancing selection in toll-like receptor (TLRs) genes–novel insights from a free-living rodent. Sci. Rep. 8, 1–10.

Kutschera, V.E., Poelstra, J.W., Botero-Castro, F., Dussex, N., Gemmell, N., Hunt, G.R., Ritchie, M.G., Rutz, C., Wiberg, R.A.W., Wolf, J.B.W., 2020. Purifying Selection in Corvids Is Less Efficient on Islands. Mol. Biol. Evol. https://doi.org/10.1093/molbev/msz233

Kuznetsova, A., Brockhoff, P.B., Christensen, R.H., 2017. lmerTest package: tests in linear mixed effects models. J Stat Softw 82, 1–26.

Laine, V.N., Gossmann, T.I., Schachtschneider, K.M., Garroway, C.J., Madsen, O., Verhoeven, K.J., De Jager, V., Megens, H.-J., Warren, W.C., Minx, P., 2016. Evolutionary signals of selection on cognition from the great tit genome and methylome. Nat. Commun. 7, 1–9.

Lamichhaney, S., Berglund, J., Almén, M.S., Maqbool, K., Grabherr, M., Martinez-Barrio, A., Promerová, M., Rubin, C.-J., Wang, C., Zamani, N., 2015. Evolution of Darwin/'s finches and their beaks revealed by genome sequencing. Nature 518, 371–375.

Lee, J.W., Beebe, K., Nangle, L.A., Jang, J., Longo-Guess, C.M., Cook, S.A., Davisson, M.T., Sundberg, J.P., Schimmel, P., Ackerman, S.L., 2006. Editing-defective tRNA synthetase causes protein misfolding and neurodegeneration. Nature 443, 50–55. https://doi.org/10.1038/nature05096

Lee, K.A., 2006. Linking immune defenses and life history at the levels of the individual and the species. Integr Comp Biol 46, 1000–1015.

Leroy, Anselmetti, Y., Tilak, M.-K., Bérard, S., Csukonyi, L., Gabrielli, M., Scornavacca, C., Milá, B., Thébaud, C., Nabholz, B., 2021a. A bird's white-eye view on avian sex

824   chromosome evolution. Peer Community J. 1.

825 Leroy, Rousselle, M., Tilak, M.-K., Caizergues, A.E., Scornavacca, C., Recuerda, M., Fuchs,
826   J., Illera, J.C., De Swardt, D.H., Blanco, G., 2021b. Island songbirds as windows into
827   evolution in small populations. Curr. Biol. 31, 1303-1310. e4.

828 Levin, I.I., Zwiers, P., Deem, S.L., Geest, E.A., Higashiguchi, J.M., Iezhova, T.A., Jiménez-
829   Uzcátegui, G., Kim, D.H., Morton, J.P., Perlut, N.G., Renfrew, R.B., Sari, E.H.R.,
830   Valkiunas, G., Parker, P.G., 2013. Multiple Lineages of Avian Malaria Parasites
831   (Plasmodium) in the Galapagos Islands and Evidence for Arrival via Migratory Birds.
832   Conserv. Biol. 27, 1366–1377. https://doi.org/10.1111/cobi.12127

833 Levy, H., Fiddaman, S.R., Vianna, J.A., Noll, D., Clucas, G.V., Sidhu, J.K., Polito, M.J.,
834   Bost, C.A., Phillips, R.A., Crofts, S., 2020. Evidence of pathogen-induced
835   immunogenetic selection across the large geographic range of a wild seabird. Mol.
836   Biol. Evol. 37, 1708–1726.

837 Li, H., 2013. Aligning sequence reads, clone sequences and assembly contigs with BWA-
838   MEM. ArXiv Prepr. ArXiv13033997.

839 Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis,
840   G., Durbin, R., 2009. The Sequence Alignment/Map format and SAMtools.
841   Bioinforma Oxf Engl 25, 2078–2079.

842 Lindström, K.M., Foufopoulos, J., Pärn, H., Wikelski, M., 2004. Immunological investments
843   reflect parasite abundance in island populations of Darwin's finches. Proc R Soc Lond
844   B Biol Sci 271, 1513–1519.

845 Lobato, E., Doutrelant, C., Melo, M., Reis, S., Covas, R., 2017. Insularity effects on bird
846   immune parameters: A comparison between island and mainland populations in West
847   Africa. Ecol. Evol. 7, 3645–3656.

848 Loire, E., Chiari, Y., Bernard, A., Cahais, V., Romiguier, J., Nabholz, B., Lourenço, J.M.,
849   Galtier, N., 2013. Population genomics of the endangered giant Galapagos tortoise.
850   Genome Biol. 14, R136. https://doi.org/10.1186/gb-2013-14-12-r136

851 Loiseau, C., Melo, M., Lobato, E., Beadell, J.S., Fleischer, R.C., Reis, S., Doutrelant, C.,
852   Covas, R., 2017. Insularity effects on the assemblage of the blood parasite community
853   of the birds from the Gulf of Guinea. J. Biogeogr. 44, 2607–2617.

854 Lomolino, M.V., 2005. Body size evolution in insular vertebrates: generality of the island
855   rule. J. Biogeogr. 32, 1683–1699. https://doi.org/10.1111/j.1365-2699.2005.01314.x

856 Losos, J.B., Ricklefs, R.E., 2009. Adaptation and diversification on islands. Nature 457, 830–
857   836.

858 Lundberg, M., Liedvogel, M., Larson, K., Sigeman, H., Grahn, M., Wright, A., Åkesson, S.,
859   Bensch, S., 2017. Genetic differences between willow warbler migratory phenotypes
860   are few and cluster in large haplotype blocks. Evol Lett 1, 155–168.

861 Luo, R., Liu, B., Xie, Y., Li, Z., Huang, W., Yuan, J., He, G., Chen, Y., Pan, Q., Liu, Y.,
862   2012. SOAPdenovo2: an empirically improved memory-efficient short-read de novo
863   assembler. Gigascience 1.

864 MacArthur, R.H., Wilson, E.O., 1967. The theory of island biogeography, in: The Theory of
865   Island Biogeography. Princeton university press.

866 Maria, L., Svensson, E., Ricklefs, R.E., 2009. Low diversity and high intra-island variation in
867   prevalence of avian Haemoproteus parasites on Barbados, Lesser Antilles.
868   Parasitology 136, 1121–1131.

869 Martinez, J., Vasquez, R.A., Venegas, C., Merino, S., 2015. Molecular characterisation of
870   haemoparasites in forest birds from Robinson Crusoe Island: is the Austral Thrush a
871   potential threat to endemic birds? Bird Conserv. Int. 25, 139–152.

872 Matson, K.D., 2006. Are there differences in immune function between continental and
873   insular birds? Proc. Biol. Sci. 273, 2267–2274.

https://doi.org/10.1098/rspb.2006.3590

Matson, K.D., Beadell, J.S., 2010. Infection, immunity, and island adaptation in birds.

Minias, P., Pikus, E., Whittingham, L.A., Dunn, P.O., 2019. Evolution of copy number at the
MHC varies across the avian tree of life. Genome Biol Evol 11, 17–28.

Mueller, J.C., Kuhl, H., Timmermann, B., Kempenaers, B., 2016. Characterization of the
genome and transcriptome of the blue tit Cyanistes caeruleus: polymorphisms, sex-
biased expression and selection signals. Mol. Ecol. Resour. 16, 549–561.
https://doi.org/10.1111/1755-0998.12450

Munoz-Mérida, A., Viguera, E., Claros, M.G., Trelles, O., Pérez-Pulido, A.J., 2014. Sma3s: a
three-step modular annotator for large sequence datasets. DNA Res 21, 341–353.

Nguyen, L.-T., Schmidt, H.A., Haeseler, A., Minh, B.Q., 2014. IQ-TREE: a fast and effective
stochastic algorithm for estimating maximum-likelihood phylogenies. Mol Biol Evol
32, 268–274.

Nieberding, C., Morand, S., Libois, R., Michaux, J., 2006. Parasites and the island syndrome:
the colonization of the western Mediterranean islands by Heligmosomoides polygyrus
(Dujardin, 1845. J Biogeogr 33, 1212–1222.

O'Connor, E.A., Hasselquist, D., Nilsson, J.-Å., Westerdahl, H., Cornwallis, C.K., 2020.
Wetter climates select for higher immune gene diversity in resident, but not
migratory, songbirds. Proc. R. Soc. B Biol. Sci. 287, 20192675.
https://doi.org/10.1098/rspb.2019.2675

Ohta, T., 1992. The nearly neutral theory of molecular evolution. Annu Rev Ecol Syst 23,
263–286.

Ortutay, C., Vihinen, M., 2009. Identification of candidate disease genes by integrating Gene
Ontologies and protein-interaction networks: case study of primary
immunodeficiencies. Nucleic Acids Res 37, 622–628.

Padilla, D.P., Illera, J.C., Gonzalez-Quevedo, C., Villalba, M., Richardson, D.S., 2017.
Factors affecting the distribution of haemosporidian parasites within an oceanic
island. Int. J. Parasitol. 47, 225–235.

Paradis, E., Schliep, K., 2019. ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. Bioinformatics 35, 526–528.

Peona, V., Blom, M.P.K., Xu, L., Burri, R., Sullivan, S., Bunikis, I., Liachko, I., Haryoko, T.,
Jønsson, K.A., Zhou, Q., 2021. Identifying the causes and consequences of assembly
gaps using a multiplatform genome assembly of a bird-of-paradise. Mol Ecol Resour
21, 263–286.

Peona, V., Weissensteiner, M.H., Suh, A., 2018. How complete are "complete" genome
assemblies?—An avian perspective. Mol Ecol Resour 18, 1188–1195.

Pérez-Rodríguez, A., Ramírez, Á., Richardson, D.S., Pérez-Tris, J., 2013. Evolution of
parasite island syndromes without long-term host population isolation: Parasite
dynamics in Macaronesian blackcaps Sylvia atricapilla. Glob Ecol Biogeogr 22,
1272–1281.

Pinheiro, J., Bates, D., DebRoy, S., Sarkar, D., Heisterkamp, S., Willigen, B., Maintainer, R.,
2017. Package 'nlme. Linear Nonlinear Mix Eff Models Version 3.

R Core Team, 2018. R: A language and environment for statistical computing.

Rando, J.C., Alcover, J.A., Illera, J.C., 2010. Disentangling Ancient Interactions: A New
Extinct Passerine Provides Insights on Character Displacement among Extinct and
Extant Island Finches. PLOS ONE 5:e12956.

Ranwez, V., Harispe, S., Delsuc, F., Douzery, E.J., 2011. MACSE: Multiple Alignment of
Coding SEquences accounting for frameshifts and stop codons. PLoS One 6:e22594.

Recuerda, M., Vizueta, J., Cuevas-Caballé, C., Blanco, G., Rozas, J., Milá, B., 2021.
Chromosome-level genome assembly of the common chaffinch (Aves: Fringilla

coelebs): a valuable resource for evolutionary biology. Genome Biol. Evol. 13,
evab034.

Robinson, J.A., Ortega-Del Vecchyo, D., Fan, Z., Kim, B.Y., Marsden, C.D., Lohmueller,
K.E., Wayne, R.K., 2016. Genomic flatlining in the endangered island fox. Curr Biol
26, 1183–1189.

Rogers, R.L., Slatkin, M., 2017. Excess of genomic defects in a woolly mammoth on
Wrangel island. PLoS Genet 13:e1006601.

Rohland, N., Reich, D., 2012. Cost-effective, high-throughput DNA sequencing libraries for
multiplexed target capture. Genome Res. 22, 939–946.

Rousselle, M., Simion, P., Tilak, M.-K., Figuet, E., Nabholz, B., Galtier, N., 2020. Is
adaptation limited by mutation? A timescale-dependent effect of genetic diversity on
the adaptive substitution rate in animals. PLoS Genet. 16, e1008668.

Santonastaso, T., Lighten, J., Oosterhout, C., Jones, K.L., Foufopoulos, J., Anthony, N.M.,
2017. The effects of historical fragmentation on major histocompatibility complex
class II β and microsatellite variation in the Aegean island reptile, Podarcis erhardii.
Ecol Evol 7, 4568–4581.

She, R., Chu, J.S.-C., Uyar, B., Wang, J., Wang, K., Chen, N., 2011. genBlastG: using
BLAST searches to build homologous gene models. Bioinforma Oxf Engl 27, 2141–
2143.

Shultz, A.J., Sackton, T.B., 2019. Immune genes are hotspots of shared positive selection
across birds and mammals. Elife 8, e41815.

Siewert, K.M., Voight, B.F., 2020. BetaScan2: Standardized Statistics to Detect Balancing
Selection Utilizing Substitution Data. Genome Biol. Evol. 12, 3873–3877.
https://doi.org/10.1093/gbe/evaa013

Silva-Iturriza, A., Ketmaier, V., Tiedemann, R., 2012. Prevalence of avian haemosporidian
parasites and their host fidelity in the central Philippine islands. Parasitol. Int. 61,
650–657.

Simion, P., Belkhir, K., François, C., Veyssier, J., Rink, J.C., Manuel, M., Philippe, H.,
Telford, M.J., 2018. A software tool 'CroCo'detects pervasive cross-species
contamination in next generation sequencing data. BMC Biol. 16, 1–9.

Singhal, S., Leffler, E.M., Sannareddy, K., Turner, I., Venn, O., Hooper, D.M., Strand, A.I.,
Li, Q., Raney, B., Balakrishnan, C.N., 2015. Stable recombination hotspots in birds.
Science 350, 928–932.

Slade, R.W., McCallum, H.I., 1992. Overdominant vs. frequency-dependent selection at
MHC loci. Genetics 132.

Slowikowski, K., Schep, A., Hughes, S., Lukauskas, S., Irisson, J.-O., Kamvar, Z.N., Ryan,
T., Christophe, D., Hiroaki, Y., Gramme, P., 2018. Package ggrepel. Autom Position
Non-Overlapping Text Labels ggplot2.

Smeds, L., Qvarnstrom, A., Ellegren, H., 2016. Direct estimate of the rate of germline
mutation in a bird. Genome Res. gr-204669.

Spiess, A.-N., Spiess, M.A.-N., 2018. Package 'qpcR, in: Model. Anal. Real-Time PCRdata
Httpscran R-Proj. OrgwebpackagesqpcRqpcR Pdf.

Spurgin, L.G., Van Oosterhout, C., Illera, J.C., Bridgett, S., Gharbi, K., Emerson, B.C.,
Richardson, D.S., 2011. Gene conversion rapidly generates major histocompatibility
complex diversity in recently founded bird populations. Mol. Ecol. 20, 5213–5225.
https://doi.org/10.1111/j.1365-294X.2011.05367.x

Tange, O., 2018. GNU parallel 2018.

van Dijk, A., Veldhuizen, E.J., Haagsman, H.P., 2008. Avian defensins. Vet. Immunol.
Immunopathol. 124, 1–18.

Van Riper III, C., Van Riper, S.G., Goff, M.L., Laird, M., 1986. The epizootiology and

ecological significance of malaria in Hawaiian land birds. Ecol. Monogr. 56, 327–
344.
Velová, H., Gutowska-Ding, M.W., Burt, D.W., Vinkler, M., 2018. Toll-like receptor
evolution in birds: gene duplication, pseudogenization, and diversifying selection.
Mol Biol Evol 35, 2170–2184.
Warren, B.H., Simberloff, D., Ricklefs, R.E., Aguilée, R., Condamine, F.L., Gravel, D.,
Morlon, H., Mouquet, N., Rosindell, J., Casquet, J., 2015. Islands as model systems in
ecology and evolution: Prospects fifty years after MacArthur-Wilson. Ecol. Lett. 18,
200–217.
Warren, W.C., Clayton, D.F., Ellegren, H., Arnold, A.P., Hillier, L.W., Künstner, A., Searle,
S., White, S., Vilella, A.J., Fairley, S., 2010. The genome of a songbird. Nature 464,
757–762.
Welch, J.J., Eyre-Walker, A., Waxman, D., 2008. Divergence and Polymorphism Under the
Nearly Neutral Theory of Molecular Evolution. J. Mol. Evol. 67, 418–426.
https://doi.org/10.1007/s00239-008-9146-9
Wickham, H., 2016. ggplot2: Elegant Graphics for Data Analysis.
Wikelski, M., Foufopoulos, J., Vargas, H., Snell, H., 2004. Galápagos birds and diseases:
invasive pathogens as threats for island species. Ecol. Soc. 9.
Wolf, J.B.W., Künstner, A., Nam, K., Jakobsson, M., Ellegren, H., 2009. Nonlinear
Dynamics of Nonsynonymous (dN) and Synonymous (dS) Substitution Rates Affects
Inference of Selection. Genome Biol Evol 1, 308–319.
Zhang, G., Parker, P., Li, B., Li, H., Wang, J., 2012. The genome of Darwin's Finch
(Geospiza fortis). https://doi.org/10.5524/100040