Bovine serum albumin as a resuscitation promoting factor for viable but non-culturable *Mycobacterium tuberculosis* via the activation of protein kinase A-dependent cellular processes

Yuta Morishige[1*], Yoshiro Murase[1], Kinuyo Chikamatsu[1], Akio Aono[1], Yuriko Igarashi[1],

Yoshiko Shimomura[1], Makiko Hosoya[1], Keisuke Kamada[1], Hiroyuki Yamada[1], Akiko

Takaki[1], and Satoshi Mitarai[1,2]

[1]Department of Mycobacterial Reference and Research, The Research Institute of Tuberculosis,

Japan Anti-Tuberculosis Association, Kiyose, Tokyo, Japan.

[2]Department of Basic Mycobacteriology, Graduate School of Biomedical Sciences, Nagasaki

University, Nagasaki, Japan.

* Correspondence:

Yuta Morishige

ymorishige@jata.or.jp

Keywords: VBNC, Mycobacterium, Albumin, Resuscitation, Dormancy

Abstract

Objective: *Mycobacterium tuberculosis* (Mtb) H37Ra strain has been reported to rapidly enter

the viable but non-culturable (VBNC) state following treatment with an NADH oxidase

inhibitor (diphenyleneiodonium [DPI]) and to be resuscitated by fetal bovine serum (FBS).

However, the mechanism underlying FBS-induced resuscitation is currently unclear. We tried

to reveal the underlying mechanism of FBS-induced resuscitation using *M. tuberculosis* H37Rv.

Methods: First, we evaluated the effect of DPI on culturability, viability and changes of cellular

phenotypes toward H37Rv. Secondly, we measured the resuscitation-promoting effects of

human serum albumin, egg-white albumin, N-acetyl-*L*-cysteine, and D-mannitol in DPI-

induced VBNC cells, as antioxidative agents have been reported to be key molecules for

resuscitation of other microbes. We also evaluated the effect of inhibition of cAMP production

and protein kinase A on BSA-induced resuscitation.

Results: DPI treatment successfully induced a VBNC state in H37Rv, resulting in a low

proportion of culturable cells, loss of acid-fastness and lipid-accumulation but a high

proportion of viable cells. Not only FBS but also bovine serum albumin (BSA) alone could

resuscitate H37Rv. Contrary to our expectation, only human serum albumin had a similar

resuscitative effect to BSA. The inhibition of adenylyl cyclase by SQ22536 did not have a

significant effect on resuscitation; however, the inhibition of protein kinase A by H89 strongly

suppressed the BSA-induced resuscitation.

Conclusion: DPI-induced VBNC Mtb cells may be resuscitated via the activation of protein

kinase A-dependent processes through interaction with BSA.

Introduction

Mycobacterium tuberculosis (Mtb) is known to as one of the intracellular parasitic bacteria

that can survive inside host macrophages. It develops a latent phenotype, dormancy, mainly

due to various stresses from host immunity. The dormant Mtb cells can persist inside the host

and cause latent tuberculosis infection (LTBI), which seeks reactivation. Lowering the risk of

Mtb reactivation in LTBI patients is considered to be one of the important strategies to reduce

the incidence of TB (World Health Organization, 2020). Dormancy in non-sporulating bacteria

that can regain culturability by certain stimuli is referred to as the viable but non-culturable

(VBNC) state, which is first reported in *Escherichia coli* and *Vibrio cholerae* (Xu et al., 1982).

Many bacteria, including Mtb, can enter into the VBNC state, and it is considered to be their

natural state in the environment (Oliver, 2010). Various *in vitro* models, such as models of

hypoxia, nutritional starvation (including potassium-deficiency), a lipid-rich environment, and

other multiple stresses are reported to to induce a dormant state in Mtb cells (Wayne and Hayes,

1996; Betts et al., 2002; Deb et al., 2009; Salina et al., 2014; Aguilar-Ayala et al., 2018). A

detailed understanding of the mechanisms of reactivation is necessary for the prevention of

active tuberculosis.

One of the major hypotheses regarding the mechanism underlying the formation of VBNC

bacteria involves oxidative damage generated by harsh external conditions (Desnues et al.,

2003). Thus, some studies have reported that the detoxification of oxidative stress during

culture is one of the key mechanisms through which VBNC bacteria are resuscitated (Mizunoe

et al., 2000; Imazaki and Nakaho, 2009). Some mycobacterial culture media contain bovine

serum albumin (BSA) and BSA on mycobacterial culture has been considered to be growth-

supporting agent that is involved in detoxification by the absorption of free fatty acid and other

growth-arresting agents that are spontaneously generated during culturing (Davis and Dubos,

1947; Dubos, 1947; Lynn et al., 1979). For this reason, the mycobacterial culture media that

are most frequently used at the present time are protein-containing systems, such as

Middlebrook 7H series or Dubos medium. However, few studies have focused on the effect of

such media on VBNC cells.

Mukamolova et al. revealed that among patients with active TB, there is certain

subpopulation for whom sputum is not culturable in conventional medium, but which becomes

to be culturable in the presence of resuscitation promoting factor (Rpf), which is a

peptidoglycan-hydrolyzing protein that is highly conserved in actinobacteria (Mukamolova et

al., 2006). Rpf reactivates non-culturable Mtb cells both *in vitro* and *in vivo* (Mukamolova et

al., 2010), suggesting the presence of VBNC Mtb cells in clinical specimens.

Previously, NADH oxidase inhibitor diphenyleneiodonium chloride (DPI) was reported to

simply and rapidly induce a VBNC state in the Mtb H37Ra strain, and incubation with fetal

bovine serum, which is a common supplement for mammalian cell cultures, could facilitate the

resuscitation of VBNC cells, while incubation with OADC (oleate-albumin-dextrose-catalase)

supplementation could not (Yeware et al., 2019). This report gave us an important clue for

constructing a simple and rapid assay system to induce and resuscitate VBNC Mtb cells.

However, when we tried to apply this system to H37Rv, we happened to find that DPI-induced

VBNC cells that could be resuscitated by not only fetal bovine serum but also by the addition

of BSA—this was observed for both H37Rv and H37Ra (internal data). Then, we hypothesized

that BSA would act as a resuscitation-promoting factor toward DPI-induced VBNC cells.

To support this hypothesis, there are some reports showing that the removal of oxidative

stress is one of the key mechanisms of resuscitation from the VBNC state (Dong et al., 2020).

The antioxidative effect of serum albumin has been well-studied (Roche et al., 2008).

Accordingly, it seems easy to hypothesize that the resuscitation-promoting effect and

antioxidative property of albumin may have a close relationship. Another possibility is that the

cellular processes involving cyclic AMP, an important second messenger of the cell. Shleeva

et al. showed that the presence of high amounts of cAMP provided by adenylyl cyclase is also

essential for resuscitation of *M. smegmatis* and Mtb from a dormant state (Shleeva et al., 2013,

2017). Notably, Mtb *Rv2212*, the major gene that encodes adenylyl cyclase (Knapp et al., 2015),

its overexpression, significantly affects entry into the dormant state, while its overexpression

causes resuscitation (Shleeva et al., 2017).

Understanding the detailed mechanism of resuscitation from the VBNC state is crucial to

reduce the risk of reactivation of Mtb cells in LTBI patients. In this study, we examined the

effects of antioxidative agents and the inhibition of adenylyl cyclase to reveal the roles of these

factors in the resuscitation of VBNC Mtb cells.

Materials and Methods

Bacterial strains, growth media and culture conditions

This work was carried out using *Mycobacterium tuberculosis* H37Rv (ATCC 27294) and

H37Ra (ATCC 25177). Both cells were cultured for 3–5 days in a 60 mL glycol-modified

polyethylene terephthalate (PETG) bottle containing 20 mL Dubos broth (2.5 g/L Na_2HPO_4,

2.0 g/L L-asparagine, 1.0 g/L KH_2PO_4, 0.5 g/L pancreatic digest of casein, 0.2 g/L Tween 80,

0.5 mg/mL $CaCl_2 \cdot 2H_2O$, 0.1 mg/mL $CuSO_4$, 0.1 mg/mL $ZnSO_4 \cdot 7H_2O$, 0.05 g/L ferric

ammonium citrate, 0.01g $MgSO_4 \cdot 7H_2O$) with 5% (v/v) glycerol, and 10% (v/v) ADC

supplementation (5.0 g bovine serum albumin fraction V (BSA), 2.0 g dextrose, 0.003 g

catalase in 100 mL distilled water) at 37°C in an orbital shaker (100 rpm) until the OD_{600}

reached up to 0.35. All procedures were performed at a BSL-3 facility. For further analysis, we

used commercially available BSA Cohn fraction V (Merck, Darmstadt, Germany), unless

otherwise stated. In addition, we used molecular biology grade Tween 80 (Merck product

number P5188-100ML) without any predilution.

Counting the number of culturable cells

The cultured bacterial suspension was serially diluted 10-fold in phosphate buffered saline (pH

6.8) with 0.1%(w/v) Tween 80 (PBS-T) and 25 µL of each diluent was inoculated onto a

Middlebrook 7H10 agar plate supplemented with oleic-albumin-dextrose-catalase (OADC)

(Becton Dickinson, Sparks, MD) in duplication. Colonies were counted after at least 3 weeks'

incubation at 37°C under 5% CO_2. The limit of detection was determined to be 20 CFU/mL

because of the diluent factor.

Effect of BSA on the growth of Mtb cells in Dubos medium

Ten milliliters of log phase culture of Mtb H37Rv that reached an OD_{600} of up to 0.35 in Dubos

medium, as described previously, was sedimented at 3,000 ×*g* for 10 minutes at 4°C and the

supernatant was discarded. The sediment was washed twice with fresh Dubos broth and

resuspended with 10 mL of Dubos broth. The suspension was diluted to 1:10 with Dubos broth

with or without the addition of BSA solution (5.0 g BSA fraction V in 100 mL distilled water)

at the final concentration of 0.1% (w/v) BSA. The diluted suspension was dispensed into 24-

well plates (Sarstedt, Nümbrecht, Germany) in 1.5 mL increments and sealed with gas-

permeable film (Breathe-Easy, Diversified Biotech Inc, Dedham, MA, United States). Cultures

were incubated for 5, 10, 15 and 20 days at 37°C under 5% CO_2. At each time-point, the number

of culturable cells was measured as described above.

Induction of the VBNC cells

Induction of the VBNC cells were performed according to the previous report (Yeware et al.,

2019) with some modification. In short, the log phase culture of Mtb H37Rv or H37Ra that

reached an OD_{600} of up to 0.35 in Dubos medium, as previously described, was evenly divided

into 30 mL PETG bottles and 5 mg/mL diphenyleneiodonium chloride (DPI) in dimethyl

sulfoxide (DMSO) solution was directly added at the final concentration of 4 µg/mL with

immediate agitation. The same volume of DMSO was added to the untreated control sample.

Cultures were further incubated for 24 h at 37°C without shaking. After incubation, their

culturability and viability were measured as described above.

Cell viability assay

Mtb cell viability was measured using esterase activity and membrane integrity of the cells

using dual staining with 10 mg/mL carboxyfluorescein diacetate (CFDA, Dojindo, Kumamoto,

Japan) and ethidium bromide (EB, Dojindo). Briefly, 500 µL of DPI-treated or untreated Mtb

culture was sedimented by centrifugation at 15,000 ×*g* for 5 minutes at 4°C and washed twice

with the same volume of PBS-T. The washed bacterial sediment was resuspended in 200 µL of

PBS-T with CFDA and EB incubated at room temperature (RT) for 30 minutes in shade. The

final concentration of CFDA and EB were 60 µg/mL and 2 µg/mL, respectively. Stained Mtb

cells were washed twice with PBS-T and fixed with 3.7% formaldehyde solution at RT for at

least 2 h. After fixation, 10 µL of bacterial suspension was smeared on three APS-coated 8-

well slides (Matsunami glass, Osaka, Japan) and air-dried. Then the slides were mounted with

DPX new non aqueous mounting medium (Merck, Darmstadt, Germany) and a 150 µm-thick

cover glass (Matsunami glass, Osaka, Japan) prior to the microscopic observation. Specimens

were examined with a BX53 fluorescence microscope (Olympus, Tokyo, Japan) at a

magnification of 100× for green (esterase-active with intact membrane; live) or red (esterase-

negative with damaged membrane; dead) fluorescing cells. The excitation source for CFDA

and EB was generated from a mercury-arc lamp (blue beam, 470–495 nm; green beam, 530–

550nm) using a corresponding band-pass filter. The green-fluorescence emission from the

CFDA-positive cells was collected through a 510–550 nm band-pass filter. The red-

fluorescence emission from EB-positive cells was collected through a 575 nm long-pass filter.

At least 10 random fields were observed for each sample and the images (TIFF format) were

analyzed using the Fiji/ImageJ software program (Schindelin et al., 2012). The live/dead ratio

was calculated by direct counting of green or red fluorescing cells.

Auramine-O/Nile Red dual staining

Loss of acid-fastness and accumulation of neutral lipids are distinctive features of dormant

mycobacteria (Garton et al., 2002). To detect the phenotype, fluorescent acid-fast staining with

auramine-O and neutral lipid staining with Nile Red (9-dimethylamino-5H-benzo-α-

phenoxadine-5-one) were performed using a previously described method (Deb et al., 2009)

with some modification. Briefly, 10 µL of DPI-treated or untreated culture were smeared onto

an APS-coated slide (Matsunami glass, Osaka, Japan) and air-dried. Then, the slide was heat-

fixed and cooled down to room temperature before staining. The smear was flooded with

fluorochrome staining solution (10 µg/mL auramine-O in 5% [w/v] phenol solution) and

incubated at room temperature for 20 min in shade. Excessive dye was removed using 3%

hydrochloric acid ethanol for 15 minutes. Next, the smear was covered with 10 µg/mL Nile

Red in ethanol and incubated at room temperature for 15 minutes. Finally, the smear was

counterstained with 0.1% (w/v) potassium permanganate solution with 1 minute for

counterstaining. Stained slides were air-dried and mounted as described previously.

Fluorescence microscopy was performed at 100× magnification for green (acid-fastness

positive, lipid accumulation negative; active) or red (acid-fastness negative, lipid accumulation

positive; dormant) fluorescing cells. The excitation source for auramine-O and Nile Red was

generated from a mercury-arc lamp (blue beam, 470–495 nm; green beam, 530–550 nm) using

a corresponding band-pass filter. The green-fluorescence emission from auramine-O-positive

cells was collected through a 510–550 nm band-pass filter. The red-fluorescence emission from

Nile Red-positive cells was collected through 575 nm long-pass filter. At least 5 random fields

were observed for each sample and images (TIFF format) were analyzed by the Fiji/ImageJ

software program and the proportion of acid-fast-positive cells and lipid-accumulation-positive

cells was calculated by direct counting of green or red fluorescing cells.

Ziehl-Neelsen staining

Ziehl-Neelsen staining was also performed to confirm the loss of acid-fastness resulting in the

VBNC state by DPI treatment, according to the standard method. Stained slides were air-dried

and observed by light microscope under 100x magnification with an oil immersion lens (BX53,

Olympus, Tokyo, Japan).

Airyscan super-resolution microscopy

Detailed images of accumulated lipids were also obtained by a confocal laser-scanning

microscope (LSM900 with Airyscan 2, Carl Zeiss Microscopy, Jena, Germany). Briefly, culture

slides of DPI-treated cells dual stained with auramine-O and Nile Red, as described above,

were observed at 63x magnification with oil immersion lens. The excitation source of

auramine-O and Nile Red was 488 nm (blue beam) and 563 nm (green beam), from a solid-

state laser unit. The green-fluorescence emission from auramine-O and red fluorescence from

Nile Red were sequentially collected through a variable dichroic beamsplitter and an Airyscan

2 detector. Super-resolution images were processed with the ZEN Blue software program (Carl

Zeiss, Jena, Germany).

Resuscitation assay

The resuscitation assay was performed as described elsewhere (Yeware et al., 2019) with some

modification. DPI-treated culture was sedimented by centrifugation at 3,000 xg for 10 minutes

at 4˚C and washed twice with fresh Dubos broth, then resuspended in Dubos broth. The

bacterial suspension was diluted to 1:10 with fresh Dubos medium with or without 0.1% (w/v)

BSA. For both BSA-containing and BSA-free series, OADC supplementation (Becton

Dickinson, Sparks, MD), fetal bovine serum (FBS; Moregate BioTech, Bulimba, Queensland,

Australia), or sodium pyruvate (PA; Merck) was added. The final concentration of each reagent

was as follows: 10% (v/v) for OADC supplementation, 2% (v/v) for FBS, and 3 mM for PA.

Then, the diluted suspension was dispensed into 24-well plates in 1.5 mL increments and sealed

with gas-permeable film. Cultures were incubated for 5, 10, 15 and 20 days at 37˚C under 5%

CO_2. At each time-point, the number of culturable cells was measured as described previously.

Evaluation of the resuscitation-promoting effect of human serum albumin, egg-white

albumin (ovalbumin) and antioxidative agents

To determine whether the resuscitation-promoting effect of BSA is its antioxidative property,

we measured the resuscitation-promoting activity of other albumins and antioxidative agents

toward DPI-treated Mtb cells. Washed DPI-treated cells were resuspended with fresh Dubos

broth without BSA and the suspension was diluted to 1:10 with Dubos broth containing BSA,

human serum albumin (HSA), ovalbumin (OVA), N-acetyl-L-cysteine (NAC) or D-mannitol

(MAN). The final concentration of each reagent was 0.1% (w/v) for BSA, HSA and OVA, 500

μM for NAC, and 50 mM for MAN, respectively. Then, the diluted suspension was dispensed

into 24-well plates in 1.5 mL increments and sealed with gas-permeable film. The plate was

incubated for 20 days at 37˚C under 5% CO_2. At the end of incubation, the number of colonies

grown was measured as described previously.

Evaluation of the resuscitation-promoting effect of fatty acid and globulin-free BSA

To determine the effect of BSA Cohn fraction V contaminants, we measured the resuscitation-

promoting activity of fatty acid and globulin-free BSA toward DPI-treated VBNC Mtb cells.

Washed DPI-treated cells were resuspended with fresh Dubos broth without BSA and the

suspension was diluted to 1:10 with Dubos broth containing BSA Cohn fraction V or fatty acid

and globulin-free BSA. Fatty acid and globulin-free BSA were acquired from Merck. The final

concentration of both BSAs was 0.1% (w/v). The number of culturable cells was measured as

described previously.

Effects of adenylyl cyclase or protein kinase A on BSA-induced resuscitation

Washed DPI-treated VBNC Mtb cells were resuspended with fresh Dubos broth and the

suspension was diluted to 1:10 with Dubos broth with 0.1% (w/v) BSA containing adenylyl

cyclase inhibitor SQ22536 (Merck, Darmstadt, Germany) or protein kinase A inhibitor H89

(Abcam, Cambridge, United Kingdom). The final concentration of each inhibitor was as

follows: 0.1 to 10 mM for SQ22536 and 1 to 30 μM for H89. Then, the number of culturable

cells were measured as described previously.

Statistical analysis

The statistical analysis was performed with R (R Core Team, 2021) using the EZR on R

package (Kanda, 2013). *P* values of < 0.05 were considered statistically significant.

Results

I.   BSA is not essential for the growth of MTB in Dubos medium with purified non-

prediluted Tween 80

We first analyzed the different effects of BSA-containing and BSA-free Dubos medium on

the growth of Mtb H37Rv cells. As shown in Fig. 1(A), there was little difference in the rate

of growth between the two conditions. Fig. 1(B) also shows the sufficient growth of cells. It

243 seemed that the cells may have been more aggregative without BSA, however, the aggregation

244 was easily homogenized by pipetting and there was no significant difference in the OD_{600} value

245 (data not shown). These results suggest that Mtb could grow normally regardless of the albumin

246 component.

247 **II. DPI treatment could induce a VBNC state in Mtb H37Rv**

248 Next, we confirmed that DPI treatment could induce a VBNC state in the H37Rv strain, in

249 addition to H37Ra, which was previously reported (Yeware et al., 2019). As shown in Fig.

250 2(A), the numbers of culturable cells of untreated control population and DPI-treated

251 population were significantly different: $4.43 \pm 1.43 \times 10^8$ CFU/mL for the untreated control

252 population and $4.64 \pm 6.10 \times 10^3$ CFU/mL for DPI-treated population. Fig. 2(B) shows that the

253 CFDA-positive cells in the untreated control and DPI-treated populations were $81.8 \pm 11.4\%$

254 and $65.9 \pm 11.8\%$, respectively. These results suggested that the majority of DPI-treated H37Rv

255 cells were alive, while their culturability was significantly reduced ($p = 0.033$). Representative

256 images of CFDA/EB dual-stained cells of the untreated control and DPI-treated population also

257 showed that there was little difference in the proportions of CFDA-positive and EB-positive

258 cells. (Fig. 2[C])

259 Auramine-O/Nile Red dual staining showed the reduction of acid-fastness and the

260 accumulation of the neutral lipid, triacylglycerol, which are common morphological features

261 of dormant Mtb cells.(Garton et al., 2002; Bhatt et al., 2007; Deb et al., 2009; Kapoor et al.,

262 2013) As shown in Fig. 2(D), the proportion of acid-fast-negative/lipid accumulation-positive

263 cells was $84.0 \pm 14.0\%$ in the DPI-treated population, and $12.1 \pm 11.9\%$ in the untreated control

264 population. Representative images of auramine-O/Nile Red dual-stained cells of the untreated

265 control and DPI-treated populations also showed that there were distinctive differences in the

266 proportions of acid-fast-positive cells and lipid-accumulation-positive cells (Fig. 2[E]). The

267 loss of acid-fastness was also confirmed by Ziehl-Neelsen staining; the majority of DPI-treated

population were found to have lost their acid-fastness (Suppl. Fig. 1). Interestingly, Airyscan

super-resolution microscopy revealed that were some DPI-treated cells that were both acid-

fast-positive and neutral lipid stain-positive (Fig. 2[F]). This could also reveal the distribution

of the lipid body in the cell as some foci of relatively strong signals of Nile Red, suggesting

that the transition from culturable to VBNC state. These results indicated that DPI treatment

successfully induced the VBNC state in H37Rv as well as in H37Ra.

III.   The effects of BSA, FBS, OADC and sodium pyruvate on the resuscitation of

DPI-treated Mtb cells

As described elsewhere, Yeware et al. first reported that the resuscitation of DPI-induced

VBNC Mtb cells was only achieved by incubation with FBS; OADC could not facilitate

resuscitation. As shown in Fig. 3(A) and (B), our results partially support their study with some

differences; although the culturability of DPI-treated H37Rv cells was regained by incubation

with FBS-supplemented medium, OADC-supplemented medium could also induce

culturability. Surprisingly, the addition of albumin alone into the medium could also induce

reactivation. The addition of FBS in BSA-free Dubos medium slightly reduced the regrowth

rate; however, cells were successfully resuscitated at the end of incubation with or without FBS.

These phenomena were also observed in H37Ra (Suppl. Fig 2). Thus, we used the H37Rv strain

for the further analyses in this study.

Incubation with sodium pyruvate, which was reported to have a resuscitation-promoting

effect on VBNC cells elsewhere, led to transient regrowth by day 15 and a slight reduction at

day 20. (Suppl. Fig. 3[A] and [B]).

IV.   The resuscitation-promoting effect of albumin is not due to its antioxidative

property or the supplementation of fatty acid from BSA.

As shown in Fig. 4(A) and (B), the resuscitation-promoting effect of albumin was specific

to bovine and human serum albumin. Ovalbumin did not show a resuscitation-promoting effect

but maintained the number of culturable cells in this system. Furthermore, N-acetyl-*L*-cysteine

(antioxidative agent) and D-mannitol (free radical scavenger) did not show any such effects.

At the end of incubation, there were no detectable culturable cells in either the NAC-treated or

MAN-treated cells.

We also checked whether the purity of albumin affects the promotion of resuscitation using

ethanol and heat-treated albumin and confirmed that there was no significant difference in

resuscitation-promoting effects of normal, ethanol-fractionated albumin, and fatty acid-free

albumin (Suppl. Fig 4). These results suggest that impurities of commercially available

albumin including fatty acids do not affect the regrowth of DPI-treated Mtb.

V.   The resuscitation-promoting effect of albumin is canceled by treatment with

protein kinase A inhibitor.

As shown in Fig. 5(A) and (B), SQ22536, adenylyl cyclase inhibitor, only suppressed

resuscitation at a very high concentration (10 mM) in this study. However, H89, a protein

kinase A inhibitor, suppressed regrowth in a dose-responsive manner. In detail, incubation with

10 µM or 30 µM H89 resulted in a CFU/mL value for 3.42 $\times$ $10^6 \pm 7.45 \times 10^5$ CFU/mL and

$3.63 \times 10^2 \pm 1.70 \times 10^2$ CFU/mL, while incubation without H89 resulted in $3.00 \times 10^8 \pm 5.07 \times$

10^7 CFU/mL. We should note that neither SQ22536 nor H89 caused a drastic reduction of the

growth of intact MTB cells. (Suppl. Fig. 5A and 5B). These results suggest that protein kinase

A might mainly be involved with the resuscitation of VBNC Mtb cells triggered by BSA and

the contribution of adenylyl cyclase may be smaller than that of protein kinase A.

Discussion

In this study, we demonstrated that the resuscitation of DPI-induced VBNC Mtb cells could

be facilitated not only by incubation with FBS but also by incubation with albumin-containing

supplements like OADC supplementation or BSA alone. We also showed that the resuscitation-

promoting effect of albumin may not be due to its antioxidative property. Furthermore, we

suggested that the BSA-induced resuscitation may occur via the activation of protein kinase A-

dependent processes.

Firstly, we found that DPI-induced VBNC could facilitate resuscitation not only by

incubation with FBS but also with OADC supplementation and BSA alone, suggesting albumin

might act as a resuscitation-promoting agent in both H37Rv and H37Ra (Fig .3 and Suppl. Fig.

2). These findings were contrary to the findings of our previous study. We considered the

reason for the difference may be due to contaminant in Tween 80. As described previously, the

purity of Tween 80 seemed to be important for the growth of Mtb cells. We also confirmed that

the resuscitation was not facilitated by the use of prediluted Tween 80 for the preparation of

Dubos medium (internal data) for either H37Rv and H37Ra. Thus, our results suggested that

the key component of resuscitation-promoting agent in FBS might be serum albumin, and

further analyses were performed using H37Rv. We also tested this effect in Wayne's hypoxic

culture, which is widely used for inducing a dormant state of Mtb, and found that there was no

significant difference according to the presence or absence of BSA (data not shown).

Pyruvate, which is known to act as resuscitation promoting agent for both gram-negative

bacteria, such as *Eshcerichia, Legionella, Vibrio* and gram-positive bacteria such as

Staphylococcus (Pasquaroli et al.; Asakura et al., 2005, 2007; Ducret et al., 2014), did not show

any resuscitation-promoting effect in our experiment (Suppl. Fig.3). Vilhena et al. reported that

their proteomic analysis of VBNC *E. coli* cells during resuscitation revealed that enzymes

involved in pyruvate metabolism, pyruvate formate-lyase-activating protein (PflA),

phosphoenolpyruvate carboxykinase (PckA) and lactate dehydrogenase (LldD), were

significantly upregulated (Vilhena et al., 2019), while Wagley et al. also reported that

VPA1499, an LldD homologue of Vibrio parahaemolyticus, was significantly upregulated in

VBNC *V. parahaemolyticus* and that depletion of LldD led to quicker entry to the VBNC state

(Wagley et al., 2021). Interestingly, they showed that the addition of pyruvate, the product of

lactate oxidation, to LldD-depleted mutant restored the wild-type characteristics of cells in the

VBNC state. Their studies might give some crew of the lack of resuscitation-promoting effect

toward mycobacterial VBNC. Since Mtb does not have a catalytic center of pyruvate formate-

lyase (PflB), pyruvate might not be able to act as resuscitation-promoting agent, although

pyruvate enhances the growth of Mtb and *M. bovis* both intracellular and extracellular growth

(Schaefer, 1952; Osada-Oka et al., 2019).

Our results showed that the resuscitation promoting effect of albumin might be structure-

specific because ovalbumin (OVA) did not facilitate resuscitation, but bovine (BSA) and

human serum albumin (HSA) did. Focusing on their amino acid sequences, BSA and HSA

share 76% sequence homology (Huang et al., 2004); however, BSA and OVA do not have

sequence homology. One of the considerable roles of albumin may be as an antioxidative agent

due to its cysteine residues with free thiols (Roche et al., 2008; Rombouts et al., 2015). BSA

and HSA have only one cysteine residue with free thiol, while OVA has four cysteines with

free thiols (Broersen et al., 2006). It therefore seems to be more reductive than BSA and HSA.

Our results were paradoxical in relation to this hypothesis; however, Davis and Dubos showed

the differential growth-promoting effect of albumin toward Mtb. Their study supplied some

interesting clues for our present study; ovalbumin does not show a growth-promoting effect,

while bovine and human serum albumin showed the effect (Davis and Dubos, 1947). Moreover,

no antioxidative agents, such as N-acetyl-*L*-cysteine and D-mannitol, showed any resuscitative

effects (Fig. 4). We also examined catalase, a component of OADC supplementation, and its

resuscitation promoting effect toward VBNC *Corynebacterium, Lactobacillus, Salmonella,*

Ralstonia and *Vibrio* (Kong et al., 2004, 2014; Senoh et al., 2015; Liu et al., 2017; Morishige

et al., 2017; Hamabata et al., 2021); no resuscitative effects were observed (data not shown).

These results suggest that the effect of albumin may be due to complex functions rather than

its antioxidative property.

Serum albumin is known to act as a good carrier of many kinds of biologically active

substances, including free fatty acid. In general, BSA is added to medium for mycobacterial

culture as a carrier of growth-supporting agents, including fatty acids. Shleeva et al. showed

that several types of unsaturated free fatty acid, such as oleic, linoleic, and arachidonic acids,

can promote the resuscitation of VBNC *M. smegmatis* in modified Sauton's medium and oleic

acid also works for Mtb (Shleeva et al., 2013). In this study, we should note that impurities of

albumin did not affect resuscitation. Although the underlying mechanism is still unclear, fatty

acid and globulin-free BSA and BSA Cohn fraction V did not show any difference toward

resuscitation of DPI-treated Mtb (Suppl. Fig. 4).

The assay of inhibition of resuscitation by SQ22536 and H89 gave us an important clue for

understanding resuscitation. Shleeva et al. showed that the presence of high amounts of cAMP

provided by adenylyl cyclase is essential for resuscitation from a dormant state in *M. smegmatis*

and *M. tuberculosis* (Shleeva et al., 2013, 2017). Especially in *M. tuberculosis*, the

overexpression of *Rv2212*, the major gene that encodes adenylyl cyclase (Knapp et al., 2015),

significantly affects both entry into the dormant state and resuscitation (Shleeva et al., 2017).

In the present study, we could suppress the resuscitation of DPI-induced VBNC cells by

inhibiting adenylyl cyclase only with a very high concentration of SQ22536. Although the

underlying mechanism is unclear, adenylyl cyclase might not play a key role in resuscitation

from a DPI-induced VBNC state. However, the inhibition of protein kinase A by H89 clearly

suppressed resuscitation. Mtb has the eukaryotic-type Ser/Thr protein kinase PknA, which

regulates major cell processes, regulates several types of proteins involved with mycobacterial

cell division and morphogenesis via phosphorylation of such proteins in coordination with

protein kinase B (PknB; Manuse et al., 2016; Carette et al., 2018). For example,

phosphorylation of GlmU (*Rv1018c*) and MurD (*Rv2155c*), Wag31 (*Rv2145c*), HupB

(*Rv2986c*) and ParB (*Rv3917c*), influence the production and cross-linking of peptidoglycan

precursors (Thakur and Chakraborti, 2008; Parikh et al., 2009; Gee et al., 2012; Jagtap et al.,

2012), the spatial localization of peptidoglycan synthesis (Kang et al., 2008), DNA

condensation and partitioning (Gupta et al., 2014; Baronian et al., 2015), respectively. It should

be noted that the phosphorylation of FtsZ and FipA by PknA controls cell division under

oxidative stress (Thakur and Chakraborti, 2006; Sureka et al., 2010). Our study suggested that

the inhibition of PknA by H89 seems to critically affect several important cellular processes,

followed by resuscitation (Fig. 6). To confirm the essentiality of PknA and other involved

proteins toward resuscitation from this DPI-induced VBNC state by albumin, further studies

using experiments such as knock-down of genes using the CRISPR interference (CRISPRi)

technique (Rock et al., 2017), because such proteins—with the exception of FipA (*Rv0019c*)—

are known to be essential for *in vitro* growth (Sassetti et al., 2003; Griffin et al., 2011; DeJesus

et al., 2017; Minato et al., 2019); thus, manipulating their coding genes is not permissible.

We also confirmed that the absence of BSA in Dubos medium did not show any significant

effect on the growth of Mtb (Fig.1[A]). This may be due to the purity of Tween 80 because

Dubos et al. previously reported that the presence of unpurified Tween 80 may inhibit growth

due to release of free oleic acid (Davis and Dubos, 1947). Our result was in good concordance

with that study. We also confirmed that the use of prediluted Tween 80 (20% w/v) in BSA-free

Dubos medium clearly arrested the growth of Mtb (internal data). The concentration of free

fatty acid in the medium with prediluted Tween 80 was approximately 80 µM. On the other

hand, the concentration in the medium with non-prediluted Tween 80 was only approximately

7 µM (unpublished data), suggesting that excessive concentrations of fatty acid are toxic for

Mtb (Dubos, 1950).

In addition, we confirmed that DPI could successfully induce a VBNC state in H37Rv as

well as H37Ra. The mechanism underlying the effect of DPI is considered to involve the

inhibitory effect of NADH oxidase, which results in the inhibition of the electron transport

system of Mtb. This may induce a hypoxic state in the liquid culture, resulting in significant

decrease of culturability (Fig.2[A]). However, viability was highly retained (65.9%),

suggesting that majority of the cells transitioned to VBNC state (Fig. 2[B]). Additionally,

transition to VBNC state was also confirmed by auramine-O and Nile Red staining. The loss

of acid-fastness and the accumulation of neutral lipids are distinctive features of dormant

mycobacterial cells, and can be detected by acid-fast staining and neutral lipid staining. Our

data also demonstrated that the majority of DPI-treated Mtb H37Rv cells were stained with

Nile Red alone (Fig. 2[D]). Airyscan super-resolution microscopy showed the presence of cells

that were stained with auramine-O and which contained an intracellular lipid body stained with

Nile Red (Fig. 2[F]), which may indicate that the cells were on the way to the dormant state

from an actively growing state, with drastic alteration of the lipid metabolism.

To the best of our knowledge, few studies have focused on the resuscitation-promoting effect

of albumin toward VBNC Mtb. Taken together, our findings indicate that the resuscitation of

MTB from a DPI-induced VBNC state could be obtained by interaction with bovine serum

albumin and the mechanism of albumin-induced resuscitation may involve the activation of

protein kinase A, which regulates several important processes of cellular component

construction and cell division. Although it remains unclear how albumin interacts with VBNC

cells, our study may supply some important clues to understanding the physiology of the

mycobacterial VBNC state.

Conclusion

Our data showed that the presence of BSA in Dubos medium could trigger the resuscitation

of the DPI-induced VBNC state in both H37Rv and H37Ra strains. The resuscitation-

promoting effect of other kinds of albumins and antioxidative agents, including N-acetyl-*L*-

cysteine and D-mannitol was tested; however, only human serum albumin showed a

resuscitation-promoting effect. The results of the assay of resuscitation using SQ22536 and

H89 demonstrated that the inhibition of protein kinase A suppressed resuscitation to a much

greater extent than adenylyl cyclase, indicating the presence of a protein kinase A-dependent

resuscitation-promoting pathway.

Author Contributions

YMori, YMura and SM conceived and designed the experiments. YMori performed

experiments. YMura, KC, AA, YI, YS, MH, KK, HY, AT, and SM assisted in the experiments.

YMori acquired data and YMori and YMura analyzed data. YMori performed the statistical

analysis. YMori and SM wrote the manuscript. All authors contributed to the article and

approved the submitted version.

Funding

This work was supported by Research Program on Emerging and Re-emerging Infectious

Diseases of Japan Agency for Medical Research and Development (AMED) and Grant-in-Aid

for Early-Career Scientists of Japan Society for the Promotion of Science (JSPS) KAKENHI.

Grant Numbers are JP21fk0108090 and 21K15442, respectively.

Acknowledgments

Authors thank K. Suenaga and H. Sakuma (Carl Zeiss Co., Ltd., Tokyo, Japan) for their kind

assistance in obtaining super-resolution images. Authors also thank Dr. A. Osugi (The Research

Institute of Tuberculosis, Japan Anti-Tuberculosis Association) for valuable discussions.

Authors also thank Dr. Brian Quinn (Japan Medical Communication. Inc.) for the help in the

preparation of this manuscript.

Conflict of Interest Statement

The authors declare that the research was conducted in the absence of any commercial or

financial relationships that could be construed as a potential conflict of interest.

References

Aguilar-Ayala, D. A., Cnockaert, M., Vandamme, P., Palomino, J. C., Martin, A., and

Gonzalez-Y-Merchand, J. (2018). Antimicrobial activity against *Mycobacterium*

tuberculosis under in vitro lipid-rich dormancy conditions. *J. Med. Microbiol.* 67, 282–

285. doi:10.1099/jmm.0.000681.

Asakura, H., Igimi, S., Kawamoto, K., Yamamoto, S., and Makino, S.-I. (2005). Role of *in*

vivo passage on the environmental adaptation of enterohemorrhagic *Escherichia coli*

O157:H7: cross-induction of the viable but nonculturable state by osmotic and oxidative

stresses. *FEMS Microbiol. Lett.* 253, 243–9. doi:10.1016/j.femsle.2005.09.039.

Asakura, H., Ishiwa, A., Arakawa, E., Makino, S., Okada, Y., Yamamoto, S., et al. (2007).

Gene expression profile of *Vibrio cholerae* in the cold stress-induced viable but non-

culturable state. *Environ. Microbiol.* 9, 869–79. doi:10.1111/j.1462-2920.2006.01206.x.

Baronian, G., Ginda, K., Berry, L., Cohen-Gonsaud, M., Zakrzewska-Czerwińska, J.,

Jakimowicz, D., et al. (2015). Phosphorylation of *Mycobacterium tuberculosis* ParB

participates in regulating the ParABS chromosome segregation system. *PLoS One* 10,

e0119907. doi:10.1371/journal.pone.0119907.

Betts, J. C., Lukey, P. T., Robb, L. C., McAdam, R. A., and Duncan, K. (2002). Evaluation of

a nutrient starvation model of *Mycobacterium tuberculosis* persistence by gene and

protein expression profiling. *Mol. Microbiol.* 43, 717–31. doi:10.1046/j.1365-

2958.2002.02779.x.

Bhatt, A., Fujiwara, N., Bhatt, K., Gurcha, S. S., Kremer, L., Chen, B., et al. (2007). Deletion

of kasB in *Mycobacterium tuberculosis* causes loss of acid-fastness and subclinical

latent tuberculosis in immunocompetent mice. *Proc. Natl. Acad. Sci. U. S. A.* 104, 5157–

62. doi:10.1073/pnas.0608654104.

Broersen, K., Van Teeffelen, A. M. M., Vries, A., Voragen, A. G. J., Hamer, R. J., and De

Jongh, H. H. J. (2006). Do sulfhydryl groups affect aggregation and gelation properties

of ovalbumin? *J. Agric. Food Chem.* 54, 5166–74. doi:10.1021/jf0601923.

Carette, X., Platig, J., Young, D. C., Helmel, M., Young, A. T., Wang, Z., et al. (2018).

Multisystem Analysis of *Mycobacterium tuberculosis* Reveals Kinase-Dependent

Remodeling of the Pathogen-Environment Interface. *MBio* 9. doi:10.1128/mBio.02333-

17.

Davis, B. D., and Dubos, R. J. (1947). THE BINDING OF FATTY ACIDS BY SERUM

ALBUMIN, A PROTECTIVE GROWTH FACTOR IN BACTERIOLOGICAL

MEDIA. *J. Exp. Med.* 86, 215–228. doi:10.1084/jem.86.3.215.

Deb, C., Lee, C. M., Dubey, V. S., Daniel, J., Abomoelak, B., Sirakova, T. D., et al. (2009).

A novel in vitro multiple-stress dormancy model for *Mycobacterium tuberculosis*

generates a lipid-loaded, drug-tolerant, dormant pathogen. *PLoS One* 4, e6077.

doi:10.1371/journal.pone.0006077.

DeJesus, M. A., Gerrick, E. R., Xu, W., Park, S. W., Long, J. E., Boutte, C. C., et al. (2017).

Comprehensive Essentiality Analysis of the *Mycobacterium tuberculosis* Genome via

Saturating Transposon Mutagenesis. *MBio* 8. doi:10.1128/mBio.02133-16.

Desnues, B., Cuny, C., Grégori, G., Dukan, S., Aguilaniu, H., and Nyström, T. (2003).

Differential oxidative damage and expression of stress defence regulons in culturable

and non-culturable *Escherichia coli* cells. *EMBO Rep.* 4, 400–404.

doi:10.1038/SJ.EMBOR.EMBOR799.

Dong, K., Pan, H., Yang, D., Rao, L., Zhao, L., Wang, Y., et al. (2020). Induction, detection,

formation, and resuscitation of viable but non-culturable state microorganisms. *Compr.*

Rev. Food Sci. Food Saf. 19, 149–183. doi:10.1111/1541-4337.12513.

Dubos, R. J. (1947). THE EFFECT OF LIPIDS AND SERUM ALBUMIN ON

BACTERIAL GROWTH. *J. Exp. Med.* 85, 9–22. doi:10.1084/jem.85.1.9.

Dubos, R. J. (1950). THE EFFECT OF ORGANIC ACIDS ON MAMMALIAN TUBERCLE

BACILLI. *J. Exp. Med.* 92, 319–332. doi:10.1084/JEM.92.4.319.

Ducret, A., Chabalier, M., and Dukan, S. (2014). Characterization and resuscitation of "non-

culturable" cells of *Legionella pneumophila*. *BMC Microbiol.* 14, 3. doi:10.1186/1471-

2180-14-3.

Garton, N. J., Christensen, H., Minnikin, D. E., Adegbola, R. A., and Barer, M. R. (2002).

Intracellular lipophilic inclusions of mycobacteria in vitro and in sputum. *Microbiology*

148, 2951–2958. doi:10.1099/00221287-148-10-2951.

Gee, C. L., Papavinasasundaram, K. G., Blair, S. R., Baer, C. E., Falick, A. M., King, D. S.,

et al. (2012). A phosphorylated pseudokinase complex controls cell wall synthesis in

mycobacteria. *Sci. Signal.* 5, ra7. doi:10.1126/scisignal.2002525.

Griffin, J. E., Gawronski, J. D., Dejesus, M. A., Ioerger, T. R., Akerley, B. J., and Sassetti, C.

M. (2011). High-resolution phenotypic profiling defines genes essential for

mycobacterial growth and cholesterol catabolism. *PLoS Pathog.* 7, e1002251.

doi:10.1371/journal.ppat.1002251.

Gupta, M., Sajid, A., Sharma, K., Ghosh, S., Arora, G., Singh, R., et al. (2014). HupB, a

nucleoid-associated protein of *Mycobacterium tuberculosis*, is modified by

serine/threonine protein kinases in vivo. *J. Bacteriol.* 196, 2646–57.

doi:10.1128/JB.01625-14.

Hamabata, T., Senoh, M., Iwaki, M., Nishiyama, A., Yamamoto, A., and Shibayama, K.

(2021). Induction and Resuscitation of Viable but Nonculturable *Corynebacterium*

diphtheriae. *Microorganisms* 9. doi:10.3390/microorganisms9050927.

Huang, B. X., Kim, H. Y., and Dass, C. (2004). Probing three-dimensional structure of

bovine serum albumin by chemical cross-linking and mass spectrometry. *J. Am. Soc.*

Mass Spectrom. 15, 1237–1247. doi:10.1016/J.JASMS.2004.05.004.

Imazaki, I., and Nakaho, K. (2009). Temperature-upshift-mediated revival from the sodium-
pyruvate-recoverable viable but nonculturable state induced by low temperature in
Ralstonia solanacearum: linear regression analysis. *J. Gen. Plant Pathol. 2009 753* 75,
213–226. doi:10.1007/S10327-009-0166-0.

Jagtap, P. K. A., Soni, V., Vithani, N., Jhingan, G. D., Bais, V. S., Nandicoori, V. K., et al.
(2012). Substrate-bound crystal structures reveal features unique to *Mycobacterium*
tuberculosis N-acetyl-glucosamine 1-phosphate uridyltransferase and a catalytic
mechanism for acetyl transfer. *J. Biol. Chem.* 287, 39524–37.
doi:10.1074/jbc.M112.390765.

Kanda, Y. (2013). Investigation of the freely available easy-to-use software "EZR" for
medical statistics. *Bone Marrow Transplant.* 48, 452–8. doi:10.1038/bmt.2012.244.

Kang, C.-M., Nyayapathy, S., Lee, J.-Y., Suh, J.-W., and Husson, R. N. (2008). Wag31, a
homologue of the cell division protein DivIVA, regulates growth, morphology and polar
cell wall synthesis in mycobacteria. *Microbiology* 154, 725–735.
doi:10.1099/mic.0.2007/014076-0.

Kapoor, N., Pawar, S., Sirakova, T. D., Deb, C., Warren, W. L., and Kolattukudy, P. E.
(2013). Human granuloma in vitro model, for TB dormancy and resuscitation. *PLoS One*
8, e53657. doi:10.1371/journal.pone.0053657.

Knapp, G. S., Lyubetskaya, A., Peterson, M. W., Gomes, A. L. C., Ma, Z., Galagan, J. E., et
al. (2015). Role of intragenic binding of cAMP responsive protein (CRP) in regulation
of the succinate dehydrogenase genes Rv0249c-Rv0247c in TB complex mycobacteria.
Nucleic Acids Res. 43, 5377–93. doi:10.1093/nar/gkv420.

Kong, H. G., Bae, J. Y., Lee, H. J., Joo, H. J., Jung, E. J., Chung, E., et al. (2014). Induction
of the viable but nonculturable state of *Ralstonia solanacearum* by low temperature in
the soil microcosm and its resuscitation by catalase. *PLoS One* 9, e109792.
doi:10.1371/journal.pone.0109792.

Kong, I.-S., Bates, T. C., Hülsmann, A., Hassan, H., Smith, B. E., and Oliver, J. D. (2004).

Role of catalase and oxyR in the viable but nonculturable state of *Vibrio vulnificus*.

FEMS Microbiol. Ecol. 50, 133–142. doi:10.1016/j.femsec.2004.06.004.

Liu, J., Li, L., Peters, B. M., Li, B., Chen, L., Deng, Y., et al. (2017). The viable but

nonculturable state induction and genomic analyses of *Lactobacillus casei* BM-

LC14617, a beer-spoilage bacterium. *Microbiologyopen* 6. doi:10.1002/mbo3.506.

Lynn, M., Wilson, A. R., and Solotorovsky, M. (1979). Role of bovine serum albumin in the

nutrition of *Mycobacterium tuberculosis. Appl. Environ. Microbiol.* 38, 806–810.

doi:10.1128/aem.38.5.806-810.1979.

Manuse, S., Fleurie, A., Zucchini, L., Lesterlin, C., and Grangeasse, C. (2016). Role of

eukaryotic-like serine/threonine kinases in bacterial cell division and morphogenesis.

FEMS Microbiol. Rev. 40, 41–56. doi:10.1093/femsre/fuv041.

Minato, Y., Gohl, D. M., Thiede, J. M., Chacón, J. M., Harcombe, W. R., Maruyama, F., et

al. (2019). Genomewide Assessment of *Mycobacterium tuberculosis* Conditionally

Essential Metabolic Pathways. *mSystems* 4. doi:10.1128/mSystems.00070-19.

Mizunoe, Y., Wai, S. N., Ishikawa, T., Takade, A., and Yoshida, S. (2000). Resuscitation of

viable but nonculturable cells of *Vibrio parahaemolyticus* induced at low temperature

under starvation. *FEMS Microbiol. Lett.* 186, 115–120. doi:10.1111/J.1574-

6968.2000.TB09091.X.

Morishige, Y., Koike, A., Tamura-Ueyama, A., and Amano, F. (2017). Induction of viable

but nonculturable *Salmonella* in exponentially grown cells by exposure to a low-

humidity environment and their resuscitation by catalase. *J. Food Prot.* 80, 288–294.

doi:10.4315/0362-028X.JFP-16-183.

Mukamolova, G. V., Turapov, O., Malkin, J., Woltmann, G., and Barer, M. R. (2010).

Resuscitation-promoting factors reveal an occult population of tubercle bacilli in

sputum. *Am. J. Respir. Crit. Care Med.* 181, 174–180. doi:10.1164/rccm.200905-

0661OC.

Mukamolova, G. V, Murzin, A. G., Salina, E. G., Demina, G. R., Kell, D. B., Kaprelyants, A.

S., et al. (2006). Muralytic activity of *Micrococcus luteus* Rpf and its relationship to

physiological activity in promoting bacterial growth and resuscitation. *Mol. Microbiol.*

59, 84–98. doi:10.1111/j.1365-2958.2005.04930.x.

Oliver, J. D. (2010). Recent findings on the viable but nonculturable state in pathogenic

bacteria. *FEMS Microbiol. Rev.* 34, 415–425. doi:10.1111/j.1574-6976.2009.00200.x.

Osada-Oka, M., Goda, N., Saiga, H., Yamamoto, M., Takeda, K., Ozeki, Y., et al. (2019).

Metabolic adaptation to glycolysis is a basic defense mechanism of macrophages for

Mycobacterium tuberculosis infection. *Int. Immunol.* 31, 781–793.

doi:10.1093/INTIMM/DXZ048.

Parikh, A., Verma, S. K., Khan, S., Prakash, B., and Nandicoori, V. K. (2009). PknB-

mediated phosphorylation of a novel substrate, N-acetylglucosamine-1-phosphate

uridyltransferase, modulates its acetyltransferase activity. *J. Mol. Biol.* 386, 451–64.

doi:10.1016/j.jmb.2008.12.031.

Pasquaroli, S., Zandri, G., Vignaroli, C., Vuotto, C., Donelli, G., and Biavasco, F. Antibiotic

pressure can induce the viable but non-culturable state in *Staphylococcus aureus*

growing in biofilms. doi:10.1093/jac/dkt086.

R Core Team (2021). R: A Language and Environment for Statistical Computing. Available

at: https://www.r-project.org.

Roche, M., Rondeau, P., Singh, N. R., Tarnus, E., and Bourdon, E. (2008). The antioxidant

properties of serum albumin. *FEBS Lett.* 582, 1783–1787.

doi:10.1016/J.FEBSLET.2008.04.057.

Rock, J. M., Hopkins, F. F., Chavez, A., Diallo, M., Chase, M. R., Gerrick, E. R., et al.

(2017). Programmable transcriptional repression in mycobacteria using an orthogonal

CRISPR interference platform. *Nat. Microbiol.* 2, 16274.

doi:10.1038/nmicrobiol.2016.274.

Rombouts, I., Lagrain, B., Scherf, K. A., Lambrecht, M. A., Koehler, P., and Delcour, J. A.

(2015). Formation and reshuffling of disulfide bonds in bovine serum albumin

demonstrated using tandem mass spectrometry with collision-induced and electron-

transfer dissociation. *Sci. Rep.* 5, 12210. doi:10.1038/srep12210.

Salina, E. G., Waddell, S. J., Hoffmann, N., Rosenkrands, I., Butcher, P. D., and Kaprelyants,

A. S. (2014). Potassium availability triggers *Mycobacterium tuberculosis* transition to,

and resuscitation from, non-culturable (dormant) states. *Open Biol.* 4.

doi:10.1098/rsob.140106.

Sassetti, C. M., Boyd, D. H., and Rubin, E. J. (2003). Genes required for mycobacterial

growth defined by high density mutagenesis. *Mol. Microbiol.* 48, 77–84.

doi:10.1046/j.1365-2958.2003.03425.x.

Schaefer, W. B. (1952). GROWTH REQUIREMENTS OF DYSGONIC AND EUGONIC

STRAINS OF *MYCOBACTERIUM TUBERCULOSIS VAR. BOVIS*. *J. Exp. Med.* 96,

207–219. doi:10.1084/JEM.96.3.207.

Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., et al.

(2012). Fiji: an open-source platform for biological-image analysis. *Nat. Methods 2012*

97 9, 676–682. doi:10.1038/nmeth.2019.

Senoh, M., Hamabata, T., and Takeda, Y. (2015). A factor converting viable but

nonculturable *Vibrio cholerae* to a culturable state in eukaryotic cells is a human

catalase. *Microbiologyopen* 4, 589–96. doi:10.1002/mbo3.264.

Shleeva, M., Goncharenko, A., Kudykina, Y., Young, D., Young, M., and Kaprelyants, A.

(2013). Cyclic amp-dependent resuscitation of dormant mycobacteria by exogenous free

fatty acids. *PLoS One* 8, e82914. doi:10.1371/journal.pone.0082914.

Shleeva, M. O., Kondratieva, T. K., Demina, G. R., Rubakova, E. I., Goncharenko, A. V.,

Apt, A. S., et al. (2017). OVerexpression Of adenylyl cyclase encoded by the

Mycobacterium tuberculosis Rv2212 gene confers improved fitness, accelerated

recovery from dormancy and enhanced virulence in mice. *Front. Cell. Infect. Microbiol.*

7, 370. doi:10.3389/fcimb.2017.00370.

Sureka, K., Hossain, T., Mukherjee, P., Chatterjee, P., Datta, P., Kundu, M., et al. (2010).

Novel role of phosphorylation-dependent interaction between FtsZ and FipA in

mycobacterial cell division. *PLoS One* 5, e8590. doi:10.1371/journal.pone.0008590.

Thakur, M., and Chakraborti, P. K. (2006). GTPase activity of mycobacterial FtsZ is

impaired due to its transphosphorylation by the eukaryotic-type Ser/Thr kinase, PknA. *J.*

Biol. Chem. 281, 40107–13. doi:10.1074/jbc.M607216200.

Thakur, M., and Chakraborti, P. K. (2008). Ability of PknA, a mycobacterial eukaryotic-type

serine/threonine kinase, to transphosphorylate MurD, a ligase involved in the process of

peptidoglycan biosynthesis. *Biochem. J.* 415, 27–33. doi:10.1042/BJ20080234.

Vilhena, C., Kaganovitch, E., Grünberger, A., Motz, M., Forné, I., Kohlheyer, D., et al.

(2019). Importance of Pyruvate Sensing and Transport for the Resuscitation of Viable

but Nonculturable *Escherichia coli* K-12. *J. Bacteriol.* 201. doi:10.1128/JB.00610-18.

Wagley, S., Morcrette, H., Kovacs-Simon, A., Yang, Z. R., Power, A., Tennant, R. K., et al.

(2021). Bacterial dormancy: A subpopulation of viable but non-culturable cells

demonstrates better fitness for revival. *PLoS Pathog.* 17, e1009194.

doi:10.1371/journal.ppat.1009194.

Wayne, L. G., and Hayes, L. G. (1996). An in vitro model for sequential study of shiftdown

of *Mycobacterium tuberculosis* through two stages of nonreplicating persistence. *Infect.*

Immun. 64, 2062–2069. doi:10.1128/iai.64.6.2062-2069.1996.

World Health Organization (2020). "TB disease burden," in *Global Tuberculosis Report*

2020, Chapter 4, 23.

Xu, H. S., Roberts, N., Singleton, F. L., Attwell, R. W., Grimes, D. J., and Colwell, R. R.

(1982). Survival and viability of nonculturable *Escherichia coli* and *Vibrio cholerae* in

the estuarine and marine environment. *Microb. Ecol.* 8, 313–23.

doi:10.1007/BF02010671.

Yeware, A., Gample, S., Agrawal, S., and Sarkar, D. (2019). Using diphenyleneiodonium to

induce a viable but non-culturable phenotype in *Mycobacterium tuberculosis* and its

metabolomics analysis. *PLoS One* 14, e0220628. doi:10.1371/journal.pone.0220628.

Figure legends

Figure 1. Growth curves of Mtb H37Rv cells in Dubos medium in the presence or absence of

BSA

(A) Number of culturable cells in Dubos medium in the presence (closed circle) or

absence (open circle) of 0.1% BSA at the indicated days of incubation. The CFU/mL

values were determined by plating cells onto 7H10 plates in duplicate. Data represent

the mean ± SD from three independent experiments.

(B) Representative images of Mtb growth captured at the end of incubation.

Figure 2. Viability analyses of DPI-treated Mtb

(A) Number of culturable cells after DPI treatment. The CFU/mL values were determined

by plating cells onto 7H10 plates in duplicate. Asterisk indicates a statistically

significant difference (*$p < 0.05$, Welch's *t*-test). Data represent the mean ± SD from

three independent experiments.

(B) Percentages of esterase active cells after DPI treatment. Around 850 cells were

directly counted under 100× objective lens. Data represent the mean ± SD from three

independent experiments.

(C) Merged fluorescence micrographs of Mtb H37Rv cells stained with

carboxyfluorescein diacetate (CFDA) and ethidium bromide (EtBr). Cells stained with

CFDA (green) are esterase-positive; those stained with EtBr (red) are membrane-

damaged. Images are representative images from three individual experiments. Scale

bar: 2μm

(D) Percentage of acid-fast-positive cells (open column) or acid-fast-negative but lipid-

accumulation-positive cells (gray column). Approximately 1,100 cells were directly

counted under 100× objective lens. Data represent the mean ± SD from four

independent experiments. Asterisk indicates significant difference (***$p < 0.005$,

Welch's *t*-test).

(E) Merged fluorescence micrographs of Mtb H37Rv cells stained with auramine-O and

Nile Red. Cells stained with auramine-O (green) are acid-fast-positive; those stained

with Nile Red (red) are acid-fast-negative but lipid accumulation-positive. Images are

representative images from three individual experiments. Scale bar: 2μm

(F) Magnified views of Airyscan confocal super-resolution microscopy observation of

three different Mtb cells obtained from DPI-treated culture. Only acid-fast-positive

cells (left), both acid-fast-positive and lipid accumulation-positive cells (center,

counted as acid-fast-positive) and acid-fast-negative cells with only lipid

accumulation-positive cells (right). Dual-positive cells probably represent the

transition to a non-culturable state. Indicated foci with arrowheads represent a neutral

lipid body. Scale bar: 1 μm

Figure 3. Resuscitation of DPI-treated Mtb H37Rv cells by FBS and OADC supplementation

(A) Number of culturable cells at the indicated days of incubation, without

supplementation or those cells supplemented with 2% FBS or 10% OADC in the

presence (closed circle) or absence (open circle) of 0.1% BSA. The CFU/mL values

were determined by plating cells onto 7H10 plates in duplicate. Data represent the

mean ± SD from three independent experiments. U.D. means under the limit of

detection (20 CFU/mL).

(B) Representative images of regrowth under the indicated conditions captured at the end

of incubation. None, FBS, and OADC represent no supplementation, 2% FBS and

10% OADC supplementation, respectively.

Figure 4. Effect of human serum albumin, ovalbumin and antioxidants on the resuscitation of

DPI-treated Mtb.

(A) Number of culturable cells at the end of incubation, supplemented with albumins or

740   antioxidative agents. The CFU/mL values were determined by plating cells onto 7H10

741   plates in duplicate. Data represent the mean ± SD from three independent

742   experiments. Asterisk indicates a statistically significant difference between Day 0

743   and Day 20 by Student's *t*-test. (*p<0.05)

(B) Representative images of regrowth captured at the end of incubation.

Abbreviations:

BSA, bovine serum albumin; HSA, human serum albumin; OVA, ovalbumin; NAC, N-

acetyl-L-cysteine; MAN, D-mannitol.

The final concentrations of each reagent were as follows:

BSA, HSA and OVA: 0.1% (w/v), NAC: 500 µM, MAN: 50 mM.

Figure 5. The effect of SQ22536 or H89 on the resuscitation of DPI-treated Mtb

(A) Number of culturable cells at the end of incubation, supplemented with 0.1% BSA

752   and SQ22536 or H89. Inhibitor was added at the indicated concentration.

(B) Representative images of regrowth captured at the end of incubation.

The CFU/mL values were determined by plating cells onto 7H10 plates in duplicate. Data

represent the mean ± SD from three independent experiments.

Figure 6. Schematic representation of the mechanism of BSA-induced resuscitation of Mtb.

Although the detailed mechanism of interaction between BSA and PknA is not yet clear, BSA

may— in coordination with PknB—activate PknA and positively regulate target proteins that

are involved in the localized synthesis of peptidoglycan, cell division control, and DNA

condensation and partitioning through their phosphorylation, resulting in the resuscitation of

VBNC Mtb cells.

Figures

Figure 1.

773   Figure 2.

778   Figure 3.

780   Figure 4.

784   Figure 5.

786   Figure 6.

787