An evolutionary ecological mechanism of CyanoHABs: unavoidable consequences when low nutrient requirements meet eutrophication

Wenduo Cheng[†b], Somin Hwang[†b], Qisen Guo[†b], Haibei Zhang[†b], Leyuan Qian[b], Weile Liu[b], Yang Yu[b], Yi Tao[*a], and Huansheng Cao[*b]

[a]Guangdong Provincial Engineering Research Center for Urban Water Recycling and Environmental Safety, Tsinghua Shenzhen International Graduate School, Tsinghua University, Shenzhen, 518055, China

[b]Division of Natural and Applied Sciences, Duke Kunshan University, 8 Duke Ave, Kunshan, 215316, China

[†]Equal Contribution

Correspondence: Huansheng Cao (hc284@duke.edu) and Yi Tao (tao.yi@sz.tsinghua.edu.cn)

Abstract

The mechanism of cyanobacterial harmful algal blooms (CyanoHABs) is complicated and confusing. One major reason is they are studied primarily from an ecological perspective and on bloom-forming species only. This narrow angle loses a broader evolutionary and ecological context in which CyanoHABs occur and fails to provide information on relevant components to achieve a wholistic understanding.

To derive a comprehensive mechanism of CyanoHABs, we examine CyanoHABs through the overlooked evolutionary and ecological lenses: evolutionary radiation, ecological comparison with co-living algae, and recently identified genomic functional repertoire between blooming and non-blooming species. We found four key factors: (1) elaborate diverse functional repertoire and low nutrient requirement in cyanobacteria molded by early adaptive evolution, (2) cyanobacteria having lower nutrient requirements than co-living algae, (3) there is no directed evolution in biological functions toward water eutrophication in cyanobacteria, (4) the CyanoHAB-associated functional repertoire are more abundant and complete in blooming than non-blooming species. These factors lead us to postulate a preliminary mechanism of CyanoHABs as a synergistic quad: superior functional repertoire, established with long adaptive radiation under nutrient-deficient conditions and not evolved toward eutrophic conditions, enables cyanobacteria to efficiently utilize elevated nutrients under current eutrophic levels for excess growth and CyanoHABs thereof, due to their lower nutrient requirements than co-living algae. This working comprehensive mechanism without doubt needs further empirical testing, which can be undertaken with more comparative studies of multiple species using integrated systems biology approaches.

Key words:

I. Introduction

CyanoHABs are one of the most profound environmental hazards in modern human history in terms of their global geographical scale (Paerl et al., 2020), longstanding duration (over a century) (Francis, 1878; Lathrop & Carpenter, 1992), and tremendous economic loss (Steffensen, 2008). The mere fact that they are still intensifying and expanding under the global climate change (Chapra et al., 2017; Gobler, 2020; Paerl et al., 2020) attests CyanoHABs are still not fully understood such that their ecology remains 'complicated and confusing' (Wilhelm et al., 2020).

Are CyanoHABs an evolutionary consequence or ecological outcome in the era of water eutrophication? Nearly all research on CyanoHABs until now has focused on ecological factors, including nutrients, temperature, and climate change (Adrian et al., 2009; Beaulieu et al., 2013b; O'Neil et al., 2012). A major recent advance is the identification of genomic functional repertoire enabling the elevated nutrients for cyanobacterial growth and bloom formation (Cao et al., 2020a; Du et al., 2020). Now it becomes clear that CyanoHABs are immediate outcomes of synergistic interactions between superior cyanobacterial functions and eutrophic conditions. Is this all there is to it? In any eutrophic water, why it is primarily cyanobacteria, not other algae, that form blooms? Apparently, there ought to be more factors involved in causing this differential dominance in phytoplankton communities. One fact is the current eutrophic levels are sufficient to support CyanoHABs but not blooms of other algae.

To address this question of differential dominance, one can take a step back and use a wholistic approach by considering the fundamental principles of biology. For example, many seemingly complex genomic questions are often reconciled in the light of evolution—"Nothing in biology makes sense except in the light of evolution" (Dobzhansky, 1973)—as shown previously (Avise & Ayala, 2007; Lynch, 2007; Meyer, 2008). Cyanobacteria, the most ancient autotrophs (Boden et al., 2021) with the broadest geographical distribution on modern Earth (Schopf, 2002), have come to this success through various trajectories of adaptive evolution. By examining these

trajectories of geological timescales in the context of each niche, one can deduce the

key biological functions for their success. More importantly, comparing the early

adaptive radiation and present-day ecological dominance of cyanobacteria in eutrophic

conditions enables identification of different factors contributing to either type of

success, respectively.

Following this line of reasoning, we first reexamined the biological abilities of early

cyanobacteria to survive and thrive in habitats which are physically harsh and

nutritiously deficient during evolution on geological timescales. Next, we revisited the

water eutrophication processes to establish elevated nutrients as the necessary

condition for CyanoHABs and highlighted the overlooked fact that cyanobacteria have

lower nutrient requirements than green algae, which have never bloomed in the current

regime of eutrophication. Third, we summarized the biological functions driving bloom

formation in eutrophic waters and showed the genes encoding these functions are not

under positive (directed) selection, but purifying selection instead. This finding suggests

their biological capabilities are well established before the era of water eutrophication.

All these considerations above led us to postulate an evolutionary ecological

mechanism of CyanoHABs. This new synthesis requires expanding biological

dimensions in future CyanoHAB research.

51   II. Cyanobacteria: evolutionary champions

Originating in an anoxic biosphere of early Earth, cyanobacteria are likely to evolve in

scarcity of macronutrients such as phosphorus and nitrogen (Bjerrum & Canfield, 2002;

Falkowski, 1997; Kipp & Stüeken, 2017; Reinhard et al., 2017) and trace metals (Anbar

& Knoll, 2002; Saito et al., 2003). This humble beginning of life leads to primitive

metabolic networks (Goldford & Segrè, 2018), which suggests that cyanobacteria could

only obtain minimal energy and matter in both variety and amount to support cell

survival and division (Schopf, 2002). Consequently, their nutrient requirements for life

are likely to be very low. Besides nutrient deficiency, early cyanobacteria also faced

various types of stresses which are 'unforeseeable' for them. One type is the rising

oxygen level they caused themselves through oxygenic photosynthesis, for which they

developed various counterattack strategies observed in modern cyanobacteria (Moore

et al., 2017; Raymond & Segrè, 2006). Ultraviolet light is another early stress which has

persisted until now, and some cyanobacteria evolved against it by producing

sunscreens (Garcia-Pichel et al., 2019). These global stresses were also intertwined

with landscape changes through geophysical processes (e.g., volcanism and climatic

shifts). Here, we demonstrate the remarkable adaptive ability of cyanobacteria by

showcasing their evolution of biological capability and morphology under the rising

oxygen levels.

Since origin, particularly after the Great Oxidation Event (GOE) (Figure 1), ancestral

cyanobacteria quickly acquired versatile antioxidant capabilities against reactive oxygen

species (ROSs), derived from O_2 produced as a result of oxygenic photosynthesis.

Different classes of superoxide dismutases (SODs) and other types of antioxidant

enzymes are evolved to deal with ROSs throughout their entire history (Boden et al.,

2021). Based on the availability of metal cofactors (another case of adaptation), three

types of SODs—NiSOD, CuZnSOD, and MnSOD/FeSOD—have been found in four

types of aquatic environments (Figure 1A). Besides direct enzymatic removal of ROSs,

cyanobacteria have also evolved other strategies, e.g., in photosynthetic reaction center

(RC) which split into two types as a response to the rising oxygen (Orf et al., 2018).

Another adaptation lies in the redundancy (heavy investment) of key functions. For

example, in photosystem I, cyanobacteria have two electron transporters that evolved

separately: copper-dependent plastocyanin (PC) and iron-dependent cytochrome c_6

(Cytc_6). Distinct in the primary sequence and tertiary structure of protein/amino acid,

their expression is controlled by a BlaI/CopY-family transcription factor (PetR) and a

BlaR-membrane protease (PetP), depending on the availability of the metal cofactors

(García-Cañas et al., 2021).

With a continuous rise in oxygen level, morphological diversification was called upon to

provide an extra dimension for functional expansion. It has been confirmed that the

evolution of multicellular morphotypes and the rate of morphological diversification

coincide with the GOE onset (Schirrmeister et al., 2013). Heterocyst became a

specialized cell form for nitrogen fixation, in which the oxygen level is reduced to pre-
GOE levels (Figure 1B). Nitrogen fixation is viewed as a leap forward in promoting
marine primary production and contributed to increased O_2 levels, coinciding with the
rise of animals (Lyons et al., 2014). Similarly, cyanobacteria also obtained a specific
photosynthetic structure for carbon fixation, carboxysome, in which CO_2 is increased
from 10-15μM to 40mM outside and greatly improved the efficiency of enzymatic fixation
of carbon dioxide (Mangan & Brenner, 2014). After the evolution of geological
timescales (Figure 1C and 1D), modern cyanobacteria are among the most diverse
prokaryotic phyla, with morphotypes ranging from unicellular to multicellular filamentous
forms, including those able to terminally (i.e., irreversibly) differentiate in form and
function (Hammerschmidt et al., 2020; Schirrmeister et al., 2013).
104
Billions of years of adaptive evolution have allowed structural elaboration and functional
diversification in coping with nutrient deficiency, environmental extremes and constraints
(Paerl, 2014). This evolutionary divergence has enabled cyanobacteria to occupy all
geographic habitats in modern Earth, including terrestrial and aquatic ecosystems,
ranging from deserts to tropical rain forests, soils and limestones, and from open
oceans and brackish waters to freshwaters and hydrothermal vents (Figure 2). This
broad array of biological functions underlying the adaptive radiation in cyanobacteria is
recorded in their genomes (Chen et al., 2021). A large proportion of the genes in their
genomes are associated with adaption to the specific habitats they occupy, but not to
any other habitats. Therefore, the genomes produce a pangenome with most genes
being unique and only a small set of 323 genes being common among cyanobacteria
(Shi & Falkowski, 2008). These core genes are mainly involved in housekeeping
functions, as opposed to interacting with environment, e.g., ribosomal proteins,
photosynthetic apparatus, ATP synthesis, chlorophyll biosynthesis, and the Calvin cycle.
This small stable core and large variable shell of genomes suggest most of the non-
overlapping functions are specialized in adaptation to distinct habitats.
121

Figure 1. The evolutionary trajectory of cyanobacteria in the presence of oxygen. (A) The origin and

distribution of SODs in different habitats. The timing of origin and habitat distribution of SOD genes

among taxa is created based on Boden et al (Boden et al., 2021). Starting points of horizontal bars

represent time of origin and color SOD types: CuZnSOD (blue), NiSOD (green), and Fe- and Mn-utilizing

SODs (yellow). (B) The oxygen level in atmosphere and nitrogen fixation in heterocysts in cyanobacteria

in the presence of elevated oxygen levels. The level of oxygen is based on (Tomitani et al., 2006). (C) and

(D) The temporal morphological diversification cyanobacteria is based on Schirrmeistera et al

(Schirrmeister et al., 2013).

1. Soil
- Absorbs light crossing sands and soils
- Accumulates a high concentration of sugar to face soil desiccation

2. Desert
- Extracellular Polymeric Substances (EPS) in cyanobacteria resists the major diurnal fluctuations in temperatures and retain water in a long time
- Microbial tissue and expansion of EPS pioneer the development of microphytic soil crust

3. Forest
- Protects nitrogenase from oxygen inactivation as a consequence of oxygenic photosynthesis
- Motile filaments (hormogonia) serve as infective agents in many forms of symbiosis
- Responds to plant signals, resulting in reduced growth rate, reduced carbon dioxide and enhanced nitrogen by morphological and physiological changes

4. Salty Area
- Utilizes sulfide: apply the anoxygenic photosynthetic electron transport chain
- Applies organic osmolytes such as disaccharides sucrose and trehalose (Osmolarity Balance)

5. Oil-spilled Ocean
- Degrades aromatic and aliphatic oil constituents
- N-fixing cyanobacteria's growth is enhanced by hydrocarbon

6. Ocean
- Uses gas vesicles to adjust buoyancy and positions
- Conducts positive phototropism and fast growth in endoliths of live coral
- Utilize swimming in chemotactic response to N compounds

7. Limestone
- Deposites Calcium carbonate by trapping particles or contributing to communities where calcite and aragonite
- Etches limestones and grow within substrates

8. Geothermal Habitat
- Disperses barriers to migratory gene flow
- Occupies different niches depending on different temperatures, pH, salinity of the hotsprings

9. Polar Region
- Benthic cyanobacterial mats: covers by a thin layer of concentrated brine (salt) with salinity of six times more than sea water at the liquid water temperature of -12℃
- Minimizes self-shading with small diameter of mats that enhances light-capturing abilities under dim PAP conditions

10. Ultraviolet Light
- Chooses habitats such as beneath rock surfaces or microbial mat matrices; position themselves vertically in water columns
- Actively conducts repair for UV tolerance: excision repair and photoreactivation activity by the DNA repair mechanism
- Produces UV-absorbing sheath pigments to protect from UV light and UV-screening compounds such as MMA

Figure 2. Cyanobacterial adaptation strategies in nine environments, namely soil, desert, forest, salty area, oil-spilled ocean, open ocean, limestone, geothermal water, and polar region. Cyanobacteria also cope with the damage caused by ultraviolet light.

Figure 3. The phylogeny of cyanobacteria based on 16S ribosomal genes. Blue labels: bloom-
forming cyanobacteria; and red labels: non-bloom-forming cyanobacteria. The shapes of
different colors represent they types of morphology.

III. CyanoHABs: an unavoidable consequence when low nutrient-requiring cyanobacteria meet eutrophic waters

Anthropogenic eutrophication, elevated loading of nutrients into waters from human activities on a global scale, is an 'unforeseeable' surprise to cyanobacteria which have been evolving, but not forming blooms, in physically harsh and nutrient-deficient waters. Apparently, eutrophic waters present themselves as a new but favorable condition (a surprise), in which cyanobacteria dominate over co-living algae in phytoplankton communities and form CyanoHABs. To achieve a wholistic mechanism of CyanoHABs, three types of questions should be addressed: First, is there direct causality between eutrophication and CyanoHABs to establish eutrophication as the necessary condition for CyanoHABs? Second, why is it cyanobacteria, not other co-living algae (e.g., green algae), that form blooms? Do other algal species have higher nutrient requirements than cyanobacteria, which cannot be met by current eutrophic levels? This is another necessary condition for CyanoHABs. Last, what are the superior biological functions of blooming cyanobacteria, the sufficient condition of CyanoHABs, that take advantage of eutrophic conditions compared to the non-blooming cyanobacteria?

3.1 A first necessary factor: water eutrophication from oligotrophic state

Despite natural eutrophication, few CyanoHABs in natural waters were reported before the era of anthropogenic eutrophication. Water eutrophication is not only a result of nutrient loading but also a process during which favorable factors such as hydrology are brought about by anthropogenic landscape changes. The driving role of eutrophication in CyanoHABs is well established (Heisler et al., 2008). Its causality is manifested in two ways: fast bloom formation in response to nutrient elevation and bloom decline with de-eutrophication. Immediate bloom formation has been repeatedly observed in laboratory simulations (Lürling et al., 2018) and field studies (Molot et al., 2021; Wilkinson et al., 2018). Here we provide two examples to show that the transition from oligotrophic to eutrophic state is swift and CyanoHABs follow in a short period of time.

One example is the 'most studied lake in the world', Lake Mendota (Brock, 1985; Kitchell, 1992). Started as a clear oligotrophic lake prior to Euro-American colonization

in the 1830s in Madison, Wisconsin (Riera et al., 2001), Lake Mendota had CyanoHABs

in 1882 and later 1890s (Lathrop & Carpenter, 1992) due to eutrophication from

combined landscape changes of agricultural expansion in area and crops (corn in place

of wheat and oats), wetlands drainage, lake-level rise and slow water flow, and soil

erosion due to the damming lake outlet. Arguably the most convincing evidence for the

cause of eutrophication for CyanoHABs is from Lake Erie, where CyanoHABs not only

form following eutrophication but also decline with de-eutrophication, as summarized by

Steffen et al (2014). Briefly, an increase in loading of phosphorus and other

macronutrients to Lake Erie led to CyanoHABs (so-called autumnal maximum)

dominated by filamentous cyanobacteria (*Anabaena* spp., *Aphanizomenon* spp.,

Lyngbya spp.), which quickly succeeded to *Microcystis* sp. With the phosphorus

identified as the key nutrient and subsequent implementation of the Great Lakes Water

Quality Agreement in 1972, the total phosphorus loading goal was reached in 1981 and

continued to mid-1990s. Accordingly, a consistent decline in phytoplankton was

observed. Unfortunately, a return of *Microcystis aeruginosa* blooms was observed in

1995 (Conroy et al., 2005) due to external loading of soluble reactive phosphorus

(Joosse & Baker, 2011) and CyanoHABs have persisted until now.

Besides freshwater ecosystems, eutrophication-led CyanoHABs were also found in

brackish waters, e.g., Baltic Sea (Bianchi et al., 2000; Gustafsson et al., 2012) and

coastal waters and seas (Anderson et al., 2008; Malone & Newton, 2020; Rahav & Bar-

Zeev, 2017; Sabeur et al., 2016). The fact that CyanoHABs form on the entire spectrum

of salinity indicates that salinity is not an effective deterring factor, which is a

confirmation that cyanobacteria have long adapted to each of these habitats under

oligotrophic conditions (Figure 2).

3.2 A second necessary factor: low nutrient-requirements in cyanobacteria

As observed in the studies of causality between eutrophication and CyanoHABs, it is

cyanobacteria, not green algae, that form blooms in response to nutrient addition.

Considering the adaptive radiation of cyanobacteria, one can reason that cyanobacteria

have lower requirements for nutrients than green algae. As nutrients increase in

originally oligotrophic waters, they meet the (lower) optimal needs of cyanobacteria first

and further increases will satisfy the (higher) optimal demand of green algae to support

their mass growth for bloom formation.

To test this hypothesis, four comparisons were performed. First, we compared the

recipes of synthetic media for cyanobacteria and green algae and found those for green

algae have much higher ionic concentrations than those for cyanobacteria (Figure 4).

Second, we summarized the results which directly compared the growth of

cyanobacteria and green algae in the field, laboratory, and cosmos by supplying

external nutrients (Table 1). Five groups of studies support our hypothesis: low nutrients

(TP, TN, DIN, urea-N, Ca, etc.) support dominance of cyanobacteria and high nutrients

support dominance of green algae and even brown algae, as summarized in Table 1.

Third, cyanobacteria have fewer genes, about 50% of those encoded in green algae.

More importantly, based on the genome-scale metabolic networks (GSMs), the size of

the GSMs of cyanobacteria is about half of those of green algae, having on average 780

metabolites and 820 reactions, which are about half of those in green algal GSMs

(Table 2). Last, cyanobacteria tend to have a higher stress tolerance to N limitation than

green algae. For example, *M. aeruginosa* can increase their photosynthetic

performance which helps maintain their population size during seasonal N limitation,

while photosynthetic performance is decreased in green alga *Porphyridium Cruentum*

under the same N limitation (Jin *et al.*, 2020). All these evidence clearly shows that

cyanobacteria indeed have low requirements than green algae and thus bloom first

when nutrient levels increase.

Figure 4. Lower nutrient requirements by cyanobacteria than green algae in synthetic
culture media. Among seven selected media, the recipes of mAC, MJ, VTAC, and CHU-
11 (left panel) are used for culturing green algae and the recipes of M11, BG11 and CT
(right panel) are used for culturing cyanobacteria.

Table 1. A list of case studies which show cyanobacteria have lower nutrient requirements than green algae. B: cyanobacteria; G: green algae. DIN: dissolved inorganic N; SRP: soluble reactive phosphorus; TN: total nitrogen; and TP: total phosphorus.

	Study Type	Study Site(s)	Indicator(s)	Concentration(s)	Dominance	Dominant Species	References
Group 1	Mesocosm	Wascana Lake, Canada	Urea-N	1-8 mg N L^{-1}	B	Colonial cyanobacteria	(Bogard *et al.*, 2020)
				>8 mg N L^{-1}	G	Chlorophytes	
	Laboratory	N/A	NH$_4^+$ and NO$_3^-$	32μM (NO$_3$) after 48h; 32μM NH$_4^+$ after 48h	B	*Synechococcus 6301*	(Hyenstrand *et al.*, 2000)
				32μM NO$_3$ after 25h; 32μM NH$_4$ after 18h	G	*Scenedesmus sp.*	
Group 2	*In situ*	210 Danish lakes	TP	0.25-0.8 mg P L^{-1}	B	Non-heterocystous cyanobacteria	(Jensen *et al.*, 1994)
				>1.0 mg P L^{-1}	G	Chlorophytes	
Group 3	*In situ*	Vela Lake, Central Portugal	DIN and SRP	0.66-2.42 mg L (DIN); 0-0.24 mg L (SRP)	B	*Aphanizomenon flos-aquae, Ch roococcus limneticus, M. aeruginosa* and *Pseudanabaena sp.*	(Figueiredo *et al.*, 2006)
				0.7-3.3 mg L (DIN); 0.19-0.93 mg L (SRP)	G	*Coelastrum reticulatum* var. *reticulatum, Kirchneriella lunaris, Monoraphidium contortum, Scenedesmus acuminatus* var. *acuminatus* and *Pediastrum boryanum* var. *boryanum*	
	In situ	Murchison Bay, Lake Victoria	TN and TP	1127 ug L (TN); 91 ug L(TP)	B	*Anabaena sp., M. wesenbergii, Gomphosphaeria aponina* and *Microcystis aeruginosa, Planktolyngbya circumcreta*	(Haande *et al.*, 2011)
	Laboratory and *in situ*	Lake Tai, China	N and P	3 mg/L (TN) and 0.2 mg/L (TP)	B	*M. aeruginosa*	(Zhu, Wan, & Zhao, 2010)
				10, 15, or 20 mg/L (TN), and 1 mg/L (TP)	G	*Scenedesmus quadricauda*	
Group 4	Mesocosm	Archipelago Sea	N:P Ratio	1N:1P 1.7 μg L^{-1}: 1.7 μg L^{-1}	B	*Aphanizomenon* sp., *Nodularia spumigena, Anabaena* spp., *Synechococcus* spp.	(Vuorio *et al.*, 2005)
				7N:1P 12 μg L^{-1}:1.7 μg L^{-1}	G	*Dictyosphaerium subsolitarium, Monoraphidium contortum*, and *Monoraphidium contortum, Oocystis* spp..	
	Mesocosm	Wascana Lake, Canada	N:P ratio	15:1 to >24:1	G	*Micractinium, Oocystis*	(Finlay *et al.*, 2010)
Group 5	*In situ* and laboratory	Taipei Feitsui Reservoir	Ca	7-9 mg L^{-1}	B	*Microcystis* spp. and *Aphanocapsa delicatissima.*	(Wu & Kow, 2010)
				9,11,13,15,18 mg L^{-1}	G	*Eutetramorus, Coelastrum, Coenocystis*, and *Dictyosphaerium*	

234   Table 2. Comparison of genome-scale metabolic networks of cyanobacteria and green

235   algae.

	Organism	Genes	Model Genes	Metabolites	Reactions	Reference
Cyanobacteria	Arthrospira platensis	6631	620	673	746	(Yoshikawa et al., 2015)
	Cyanothece sp. ATCC 51142	4912	806	651	667	(Vu et al., 2012)
	Spirulina platensis C1	6153	692	837	875	(Klanchui et al., 2012)
	Synechococcus elongatus	2723	785	768	850	(Broddrick et al., 2016)
	Synechococcus sp PCC 7002	3179	821	777	792	(Qian et al., 2017)
	Synechocystis sp. PCC6803	3221	678	795	863	(Nogales et al., 2012)
	Synechocystis sp. PCC6803	3221	669	790	882	(Montagud et al., 2010)
Green algae	Chlamydomonas reinhardtii	16709	866	1862	1725	(Gomes de Oliveira Dal'Molin et al., 2011)
	Chlamydomonas reinhardtii	16709	1080	1068	2190	(Chang et al., 2011)
	Chlorella variabilis	9879	526	1236	1455	(Juneja et al., 2016)
	Chlorella vulgaris	7100	843	1770	2286	(Zuñiga et al., 2016)

3.3 A sufficient factor: superior functional repertoire without prior local adaptation

Given their global presence and persistence for over a century (Francis, 1878; Lathrop

& Carpenter, 1992), one may wonder whether there is prior adaptive evolution in

cyanobacteria to form bloom in eutrophic waters, or CyanoHABs are simply ecological

consequences of synergistic interaction between pre-equipped (before eutrophication

era) superior biological functions and elevated nutrients. We think the latter is more

likely than the former. First, short-time whole-lake experiments in fertilization or nutrient

reduction (Molot et al., 2021; Schindler et al., 2016; Wilkinson et al., 2018) have

indisputably shown the causality of eutrophication for CyanoHABs on an ecological time

scale, allowing no time for evolution. Second, anthropogenic eutrophication is

unforeseeable to cyanobacteria, and therefore they could not proactively prepare

themselves through directed evolution.

Third, we confirmed there is no directed evolution toward CynaoHABs, by calculating

the metric, dN/dS (ratio of non-synonymous to synonymous mutations), for all

homologous genes in 20 blooming species of *M. aeruginosa* genomes from different

countries. Indeed, almost all homologous genes display dN/dS < 0.8, with a median of

0.32 (Figure 5A). When grouped into the functional pathways curated previously (Cao et

al., 2020b), the homologous genes associated with CyanoHABs displayed the levels of

purifying selection pressure as those in core metabolism and housekeeping processes

(e.g., translation and transcription regulation) (Figure 5B). This strong purifying selection

suggests these gene functions not only are important functions such that cells cannot

afford them to be comprised by mutations. The levels of function are sufficient in the

metabolic networks for blooming cyanobacteria to thrive in oligotrophic conditions and

dominate in eutrophic conditions.

Last, the timescales observed between eutrophication and first blooms permit no

evolution. We can calculate the positive mutation generated by mutations in one annual

cycle. The spontaneous mutation rate (μ) in cyanobacteria is estimated at 10^{-7} per

genome per generation (García-Villada et al., 2004; López-Rodas et al., 2007; Osburne

et al., 2011; Segawa et al., 2018). For a lake of a million m^3, with cyanobacterial 20

cells/L growing to 10^8 cells/L after 25 doublings (t) at a growth rate 0.05-1.1 d^{-1} in an

annual cycle bottlenecked in fall and winter, this gives an effective population size (N_e)

of 10^8 cells, based on the equation 15 in (Wilson Alan et al., 2006). So, the number of

beneficial mutations to be fixed can be approximated by, assuming a beneficial rate (s)

at 0.01 and a rate of fixation (2s) at 0.05, $N_e \mu t 2s$ ($10^8 \times 10^{-7} \times 25 \times 2 \times 0.05 = 0.12$).

Therefore, each year there is less than one beneficial mutant will occur, which makes

prior adaptive evolution highly unlikely.

Figure 5. dN/dS for all homologous genes in 20 genomes of *M. aeruginosa*. (A) An overall

distribution of dN/dS for all homologous genes; and (B) the dN/dS for genes in different groups

of functions. dN/dS values were calculated by CODEML from the PAML 4.9j package with

default settings (Yang, 2007). AAPEP: uptake of amino acids and peptides; CCM: CO_2-

concentrating mechanism; DrugR: antibiotics resistance; MAA: UV radiation; MetalR: heavy

metal resistance; Nfix: nitrogen fixation; Nitrogen: nitrogen utilization; Osmos: osmos

homeostasis; OSR: oxidative stress resistance; PBS: phycobilisome; Phosphorus:

inorganic/organic utilization; PS-I/PS-II: photosystem I and photosystem II; Sugar: sugar

assimilation; TEVit: assimilation of trace metals and vitamins; Toxins: cyanotoxins; UFA:

unsaturated fatty acids; Vesicle: gas vesicles. Purple bars are pathways strongly associated

with CyanoHABs; green bars are pathways not specifically related to CyanoHABs; the last

seven blue bars represent essential biological functions (Rubin *et al.*, 2015).

IV. Biological functions associated with CyanoHABs

The relevant biological functions assimilating and converting them into biomass have been recently identified by comparative genomics between blooming and non-blooming species (Cao et al., 2020b). They are also made available as a database, CyanoPATH (Du et al., 2020) (Figure 6). These pathways have one-on-one correspondence with nutrients or stressors, which also show varying degrees of hierarchical structure. Below we detail the major pathways associated with CyanoHABs.

Figure 6. The core and query pathways in *Aphanizomenon flos-aquae* NIES-81. The pathways labeled with dashed borders are not complete (due to the absence of required components) in this strain but may be complete in others.

4.1 Gas vesicles

Gas vesicles are hollow protein structures made of gene products of *gvpA* and *gvpC* of the *gvp* gene cluster, consisting of 14 genes in several operons (Walsby, 1972, 1994). Gas vesicles are well-known for their roles in buoyancy regulation, allowing cyanobacteria to position themselves in the water column for optimal resource utilization (Carey et al., 2012; Walsby et al., 1997). This pathway is more enriched in blooming than non-blooming species and correlated with the blooming incidence of cyanobacteria recorded in the literature (Cao et al., 2020b): nearly all blooming species have at least one copy of the *gvpA* and *gvpC* genes, and the genes *gvpFGJKNVWX*, while all but one non-blooming species do not have any of these genes (Figure 7A). This pathway operates independently of other pathways, without a direct association with central metabolism.

4.2 Functions for macronutrient utilization

Phosphorus and nitrogen are recognized as the top two macronutrients for CyanoHABs by empirical studies (Heisler et al., 2008; Molot et al., 2021; Schindler et al., 2016) and statistical analyses (Beaulieu et al., 2013a). Indeed, the reconstructed pathways show that bloom-forming cyanobacteria invest heavily in these two functions, as they are limiting in the history of Earth (Kipp & Stüeken, 2017; Lyons et al., 2014; Reinhard et al., 2017). For phosphorus, both organic and inorganic phosphorus can be utilized. Inorganic phosphate can be internalized via ABC transporter *pstSCAB* or low facilitator affinity transporter *pitA*/*pitB* (Harris et al., 2001). Strikingly, nearly all cyanobacteria have one or two *pstSCAB* operons, and the globally dominant species *M. aeruginosa* has three. Organic phosphonate is imported via an ABC transporter PhnCDE and converted into phosphate by aminotransferase PhnW. The phosphate obtained has two destinations: incorporated into ATP as an energy molecule or converted to polyphosphate by polyphosphate kinase (PPK). Additionally, phosphate can be internally recycled, released from phosphate-containing compounds by inorganic pyrophosphatase and alkaline phosphatase. Interestingly, despite its verified driving role in CyanoHABs, the phosphate utilization pathway is not significantly more enriched (abundant or complex) in blooming than non-blooming species (Cao et al., 2020b). This

underscores two points. First, phosphorus has been a global limiting factor historically

(Elser et al., 2007; Maberly et al., 2020; Schindler, 1977) such that all cyanobacteria

have at least one and often two ABC transporters; phosphate appears to be only a

necessary but not sufficient driver for CyanoHABs, this is probably why total nitrogen

rather than total phosphorus is a better CyanoHAB predictor (Beaulieu et al., 2013a).

Second, this suggests the biological machinery for CyanHABs is complicated and more

likely to operate in a systems manner (Kirschner, 2005).

Total nitrogen and water temperature are recognized as the best predictors of

cyanobacterial biomass (Beaulieu et al., 2013a). Indeed, as the nitrogen utilization

pathway reconstructed with transporters and enzymes shows, cyanobacteria can use

four forms of nitrogen: nitrogen gas, inorganic nitrate/nitrite, ammonium, and organic

urea (Figure 7C). Key transporters for nitrate/nitrite (*nrtABCD*), ammonium (amt), urea

(*urtABCDE*) and key nitrate/nitrite reductive enzymes *narB*, *nirB*, and urease

(*ureABCDE*) are more enriched in blooming than non-blooming cyanobacteria (Figure

7D). Interestingly, the genes involved in incorporating nitrogen into amino acids (e.g.,

gdhA or gltB/gltD) are not significantly more enriched in blooming species; this suggests

that superior nitrogen utilizing ability mainly lies in the uptake from the extracellular

environment, other than intracellular incorporation. Other metabolic pathways for

macronutrients include those for sulfur (Figure 6), which is also enriched in blooming

than non-blooming species (Cao et al., 2020b).

4.3 Utilization of organic compounds

Organic compounds were not initially considered, even though cyanobacteria were

known to utilize organic nutrients under light (Baalen et al., 1971; Rippka, 1972; Stanier,

1973). As CyanoHAB research progresses, organics gained some attention but were

limited to organic nitrogen and phosphorus in a few species (Berman & Chava, 1999;

Chaffin & Bridgeman, 2014). The transporters internalizing organic compounds, such as

amino acids and sugars, have been identified systematically (Figure 6) and ABC amino

acid transporters are found more enriched in blooming than non-blooming species (Cao

et al., 2020b). Two transporters, AaPJMPQ for L-amino acids and NatABCDE for neutral

amino acids, are found to be encoded in nearly all blooming species. Strikingly, the top

blooming species, *M. aeruginosa* have additional transporters, HisJMPQ (for histidine),

ArtIJMQP (arginine), DppABCDF (d,d-dipeptide), GlnHPQ (Glutamine), GltIKJL

(Glutamate/aspartate), LivFGHKM (leucine, isoleucine, and valine), GluABCD

(glutamate), and AotJMPQ (arginine/ornithine). Experimentally, this has been verified:

glutamic acid, aspartic acid, leucine, alanine and arginine can be assimilated by *M.*

aeruginosa at varying rates (Dai et al., 2009). Besides, some sugars can also be

internalized by blooming cyanobacteria, e.g., lipopolysaccharides internalized by

transporter RfbAB and chitobiose by ChiEFG.

4.4 CO_2-Concentrating Mechanism (CCM)

CCM concentrates CO_2 in carboxysomes to high levels (~ 40μM) for effective

incorporation into ribulose bisphosphate in the Calvin cycle. CCM is more enriched in

blooming than non-blooming species (Cao et al., 2020b) *et al.*, 2020). The CCM genes

encode four types of functions: *ccmKLMNO* (structural proteins forming the

carboxysome), *icfA* (carbonic anhydrase), *bicA* and *sbtA* (bicarbonate facilitator),

cdnF$_3$D$_3$chY, *cdnF$_4$D$_4$chX*, and *cmpABCD* (ABC transporter for HCO_3^-), and *rbcLS*

(ribulose bisphosphate carboxylase/oxygenase). All but one (*bicA*) have more gene

copies in blooming than non-blooming species (Figure 7E). This enrichment is

understandable because heavy drawdown of CO_2 can easily lead to carbon limitation

during a bloom, particularly when a diffusion limitation effect is associated with high

population density of cyanobacteria in the water. Meanwhile, elevated CO_2 level has

been found to intensify CyanoHABs (Verspagen et al., 2014) and CCM transporters and

regulators respond quickly (Ma et al., 2019; Sandrini et al., 2016).

Figure 7. Some important CyanoPATHs associated with CyanoHABs in blooming vs. non-blooming cyanobacteria. (A): gas vesicle genes; (B): genes for phosphorus utilization; (C): nitrogen utilization pathway; (D): nitrogen utilization genes; and (E): CCM genes.

4.5 Disputed roles of cyanotoxins

Cyanotoxins are conventionally viewed as deterrents for anti-predation/anti-grazing of cyanobacteria (Zanchett & Oliveira-Filho, 2013). Recent research suggests their

functions are not limited to deterrents: their toxicity may as well be a coincidence, and
its real roles are rather different. First, a new study suggests a direct role in antioxidant
response leading to reduced ROSs outside and inside of *M. aeruginosa* cells (Malanga
et al., 2019). Second, a protein-binding role has been identified for microcystin-LR (MC-
LR), which binds several enzymes of the Calvin cycle, phycobiliproteins and two
NADPH-dependent reductases. The binding, strongly enhanced under high irradiance
and oxidative stress, is a protection and increases cell growth (Zilliges et al., 2011).
Third, they act as a signaling molecule in maintaining algal colonies and the dominance
of *Microcystis* spp. (Gan et al., 2012). Last, some cyanotoxins, e.g., microcystins, has a
deep evolutionary origin, which predates that of predators/grazers and eliminates the
possibility of directed evolution for cyanotoxins toward intoxicating animals (Rantala et
al., 2004).
408

409 **4.6 Other functions**
410 There are two additional types of functions associated with CyanoHABs: assimilation of
411 trace elements and stress resistance.
412

Trace elements and vitamins. This pathway mainly consists of various classes of
transporters for their uptake. The utilization pathways are for magnesium, ferric/ferrous
ions, copper, nickel, manganese, and cobalt (Ahern et al., 2008; Kosakowska et al.,
2007). Among them, iron (ferric and ferrous) has the most types of transporters for
siderophore-based high-affinity iron-uptake *fecBCDE*, *feoABC* (for ferrous iron),
fhuBCD, *futABC*, *sfuABC*, and *sufBCD*. Despite these driving roles reported for some
trace elements (Berman-Frank et al., 2001; Facey et al., 2019; Gu et al., 2020; Molot et
al., 2014), there are no significant changes in gene copy number between blooming and
non-blooming species (Cao et al., 2020b). For vitamins, the transporters are for
riboflavin, nicotinamide riboside, biotin (vitamin B-7), and cobalamin (vitamin B12). Most
bloom-forming species are vitamin B1 and B12 auxotrophs and their transporters have
been identified BioYMN (biotin), YbaA/PnuX/ImpX (for riboflavin), PnuC (for
nicotinamide riboside), and BtuFCD (cobalamin).
426

***Stress resistance*.** Some stressors in eutrophic waters include heavy metals,

antibiotics, and dyes (Devarajan et al., 2015; Guo et al., 2020; Li et al., 2020). Despite

the expected inhibitive effects of heavy metals on cyanobacterial growth (Richardson et

al., 2018), some antibiotics are found to be promoting in two recent studies (Li et al.,

2021; Xu et al., 2021). Here, three types of stress resistance are summarized in

CyanoPATH: heavy metal and antibiotics, and mycosporine-like amino acids (MAAs) to

shield ultraviolet light (Fig. 6). For heavy metals and antibiotics resistance, the main

metabolic functions identified so far are either ATP-powered efflux ABC transporters or

simple facilitator outward pumps relying on cross-membrane gradients as drive. The

identified efflux transporters include ChrA (chromate efflux), CnrABC (cobalt/nickel),

CopABC (copper efflux ATPase), CueO (Multicopper oxidase), CusABCF (Cu^+/Ag^+

efflux), CzcABC and ZitB (Co^{2+}, Zn^{2+}, Cd^{2+}, and Ni^{2+} efflux), CzcD (Co^{2+} and Zn^{2+} efflux),

NccABC and NreB (Ni^{2+}), and ZntA (Zn^{2+} and Cd^{2+} P-type ATPase exporter) (Cao et al.,

2020b). The antibiotic resistance genes are mostly responsible for exporting toxic

compounds and antibiotics out of cells. These transporters are EmrBA-TolC (uncouplers

and antibiotics (nalidixic acid)), LmrP (cationic dyes, tetracycline, and

daunomycin/macrolides), MdfA (cationic dyes, chloramphenicol/fluoroquinolone), NorA

(fluoroquinolones/cationic dye/inhibitors), QacA/B (monocationic biocide and dyes),

EmrE (lipophilic cations), AcrB/D-A-TolC, MexBA-OprM, MexDC-OprJ, and MexFE-

OprN (multidrugs, e.g., tetracycline, chloramphenicol, fluoroquinolones, novobiocin,

erythromycin), and MexYX-OprM (aminoglycosides).

Another type of stress is universal, UV light, since early Earth (Garcia-Pichel et al.,

2019). Cyanobacteria have been equipped with the capabilities to produce sunscreen

compounds MAAs (Jain et al., 2017; Sinha et al., 2001). The operons encoding the

biosynthetic genes have been identified in *Anabaena variabilis* and other species

(Balskus & Walsh, 2010; Miyamoto et al., 2014). A gene cluster of four synthetases

catalyzes the formation of shinorine from sedoheptulose 7-phosphate in the pentose

phosphate pathway. The synthetases are nonribosomal peptide synthetase (ava-3855),

ATP-grasp family enzyme (ava-3856), O-methyltransferase (ava_3857), and a

dehydroquinate synthase (ava_3858). Interestingly, only some of the blooming species

have one cluster whereas non-blooming species do not contain any (Cao et al., 2020b).

V. Synthesis of an CyanoHAB mechanism and outlook

Through the lens of adaptive radiation, four factors of the ecology of CyanoHABs stand

out: (1) elaborate diverse functional repertoire and low nutrient requirement in

cyanobacteria molded by early adaptive evolution, (2) cyanobacteria having lower

nutrient requirements than green algae indeed, (3) there is no directed evolution in

biological functions toward water eutrophication in cyanobacteria, (4) the CyanoHAB-

associated functional repertoire are more abundant and complete in blooming than non-

blooming species. Based on all the discussions above, we propose a comprehensive

working mechanism of CyanoHABs as a synergistic quad: superior functional repertoire,

established with long adaptive radiation under nutrient-deficient conditions and not

having evolved toward eutrophic conditions, enables cyanobacteria to efficiently utilize

elevated nutrients under current eutrophic regime for excess growth and CyanoHABs

thereof, due to their lower nutrient requirements than co-living algae. Low nutrient

requirement in cyanobacteria and higher requirements in green algae are necessary

conditions, and superior functional repertoire in cyanobacteria and water eutrophication

are sufficient conditions. One prominent feature of the quad mechanism is the relative

nutrient requirement of different algae. This boils down to the chemical composition of

the cellular machinery and how the composition can be achieved under various growth

conditions. In this sense, this quad mechanism has profound implications for essentially

all fields of life sciences.

The significance of the synthesized mechanism lies in the fact it places CyanoHABs in

the continuum of cyanobacterial evolution in aquatic environment and in the context of

community. This replacement underscores the wholistic requirement in fully

understanding occurrence and control of CyanoHABs, and in turn calls for a shift in

study approach. Here we identify three types of approaches. First, comparative

research of multiple factors in nutrient requirements and their abilities to compete for

available resources in the same water. Second, integrated systems biology assessment

of the relative roles in promoting growth in cyanobacteria, which should be based on

genome-scale metabolic networks. Generally speaking, the evolutionary ecological

perspective is applicable to various of life science areas, such as pathogens.

Acknowledgements

493 This work was supported by National Natural Science Foundation of China (32171565

494 and 52070117) and Duke Kunshan University Summer Research Scholarships and

495 Signature Work Research Grants.

References

Adrian, R., O'Reilly, C. M., Zagarese, H., Baines, S. B., Hessen, D. O., Keller, W., . . . Winder,
M. (2009). Lakes as sentinels of climate change. *Limnology and Oceanography,*
54(6part2), 2283-2297. doi:10.4319/lo.2009.54.6_part_2.2283
Ahern, K. S., Ahern, C. R., & Udy, J. W. (2008). In situ field experiment shows *Lyngbya*
majuscula (cyanobacterium) growth stimulated by added iron, phosphorus and nitrogen.
Harmful Algae, 7(4), 389-404. doi:http://dx.doi.org/10.1016/j.hal.2007.08.006
Anbar, A. D., & Knoll, A. H. (2002). Proterozoic ocean chemistry and evolution: a bioinorganic
bridge? *Science, 297*(5584), 1137-1142. doi:doi:10.1126/science.1069651
Anderson, D. M., Burkholder, J. M., Cochlan, W. P., Glibert, P. M., Gobler, C. J., Heil, C. A., . . .
Vargo, G. A. (2008). Harmful algal blooms and eutrophication: Examining linkages from
selected coastal regions of the United States. *Harmful Algae, 8*(1), 39-53.
doi:https://doi.org/10.1016/j.hal.2008.08.017
Avise, J. C., & Ayala, F. J. (2007). In the light of evolution I: Adaptation and complex design.
Proceedings of the National Academy of Sciences, 104(suppl 1), 8563.
doi:10.1073/pnas.0702066104
Baalen, C. V., Hoare, D. S., & Brandt, E. (1971). Heterotrophic Growth of Blue-Green Algae in
Dim Light. *Journal of Bacteriology, 105*(3), 685-689. doi:doi:10.1128/jb.105.3.685-
689.1971
Balskus, E. P., & Walsh, C. T. (2010). The Genetic and Molecular Basis for Sunscreen
Biosynthesis in Cyanobacteria. *Science, 329*(5999), 1653-1656.
doi:doi:10.1126/science.1193637
Beaulieu, M., Pick, F., & Gregory-Eaves, I. (2013a). Nutrients and water temperature are
significant predictors of cyanobacterial biomass in a 1147 lakes data set. *Limnology and*
Oceanography, 58(5), 1736-1746. doi:https://doi.org/10.4319/lo.2013.58.5.1736
Beaulieu, M., Pick, F., & Gregory-Eaves, I. (2013b). Nutrients and water temperature are
significant predictors of cyanobacterial biomass in a 1147 lakes data set. *Limnology and*
Oceanography, 58(5), 1736–1746.
Berman, T., & Chava, S. (1999). Algal growth on organic compounds as nitrogen sources.
Journal of Plankton Research, 21(8), 1423-1437. doi:10.1093/plankt/21.8.1423
Berman-Frank, I., Cullen, J. T., Shaked, Y., Sherrell, R. M., & Falkowski, P. G. (2001). Iron
availability, cellular iron quotas, and nitrogen fixation in Trichodesmium. *Limnology and*
Oceanography, 46(6), 1249-1260. doi:doi:10.4319/lo.2001.46.6.1249
Bianchi, T. S., Engelhaupt, E., Westman, P., Andrén, T., Rolff, C., & Elmgren, R. (2000).
Cyanobacterial blooms in the Baltic Sea: Natural or human-induced? *Limnology and*
Oceanography, 45(3), 716-726. doi:https://doi.org/10.4319/lo.2000.45.3.0716

Bjerrum, C. J., & Canfield, D. E. (2002). Ocean productivity before about 1.9 Gyr ago limited by phosphorus adsorption onto iron oxides. *Nature, 417*(6885), 159-162. doi:10.1038/417159a

Boden, J. S., Konhauser, K. O., Robbins, L. J., & Sánchez-Baracaldo, P. (2021). Timing the evolution of antioxidant enzymes in cyanobacteria. *Nature Communications, 12*(1). doi:10.1038/s41467-021-24396-y

Brock, T. D. (1985). Long-term Change in Lake Mendota. In *A eutrophic lake: Lake Mendota, Wisconsin* (pp. 203-215). New York, NY: Springer New York.

Broddrick, J. T., Rubin, B. E., Welkie, D. G., Du, N., Mih, N., Diamond, S., . . . Palsson, B. O. (2016). Unique attributes of cyanobacterial metabolism revealed by improved genome-scale metabolic modeling and essential gene analysis. *Proceedings of the National Academy of Sciences, 113*(51), E8344. doi:10.1073/pnas.1613446113

Cao, H., Shimura, Y., Steffen, M., Yang, Z., Lu, J., Joel, A., . . . Garcia-Pichel, F. (2020a). The trait repertoire enabling cyanobacteria to bloom assessed through comparative genomic complexity and metatranscriptomics. *mBio, Accepted*.

Cao, H., Shimura, Y., Steffen, M. M., Yang, Z., Lu, J., Joel, A., . . . Garcia-Pichel, F. (2020b). The Trait Repertoire Enabling Cyanobacteria to Bloom Assessed through Comparative Genomic Complexity and Metatranscriptomics. *mBio, 11*(3), e01155-01120. doi:doi:10.1128/mBio.01155-20

Carey, C. C., Ibelings, B. W., Hoffmann, E. P., Hamilton, D. P., & Brookes, J. D. (2012). Eco-physiological adaptations that favour freshwater cyanobacteria in a changing climate. *Water Research, 46*(5), 1394-1407. doi:http://dx.doi.org/10.1016/j.watres.2011.12.016

Chaffin, J. D., & Bridgeman, T. B. (2014). Organic and inorganic nitrogen utilization by nitrogen-stressed cyanobacteria during bloom conditions. *Journal of Applied Phycology, 26*(1), 299-309. doi:10.1007/s10811-013-0118-0

Chang, R. L., Ghamsari, L., Manichaikul, A., Hom, E. F. Y., Balaji, S., Fu, W., . . . Papin, J. A. (2011). Metabolic network reconstruction of Chlamydomonas offers insight into light-driven algal metabolism. *Molecular Systems Biology, 7*(1), 518. doi:https://doi.org/10.1038/msb.2011.52

Chapra, S. C., Boehlert, B., Fant, C., Bierman, V. J., Henderson, J., Mills, D., . . . Paerl, H. W. (2017). Climate Change Impacts on Harmful Algal Blooms in U.S. Freshwaters: A Screening-Level Assessment. *Environmental Science & Technology, 51*(16), 8933-8943. doi:10.1021/acs.est.7b01498

Chen, M.-Y., Teng, W.-K., Zhao, L., Hu, C.-X., Zhou, Y.-K., Han, B.-P., . . . Shu, W.-S. (2021). Comparative genomics reveals insights into cyanobacterial evolution and habitat adaptation. *The ISME Journal, 15*(1), 211-227. doi:10.1038/s41396-020-00775-z

Conroy, J. D., Kane, D. D., Dolan, D. M., Edwards, W. J., Charlton, M. N., & Culver, D. A. (2005). Temporal Trends in Lake Erie Plankton Biomass: Roles of External Phosphorus Loading and Dreissenid Mussels. *Journal of Great Lakes Research, 31*, 89-110. doi:https://doi.org/10.1016/S0380-1330(05)70307-5

Dai, R., Liu, H., Qu, J., Zhao, X., & Hou, Y. (2009). Effects of amino acids on microcystin production of the Microcystis aeruginosa. *Journal of Hazardous Materials, 161*(2), 730-736. doi:https://doi.org/10.1016/j.jhazmat.2008.04.015

Devarajan, N., Laffite, A., Graham, N. D., Meijer, M., Prabakar, K., Mubedi, J. I., . . . Poté, J. (2015). Accumulation of Clinically Relevant Antibiotic-Resistance Genes, Bacterial Load, and Metals in Freshwater Lake Sediments in Central Europe. *Environmental Science & Technology, 49*(11), 6528-6537. doi:10.1021/acs.est.5b01031

Dobzhansky, T. (1973). Nothing in Biology Makes Sense Except in the Light of Evolution. *American Biology Teacher, 35*, 125-129.

Du, W., Li, G., Ho, N., Jenkins, L., Hockaday, D., Tan, J., & Cao, H. (2020). CyanoPATH: a knowledgebase of genome-scale functional repertoire for toxic cyanobacterial blooms.

584   *Briefings in Bioinformatics, 22*(4). doi:10.1093/bib/bbaa375

585 Elser, J. J., Bracken, M. E. S., Cleland, E. E., Gruner, D. S., Harpole, W. S., Hillebrand, H., . . .
586   Smith, J. E. (2007). Global analysis of nitrogen and phosphorus limitation of primary
587   producers in freshwater, marine and terrestrial ecosystems. *Ecology Letters, 10*(12),
588   1135-1142. doi:https://doi.org/10.1111/j.1461-0248.2007.01113.x

589 Facey, J. A., Apte, S. C., & Mitrovic, S. M. (2019). A Review of the Effect of Trace Metals on
590   Freshwater Cyanobacterial Growth and Toxin Production. *Toxins, 11*(11), 643.
591   doi:10.3390/toxins11110643

592 Falkowski, P. G. (1997). Evolution of the nitrogen cycle and its influence on the biological
593   sequestration of CO2 in the ocean. *Nature, 387*(6630), 272-275. doi:10.1038/387272a0

594 Francis, G. (1878). Poisonous Australian Lake. *Nature, 18*(444), 11-12. doi:10.1038/018011d0

595 Gan, N., Xiao, Y., Zhu, L., Wu, Z., Liu, J., Hu, C., & Song, L. (2012). The role of microcystins in
596   maintaining colonies of bloom-forming Microcystis spp. *Environmental Microbiology,*
597   *14*(3), 730-742. doi:10.1111/j.1462-2920.2011.02624.x

598 García-Cañas, R., Giner-Lamia, J., Florencio, F. J., & López-Maury, L. (2021). A protease-
599   mediated mechanism regulates the cytochrome
600   c₆/plastocyanin switch in
601   Synechocystis sp. PCC 6803. *Proceedings of the National*
602   *Academy of Sciences, 118*(5), e2017898118. doi:10.1073/pnas.2017898118

603 Garcia-Pichel, F., Lombard, J., Soule, T., Dunaj, S., Wu, S. H., Wojciechowski, M. F., &
604   Giovannoni, S. J. (2019). Timing the Evolutionary Advent of Cyanobacteria and the Later
605   Great Oxidation Event Using Gene Phylogenies of a Sunscreen. *mBio, 10*(3), e00561-
606   00519. doi:doi:10.1128/mBio.00561-19

607 García-Villada, L., Rico, M., Altamirano, M. a., Sánchez-Martín, L., López-Rodas, V., & Costas,
608   E. (2004). Occurrence of copper resistant mutants in the toxic cyanobacteria Microcystis
609   aeruginosa: characterisation and future implications in the use of copper sulphate as
610   algaecide. *Water Research, 38*(8), 2207-2213.
611   doi:https://doi.org/10.1016/j.watres.2004.01.036

612 Gobler, C. J. (2020). Climate Change and Harmful Algal Blooms: Insights and perspective.
613   *Harmful Algae, 91*, 101731. doi:https://doi.org/10.1016/j.hal.2019.101731

614 Goldford, J. E., & Segrè, D. (2018). Modern views of ancient metabolic networks. *Current*
615   *Opinion in Systems Biology, 8*, 117-124. doi:https://doi.org/10.1016/j.coisb.2018.01.004

616 Gomes de Oliveira Dal'Molin, C., Quek, L.-E., Palfreyman, R. W., & Nielsen, L. K. (2011).
617   AlgaGEM – a genome-scale metabolic reconstruction of algae based on the
618   Chlamydomonas reinhardtii genome. *BMC Genomics, 12*(4), S5. doi:10.1186/1471-
619   2164-12-S4-S5

620 Gu, P., Li, Q., Zhang, W., Zheng, Z., & Luo, X. (2020). Effects of different metal ions (Ca, Cu,
621   Pb, Cd) on formation of cyanobacterial blooms. *Ecotoxicology and Environmental Safety,*
622   *189*, 109976. doi:https://doi.org/10.1016/j.ecoenv.2019.109976

623 Guo, C., Chen, Y., Xia, W., Qu, X., Yuan, H., Xie, S., & Lin, L.-S. (2020). Eutrophication and
624   heavy metal pollution patterns in the water supling lakes of China's south-to-north
625   water diversion project. *Science of The Total Environment, 711*, 134543.
626   doi:https://doi.org/10.1016/j.scitotenv.2019.134543

627 Gustafsson, B. G., Schenk, F., Blenckner, T., Eilola, K., Meier, H. E. M., Müller-Karulis, B., . . .
628   Zorita, E. (2012). Reconstructing the Development of Baltic Sea Eutrophication 1850–
629   2006. *AMBIO, 41*(6), 534-548. doi:10.1007/s13280-012-0318-x

630 Hammerschmidt, K., Landan, G., Domingues Kümmel Tria, F., Alcorta, J., & Dagan, T. (2020).
631   The Order of Trait Emergence in the Evolution of Cyanobacterial Multicellularity.
632   *Genome Biology and Evolution, 13*(2). doi:10.1093/gbe/evaa249

633 Harris, R. M., Webb, D. C., Howitt, S. M., & Cox, G. B. (2001). Characterization of PitA and PitB
634   from *Escherichia coli*. *Journal of Bacteriology, 183*(17), 5008-5014.

doi:doi:10.1128/JB.183.17.5008-5014.2001

Heisler, J., Glibert, P. M., Burkholder, J. M., Anderson, D. M., Cochlan, W., Dennison, W. C., . . . Suddleson, M. (2008). Eutrophication and harmful algal blooms: A scientific consensus. *Harmful Algae, 8*(1), 3-13. doi:http://dx.doi.org/10.1016/j.hal.2008.08.006

Jain, S., Prajapat, G., Abrar, M., Ledwani, L., Singh, A., & Agrawal, A. (2017). Cyanobacteria as efficient producers of mycosporine-like amino acids. *Journal of Basic Microbiology, 57*(9), 715-727. doi:https://doi.org/10.1002/jobm.201700044

Joosse, P. J., & Baker, D. B. (2011). Context for re-evaluating agricultural source phosphorus loadings to the Great Lakes. *Canadian Journal of Soil Science, 91*(3), 317-327. doi:10.4141/cjss10005

Juneja, A., Chaplen, F. W. R., & Murthy, G. S. (2016). Genome scale metabolic reconstruction of Chlorella variabilis for exploring its metabolic potential for biofuels. *Bioresource Technology, 213*, 103-110. doi:https://doi.org/10.1016/j.biortech.2016.02.118

Kipp, M. A., & Stüeken, E. E. (2017). Biomass recycling and Earth's early phosphorus cycle. *Science Advances, 3*(11), eaao4795. doi:doi:10.1126/sciadv.aao4795

Kirschner, M. W. (2005). The Meaning of Systems Biology. *Cell, 121*(4), 503-504. doi:10.1016/j.cell.2005.05.005

Kitchell, J. E. (Ed.) (1992). *Food Web Management: A Case Study of Lake Mendota*. New York: Springer-Verlag New York.

Klanchui, A., Khannapho, C., Phodee, A., Cheevadhanarak, S., & Meechai, A. (2012). i AK692: A genome-scale metabolic model of Spirulina platensis C1. *BMC Systems Biology, 6*(1), 71. doi:10.1186/1752-0509-6-71

Kosakowska, A., Nędzi, M., & Pempkowiak, J. (2007). Responses of the toxic cyanobacterium *Microcystis aeruginosa* to iron and humic substances. *Plant Physiology and Biochemistry, 45*(5), 365-370. doi:http://dx.doi.org/10.1016/j.plaphy.2007.03.024

Lathrop, R. C., & Carpenter, S. R. (1992). Phytoplankton and their relationship to nutrients. In J. F. Kitchell (Ed.), *Food Web Management: A Case Study of Lake Mendota* (pp. 97-126). New York, NY: Springer New York.

Li, J.-J., Chao, J.-J., McKay, R. M. L., Xu, R.-B., Wang, T., Xu, J., . . . Chang, X.-X. (2021). Antibiotic pollution promotes dominance by harmful cyanobacteria: A case study examining norfloxacin exposure in competition experiments. *Journal of Phycology, 57*(2), 677-688. doi:https://doi.org/10.1111/jpy.13133

Li, X.-d., Chen, Y.-h., Liu, C., Hong, J., Deng, H., & Yu, D.-j. (2020). Eutrophication and Related Antibiotic Resistance of Enterococci in the Minjiang River, China. *Microbial Ecology, 80*(1), 1-13. doi:10.1007/s00248-019-01464-x

López-Rodas, V., Flores-Moya, A., Maneiro, E., Perdigones, N., Marva, F., García, M. E., & Costas, E. (2007). Resistance to glyphosate in the cyanobacterium Microcystis aeruginosa as result of pre-selective mutations. *Evolutionary Ecology, 21*(4), 535-547. doi:10.1007/s10682-006-9134-8

Lürling, M., Mello, M. M. e., van Oosterhout, F., de Senerpont Domis, L., & Marinho, M. M. (2018). Response of natural cyanobacteria and algae assemblages to a nutrient pulse and elevated temperature. *Frontiers in Microbiology, 9*(1851). doi:10.3389/fmicb.2018.01851

Lynch, M. (2007). The frailty of adaptive hypotheses for the origins of organismal complexity. *Proceedings of the National Academy of Sciences, 104*(suppl 1), 8597. doi:10.1073/pnas.0702207104

Lyons, T. W., Reinhard, Christopher T., & Planavsky, Noah J. (2014). Evolution: a fixed-nitrogen fix in the early ocean? *Current Biology, 24*(7), R276-R278. doi:10.1016/j.cub.2014.02.034

Ma, J., Wang, P., Wang, X., Xu, Y., & Paerl, H. W. (2019). Cyanobacteria in eutrophic waters benefit from rising atmospheric CO_2 concentrations. *Science of The Total Environment,*

686   *691*, 1144-1154. doi:https://doi.org/10.1016/j.scitotenv.2019.07.056
687 Maberly, S. C., Pitt, J.-A., Davies, P. S., & Carvalho, L. (2020). Nitrogen and phosphorus
688   limitation and the management of small productive lakes. *Inland Waters, 10*(2), 159-172.
689   doi:10.1080/20442041.2020.1714384
690 Malanga, G., Giannuzzi, L., & Hernando, M. (2019). The possible role of microcystin (D-Leu1
691   MC-LR) as an antioxidant on Microcystis aeruginosa (Cyanophyceae). In vitro and in
692   vivo evidence. *Comparative Biochemistry and Physiology Part C: Toxicology &*
693   *Pharmacology, 225*, 108575. doi:https://doi.org/10.1016/j.cbpc.2019.108575
694 Malone, T. C., & Newton, A. (2020). The Globalization of Cultural Eutrophication in the Coastal
695   Ocean: Causes and Consequences. *Frontiers in Marine Science, 7*(670).
696   doi:10.3389/fmars.2020.00670
697 Mangan, N. M., & Brenner, M. P. (2014). Systems analysis of the CO2 concentrating
698   mechanism in cyanobacteria. *eLife, 3*, e02043. doi:10.7554/eLife.02043
699 Meyer, A. (2008). Genomes evolve, but how? *Nature, 451*(7180), 771-772.
700   doi:10.1038/451771b
701 Miyamoto, K. T., Komatsu, M., Ikeda, H., & Nojiri, H. (2014). Discovery of Gene Cluster for
702   Mycosporine-Like Amino Acid Biosynthesis from Actinomycetales Microorganisms and
703   Production of a Novel Mycosporine-Like Amino Acid by Heterologous Expression.
704   *Applied and Environmental Microbiology, 80*(16), 5028-5036.
705   doi:doi:10.1128/AEM.00727-14
706 Molot, L. A., Higgins, S. N., Schiff, S. L., Venkiteswaran, J. J., Paterson, M. J., & Baulch, H. M.
707   (2021). Phosphorus-only fertilization rapidly initiates large nitrogen-fixing cyanobacteria
708   blooms in two oligotrophic lakes. *Environmental Research Letters, 16*(6), 064078.
709   doi:10.1088/1748-9326/ac0564
710 Molot, L. A., Watson, S. B., Creed, I. F., Trick, C. G., McCabe, S. K., Verschoor, M. J., . . . Schiff,
711   S. L. (2014). A novel model for cyanobacteria bloom formation: the critical role of anoxia
712   and ferrous iron. *Freshwater Biology, 59*(6), 1323-1340.
713   doi:https://doi.org/10.1111/fwb.12334
714 Montagud, A., Navarro, E., Fernández de Córdoba, P., Urchueguía, J. F., & Patil, K. R. (2010).
715   Reconstruction and analysis of genome-scale metabolic model of a photosynthetic
716   bacterium. *BMC Systems Biology, 4*(1), 156. doi:10.1186/1752-0509-4-156
717 Moore, E. K., Jelen, B. I., Giovannelli, D., Raanan, H., & Falkowski, P. G. (2017). Metal
718   availability and the expanding network of microbial metabolisms in the Archaean eon.
719   *Nature Geoscience, 10*(9), 629-636. doi:10.1038/ngeo3006
720 Nogales, J., Gudmundsson, S., Knight, E. M., Palsson, B. O., & Thiele, I. (2012). Detailing the
721   optimality of photosynthesis in cyanobacteria through systems biology analysis.
722   *Proceedings of the National Academy of Sciences, 109*(7), 2678.
723   doi:10.1073/pnas.1117907109
724 O'Neil, J. M., Davis, T. W., Burford, M. A., & Gobler, C. J. (2012). The rise of harmful
725   cyanobacteria blooms: The potential roles of eutrophication and climate change. *Harmful*
726   *Algae, 14*(0), 313-334. doi:http://dx.doi.org/10.1016/j.hal.2011.10.027
727 Orf, G. S., Gisriel, C., & Redding, K. E. (2018). Evolution of photosynthetic reaction centers:
728   insights from the structure of the heliobacterial reaction center. *Photosynthesis*
729   *Research, 138*(1), 11-37. doi:10.1007/s11120-018-0503-2
730 Osburne, M. S., Holmbeck, B. M., Coe, A., & Chisholm, S. W. (2011). The spontaneous mutation
731   frequencies of Prochlorococcus strains are commensurate with those of other bacteria.
732   *Environmental Microbiology Reports, 3*(6), 744-749. doi:doi:10.1111/j.1758-
733   2229.2011.00293.x
734 Paerl, H. W. (2014). Mitigating Harmful Cyanobacterial Blooms in a Human- and Climatically-
735   Impacted World. *Life, 4*(4), 988-1012.
736 Paerl, H. W., Havens, K. E., Hall, N. S., Otten, T. G., Zhu, M., Xu, H., . . . Qin, B. (2020).

Mitigating a global expansion of toxic cyanobacterial blooms: confounding effects and challenges posed by climate change. *Marine and Freshwater Research, 71*(5), 579-592.

Qian, X., Kim, M. K., Kumaraswamy, G. K., Agarwal, A., Lun, D. S., & Dismukes, G. C. (2017). Flux balance analysis of photoautotrophic metabolism: Uncovering new biological details of subsystems involved in cyanobacterial photosynthesis. *Biochimica et Biophysica Acta (BBA) - Bioenergetics, 1858*(4), 276-287. doi:https://doi.org/10.1016/j.bbabio.2016.12.007

Rahav, E., & Bar-Zeev, E. (2017). Sewage outburst triggers Trichodesmium bloom and enhance N2 fixation rates. *Scientific Reports, 7*(1), 4367. doi:10.1038/s41598-017-04622-8

Rantala, A., Fewer, D. P., Hisbergues, M., Rouhiainen, L., Vaitomaa, J., Börner, T., & Sivonen, K. (2004). Phylogenetic evidence for the early evolution of microcystin synthesis. *Proceedings of the National Academy of Sciences of the United States of America, 101*(2), 568. doi:10.1073/pnas.0304489101

Raymond, J., & Segrè, D. (2006). The Effect of Oxygen on Biochemical Networks and the Evolution of Complex Life. *Science, 311*(5768), 1764-1767. doi:doi:10.1126/science.1118439

Reinhard, C. T., Planavsky, N. J., Gill, B. C., Ozaki, K., Robbins, L. J., Lyons, T. W., . . . Konhauser, K. O. (2017). Evolution of the global phosphorus cycle. *Nature, 541*(7637), 386-389. doi:10.1038/nature20772

Richardson, J., Miller, C., Maberly, S. C., Taylor, P., Globevnik, L., Hunter, P., . . . Carvalho, L. (2018). Effects of multiple stressors on cyanobacteria abundance vary with lake type. *Global Change Biology, 24*(11), 5044-5055. doi:https://doi.org/10.1111/gcb.14396

Riera, J., Voss, P. R., Carpenter, S. R., Kratz, T. K., Lillesand, T. M., Schnaiberg, J. A., . . . Wegener, M. W. (2001). Nature, society and history in two contrasting landscapes in Wisconsin, USA: Interactions between lakes and humans during the twentieth century. *Land Use Policy, 18*(1), 41-51. doi:https://doi.org/10.1016/S0264-8377(00)00044-2

Rippka, R. (1972). Photoheterotrophy and chemoheterotrophy among unicellular blue-green algae. *Archiv für Mikrobiologie, 87*(1), 93-98. doi:10.1007/BF00424781

Sabeur, H. I., Wafa, F.-S., Asma, H., & Malika, B. H. (2016). Long term characterization of Trichodesmium erythraeum blooms in Gabès Gulf (Tunisia). *Continental Shelf Research, 124*, 95-103. doi:https://doi.org/10.1016/j.csr.2016.05.007

Saito, M. A., Sigman, D. M., & Morel, F. M. M. (2003). The bioinorganic chemistry of the ancient ocean: the co-evolution of cyanobacterial metal requirements and biogeochemical cycles at the Archean–Proterozoic boundary? *Inorganica Chimica Acta, 356*, 308-318. doi:https://doi.org/10.1016/S0020-1693(03)00442-0

Sandrini, G., Tann, R. P., Schuurmans, J. M., van Beusekom, S. A. M., Matthijs, H. C. P., & Huisman, J. (2016). Diel variation in gene expression of the CO_2-concentrating mechanism during a harmful cyanobacterial bloom. *Frontiers in Microbiology, 7*(551). doi:10.3389/fmicb.2016.00551

Schindler, D. W. (1977). Evolution of phosphorus limitation in lakes. *Science, 195*(4275), 260-262. doi:10.1126/science.195.4275.260

Schindler, D. W., Carpenter, S. R., Chapra, S. C., Hecky, R. E., & Orihel, D. M. (2016). Reducing Phosphorus to Curb Lake Eutrophication is a Success. *Environmental Science & Technology, 50*(17), 8923-8929. doi:10.1021/acs.est.6b02204

Schirrmeister, B. E., de Vos, J. M., Antonelli, A., & Bagheri, H. C. (2013). Evolution of multicellularity coincided with increased diversification of cyanobacteria and the Great Oxidation Event. *Proceedings of the National Academy of Sciences, 110*(5), 1791. doi:10.1073/pnas.1209927110

Schopf, J. W. (2002). The Fossil Record: Tracing the Roots of the Cyanobacterial Lineage. In B. A. Whitton & M. Potts (Eds.), *The Ecology of Cyanobacteria: Their Diversity in Time and Space* (pp. 13-35). Dordrecht: Springer Netherlands.

Segawa, T., Takeuchi, N., Fujita, K., Aizen, V. B., Willerslev, E., & Yonezawa, T. (2018).
Demographic analysis of cyanobacteria based on the mutation rates estimated from an
ancient ice core. *Heredity, 120*(6), 562-573. doi:10.1038/s41437-017-0040-3
Shi, T., & Falkowski, P. G. (2008). Genome evolution in cyanobacteria: The stable core and the
variable shell. *Proceedings of the National Academy of Sciences, 105*(7), 2510.
doi:10.1073/pnas.0711165105
Sinha, R. P., Klisch, M., Walter Helbling, E., & Häder, D.-P. (2001). Induction of mycosporine-like
amino acids (MAAs) in cyanobacteria by solar ultraviolet-B radiation. *Journal of
Photochemistry and Photobiology B: Biology, 60*(2), 129-135.
doi:https://doi.org/10.1016/S1011-1344(01)00137-3
Stanier, R. (1973). Autotrophy and heterotrophy in unicellular blue-green algae. In N. Carr & B.
Whitton (Eds.), *The biology of blue-green algae* (pp. 501-508). Berkeley, CA.: University
of California Press.
Steffen, M. M., Belisle, B. S., Watson, S. B., Boyer, G. L., & Wilhelm, S. W. (2014). Status,
causes and controls of cyanobacterial blooms in Lake Erie. *Journal of Great Lakes
Research, 40*(2), 215-225. doi:https://doi.org/10.1016/j.jglr.2013.12.012
Steffensen, D. A. (2008). Economic cost of cyanobacterial blooms. In H. K. Hudnell (Ed.),
Cyanobacterial Harmful Algal Blooms: State of the Science and Research Needs (pp.
855-865). New York, NY: Springer New York.
Tomitani, A., Knoll, A. H., Cavanaugh, C. M., & Ohno, T. (2006). The evolutionary diversification
of cyanobacteria: Molecular–phylogenetic and paleontological perspectives.
Proceedings of the National Academy of Sciences, 103(14), 5442.
doi:10.1073/pnas.0600999103
Verspagen, J. M. H., Van de Waal, D. B., Finke, J. F., Visser, P. M., Van Donk, E., & Huisman, J.
(2014). Rising CO_2 levels will intensify phytoplankton blooms in eutrophic and
hypertrophic lakes. *PloS One, 9*(8), e104325. doi:10.1371/journal.pone.0104325
Vu, T. T., Stolyar, S. M., Pinchuk, G. E., Hill, E. A., Kucek, L. A., Brown, R. N., . . . Reed, J. L.
(2012). Genome-Scale Modeling of Light-Driven Reductant Partitioning and Carbon
Fluxes in Diazotrophic Unicellular Cyanobacterium Cyanothece sp. ATCC 51142. *PLoS
Computational Biology, 8*(4), e1002460. doi:10.1371/journal.pcbi.1002460
Walsby, A. E. (1972). Structure and function of gas vacuoles. *Bacteriological Reviews, 36*(1), 1-
32. doi:doi:10.1128/br.36.1.1-32.1972
Walsby, A. E. (1994). Gas vesicles. *Microbiological Reviews, 58*(1), 94-144.
doi:doi:10.1128/mr.58.1.94-144.1994
Walsby, A. E., Hayes, P. K., Boje, R., & Stal, L. J. (1997). The selective advantage of buoyancy
provided by gas vesicles for planktonic cyanobacteria in the Baltic Sea. *New Phytologist,
136*(3), 407-417. doi:10.1046/j.1469-8137.1997.00754.x
Wilhelm, S. W., Bullerjahn, G. S., McKay, R. M. L., & Moran, M. A. (2020). The Complicated and
Confusing Ecology of *Microcystis* Blooms. *mBio, 11*(3), e00529-00520.
doi:doi:10.1128/mBio.00529-20
Wilkinson, G. M., Carpenter, S. R., Cole, J. J., Pace, M. L., Batt, R. D., Buelo, C. D., &
Kurtzweil, J. T. (2018). Early warning signals precede cyanobacterial blooms in multiple
whole-lake experiments. *Ecological Monographs, 88*(2), 188-203.
doi:https://doi.org/10.1002/ecm.1286
Wilson Alan, E., Wilson Whitney, A., & Hay Mark, E. (2006). Intraspecific Variation in Growth and
Morphology of the Bloom-Forming Cyanobacterium Microcystis aeruginosa. *Applied and
Environmental Microbiology, 72*(11), 7386-7389. doi:10.1128/AEM.00834-06
Xu, S., Jiang, Y., Liu, Y., & Zhang, J. (2021). Antibiotic-accelerated cyanobacterial growth and
aquatic community succession towards the formation of cyanobacterial bloom in
eutrophic lake water. *Environmental Pollution, 290*, 118057.
doi:https://doi.org/10.1016/j.envpol.2021.118057

Yoshikawa, K., Aikawa, S., Kojima, Y., Toya, Y., Furusawa, C., Kondo, A., & Shimizu, H. (2015).
Construction of a Genome-Scale Metabolic Model of Arthrospira platensis NIES-39 and
Metabolic Design for Cyanobacterial Bioproduction. *PloS One, 10*(12), e0144430.
doi:10.1371/journal.pone.0144430
Zanchett, G., & Oliveira-Filho, E. C. (2013). Cyanobacteria and Cyanotoxins: From Impacts on
Aquatic Ecosystems and Human Health to Anticarcinogenic Effects. *Toxins, 5*(10), 1896-
1917.
Zilliges, Y., Kehr, J.-C., Meissner, S., Ishida, K., Mikkat, S., Hagemann, M., . . . Dittmann, E.
(2011). The Cyanobacterial Hepatotoxin Microcystin Binds to Proteins and Increases the
Fitness of Microcystis under Oxidative Stress Conditions. *PLOS ONE, 6*(3), e17615.
doi:10.1371/journal.pone.0017615
Zuñiga, C., Li, C.-T., Huelsman, T., Levering, J., Zielinski, D. C., McConnell, B. O., . . . Zengler,
K. (2016). Genome-Scale Metabolic Model for the Green Alga Chlorella vulgaris UTEX
395 Accurately Predicts Phenotypes under Autotrophic, Heterotrophic, and Mixotrophic
Growth Conditions. *Plant Physiology, 172*(1), 589-602. doi:10.1104/pp.16.00593
Zhu, W., Wan, L. & Zhao, L. (2010) Effect of nutrient level on phytoplankton community
structure in different water bodies. *Journal of Environmental Sciences 22*, 32–39.
856

Figure Legend

Figure 1

Figure 1. The evolutionary trajectory of cyanobacteria in the presence of oxygen. (A) The origin

and distribution of SODs in different habitats. The timing of origin and habitat distribution of SOD

genes among taxa is created based on Boden et al (Boden et al., 2021). Starting points of

horizontal bars represent time of origin and color SOD types: CuZnSOD (blue), NiSOD (green),

and Fe- and Mn-utilizing SODs (yellow). (B) The oxygen level in atmosphere and nitrogen

fixation in heterocysts in cyanobacteria in the presence of elevated oxygen levels. The level of

oxygen is based on (Tomitani et al., 2006). (C) and (D) The temporal morphological

diversification cyanobacteria is based on Schirrmeistera et al (Schirrmeister et al., 2013).

Figure 2

Figure 2. Cyanobacterial adaptation strategies in nine environments, namely soil, desert, forest,

salty area, oil-spilled ocean, open ocean, limestone, geothermal water, and polar region.

Cyanobacteria also cope with the damage caused by ultraviolet light.

Figure 3

Figure 3. The phylogeny of cyanobacteria based on 16S ribosomal genes. Blue labels: bloom-

forming cyanobacteria; and red labels: non-bloom-forming cyanobacteria. The shapes of

different colors represent they types of morphology.

Figure 4

Figure 4. Figure 4. Lower nutrient requirements by cyanobacteria than green algae in aquatic

environment. Among seven selected media, the recipes of mAC, MJ, VTAC, and CHU-11 (left

panel) are used for culturing green algae and the recipes of M11, BG11 and CT (right panel) are

used for culturing cyanobacteria.

Figure 5

Figure 5. dN/dS for all homologous genes in 20 genomes of *M. aeruginosa*. (A) An overall

distribution of dN/dS for all homologous genes; and (B) the dN/dS for genes in different groups

of functions. dN/dS values were calculated by CODEML from the PAML 4.9j package with

default settings (Yang, 2007). AAPEP: uptake of amino acids and peptides; CCM: CO_2-

concentrating mechanism; DrugR: antibiotics resistance; MAA: UV radiation; MetalR: heavy

metal resistance; Nfix: nitrogen fixation; Nitrogen: nitrogen utilization; Osmos: osmos

homeostasis; OSR: oxidative stress resistance; PBS: phycobilisome; Phosphorus:

inorganic/organic utilization; PS-I/PS-II: photosystem I and photosystem II; Sugar: sugar

assimilation; TEVit: assimilation of trace metals and vitamins; Toxins: cyanotoxins; UFA:

unsaturated fatty acids; Vesicle: gas vesicles. Purple bars are pathways strongly associated

with CyanoHABs; green bars are pathways not specifically related to CyanoHABs; the last

seven blue bars represent essential biological functions (Rubin *et al.*, 2015).

Figure 6

Figure 6. The core and query pathways in *Aphanizomenon flos-aquae* NIES-81. The pathways

labeled with dashed borders are not complete (due to the absence of required components) in

this strain but may be complete in others.

Figure 7

Figure 7. Some important CyanoPATHs associated with CyanoHABs in blooming vs. non-

blooming cyanobacteria. (A): gas vesicle genes; (B): genes for phosphorus utilization; (C):

nitrogen utilization pathway; (D): nitrogen utilization genes; and (E): CCM genes.