Cooperative binding of TCR and CD4 to pMHC enhances TCR sensitivity

Muaz Nik Rushdi[1,2§†], Victor Pan[1,2§‡], Kaitao Li[1,2§], Stefano Travaglino[1,2], Hyun-Kyu Choi[1,2],

Jinsung Hong[1,2¶], Fletcher Griffitts[2,3], Pragati Agnihotri[4,5#], Roy A. Mariuzza[4,5], Yonggang

Ke[1,6*], Cheng Zhu[1,2,3*]

[1]Wallace H. Coulter Department of Biomedical Engineering, Georgia Institute of Technology and

Emory University, Atlanta, GA, USA; [2]Parker H. Petit Institute for Bioengineering and

Biosciences, [3]Georgia W. Woodruff School of Mechanical Engineering, Georgia Institute of

Technology, Atlanta, GA, USA; [4]W. M. Keck Laboratory for Structural Biology, Institute for

Bioscience and Biotechnology Research, University of Maryland, Rockville, MD, USA;

[5]Department of Cell Biology and Molecular Genetics, University of Maryland, College Park, MD,

USA; [6]Medical Scientist Training Program, Emory University School of Medicine, Atlanta, GA,

USA;

[§]These authors contributed equally

[*]Corresponding authors Emails: cheng.zhu@bme.gatech.edu, yonggang.ke@emory.edu

[†]Present Address: Medtronic CO., Minneapolis, MN, USA

[‡]Present Address: Parabon Nanolabs, Huntington, WV, USA

[¶]Present Address: Food and Drug Administration, Silver Spring, MD, USA

[#]Present Address: Advanced Bioscience Laboratories, Rockville, MD, USA

Abstract

Antigen recognition of CD4[+] T cells by the T cell receptor (TCR) can be greatly enhanced by the coreceptor CD4[1-7]. Yet, understanding of the molecular mechanism is hindered by the ultra-low affinity of CD4 binding to class-II peptide-major histocompatibility complexes (pMHC)[1,7-10]. Using two-dimensional (2D) mechanical-based assays, we determined a CD4–pMHC interaction to have 3-4 logs lower affinity than cognate TCR–pMHC interactions[8], and to be susceptible to increased dissociation by forces (slip bond)[5,8,11]. In contrast, CD4 binds TCR-prebound pMHC at 5-6 logs higher affinity, forming TCR–pMHC–CD4 trimolecular bonds that are prolonged by force (catch bond)[5,8,11] and modulated by protein mobility on the cell membrane, indicating profound TCR–CD4 cooperativity. Consistent with a tri-crystal structure[12], using DNA origami as a molecular ruler to titrate spacing between TCR and CD4 indicates that 7-nm proximity optimizes trimolecular bond formation with pMHC. Our results reveal how CD4 augments TCR antigen recognition.

Main

CD4[+] T cells play vital and versatile roles in the adaptive immunity by differentiating

into lineages of various functions upon activation by antigen. Antigen recognition is achieved by

the T cell receptor (TCR) interacting with its cognate peptide bound to major histocompatibility

complex (MHC, or pMHC when including the peptide) expressed on an antigen-presenting cell

(APC) or target cell. TCR–pMHC binding triggers phosphorylation of CD3 immunoreceptor

tyrosine-based activation motifs (ITAMs), leading to a cascade of signaling events that result in

T cell activation and effector function[13]. Expressed on the surface of their respective T cell

subsets, CD8 and CD4 coreceptors are thought to facilitate the initiation of the intracellular

signaling cascades by stabilizing TCR–pMHC binding and delivering its associated Lck to

TCR[14,15]. However, unlike CD8, CD4's extremely low affinity for MHC hinders our

understanding of these proposed mechanisms that require CD4 binding to pMHC[9,16]. Indeed,

surface plasmon resonance (SPR) was unable to measure CD4–MHC interactions in three-

dimensions (3D)[7,17], except for one report of a ~200 μM K_d[10]. Although affinity measurements

were successfully attained using B cells and a supported lipid bilayer reconstituted with CD4[7], it

required additional CD2–CD58 interactions to stabilize the 2D interface. The estimated 2D K_d of

~5,000 μm^{-2} is >1-log higher than the K_d of TCR–pMHC interactions[18] or the CD4 density of T

cell surface[8], neither of which predicts CD4–MHC interaction *in vivo*.

Also due to the extremely low affinity of CD4–MHC, the ectodomain crystal structure of

TCR–pMHC–CD4 tri-molecular complex was not available until a gain-of-function CD4 mutant

with substantially higher affinity toward MHC was engineered[12]. Fascinatingly, the rigid

coreceptor maintains a membrane proximal spacing of ~7 nm apart from the TCR while binding

to the MHC, providing an adequate gap for the missing CD3 proteins to amalgamate around the

TCR[17,19]. The co-crystal structure suggests that the membrane proximal positioning of TCR and CD4 ectodomains is energetically favorable to form a trimolecular complex with pMHC. Due to the differential affinities to pMHC, it is unlikely driven by the CD4 binding, but instead needs the TCR binding to pMHC to initiate the trimolecular complex formation. The above reasoning provides a rationale of our hypothesis that CD4 augmentation comes into play via its synergy with TCR, a mechanism similar to that in TCR-CD8 cooperation[3-5,20]. However, such cooperativity remains elusive with conflicting observations from previous studies. Our previous study using a mechanical 2D assay showed no effect of CD4 on TCR–pMHC binding under force-free conditions[8]. Studies of others using Forster Resonance Energy Transfer (FRET)[18] and pMHC tetramer staining techniques[21] failed to detect CD4 enhancement, whereas another study using Fluorescence Recovery After Photobleaching (FRAP) observed that TCR and CD4 reciprocally enhance each other's binding to pMHC[22]. Most importantly, CD4's ability to bind pMHC is still required for TCR-mediated IL-2 production even when its association with Lck is abolished[22].

Combining our 2D kinetic assays and mathematical modeling, we analyzed the bimolecular CD4–MHC interaction and TCR-CD4 cooperativity using both purified protein ectodomains and cell systems. We identified enormous cooperative binding of CD4 to TCR-prebound pMHC, which has 5-6 logs higher affinity than binding to free pMHC. This cooperativity manifests substantially greater number and longer-lived bonds of pMHC with both TCR and CD4 than the sum of separate pMHC bonds with TCR or CD4 as caused by the emergence of a dominating TCR–pMHC–CD4 trimolecular bond species that is dynamically regulated by mechanical force applied on the molecular complex. By presenting TCR and CD4 on different surfaces to modulate molecular diffusion or DNA origami to modulate molecular

spacing, we showed that the TCR-CD4 cooperativity requires molecular mobility to allow for their colocalization within a 7-nm proximity, consistent with that reported in TCR–pMHC–CD4 tri-crystal structure. Our results explain how CD4 could efficiently augment TCR antigen recognition via cooperative binding to TCR-prebound pMHC, thereby boosting TCR sensitivity. They also underline the importance of physical factors in regulating T-cell antigen recognition, which may be imposed on molecular organization at the cell membrane via signaling dependent and/or independent mechanisms.

Results

CD4–MHC binding displays low 2D binding affinity

We used our adhesion frequency assay[23,24] (Fig. 1a), which analyzes cross-junctional interactions upon initial cell-cell encounters, to quantify the 2D binding between HLA-DR1 presenting a melanoma antigenic peptide derived from the glycolytic enzyme triosephosphate isomerase (TPI:HLA-DR1, referred hereafter as pMHC, unless specified)[25] and its cognate E8 TCR, wild-type (WT) CD4, or a high-affinity mutant (MT) CD4 generated via directed evolution[17]. Ectodomains of each binding pair were respectively coated on two opposing RBCs, which were brought into contact and then separated to detect adhesion by RBC elongation. Results were presented as adhesion frequency (P_a) of many repeated contact-separation cycles, which was higher between RBCs functionalized with designated molecules at appropriate densities than between blank ones at the same contact time t_c, indicating specific interactions (Fig. 1b). P_a increased initially and plateaued as the probability of bond formation was balanced by that of dissociation with prolonged t_c, resulting in a steady-state (Fig. 1c). We evaluated the effective 2D affinity (A_cK_a) and off-rate (k_{-1}) by fitting the P_a vs t_c curve (Fig. 1c) or the log-transformed and molecular density-normalized counterparts (Fig. 1d) to a published model[23,24]

(Table 1). The log transformation converts P_a to the average number of bonds per contact, $\langle n \rangle =$

$- \ln(1 - P_a)$, which, after normalization by densities of the receptor (m_r) and ligand (m_l) collapses

three curves measured at different $m_r \times m_l$ values (indicated by different symbols in Fig. 1d) and

approaches to $A_c K_a$ at large t_c. This model was well supported as the steady-state $\langle n \rangle$ increased

linearly with $m_r \times m_l$ (Supplementary Fig. 1a-c) and collapsed into a single plateau upon

normalization by $m_r \times m_l$ (Fig. 1e), suggesting the lack of multimeric binding and supporting the

reliability of our measurements[23,24]. The effective 2D affinity of E8 TCR for pMHC ($A_c K_{a,TCR} =$

$7.70 \pm 0.40 \times 10^{-4}$ μm^4) is comparable to, but off-rate ($k_{-1,TCR} = 0.48 \pm 0.07$ s^{-1}) is slower than,

other TCR–pMHC-II interactions[8]. Remarkably, the effective 2D affinity of CD4–MHC-II

binding ($A_c K_{a,CD4} = 4.35 \pm 0.49 \times 10^{-7}$ μm^4) is >3 logs lower and among the lowest values

recorded[26] (Fig. 1e). Consistent with its 3D SPR measurement[17], the affinity of MT CD4 for

MHC ($2.48 \pm 0.15 \times 10^{-3}$ μm^4) is 3-4 logs higher than that of WT CD4 (Fig. 1e, Table 1).

TCR and CD4 cooperatively binds to pMHC

To explain how CD4–MHC binding could disproportionally recruit Lck to TCR[7], we

tested the hypothesis that TCR and CD4 bind pMHC synergistically as observed for TCR and

CD8[3-5]. E8 TCR and CD4 were co-functionalized on RBCs at different density ratios

(m_{TCR}/m_{CD4}) while keeping their sum constant and tested against RBCs bearing pMHC with the

same density (m_{pMHC}). We first analyzed the P_a vs t_c curves so measured (Fig. 2a) using a dual-

species concurrent and independent binding model[27-29], opposite to our synergistic binding

hypothesis. Without synergy, this model predicts that the average whole number of bonds $\langle n \rangle_w$

is equal to the sum of its parts, i.e., the average numbers of TCR–pMHC bonds $\langle n \rangle_{TCR}$ plus

CD4–MHC bonds $\langle n \rangle_{CD4}$. Given the low CD4 density, which was far from enough to make up

its low affinity, the measured $\langle n \rangle_w$ should have yielded $\langle n \rangle_{TCR}$ as its only detectable component

and normalizing it by m_{TCR} and m_{pMHC} should have collapsed the entire family of curves as did

Fig. 1d. To the contrary, the $<n>_W/(m_{TCR}\times m_{pMHC})$ vs t_c curve upward shifted as m_{CD4} increased

(Fig. 2b and Supplementary Fig. 2a), contradicting the independent binding assumption[27-29].

Instead of increasing linearly as in Supplementary Fig. 1, the steady-state $<n>_W$ increased

nonlinearly with increasing $m_{TCR}\times m_{pMHC}$ (Supplementary Fig. 2b). Instead of being a constant as

Fig. 1e, the TCR- and pMHC-density normalized whole bond number, $<n>_W/m_{TCR}\times m_{pMHC}$,

increased linearly with increasing CD4 density m_{CD4} (Supplementary Fig. 2c). These results

support our hypothesis and indicate that TCR and CD4 cooperatively bind to pMHC, manifesting

higher number of whole bonds than the sum of TCR–pMHC and CD4–pMHC bonds as parts.

From the difference between the whole and the sum of parts, we identified a new bond

species – the emerging TCR–pMHC–CD4 trimolecular bonds whose average number is $<n>_{tri} =$

$\Delta<n> = <n>_W - (<n>_{TCR} + <n>_{CD4})$. We developed a kinetics model in which CD4 binding to

free pMHC is assumed negligible compared to TCR, but CD4 can bind pMHC prebound by TCR

with a higher affinity $K^*_{a,CD4}(= k_{+2,CD4}/k_{-2,CD4})$. This model predicts:

$$\frac{\langle n \rangle_W}{m_{TCR}m_{pMHC}} = A_c K_{a,TCR}\left(1 + m_{CD4}K^*_{a,CD4}\right)\left[1 + \frac{\lambda_2(\lambda_2 - k_{-1,TCR})e^{-\lambda_1 t_c} - \lambda_1(\lambda_1 - k_{-1,TCR})e^{-\lambda_2 t_c}}{(\lambda_1 - \lambda_2)(m_{CD4}k_{+2,CD4} + k_{-2,CD4})}\right] \quad (1)$$

where λ_1 and λ_2 are positive functions of off-rate of TCR–pMHC bond ($k_{-1,TCR}$), on- and off-rate
of CD4 binding to ($k_{+2,CD4}$) and dissociating from ($k_{-2,CD4}$) TCR-prebound pMHC, and CD4 density
(m_{CD4}):

$$\lambda_{1,2} = \frac{1}{2}\left(k_{-1,TCR} + k_{-2,CD4} + m_{CD4}k_{+2,CD4}\right)$$

$$\pm \frac{1}{2}\left[(k_{-1,TCR} + k_{-2,CD4} + m_{CD4}k_{+2,CD4})^2 - 4k_{-1,TCR}k_{-2,CD4}\right]^{1/2}. \quad (2)$$

The steady-state solution as $t_c \rightarrow \infty$ is

$$\frac{\langle n \rangle_{W,\infty}}{m_{TCR}m_{pMHC}} = A_c K_{a,TCR} + m_{CD4}A_c K_{a,TCR}K^*_{a,CD4} \quad (3)$$

The second term on the right-hand side of Eq. 3 can be identified as the steady-state average

number of trimolecular bonds $<n>_{tri}$ per densities of TCR and pMHC. Both the original (Fig. 2a)

and normalized (Fig. 2b) data were well fitted by Eqs. 1&2, which returned values of $A_cK_{a,TCR}$ =

$8.31 \pm 0.46 \times 10^{-4}$ μm^4 and $k_{-1,TCR} = 0.53 \pm 0.05 \times$ s^{-1} for the TCR–pMHC interaction that agree

well with previous single receptor–ligand species measurements without CD4 (Table 1).

Remarkably, the value of $K^*_{a,CD4}$ is 0.0526 ± 0.0072 μm^2, >250 times higher than the previous

value obtained using CD4-reconstituted supported lipid bilayers[7] and >350,000 times higher than

the affinity for free pMHC calculated by dividing the $A_cK_{a,CD4}$ value determined from Fig. 1d by

the ~3 μm^2 contact area A_c estimated from the side-view microscopic image.

To visualize the emergence of TCR–pMHC–CD4 trimolecular bonds, we transformed the

data according to Eq. 3: subtracting the black data set ($m_{CD4} = 0$) from the colored data sets (m_{CD4}

> 0) and normalizing them by their corresponding CD4 density. This nearly collapses the three

colored data sets into the average number of trimolecular bonds per unit densities of TCR,

pMHC, and CD4. This $<n>_{tri}/(m_{TCR} \times m_{pMHC} \times m_{CD4})$ data vs contact time t_c plot compares well

with our model prediction using the best-fit kinetic parameters obtained earlier, showing how the

degree of cooperativity increase in time until reaching the steady-state value $A_cK_{a,TCR}K^*_{a,CD4}$

(Fig. 2c). Thus, the steady-state trimolecular bond number per unit density of TCR, CD4, and

pMHC is equal to the product of the effective 2D affinity of the TCR–pMHC bond and the

affinity of CD4 for TCR-prebound pMHC, an intuitive result that make very good sense.

TCR–pMHC–CD4 trimolecular bond shows a much greater catch than TCR–pMHC

bimolecular bond

To examine whether the trimolecular bond can better sustain mechanical force, we used

the biomembrane force probe (BFP) thermal fluctuation[24,30] and force-clamp[5,8,11] assays to

measure respective force-free and force-dependent lifetimes of single bonds between probes

bearing pMHC and targets presenting ectodomains of E8 TCR, CD4, or both mixed at a 1:1

concentration ratio. The thermal fluctuation assay yielded fairly linear survival probability vs bond lifetime semi-log plots for dissociation of single-receptor species (TCR or CD4) and dual-receptor species (TCR and CD4) from TPI:HLA-DR1, which were much longer lived than the negative control 9V:HLA-A2 (Fig. 2d), conforming binding specificity of TCR and/or CD4.

The force-clamp assay found the CD4–MHC bond lifetimes to decrease with increasing force, displaying a slip-only bond behavior[5,8,11,31] (Fig. 2e). The TCR–pMHC interaction, in contrast, behaved as a catch-slip bond where the lifetime increased, reached a peak, and decreased in a biphasic fashion as force increased[5,8,11,31] (Fig. 2e). Remarkably, the interactions between pMHC probes and targets co-presenting TCR and CD4 exhibited a much more pronounced catch-slip bond (Fig. 2e). Since lifetimes were measured at infrequent binding (<20%) to ensure >89% single-bond probability, the "whole" lifetime ensemble included subpopulations of single TCR–pMHC, CD4–MHC, and TCR–pMHC–CD4 bond lifetimes, where the whole lifetime represented a weighted sum of all three contributions. Average lifetimes of whole bonds were much longer than the weighted sum of two parts – the average lifetimes of TCR–pMHC and CD4–MHC bonds – regardless of how their relative weights were adjusted (Fig. 2e), as it cannot be longer than the longer component lifetime of the two. This demonstrates the existence of a more "catchy" and longer-lived trimolecular bond species.

We used a single-state, single-pathway dissociation model for the slip-only bonds of CD4–MHC interaction and two-state, two-pathway dissociation models for the catch-slip bonds of respective TCR–pMHC and TCR–pMHC–CD4 interactions (Supplementary Fig. 3a) to globally fit the measured lifetime vs force scattergrams (Supplementary Fig. 3b). The models fit the mean ± SE bond lifetime vs force data well (Fig. 2e) and predict the probability and fraction of strong state of TCR–pMHC and TCR–pMHC–CD4 bonds (Supplementary Fig. 3e-h) as well

as the average bond lifetime vs force curves of the strong state, weak state, and their sum (Fig.

2f), which reveals how the TCR–pMHC–CD4 bond is activated to a strong state by force to form

a cooperative catch bond.

These results are similar to our previous data of class-I pMHC interacting with TCR and

CD8 on OT1 thymocytes[5], but were obtained using purified proteins that lack transmembrane

and cytoplasmic domains as well as the CD3 signaling chains, which were shown important in

our previous studies[3,5]. To test the results of the cell-free system in a live-cell system, we

repeated the experiments using Jurkat cells transfected with, and sorted to express comparable

levels of E8 TCR, CD4, or both. We first analyzed the P_a vs t_c curves of these Jurkat cells

binding to TPI:HLA-DR1 coated bead using the BFP setup (Fig. 3a). Comparing to purified

TCR$\alpha\beta$ coated on RBCs, cellular TCR-CD3 complex exhibits a ~1 log lower effective 2D

affinity ($A_cK_{a,TCR} = 3.59 \pm 0.37 \times 10^{-5} \mu m^4$), which may be due to the much smaller functional

contact area formed between bead and rough Jurkat cell surface vs between two smooth RBC

surfaces[32]. In contrast, despite the smaller functional contact area, cellular CD4 has a ~1 log

higher effective 2D affinity ($A_cK_{a,CD4} = 6.57 \pm 0.64 \times 10^{-6} \mu m^4$) for pMHC than purified CD4,

suggesting that the cellular environment may enhance its 2D interactions on the cell membrane.

Similar to the cell-free system, measurements with cells expressing both TCR and CD4 showed

significantly higher adhesion frequency data (Fig. 3a, black) than the predicted P_a vs t_c curve

mediated by the sum of TCR–pMHC and CD4–MHC interactions calculated using their

measured kinetic rates and molecular densities (Fig. 3a, dashed curve), demonstrating synergistic

binding of TCR and CD4 to pMHC on the cell system.

To compensate the >1 log lower affinity of CD4 than TCR, a >1 log higher pMHC

density was used to measure binding of CD4 than TCR (Supplementary Table 1). To better

visualize TCR-CD4 cooperativity, we applied the log transformation and site density

normalization to enable a more objective comparison of the TCR and CD4 bonds (Fig. 3b). It is

evident that their normalized predicted sum, $(<n>_{TCR} + <n>_{CD4})/(m_{TCR} \times m_{pMHC})$, is significantly

lower than the measured normalized whole bond number, $<n>_W/m_{TCR} \times m_{pMHC}$, at all contact time

points (compare the dashed curve to black circles in Fig. 3b), confirming TCR-CD4 cooperative

binding of pMHC.

7   Comparing the prediction of Eqs. 1&2 with the $<n>_W/m_{TCR} \times m_{pMHC}$ vs t_c measurement

showed excellent agreement between the model and data in both the logarithmic transformed

variable (Fig. 3b) and the directly measured variable (Fig. 3a). Fitting the data returned values of

$K^*_{a,CD4} = 0.269 \pm 0.114$ μm^2, ~5 times the value estimated from the cell-free system, and $k_{-2, CD4} =$

0.131 ± 0.117 s^{-1}, ~1 log smaller than the value estimated from the cell-free system (Table 1).

Interestingly, the values of $A_c K_{a,TCR} = 1.42 \pm 0.463 \times 10^{-4}$ μm^4 and $k_{-1,TCR} = 1.60 \pm 1.06$ s^{-1},

estimated from fitting the $<n>_W/m_{TCR} \times m_{pMHC}$ data are, respectively, 4 and 4.7 times of the

counterpart values estimated from fitting the respective $<n>_{TCR}/m_{TCR} \times m_{pMHC}$ and

$<n>_{CD4}/m_{TCR} \times m_{pMHC}$ data, which is in contrast to the cell-free system where the corresponding

numbers match (Table 1). Indeed, fitting the same whole bond data by Eqs. 1&2 using the

$A_c K_{a,TCR}$ and $k_{-1,TCR}$ values estimated from single bond species measurements as given rather than

free fitting parameters resulted in a poor fit of significantly slower kinetics than the actual data

for contact times ≤ 4s (Fig. 3c), in sharp contrast to the excellent fit in Fig. 2b that returned

higher affinity and faster off-rate for the TCR. To compensate for the pre-determined lower

$A_c K_{a,TCR}$ and slower $k_{-1,TCR}$ values, the two-parameter fit (Fig. 3c) required much higher affinity

$(K^*_{a,CD4} = 0.726 \pm 0.152$ $\mu m^2)$ and off-rate $(k_{-2, CD4} = 10.0 \pm 16.9$ s$^{-1})$ for CD4 to respectively bind

to and dissociate from TCR-prebound pMHC as compared to the four-parameter fit (Fig. 3b).

These results reveal similarities and distinctions between the cell and cell-free systems and

suggest that intracellular regulations, which are absent in the cell-free system but present in the

cell system, may speed up the first-stage kinetics to allow CD4 to cooperate in pMHC binding at

an earlier time, a contention consistent with our previous results that intracellular regulations is

important to the TCR-CD8 cooperative binding of class-I pMHC[3,5].

Cellular TCR-CD4 cooperativity was also confirmed in bond lifetime analysis: TPI:HLA-

DR1 formed a slip-only bond, weak catch-slip bond, and strong catch-slip bond with cells

expressing CD4 only, E8 TCR only, and both E8 TCR and CD4, respectively, similar to their

purified protein counterparts (Fig. 3d). In addition, Jurkat cells expressing a higher level of TCR

than CD4 formed a more pronounced catch bond than those express similar levels of TCR and

CD4 (Fig. 3d), suggesting a higher degree of cooperativity.

As an additional confirmation, we analyzed $CD4^+CD8^-$ and $CD4^-CD8^+$ single-positive T

cells[33] from 3.L2 TCR transgenic mice, which recognizes a hemoglobulin (Hb) epitope

presented by mouse class-II pMHC I-E^k (Hb:I-E^k)[8]. Our previous adhesion frequency experiment

was unable to detect mouse CD4–Hb:I-E^k binding, indicating an $A_cK_{a,CD4} < 7.0 \times 10^{-8}$ μm^4 (the

detection limit), a log lower than the human CD4–HLA-DR1 effective 2D affinity, and observed

a TCR–pMHC catch-slip bond between BFP probes bearing Hb:I-E^k and $CD4^-CD8^+$ 3.L2 T cells

by force-clamp BFP experiment[8]. Interestingly, when using $CD4^+CD8^-$ T cells we observed a

much more pronounced catch-slip bonds, which was suppressed by anti-CD4 (GK1.5) blocking

to the level of using $CD4^-CD8^+$ T cells (Fig. 3e). These data confirm that the prolonged lifetimes

were CD4-dependent and further support the formation of more durable and force-resistant

TCR–pMHC–CD4 trimolecular bonds by TCR-CD4 cooperation.

TCR-CD4 cooperativity requires molecular mobility

The faster cooperativity observed in cells (Fig. 3c) may suggest enhanced CD4 recruitment to TCR–pMHC, a process that requires both molecular mobility and proximity. To test whether cooperativity depends on molecular diffusion, we used four different pairings of surfaces to present proteins: 1) pMHC on RBCs interacting with TCR only on RBCs; 2) pMHC on beads interacting with TCR and CD4 on RBCs; 3) pMHC on RBCs interacting with TCR and CD4 on beads; and 4) pMHC on RBCs interacting with TCR and CD4 on beads. The idea is to allow lateral diffusion of TCR and CD4 on the RBC membrane but prevent them from doing so on the bead surface. The data were presented as average number of whole bonds per unit densities of TCR and pMHC as in Fig. 2b. When interacting proteins on both sides were coated on RBCs, pMHC interaction with TCR-CD4 dual-receptor species (Fig. 3f, black) generated a much higher $\langle n \rangle_w/(m_{TCR} \times m_{pMHC})$ curve than did with TCR single-receptor species (Fig. 3f, green), indicating cooperative binding as in Fig. 2b. In sharp contrast, when pMHC was coated on RBCs and TCR and CD4 were co-functionalized on beads, the binding curve was substantially downward shifted to the level indistinguishable to the curve generated by coating TCR-only on RBCs binding to RBCs bearing pMHC (Fig. 3f, orange). This indicates that the cooperativity between TCR and CD4 vanished when they were immobilized at low densities on glass bead surface. When pMHC was coated on beads and the TCR/CD4 mixture was coated on RBCs, the binding curve was upward shifted to a level higher than that resulted from coating proteins from both sides on RBCs, as predicted by the lower CD4 density (Fig. 3f, blue).

Using Eqs. 1&2 to globally fit the green and black data (Fig. 3f) returns $A_cK_{a,TCR}$ and $k_{-1,TCR}$ values comparable to, and $K_a^*{}_{,CD4}$ and $k_{-2,CD4}$ values even larger than, those obtained by fitting Fig. 2b (Table 1). In sharp contrast, fitting the orange data (Fig. 3f) returns a vanishingly small 2D affinity of CD4 for TCR-prebound pMHC $K_a^*{}_{,CD4}$ (1.0×10^{-9} μm^2) with a 4-log larger

standard error (4.2×10^{-5} μm^2) despite reasonable values of $A_cK_{a,TCR}$, $k_{-1,TCR}$, and $k_{-2,CD4}$ (Table

1). Fitting the blue data (Fig. 3f) after constraining $k_{-1,TCR}$, and $k_{-2,CD4}$ to prevent over-fitting

generated $A_cK_{a,TCR}$ and $K_a^*{}_{,CD4}$ values similar to those obtained by globally fitting the green and

black data (Table 1). Together, these results indicate that cooperative trimolecular interaction

was modulated by the mobility of TCR and CD4 on the cell surface.

Optimal TCR-CD4 cooperativity is achieved at 7 nm spacing

7   We hypothesized that the mobility effect might stem from the requirement for sparsely

distributed TCR and CD4 to diffuse into close proximity for them to physically reach the same

pMHC. To test this hypothesis, we designed a DNA nanostructure – a ~250-nm 10-helix bundle

(10HB) DNA origami nanorod[34], which can be used as a molecular ruler for precise control of

the arrangement and relative locations of proteins[34]. 10HB nanorods functionalized with TCR

and CD4 were captured on a magnetic bead to form a surrogate T cell (Fig. 4a) to test against a

pMHC bearing RBC in the 2D kinetic assay (Fig. 4b). Many poly-A DNA strands protrude from

one side of 10HB to bind poly-T strands on the bead surface, while TCR and CD4 are anchored

by "capture" DNA strands on the other side of 10HB (Fig. 4c). The position of the capture DNA

strands was changed to adjust the TCR-CD4 intermolecular spacing, which was designed to be 6,

13, 20, and 100 nm for this study (Fig. 4d). To bind the proteins to their complementary capture

DNA strands on 10HB, CD4 was covalently linked with a DNA strand, verified by Sodium

dodecyl-sulfate polyacrylamide gel electrophoresis (SDS PAGE) (Fig. 4e). C-terminally

biotinylated E8 TCR was bound to a streptavidin (SA), which was in turn bound to a biotinylated

DNA strand on the 10HB (Fig. 4c). Successful assembly of 10HB nanorods was confirmed via

native agarose gel electrophoresis (Fig. 4f). A slight decrease in mobility was observed after the

addition of single-stranded extensions (poly-T strands and capture strands), compared to naked

10HB nanorods. The nanorods were extracted from the gel bands, purified, and used for

attachment of TCR and CD4. Subsequent transmission electron microscopy imaging on

negatively stained samples clearly showed the binding of TCR and CD4 and their 13-nm and 20-

nm designed intermolecular spacings (Fig. 4g). We should note that the designed spacing are

nominal, as the actual spacing can be affected by the length of DNA handles, the size of

streptavidin, and the flexibility of the linkage.

To examine how TCR-CD4 spacing impacted their synergistic binding to pMHC, we

measured adhesion frequencies between RBCs bearing pMHC and origami beads coated with E8

TCR, CD4, or both with defined spacing. CD4-alone origami adhered negligibly as did

protein/handle negative or origami negative controls, whereas the same density of TCR-alone

origami yielded ~20% of P_a at steady-state ($t_c = 4$ s), consistent with its >3 logs higher affinity

(Supplementary Fig. 4). Multiplexing TCR and CD4 on the origami 100 and 20 nm apart

generated indistinguishable adhesions comparable to the TCR-alone group. Strikingly, further

decreasing the spacing to 13 and 6 nm profoundly increased P_a (Supplementary Fig. 4) and its

density-normalized log-transformation (Fig. 5a), indicating strong cooperativity as in Figs. 2b

and 3f.

We extended Eq. 3 by assuming that increasing separation between TCR and CD4 would

decrease the trimolecular bond formation exponentially, obtaining a phenomenological model

that fits the data well:

$$\frac{\langle n \rangle_W}{m_{TCR}m_{pMHC}} = A_c K_{a,TCR}\left[1 + m_{CD4}K_{a,CD4}^*\exp\left(-\frac{x-x_0}{l}\right)\right] \tag{4}$$

where $A_c K_{a,TCR}$ ($1.29 \pm 0.52 \times 10^{-5}$ μm^4) was obtained from the TCR-alone measurement. Its

smaller than other values in Table 1 may be explained by presenting TCR on different surfaces

(RBCs vs origami-coated beads)[32]. Fitting returned an apparent separation x_0 (7.0 ± 1.7 nm)

required for optimal cooperativity, decay length scale l (7.7 ± 2.1 nm), and CD4 affinity for TCR

pre-engaged pMHC $K^*_{a,CD4}$ (0.050 ± 0.015 μm^2). The first value agrees well with the 7-nm gap

between the membrane anchors of TCR and CD4 found in their ternary structure complexed with

pMHC[12], while values <7 nm lead to worse fits. The second value, l describes the exponential

decay in the ability of CD4 to bind TCR-engaged pMHC, underscoring the requirement for

spatial proximity. The last value ($K^*_{a,CD4}$) is statistically indistinguishable from the value

estimated from Fig. 2b using Eqs. 1 and 2 (Table 1).

To obtain independent evidence for the existence of trimolecular bonds, we analyzed

single-bond stiffness to distinguish bimolecular vs trimolecular interactions[5] (Fig. 5b,c). When

TCR and CD4 were spaced 20 or 100 nm apart, the histograms of molecular stiffness exhibited a

single peak well fitted by a Gaussian distribution, yielding a mean stiffness of 0.32 pN/nm for

both samples (Fig. 5c). Spacing the molecules 13 nm apart gave rise to a double-peak histogram

(Fig. 5c). The first peak has nearly the same mean stiffness of 0.31 pN/nm but the second peak

represents a subpopulation (38% of total) of nearly doubling the stiffness (0.58 pN/nm),

indicating the presence of trimolecular bonds. The 6-nm spacing further populates the second

peak (51%) but maintains the mean stiffness values for the two peaks (Fig. 5c). Interestingly,

these fractions are consistent with the corresponding fractions with low and high average

numbers of whole bonds ($<n>_w$) normalized by the TCR and pMHC densities (m_{TCR} and m_{pMHC})

measured at 13 and 6 nm using the 2D assay (Fig. 5a). Overall, the data indicate that a threshold

distance is required for TCR and CD4 to form trimolecular bonds with pMHC.

Discussion

A previous study using FRAP to analyze the mobility of TCR-CD3 or CD4 at the

interface with pMHC observed reduced mobility of CD4 in the presence of TCR–pMHC

interaction and reduced TCR-CD3 mobility in the presence of CD4–MHC interaction[22],

suggesting the binding of cellular TCR and CD4 to the same pMHC. Here we showed in detail

how and why this occurs despite the extremely low CD4–MHC affinity. By demonstrating that

CD4 binds TCR-prebound pMHC with 5-6 logs higher affinity than it binds free pMHC, our

results explain how CD4 could efficiently augment TCR antigen recognition via stabilizing

TCR–pMHC bonds and recruiting Lck to TCR-CD3. This also answered a perplexing question

of why anti-CD4 blocking antibody would suppress Lck phosphorylation of CD3[35], if the CD4

could not even bind pMHC. While CD4 alone cannot initiate T cell interaction with APC, it is

able to provide adequate "help" to TCR in recognition of cognate pMHC via enormous synergy

and cooperation, a mechanism that greatly enhances TCR sensitivity.

Our work highlights the regulatory effects of force on dynamic cooperative molecular

bonds. The cooperativity of mouse CD4 and TCR binding to Hb:I-E^k could only be detected by

the BFP force-clamp assay but not by the force-free adhesion frequency assay[8], suggesting that

force on pMHC may enhance its on-rate for CD4. Our model also suggests that force-elicited

catch bond enhances the sustainability of the pMHC–CD4 arm of the trimolecular bond by

prolonging its lifetime.

An intriguing question raised by our findings is the molecular basis for the enhanced

affinity of CD4 for TCR-prebound pMHC. One possibility is that TCR binding to pMHC

induces conformational changes in the CD4 binding site of MHC that increase affinity for CD4.

However, crystallographic studies of numerous TCR and pMHC molecules in free and bound

form provide no evidence of TCR-induced structural changes at the site on the MHC class II β2

domain where CD4 binds. This would argue against an allosteric mechanism to explain increased

CD4 affinity for TCR-prebound MHC. However, it is also possible that the relevant changes may

be in protein dynamics, a parameter that cannot be addressed via the static snapshots provided by

X-ray crystallography. Indeed, several NMR studies have found that pMHC binding to TCR

induces long-range allosteric changes in TCR dynamics that could be relayed to CD3, even in the

absence of obvious structural changes in the TCR[36-38]. However, the reverse situation (i.e. long-

range allosteric changes in pMHC dynamics upon TCR binding) has not been examined by

NMR, but probably should be given our results here.

The proposed structural mechanisms are supported by our modeling analysis, which was

used to fit the bond lifetime data measured by the BFP experiments (Figs. 2d-f, Supplementary

Fig. 3). To show this, we first compared the abilities of three models to simultaneously fit the

three specific interaction data measured by thermal fluctuation (Fig. 2d). The first model is

depicted in Supplementary Fig. 3a except that the activation to and dissociation from the strong

state (TCR–pMHC)[*] is neglected because force was zero. Also, the weak (TCR–pMHC–CD4)

and strong (TCR–pMHC–CD4)[*] trimolecular bonds are assumed to be in equilibrium and the

two states dissociate concurrently to free molecules TCR + pMHC + CD4 in a single step (hence

called concurrent model). The second model is a single-pathway, two-step, sequential

dissociation model (sequential model I). It differs from the concurrent model in that the TCR–

pMHC–CD4 trimolecular bond has only a single state that dissociates sequentially from right to

left in Eq. 9 (Methods), first the pMHC–CD4 arm reversibly to generate TCR–pMHC complex +

CD4, and then the TCR–pMHC bimolecular bond to free molecules TCR + pMHC + CD4. The

third model is also a single-pathway, two-step, sequential dissociation model (sequential model

II). It is similar to the sequential model I except that the order of sequential dissociation of the

TCR–pMHC–CD4 trimolecular bond is reversed, which occurs first at the TCR–pMHC arm

reversibly to generate pMHC–CD4 complex + TCR, and then at the pMHC–CD4 bimolecular

bond to generate free molecules TCR + pMHC + CD4. Using the Akaike Information Criterion

(AIC) to choose between two non-nested models, we found that the concurrent model is more

likely to be correct than sequential model I by >58,000 times and even much more times than

sequential model II. This conclusion still holds even when we analyzed data measured by force-

clamp assay (Supplementary Fig. 3b) to evaluate their force-dependent kinetic parameters

(Supplementary Fig. 3c,d). The concurrent model suggests a strong functional coupling between

the spatially separate TCR binding site and the CD4 binding site on the pMHC, supporting our

allostery hypothesis. Future studies are required to test this hypothesis. Moreover, to confirm that

association and dissociation happen preferentially via pathway I instead of pathway II, we

considered a more general two-pathway, two-step model that combines both sequential models I

and II, and globally fit the adhesion frequency data from Fig. 2a-b. We compared it to the

sequential model I by AIC, and found little difference statistically, suggesting that both

association and dissociation along the first pathway (i.e., sequential model I) dominates over the

second pathway (i.e., sequential model II), which only happens with very low probability (~0.99

and ~0.01 for pathway I and two-pathway, two-step model respectively).

This work builds upon our previous findings of TCR-CD8 cooperativity for class-I

pMHC binding[3-5,7,20] and adds to our previously proposed mechanistic model[5,39]. In live-cell

systems, cooperativity relies on the recruitment of coreceptor to phosphorylated CD3 via a Lck

"bridge", where both the adaptor function and kinase function of Lck are required[3,5,20,39,40] as

seen with co-immunoprecipitation studies that identify similar molecular assembly connecting

CD4 or CD8 to triggered TCR/CD3[15,41-44]. Here we showed that using purified TCR and CD4

lacking cytoplasmic domains and CD3, the synergistic cooperation could still occur if both

molecules are allowed to diffuse on the RBCs or packed closed enough on the bead surface,

suggesting that modulation of ectodomain cooperation could be decoupled from intracellular regulation. This is also consistent with the observation that CD4 lacking the ability of binding Lck is still required for TCR-mediated IL-2 production[22]. On the other hand, our data of cellular TCR and CD4 cooperating with a faster kinetics than their purified ectodomain counterparts suggest that there may be cellular regulation(s) to expedite TCR-CD4 cooperativity. Modulation of ectodomain by physical factors such as mobility and colocalization are intuitive, but require 2D analyses to decipher. Indeed, by limiting mobility using different surface and controlling TCR-CD4 proximity using DNA origami, we found a 7-nm apparent spacing for TCR-CD4 cooperativity, such that increasing the spacing resulted in an exponential decay in the capacity to form TCR–pMHC–CD4 trimolecular bonds. Such physical regulatory mechanisms revealed by the in vitro system further advance our understanding of how interactions between membrane receptors and ligands critically depend on their cell surface organization. Technically, our DNA origami-based 2D kinetic assays hold great potential to dissect the spatial requirement of the cooperation of single- or multi-receptor–ligand species by ectodomain binding and/or intracellular crosstalk.

Methods

Proteins

Biotinylated CD4 and biotinylated αβ E8 TCR were prepared as previously described[15,19].

Briefly, a 17-amino acid tag (TPI, GGGLNDIFEAQKIEWHE) was added to the C-terminus of

CD4 ectodomain and to the C-terminus of the E8 TCRα ectodomain. CD4 was expressed in

baculovirus-infected Sf9 insect cells[15]. E8 TCR was produced by *in vitro* folding from inclusion

bodies expressed in *E. coli*[19]. The purified proteins were biotinylated using biotin protein ligase

(Avidity); excess biotin and ligase were removed with a Superdex 200 column (GE Healthcare).

The control MHC-I (HLA-A2) presenting the melanoma antigen peptide NY-ESO-1$_{157-165}$ (9V,

SLLMWITQV) was produced by the NIH Tetramer Facility at Emory University.

Biotinylated TPI: HLA-DR1 was made by the NIH Tetramer Core facility (Atlanta, GA)

with peptide (GELIGTLNAAKVPAD) purchased from GenScript. Divalent streptavidin was

generously gifted by Dr. Baoyu Liu (University of Utah).

Cells

Human RBCs were isolated from blood of healthy donors according to a protocol

approved by the Institutional Review Board of Georgia Institute of Technology. For adhesion

frequency assay, RBCs were purified by Histopaque-1077, washed with ice cold PBS, and

resuspended in EAS-45 buffer (2 mM Adenine, 110 mM D-glucose, 55 mM D-Mannitol, 50 mM

Sodium Chloride, 20 mM Sodium Phosphate, and 10 mM L-glutamine). Equal aliquots of RBCs

were then mixed with various concentrations of EZ-Link Sulfo-NHS-LC-Biotin (Thermo

Scientific) at a pH of 7.2 for 30 min at room temperature, yielding different densities of biotin

sites on RBC surfaces. Biotinylated RBCs were washed with EAS-45 buffer and stored at 4°C.

For BFP experiments, freshly isolated human RBCs were biotinylated with biotin-PEG3500-

NHS (Jenkem Technology) and then incubated with nystatin in N2 buffer (265.2 mM KCl, 38.8 mM NaCl, 0.94 mM KH2PO4, 4.74 mM Na2HPO4, and 27 mM sucrose; pH 7.2 at 588 mOsm) for 30 min on ice. Nystatin-treated biotinylated RBCs were washed twice with N2 buffer and stored at 4°C for BFP experiments.

HEK 293T cells were purchased from ATCC (Manassas, VA) and cultured in DMEM supplemented with 10% FBS, 6 mM L-glutamine, 0.1 mM MEM non-essential amino acids, and 1 mM sodium pyruvate. TCR β-chain deficient Jurkat J.RT3 cells were purchased from ATCC (Manassas, VA) and cultured in RPMI 1640 supplemented with 10% FBS, 100 U/mL penicillin, 100 μg/mL streptomycin, 2 mM L-glutamine, and 20 mM HEPES. J.RT3 were transduced by lentivirus to express the E8 TCR. Briefly, E8 TCR α and β chains joined by a P2A element were subcloned into pLenti6.3 vector with a T2A-rat CD2 reporter. Lentivirus encoding E8 TCR were produced by co-transfection of HEK 293T cells with E8TCR-pLenti6.3, pMD2.G (Addgene #12259), and psPAX2 (Addgene #12260) using lipofectamine 3000 (ThermoFisher Scientific). J.RT3 cells were transduced by incubating overnight with supernatant containing lentivirus and FACS sorted using Aria cell sorter (BD Biosciences) based on the surface expression E8 TCR. E8TCR J.RT3 cells were then transfected using Nucleofection (Lonza, Morristown, NJ) to express full-length human CD4 and FACS sorted using Aria cell sorter (BD Biosciences) for stable CD4 expression.

Site density measurements:

Site densities of TCR, CD4, and pMHC on RBCs or beads were measured by flow cytometry using PE-conjugated antibodies from BD Biosciences (San Jose, CA): anti-human TCR-β1 (clone JOVI.1, 1:20), anti-human CD4 (clone OKT4, 1:20), and anti-human HLA-DR (clone L243, 1:5). Isotype control antibodies were PE mouse IgG2a κ (clone G155-178, 1:20) for

TCR and HLA-DR, and PE mouse IgG2b, κ (Clone 27-35, 1:20) for CD4. RBCs or beads were

incubated with corresponding antibodies in 1X PBS + 2% BSA for 30 min at 25 °C and washed

three times before being analyzed using the BD Accuri Flow Cytometer. Site density was

calculated based on antibody-conjugated PE fluorescence relative to the QuantiBRITE PE

standard beads (BD Biosciences). For micropipette experiments, RBCs bearing protein of

interest (TCR, CD4, or MHC) were prepared for flow cytometry using three separate samples:

one saturating the RBC surface, one of the experimental conditions, and one isotype control.

Brightness across all antibodies was normalized against the CD4 staining which was brighter

than the TCR staining.

Micropipette Adhesion Frequency Assay

The theoretical framework and detailed procedures of this assay have been reported

previously[23,24,45]. This assay leverages the ultra-soft RBC membrane to detect interactions with

single-bond sensitivity. Briefly, RBCs with controlled biotin sites were first incubated with 200

µg/mL divalent streptavidin and washed, followed by subsequent coating with biotinylated

pMHC, TCR, CD4, or a mixture of TCR and CD4. After washing, RBCs coated with receptor-

ligand pairs were injected into an imaging chamber containing non-CO2 dependent L15 media

supplemented with HEPES and 2% BSA. During experiment, two RBCs were aspirated by

micropipette and repeatedly brought into contact of defined duration by a program-controlled

piezo. Adhesion between two RBCs were detected by the deflection of RBC membrane during

the separation after each contact with '1' indicating an adhesive event and '0' indicating no

adhesion. Each pair of RBCs were tested for 50 cycles of the same contact duration to yield the

adhesion frequency (P_a), which is as the average score from the 50 cycles. The same process was

repeated for multiple pairs of RBC of each contact duration (t_c), yielding a P_a vs t_c curve that is

fitted to a published model[23,24] (see below).

Non-specific controls were conducted for all sets of species coatings. To do so, RBCs

coated with divalent streptavidin anchored irrelevant protein (OT1 TCR) were brought into

contact with RBCs coated with proteins of interest. An adhesion frequency curve would be

obtained for nonspecific adhesion. Then, for each contact time, the distribution of nonspecific

adhesion frequency would be removed from the measured distribution of adhesion frequency[27-

29].

Biomembrane Force Probe Assay

The principle and experimental procedures for BFP assays have been described

previously[11]. To coat proteins on beads, borosilicate glass beads were mercapto-propyl silanated

and covalently functionalized to monovalent streptavidin-maleimide in phosphate buffer saline

(pH 6.8) by overnight incubation at 25°C. Streptavidinylated beads were then incubated with

sub-saturating concentrations of biotinylated pMHC, TCR, CD4, or 1:1 mixture of TCR and

CD4 for 2 hours in HBSS + 2% BSA at 4°C. During the experiment, a protein-coated bead

(tracking bead) was attached to the apex of a micropipette-aspirated RBC that serves as a spring

with pre-adjusted spring constant of 0.1-0.3 pN/nm. For force clamp assay, the tracking bead was

repeatedly contacted by a piezo-driven target bead coated with (or target cell expressing) the

corresponding binding partner(s). The displacement of tracking bead was monitored with a high-

speed camera at 1000 fps and was translated into force by applying the pre-defined spring

constant. During separation, bond formation between tracking bead and target bead/cell pulled

the tracking bead away from its baseline, manifesting positive force loading on the molecular

bond. Bond lifetime was defined as the duration from the start of clamp at the preset force level

to bond rupture. Several hundred bond lifetime events were collected and pooled for various

clamp forces using multiple bead-bead or bead-cell pairs. The stiffnesses of molecular bonds

were modeled as Hookean spring constants which is determined by the difference between the

displacements of the bead tracking system (F) and the displacements of the piezoelectric

actuator-capacitance sensor feedback system (Δx). Two straight lines were fitted to the piecewise

data: One was fitted to the compressive force regime where the bead was slowly ramping away

from cell impingement (force < 0), where the slop represents the spring constant of the cell

membrane (compression). The other was fitted to the tensile force regime where the bead would

ramp away from the cell beyond the point of initial contact (force > 0), where the slop represents

the resultant spring constant of cell membrane (extension) and molecular bond connected in

series. By assuming the spring constant of cell membrane is identical during compression as

during extension, the spring constant of molecular bond was extracted by applying Hooke's law

for springs in series. An ensemble of molecular spring constant was collected as histograms

using similar force bins and fitted to single or double Gaussian distributions.

For thermal fluctuation assays[24,30], a target bead was driven by piezo to the proximity

(~10 nm) of the probe bead, allowing bond formation between the two during intermittent

contacts caused by thermal fluctuation, which would decrease the fluctuation of probe bead

position. By analyzing the standard deviation (SD) of the tracking bead position, bond formation

and rupture were identified by the decrease of SD below a cutoff of 3 nm and the subsequent

resumption SD to baseline level. The cutoff baseline was determined as the lowest SD value of

the probe bead before contact (*i.e.*, SD of bead without bonds).

For adhesion frequency assays using BFP beads, the same process of RBC-based

adhesion frequency assay was performed except that a target cell was repeatedly brought into

1   contact with a pMHC-coated bead, held for a defined duration, and then separated. Adhesion was

2   defined as the deflection of bead away from its equilibrium position.

3   **Kinetic models**

4   Single-step model for bimolecular interaction- The bimolecular interactions of either TCR or

5   CD4 with pMHC are modeled as a single-step reversible reaction:

$$R + L \underset{k_{-1}}{\overset{k_{+1}}{\rightleftharpoons}} B \tag{5}$$

7   where R, L, and B denote, respectively, receptor, ligand, and bond; k_{+1} and k_{-1} denote,

8   respectively, on- and off-rates. Their ratio is the binding affinity, $K_a = k_{+1}/k_{-1}$. Our published

9   probabilistic kinetic model for the above reaction states that the probability of adhesion (P_a)

10 depends on the contact time (t_c) and area (A_c) as well as the densities of receptors (m_r) and

11 ligands (m_l)[23,24].

$$P_a = 1 - \exp\{-m_r m_l A_c K_a [1 - \exp(-k_{-1} t_c)]\} \tag{6}$$

13 On the right-hand side, the term inside the braces after the minus sign is the average number of

14 bonds per contact $<n>$.

$$\langle n \rangle = m_r m_l A_c K_a [1 - \exp(-k_{-1} t_c)] \tag{7}$$

16 After normalizing $<n>$ by $m_r m_l$, the right-hand should be independent of the molecular densities

17 and approaches $A_c K_a$, the effective 2D affinity, as t_c becomes large.

$$\frac{\langle n \rangle}{m_r m_l} = A_c K_a [1 - \exp(-k_{-1} t_c)] \rightarrow A_c K_a \qquad \text{as } t_c \gg 1/k_{-1} \tag{8}$$

It follows from Eq. 8 that $<n>$ is proportional to $m_r m_l$ if measured at large contact time.

We use TCR or CD4 in the subscript to identify the interaction with which the kinetic

parameters are associated, e.g., $k_{+1,TCR}$, $k_{-1,TCR}$, and $K_{a,TCR}$ are kinetic rates and binding affinity

for the TCR–pMHC interaction, which are different from $k_{+1,CD4}$, $k_{-1,CD4}$, and $K_{a,CD4}$, which are

corresponding parameters for the CD4–MHC interaction. Also, m_{TCR}, m_{CD4}, and m_{pMHC} are used

to designate the densities of TCR, CD4, and pMHC.

<u>Two-step model for trimolecular interaction-</u> We propose a two-step model for the

formation of trimolecular bonds. Since the affinity and on-rate for the CD4–pMHC interaction

are so much smaller than those of the TCR–pMHC interaction, it seems reasonable to assume

that TCR interacts with pMHC at the same kinetic rates as bimolecular interaction in the first

step as if CD4 does not interact with free pMHC. However, CD4 is able to interact with TCR-

stabilized pMHC with a much higher affinity, thereby forming trimolecular bonds in the second

step.

10
$$\text{TCR} + \text{pMHC} + \text{CD4} \underset{k_{-1,\text{TCR}}}{\overset{k_{+1,\text{TCR}}}{\rightleftharpoons}} \text{TCR–pMHC} + \text{CD4} \underset{k_{-2,\text{CD4}}}{\overset{k_{+2,\text{CD4}}}{\rightleftharpoons}} \text{TCR–pMHC–CD4} \quad (9)$$

11   Let $p_{m,n}$ be the probability of having m TCR–pMHC bonds and n TCR–pMHC–CD4 bonds,

12   which are governed by the following master equations:

13
$$\frac{dp_{m,n}}{dt} = m_{\text{TCR}} m_{\text{pMHC}} A_c k_{+1,\text{TCR}} p_{m-1,n} + k_{-1,\text{TCR}}(m+1) p_{m+1,n}$$

14
$$+ m_{\text{CD4}} k_{+2,\text{CD4}}(m+1) p_{m+1,n-1} + k_{-2,\text{CD4}}(n+1) p_{m-1,n+1}$$

15
$$- [m_{\text{TCR}} m_{\text{pMHC}} A_c k_{+1,\text{TCR}} + (k_{-1,\text{TCR}} + m_{\text{CD4}} k_{+2,\text{CD4}})m + k_{-2,\text{CD4}} n] p_{m,n} \quad (10)$$

16   We solved these master equations using a probability generating function,

17
$$g(x,y\,;t) \equiv \sum_{m,n=0}^{\infty} x^m y^n p_{m,n}(t), \quad (11)$$

18   which converts Eq. 10 to a single first-order linear partial differential equation,

19
$$\frac{\partial g}{\partial t} + \left[\left(k_{-1,\text{TCR}} + m_{\text{CD4}} k_{+2,\text{CD4}}\right)x - m_{\text{CD4}} k_{+2,\text{CD4}} y - k_{-1,\text{TCR}} \right] \frac{\partial g}{\partial x}$$

20
$$+ k_{-2,\text{CD4}}(y-x)\frac{\partial g}{\partial y} + m_{\text{TCR}} m_{\text{pMHC}} A_c k_{+1,\text{TCR}}(1-x)g = 0. \quad (12)$$

We found a general solution to Eq. 12 using the method of characteristics, which is

$g = J(u_1 e^{-\lambda_1 t}, u_2 e^{-\lambda_2 t}) \exp\{ m_{\text{TCR}} m_{\text{pMHC}} A_c K_{a,\text{TCR}}[x - y + (1 + m_{\text{CD4}} K^*_{a,\text{CD4}})(y - 1)]\}$ (13)

where J is an arbitrary function of its two arguments. u_1 and u_2 are functions of x and y:

$$u_{1,2}(x, y) = \pm[\lambda_{2,1}(1 - y) + k_{-2,\text{CD4}}(y - x)]/(\lambda_1 - \lambda_2), \qquad (14)$$

λ_1 and λ_2 are given by Eq. 2. They can be viewed as the fast (λ_1) and slow (λ_2) rates that control

the two p44hases of the two-step interaction. To determine the function of integration J requires

initial conditions. If there is no TCR–pMHC or TCR–pMHC–CD4 bonds at $t = 0$, the initial

conditions on $p_{\text{m,n}}$ are:

$$p_{\text{m,n}}(0) = \begin{cases} 1, & m = n = 0 \\ 0, & m, n > 0 \end{cases}. \qquad (15)$$

The corresponding initial condition of g, obtained by substituting Eq. 15 into Eq. 11, is

$$g_{00}(x, y; 0) = 1. \qquad (16)$$

Solving x and y in terms of $u_{1,2}$ from Eq. 14, substituting them into Eq. 16, and comparing the

resulting equation to Eq. 13 evaluated at $t = 0$ allows us to solve for a particular solution for J:

$J_{00}(u_1, u_2) = \exp\{ m_{\text{TCR}} m_{\text{pMHC}} A_c K_{a,\text{TCR}}[(\lambda_1/k_{-2,\text{CD4}})u_1 + (\lambda_2/k_{-2,\text{CD4}})u_2$

$$- (1 + m_{\text{CD4}} K^*_{a,\text{CD4}})(u_1 + u_2)]\}. \quad (17)$$

Substituting Eq. 17 into Eq. 13, we have

$g_{00}(x, y; t) = \exp\Big(m_{\text{TCR}} m_{\text{pMHC}} A_c K_{a,\text{TCR}}\{ k_{-1,\text{TCR}} \dfrac{e^{-\lambda_1 t} - e^{-\lambda_2 t}}{\lambda_1 - \lambda_2}$

$-\Big(1 + m_{\text{CD4}} K^*_{a,\text{CD4}}\Big)\Big(1 + \dfrac{\lambda_2 e^{-\lambda_1 t} - \lambda_1 e^{-\lambda_2 t}}{\lambda_1 - \lambda_2}\Big)$

$+[1 + \dfrac{(\lambda_2 - k_{-1,\text{TCR}})e^{-\lambda_1 t} - (\lambda_1 - k_{-1,\text{TCR}})e^{-\lambda_2 t}}{\lambda_1 - \lambda_2}]x + m_{\text{CD4}} K^*_{a,\text{CD4}}(1 + \dfrac{\lambda_2 e^{-\lambda_1 t} - \lambda_1 e^{-\lambda_2 t}}{\lambda_1 - \lambda_2})y\}\Big)$ (18)

Expanding the right-hand side of Eq. 18 into Taylor series of x and y, and comparing the

coefficients of the $x^m y^n$ terms to those of the right-hand side of Eq. 11, we found

$$p_{\text{m,n}}(t) = \left[\frac{\langle m \rangle^m}{m!} \exp(-\langle m \rangle)\right]\left[\frac{\langle n \rangle^n}{n!} \exp(-\langle n \rangle)\right] \qquad (19a)$$

where

$$\langle m \rangle = \langle n \rangle_{\text{TCR}} = m_{\text{TCR}} m_{\text{pMHC}} A_c K_{a,\text{TCR}} [1 + \frac{(\lambda_2 - k_{-1,\text{TCR}})e^{-\lambda_1 t} - (\lambda_1 - k_{-1,\text{TCR}})e^{-\lambda_2 t}}{\lambda_1 - \lambda_2}] \quad \text{(19b)}$$

$$\langle n \rangle = \langle n \rangle_{\text{tri}} = m_{\text{TCR}} m_{\text{pMHC}} A_c K_{a,\text{TCR}} m_{\text{CD4}} K_{a,\text{CD4}}^* \left(1 + \frac{\lambda_2 e^{-\lambda_1 t} - \lambda_1 e^{-\lambda_2 t}}{\lambda_1 - \lambda_2} \right) \quad \text{(19c)}$$

are the average numbers of TCR–pMHC bonds and TCR–pMHC–CD4 bonds. Thus, the solution

is of the form of the product of two Poisson distributions. Except for the expressions for <m>

and <n>, this is the same as the bond distribution of the dual species concurrent independent

binding model described previously[27-29].

The adhesion probability is the probability of having at least one bond of either type, or

one minus the probability of having no bond, which can be found from Eq. 19:

$$P_a(t) = 1 - \exp\{ - m_{\text{TCR}} m_{\text{pMHC}} A_c K_{a,\text{TCR}} \left(1 + m_{\text{CD4}} K_{a,\text{CD4}}^* \right)$$

$$\times [1 + \frac{\lambda_2(\lambda_2 - k_{-1,\text{TCR}})e^{-\lambda_1 t} - \lambda_1(\lambda_1 - k_{-1,\text{TCR}})e^{-\lambda_2 t}}{(\lambda_1 - \lambda_2)(m_{\text{CD4}} k_{+2,\text{CD4}} + k_{-2,\text{CD4}})}]\} \quad \text{(20)}$$

Taking a log transformation, we obtain the average number of whole bonds $\langle n \rangle_W =$

$-\ln(1 - P_a)$. Normalizing $\langle n \rangle_W$ by the densities of TCR and pMHC, we obtain Eq. 1. Setting

$k_{+2,\text{CD4}} = k_{-2,\text{CD4}} = 0$ reduces Eq. 2 to $\lambda_1 = k_{-1}$ and $\lambda_2 = 0$, which also reduces Eqs. 20 and 1 to Eqs.

6 and 8, respectively, as expected.

DNA Origami preparation

Origami were created at 20 nM using p8064 scaffold, 5× staple excess, and 25× biotin

strand for TCR capture. Following agarose gel electrophoresis in 1% agarose at 60 V for 90 min

on ice bath, origami were purified with Freeze 'n' Squeeze spin columns. The four samples with

different protein separation distances were indistinguishable based on gel mobility. However, a

control sample with only handles for surrogate cell attachment ran with slightly increased

mobility, while a complete blank 10 HB sample with no handle extensions whatsoever ran markedly faster through the gel. To confirm proper assembly of protein onto the DNA origami, purified samples were incubated with excesses of CD4–DNA and wild type streptavidin at 4 °C overnight. After another agarose gel purification, a 10-fold molar excess of biotinylated TCR was added and allowed to incubate at 4 °C for 3 h. Excess protein was removed using 100 kD MWCO spin filters, then the samples were deposited on 400 mesh carbon formvar grids and stained with 1% uranyl formate solution for 30 s. During TEM imaging, the identity of blinded samples could be reliably divined by observing relative CD4, SA–TCR location.

An amine-modified DNA strand was conjugated to lysine residues in CD4. CD4 was reacted with 50-fold excess succinimidyl-6-hydrazino-nicotinamide (S-HyNic) (Solulink) for 4 h, and terminal amine DNA was reacted with 20-fold excess succinimidyl-4-formylbenzene (S-4FB) (Solulink) for 4 h. Amicon Ultra spin filters (Millipore) were used to remove excess linker and buffer exchange into citrate buffer (50nM sodium citrate, 150 mM NaCl, pH 6). Functionalized DNA was combined with CD4 at a 3:1 ratio, and reacted overnight. CD4-DNA conjugates were purified on a Superdex 200 Increase 10/300 GL column using an AKTA Pure FPLC (GE Healthcare). Conjugation was verified with SDS/PAGE and Coomassie blue staining.

In order to examine the effect of geometric docking on the simultaneous interaction of two molecules with a third, origami structures differentially spacing TCR and CD4 were coated on beads to represent a simplified T cell surface. Beads were respectively coated with several different origami structures: those with handles for TCR and CD4 which were differentially spaced apart at 6, 13, 20, and 100 nm; those lacking protein handles; and origami lacking connecting strands to the beads. The purified samples were loaded onto surrogate T cells – 3 μm diameter magnetic dt Dynabeads. For each sample, 15 μL of beads were washed 3× with PBS

then added to 1 µg (~30 µL) of purified DNA origami. A large excess of CD4–DNA and streptavidin were added, and the samples were left to incubate on a rotating shaker at 4 ℃ overnight. The magnetically pelleted beads were washed with PBS supplemented with calcium and magnesium, and 2% BSA five times. Biotinylated (at the C-terminus of the α-chain) E8 TCR was added to the beads and incubated for 30 min. To this solution, blocking DNA strand (25 dA) was added and the reaction incubated at 4 °C for an additional 1 h.

Fitting models to data

Linear fittings of data in Fig. 2d, Fig. 5b, Supplementary Fig. 1, and Supplementary Fig. 2c were implemented in Prism using linear regression to estimate the slope and y-intercept of the trend. Non-linear fittings of bimolecular kinetic equations to data in Figs. 1c,d were implemented in Prism using Non-linear regression to estimate the corresponding kinetic parameters. Gaussian fitting of data in Fig. 5c was implemented in Prism using non-linear regression to estimate the Gaussian distribution and frequency of each distribution. For all fitted parameters, fitting error was presented as Standard Error (SE) assuming symmetrical confidence intervals.

The model predicted curves in Figs. 2a-c and Supplementary Fig. 2a were obtained by globally fitting all data to Eqs. 1&2, which were also fit to data in Fig. 3a,b,c,f. For the Fig. 3f case, the green and black data were globally fit, the orange data was fit on its own, and the blue data was fit after constraining the off-rates obtained from the orange data fit to avoid overfitting. The separation distance-dependent adhesion frequency of the DNA origami assay (Fig. 5a) was fit to Eq. 4. For Figs. 2a-c, 3a,b,c,f , and 5a, a two-step optimization procedure was employed to minimize the mean squared error (MSE) of the fits with respect to the data, which explored a large parameter space by Differential Evolution to obtain an approximate solution, followed by

L-BFGS to hone in the minimum. For Figs. 2a-c and 3a,b,c,f the parameters Standard Errors

(SE) were calculated from the covariance matrix obtained by inverting the Hessian matrix of the

loss function at the minimum found. The SEs were then propagated through the model to display

error bands of the fits as mean ± SE. For Fig. 5a, the parameters SE were obtained via

bootstrapping, with 1000 bootstrap samples.

The force-dependent lifetimes of pMHC bonds with TCR alone or co-presented TCR and

CD4 behaved as catch-slip bonds, while pMHC bonds with CD4 alone behaved as a slip bond.

Although at zero force the two catch-slip bonds had mono-exponentially decaying survival

probabilities (Fig. 2d), a second population appeared at forces >0, displaying biexponentially

decaying survival probabilities, suggesting the presence of two bound species, even for the

bimolecular TCR–pMHC. Therefore, while the CD4–pMHC bonds were modelled to dissociate

along a single pathway, both TCR–pMHC and TCR–pMHC–CD4 bonds were modelled to

dissociate along two pathways – a fast dissociating pathway from a weak state or a slow

dissociating pathway from a strong state (indicated by *) allowed to exchange from each other as

depicted in Supplementary Fig. 3a (left and right panels) similarly to previous catch bond models

[46,47]. The force dependence of all kinetic rates follows Bell's equation[48]:

$$k_{i,j}(f) = k_{i,j}^0 \exp\left(\frac{f \cdot x_{i,j}}{k_B T}\right) \tag{21}$$

where f is force, the superscript 0 denotes the zero-force kinetic rate constant, the subscript i

denotes rates of dissociation along fast (-f) or slow (-s) pathways and of activation (a) or

deactivation (-a) transition between the weak and strong states, the subscript j denotes different

bonds (TCR–pMHC or TCR–pMHC–CD4), $x_{i,j}$ denotes distance to the transition state, or width

of the energy well that traps the system in the bound state, k_B is Boltzmann constant, and T is

absolute temperature. The weak state dissociates more rapidly than the strong state, so $k_{-f,i} > k_{-s,i}$.

1   The two-states, two-path catch-slip model can be expressed in terms of ordinary differential

2   equations that govern the probability of having bonds P at a given time t:

3
$$\begin{cases} \frac{dP_j}{dt} = -k_{-f}P_j - k_aP_j + k_{-a}P_{(j)^*} \\ \frac{dP_{(j)^*}}{dt} = -k_{-s}P_{(j)^*} + k_aP_j - k_{-a}P_{(j)^*} \end{cases} \qquad (22)$$

where the subscript j, as above, stands for TCR–pMHC or TCR–pMHC–CD4 bonds in the weak

(without *) or strong (with *) state, and all rates k_i vary as a function of force as given by Eq. 21.

This two-state catch-bond model could recapitulate the average bond lifetime as well as the

biexponentially distributed bond survival probabilities of both TCR–pMHC and TCR–pMHC–

CD4 interactions. All kinetic parameters were determined by globally fitting the models to the

force-lifetime distributions as a whole via maximum likelihood estimation (MLE). The initial

proportions of the two states a $t = 0$ were determined from the ratios of the kinetic rates assuming

equilibrium. A two-step optimization procedure was employed to minimize the MLE, which

explored a large parameter space by Differential Evolution to obtain an approximate solution,

followed by L-BFGS to home in the minimum. This returns the best-fit parameters summarized

in Supplementary Fig. 3d. The force dependence of the kinetic rates is graphically depicted in

Supplementary Fig. 3c. Using Eqs. 21&22 with the best-fit parameters we calculated the

probability of the strong state and its fraction as functions of time and force for both TCR–

pMHC and TCR–pMHC–CD4 interactions (Supplementary Fig. 3e-f). The same equations and

best-fit parameters predict the mean ± SE of lifetime vs force of the TCR–pMHC, pMHC–CD4

and TCR–pMHC–CD4 bonds that agree well with the mean ± SEM data of these interactions

(Fig. 2e). Also predicted are the average lifetime vs force of weak state, strong state, and their

sum of the trimolecular bond that shows the force enhancement of cooperativity (Fig. 2f). It is

important to note that although the two-state, two-path model fits the data as a catch-slip bond,

this does not need to always be the case. In fact, the two-state, two-path model could in theory

also yield slip bond or slip-catch-slip bond behavior given specific sets of parameters.

Nevertheless, the catch-slip behavior naturally arises from maximizing the likelihood of the

TCR–pMHC and TCR–pMHC–CD4 force-lifetime distributions, in good agreement with the

average lifetimes of each bin obtained independently from the fitting. This further adds

confidence both in the existence of the two catch-bonds as well as the reliability of the model.

The standard error for each parameter was obtained by inverting the Hessian matrix of the

negative log likelihood at the MLE to obtain the approximate covariance matrix whose diagonal

entries correspond to the variance of the fitted parameters. All kinetic modeling and associated

statistics were programmatically implemented in Julia and Python.

Data Availability

Data supporting the findings of this study are presented in the article and supplementary

materials and are available from the corresponding authors upon reasonable request.

Code Availability

The code used to fit the data presented in this study is available at

https://github.com/vaglino/TCR-MHC-CD4_kinetic_modeling under MIT license.

Acknowledgements

We thank the NIH Tetramer Facility at Emory University for providing the pMHC

molecules. This work was supported by NIH grants R21AI135753 (to Y.K and C.Z.),

R01GM124489, U01CA214354, and U01CA250040 (to C.Z.), and R01AI129893 (to R.A.M.).

Author Contributions

M.N.R., V.P., C.Z., and Y.K. conceived and designed the study; C.Z. and Y.K. directed

the study; M.N.R., V.P., K.L., H.-K.C., J.H., F.G., and P.A. performed experiments; C.Z. and

S.T. developed mathematical models; S.T. analyzed data using mathematical model; R.A.M.

provided critical reagents; M.N.R., V.P., K.L. Y.K., and C.Z. wrote the paper with contributions

from other authors.

Statement of competing interests

The authors declare no competing interests.

References

Doyle, C. & Strominger, J. L. Interaction between CD4 and class II MHC molecules mediates cell adhesion. *Nature* **330**, 256-259, doi:10.1038/330256a0 (1987).

Norment, A. M., Salter, R. D., Parham, P., Engelhard, V. H. & Littman, D. R. Cell-cell adhesion mediated by CD8 and MHC class I molecules. *Nature* **336**, 79-81, doi:10.1038/336079a0 (1988).

Jiang, N. *et al.* Two-stage cooperative T cell receptor-peptide major histocompatibility complex-CD8 trimolecular interactions amplify antigen discrimination. *Immunity* **34**, 13-23, doi:10.1016/j.immuni.2010.12.017 (2011).

Liu, B. *et al.* 2D TCR-pMHC-CD8 kinetics determines T-cell responses in a self-antigen-specific TCR system. *European journal of immunology* **44**, 239-250, doi:10.1002/eji.201343774 (2014).

Hong, J. *et al.* A TCR mechanotransduction signaling loop induces negative selection in the thymus. *Nature immunology* **19**, 1379-1390, doi:10.1038/s41590-018-0259-z (2018).

Stepanek, O. *et al.* Coreceptor Scanning by the T Cell Receptor Provides a Mechanism for T Cell Tolerance. *Cell* **159**, 333-345, doi:10.1016/j.cell.2014.08.042 (2014).

Jonsson, P. *et al.* Remarkably low affinity of CD4/peptide-major histocompatibility complex class II protein interactions. *P Natl Acad Sci USA* **113**, 5682-5687, doi:10.1073/pnas.1513918113 (2016).

Hong, J. *et al.* Force-Regulated In Situ TCR-Peptide-Bound MHC Class II Kinetics Determine Functions of CD4+ T Cells. *Journal of immunology* **195**, 3557-3564, doi:10.4049/jimmunol.1501407 (2015).

Mørch, A. M., Bálint, Š., Santos, A. M., Davis, S. J. & Dustin, M. L. Coreceptors and TCR Signaling – the Strong and the Weak of It. *Frontiers in Cell and Developmental Biology* **8**, doi:10.3389/fcell.2020.597627 (2020).

Xiong, Y., Kern, P., Chang, H. & Reinherz, E. T Cell Receptor Binding to a pMHCII Ligand Is Kinetically Distinct from and Independent of CD4. *The Journal of biological chemistry* **276**, 5659-5667, doi:10.1074/jbc.M009580200 (2001).

Liu, B., Chen, W., Evavold, B. D. & Zhu, C. Accumulation of Dynamic Catch Bonds between TCR and Agonist Peptide-MHC Triggers T Cell Signaling. *Cell* **157**, 357-368, doi:10.1016/j.cell.2014.02.053 (2014).

Yin, Y., Wang, X. X. & Mariuzza, R. A. Crystal structure of a complete ternary complex of T-cell receptor, peptide-MHC, and CD4. *Proceedings of the National Academy of Sciences* **109**, 5405-5410, doi:10.1073/pnas.1118801109 (2012).

Courtney, A. H., Lo, W. L. & Weiss, A. TCR Signaling: Mechanisms of Initiation and Propagation. *Trends in biochemical sciences*, doi:10.1016/j.tibs.2017.11.008 (2017).

Rudd, C. E. CD4, CD8 and the TCR-CD3 complex: a novel class of protein-tyrosine kinase receptor. *Immunology Today* **11**, 400-406, doi:https://doi.org/10.1016/0167-5699(90)90159-7 (1990).

Xu, H. & Littman, D. R. A kinase-independent function of Lck in potentiating antigen-specific T cell activation. *Cell* **74**, 633-643, doi:10.1016/0092-8674(93)90511-n (1993).

Glatzova, D. & Cebecauer, M. Dual Role of CD4 in Peripheral T Lymphocytes. *Frontiers in immunology* **10**, 618, doi:10.3389/fimmu.2019.00618 (2019).

Wang, X. X. *et al.* Affinity maturation of human CD4 by yeast surface display and crystal structure of a CD4-HLA-DR1 complex. *P Natl Acad Sci USA* **108**, 15960-15965, doi:10.1073/pnas.1109438108 (2011).

Huppa, J. B. *et al.* TCR-peptide-MHC interactions in situ show accelerated kinetics and increased affinity. *Nature* **463**, 963-967, doi:10.1038/nature08746 (2010).

Dong *et al.* Structural basis of assembly of the human T cell receptor-CD3 complex. *Nature*, doi:10.1038/s41586-019-1537-0 (2019).

Casas, J. *et al.* Ligand-engaged TCR is triggered by Lck not associated with CD8 coreceptor. *Nature communications* **5**, 5624, doi:10.1038/ncomms6624 (2014).

Crawford, F., Kozono, H., White, J., Marrack, P. & Kappler, J. Detection of Antigen-Specific T Cells with Multivalent Soluble Class II MHC Covalent Peptide Complexes. *Immunity* **8**, 675-682, doi:10.1016/s1074-7613(00)80572-5 (1998).

Glassman, C. R., Parrish, H. L., Lee, M. S. & Kuhns, M. S. Reciprocal TCR-CD3 and CD4 Engagement of a Nucleating pMHCII Stabilizes a Functional Receptor Macrocomplex. *Cell Reports* **22**, 1263-1275, doi:10.1016/j.celrep.2017.12.104 (2018).

Chesla, S. E., Selvaraj, P. & Zhu, C. Measuring Two-Dimensional Receptor-Ligand Binding Kinetics by Micropipette. *Biophys. J.* **75**, 1553-1572 (1998).

Huang, J. *et al.* The kinetics of two-dimensional TCR and pMHC interactions determine T-cell responsiveness. *Nature* **464**, 932-U156, doi:10.1038/nature08944 (2010).

Deng, L. *et al.* Structural basis for the recognition of mutant self by a tumor-specific, MHC class II-restricted T cell receptor. *Nat. Immunol.* **8**, 398-408, doi:10.1038/ni1447 (2007).

Huang, J., Edwards, L. J., Evavold, B. D. & Zhu, C. Kinetics of MHC-CD8 interaction at the T cell membrane. *J. Immunol.* **179**, 7653-7662 (2007).

Zhu, C. & Williams, T. E. Modeling concurrent binding of multiple molecular species in cell adhesion. *Biophysical journal* **79**, 1850-1857, doi:10.1016/S0006-3495(00)76434-4 (2000).

Williams, T. E., Selvaraj, P. & Zhu, C. Concurrent binding to multiple ligands: kinetic rates of CD16b for membrane-bound IgG1 and IgG2. *Biophysical journal* **79**, 1858-1866, doi:10.1016/S0006-3495(00)76435-6 (2000).

Williams, T. E., Nagarajan, S., Selvaraj, P. & Zhu, C. Concurrent and independent binding of Fcgamma receptors IIa and IIIb to surface-bound IgG. *Biophys J* **79**, 1867-1875, doi:10.1016/S0006-3495(00)76436-8 (2000).

Chen, W., Evans, E. A., McEver, R. P. & Zhu, C. Monitoring receptor-ligand interactions between surfaces by thermal fluctuations. *Biophys. J.* **94**, 694-701 (2008).

Marshall, B. T. *et al.* Direct observation of catch bonds involving cell-adhesion molecules. *Nature* **423**, 190-193, doi:10.1038/nature01605 (2003).

Williams, T. E., Nagarajan, S., Selvaraj, P. & Zhu, C. Quantifying the impact of membrane microtopology on effective two-dimensional affinity. *J Biol Chem* **276**, 13283-13288, doi:10.1074/jbc.M010427200 (2001).

Kao, H. & Allen, P. M. An antagonist peptide mediates positive selection and CD4 lineage commitment of MHC class II-restricted T cells in the absence of CD4. *J. Exp. Med.* **201**, 149-158, doi:10.1084/jem.20041574 (2005).

Rothemund, P. W. Folding DNA to create nanoscale shapes and patterns. *Nature* **440**, 297-302, doi:10.1038/nature04586 (2006).

Li, Q.-J. *et al.* CD4 enhances T cell sensitivity to antigen by coordinating Lck accumulation at the immunological synapse. *Nat Immunol* **5**, 791-799 (2004).

Natarajan, K. *et al.* An allosteric site in the T-cell receptor Cbeta domain plays a critical signalling role. *Nature communications* **8**, 15260, doi:10.1038/ncomms15260 (2017).

He, Y. *et al.* Peptide-MHC Binding Reveals Conserved Allosteric Sites in MHC Class I- and Class II-Restricted T Cell Receptors (TCRs). *J Mol Biol* **432**, 166697, doi:10.1016/j.jmb.2020.10.031 (2020).

Rangarajan, S. *et al.* Peptide-MHC (pMHC) binding to a human antiviral T cell receptor induces long-range allosteric communication between pMHC- and CD3-binding sites. *The Journal of biological chemistry* **293**, 15991-16005, doi:10.1074/jbc.RA118.003832 (2018).

Zhu, C., Chen, W., Lou, J., Rittase, W. & Li, K. Mechanosensing through immunoreceptors. *Nature immunology*, doi:10.1038/s41590-019-0491-1 (2019).

Li, K. *et al.* PD-1 suppresses TCR-CD8 cooperativity during T-cell antigen recognition. *Nature communications* **12**, 2746, doi:10.1038/s41467-021-22965-9 (2021).

Rudd, C. E., Trevillyan, J. M., Dasgupta, J. D., Wong, L. L. & Schlossman, S. F. The CD4 receptor is complexed in detergent lysates to a protein-tyrosine kinase (pp58) from human T lymphocytes. *Proc Natl Acad Sci U S A* **85**, 5190-5194, doi:10.1073/pnas.85.14.5190 (1988).

Burgess, K. E. *et al.* Biochemical identification of a direct physical interaction between the CD4:p56lck and Ti(TcR)/CD3 complexes. *European journal of immunology* **21**, 1663-1668, doi:10.1002/eji.1830210712 (1991).

Duplay, P., Thome, M., Herve, F. & Acuto, O. P56(Lck) Interacts Via Its Src Homology-2 Domain with the Zap-70 Kinase. *Journal of Experimental Medicine* **179**, 1163-1172, doi:DOI 10.1084/jem.179.4.1163 (1994).

Thome, M., Germain, V., DiSanto, J. P. & Acuto, O. The p56lck SH2 domain mediates recruitment of CD8/p56lck to the activated T cell receptor/CD3/zeta complex. *European journal of immunology* **26**, 2093-2100, doi:10.1002/eji.1830260920 (1996).

Chen, W., Zarnitsyna, V. I., Sarangapani, K. K., Huang, J. & Zhu, C. Measuring Receptor-Ligand Binding Kinetics on Cell Surfaces: From Adhesion Frequency to Thermal Fluctuation Methods. *Cell Mol Bioeng* **1**, 276-288, doi:10.1007/s12195-008-0024-8 (2008).

Evans, E., Leung, A., Heinrich, V. & Zhu, C. Mechanical switching and coupling between two dissociation pathways in a P-selectin adhesion bond. *Proc Natl Acad Sci U S A* **101**, 11281-11286, doi:10.1073/pnas.0401870101 (2004).

Buckley, C. D. *et al.* Cell adhesion. The minimal cadherin-catenin complex binds to actin filaments under force. *Science* **346**, 1254211, doi:10.1126/science.1254211 (2014).

Bell, G. Models for the specific adhesion of cells to cells. *Science* **200**, 618-627, doi:10.1126/science.347575 (1978).

Figure 1

Figure 1 | 2D kinetic analysis of TPI:HLA-DR1 interaction with E8 TCR or CD4. a,

Schematics of micropipette binding frequency assay (*top*) and molecular coating (*bottom*). A

pMHC-coated RBC (*left*) was repeatedly brought into contact with an opposing RBC coated with

E8 TCR, or WT/MT CD4 (right) and separated to detect bond formation across the contact

interface. **b,** Specificity is shown by the significant higher (p-value indicated, by Mann-Whitney

test) of steady-state (contact time t_c = 4 s) adhesion frequencies (P_a, mean ± SEM (n = 6, 6, 9, 3

cell pairs) of indicated points each determined from 50 cycles of contact) between pMHC-coated

RBCs and RBCs coated with E8 TCR, WT CD4, or MT CD4, than that without ligand. **c,** Mean ±

SEM (n = 3 cell pairs per data point) P_a vs t_c curves for pMHC interacting with MT CD4 at various

coating densities as indicated by the number in legend. **d,** The normalized bond number $<n>/m_r m_l$

vs t_c curve for pMHC interaction with E8 TCR (green), WT CD4 (purple), or MT CD4 (orange).

Different symbols represent independent experiments using different molecular densities that are

shown in Supplementary Table 1. **e,** Normalized bond number at steady-state calculated from **d**

was plotted vs receptor and ligand densities.

Figure 2

Figure 2 | Ectodomains of TCR and CD4 bind cooperatively to pMHC. a, P_a vs t_c data and their model fits of pMHC-coated RBCs (38 μm^{-2}) binding to RBCs coated with TCR alone, or a mixture of TCR and CD4 at densities (# of molecules/μm^2) indicated by the number in legend. **b,** Normalized whole bond number $<n>_w/m_{TCR}m_{pMHC}$ vs t_c curves and their respective model fits for interactions in **a**. Arrow indicates decreasing density of TCR and increasing density of CD4. **c,** Normalized trimolecular bond number $<n>_{tri}/m_{TCR}m_{pMHC}m_{CD4}$ vs t_c curves calculated from **b** by subtracting the black data from colored data followed by further normalization by m_{CD4}. **d,** Survival probabilities plotted as ln(# of events with lifetime greater than t) vs bond lifetime for pMHC-coated beads interacting with beads coated with TCR, CD4, or a 1:1 mixture of TCR and CD4 measured by BFP thermal fluctuation at zero force. Each data set (points) was fitted by a straight line with the negative slope representing off-rate for TCR ($k_{-1,TCR}$, n = 71) or CD4 ($k_{-1, CD4}$, n = 92), or apparent off-rate for the mixed TCR and CD4 (n = 97). As a negative control, class-I MHC 9V:HLA-A2 was used instead of TPI:HLA-DR1 to measure non-specific binding (n = 42). **e,** Mean ± SEM bond lifetime vs force data (points), measured by BFP force-clamp experiment using beads bearing TPI:HLA-DR1 interacting with E8 TCR (n = 230), CD4 (n = 355), or a 1:1 mixture of E8 TCR and CD4 (n = 424) coated on beads, are compared with theoretical curves (mean ± SE) predicted using the model and best-fit parameters in Supplementary Fig. 3. **f,** Model predicted trimolecular bond lifetime vs force plots for strong (red) and weak (blue) states as well as their sum (black). Data (points) in **a** and **b** are presented as mean ± SEM of two independent experiments for each curve. Their model fits (mean ± SE) are calculated using Eqs. 1 and 2 based on the best-fit parameters listed in Table 1.

Figure 3

Figure 3 | Cellular TCR and CD4 bind cooperatively to pMHC. a, P_a vs t_c curves (Mean ± SEM of 5 or more cells per data point) and their model fits of pMHC-coated beads binding to Jurkat cells expressing TCR alone, CD4 alone, or both at densities (μm^{-2}) indicated by the number in legend. The dashed curve represents predict sum of two concurrent and independent bimolecular interactions, which used the kinetic parameters of the two bimolecular interactions with molecular densities matched those used in the experiment measuring total interactions. Best-fit P_a ± SE was calculated from best-fit $<n>$ ± SE in **b**. p = 0.0016, 0.0004, 0.0013, 0.00005, 0.0004, and 0.0066 for comparing measured P_a of whole interactions to predicted sum at t_c = 0.5, 1, 2, 4, 8, and 16 s. **b&c,** Normalized whole bond number $<n>_w/m_{TCR}m_{pMHC}$ vs t_c curves and their respective model fits for interactions in **a**. Predications were made after normalizing the densities in different curves. TCR + CD4 curve was fitted with (**c**) or without (**b**) keeping the same TCR affinity and off-rate as in the bimolecular interaction. p = 0.0022, 0.0006, 0.0019, 0.0001, 0.0011, and 0.0140 for comparing measured whole bonds to predicted sum at t_c = 0.5, 1, 2, 4, 8, and 16 s in **b**. p = 0.0044, 0.0013, 0.0066, 0.0029, 0.3089, and 0.5179 for comparing measured whole bonds to its fitted curve at t_c = 0.5, 1, 2, 4, 8, and 16 s in **c**. **d,** Mean ± SEM bond lifetime vs force data of BFP beads bearing TPI:HLA-DR1 interacting with Jurkat cells expressing E8 TCR (n = 598), CD4 (n = 283), E8 TCR and CD4 of similar density (n = 279) or more TCR than CD4 (n = 288). expressed on Jurkat cells. **e,** Mean ± SEM bond lifetime vs force data of BFP beads bearing Hb:I-E^k interacting with 3.L2 CD8$^+$CD4$^-$ T cells (n = 298), CD8$^-$CD4$^+$ T cells (n = 711), or CD8$^-$CD4$^+$ T cells with CD4 blocking antibody clone GK1.5 (n = 240). **f,** Comparison among normalized bond numbers for pMHC interaction with TCR or mixed TCR and CD4 coated in four ways at densities (# of molecules/μm^2) indicated by the number in legend: 1) pMHC on RBCs vs TCR on RBCs (green), 2) pMHC on RBCs vs mixed TCR and CD4 on RBCs (black), 3) pMHC on RBCs vs mixed TCR

1   and CD4 on beads (orange), 4) pMHC on beads vs mixed TCR and CD4 on RBCs (blue). Data

2   (points) are presented as mean ± SEM of two independent experiments for each curve. Their model

3   fits (mean ± SE) are calculated using Eqs. 1 and 2 based on the best-bit parameters listed in Table

4   1.

Figure 4

a

100nm

b

RBC

pMHC

Bead

TCR and CD4
on origami

c

Adjustable distance (d)

TCR   CD4

Capturing
DNA

Bead surface

d

d = 6nm

d = 13nm

d = 20nm

d = 100nm

e

Ladder   CD4   CD4-DNA

70k

60k

50k

f

1   2   3   4   5   6

g

STV   TCR   CD4

d=13nm

STV   TCR   CD4

50nm

d=20nm

Figure 4 | DNA origami-based platform for spacing control of TCR and CD4. a, Schematic of "surrogate" T cell – magnetic bead bearing 10 HB origami functionalized with precisely spaced TCR and CD4. **b,** Schematic of micropipette adhesion frequency assay combined with ligand presentation by origami. **c,** 10 HB origami controls separation distance of TCR and CD4 and attaches to magnetic bead via DNA strand hybridization. TCR binds to a streptavidin (SA), which also binds to a biotinylated DNA strand that hybridizes with a TCR-capturing DNA strand on 10HB origami. CD4 is covalently conjugated with a DNA stand that binds to a CD4-captruing strand. **d,** Four distances were employed, with either 6, 13, 20, or 100 nm TCR–CD4 spacing. **e,** SDS PAGE of CD4 conjugated to oligonucleotide for loading onto 10HB origami. **f,** Native agarose gel electrophoresis of 10 HB origami with 1) no poly-A extensions or protein handles, 2) poly-A extensions but no protein handles, 3-6) poly-A extensions with protein handles spaced at 6, 13, 20 and 100 nm respectively. **g,** TEM image of 10HB origami bearing CD4 and SA-coupled TCR with 13 nm (top) and 20 nm (bottom) spacing.

Figure 5

a

$$y = A_C K_{a,TCR} \left(1 + m_{CD4} K^*_{a,CD4} e^{\frac{-(x-x_0)}{l}} \right)$$

$p = 0.0022$
$p = 0.0402$
$p = 0.4168$
$p = 0.8619$
$p = 0.0002$

y-axis: $\langle n \rangle_w / m_{TCR} m_{pMHC}$ (μm^4)

x-axis: TCR-CD4 Separation Distance x (nm)

6   13   20   100   TCR only   CD4 only

b

Force (pN)

slope = 0.6212 pN/nm
slope = 0.3155 pN/nm

Extension (nm)

c

6 nm
51%
0.61 pN/nm
49%
0.32 pN/nm
$R^2 = 0.96$

13 nm
62%
0.31 pN/nm
38%
0.58 pN/nm
$R^2 = 0.99$

20 nm
100%
0.32 pN/nm
$R^2 = 0.99$

100 nm
100%
0.32 pN/nm
$R^2 = 0.97$

Frequency

Stiffness (pN/nm)

Figure 5 | TCR-CD4 cooperativity is optimal at 7-nm spacing. a, Normalized whole number of

bonds $\langle n \rangle_w / m_{TCR} m_{pMHC}$ for pMHC-bearing RBCs interacting with DNA origami beads presenting

TCR-CD4 pairs at 6, 13, 20, or 100 nm spacing, or presenting TCR alone or CD4 along (points, n

= 21, 17, 15, 15, 8, and 7 cell-bead pairs). In the violin plots that show data densities, the solid

lines in the middle represent mean and the lower and upper dotted lines represent the first and third

quartiles. Eq. 4 was fit (mean $\pm$ SE curves) to the $\langle n \rangle_w / m_{TCR} m_{pMHC}$ vs separation distance (x) data

using the CD4 density (m_{CD4} = 31 μm^{-2}) measured by flow cytometry. P-values indicate

comparison with TCR only group using the Mann-Whitney test. **b,** Representative force vs

molecular extension plot illustrating the determination of molecular stiffness using the Hook's law.

Blue: compression phase, where slope represents the spring constant of the cell. Orange: extension

phase, where the slope represents the resultant spring constant of the cell the molecule connected

in series. **c,** Histograms of stiffness of bonds between pMHC and TCR and CD4 spaced at 6, 13,

20, or 100 nm. Histograms were fitted by single gaussian distribution (20 and 100 nm) or double

gaussian distribution (6 and 13 nm) to identify the peak value(s) and the associated fractions.

Table 1 | Kinetic parameters of the purified protein system and the measurement methods.

E8 TCR		CD4		Mutant CD4	
$A_c K_{a,TCR}$ (μm⁴)	Source	$A_c K_{a,CD4}$ (μm⁴)	Source	$A_c K_{a,MTCD4}$ (μm⁴)	Source
$7.70 \pm 0.40 \times 10^{-4}$	Fig. 1d	$4.35 \pm 0.49 \times 10^{-7}$	Fig. 1d	$2.48 \pm 0.15 \times 10^{-3}$	Fig. 1d
$3.01 \pm 0.69 \times 10^{-4}$	Fig. S1a	$7.74 \pm 0.60 \times 10^{-7}$	Fig. S1b	$1.30 \pm 0.26 \times 10^{-3}$	Fig. S1c
$8.31 \pm 0.46 \times 10^{-4}$	Fig. 2a,b	$6.57 \pm 0.64 \times 10^{-6}$	Fig. 3a purple	$k_{-1,MTCD4}$ (s⁻¹)	Source
$3.59 \pm 0.37 \times 10^{-5}$	Fig. 3a, green	$k_{-1,CD4}$ (s⁻¹)	Source	0.80 ± 0.15	Fig. 1d
$1.42 \pm 0.46 \times 10^{-4}$	Fig. 3a, black	0.37 ± 0.11	Fig. 1d	$A_c k_{+1,MTCD4}$ (μm⁴s⁻¹)	Source
$6.45 \pm 0.58 \times 10^{-4}$	Fig. 3f, green	4.0 ± 0.03	Fig. 2d	$2.00 \pm 0.30 \times 10^{-3}$	Fig. 1d
$5.30 \pm 0.33 \times 10^{-4}$	Fig. 3f, black	0.66 ± 0.08	Fig. 3a purple		
$6.09 \pm 0.62 \times 10^{-4}$	Fig. 3f, orange	$A_c k_{+1,CD4}$ (μm⁴s⁻¹)	Source		
$20.2 \pm 1.8 \times 10^{-4}$	Fig. 3f, blue	$1.62 \pm 0.35 \times 10^{-7}$	Fig. 1d		
$8.39 \pm 1.07 \times 10^{-4}$	Fig. S2c	$K_{a,CD4}^*$ (μm²)	Source		
$1.29 \pm 0.52 \times 10^{-5}$	Fig. 5a	$5.26 \pm 0.72 \times 10^{-2}$	Fig. 2a,b		
$k_{-1,TCR}$ (s⁻¹)	Source	$2.69 \pm 1.14 \times 10^{-1}$	Fig. 3b, black		
0.48 ± 0.07	Fig. 1d	$(7.26 \pm 1.52 \times 10^{-1})^{§§}$	Fig. 3c, black		
0.53 ± 0.05	Fig. 2a,b	$29.9 \pm 2.7 \times 10^{-2}$	Fig. 3f, black		
11 ± 0.19	Fig. 2d	$1.0 \pm 42{,}000 \times 10^{-9}$	Fig. 3f, orange		
0.34 ± 0.05	Fig. 3a, green	$37.5 \pm 9.7 \times 10^{-2}$	Fig. 3f, blue		
1.60 ± 1.06	Fig. 3a, black	$4.87 \pm 1.07 \times 10^{-2}$	Fig. S2c		
0.41 ± 0.09	Fig. 3f, green	$5.0 \pm 1.5 \times 10^{-2}$	Fig. 5a		
1.11 ± 0.11	Fig. 3f, black	$k_{-2,CD4}$ (s⁻¹)	Source		
0.99 ± 0.34	Fig. 3f, orange	1.99 ± 1.6	Fig. 2a,b		
$(0.99 \pm 0.34)^{§}$	Fig. 3f, blue	0.13 ± 0.12	Fig. 3b, black		
$A_c k_{+1,TCR}$ (μm⁴s⁻¹)	Source	$(10 \pm 17)^{§§}$	Fig. 3c, black		
$3.66 \pm 0.42 \times 10^{-4}$	Fig. 1d	4.2 ± 2.5	Fig. 3f, black		
$4.38 \pm 0.48 \times 10^{-4}$	Fig. 2a,b	2.0 ± 3.8	Fig. 3f, orange		
$2.66 \pm 0.46 \times 10^{-4}$	Fig. 3f, green	$(2.0 \pm 3.8)^{§}$	Fig. 3f, blue		
$5.88 \pm 0.69 \times 10^{-4}$	Fig. 3f, black	$k_{+2,CD4}$ (μm² s⁻¹)	Source		
$6.03 \pm 2.2 \times 10^{-4}$	Fig. 3f, orange	0.11 ± 0.093	Fig. 2a,b		
$20.0 \pm 7.1 \times 10^{-4}$	Fig. 3f, blue	0.35 ± 0.35	Fig. 3b, black		
		$(7.2 \pm 12)^{§§}$	Fig. 3c, black		
		1.26 ± 0.76	Fig. 3f, black		
		$2.0 \pm 84{,}000 \times 10^{-9}$	Fig. 3f, orange		
		0.75 ± 1.4	Fig. 3f, blue		

§Use the same value as Fig. 3f, orange.

§§Two-parameter fit values.

Methods:

Fig. 1d: Micropipette adhesion frequency assay with fitting to Eq. 8.

Fig. 2a,b: Micropipette adhesion frequency assay with fitting to Eqs. 1 and 2.

Fig. 2d: BFP thermal fluctuation assay with fitting to $\ln(p) = -k_{off} t$.

Fig. 3a,b: BFP adhesion frequency assay with fitting to Eqs. 1 and 2.

Fig. 3c: BFP adhesion frequency assay with the green and purple data fitting to Eq. 1 and the black data fitting to Eq. 2 with only two parameters ($K_{a,CD4}^*$ and $k_{-2,CD4}$) after fixing $A_c K_{a,TCR}$ and $k_{-1,TCR}$ at values obtained from fitting the green and purple data.

Fig 3f: Micropipette adhesion frequency assay with fitting to Eq. 8 (green curve) and Eqs. 1 and 2 (green and black curves, orange curve, and blue curve with $k_{-1,TCR}$ and $k_{-2,CD4}$ fixed at the values obtained from fitting the orange curve).

Fig. 5a: Micropipette adhesion frequency assay with fitting to Eq. 4. at steady-state.

Fig. S1a-c: Micropipette adhesion frequency assay with fitting to Eq. 7 at steady-state.

Fig. S2c: Micropipette adhesion frequency assay with fitting to Eq. 3. at steady-state.