1 **Induction and inhibition of *Drosophila* X chromosome gene expression are both impeded**
2 **by the dosage compensation complex**
3

4 Richard P. Meisel[1]*, Danial Asgari[1], Florencia Schlamp[2], Robert L. Unckless[3]
5
1. Department of Biology and Biochemistry, University of Houston, 3455 Cullen Blvd, Houston,
TX 77204-5001, USA
2. Department of Medicine, NYU Grossman School of Medicine, 435 E 30th St, New York, NY
10016, USA
3. Department of Molecular Biosciences, University of Kansas, 4055 Haworth Hall, 1200
Sunnyside Avenue, Lawrence, KS 66045, USA
12
* To whom correspondence should be addressed:
Richard P. Meisel
3455 Cullen Blvd.
Houston, TX 77204
USA
rpmeisel@uh.edu

19
20 **Running title:** X chromosome induction and inhibition

## ABSTRACT

Sex chromosomes frequently differ from the autosomes in the frequencies of genes with sexually dimorphic or tissue-specific expression. Multiple hypotheses have been put forth to explain the unique gene content of the X chromosome, including selection against male-beneficial X-linked alleles, expression limits imposed by the haploid dosage of the X in males, and interference by the dosage compensation complex (DCC) on expression in males. Here, we investigate these hypotheses by examining differential gene expression in *Drosophila melanogaster* following several treatments that have widespread transcriptomic effects: bacterial infection, viral infection, and abiotic stress. We found that genes that are induced (up-regulated) by these biotic and abiotic treatments are frequently under-represented on the X chromosome, but so are those that are repressed (down-regulated) following treatment. We further show that whether a gene is bound by the DCC in males can largely explain the paucity of both up- and down-regulated genes on the X chromosome. Specifically, genes that are bound by the DCC, or close to a DCC high-affinity site, are unlikely to be up- or down-regulated after treatment. This relationship, however, could partially be explained by a correlation between differential expression and breadth of expression across tissues. Nonetheless, our results suggest that DCC binding, or the associated chromatin modifications, inhibit both up- and down-regulation of X chromosome gene expression within specific contexts. We propose multiple possible mechanisms of action for the effect, including a role of Males absent on the first (Mof), a component of the DCC, as a dampener of gene expression variance in both males and females. This effect could explain why the *Drosophila* X chromosome is depauperate in genes with tissue-specific or induced expression, while the mammalian X has an excess of genes with tissue-specific expression.

**INTRODUCTION**

Many animal species, as well as some plants and other eukaryotes, have sex
chromosomes, which are often under different transcriptional regulation than the autosomes.
Sex chromosomes can be grouped into several different categories, with XY and ZW systems
amongst the most common in animals (Bachtrog *et al.* 2014). X and Z chromosome gene
expression can be controlled by transcriptional regulators and histone modifications that are
unique from the autosomes (Lucchesi *et al.* 2005; Ferrari *et al.* 2014; Gu *et al.* 2019). For
example, one copy of the mammalian X chromosome is silenced (via the recruitment of
facultative heterochromatin) in somatic tissues of XX females by a combination of non-coding
RNAs and proteins (Lyon 1961; Brown *et al.* 1991; Chow *et al.* 2005). In contrast, the
*Drosophila* dosage compensation complex (DCC) up-regulates gene expression on the X
chromosome in males using a combination of RNAs and proteins (Lucchesi and Kuroda 2015).
The DCC only assembles in male somatic tissues, where it initiates the acetylation of lysine 16
in histone H4 (H4K16ac) specifically on the X chromosome, compensating for the haploid dose
(Gelbart *et al.* 2009). Furthermore, in the male germline of some animal species, there is
evidence for silencing of the single X chromosome (Lifschytz and Lindsley 1972; Turner 2007),
although the extent of this meiotic sex chromosome inactivation (MSCI) varies across taxa
(Bean *et al.* 2004; Meiklejohn *et al.* 2011; Turner 2015).
The unique transcriptional and chromatin environments of X chromosomes, along with
their hemizygosity in males, create selection pressures on X-linked genes that differ from the
autosomes, resulting in X-autosome differences in gene content that are taxon-specific. For
example, the mammalian X chromosome is enriched for genes that are expressed specifically in
male reproductive tissues, such as the prostate and testis (Wang *et al.* 2001; Lercher *et al.*
2003; Mueller *et al.* 2008, 2013; Meisel *et al.* 2012a). In contrast, the *Drosophila melanogaster*
X chromosome contains very few genes that are expressed primarily in the male-specific
accessory gland, a reproductive organ analogous to the mammalian prostate (Swanson *et al.*
2001; Ravi Ram and Wolfner 2007; Meisel *et al.* 2012a). The *Drosophila* X chromosome also
contains a paucity of genes with male-biased expression (i.e., up-regulated in males relative to
females) relative to the autosomes (Parisi *et al.* 2003; Sturgill *et al.* 2007). Taxon-specific X-
autosome differences in gene content further extend to genes with non-reproductive functions.
In *D. melanogaster*, for instance, the X chromosome is deficient for genes that have narrow
expression in non-reproductive tissues, whereas the mammalian X is enriched for genes with
tissue-specific expression (Lercher *et al.* 2003; Mikhaylova and Nurminsky 2011; Meisel *et al.*
2012a).
Multiple hypotheses have been proposed to explain the differences in gene content
between X chromosomes and autosomes (Table 1). One of these hypotheses is based upon the
prediction that sexually antagonistic selection will favor recessive male-beneficial mutations (or
dominant female-beneficial alleles) on the X chromosome (Rice 1984; Charlesworth *et al.*
1987). This sexual antagonism hypothesis has numerous limitations (Fry 2010), including the
inability to explain differences between *Drosophila* and mammalian X chromosomes in their
deficiency or enrichment, respectively, of genes expressed in male reproductive tissues (Meisel
*et al.* 2012a). A second hypothesis focuses specifically on the male germline, where MSCI
silences the X chromosome (Lifschytz and Lindsley 1972) and may favor duplication of genes to
the autosomes (Betrán *et al.* 2002; Emerson *et al.* 2004; Potrzebowski *et al.* 2008; Vibranovski

*et al.* 2009). However, there is not a deficiency of testis-biased genes on the *D. melanogaster* X
chromosome (Meisel *et al.* 2012a; Meiklejohn and Presgraves 2012), limiting the ability of MSCI
to explain the unique gene content of the *Drosophila* X chromosome. Third, the haploid dose of
the X in males may impose a maximal gene expression level lower than the autosomes,
selecting against X-linked genes with high expression (Wolfner *et al.* 1997; Vicoso and
Charlesworth 2009; Hurst *et al.* 2015). This "dosage limit" hypothesis may even apply in species
where the haploid X is dosage compensated by up-regulation of X-linked expression. For
example, in *D. melanogaster* there may be a transcriptional limit beyond which expression
cannot be exceeded or some genes may not be dosage compensated in males (Meisel *et al.*
2012a).
98
99 **Table 1.** Hypotheses to explain the unique gene content of the *Drosophila* X chromosome.

| Hypothesis | Predicted X-autosome differences |
| --- | --- |
| Sexual antagonism | Excess or deficiency of X-linked genes with sex-specific functions and/or expressed in reproductive tissues |
| MSCI | Deficiency of X-linked testis-expressed genes |
| Dosage limit | Deficiency of highly expressed or induced X-linked genes |
| DCC-interference | Deficiency of DCC-bound genes highly expressed in males |
| Variance dampening | Deficiency of X-linked genes up- or down-regulated in specific contexts (e.g., tissues, infections, abiotic treatments) |

100
Here, we focus on the effect of the DCC on X chromosome expression and gene content
in *D. melanogaster*. The DCC most strongly binds to more than 100 so-called chromatin entry or
high affinity sites (HAS), from which it is thought to spread across the X chromosome (Kelley *et*
*al.* 1999; Alekseyenko *et al.* 2008; Straub *et al.* 2008). Bachtrog *et al.* (2010) observed that
genes near an HAS or bound by the DCC are less likely to have male-biased expression, and
genes further from an HAS have a larger magnitude of male-biased expression. This led them
to hypothesize that the DCC interferes with acquisition of male-biased expression on the X
chromosome.
There is mixed evidence for the hypothesis that DCC-interference is responsible for the
unique gene content of the X chromosome. Consistent with the DCC-interference hypothesis,
when Belyi *et al.* (2020) measured expression of a reporter construct that was inserted at
random locations on the X chromosome, they found reduced expression in male somatic tissues
for transgenes inserted at chromosomal loci closer to endogenous DCC binding sites. However,
when genes with testis-biased expression are excluded or when somatic tissues are analyzed
separately, there is no relationship between male-biased expression and distance from an HAS
for endogenous genes (Vensko and Stone 2014; Gallach and Betrán 2016). In addition, genes
with male-biased expression in brain or head are over-represented on the *D. melanogaster* X
chromosome and closer to DCC binding sites (Huylmans and Parsch 2015), which is opposite
of what is predicted by the DCC-interference hypothesis.

Our analysis addresses a fifth hypothesis, specifically whether the *Drosophila* DCC creates an unfavorable environment for X-linked genes that are differentially expressed in specific contexts. The *D. melanogaster* X chromosome is depauperate in genes with narrow expression in specific tissues (Mikhaylova and Nurminsky 2011; Meisel *et al.* 2012a), and X-linked genes further from an HAS or not bound by the DCC have more tissue-specific expression (Meisel *et al.* 2012b). In contrast, X-linked genes with female-biased expression, which also tend to be broadly expressed (Meisel 2011), are more likely to be bound by the DCC (Gallach and Betrán 2016). This suggests that the DCC creates an unfavorable environment for X-linked genes that are up- or down-regulated in specific tissues, possibly because the DCC prevents the differential regulation of gene expression across contexts. Consistent with this hypothesis, there is evidence that Mof, one of the proteins in the DCC, dampens transcriptional variation on the *D. melanogaster* X chromosome (Lee *et al.* 2018). Moreover, genes that are bound by the DCC have less genetic variation for gene expression than X-linked unbound genes (Meisel *et al.* 2012b), and transgenes inserted on the *D. melanogaster* X chromosome have less intra-locus expression variation in males than females (Belyi *et al.* 2020). Both of these observations are also consistent with the DCC dampening transcriptional variance. This "variance dampening" by the DCC, or Mof specifically, may inhibit context-dependent gene expression by reducing the ability of transcription factors to regulate expression subsequent to DCC-associated chromatin modifications that are already up-regulating expression (Table 1).

We used bacterial infection, viral infection, and abiotic stressors as model systems to test the hypothesis that the *Drosophila* DCC is a variance dampener that reduces differential expression of X-linked genes in specific contexts. Biotic and abiotic stress represents a notable contrast to previous studies of context-dependent expression involving X-autosome comparisons of genes with tissue-specific expression (e.g., Mikhaylova and Nurminsky 2011; Meisel *et al.* 2012a). Bacterial infection, for example, causes the dramatic induction of gene expression, including effectors of the humoral immune system that are expressed more than 100 times higher within 12 hours (De Gregorio *et al.* 2001; Troha *et al.* 2018; Schlamp *et al.* 2021). Curiously, none of the 30–40 *D. melanogaster* genes encoding antimicrobial peptides (AMPs, a class of effectors) are found on the X chromosome (Hill-Burns and Clark 2009), providing *a priori* evidence consistent with selection against X-linked genes induced by infection. We analyzed multiple RNA-seq studies of gene expression after biotic and abiotic treatments to test the hypothesis that the DCC inhibits context-dependent differential expression, which would explain the paucity of genes with tissue- or environment-specific expression on the *Drosophila* X chromosome.

## RESULTS

### Genes differentially expressed after infection are under-represented on the Drosophila melanogaster *X chromosome*

We tested if the *D. melanogaster* X chromosome is depauperate for genes induced (i.e., up-regulated) by bacterial infection regardless of functional annotation. To those ends, we analyzed RNA-seq data in which *D. melanogaster* males were infected with one of 10 different bacteria versus a control (Troha *et al.* 2018). From those infection experiments, we selected results from the five bacterial treatments with >50 differentially expressed (DE) genes, in order to have sufficient power to detect X-autosome differences. For three out of five bacterial

infections we considered, there was a significant deficiency of induced genes on the X
chromosome (Figure 1A). For the remaining two bacterial infections, the observed number of
induced genes on the X chromosome was less than expected, although the difference was not
significant. It is unlikely to observe fewer induced X-linked genes than expected for all five
treatments, assuming a null hypothesis of equal proportions above and below the expectation
($p$=0.031 in a binomial exact test). In addition, when we considered all Gram-positive or Gram-
negative bacteria (with a statistical model that has bacterial strain nested in treatment) from the
experiment together, there was a significant deficiency of induced genes on the X chromosome
in both cases (Figure 1A). Moreover, genes that were up-regulated by at least one, two, three,
or four different bacteria were also significantly under-represented on the X chromosome
(Figure 1B). Therefore, genes that are induced by bacterial infection are generally under-
represented on the *D. melanogaster* X chromosome regardless of the criteria used to categorize
induction. This is consistent with the deficiency of X-linked AMP genes (Hill-Burns and Clark
2009).
There was not a significant deficiency of X-linked *D. melanogaster* genes down-
regulated after infection with any of the five bacterial treatments or when we considered all
Gram-positive or Gram-negative bacteria (Figure 1C). Similarly, genes that were downregulated
by one or more different bacteria were not significantly under-represented on the X
chromosome (Figure 1D). However, in most cases, the number of down-regulated X-linked
genes was less than the expectation, albeit not significant. The failure to detect a significant
deficiency of down-regulated X-linked genes may have been caused by low statistical power—
there were fewer down-regulated genes than up-regulated genes in most bacterial treatments.
We examine this further by considering other experiments with more down-regulated genes
below.
The total number of DE genes (either induced or repressed after infection) on the X
chromosome was significantly less than the expectation for 4 of 5 individual bacterial
treatments, Gram-positive bacteria, and Gram-negative bacteria (Figure 1E). The X
chromosome also has significantly fewer genes that were DE in at least one or more of the
bacterial treatments (Figure 1F). Therefore, both up-regulated and DE genes are under-
represented on the X chromosome.
To further evaluate if the X chromosome is anomalous, we tested if any individual
autosomes had a deficiency (or excess) of induced, repressed, or DE genes following bacterial
infection. Notably, the left arm of the second chromosome (2L) had an excess of induced genes
in every treatment (Supplemental Figures S1-S2). This is surprising because none of the
annotated *D. melanogaster* AMP genes are on chromosome 2L, and only 10/74 immune
effector genes are found on 2L (Sackton *et al.* 2007). Chromosome 2L therefore has an excess
of genes induced by bacterial infection (Supplemental Figures S1-S2), despite having a
significant deficiency of effector genes ($p$=0.006 comparing effector and non-effector immune
genes on chromosome 2L with the other chromosomes in a Fisher's exact test). The right arm
of the third chromosome (3R), in contrast, has a deficiency of DE genes (Supplemental Figures
S1-S2), even though it contains at least five AMP genes and has neither an excess nor a
deficiency of effector genes ($p$=0.7 comparing effectors and non-effectors between 3R and
other chromosomes in Fisher's exact test). We next ranked each chromosome arm within each
treatment by the percent of induced, repressed, or DE genes (excluding the diminutive

chromosome 4 because it has <100 genes). On average, the X chromosome has the lowest
percentage of induced, repressed, or DE genes across all treatments, and chromosome 3R has
the second lowest (Supplemental Figures S3-S4). Therefore, the X chromosome is anomalous
from each of the autosomes in its deficiency of up-regulated and DE genes after bacterial
infection.
212

213
**Figure 1.** Genes that were differentially expressed after bacterial infection in *D. melanogaster* males are
under-represented on the X chromosome. Bar graphs show the number of X-linked genes that were up-
regulated (A-B), down-regulated (C-D), or differentially expressed (E-F) after infection with individual
bacteria, combined Gram-negative (Gram–) bacteria, combined Gram-positive (Gram+) bacteria, (A, C,
E), or across multiple bacterial treatments (B, D, F). Dots connected by broken lines show the expected
number of X-linked genes in each category based on the fraction of autosomal genes that are
differentially expressed after infection. Black asterisks represent significant differences between observed
and expected counts in a Fisher's exact test (*$P<0.05$, **$P<0.005$, ***$P<0.0005$).
222
If the X chromosome has a maximal expression that prevents up-regulation of individual
genes (i.e., a dosage limit), we should observe a difference in the distribution of $\log_2$FC values
between X-linked and autosomal genes (Meiklejohn *et al.* 2011; Meiklejohn and Presgraves
2012). If we do not observe such a difference, it would suggest that there is not a dosage limit
that selects against induced X-linked genes (Table 1). The X chromosome did not have a

significantly lower log$_2$FC than the autosomes in any of the five bacterial treatments
(Supplemental Figure S5). In two of the five bacterial treatments (*M. luteus* and *S. marcescens*
*Db11*), X-linked genes possess a higher median log$_2$FC than autosomal genes (Supplemental
Figure S5), which is opposite of the direction predicted by the dosage limit hypothesis.
Therefore, the paucity of X-linked genes up-regulated after infection cannot be explained by an
overall reduced dose of X chromosome gene expression.
234

235
**Figure 2.** Genes that were differentially expressed after bacterial infection in *D. melanogaster* are under-
represented on the X chromosome. Bar graphs show the number of X-linked genes that were up-
regulated (A), down-regulated (B), or differentially expressed (C) after infection with *P. rettgeri* in either
males (top) or females (bottom). Dots connected by broken lines show the expected number of X-linked
genes in each category based on the fraction of autosomal genes that are differentially expressed after
infection. Black asterisks represent significant differences between observed and expected counts in a
Fisher's exact test (*$P<0.05$).
243
We next tested if genes that are induced or repressed in female *D. melanogaster*
following infection are also under-represented on the X chromosome. To those ends, we
identified DE genes in *D. melanogaster* males and females 8 h after infection with *P. rettgeri*
(Duneau *et al.* 2017), one of the bacteria that induced a deficiency of X-linked genes in males
(Figure 1A). We report results for multiple log$_2$FC cutoffs because raw data or *p*-values are not
available for this experiment. Surprisingly, we observed a significant deficiency of X-linked
genes up-regulated in males at only one of the eight log$_2$FC cutoffs we considered (Figure 2A).
In contrast, at five of the eight log$_2$FC cutoffs, there was a significant deficiency of X-linked
genes up-regulated in females after infection (Figure 2A). Similarly, at 5 of 8 log$_2$FC cutoffs
there was a deficiency of X-linked genes down-regulated in females (Figure 2B). Considering
genes that were DE after infection, regardless of up- or down-regulation, there was a significant
deficiency on the X chromosome at 7 and 4 log fold-change cutoffs in females and males,
respectively (Figure 2C). Therefore, there is a significant deficiency of X-linked DE genes after
infection in both male and female *D. melanogaster*.

259
**Figure 3.** Genes that were differentially expressed at different time points after immune challenge in
*D. melanogaster* males are under-represented on the X chromosome. Bar graphs show the number of X-
linked genes that were up-regulated (A), down-regulated (B), or differentially expressed (C) after
*D. melanogaster* males were injected with *E. coli*-derived crude lipopolysaccharide. Dots connected by
broken lines show the expected number of X-linked genes in each category based on the fraction of
autosomal genes that are differentially expressed after infection. Black asterisks represent significant
differences between observed and expected counts in a Fisher's exact test (*P<0.05).

267
We also tested if there is a paucity of X-linked genes induced at different time points
following immune challenge by analyzing RNA-seq data from 1–120 h after *D. melanogaster*
males were injected with *E. coli*-derived crude lipopolysaccharide (Schlamp *et al.* 2021). There
was a significant deficiency of X-linked genes up-regulated at 16 of 19 time points (Figure 3A).
Similarly, at 9 of 19 timepoints, there was a significant deficiency of X-linked genes
downregulated (Figure 3B). Furthermore, there was a significant deficiency of X-linked DE
genes (regardless of up- or down-regulation) at all time points (Figure 3C). Therefore, both up-
and down-regulated genes are under-represented on the X chromosome across the full
temporal spectrum during the response to infection.
These results show that there is a paucity of DE genes on the X chromosome after
infection across a wide range of bacterial pathogens, in both sexes, and across a dense

sampling of timepoints. The paucity of X-linked DE genes can be attributed to a deficiency of
both up- and down-regulated genes. This suggests the deficiency of X-linked DE genes is
robust to experimental variation, and also that gene dosage in males cannot fully explain the
pattern.
283
284

285
**Figure 4.** Genes that are differentially expressed after viral or abiotic treatments can be under- or over-
represented on the X chromosome. Bar graphs show the number of X-linked genes that were up-
regulated (A), down-regulated (B), or differentially expressed (C) after *D. melanogaster* were subjected to
a viral or abiotic treatment. The treatments are listed on the x-axis, and bars are colored based on the sex
of the flies used in the experiment. Dots connected by broken lines show the expected number of X-linked
genes in each category based on the fraction of autosomal genes that are differentially expressed after
treatment. Black asterisks represent significant differences between observed and expected counts in a
Fisher's exact test (*$P<0.05$; *****$P<0.000005$).
294

***Genes differentially expressed under viral or abiotic stress are usually under-represented***
***on the* Drosophila melanogaster *X chromosome***
We next tested if genes induced by viral infection are under-represented on the
*D. melanogaster* X chromosome. We found fewer X-linked genes than expected were induced
by Zika or Kallithea virus, albeit at insignificant differences (Figure 4A). Neither viral infection

resulted in a significant deviation from the expected number of X-linked down-regulated genes
either (Figure 4B). The Kallithea virus data were collected from both males and females, and
genes that were up-regulated in females after Kallithea infection were significantly under-
represented on the X chromosome (Figure 4A). There was also a deficiency of X-linked down-
regulated genes following Kallithea virus infection in males (Figure 4B). In addition, genes that
were DE after Zika virus infection were under-represented on the X chromosome (Figure 4C).
We also tested if genes induced by four different abiotic treatments are under-
represented on the X chromosome. These data include copper treatment for genotypes that are
sensitive to copper and those that are resistant, which we analyzed separately. The data also
included starvation treatments in which gene expression was measured in whole flies, adult
brains after complete starvation, and adult brains after sugar starvation. Both radiation and
sugar starvation resulted in a significant deficiency of X-linked induced genes (Figure 4A).
Complete starvation (with expression measured in the brain) was the only abiotic treatment that
resulted in a significant deficiency of X-linked down-regulated genes (Figure 4B). There was
also a deficiency of X-linked DE genes after both radiation and starvation (Figure 4C). In
contrast to all other biotic and abiotic treatments, there was an excess of X-linked genes up-
regulated after complete starvation and cocaine treatment (Figure 4A). There was also an
excesss of X-linked down-regulated and DE genes after exposure to cocaine, regardless of the
sex of the flies (Figure 4). Both the complete starvation and cocaine treatments measured gene
expression in the brain, suggesting that the brain may be an outlier with an excess, rather than
a deficiency, of X-linked up-regulated genes (or DE genes in general) following abiotic stress.
We further tested if any individual autosomes had a deficiency (or excess) of DE genes
following viral infection or abiotic treatments. None of the autosomal chromosome arms had a
consistent excess or deficiency of induced, repressed, or DE genes after viral or abiotic
treatment (Supplemental Figure S6). When we ranked each chromosome arm within each
treatment by the percent of induced, repressed, or DE genes, the X chromosome had the lowest
percentage of induced, repressed, or DE genes when averaged across all treatments
(Supplemental Figure S7). This provides additional evidence that the X chromosome is an
outlier with a deficiency of DE genes.
In summary, out of six total viral and abiotic treatments, the observed number of X-linked
up-regulated genes was less than expected in most treatments, and significantly so in three
treatments (Figure 4A). Down-regulated and DE genes were also under-represented on the X
chromosome in some treatments (Figure 4B-C). In addition, we observed similar patterns
regardless of sex or genotype of the flies used in the experiments, although sex did affect
whether differences were statistically significant (Figure 4). The notable exceptions to this
pattern are the effects of starvation or cocaine on gene expression in the brain, which were the
only treatments (biotic or abiotic) that resulted in a significant excess of X-linked genes that
were up-regulated (Figure 4).

***DE genes are less likely to be bound by the DCC***
We evaluated the hypothesis that the DCC prevents the induction of X-linked genes by
testing if there is a relationship between induction and DCC binding. In total, a smaller fraction
of DCC-bound genes were up-regulated than X-linked unbound genes for nearly all bacterial,
viral, and abiotic treatments (Figure 5A), if we do not consider whether the difference is
significant. Within individual treatments, DCC-bound genes were significantly less likely to be

up-regulated, relative to X-linked unbound genes, following *P. entomophila* infection, radiation
treatment, starvation (in brain), copper exposure (for resistant flies), and cocaine feeding
(Supplemental Figure S8A). The negative effect of DCC binding on up-regulation following
cocaine was observed for both male and female flies. These results are in accordance with what
would be expected if the DCC prevents induction of X-linked genes.
349

350
**Figure 5.** Genes that are differentially expressed (DE) after treatment are less likely to be bound by the
dosage compensation complex (DCC). The percent of genes that are DE given that they are autosomal
(white), X-linked and bound by the DCC (back), or X-linked and unbound (gray) is plotted for each
treatment. Data are shown for genes up-regulated after a treatment (A), genes down-regulated after a
treatment (B), and all DE genes (C). Red asterisks show significant differences between the number of
genes that are DE and not DE when comparing autosomal vs DCC bound or unbound X-linked genes
(*$P$<0.05 in Fisher's exact test).
358

We next tested if DCC binding could explain the paucity of up-regulated genes on the X
chromosome. To those ends, we determined if there is a difference in the proportion of up-
regulated genes when we compare the autosomes with either DCC-bound or unbound genes on
the X chromosome. If DCC binding explains the paucity of up-regulated genes, we expect a "V-
shaped" pattern when we plot the %DE genes amongst autosomes, DCC-bound X-linked
genes, and unbound X-linked genes (Figure 5). There was a significant deficiency of DCC-
bound up-regulated genes relative to the autosomes across 7 different treatments
(*S. marcescens* Db11, *P. rettgeri*, *P. entomophila*, *S. aureus*, radiation, copper, and cocaine),
among genes up-regulated in at least one bacterial treatment, and for genes up-regulated by
Gram-positive bacteria (Figure 5A). This deficiency is the bottom of the V-shape. In contrast,
there was only one treatment (Kallithea virus in females) in which there was a significant
deficiency of up-regulated X-linked genes unbound by the DCC relative to autosomal genes
(Figure 5A). Most other treatments had the V-shape, with no significant differences in the
fraction of DE genes between autosomal and unbound X-linked genes. In addition, unbound X-
linked genes were significantly more likely to be up-regulated by cocaine than autosomal genes
(Figure 5A). Therefore, the evidence for a paucity of X-linked up-regulated genes is much
greater for DCC-bound than unbound genes (creating the V-shape in Figure 5A), which is
consistent with the expectation if the DCC prevents the induction of X-linked genes.
We also observed that X-linked genes bound by the DCC were less likely to be down-
regulated after treatment than X-linked unbound genes (Figure 5B). In three different viral or
abiotic treatments (Kallithea virus, radiation, and copper), X-linked down-regulated genes were
significantly less likely to be DCC-bound than unbound (Supplemental Figure S8B). The same
results were observed for Kallithea viruses when we considered female samples only, and
similar trends were observed for males (although not significant because of small sample sizes
of down-regulated genes). Therefore, the DCC appears to interfere with both up- and down-
regulation of gene expression. The cumulative effect of the DCC on both up- and down-
regulation can be seen in the significant deficiency of DCC-bound DE genes for five unique
treatments (Supplemental Figure S8C).
DCC-bound genes were also significantly less likely to be down-regulated than
autosomal genes in four different treatments—Kallithea virus, radiation, starvation, and copper
(Figure 5B). Similarly, DCC-bound genes were less likely to be DE than autosomal genes after
most treatments (Figure 5C). In contrast, X-linked unbound genes were significantly more likely
to be down-regulated or DE than autosomal genes after Kallithea virus infection, copper
treatment, or cocaine (Figure 5). Therefore, down-regulated and DE genes also tend to have the
V-shaped distribution. These results are all consistent with the expectations if the DCC prevents
down-regulation of X-linked genes.
Our results provide consistent evidence that DCC binding can largely explain the paucity
of X-linked up-regulated, down-regulated, and DE genes. Notably, we observed much stronger
evidence for a deficiency of X-linked DE genes when we considered DCC-bound genes, and
only weak (or no) evidence for unbound genes (the V-shapes in Figure 5). This is consistent
with the hypothesis that the DCC is a variance dampener that prevents both up- and down-
regulation of X-linked genes. However, there is a deficiency of up-regulated X-linked unbound
genes relative to autosomal genes in some treatments (Figure 5A), suggesting that DCC
binding alone cannot completely explain the paucity of up-regulated genes on the X

chromosome. Therefore, other factors, such as a dosage limit, may also be necessary to
explain the exclusion of up-regulated genes from the *Drosophila* X chromosome.

### Differentially expressed genes are further from DCC high affinity sites

A complementary way to assess the effect of the DCC on differential gene expression is
to measure the distance to the nearest DCC high affinity site (HAS) for each gene. We cannot
test for differences in distance to HAS between DE genes and non-DE genes because there are
too few X-linked DE genes for statistical testing. Instead, we calculated the correlation between
distance to the nearest HAS and $\log_2$ fold-change between treatment and control ($\log_2$FC) for all
genes, regardless of whether they are significantly DE.

412
413

414
**Figure 6.** Correlations between distance from a dosage compensation complex high affinity site (HAS)
and the $\log_2$ fold-change in expression between treatment and control ($\log_2$FC). Each dot is the rank order
correlation ($\rho$) between distance to the nearest HAS and $\log_2$FC. Error bars show the 95% confidence
interval determined by bootstrap resampling the data 1,000 times. The X-axis shows the specific
treatment. Dots and error bars are colored based on the sex of the flies used in the experiment (see
legend). HAS were obtained from two different data sets (Alekseyenko *et al.* 2008; Straub *et al.* 2008),
with results from the two different data sets shown separately in the two columns. Correlations are plotted
with |$\log_2$FC| values for all genes (A), only genes with $\log_2$FC>0 (B), and only genes with $\log_2$FC<0 (C).
423

First, we considered |log$_2$FC| as a measure of the extent of differential expression, regardless of up- or down-regulation. In nearly all combinations of treatments, HAS data sets, and sexes, there was a positive correlation between |log$_2$FC| and distance from an HAS (Figure 6A). Therefore, X-linked genes further from an HAS were more differentially expressed following bacterial infection, viral infection, or abiotic treatment. This is consistent with the hypothesis that the DCC inhibits differential expression (i.e., both up- and down-regulation).

To specifically evaluate if proximity to an HAS affects up-regulation, down-regulation, or both, we separately considered genes with log$_2$FC>0 and log$_2$FC<0 (regardless of whether the deviation from 0 is significant). When we considered only genes with log$_2$FC>0, there was evidence for a positive correlation between log$_2$FC and distance from an HAS for most treatments (Figure 6B). In comparison, when we considered genes with log$_2$FC<0, there was a negative correlation between log$_2$FC and distance from an HAS for most treatments (Figure 6C). Both of these correlations indicate that genes further from an HAS were more likely to be either up- or down-regulated after bacterial infection, viral infection, or abiotic treatment. These results are consistent with the hypothesis that the DCC inhibits both up- and down-regulation of gene expression.

### Differential expression, dosage compensation, and expression breadth

We next considered if expression breadth could explain the correlations between log$_2$FC and distance from an HAS. This analysis was motivated by the previously described observation that genes bound by the DCC or closer to an HAS are narrowly expressed in fewer tissues than X-linked unbound genes and those further from an HAS (Meisel *et al.* 2012b). Here, we calculated partial correlations (Schäfer and Strimmer 2005) between log$_2$FC, distance from an HAS, and expression breadth. We quantified expression breadth using $\tau$, which ranges from 0 (for genes expressed in many tissues ) to 1 (for genes highly expressed in a single tissue) (Yanai *et al.* 2005). We confirmed the positive correlation between $\tau$ and distance from an HAS, even when log$_2$FC is included in the analysis (Supplemental Figures S9–S12). We also found that |log$_2$FC| was positively correlated with $\tau$ (Supplemental Figures S9–S12). Moreover, there was a positive correlation between log$_2$FC and $\tau$ for genes with log$_2$FC>0, and there was a negative correlation for genes with log$_2$FC<0 (Supplemental Figures S9–S12). Therefore, genes that were more up- or down-regulated tended to also be more narrowly expressed. Unsurprisingly, genes induced by bacterial infection were narrowly expressed in the fat body, which is the primary organ of the humoral immune response (Lemaitre and Hoffmann 2007).

When we considered the correlation with $\tau$, many partial correlations between log$_2$FC and distance from an HAS were no longer significantly different from zero (Figure 7; Supplemental Figure S13). This is true regardless of whether sex-specific reproductive tissues are included in the calculation of $\tau$. Therefore, the correlations between log$_2$FC and distance from an HAS could often be explained by the correlations between $\tau$ and both log$_2$FC and distance from an HAS. However, some partial correlations between log$_2$FC and distance from an HAS still remained significantly different from 0 (Figure 7). These significant correlations were almost always in a direction consistent with genes further from an HAS being more up-regulated or more down-regulated (i.e., a positive correlation for |log$_2$FC| or log$_2$FC>0, or a negative correlation for log$_2$FC<0). The one exception to this rule was a positive partial correlation between log$_2$FC and distance from an HAS for genes with log$_2$FC<0 after copper treatment in resistant flies (Figure 7D). This negative correlation is suggestive that genes closer

468    to an HAS are more down-regulated following copper treatment. However, there were very few
469    genes down-regulated after copper treatment in resistant flies (Figures 4–5), suggesting that
470    this positive correlation may be an artifact of a small range of negative log$_2$FC values.
471

472
**Figure 7.** Partial correlations between log$_2$ fold-change in expression between treatment and
control (log$_2$FC) and distance from a dosage compensation complex high affinity site (HAS).
Partial correlations were calculated based on rank order correlations between log$_2$FC, distance
from an HAS, and tissue expression breadth. Each dot shows the partial correlation between
log$_2$FC and distance from an HAS, with the error bars representing 95% confidence intervals from
1,000 bootstrap replicates of the data. The X-axis shows the specific treatment. Dots and error
bars are colored based on the sex of the flies used in the experiment (see legend). HAS were
obtained from the Alekseyenko *et al.* (2008) data set; results from the Straub *et al.* (2008) data
are shown in the supplemental material. Expression breadth was calculated using microarray
data from either 14 unique adult tissues (left) or 10 adult tissues that are not sex-specific (right).
Partial correlations are plotted with log$_2$FC values for all genes (A), |log$_2$FC| values for all genes
(B), only genes with log$_2$FC>0 (C), and only genes with log$_2$FC<0 (D).

## DISCUSSION

We analyzed RNA-seq data from *D. melanogaster* subjected to bacterial, viral, or abiotic treatments. We found that DE genes—both up- and down-regulated—were often depauperate on the X chromosome regardless of the sex of the flies or the type of treatment (Figures 1–4). We additionally determined that DE genes were less likely to be bound by the DCC, and that DCC binding can largely explain the deficiency of X-linked DE genes (Figure 5). In addition, genes that are further from an HAS were more differentially expressed, regardless of whether the genes were up- or down-regulated (Figure 6). However, much of the relationship between differential expression and distance from an HAS could be explained by both variables being correlated with tissue-specific gene expression (Supplemental Figures S9–S12). Nonetheless, a significant correlation between $\log_2$FC and distance from an HAS remained for some treatments after controlling for expression breadth across tissues (Figure 7).

### *Chromosome 2L has a complementary gene content to the X chromosome*

An excess of chromosome 2L genes were up-regulated in many conditions, in contrast to the deficiency of X-linked DE genes. We found an excess of chromosome 2L genes induced in every bacterial treatment (Supplemental Figures S1-S4), even though no AMP genes (and only 10 of 74 immune effectors) are on chromosome 2L (Sackton *et al.* 2007). An excess of chromosome 2L genes were also induced after copper treatment (Supplementary Figure S6). Previous analyses revealed that chromosome 2L (also known as Muller element B) has an excess of genes with male-biased expression in multiple *Drosophila* species, in contrast to the paucity of X-linked genes up-regulated in males (Parisi *et al.* 2003; Meisel *et al.* 2012a). In addition, chromosome 2L has an excess of genes encoding accessory gland proteins, while the X chromosome is depauperate (Ravi Ram and Wolfner 2007). The expression level in accessory gland is also higher for chromosome 2L genes, and lower for X-linked genes, than other chromosomes (Meisel *et al.* 2012a). Moreover, chromosome 2L genes have narrower expression in specific tissues than genes on other chromosomes, in contrast to X-linked genes that tend to be more broadly expressed (Meisel *et al.* 2012a). Future work could address why chromosome 2L has a complementary pattern to the X chromosome.

### *Minimal support for the dosage limit hypothesis*

Our analysis allowed us to test the dosage limit hypothesis, which predicts that highly expressed genes (i.e., those that are up-regulated in specific contexts) are under-represented on the X chromosome because of its haploid dose in males (Vicoso and Charlesworth 2009; Meisel *et al.* 2012a; Hurst *et al.* 2015). We determined that *D. melanogaster* genes that are either up- or down-regulated following biotic or abiotic treatments are frequently under-represented on the X chromosome (Figures 1–4). The paucity of down-regulated X-linked genes is inconsistent with the dosage limit hypothesis, which only predicts that up-regulated genes will be under-represented on the X chromosome (Table 1). Moreover, we frequently observed a deficiency of up-regulated X-linked genes in females, but not males, when data from both sexes were available (Figures 2 and 4). A female-specific paucity of X-linked up-regulated genes is also not predicted by the dosage limit hypothesis. Furthermore, there was an excess of X-linked up-regulated genes following cocaine treatment in both males and females (Figure 4), which is also inconsistent with the dosage limit hypothesis.

Our results thus provide strong evidence that the dosage limit hypothesis cannot
completely explain the unique gene content of the *Drosophila* X chromosome. However, we also
cannot explain the paucity of X-linked up-regulated genes based on proximity to an HAS alone
(Figure 5). A dosage limit may therefore be partially responsible for some aspects of the unique
gene content of the X chromosome. The dosage limit hypothesis may be especially important in
testis, where the haploid dose of the X chromosome does not appear to be compensated
(Meiklejohn *et al.* 2011). Specifically, the reduced dosage of the X chromosome could explain
the biased duplication of genes from the X to the autosomes, with the autosomal derived
paralogs expressed primarily in testis in order to compensate for under-expression of the X
(Betrán *et al.* 2002; Meisel *et al.* 2009, 2010). In addition, the dosage limit hypothesis is also
consistent with the observation that genes expressed specifically in the male accessory gland
are under-represented on the X chromosome (Swanson *et al.* 2001; Ravi Ram and Wolfner
2007; Meisel *et al.* 2012a). Therefore, both dosage limits and DCC binding may act in concert to
exclude up-regulated genes from the *Drosophila* X chromosome.

***Evidence for the DCC as a variance dampener that inhibits X-linked DE genes***
Our results add evidence in support of the hypothesis that the DCC is a variance
dampener (Lee *et al.* 2018), which prevents both up- and down-regulation of gene expression
on the *Drosophila* X chromosome. First, there is a significant deficiency of down-regulated
genes on the X chromosome, in addition to a paucity of X-linked up-regulated genes, in a
variety of treatments (Figures 2–4). Second, the X-linked genes that are up- or down-regulated
tend not to be bound by the DCC, and DCC-binding largely explains the X-autosome differences
in up-regulated, down-regulated, and DE genes (Figure 5). Lastly, the magnitude of differential
expression is correlated with distance from an HAS for both up- and down-regulated genes
(Figures 6 and 7). Because we observe similar results for both up- and down-regulated genes,
we hypothesize that the DCC is a variance dampener that inhibits both induction and repression
of gene expression (Table 1).
We further hypothesize that the variance dampening effect of the DCC can explain other
aspects of X chromosome gene content. For example, there is a paucity of genes with tissue-
specific expression on the *Drosophila* X chromosome (Mikhaylova and Nurminsky 2011; Meisel
*et al.* 2012a). The genes with tissue-specific expression that are on the X chromosome are less
likely to be DCC-bound than broadly expressed X-linked genes (Meisel *et al.* 2012b). Tissue-
specific expression can be thought of as a form of context-dependence in which the context is
developmental, rather than environmental. Therefore, the variance dampening effect of the DCC
may prevent tissue-specific induction, or repression, of gene expression.
There are at least two explanations for how the variance dampening effect of the DCC
could lead to a deficiency of genes with context-specific expression on the X chromosome. First,
in what we will call the mechanistic explanation, the DCC itself may affect gene expression in
the experiments from which we obtained data. While this mechanism is feasible for gene
expression in males, it is less obvious how it could explain the deficiency of X-linked DE genes
in females (Figures 2 and 4) because the DCC does not assemble in females (Lucchesi and
Kuroda 2015). The lack of a complete DCC in females occurs because the Msl-2 protein is only
expressed in males (Kelley *et al.* 1997). However, Mof, the component of the DCC responsible
for H4K16ac, is expressed in both sexes, and there is evidence that it serves as a variance
dampener in both males and females (Lee *et al.* 2018). In addition, the *Drosophila* X

chromosome has a distinct chromatin environment from the autosomes in females, including
histone marks associated with dosage compensation (Zhang and Oliver 2010). These chromatin
modifications, possibly mediated by Mof, may serve to dampen variance in X-linked gene
expression in both sexes, and therefore reduce differential expression on the X chromosome.
Furthermore, there is a complex interplay between heterochromatin and dosage compensation
in *Drosophila (Makki and Meller 2021),* which could also contribute to a variance dampening
effect in both sexes.
Alternatively, in what we call the evolutionary explanation, we may have observed an
evolved difference between the X and autosomes in the experimental data we analyzed. These
evolved differences would be a result of selection against genes with context-dependent gene
expression on the X chromosome as a result of the variance dampening effect of the DCC, or
perhaps only Mof. This explanation is attractive because it allows for selection in males only
(where the DCC assembles) to shape the gene expression profile of the X chromosome in both
sexes, where we observe the paucity of X-linked genes with context-dependent expression.
This is analogous to how a shared genome prevents the evolution of sexual dimorphism
because of inter-sexual phenotypic correlations (Rowe *et al.* 2018). Consistent with the
evolutionary explanation, DCC bound genes have slower evolving gene expression levels than
X-linked unbound genes (Meisel *et al.* 2012b), suggesting that the DCC may inhibit the
evolution of gene expression. Similarly, X-linked genes whose expression levels changed more
during a laboratory evolution experiment were further from an HAS (Abbott *et al.* 2020). It is
worth noting, however, that the mechanistic and evolutionary explanations are not mutually
exclusive, and additional work is necessary to evaluate how well each can explain the effect of
the DCC, or Mof in particular, on the unique gene content of the *Drosophila* X chromosome.
Regardless of the mode of action, the variance dampening hypothesis could also explain
the differences in gene content between mammalian and *Drosophila* X chromosomes. The
mammalian X has an excess of genes with tissue-specific expression (Lercher *et al.* 2003;
Meisel *et al.* 2012a), in contrast to the deficiency of genes with context-dependent expression
on the *D. melanogaster* X chromosome. A dosage limit hypothesis has been proposed to
explain the excess of narrowly expressed genes on the mammalian X chromosome because
broadly expressed genes have a higher maximal expression (Hurst *et al.* 2015). Evidence for
large-scale up-regulation of the mammalian X is much weaker than the evidence that the
*Drosophila* DCC up-regulates X-linked expression (Xiong *et al.* 2010; Pessia *et al.* 2012; Lin *et*
*al.* 2012; Deng *et al.* 2013; Mank 2013; Gu and Walters 2017; Deng and Disteche 2019).
Therefore, the taxon-specific peculiarities of dosage compensation could explain the differences
in X chromosome gene content between *Drosophila* and mammals. Specifically, only in
*Drosophila* is there evidence for a variance dampener that targets X-linked genes, whereas
dosage limits may be more pronounced in mammals.

**608 *Exceptions to the rules are also evidence for the DCC as a variance dampener***
We observed exceptions to the general relationships between DE genes, X-linkage, and
the DCC within specific treatments. For example, there was an excess of DE genes on the X
chromosome after cocaine or starvation treatments (Figure 4). In addition, some of the
correlations between $\log_2$FC and distance from an HAS were in opposite directions depending
on the treatment (Figures 6 and 7). We discuss these exceptions below, explain how they are
consistent with the variance dampener hypothesis despite the atypical patterns, and describe

how they can help us discriminate between the mechanistic and evolutionary explanations for
the reduced DE on the X chromosome.
*Excess of X-linked DE genes in brain:* In both of the brain samples analyzed (starvation
and cocaine treatments), we observed an excess of X-linked DE genes (Figure 4). The cocaine
treatment resulted in an excess of both up- and down-regulated genes, whereas the starvation
treatment only caused an excess of up-regulated genes. There is a similar enrichment of X-
linked genes with male-biased expression in *D. melanogaster* brain (Huylmans and Parsch
2015). Multiple genes that encode DCC proteins, including *msl-2* and *mle*, are highly expressed
in brain (Chintapalli *et al.* 2007; Straub *et al.* 2013; Vensko and Stone 2014; Huylmans and
Parsch 2015)*,* as are the non-coding RNAs that assemble with the DCC (Amrein and Axel 1997;
Meller *et al.* 1997)*.* If the DCC inhibits context-dependent differential expression, we may expect
fewer X-linked DE genes in the brain because of the high expression of the DCC components. It
is therefore surprising that we observe the opposite pattern in the brain.
Despite the excess of X-linked up-regulated genes in the brain after either starvation or
cocaine, we observed that DE genes in both treatments are less likely to be bound by the DCC,
consistent with most other treatments (Figure 5). Moreover, X-linked DE genes after cocaine
treatment were further from an HAS, similar to other treatments (Figure 7). These observations
are consistent with the hypothesis that the DCC is a variance dampener, but they do not explain
why the unbound X-linked genes are more likely to be DE in the brain than in other tissues.
One explanation for an excess of X-linked DE genes in the brain is the heterogeneity of
dosage compensation across the X chromosome in brain cells. Belyi et al. (2020) observed
greater position effects on brain gene expression than expression in non-brain head tissus for
transgenes inserted on the X chromosome. This suggests that the X chromosome is more of a
patchwork of DCC-bound and unbound regions in the brain than in other tissues. The unbound
regions may allow for context-dependent transcriptional regulation of X-linked genes in the brain
more than in other tissues where DCC-binding is more uniform.
*Excess of X-linked DE genes after cocaine treatment:* The excess of X-linked DE genes
after cocaine treatment points to a possible mechanism to explain the context-dependent
transcriptional regulation of X-linked genes unbound by the DCC. Cocaine affects chromatin in
neural cells by reversing trimethylation of histone H3 at lysine 9 (H3K9me3), a repressive
chromatin mark, leading to de-repression of genes and repetitive elements that would normally
be silenced (Kumar *et al.* 2005; Renthal *et al.* 2007; Maze *et al.* 2011; Covington *et al.* 2011).
We observed that the majority of X-linked genes that were up-regulated following cocaine
treatment in *D. melanogaster* brain were not bound by the DCC (Figure 5A), and previous work
showed that unbound genes are more likely to be associated with repressive chromatin marks
than DCC-bound genes (Meisel *et al.* 2012b). Therefore, the X-linked DE genes after cocaine
treatment are probably located in chromatin that was unbound by the DCC, and cocaine
induced a conversion from repressive to transcriptionally active chromatin. Importantly, this
interpretation is consistent with the hypothesis that the DCC is a variance dampener and X-
linked genes unbound by the DCC are more likely to have context-dependent expression.
Future work should evaluate if chromatin state was in fact altered in X-linked unbound genes in
neuronal cells following cocaine treatment.
*Genotypic differences help discriminate between mechanistic and evolutionary*
*explanations:* We observed different patterns following copper treatment depending on whether

the flies were sensitive or resistant. Many more X-linked genes were down-regulated in
sensitive flies, and this was accompanied by an equivalent enrichment of down-regulated
autosomal genes (Figure 4B). The down-regulated X-linked genes were extremely biased
toward those unbound by the DCC (Figure 5B), consistent with our hypothesis that the DCC is a
variance dampener. However, there was a difference between sensitive and resistant flies in the
correlation of $\log_2$FC and distance from an HAS. In sensitive flies, the correlation was positive
when we considered $|\log_2$FC$|$ or genes with $\log_2$FC$>0$ (Figure 7A–B), consistent with more
differential expression further from an HAS. In resistant flies, the correlation was positive for
genes with $\log_2$FC$<0$ (Figure 7C), suggesting that genes further from an HAS were less down-
regulated. One possible explanation for this discrepancy is that, because there were so few DE
genes in resistant flies, the correlation between $\log_2$FC and distance from an HAS differed from
other treatments (possibly because few genes have extreme negative $\log_2$FC values). This
explanation is not well supported because other treatments had even fewer DE genes (e.g.,
Zika virus and starvation), yet the correlations between $\log_2$FC and distance from an HAS in
those other treatments were in the same directions as in the treatments with more DE genes
(Figures 6 and 7). Alternatively, resistance to copper (or any treatment) may affect the
relationship between gene expression and the DCC.
Regardless of the causes of the differences between copper sensitive and resistant flies,
the comparison is informative for evaluating the mechanistic and evolutionary explanations for
the relationship between DE genes and DCC binding. Copper resistant flies are likely to
resemble genotypes that would emerge following adaptation to copper exposure. As described
above, X-linked DE genes in copper resistant flies were not necessarily further from an HAS
(Figure 7). In contrast, X-linked DE genes in copper sensitive flies were further from an HAS,
similar to what was observed in most other treatments (Figure 7). Therefore, adaptation may not
necessarily lead to the evolved relationship between X-linked DE genes and the DCC that we
observed, which could be interpreted as support for the mechanistic explanation for X-autosome
differences. However, there are clear evolved differences between the X and autosomes, such
as the absence of X-linked AMP genes (Hill-Burns and Clark 2009) and deficiency of X-linked
accessory gland expressed genes (Swanson *et al.* 2001; Ravi Ram and Wolfner 2007; Meisel *et*
*al.* 2012a), which cannot be explained by the mechanistic explanation. This leads us to
conclude that both mechanistic effects of the DCC (or Mof) and evolved X-autosome differences
are responsible for the paucity of X-linked DE genes. The evolved X-autosome differences are
likely the result of selection against X-linked genes with context-dependent expression in
response to both the variance dampening effects of the DCC/Mof and dosage limits of the
haploid X chromosome in males.

**MATERIALS AND METHODS**

***General statistical analysis***
The analyses and figure generation were performed in R (R Core Team 2019) using the
following packages: boot (Davison and Hinkley 1997; Canty and Ripley 2021), corpcor (Schäfer
and Strimmer 2005; Schafer *et al.* 2017), ggplot2 (Wickham 2016), ggridges (Wilke 2021), and
cowplot (Wilke 2020). Additional R packages were used for specific analyses, as described
below.

 ***RNA-seq data analysis***

We analyzed available RNA-seq data to test for differential expression between control
*D. melanogaster* and flies that received a bacterial, viral, or abiotic treatment (Supplementary
File 1). In one dataset, *D. melanogaster* adult males were infected with one of ten different
bacteria or a control treatment (Troha *et al.* 2018). For that experiment, we used RNA-seq data
from 12 hours post treatment for only live (not heat-killed) bacteria, which we downloaded as
fastq files from the NCBI sequence read archive (BioProject PRJNA428174). We only included
bacterial treatments with at least 50 DE genes relative to the control condition (see below for
methods used to identify DE genes), which was 5/10 treatments. In another data set, we
compared gene expression 8 h following injection of *Providencia rettgeri* with uninfected flies, in
males and females separately (Duneau *et al.* 2017). In a third data set, we analyzed the
response to infection at 19 different timepoints from 1 h to 120 h after injection of *E. coli*-derived
crude lipopolysaccharide (Schlamp *et al.* 2021). Other datasets include exposure to one of two
different viral infections, copper, starvation, radiation, and cocaine (Palmer *et al.* 2018; Prasad
and Hens 2018; Harsh *et al.* 2020; de Oliveira *et al.* 2021; Green *et al.* 2021; Baker *et al.* 2021).
A full list of accession numbers is provided in Supplementary File 1.

Raw RNA-seq reads were assigned to annotated *D. melanogaster* transcripts (r6.22)
using kallisto v0.44.0 (Bray *et al.* 2016). Read counts for all transcripts from each gene were
summed to obtain gene-level expression estimates, and the counts per gene were then rounded
to the nearest integer. For a given treatment, we only considered genes with at least 10 mapped
reads total across all replicates from control and treatment samples. The integer counts were
used as input into DESeq2 (Love *et al.* 2014), which we used to identify DE genes between the
treatment and control samples (see below). We performed a principal component analysis
(PCA) on regularized log transformed read counts to identify replicate samples that were
outliers relative to other replicates of the same sample type. We identified one outlier male
control replicate in the Kallithea virus data, which we excluded from all subsequent analyses.

We identified DE genes based on the false-discovery rate (FDR) corrected p-value ($p_{ADJ}$)
and $\log_2 FC$ of treatment to control expression. Genes were considered up-regulated in a
treatment if $p_{ADJ} < 0.05$ and $\log_2 FC > 1$ (i.e., a significant 2-fold increase upon treatment).
Similarly, genes were considered downregulated if $p_{ADJ} < 0.05$ and $\log_2 FC < -1$ (i.e., a significant
2-fold decrease upon treatment). DE genes are those that are either up- or down-regulated in a
given treatment (i.e., $p_{ADJ} < 0.05$ and $|\log_2 FC| > 1$). We only included datasets with at least 50
DE genes between treatment and control samples.

We also analyzed the bacterial infection data considering all Gram-negative or Gram-
positive bacteria from Troha *et al.* (2018) as a treatment group. In this case, we used a
statistical model that examined the effect of treatment (bacteria or control), with strain nested
within bacterial treatment. This was done separately for all Gram-negative bacteria and all
Gram-positive bacteria. DE genes, as well as up- and down-regulated genes, were identified
using the same $p_{ADJ}$ and $\log_2 FC$ criteria described above.

In cases where both female and male RNA-seq data were available (cocaine and
Kallithea virus), we performed two separate analyses focusing on the effect of the treatment
(i.e., not on the effect of sex) on gene expression. First, we analyzed the male and female data
together using a linear model that included the effect of treatment and the interaction of
treatment and sex. From this analysis, we extracted genes that were DE based on the treatment

effect. Second, we analyzed data from the two sexes separately using a model that only
included the effect of treatment (i.e., the same way we analyzed data from other treatments
without separate sex samples).
748           We did not analyze raw RNA-seq data (i.e., Illumina sequence reads) for two of the
infection datasets. First, we obtained previously identified DE genes from the time course
analysis of expression following infection (Schlamp *et al.* 2021). Second, raw data were not
available from an experiment in which male and female *D. melanogaster* were infected with
bacteria (Duneau *et al.* 2017), but processed data were available from FlySexsick-*seq*
(http://flysexsick.buchonlab.com). For those data, we compared gene expression between
unchallenged flies and 8 h following injection of *Providencia rettgeri*. Only expression levels
(and no p-values) were provided for these data, and we therefore considered genes to be
differentially expressed based on a variety of $\log_2$FC cutoffs.
757           We determined a null expectation for the number of X-linked DE genes by multiplying
758     the fraction of autosomal genes that are DE by the total number of X-linked genes with
759     expression measurements. Similar calculations were performed to determine a null expectation
760     for up- or down-regulated X-linked genes.

761 ### *DCC binding*
762           Data on DCC binding in the *D. melanogaster* genome was obtained from a published
chromatin immunoprecipitation followed by microarray (ChIP-chip) experiment in which genes
were classified as bound by the DCC in SL2 embryonic cells, clone 8 wing imaginal disc cells,
and embryos (Alekseyenko *et al.* 2006). For the purpose of our analysis, we considered a gene
to be bound by the DCC if it was bound in at least one of the three samples. We also obtained
HAS locations from two different published data sets (Alekseyenko *et al.* 2008; Straub *et al.*
2008). Using these HAS locations, we calculated the distance of each X chromosome gene to
the nearest HAS in nucleotides.

### *Tissue-specific expression*
771           We obtained microarray measurements of gene expression from 14 unique adult
772     *D. melanogaster* tissues from FlyAtlas (Chintapalli *et al.* 2007). Of the 14 tissues, 4 are sex-
773     specific (testis and accessory gland from males, and ovary and spermatheca from females), and
774     the remaining 10 tissues (brain, eye, thoracicoabdominal ganglion, salivary gland, crop, midgut,
775     malpighian tubule, hindgut, heart, and fat body) are shared by both males and females (i.e.,
776     non-sex-specific). We used these data to calculate expression breadth ($\tau$) for each gene:
777

778
$$\tau = \frac{\Sigma_{i=1}^{N} 1 - \frac{\log S_i}{\log S_{max}}}{N-1},$$

779
where $N$ is the number of tissues analyzed (10 for the non-sex-specific tissues, and 14 for all
adult tissues), $S_i$ is the gene expression level in tissue $i$ (measured as average signal intensity
for all microarray probes assigned to that gene), and $S_{max}$ is the maximum $S_i$ across all $N$
tissues (Yanai *et al.* 2005). All $S_i < 1$ were set to 1 so that $\log S_i \geq 0$, as done previously
(Larracuente *et al.* 2008; Meisel 2011; Meisel *et al.* 2012a). Values for spermatheca from mated
and unmated females were averaged to create a single $S_i$ for spermatheca (Meisel 2011). We
calculated $\tau$ separately for all 14 unique adult tissues and for the 10 non-sex-specific tissues.

We also identified the tissue where expression is highest for every gene that has $S_{max} > 100$
and where expression was detected for at least one probe in all 4 replicate arrays in that tissue.

## DATA AVAILABILITY

791 No new data were generated or analyzed in support of this research.

## ACKNOWLEDGEMENTS

793 We thank Wen-Juan Ma for feedback and suggestions on previous versions of this manuscript.
794 This work was completed in part with resources provided by the Research Computing Data
795 Core at the University of Houston.

## FUNDING

This material is based upon work supported by the National Science Foundation under Grant
No. DEB-1845686 to RPM; the United States Department of Agriculture under cooperative
agreement No. 58-3020-8-016 to RPM; National Institutes of Health under grant No. R01-
AI139154 to RLU; and the KU Center for Genomics. Any opinions, findings, and conclusions or
recommendations expressed in this material are those of the authors and do not necessarily
reflect the views of the National Science Foundation, National Institutes of Health, or United
States Department of Agriculture.

**References Cited**

Abbott J. K., A. K. Chippindale, and E. H. Morrow, 2020 The microevolutionary response to

male-limited X-chromosome evolution in Drosophila melanogaster reflects

macroevolutionary patterns. J. Evol. Biol. 33: 738–750.

Alekseyenko A. A., E. Larschan, W. R. Lai, P. J. Park, and M. I. Kuroda, 2006 High-resolution

ChIP–chip analysis reveals that the Drosophila MSL complex selectively identifies active

genes on the male X chromosome. Genes Dev 20: 848–857.

Alekseyenko A. A., S. Peng, E. Larschan, A. A. Gorchakov, O.-K. Lee, *et al.*, 2008 A sequence

motif within chromatin entry sites directs MSL establishment on the Drosophila X

chromosome. Cell 134: 599–609.

Amrein H., and R. Axel, 1997 Genes expressed in neurons of adult male Drosophila. Cell 88:

459–469.

Bachtrog D., N. R. T. Toda, and S. Lockton, 2010 Dosage compensation and demasculinization

of X chromosomes in Drosophila. Curr. Biol. 20: 1476–1481.

Bachtrog D., J. E. Mank, C. L. Peichel, M. Kirkpatrick, S. P. Otto, *et al.*, 2014 Sex determination:

why so many ways of doing it? PLoS Biol. 12: e1001899.

Baker B. M., S. S. Mokashi, V. Shankar, J. S. Hatfield, R. C. Hannah, *et al.*, 2021 The

Drosophila brain on cocaine at single-cell resolution. Genome Res.

https://doi.org/10.1101/gr.268037.120

Bean C. J., C. E. Schaner, and W. G. Kelly, 2004 Meiotic pairing and imprinted X chromatin

assembly in Caenorhabditis elegans. Nat. Genet. 36: 100–105.

Belyi A., E. Argyridou, and J. Parsch, 2020 The Influence of Chromosomal Environment on X-

Linked Gene Expression in Drosophila melanogaster. Genome Biol. Evol. 12: 2391–2402.

Betrán E., K. Thornton, and M. Long, 2002 Retroposed new genes out of the X in Drosophila.

Genome Res. 12: 1854–1859.

Bray N. L., H. Pimentel, P. Melsted, and L. Pachter, 2016 Near-optimal probabilistic RNA-seq

quantification. Nat. Biotechnol. 34: 525–527.

Brown C. J., A. Ballabio, J. L. Rupert, R. G. Lafreniere, M. Grompe, *et al.*, 1991 A gene from the

region of the human X inactivation centre is expressed exclusively from the inactive X

chromosome. Nature 349: 38–44.

Canty A., and B. D. Ripley, 2021 boot: Bootstrap R (S-Plus) Functions

Charlesworth B., J. A. Coyne, and N. H. Barton, 1987 The relative rates of evolution of sex

chromosomes and autosomes. Am. Nat. 130: 113–146.

Chintapalli V. R., J. Wang, and J. A. T. Dow, 2007 Using FlyAtlas to identify better Drosophila

melanogaster models of human disease. Nat. Genet. 39: 715–720.

Chow J. C., Z. Yen, S. M. Ziesche, and C. J. Brown, 2005 Silencing of the mammalian X

chromosome. Annu. Rev. Genomics Hum. Genet. 6: 69–92.

Covington H. E. 3rd, I. Maze, H. Sun, H. M. Bomze, K. D. DeMaio, *et al.*, 2011 A role for

repressive histone methylation in cocaine-induced vulnerability to stress. Neuron 71: 656–

670.

Davison A. C., and D. V. Hinkley, 1997 Bootstrap Methods and Their Applications

De Gregorio E., P. T. Spellman, G. M. Rubin, and B. Lemaitre, 2001 Genome-wide analysis of

the Drosophila immune response by using oligonucleotide microarrays. Proc. Natl. Acad.

Sci. U. S. A. 98: 12590–12595.

Deng X., J. B. Berletch, W. Ma, D. K. Nguyen, J. B. Hiatt, *et al.*, 2013 Mammalian X

upregulation is associated with enhanced transcription initiation, RNA half-life, and MOF-

mediated H4K16 acetylation. Dev. Cell 25: 55–68.

Deng X., and C. M. Disteche, 2019 Rapid transcriptional bursts upregulate the X chromosome.

Nat. Struct. Mol. Biol. 26: 851–853.

Duneau D. F., H. C. Kondolf, J. H. Im, G. A. Ortiz, C. Chow, *et al.*, 2017 The Toll pathway

underlies host sexual dimorphism in resistance to both Gram-negative and Gram-positive

bacteria in mated Drosophila. BMC Biol. 15: 124.

Emerson J. J., H. Kaessmann, E. Betrán, and M. Long, 2004 Extensive gene traffic on the

mammalian X chromosome. Science 303: 537–540.

Ferrari F., A. A. Alekseyenko, P. J. Park, and M. I. Kuroda, 2014 Transcriptional control of a

whole chromosome: emerging models for dosage compensation. Nat. Struct. Mol. Biol. 21:

118–125.

Fry J. D., 2010 The genomic location of sexually antagonistic variation: some cautionary

comments. Evolution 64: 1510–1516.

Gallach M., and E. Betrán, 2016 Dosage Compensation and the Distribution of Sex-Biased

Gene Expression in Drosophila: Considerations and Genomic Constraints. J. Mol. Evol. 82:

199–206.

Gelbart M. E., E. Larschan, S. Peng, P. J. Park, and M. I. Kuroda, 2009 Drosophila MSL

complex globally acetylates H4K16 on the male X chromosome for dosage compensation.

Nat. Struct. Mol. Biol. 16: 825–832.

Green L., S. Radio, G. E. Rech, J. Salces-Ortiz, and J. González, 2021 Natural variation in

copper tolerance in Drosophila melanogaster is shaped by transcriptional and physiological

changes in the gut. bioRxiv.

Gu L., and J. R. Walters, 2017 Evolution of Sex Chromosome Dosage Compensation in

Animals: A Beautiful Theory, Undermined by Facts and Bedeviled by Details. Genome Biol.

Evol. 9: 2461–2476.

Gu L., P. F. Reilly, J. J. Lewis, R. D. Reed, P. Andolfatto, *et al.*, 2019 Dichotomy of Dosage

Compensation along the Neo Z Chromosome of the Monarch Butterfly. Curr. Biol. 29:

4071–4077.e3.

Harsh S., Y. Fu, E. Kenney, Z. Han, and I. Eleftherianos, 2020 Zika virus non-structural protein

NS4A restricts eye growth in Drosophila through regulation of JAK/STAT signaling. Dis.

Model. Mech. 13: dmm040816.

Hill-Burns E. M., and A. G. Clark, 2009 X-linked variation in immune response in Drosophila

melanogaster. Genetics 183: 1477–1491.

Hurst L. D., A. T. Ghanbarian, A. R. R. Forrest, FANTOM consortium, and L. Huminiecki, 2015

The Constrained Maximal Expression Level Owing to Haploidy Shapes Gene Content on

the Mammalian X Chromosome. PLoS Biol. 13: e1002315.

Huylmans A. K., and J. Parsch, 2015 Variation in the X:Autosome Distribution of Male-Biased

Genes among Drosophila melanogaster Tissues and Its Relationship with Dosage

Compensation. Genome Biol. Evol. 7: 1960–1971.

Kelley R. L., J. Wang, L. Bell, and M. I. Kuroda, 1997 Sex lethal controls dosage compensation

in Drosophila by a non-splicing mechanism. Nature 387: 195–199.

Kelley R. L., V. H. Meller, P. R. Gordadze, G. Roman, R. L. Davis, *et al.*, 1999 Epigenetic

spreading of the Drosophila dosage compensation complex from roX RNA genes into

flanking chromatin. Cell 98: 513–522.

Kumar A., K.-H. Choi, W. Renthal, N. M. Tsankova, D. E. H. Theobald, *et al.*, 2005 Chromatin

remodeling is a key mechanism underlying cocaine-induced plasticity in striatum. Neuron

48: 303–314.

Larracuente A. M., T. B. Sackton, A. J. Greenberg, A. Wong, N. D. Singh, *et al.*, 2008 Evolution

of protein-coding genes in Drosophila. Trends Genet. 24: 114–123.

Lee H., D.-Y. Cho, D. Wojtowicz, S. T. Harbison, S. Russell, *et al.*, 2018 Dosage-Dependent

Expression Variation Suppressed on the Drosophila Male X Chromosome. G3 8: 587–598.

Lemaitre B., and J. Hoffmann, 2007 The host defense of Drosophila melanogaster. Annu. Rev.

Immunol. 25: 697–743.

Lercher M. J., A. O. Urrutia, and L. D. Hurst, 2003 Evidence that the human X chromosome is

enriched for male-specific but not female-specific genes. Mol. Biol. Evol. 20: 1113–1116.

Lifschytz E., and D. L. Lindsley, 1972 The role of X-chromosome inactivation during

spermatogenesis. Proc. Natl. Acad. Sci. U. S. A. 69: 182–186.

Lin F., K. Xing, J. Zhang, and X. He, 2012 Expression reduction in mammalian X chromosome

evolution refutes Ohno's hypothesis of dosage compensation. Proc. Natl. Acad. Sci. U. S.

A. 109: 11752–11757.

Love M., S. Anders, and W. Huber, 2014 Differential analysis of count data--the DESeq2

package. Genome Biol. 15: 10–1186.

Lucchesi J. C., W. G. Kelly, and B. Panning, 2005 Chromatin remodeling in dosage

compensation. Annu. Rev. Genet. 39: 615–651.

Lucchesi J. C., and M. I. Kuroda, 2015 Dosage compensation in Drosophila. Cold Spring Harb.

Perspect. Biol. 7. https://doi.org/10.1101/cshperspect.a019398

Lyon M. F., 1961 Gene action in the X-chromosome of the mouse (Mus musculus L.). Nature

190: 372–373.

Makki R., and V. H. Meller, 2021 When Down Is Up: Heterochromatin, Nuclear Organization

and X Upregulation. Cells 10. https://doi.org/10.3390/cells10123416

Mank J. E., 2013 Sex chromosome dosage compensation: definitely not for everyone. Trends

Genet. 29: 677–683.

Maze I., J. Feng, M. B. Wilkinson, H. Sun, L. Shen, *et al.*, 2011 Cocaine dynamically regulates

heterochromatin and repetitive element unsilencing in nucleus accumbens. Proc. Natl.

Acad. Sci. U. S. A. 108: 3035–3040.

Meiklejohn C. D., E. L. Landeen, J. M. Cook, S. B. Kingan, and D. C. Presgraves, 2011 Sex

Chromosome-Specific Regulation in the Drosophila Male Germline But Little Evidence for

Chromosomal Dosage Compensation or Meiotic Inactivation. PLoS Biol. 9: e1001126.

Meiklejohn C. D., and D. C. Presgraves, 2012 Little evidence for demasculinization of the

Drosophila X chromosome among genes expressed in the male germline. Genome Biol.

Evol. 4: 1007–1016.

Meisel R. P., M. V. Han, and M. W. Hahn, 2009 A complex suite of forces drives gene traffic

from Drosophila X chromosomes. Genome Biol. Evol. 1: 176–188.

Meisel R. P., B. B. Hilldorfer, J. L. Koch, S. Lockton, and S. W. Schaeffer, 2010 Adaptive

evolution of genes duplicated from the Drosophila pseudoobscura neo-X chromosome. Mol.

Biol. Evol. 27: 1963–1978.

Meisel R. P., 2011 Towards a more nuanced understanding of the relationship between sex-

biased gene expression and rates of protein-coding sequence evolution. Mol. Biol. Evol. 28:

1893–1900.

Meisel R. P., J. H. Malone, and A. G. Clark, 2012a Disentangling the relationship between sex-

biased gene expression and X-linkage. Genome Res. 22: 1255–1265.

Meisel R. P., J. H. Malone, and A. G. Clark, 2012b Faster-X evolution of gene expression in

Drosophila. PLoS Genet. 8: e1003013.

Meller V. H., K. H. Wu, G. Roman, M. I. Kuroda, and R. L. Davis, 1997 roX1 RNA paints the X

chromosome of male Drosophila and is regulated by the dosage compensation system.

Cell 88: 445–457.

Mikhaylova L. M., and D. I. Nurminsky, 2011 Lack of global meiotic sex chromosome

inactivation, and paucity of tissue-specific gene expression on the Drosophila X

chromosome. BMC Biol. 9: 29.

Mueller J. L., S. K. Mahadevaiah, P. J. Park, P. E. Warburton, D. C. Page, *et al.*, 2008 The

mouse X chromosome is enriched for multicopy testis genes showing postmeiotic

expression. Nat. Genet. 40: 794–799.

Mueller J. L., H. Skaletsky, L. G. Brown, S. Zaghlul, S. Rock, *et al.*, 2013 Independent

specialization of the human and mouse X chromosomes for the male germ line. Nat. Genet.

45: 1083–1087.

Oliveira D. S. de, M. T. Rosa, C. Vieira, and E. L. S. Loreto, 2021 Oxidative and radiation stress

induces transposable element transcription in Drosophila melanogaster. J. Evol. Biol. 34:

628–638.

Palmer W. H., N. C. Medd, P. M. Beard, and D. J. Obbard, 2018 Isolation of a natural DNA virus

of Drosophila melanogaster, and characterisation of host resistance and immune

responses. PLoS Pathog. 14: e1007050.

Parisi M., R. Nuttall, D. Naiman, G. Bouffard, J. Malley, *et al.*, 2003 Paucity of genes on the

Drosophila X chromosome showing male-biased expression. Science 299: 697–700.

Pessia E., T. Makino, M. Bailly-Bechet, A. McLysaght, and G. A. B. Marais, 2012 Mammalian X

chromosome inactivation evolved as a dosage-compensation mechanism for dosage-

sensitive genes on the X chromosome. Proc. Natl. Acad. Sci. U. S. A. 109: 5346–5351.

Potrzebowski L., N. Vinckenbosch, A. C. Marques, F. Chalmel, B. Jégou, *et al.*, 2008

Chromosomal gene movements reflect the recent origin and biology of therian sex

chromosomes. PLoS Biol. 6: e80.

Prasad N., and K. Hens, 2018 Sugar Promotes Feeding in Flies via the Serine Protease

Homolog scarface. Cell Rep. 24: 3194–3206.e4.

Ravi Ram K., and M. F. Wolfner, 2007 Seminal influences: Drosophila Acps and the molecular

interplay between males and females during reproduction. Integr. Comp. Biol. 47: 427–445.

R Core Team, 2019 R: A Language and Environment for Statistical Computing

Renthal W., I. Maze, V. Krishnan, H. E. Covington 3rd, G. Xiao, *et al.*, 2007 Histone deacetylase

5 epigenetically controls behavioral adaptations to chronic emotional stimuli. Neuron 56:

517–529.

Rice W. R., 1984 SEX CHROMOSOMES AND THE EVOLUTION OF SEXUAL DIMORPHISM.

Evolution 38: 735–742.

Rowe L., S. F. Chenoweth, and A. F. Agrawal, 2018 The Genomics of Sexual Conflict. Am. Nat.

192: 274–286.

Sackton T. B., B. P. Lazzaro, T. A. Schlenke, J. D. Evans, D. Hultmark, *et al.*, 2007 Dynamic

evolution of the innate immune system in Drosophila. Nat. Genet. 39: 1461–1468.

Schäfer J., and K. Strimmer, 2005 A shrinkage approach to large-scale covariance matrix

estimation and implications for functional genomics. Stat. Appl. Genet. Mol. Biol. 4:

Article32.

Schafer J., R. Opgen-Rhein, V. Zuber, M. Ahdesmaki, A. P. D. Silva, *et al.*, 2017 corpcor:

Efficient Estimation of Covariance and (Partial) Correlation

Schlamp F., S. Y. N. Delbare, A. M. Early, M. T. Wells, S. Basu, *et al.*, 2021 Dense time-course

gene expression profiling of the Drosophila melanogaster innate immune response. BMC

Genomics 22: 304.

Straub T., C. Grimaud, G. D. Gilfillan, A. Mitterweger, and P. B. Becker, 2008 The chromosomal

high-affinity binding sites for the Drosophila dosage compensation complex. PLoS Genet.

4: e1000302.

Straub T., A. Zabel, G. D. Gilfillan, C. Feller, and P. B. Becker, 2013 Different chromatin

interfaces of the Drosophila dosage compensation complex revealed by high-shear ChIP-

seq. Genome Res. 23: 473–485.

Sturgill D., Y. Zhang, M. Parisi, and B. Oliver, 2007 Demasculinization of X chromosomes in the

Drosophila genus. Nature 450: 238–241.

Swanson W. J., A. G. Clark, H. M. Waldrip-Dail, M. F. Wolfner, and C. F. Aquadro, 2001

Evolutionary EST analysis identifies rapidly evolving male reproductive proteins in

Drosophila. Proc. Natl. Acad. Sci. U. S. A. 98: 7375–7379.

Troha K., J. H. Im, J. Revah, B. P. Lazzaro, and N. Buchon, 2018 Comparative transcriptomics

reveals CrebA as a novel regulator of infection tolerance in D. melanogaster. PLoS Pathog.

14: e1006847.

Turner J. M. A., 2007 Meiotic sex chromosome inactivation. Development 134: 1823–1831.

Turner J. M. A., 2015 Meiotic Silencing in Mammals. Annu. Rev. Genet. 49: 395–412.

Vensko S. P. 2nd, and E. A. Stone, 2014 No evidence for a global male-specific lethal complex-

mediated dosage compensation contribution to the demasculinization of the Drosophila

melanogaster X chromosome. PLoS One 9: e103659.

Vibranovski M. D., H. F. Lopes, T. L. Karr, and M. Long, 2009 Stage-specific expression

profiling of Drosophila spermatogenesis suggests that meiotic sex chromosome inactivation

drives genomic relocation of testis-expressed genes. PLoS Genet. 5: e1000731.

Vicoso B., and B. Charlesworth, 2009 The deficit of male-biased genes on the D. melanogaster

X chromosome is expression-dependent: a consequence of dosage compensation? J. Mol.

Evol. 68: 576–583.

Wang P. J., J. R. McCarrey, F. Yang, and D. C. Page, 2001 An abundance of X-linked genes

expressed in spermatogonia. Nat. Genet. 27: 422–426.

Wickham H., 2016 *ggplot2: Elegant Graphics for Data Analysis*. Springer-Verlag New York.

Wilke C. O., 2020 cowplot: Streamlined Plot Theme and Plot Annotations for "ggplot2"

Wilke C. O., 2021 ggridges: Ridgeline Plots in "ggplot2"

Wolfner M. F., H. A. Harada, M. J. Bertram, T. J. Stelick, K. W. Kraus, *et al.*, 1997 New genes

for male accessory gland proteins in Drosophila melanogaster. Insect Biochem. Mol. Biol.

27: 825–834.

Xiong Y., X. Chen, Z. Chen, X. Wang, S. Shi, *et al.*, 2010 RNA sequencing shows no dosage

compensation of the active X-chromosome. Nat. Genet. 42: 1043–1047.

Yanai I., H. Benjamin, M. Shmoish, V. Chalifa-Caspi, M. Shklar, *et al.*, 2005 Genome-wide

midrange transcription profiles reveal expression level relationships in human tissue

specification. Bioinformatics 21: 650–659.

Zhang Y., and B. Oliver, 2010 An evolutionary consequence of dosage compensation on

Drosophila melanogaster female X-chromatin structure? BMC Genomics 11: 6.

1032

**Supplemental Figures**

1034
1035        **Supplemental Figure S1.** The percent of genes on each chromosome arm that are differentially
1036       expressed following infection with one of five bacteria in male *D. melanogaster* is shown. Genes are
1037           either (A) up-regulated, (B) downregulated, or (C) differentially expressed (sum of up- and down-
1038     regulated). The percent of differentially expressed genes across the entire genome is shown as a dashed
1039       line. Asterisks indicate chromosomes where the percent of genes is significantly different from the rest of
1040                the genome (*$P<0.05$, **$P<0.005$, or ***$P<0.0005$ in Fisher's exact test).
1041

1042
**Supplemental Figure S2.** The percent of genes on each chromosome arm that are differentially
expressed following infection with at least 1, 2, 3, or 4 different bacteria in male *D. melanogaster* is
shown. Genes are either (A) up-regulated, (B) downregulated, or (C) differentially expressed (sum of up-
and down-regulated). The percent of differentially expressed genes across the entire genome is shown as
a dashed line. Asterisks indicate chromosomes where the percent of genes is significantly different from
the rest of the genome (*$P<0.05$, **$P<0.005$, or ***$P<0.0005$ in Fisher's exact test).
1049

1050

1051    **Supplemental Figure S3.** The rank order of chromosome arms is shown according to the percent of
1052    differentially expressed genes in each of five different bacterial treatments. The chromosome arm that
1053    has the highest percent of differentially expressed genes is ranked as 5, and the chromosome with the
1054    lowest percent is ranked as 1. Genes are either (A) up-regulated, (B) downregulated, or (C) differentially
1055    expressed (sum of up- and down-regulated). The mean ranks for each chromosome are shown as black
1056                                            dots and solid lines.

1057
**Supplemental Figure S5.** The rank order of chromosome arms is shown according to the percent of
differentially expressed genes in at least one, two, three, or four bacterial treatments. The chromosome
arm that has the highest percent of differentially expressed genes is ranked as 5, and the chromosome
with the lowest percent is ranked as 1. Genes are either (A) up-regulated, (B) downregulated, or (C)
differentially expressed (sum of up- and down-regulated). The mean ranks for each chromosome are
shown as black dots and solid lines.

1064

1065 **Supplemental Figure S5.** Distributions of log$_2$FC following bacterial infection for X-linked (X) and

1066 autosome (A) genes in *D. melanogaster* males. Asterisks show significant differences between X and

1067 autosomes within a treatment (**P<0.005; *****P<0.000005; Mann-Whitney test).

1068

1069

**Supplemental Figure S6.** The percent of genes on each chromosome arm that are differentially
expressed following viral and abiotic treatments is shown. Bars are colored by the sex of the flies used in
each treatment (see legend). Genes are either (A) up-regulated, (B) downregulated, or (C) differentially
expressed (sum of up- and down-regulated). The percent of differentially expressed genes across the
entire genome is shown as a dashed line. Asterisks indicate chromosomes where the percent of genes is
significantly different from the rest of the genome (*$P$<0.05, **$P$<0.005, or ***$P$<0.0005 in Fisher's exact
test).
1077
1078

1079
1080      **Supplemental Figure S7.** The rank order of chromosome arms is shown according to the percent of
1081     differentially expressed genes in viral and abiotic treatments. The chromosome arm that has the highest
1082     percent of differentially expressed genes is ranked as 5, and the chromosome with the lowest percent is
1083    ranked as 1. Genes are either (A) up-regulated, (B) downregulated, or (C) differentially expressed (sum of
1084     up- and down-regulated). The mean ranks for each chromosome are shown as black dots and solid lines.

1085
1086　**Supplemental Figure S8.** The percent of X-linked DCC-bound (black) and unbound (gray) genes that are
1087　up-regulated (A), down-regulated (B), or differentially expressed (C) are shown for each treatment and
1088　sample type (in parentheses). Asterisks show a significant difference in the percent between DCC-bound
1089　and unbound genes (*$P<0.05$ in Fisher's exact test).
1090

1091
1092
**Supplemental Figure S9.** Partial correlations between distance from a DCC high affinity site (HAS),
tissue-specificity (tau), and $\log_2$FC (LFC) are shown for each treatment. Error bars show the 95%
confidence interval from 1,000 bootstrap replicate samples of X-linked genes. The analysis was
performed on absolute values of $\log_2$FC of all genes (A), genes with positive $\log_2$FC>0 (B), or genes with
negative $\log_2$FC<0 (C). Tissue-specificity values ($\tau$) were calculated using all 14 adult tissues. HAS were
taken from Alekseyenko et al. (2008).
1099

1100
1101
**Supplemental Figure S10.** Partial correlations between distance from a DCC high affinity site (HAS),
tissue-specificity (tau), and $\log_2$FC (LFC) are shown for each treatment. Error bars show the 95%
confidence interval from 1,000 bootstrap replicate samples of X-linked genes. The analysis was
performed on absolute values of $\log_2$FC of all genes (A), genes with positive $\log_2$FC>0 (B), or genes with
negative $\log_2$FC<0 (C). Tissue-specificity values ($\tau$) were calculated using all 14 adult tissues. HAS were
taken from Straub et al. (2008).
1108

1109
1110
1111     **Supplemental Figure S11.** Partial correlations between distance from a DCC high affinity site (HAS),
1112         tissue-specificity (tau), and $\log_2$FC (LFC) are shown for each treatment. Error bars show the 95%
1113         confidence interval from 1,000 bootstrap replicate samples of X-linked genes. The analysis was
1114   performed on absolute values of $\log_2$FC of all genes (A), genes with positive $\log_2$FC>0 (B), or genes with
1115     negative $\log_2$FC<0 (C). Tissue-specificity values ($\tau$) were calculated using 10 non-sex-specific tissues.
1116                   HAS were taken from Alekseyenko et al. (2008).
1117

1118
1119
1120    **Supplemental Figure S12.** Partial correlations between distance from a DCC high affinity site (HAS),
1121       tissue-specificity (tau), and $log_2FC$ (LFC) are shown for each treatment. Error bars show the 95%
1122          confidence interval from 1,000 bootstrap replicate samples of X-linked genes. The analysis was
1123    performed on absolute values of $log_2FC$ of all genes (A), genes with positive $log_2FC>0$ (B), or genes with
1124       negative $log_2FC<0$ (C). Tissue-specificity values ($\tau$) were calculated using 10 non-sex-specific-tissues.
1125                                  HAS were taken from Straub et al. (2008).
1126

1127
1128
**Supplemental Figure S13.** Partial correlations between $\log_2$ fold-change in expression between
treatment and control ($\log_2$FC) and distance from a dosage compensation complex high affinity site
(HAS). Partial correlations were calculated based on rank order correlations between $\log_2$FC, distance
from an HAS, and tissue expression breadth. Each dot shows the partial correlation between $\log_2$FC and
distance from an HAS, with the error bars representing 95% confidence intervals from 1,000 bootstrap
replicates of the data. The X-axis shows the specific treatment. Dots and error bars are colored based on
the sex of the flies used in the experiment (see legend). HAS were obtained from the Straub *et al.* (2008)
data set. Expression breadth was calculated using microarray data from either 14 unique adult tissues
(left) or 10 adult tissues that are not sex-specific (right). Partial correlations are plotted with $|\log_2$FC$|$
values for all genes (A), only genes with $\log_2$FC>0 (B), and only genes with $\log_2$FC<0 (C).
1139