1 **Unambiguous detection of SARS-CoV-2 subgenomic mRNAs with single cell RNA**

2 **sequencing**

3 Phillip Cohen[1], Emma J DeGrace[1], Oded Danziger[1], Roosheel S Patel[1], Brad R

4 Rosenberg[1]#

5 [1]Department of Microbiology, Icahn School of Medicine at Mount Sinai, New York, NY

6 10035

7 #Address correspondence to Brad R Rosenberg, brad.rosenberg@mssm.edu

9 **Abstract**

Single cell RNA sequencing (scRNAseq) studies have provided critical insight

into the pathogenesis of Severe Acute Respiratory Syndrome CoronaVirus 2 (SARS-

CoV-2), the causative agent of COronaVIrus Disease 2019 (COVID-19). scRNAseq

workflows are generally designed for the detection and quantification of eukaryotic host

mRNAs and not viral RNAs. The performance of different scRNAseq methods to study

SARS-CoV-2 RNAs has not been thoroughly evaluated. Here, we compare different

scRNAseq methods for their ability to quantify and detect SARS-CoV-2 RNAs with a

focus on subgenomic mRNAs (sgmRNAs), which are produced only during active viral

replication and not present in viral particles. We present a data processing strategy,

single cell CoronaVirus sequencing (scCoVseq), which quantifies reads unambiguously

assigned to sgmRNAs or genomic RNA (gRNA). Compared to standard 10X Genomics

Chromium Next GEM Single Cell 3′ (10X 3′) and Chromium Next GEM Single Cell

V(D)J (10X 5′) sequencing, we find that 10X 5′ with an extended R1 sequencing

strategy maximizes the unambiguous detection of sgmRNAs by increasing the number

of reads spanning leader-sgmRNA junction sites. Differential gene expression testing

and KEGG enrichment analysis of infected cells compared with bystander or mock cells

showed an enrichment for COVID19-associated genes, supporting the ability of our

method to accurately identify infected cells. Our method allows for quantification of

coronavirus sgmRNA expression at single-cell resolution, and thereby supports high

resolution studies of the dynamics of coronavirus RNA synthesis.

Importance

Single cell RNA sequencing (scRNAseq) has emerged as a valuable tool to study

host-viral interactions particularly in the context of COronaVIrus Disease-2019 (COVID-

19). scRNAseq has been developed and optimized for analyzing eukaryotic mRNAs,

and the ability of scRNAseq to measure RNAs produced by Severe Acute Respiratory

Syndrome Coronavirus 2 (SARS-CoV-2) has not been fully characterized. Here we

compare the performance of different scRNAseq methods to detect and quantify SARS-

CoV-2 RNAs and develop an analysis workflow to specifically quantify unambiguous

reads derived from SARS-CoV-2 genomic RNA and subgenomic mRNAs. Our work

demonstrates the strengths and limitations of scRNAseq to measure SARS-CoV-2 RNA

and identifies experimental and analytical approaches that allow for SARS-CoV-2 RNA

detection and quantification. These developments will allow for studies of coronavirus

RNA biogenesis at single-cell resolution to improve our understanding of viral

pathogenesis.

Introduction

Severe Acute Respiratory Syndrome CoronaVirus 2 (SARS-CoV-2) is the causative agent of COronaVIrus Disease-2019 (COVID-19), which as of November 2021 has caused over 250 million cases and over 5 million deaths globally(1, 2). Global efforts to understand the pathogenesis of SARS-CoV-2 infection have led to the development of vaccines and antivirals, which have reduced morbidity and mortality(3). "Omics" methods have been instrumental in studying SARS-CoV-2 in part because they have generated large amounts of data regarding host-viral interactions at unprecedented speed(4–15). Single-cell RNA sequencing (scRNAseq) studies in particular have been used to study viral tropism(16–22), peripheral immune changes(23–33), transcriptional changes induced by infection(34, 35), and to develop cell atlases of COVID-19 pathology(23, 24, 36, 37). Of note, most scRNAseq workflows have been developed and optimized for studies of eukaryotic transcription but not viral, specifically SARS-CoV-2, transcription. The performance of different scRNAseq methods to detect and quantify viral RNAs may impact the analysis and interpretation of such studies.

SARS-CoV-2 is a betacoronavirus with a 29 kB positive-sense, single stranded RNA genome(38, 39). SARS-CoV-2 generates genomic RNA (gRNA), subgenomic mRNAs (sgmRNAs), and negative-sense antigenomic RNA during active infection(40, 41). Both gRNA and sgmRNAs are poly-adenylated, which enables detection by scRNAseq protocols that rely on poly-T primed reverse transcription(39–41). Translation of gRNA results in the production of one of two polyproteins, pp1a and pp1ab, which are

subsequently cleaved into an array of non-structural proteins involved in pathogenesis

and replication(39, 41). Translation of sgmRNAs generates structural and accessory

viral proteins critical for virion production and pathogenesis(39, 41). sgmRNAs are

produced only in cells with actively replicating virus, while gRNA is present in both

infected cells and virions(40, 41). Therefore, specific detection of sgmRNAs can allow

for: 1) specific identification of cells with actively replicating virus and 2) analysis of the

dynamics of viral gene expression within and across cells and viruses.

sgmRNAs are generated by discontinuous transcription events during negative

strand synthesis(40). Transcription Regulatory Sequences (TRS), present in the 5′

leader sequence of the virus (TRS-L) and upstream of each ORF body (TRS-B),

regulate this process(40). Template switching of the viral polymerase from a TRS-B to a

TRS-L generates sgmRNAs with the 5′ leader sequence fused to the sgmRNA ORF

body (**Figure 1A**)(40). These "nested" sgmRNAs share the viral ORF sequence

downstream of the junction site in addition to a common leader sequence upstream of

the junction site(40). This redundancy poses a challenge for standard scRNAseq data

processing pipelines because reads mapping to redundant sgmRNA sequences are

categorized as "ambiguous" and typically excluded from quantification. This problem

has been addressed in bulk RNAseq by quantifying SARS-CoV-2 reads spanning

leader-ORF junctions, which unambiguously identify sgmRNAs(12, 15). However, many

scRNAseq methods do not sequence this region of sgmRNAs at significant coverage

due to differences in library format and configuration of sequencing reads.

We hypothesized that both experimental (i.e. scRNAseq method) and data

processing decisions influence the ability to detect, resolve, and quantify SARS-CoV-2

RNA species with scRNAseq. We developed a data processing workflow, single cell

coronavirus sequencing (scCoVseq), to quantify only unambiguous SARS-CoV-2 reads

in scRNAseq data. We found that SARS-CoV-2 RNA detection differed by 10X

Genomics Chromium scRNAseq method, due in part to ambiguity of the library

fragments generated by each method. We show that 10X Chromium Next GEM Single

Cell V(D)J (10X 5′) scRNAseq with an extended read 1 (R1) sequencing strategy

maximized unambiguous SARS-CoV-2 reads and thereby increased detection of SARS-

CoV-2 RNAs. Using this method, we identify infected and uninfected "bystander" cells

within the same culture and determine differentially expressed genes between infected,

bystander, and mock cells.

Materials and Methods

Cell lines and Viral Infection

Vero-E6 cells (ATCC, CRL-1586) were maintained in Dulbecco's Modified Eagle

Medium (DMEM, Corning #10-017-CV) supplemented with 10% fetal-bovine serum

(FBS) and 1% Penicillin Streptomycin (PSN, Fisher scientific #15-140-122), and

routinely cultured at 37° C with 5% CO_2.

SARS-CoV-2 (isolate USA-WA1/2020, BEI resource NR-52281) and control media

(mock infected) stocks were grown by inoculating a confluent T175 flask of Vero-E6 cells

(passage 2). Mock and SARS-CoV-2 infected cultures were maintained in reduced serum

DMEM (2% FBS) for 72 hours, after which culture media was collected and filtered by

centrifugation (8000 x g, 15 minutes) using an Amicon Ultra-15 filter unit with a 100KDa

cutoff filter (Millipore # UFC910024). Concentrated stocks in reduced-serum media (2%

FBS), supplemented with 50mM HEPES buffer (Gibco #15630080) were stored at -80°C.

Viral titers were determined by plaque assay as previously described(42). All SARS-CoV-

2 propagations and experiments were performed in a Biosafety Level 3 facility in

compliance with institutional protocols and federal guidelines.

scRNAseq

For scRNAseq experiments, Vero-E6 cells in 6 well plates were infected with

SARS-CoV-2 at a MOI of 0.1, or with an equivalent volume of control media, in reduced-

serum media (2% FBS) for 24 hours. To prepare cells for scRNAseq, mock and SARS-

CoV-2 infected cultures were washed with calcium/magnesium-free PBS and

disassociated using TrypLE (Gibco # 12605010, 5 minutes at 37° C), after which

samples were centrifuged (200 x g, 5 minutes), resuspended in calcium/magnesium-

free PBS supplemented with 1% BSA, and counted. Mock and SARS-CoV-2 infected

cell culture samples were filtered through a 40μm FlowMi strainer (ScienceWare #

H13680-0040) and counted prior to loading on the 10X Genomics Chromium Controller

according to manufacturer's protocol. Mock and infected samples were loaded on

separate lanes of a 10X Genomics Chromium Controller for either NextGEM Single Cell

3′ v3.1 (10X 3′), or NextGEM Single Cell V(D)J v1.1, (10X 5′).

Gene expression libraries were prepared for 10X 3′ and 5′ samples according to

manufacturer's guidelines. Final 10X 3′ mock and infected gene expression libraries

and the 10X 5′ infected gene expression library were PCR amplified for 16 cycles while

the 10X 5′ mock gene expression library was amplified for 14 cycles. 10X 3′ gene

expression libraries were pooled and sequenced by short-read sequencing on an

Illumina NextSeq 500 using a high output 150 cycle reaction kit according to

manufacturers' protocol with the following read lengths: read 1 28 nt; i7 index 8 nt; and

read 2 130 nt. 10X 5′ gene expression libraries were also pooled and sequenced with

10X recommended read lengths (read 1 26 nt; i7 index 8 nt; and read 2 132 nt) or with

extended R1 protocol (read 1 158 nt; i7 index 8 nt; no read 2).

scRNAseq Pre-Processing

Conversion of Illumina BCL files to fastq

Fastq files for standard sequencing 3′ and 5′ gene expression libraries were

generated using the mkfastq command in cellranger v.3.1.0 (10X Genomics). Fastq files

for 5′ libraries sequenced with the extended R1 strategy were generated using bcl2fastq

v2.20.0 (Illumina, Inc). Extended R1 fastqs were then separated into pseudo R1 fastqs,

containing the cell barcode and UMI, and pseudo read 2 (R2) fastqs, containing cDNA

sequence, using a customized Python/3.7.3 script (available at github link pending) as

follows. The cell barcode and UMI are selected from the first 26 bp of R1. The

subsequent 13 bp derive from the template switch oligonucleotide and are ignored. The

remaining nucleotides (and corresponding quality scores) are reverse complemented

and stored as pseudo R2. The read header of the pseudo R2 fastqs are modified to

reflect the format for standard R2 fastqs.

Downsampling fastqs to control for sequencing depth

To control for differences in sequencing depth for each library, read depth per

library was downsampled to approximately 50,000 reads per cell. To generate a

whitelist of cell barcodes for downsampling while accounting for transcriptional

shutdown in SARS-CoV-2 infected cells (35), we generated preliminary gene x cell

matrices for our dataset using cellranger/3.1.0 count (10X Genomics, Inc) to quantify

and align reads to a host reference (African Green Monkey, ChlSab1.1) combined with

SARS-CoV-2 transcripts as annotated by the NCBI SARS-CoV-2 reference

(NC_045512.2) with modifications for USA/WA01 strain for each dataset. The resulting

output was analyzed in R/4.0.4 with Seurat/4.0.1(43–45) to filter out putative doublets

and empty droplets according to total UMIs/cell, number of genes/cell, and percent of

mitochondrial gene expression. After filtering, putative cell-containing cell barcodes

were output to a whitelist per library. Based on the these whitelists, the initial fastq files

were downsampled using seqtk (version 1.2)(46) to a total read depth of 50,000

multiplied by the number of cells in the library.

Preparation of empirically derived SARS-CoV-2 genome reference

Downsampled fastq files were then mapped using cellranger count/3.1.0 (10X

Genomics, Inc) to an empirically defined reference of SARS-CoV-2 sgmRNAs derived

from previously reported SARS-CoV-2 (BetaCoV/Korea/KCDC03/2020) RNAs

sequenced with long-read direct RNA Nanopore sequencing(12). These were

downloaded from the UCSC Genome Browser Table Browser(47) after filtering for TRS-

dependent transcripts and score > 900 and exporting to gtf format. Transcripts for

previously unknown ORFs were excluded from the annotation. An additional annotation

for genomic RNA was included which covered the entire length of the SARS-CoV-2

genome. Aligning the BetaCoV/Korea/KCDC03/2020 genome with USA/WA-CDC-

WA1/2020 genome showed that the USA/WA-CDC-WA1/2020 had an additional 21 3′

adenosine nucleotides annotated. To account for this in our reference, we extended any

annotations from BetaCoV/Korea/KCDC03/2020 that ended at the 3′ end of the genome

by an additional 21 bases. This SARS-CoV-2 reference was appended to the host

ChlSab1.1 Ensembl reference.

scCoVseq

To unambiguously assign and quantify scRNAseq reads to SARS-CoV-2 RNAs,

the cellranger output BAM was filtered for reads mapping to SARS-CoV-2 or ChlSab1.1

references using samtools (version 1.11)(48). SARS-CoV-2 aligned reads were then

subset to likely genomic RNA reads or sgmRNA reads. Genomic reads were defined as

those containing no gaps in their alignment and mapping upstream of the start of the

most 5′ sgmRNA, S. sgmRNA reads were defined as SARS-CoV-2 reads containing a

gap and mapping in part to the 5′ leader sequence, defined as the 5′ proximal 80

nucleotides of the SARS-CoV-2 genome, and in part 3′ to the start of S. All other reads

mapping to the SARS-CoV-2 genome were discarded. Reads passing these filtering

steps were quantified with umi_tools (version 1.0.0)(49). An R/3.5.3 script using the

Matrix (version 1.2-18)(50) and readr (version 1.3.1)(51) packages was used to convert

this to a sparse matrix and save as an rds file to decrease file size. UMIs that were

assigned to multiple genes were removed from the resulting matrix during analysis.

scRNAseq Analysis

Sashimi Plots

Reads from 10X 3′, 10X 5′, and 10X 5′ extended R1 data that were aligned to

the SARS-CoV-2 reference by cellranger were subset from the cellranger output BAM

file. Each BAM file was downsampled to approximately 1 x 10^6 reads to control for

differences in sequencing depth across libraries. Sashimi plots were generated with

ggsashimi (version 1.0.0)(52).

Classification of SARS-CoV-2 Infected Cells

scCoVseq-derived gene by cell matrices were loaded into R/4.0.4 and analyzed

with the Seurat/4.0.1(43–45) package. For each 10X method, mock and infected gene x

cell matrices were merged with the Seurat merge command. Scaled SARS-CoV-2 UMI

expression of 600 sampled cells were clustered with five methods (k means clustering,

hierarchical/Ward clustering, DIANA, mixture model-based clustering, and k medoids

clustering) using the clValid (version 0.7) package(53). Based on optimal performance

as measured by average distance, average distance between means, average

proportion of non-overlap, connectivity, Dunn index, figure of merit, and silhouette width,

k-medoids clustering implemented with the PAM algorithm and k set to 2 optimally

separated infected from uninfected cells. Therefore to identify infected and bystander

cells within SARS-CoV-2 treated cultures, euclidean distance between the z-scaled

expression of SARS-CoV-2 sgmRNA UMIs per cell was clustered using pam (k = 2)

implemented in the cluster (version 2.1.2) package(54). Output clusters were then

compared for viral UMI expression per cell, and the cluster with more viral UMIs was

classified as infected and the other as uninfected.

Comparison of SARS-CoV-2 RNA UMIs per scRNAseq Method

To examine the distribution of SARS-CoV-2 UMIs per cell by scRNAseq method,

the 25th percentile of total UMIs was quantified for all infected cells from each 10X

method. Any cells with fewer UMIs than the minimum 25th percentile of all samples were

discarded, and all cells were subsequently downsampled to this same number of total

UMIs/cell using the Seurat SampleUMI command. Each dataset was randomly

downsampled to the same number of infected cells to equalize for differences in cell

numbers, and viral sgmRNA UMIs/cell were plotted by scRNAseq method.

SARS-CoV-2 Read Distribution by scRNAseq Method

SARS-CoV-2 reads were defined as genomic or subgenomic using scCoVseq.

Reads aligning to the SARS-CoV-2 reference that were excluded from scCoVseq were

classified as ambiguous. The number of genomic, subgenomic, or ambiguous reads per

million SARS-CoV-2 reads was calculated and plotted for each scRNAseq method.

Differential Expression Analysis

To explore expression differences between infected, bystander, and mock cells,

differential expression testing with edgeR (version 3.32.1) was performed with

modifications for scRNAseq as previously described(55, 56). Viral genes were excluded

from analysis, and only host genes expressed in at least 10% of cells were tested. To

account for differences in RNA content of infected cells due to virally-induced

transcriptional shutdown, all cells were downsampled to the 25th percentile of total UMIs

of infected cells. Cells with fewer UMIs than the threshold were excluded from analysis.

Differential gene expression was performed with edgeR using a generalized linear

model quasi-likelihood F test adapted with a term for gene detection rate(55, 56). Genes

with an absolute $\log_2$ fold change greater than or equal to 1 and false discovery rate

less than 0.05 were considered significant. For KEGG enrichment analysis, pairwise

tests between mock, bystander, and infected cells were performed. Differentially

expressed genes with an absolute $\log_2$ fold change greater than or equal to 1 and false

discovery rate less than 0.05 were considered significant and subject to KEGG

enrichment analysis using the KEGG annotations for African Green Monkey as

implemented in the edgeR function kegga.

Quantification of SARS-CoV-2 sgmRNA Junction Sites

We explored the ability of our extended R1 sequencing to detect SARS-CoV-2

sgmRNA junctions using STARsolo (version 2.7.8a)(57). Aligned reads were re-mapped

to the empirical SARS-CoV-2 annotation described above and junction sites per cell

were quantified. The resulting junction per cell matrix was plotted in R/v4.0.4.

Flow Cytometry

Vero E6 cells were fixed with 4% paraformaldehyde at room temperature for a

minimum of 24 hours, washed once with PBS and permeabilized with 1X perm-wash

buffer (BDBiosciences #554723) for 5 minutes. SARS-CoV nucleocapsid (N) antibody

(clone 1C7C7) (kindly provided by Thomas Moran, Icahn School of Medicine at Mount

Sinai, New York, NY), conjugated to AlexaFluor 647 was diluted 1:400 in perm-wash

buffer, and added directly to samples. Samples were then incubated at room

temperature for 40 minutes in the dark. After staining, samples were washed once with

1X perm-wash buffer, once with PBS, resuspended in FACS buffer (PBS supplemented

with 1% FBS), and acquired on a Gallios flow cytometer (Beckman-Coulter). For all viral

infections, analysis was performed with FlowJo software (version 10.7.1, Becton

Dickinson), excluding cell doublets and debris and gating according to mock infected

populations.

Immunofluorescence microscopy

Vero E6 were seeded in 6-well plates (Falcon REF-353046) with one coverslip (Fisher

Scientific 12-550-143) per well. After 24 hours post infection, cells were washed with

PBS and fixed with 4% paraformaldehyde (Fisher Scientific AA433689M) overnight.

Fixed cells were permeabilized using 0.1% Triton-X (Fisher Scientific AC327371000) in

PBS and blocked with 4% bovine serum albumin (BSA, Fisher Scientific BP1600-100) in

PBS. Blocked coverslips were incubated with mouse anti-SARS-CoV N antibody (clone

1C7, 1:500 in 4% BSA PBS) overnight at 4C, washed three times with PBS, and

incubated for 45 minutes with 1:500 AlexaFluor 488-conjugated anti-mouse (Invitrogen

A11001, 1:500 in 4% BSA PBS) plus DAPI (Thermo Fisher Scientific D1306, 1:1000 in

4% BSA PBS) at room temperature. Coverslips were then stained with phalloidin (1:400

in PBS) for 1 hour at room temperature and washed again three times with PBS.

Coverslips were mounted using Prolong Diamond (Life Technologies P36970). Confocal

laser scanning was performed using a Leica SP5 DMI (ISMMS Microscopy CoRE and

Advanced Bioimaging Center) with a ×40/1.25 oil objective. Images were collected at a

resolution of 512 × 512 pixels in triplicate per slide. Images were processed and

analyzed using LAS X and CellProfiler v4(58).

Data Availability

Raw and processed scRNAseq data are available at (*GEO accession number*

pending) and code is available at (*github pending*).

Results

SARS-CoV-2 generates gRNA and sgmRNAs during infection, which are highly

redundant in their sequences (**Figure 1A**). Reads mapping to redundant sequences are

assigned to all genes which contain that sequence and are typically excluded from

quantification steps in scRNAseq processing pipelines. We therefore identified read

structures which could unambiguously identify gRNA or different species of sgmRNAs

to allow for their quantification (**Figure 1B**). Reads derived from gRNA should be

contiguous and could map anywhere on the SARS-CoV-2 genome. Reads derived from

sgmRNA could be either gapped or contiguous and could map to the 5′ leader and/or

downstream of the start site of S, the most 5′ sgmRNA. Because contiguous reads

mapping downstream of S could derive either from gRNA or sgmRNAs, they were

excluded from quantification. Only reads aligning in part to the 5′ leader and in part

downstream of S could be confidently derived from sgmRNAs. We therefore defined

gRNA reads as contiguous reads aligning upstream of regions contained in sgmRNAs.

sgmRNA reads were defined as discontinuous reads spanning the leader region and

regions used by sgmRNAs. Reads that did not match either of these formats could not

unambiguously be assigned to gRNA or an sgmRNA and were therefore excluded from

quantification (**Figure 1B**). With this framework, we developed scCoVseq to quantify

unambiguous genomic and subgenomic viral reads (**Figure 1C**). Using scCoVseq, we

compared the abilities of different scRNAseq methods to quantify SARS-CoV-2 RNAs.

In the widely available Chromium scRNAseq method developed by 10X

Genomics, Inc, there are two formats for droplet-based scRNAseq: 10X 3′ and 10X 5′.

10X 3′ generates library fragments derived from the 3′ regions of poly-adenylated RNAs

within a cell (**Figure 2A**). Because sgmRNAs share all viral sequence 3′ of the leader-

body junction site, 10X 3′ library fragments derived from SARS-CoV-2 heavily cover the

3′ end of the viral genome and do not contain leader-ORF junctions (**Figure 2D**). These

reads cannot differentiate gRNA from sgmRNA or distinguish different sgmRNA

species. 10X 5′ generates library fragments from the 5′ termini of poly-adenylated

RNAs (**Figure 2B**). These fragments are on average approximately 500 bp long

(according to the manufacturer's documentation) and should contain leader-ORF

junctions of SARS-CoV-2 sgmRNAs. The transcript read (R2), however, derives from

the 3′ end of these fragments and at the recommended read length of 91 bases is not

long enough to consistently sequence into the leader-sgmRNA junction site (**Figure**

2B). 10X 5′ can therefore detect some but not all junctions (**Figure 2E**). We reasoned

that we could use 10X 5′ library fragments to detect junction-spanning reads by

sequencing from the 5′ end of the fragment. To do this we extended R1, which is

normally used to sequence the cell barcode and UMI, to sequence into the leader-body

junction site (**Figure 2C**). Using 10X 5′ with Extended R1, we were able to sequence

more leader-sgmRNA junction sites and increase our ability to unambiguously quantify

sgmRNAs (**Figure 2F**). Indeed 10X 5′ Extended R1 increased the number of leader-

sgmRNA spanning reads over 10X 5′ and 10X 3′ (**Figure 2G**). When quantified with

scCoVseq, we found that 10X 5′ Extended R1 quantifies more UMIs per sgmRNA per

cell compared to 10X 5′ or 10X 3′ (**Figure 2H**). Importantly, the average host gene

expression per sample was significantly correlated across methods, suggesting that

host gene measurements were minimally affected by 10X 5′ Extended R1

(**Supplemental Figure 1**). Taken together, 10X 5′ libraries sequenced with extended

R1 sequencing results in a greater number of unambiguous reads derived from

sgmRNAs over 10X 3′ or 10X 5′, and consequently recovers more sgmRNA UMIs/cell.

Using this method, we analyzed Vero E6 cells 24 hours post infection with

SARS-CoV-2 at an MOI of 0.1 (**Figure 3A**). We were able to quantify sgmRNAs and

gRNA at single-cell resolution (**Figure 3B**). Using expression values for sgmRNAs, we

compared multiple unsupervised methods to identify infected cells. We found that a k-

medoid clustering approach implemented with the pam algorithm performed best as

indicated by multiple metrics to separate infected from uninfected cells (**Supplemental**

Figure 2A-C). We found that this classification method detected a similar percentage of

infected cells as detected using flow cytometry and immunofluorescence microscopy of

the same cultures (**Supplemental Figure 2D**). Using our infection classification, we

performed differential expression testing of infected cells compared to bystander cells

within the same culture as well as to cells from a mock culture. As previously

described(35), we observed downregulation of many host genes in infected cells

accompanied by an upregulation of cellular stress response genes (**Figure 3D, E**). We

further observed that, while bystander and mock cells had similar gene expression, a

small number of genes were upregulated in bystander cells compared to mock cells

(**Figure 3D, E**). This is especially notable given the inability of Vero E6 cells to produce

interferons in response to viral infection (59). KEGG enrichment analysis of differentially

expressed genes in pairwise comparisons of infected, mock, and bystander cells

showed that genes related to COVID19 were enriched in our infected cells supporting

our method for infection classification (**Figure 3F**).

Discussion

In this study, we examined the ability of two commonly used scRNAseq methods,

10X 3′ and 10X 5′, to detect and quantify SARS-CoV-2 derived RNAs with a focus on

sgmRNAs. Because of the redundant nature of coronavirus sgmRNA sequences, we

developed scCoVseq, which unambiguously quantifies both sgmRNAs and gRNAs in

10X data. We found that 10X methods detect unambiguous leader-sgmRNA junction-

spanning reads to different degrees. We were able to increase the detection of leader-

sgmRNA junction-spanning reads by extending the length of R1 during sequencing of

10X 5′ libraries, an approach we term 10X 5′ Extended R1 sequencing. Combining 10X

5′ Extended R1 with scCoVseq maximized quantification of sgmRNA UMIs compared to

10X 5′ or 10X 3′.

The ability to use sgmRNA expression to identify cells with actively replicating

virus may improve the utility of scRNAseq in studies of coronavirus tropism. A challenge

in many scRNAseq studies, particularly studies of primary tissues, has been identifying

cells with active infection as opposed to cells with large amounts of ambient or

extracellular viral RNA (such as phagocytic cells and/or cells not supporting active

infection)(16). Because sgmRNAs are produced only during viral replication and are

absent from virions, our method allows us to distinguish between infected and

uninfected cells associated with "background" viral RNA (**Supplemental Figure 2C**). In

downstream analyses of host transcriptomic changes induced by infection, accurate

classification of infected cells is important for robust analyses of transcriptional

differences between infected and uninfected cells because incorrect classifications may

dilute effect sizes and resultant significance values. This method also enables the

comparison of sgmRNA expression dynamics at single cell resolution. Such analyses

may be particularly relevant for comparing viral gene expression between different cell

types, coronaviruses, or between SARS-CoV-2 variants of interest, which have been

described to have different kinetics of sgmRNA expression(60). This approach could be

extended to any coronavirus or nidovirus, including potentially novel emerging

coronaviruses. Finally, scCoVseq can be used to examine differential junction site

usage within single cells (**Supplemental Figure 3**). Several groups have identified

TRS-independent SARS-CoV-2 sgmRNAs(12, 13, 15), the significance of which

remains unknown. It is possible that changes in junction site usage between cell types

or during the course of infection may play a role in pathogenesis.

It should be noted that there are some limitations to our study. With our dataset,

we are unable to know the true infection state of a cell processed for scRNAseq, and

therefore we cannot assess the true accuracy of our method to classify infected cells.

An additional limitation of our method is that quantification of viral genes with scCoVseq

is dependent on accurate annotation of viral RNAs. We derived our annotation based on

published empirically-defined TRS-dependent RNAs(12), but this does not preclude the

existence of other viral RNAs at time points or in cell types not studied. Importantly, we

explicitly exclude TRS-independent RNAs from our analyses. Methods such as

STARsolo(57) or sequencing 10X libraries with long-read sequencing(61) may allow for

detection and quantification of viral RNAs without reference annotation and irrespective

of TRSs.

Acknowledgments

This work was supported in part by NIH grants R21 AI149180, R01 AI151029,

and U01 AI150748. P.C. was supported by the Mount Sinai Medical Scientist Training

Program T32 GM007280. P.C, R.S.P., and E.J.D. were supported by the Viral Host

Pathogenesis Training Grant T32 AI07647.

We thank Randy Albrecht for BSL3 facility management and support. We also

thank Thomas Moran, Center for Therapeutic Antibody Discovery at the Icahn School of

Medicine at Mount Sinai, for kindly providing anti-SARS-CoV N antibody. We thank

Michael A Schotsaert for flow cytometry support. This work was supported in part

through the computational resources and staff expertise provided by Scientific

Computing at the Icahn School of Medicine at Mount Sinai. Research reported in this

paper was supported by the Office of Research Infrastructure of the National Institutes

of Health under award number S10OD026880. The content is solely the responsibility of

the authors and does not necessarily represent the official views of the National

Institutes of Health. Microscopy was performed at the Microscopy CoRE at the Icahn

School of Medicine at Mount Sinai.

Author Contributions

Conceptualization: P.C., B.R.R.

Data Curation: P.C., B.R.R.

Formal Analysis: P.C., B.R.R.

Funding Acquisition: B.R.R.

Investigation: P.C., E.J.D., O.D.

Methodology: P.C., B.R.R.

Project Administration: B.R.R.

Software: P.C., R.S.P., B.R.R.

Supervision: B.R.R.

Validation: P.C., R.S.P., E.J.D., O.D.

Visualization: P.C., B.R.R.

Writing – original draft: P.C., B.R.R.

Writing – review & editing: P.C., E.J.D., O.D., R.S.P., B.R.R.

References

1. Dong E, Du H, Gardner L. 2020. An interactive web-based dashboard to track

COVID-19 in real time. Lancet Infect Dis 20:533–534.

2. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R,

Niu P, Zhan F, Ma X, Wang D, Xu W, Wu G, Gao GF, Tan W, Investigating CNC,

Team R. 2020. A novel coronavirus from patients with pneumonia in china, 2019. N

Engl J Med 382:727–733.

3. Carvalho T, Krammer F, Iwasaki A. 2021. The first 12 months of COVID-19: a

timeline of immunological insights. 4. Nat Rev Immunol 21:245–256.

4. Gordon DE, Jang GM, Bouhaddou M, Xu J, Obernier K, White KM, O'Meara MJ,

Rezelj VV, Guo JZ, Swaney DL, Tummino TA, Huettenhain R, Kaake RM, Richards

AL, Tutuncuoglu B, Foussard H, Batra J, Haas K, Modak M, Kim M, Haas P,

Polacco BJ, Braberg H, Fabius JM, Eckhardt M, Soucheray M, Bennett MJ, Cakir

M, McGregor MJ, Li Q, Meyer B, Roesch F, Vallet T, Mac Kain A, Miorin L, Moreno

E, Naing ZZC, Zhou Y, Peng S, Shi Y, Zhang Z, Shen W, Kirby IT, Melnyk JE,

Chorba JS, Lou K, Dai SA, Barrio-Hernandez I, Memon D, Hernandez-Armenta C,

Lyu J, Mathy CJP, Perica T, Pilla KB, Ganesan SJ, Saltzberg DJ, Rakesh R, Liu X,

Rosenthal SB, Calviello L, Venkataramanan S, Liboy-Lugo J, Lin Y, Huang X-P, Liu

Y, Wankowicz SA, Bohn M, Safari M, Ugur FS, Koh C, Savar NS, Tran QD,

Shengjuler D, Fletcher SJ, O'Neal MC, Cai Y, Chang JCJ, Broadhurst DJ,

Klippsten S, Sharp PP, Wenzell NA, Kuzuoglu D, Wang H-Y, Trenker R, Young

JM, Cavero DA, Hiatt J, Roth TL, Rathore U, Subramanian A, Noack J, Hubert M,

Stroud RM, Frankel AD, Rosenberg OS, Verba KA, Agard DA, Ott M, Emerman M,

Jura N, von Zastrow M, Verdin E, Ashworth A, Schwartz O, d'Enfert C, Mukherjee

S, Jacobson M, Malik HS, Fujimori DG, Ideker T, Craik CS, Floor SN, Fraser JS,

Gross JD, Sali A, Roth BL, Ruggero D, Taunton J, Kortemme T, Beltrao P,

Vignuzzi M, García-Sastre A, Shokat KM, Shoichet BK, Krogan NJ. 2020. A SARS-

CoV-2 protein interaction map reveals targets for drug repurposing. Nature

https://doi.org/10.1038/s41586-020-2286-9.

5. Shen B, Yi X, Sun Y, Bi X, Du J, Zhang C, Quan S, Zhang F, Sun R, Qian L, Ge W,

Liu W, Liang S, Chen H, Zhang Y, Li J, Xu J, He Z, Chen B, Wang J, Yan H, Zheng

Y, Wang D, Zhu J, Kong Z, Kang Z, Liang X, Ding X, Ruan G, Xiang N, Cai X, Gao

H, Li L, Li S, Xiao Q, Lu T, Zhu Y, Liu H, Chen H, Guo T. 2020. Proteomic and

metabolomic characterization of COVID-19 patient sera. Cell

https://doi.org/10.1016/j.cell.2020.05.032.

6. Bojkova D, Klann K, Koch B, Widera M, Krause D, Ciesek S, Cinatl J, Münch C.

2020. SARS-CoV-2 infected host cell proteomics reveal potential therapy targets

https://doi.org/10.21203/rs.3.rs-17218/v1.

7. Finkel Y, Mizrahi O, Nachshon A, Weingarten-Gabbay S, Morgenstern D, Yahalom-

Ronen Y, Tamir H, Achdout H, Stein D, Israeli O, Beth-Din A, Melamed S, Weiss S,

Israely T, Paran N, Schwartz M, Stern-Ginossar N. 2020. The coding capacity of

SARS-CoV-2. Nature 1–9.

8. Bouhaddou M, Memon D, Meyer B, White KM, Rezelj VV, Marrero MC, Polacco BJ,

Melnyk JE, Ulferts S, Kaake RM, Batra J, Richards AL, Stevenson E, Gordon DE,

Rojc A, Obernier K, Fabius JM, Soucheray M, Miorin L, Moreno E, Koh C, Tran

QD, Hardy A, Robinot R, Vallet T, Nilsson-Payant BE, Hernandez-Armenta C,

Dunham A, Weigang S, Knerr J, Modak M, Quintero D, Zhou Y, Dugourd A,

Valdeolivas A, Patil T, Li Q, Hüttenhain R, Cakir M, Muralidharan M, Kim M, Jang

G, Tutuncuoglu B, Hiatt J, Guo JZ, Xu J, Bouhaddou S, Mathy CJP, Gaulton A,

Manners EJ, Félix E, Shi Y, Goff M, Lim JK, McBride T, O'Neal MC, Cai Y, Chang

JCJ, Broadhurst DJ, Klippsten S, Wit ED, Leach AR, Kortemme T, Shoichet B, Ott

M, Saez-Rodriguez J, tenOever BR, Mullins D, Fischer ER, Kochs G, Grosse R,

García-Sastre A, Vignuzzi M, Johnson JR, Shokat KM, Swaney DL, Beltrao P,

Krogan NJ. 2020. The Global Phosphorylation Landscape of SARS-CoV-2

Infection. Cell 0.

9. Schmidt N, Lareau CA, Keshishian H, Ganskih S, Schneider C, Hennig T, Melanson

R, Werner S, Wei Y, Zimmer M, Ade J, Kirschner L, Zielinski S, Dölken L, Lander

ES, Caliskan N, Fischer U, Vogel J, Carr SA, Bodem J, Munschauer M. 2021. The

SARS-CoV-2 RNA–protein interactome in infected human cells. 3. Nat Microbiol

6:339–353.

10. Flynn RA, Belk JA, Qi Y, Yasumoto Y, Wei J, Alfajaro MM, Shi Q, Mumbach MR,

Limaye A, DeWeirdt PC, Schmitz CO, Parker KR, Woo E, Chang HY, Horvath TL,

Carette JE, Bertozzi CR, Wilen CB, Satpathy AT. 2021. Discovery and functional

interrogation of SARS-CoV-2 RNA-host protein interactions. Cell 184:2394-

2411.e16.

11. Blanco - Melo D. 2020. Imbalanced host response to SARS - CoV - 2 drives

development of COVID - 19. Cell.

12. Kim D, Lee J-Y, Yang J-S, Kim JW, Kim VN, Chang H. 2020. The architecture of

SARS-CoV-2 transcriptome. Cell 181:914-921.e10.

13. Wang D, Jiang A, Feng J, Li G, Guo D, Sajid M, Wu K, Zhang Q, Ponty Y, Will S,

Liu F, Yu X, Li S, Liu Q, Yang X-L, Guo M, Li X, Chen M, Shi Z-L, Lan K, Chen Y,

Zhou Y. 2021. The SARS-CoV-2 Subgenome Landscape and its Novel Regulatory

Features. Mol Cell 0.

14. Ziv O, Price J, Shalamova L, Kamenova T, Goodfellow I, Weber F, Miska EA.

2020. The short- and long-range RNA-RNA Interactome of SARS-CoV-2. Mol Cell

https://doi.org/10.1016/j.molcel.2020.11.004.

15. Chang JJ-Y, Rawlinson D, Pitt ME, Taiaroa G, Gleeson J, Zhou C, Mordant FL,

Paoli-Iseppi RD, Caly L, Purcell DFJ, Stinear TP, Londrigan SL, Clark MB,

Williamson DA, Subbarao K, Coin LJM. 2021. Transcriptional and epi-

transcriptional dynamics of SARS-CoV-2 during cellular infection. Cell Rep 0.

16. Bost P, Giladi A, Liu Y, Bendjelal Y, Xu G, David E, Blecher-Gonen R, Cohen M,

Medaglia C, Li H, Deczkowska A, Zhang S, Schwikowski B, Zhang Z, Amit I. 2020.

512   Host-viral infection maps reveal signatures of severe COVID-19 patients. Cell

513   https://doi.org/10.1016/j.cell.2020.05.006.

514 17. Sungnak W, Huang N, Bécavin C, Berg M, Queen R, Litvinukova M, Talavera-

515   López C, Maatz H, Reichart D, Sampaziotis F, Worlock KB, Yoshida M, Barnes JL,

516   Network HLB. 2020. SARS-CoV-2 entry factors are highly expressed in nasal

517   epithelial cells together with innate immune genes. Nat Med 26:681–687.

518 18. Lukassen S, Chua RL, Trefzer T, Kahn NC, Schneider MA, Muley T, Winter H,

519   Meister M, Veith C, Boots AW, Hennig BP, Kreuter M, Conrad C, Eils R. 2020.

520   SARS-CoV-2 receptor ACE2 and TMPRSS2 are primarily expressed in bronchial

521   transient secretory cells. EMBO J 39:e105114.

522 19. Fodoulian L, Tuberosa J, Rossier D, Landis B, Carleton A, Rodriguez I. 2020.

523   SARS-CoV-2 receptor and entry genes are expressed by sustentacular cells in the

524   human olfactory neuroepithelium. BioRxiv

525   https://doi.org/10.1101/2020.03.31.013268.

526 20. Qi F, Qian S, Zhang S, Zhang Z. 2020. Single cell RNA sequencing of 13 human

527   tissues identify cell types and receptors of human coronaviruses. Biochem Biophys

528   Res Commun 526:135–140.

529 21. Ravindra NG, Alfajaro MM, Gasque V, Wei J, Filler RB, Huston NC, Wan H,

530   Szigeti-Buck K, Wang B, Montgomery RR, Eisenbarth SC, Williams A, Pyle AM,

531   Iwasaki A, Horvath TL, Foxman EF, van Dijk D, Wilen CB. 2020. Single-cell

longitudinal analysis of SARS-CoV-2 infection in human bronchial epithelial cells.

BioRxiv https://doi.org/10.1101/2020.05.06.081695.

22. Single-cell meta-analysis of SARS-CoV-2 entry genes across tissues and demographics | Nature Medicine.

23. Bieberich F, Vazquez-Lombardi R, Yermanos A, Ehling RA, Mason DM, Wagner B, Kapetanovic E, Roberto RBD, Weber CR, Savic M, Rudolf F, Reddy ST. 2021. A single-cell atlas of lymphocyte adaptive immune repertoires and transcriptomes reveals age-related differences in convalescent COVID-19 patients. bioRxiv 2021.02.12.430907.

24. Wilk AJ, Rustagi A, Zhao NQ, Roque J, Martínez-Colón GJ, McKechnie JL, Ivison GT, Ranganath T, Vergara R, Hollis T, Simpson LJ, Grant P, Subramanian A, Rogers AJ, Blish CA. 2020. A single-cell atlas of the peripheral immune response in patients with severe COVID-19. 7. Nat Med 26:1070–1076.

25. Yao C, Bora SA, Parimon T, Zaman T, Friedman OA, Palatinus JA, Surapaneni NS, Matusov YP, Chiang GC, Kassar AG, Patel N, Green CER, Aziz AW, Suri H, Suda J, Lopez AA, Martins GA, Stripp BR, Gharib SA, Goodridge HS, Chen P. 2021. Cell-Type-Specific Immune Dysregulation in Severely Ill COVID-19 Patients. Cell Rep 34.

26. MacDonald L, Otto TD, Elmesmari A, Tolusso B, Somma D, McSharry C, Gremese E, McInnes IB, Alivernini S, Kurowska-Stolarska M. 2020. COVID-19 and

Rheumatoid Arthritis share myeloid pathogenic and resolving pathways. bioRxiv

2020.07.26.221572.

27. Schreibing F, Hannani M, Ticconi F, Fewings E, Nagai JS, Begemann M, Kuppe C,

Kurth I, Kranz J, Frank D, Anslinger TM, Ziegler P, Kraus T, Enczmann J, Balz V,

Windhofer F, Balfanz P, Kurts C, Marx G, Marx N, Dreher M, Schneider RK, Saez-

Rodriguez J, Filho IGC, Kramann R. 2021. Dissecting CD8+ T cell pathology of

severe SARS-CoV-2 infection by single-cell epitope mapping. bioRxiv

2021.03.03.432690.

28. Wen W, Su W, Tang H, Le W, Zhang X, Zheng Y, Liu X, Xie L, Li J, Ye J, Dong L,

Cui X, Miao Y, Wang D, Dong J, Xiao C, Chen W, Wang H. 2020. Immune cell

profiling of COVID-19 patients in the recovery stage by single-cell sequencing. Cell

Discov 6:31.

29. Lee JS, Park S, Jeong HW, Ahn JY, Choi SJ, Lee H, Choi B, Nam SK, Sa M, Kwon

J-S, Jeong SJ, Lee HK, Park SH, Park S-H, Choi JY, Kim S-H, Jung I, Shin E-C.

2020. Immunophenotyping of COVID-19 and influenza highlights the role of type I

interferons in development of severe COVID-19. Sci Immunol 5.

30. Wang F-S, Zhang J-Y, Wang X, Xing X, Xu Z, Zhang C, Song J-W, Fan X, Xia P,

Fu J-L, Wang S-Y, Xu R-N, Dai X-P, Shi L, Huang L, Jiang T-J, Shi M, Zhang Y,

Zumla A, Maeurer M, Bai F. 2020. Single-cell landscape of immunological

responses in COVID-19 patients. bioRxiv 2020.07.23.217703.

31.   Zhang J-Y, Wang X-M, Xing X, Xu Z, Zhang C, Song J-W, Fan X, Xia P, Fu J-L,

Wang S-Y, Xu R-N, Dai X-P, Shi L, Huang L, Jiang T-J, Shi M, Zhang Y, Zumla A,

Maeurer M, Bai F, Wang F-S. 2020. Single-cell landscape of immunological

responses in patients with COVID-19. 9. Nat Immunol 21:1107–1118.

32.   Kalfaoglu B, Almeida-Santos J, Tye CA, Satou Y, Ono M. 2020. T-cell

hyperactivation and paralysis in severe COVID-19 infection revealed by single-cell

analysis. BioRxiv https://doi.org/10.1101/2020.05.26.115923.

33.   Wei L, Ming S, Zou B, Wu Y, Hong Z, Li Z, Zheng X, Huang M, Luo L, Liang J, Wen

X, Chen T, Liang Q, Kuang L, Shan H, Huang X. 2020. Viral invasion and type I

interferon response characterize the immunophenotypes during COVID-19

infection. Electron J https://doi.org/10.2139/ssrn.3555695.

34.   Wyler E, Mösbauer K, Franke V, Diag A, Gottula LT, Arsie R, Klironomos F,

Koppstein D, Ayoub S, Buccitelli C, Richter A, Legnini I, Ivanov A, Mari T, Del

Giudice S, Papies JP, Müller MA, Niemeyer D, Selbach M, Akalin A, Rajewsky N,

Drosten C, Landthaler M. 2020. Bulk and single-cell gene expression profiling of

SARS-CoV-2 infected human cell lines identifies molecular targets for therapeutic

intervention. BioRxiv https://doi.org/10.1101/2020.05.05.079194.

35.   Miorin L, Kehrer T, Sanchez-Aparicio MT, Zhang K, Cohen P, Patel RS, Cupic A,

Makio T, Mei M, Moreno E, Danziger O, White KM, Rathnasinghe R, Uccellini M,

Gao S, Aydillo T, Mena I, Yin X, Martin-Sancho L, Krogan NJ, Chanda SK,

Schotsaert M, Wozniak RW, Ren Y, Rosenberg BR, Fontoura BMA, García-Sastre

A. 2020. SARS-CoV-2 Orf6 hijacks Nup98 to block STAT nuclear import and

antagonize interferon signaling. Proc Natl Acad Sci 117:28344–28354.

36. Melms JC, Biermann J, Huang H, Wang Y, Nair A, Tagore S, Katsyv I, Rendeiro

AF, Amin AD, Schapiro D, Frangieh CJ, Luoma AM, Filliol A, Fang Y,

Ravichandran H, Clausi MG, Alba GA, Rogava M, Chen SW, Ho P, Montoro DT,

Kornberg AE, Han AS, Bakhoum MF, Anandasabapathy N, Suárez-Fariñas M,

Bakhoum SF, Bram Y, Borczuk A, Guo XV, Lefkowitch JH, Marboe C, Lagana SM,

Del Portillo A, Zorn E, Markowitz GS, Schwabe RF, Schwartz RE, Elemento O,

Saqi A, Hibshoosh H, Que J, Izar B. 2021. A molecular single-cell lung atlas of

lethal COVID-19. Nature 1–6.

37. Delorey TM, Ziegler CGK, Heimberg G, Normand R, Yang Y, Segerstolpe Å,

Abbondanza D, Fleming SJ, Subramanian A, Montoro DT, Jagadeesh KA, Dey KK,

Sen P, Slyper M, Pita-Juárez YH, Phillips D, Biermann J, Bloom-Ackermann Z,

Barkas N, Ganna A, Gomez J, Melms JC, Katsyv I, Normandin E, Naderi P, Popov

YV, Raju SS, Niezen S, Tsai LT-Y, Siddle KJ, Sud M, Tran VM, Vellarikkal SK,

Wang Y, Amir-Zilberstein L, Atri DS, Beechem J, Brook OR, Chen J, Divakar P,

Dorceus P, Engreitz JM, Essene A, Fitzgerald DM, Fropf R, Gazal S, Gould J,

Grzyb J, Harvey T, Hecht J, Hether T, Jané-Valbuena J, Leney-Greene M, Ma H,

McCabe C, McLoughlin DE, Miller EM, Muus C, Niemi M, Padera R, Pan L, Pant D,

Pe'er C, Pfiffner-Borges J, Pinto CJ, Plaisted J, Reeves J, Ross M, Rudy M,

Rueckert EH, Siciliano M, Sturm A, Todres E, Waghray A, Warren S, Zhang S,

Zollinger DR, Cosimi L, Gupta RM, Hacohen N, Hibshoosh H, Hide W, Price AL,

Rajagopal J, Tata PR, Riedel S, Szabo G, Tickle TL, Ellinor PT, Hung D, Sabeti

PC, Novak R, Rogers R, Ingber DE, Jiang ZG, Juric D, Babadi M, Farhi SL, Izar B,

Stone JR, Vlachos IS, Solomon IH, Ashenberg O, Porter CBM, Li B, Shalek AK,

Villani A-C, Rozenblatt-Rosen O, Regev A. 2021. COVID-19 tissue atlases reveal

SARS-CoV-2 pathology and cellular targets. Nature 1–8.

38.   Mechanisms of SARS-CoV-2 Transmission and Pathogenesis: Trends in

Immunology.

39. V'kovski P, Kratzel A, Steiner S, Stalder H, Thiel V. 2020. Coronavirus biology and

replication: implications for SARS-CoV-2. 3. Nat Rev Microbiol 1–16.

40. Sola I, Almazán F, Zúñiga S, Enjuanes L. 2015. Continuous and discontinuous

RNA synthesis in coronaviruses. Annu Rev Virol 2:265–288.

41. Perlman S, Masters PS. 2020. Coronaviridae: The Viruses and Their Replication,

p. 411–442. *In* Howley, PM, Knipe, DM, Whelan, S (eds.), Fields Virology:

Emerging Viruses, 7th ed. Wolters Kluwer Health/lippincott Williams & Wilkins,

Philadelphia, PA.

42. Daniloski Z, Jordan TX, Wessels H-H, Hoagland DA, Kasela S, Legut M, Maniatis

S, Mimitou EP, Lu L, Geller E, Danziger O, Rosenberg BR, Phatnani H, Smibert P,

Lappalainen T, tenOever BR, Sanjana NE. 2021. Identification of Required Host

Factors for SARS-CoV-2 Infection in Human Cells. Cell 184:92-105.e16.

43.   Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas

635   AR, Kamitaki N, Martersteck EM, Trombetta JJ, Weitz DA, Sanes JR, Shalek AK,

636   Regev A, McCarroll SA. 2015. Highly parallel genome-wide expression profiling of

637   individual cells using nanoliter droplets. Cell 161:1202–1214.

638   44.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. 2018. Integrating single-cell

639   transcriptomic data across different conditions, technologies, and species. Nat

640   Biotechnol 36:411–420.

641   45.   Hafemeister C, Satija R. 2019. Normalization and variance stabilization of single-

642   cell RNA-seq data using regularized negative binomial regression. Genome Biol

643   20:296.

644   46.   Li H. 2021. lh3/seqtk. C.

645   47.   Fernandes JD, Hinrichs AS, Clawson H, Navarro Gonzalez J, Lee BT, Nassar LR,

646   Raney BJ, Rosenbloom KR, Nerli S, Rao A, Schmelter D, Zweig AS, Lowe TM,

647   Ares M, Corbet-Detig R, Kent WJ, Haussler D, Haeussler M. 2020. The UCSC

648   SARS-CoV-2 genome browser. BioRxiv

649   https://doi.org/10.1101/2020.05.04.075945.

650   48.   Twelve years of SAMtools and BCFtools | GigaScience | Oxford Academic.

651   49.   UMI-tools: Modelling sequencing errors in Unique Molecular Identifiers to improve

652   quantification accuracy.

50.   2021. Sparse and Dense Matrix Classes and Methods [R package Matrix version

1.3-4]. Comprehensive R Archive Network (CRAN).

51.   2020. Read Rectangular Text Data [R package readr version 1.4.0].

Comprehensive R Archive Network (CRAN).

52.   Garrido-Martín D, Palumbo E, Guigó R, Breschi A. 2018. ggsashimi: Sashimi plot

revised for browser- and annotation-independent splicing visualization. PLOS

Comput Biol 14:e1006360.

53.   2008. clValid: An R Package for Cluster Validation by Guy Brock, Vasyl Pihur,

Susmita Datta, Somnath Datta.

54.   Maechler M, original) PR (Fortran, original) AS (S, original) MH (S, Hornik [trl K,

maintenance(1999-2000)] ctb] (port to R, Studer M, Roudier P, Gonzalez J,

Kozlowski K, pam()) ES (fastpam options for, Murphy (volume.ellipsoid({d >= 3}))

K. 2021. cluster: "Finding Groups in Data": Cluster Analysis Extended Rousseeuw

et al.

55.   Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for

differential expression analysis of digital gene expression data. Bioinformatics

26:139–140.

56.   Soneson C, Robinson MD. 2018. Bias, robustness and scalability in single-cell

differential expression analysis. 4. Nat Methods 15:255–261.

57.   Kaminow B, Yunusov D, Dobin A. 2021. STARsolo: accurate, fast and versatile

mapping/quantification of single-cell and single-nucleus RNA-seq data. bioRxiv

2021.05.05.442755.

58.   McQuin C, Goodman A, Chernyshev V, Kamentsky L, Cimini BA, Karhohs KW,

Doan M, Ding L, Rafelski SM, Thirstrup D, Wiegraebe W, Singh S, Becker T,

Caicedo JC, Carpenter AE. 2018. CellProfiler 3.0: Next-generation image

processing for biology. PLOS Biol 16:e2005970.

59.   Emeny JM, Morgan MJY 1979. Regulation of the Interferon System: Evidence that

Vero Cells have a Genetic Defect in Interferon Production. J Gen Virol 43:247–252.

60.   Parker MD, Lindsey BB, Shah DR, Hsu S, Keeley AJ, Partridge DG, Leary S, Cope

A, State A, Johnson K, Ali N, Raghei R, Heffer J, Smith N, Zhang P, Gallis M,

Louka SF, Whiteley M, Foulkes BH, Christou S, Wolverson P, Pohare M, Hansford

SE, Green LR, Evans C, Raza M, Wang D, Gaudieri S, Mallal S, Consortium TC-19

GU (COG-U, Silva TI de. 2021. Altered Sub-Genomic RNA Expression in SARS-

CoV-2 B.1.1.7 Infections. bioRxiv 2021.03.02.433156.

61.   Russell AB, Elshina E, Kowalsky JR, Te Velthuis AJW, Bloom JD. 2019. Single-cell

virus sequencing of influenza infections that trigger innate immunity. J Virol 93.

Figure Legends

Figure 1: A. Illustration of SARS-CoV-2 genomic RNA, gRNA, and subgenomic RNAs, sgmRNAs. **B.** *Top:* Reads included for analysis by scCoVseq. Either: 1) contiguous reads mapping to ORF1a/b and therefore derived from gRNA or 2) discontinuous reads spanning the leader region and ORFS transcribed by sgmRNAs *Bottom*: Reads excluded from analysis by scCoVseq. Either: 1) discontinuous reads that do not include sequence mapping to the leader region and downstream of S or 2) contiguous reads that map to ORFs other than ORF1a/b, which are ambiguous. **C.** Activity diagram of scCoVseq pipeline. Blue rectangles indicate inputs/outputs for each stage. Orange rounded rectangles indicate a process in bold with software indicated.

Figure 2: A-C. Illustration of gRNA and S and ORF3a sgmRNAs. Red box indicates regions contained in final 10X library. *Lower:* Example illustration of 10X library fragments derived from gRNA and S and ORF3a sgmRNAs with sequencing read 1 and read 2 indicated. 10X 3′ (**A**), 10X 5′ (**B**), and 10X 5′ Extended R1 (**C**) libraries are illustrated. **D-F.** Sashimi plot of 10X 3′ (**D**), 10X 5′ (**E**), and 10X 5′ Extended R1 (**F**) reads mapped to the SARS-CoV-2 genome filtered to show only junctions supported by at least 1,000 reads. Total number of reads visualized is indicated in the bottom right. **G.** Reads per million reads mapped to SARS-CoV-2 reads mapping to a single viral gene in 10X 3′, 10X 5′, or 10X 5′ with Extended R1 data. Reads are colored by their mapping with contiguous reads mapping to ORF1a/b in yellow, leader-sgmRNA junction-spanning reads in blue, and ambiguously mapped reads in grey. **H.** UMIs per cell for all sgmRNAs in infected cells in each dataset. Each dataset was downsampled to an equal

number of infected cells and each cells' total UMIs were downsampled to the same

value to control for differences in sequencing depth. The leader region is enlarged in

illustrations of the genome for visibility. L = Leader.

Figure 3: A. Experimental design. Vero E6 cells were infected or mock infected with

SARS-CoV-2 (USA-WA1/2020) at an MOI of 0.1. At 24 hours post-infection, cells were

analyzed by scRNAseq using 10X 5′ with Extended R1 sequencing. **B-C.** 3,047 mock

and infected cells embedded in tSNE space derived from euclidean distance of scaled

viral sgmRNA expression. Cells are colored by (**B**) indicated viral RNA expression, or

(**C**) experimental condition and assigned infection status of cells. **D.** Heatmap of genes

differentially expressed in infected, bystander, or mock cells. Differential expression

testing was performed on host gene expression downsampled to an equal number of

UMIs/cell across cells to account for infection-induced transcriptional shutdown. Genes

were selected for visualization based on false discovery rate of less than 0.05 and

absolute $\log_2$ fold change of at least 1. Non-downsampled gene expression data is

shown. Along the top, infection status, total viral UMIs and genomic RNA as quantified

by CellRanger and scCoVseq are indicated. Cells and genes are clustered with ward d2

clustering based on euclidean distance. **E.** Expression of selected host genes per cell

by infection status. Data shown is not downsampled. *Top:* genes induced in infected

cells. *Middle:* genes repressed in infected cells. *Bottom:* genes upregulated in bystander

cells compared to mock. **F.** KEGG pathway enrichment in genes differentially expressed

in pairwise comparisons of downsampled infected, bystander, and mock cells. Dot size

and fill indicates the $-\log_{10}$ p value of enrichment with red dots indicating enrichment in

the first infection state and blue in the second infection state noted above each panel.

Supplemental Figure 1: Average counts of host gene expression of cells analyzed by

10X 3′, 10X 5′, and 10X 5′ with Extended R1 sequencing. Each point represents the

average UMIs/cell of a single gene assayed in the indicated assays. At the top left, the

Pearson correlation coefficient and resulting p value are indicated.

Supplemental Figure 2: A. Comparison of performance metrics (average distance, AD;

average distance between means, ADM; average proportion of non-overlap, APN;

connectivity; Dunn Index; figure of merit, FOM; and silhouette index) by several

clustering methods (diana, model-based, hierarchical, kmeans, and pam) run on

sgmRNA expression of 600 randomly sampled cells analyzed with 10X 5′ Extended R1

and scCoVseq. *Left*: Performance metrics for each method across k values from 2 to 5.

Right: performance metrics for each method with k = 2. **B.** Visualization of infection

classification by different methods. **C.** Viral gene expression of cells by infection status,

determined by pam clustering method. **D.** Percent of infected cells per sample as

measured by flow cytometry, immunofluorescence, and infection classification with

unsupervised (pam) method or supervised infection classification by classifying infected

cells as those with at least 375 total viral UMIs. Because the same sample was

sequenced with 10X 5′ and 10X 5′ Extended R1, flow cytometry and

immunofluorescence results are duplicated for ease of visualization. Error bars for

immunofluorescence indicate mean ± one standard deviation of percent infected cells

based on three fields per sample.

Supplemental Figure 3: Detection of junction sites in SARS-CoV-2 reads with 10X 5′

Extended R1. Junction sites are represented by the 5′ start site and 3′ end site on the y

and x-axis, respectively. The color indicates the $\log_2$ total UMIs/junction across all cells

in the SARS-CoV-2 infected sample. Below each axis, the number of UMIs supporting a

position as a junction start or end site is indicated with a density plot.

Figure 1: A. Illustration of SARS-CoV-2 genomic RNA, gRNA, and subgenomic RNAs,

sgmRNAs. **B.** *Top:* Reads included for analysis by scCoVseq. Either: 1) contiguous

reads mapping to ORF1a/b and therefore derived from gRNA or 2) discontinuous reads

spanning the leader region and ORFS transcribed by sgmRNAs *Bottom*: Reads

769   excluded from analysis by scCoVseq. Either: 1) discontinuous reads that do not include

770   sequence mapping to the leader region and downstream of S or 2) contiguous reads

771   that map to ORFs other than ORF1a/b, which are ambiguous. **C.** Activity diagram of

772   scCoVseq pipeline. Blue rectangles indicate inputs/outputs for each stage. Orange

773   rounded rectangles indicate a process in bold with software indicated.

Figure 2: A-C. Illustration of gRNA and S and ORF3a sgmRNAs. Red box indicates

regions contained in final 10X library. *Lower:* Illustration of 10X library fragments derived

from gRNA and S and ORF3a sgmRNAs with sequencing read 1 and read 2 indicated.

10X 3′ (**A**), 10X 5′ (**B**), and 10X 5′ Extended R1 (**C**) libraries are illustrated. **D-F.**

Sashimi plot of 10X 3′ (**D**), 10X 5′ (**E**), and 10X 5′ Extended R1 (**F**) reads mapped to

the SARS-CoV-2 genome filtered to show only junctions supported by at least 1,000

reads. Total number of reads visualized is indicated in the bottom right. **G.** Reads per

million reads mapped to SARS-CoV-2 reads mapping to a single viral gene in 10X 3′,

10X 5′, or 10X 5′ with Extended R1 data. Reads are colored by their mapping with

contiguous reads mapping to ORF1a/b in yellow, leader-sgmRNA junction-spanning

reads in blue, and ambiguously mapped reads in grey. **H.** UMIs per cell for all sgmRNAs

in infected cells in each dataset. Each dataset was downsampled to an equal number of

infected cells and each cells' total UMIs were downsampled to the same value to control

for differences in sequencing depth. The leader region is enlarged in illustrations of the

genome for visibility. L = Leader.

Figure 3: A. Experimental design. Vero E6 cells were infected or mock infected with

SARS-CoV-2 (USA-WA1/2020) at an MOI of 0.1. At 24 hours post-infection, cells were

analyzed by scRNAseq using 10X 5′ with Extended R1 sequencing. **B-C.** 3,047 mock

and infected cells embedded in tSNE space derived from euclidean distance of scaled

viral sgmRNA expression. Cells are colored by (**B**) indicated viral RNA expression, or

(**C**) experimental condition and assigned infection status of cells. **D.** Heatmap of genes

differentially expressed in infected, bystander, or mock cells. Differential expression

testing was performed on host gene expression downsampled to an equal number of

UMIs/cell across cells to account for infection-induced transcriptional shutdown. Genes

were selected for visualization based on false discovery rate of less than 0.05 and

absolute $\log_2$ fold change of at least 1. Non-downsampled gene expression data is

shown. Along the top, infection status, total viral UMIs and genomic RNA as quantified

by CellRanger and scCoVseq are indicated. Cells and genes are clustered with ward d2

clustering based on euclidean distance. **E.** Expression of selected host genes per cell

by infection status. Data shown is not downsampled. *Top:* genes induced in infected

cells. *Middle:* genes repressed in infected cells. *Bottom:* genes upregulated in bystander

cells compared to mock. **F.** KEGG pathway enrichment in genes differentially expressed

in pairwise comparisons of downsampled infected, bystander, and mock cells. Dot size

and fill indicates the $-\log_{10}$ p value of enrichment with red dots indicating enrichment in

the first infection state and blue in the second infection state noted above each panel.

Supplemental Figure 1: Average counts of host gene expression of cells analyzed by

10X 3′, 10X 5′, and 10X 5′ with Extended R1 sequencing. Each point represents the

average expression of a single gene assayed in the indicated assays. At the top left, the

Pearson correlation coefficient and resulting p value are indicated.

Supplemental Figure 2: A. Comparison of performance metrics (average distance, AD;

average distance between means, ADM; average proportion of non-overlap, APN;

connectivity; Dunn Index; figure of merit, FOM; and silhouette index) by several

clustering methods (diana, model-based, hierarchical, kmeans, and pam) run on

sgmRNA expression of 600 randomly sampled cells analyzed with 10X 5′ Extended R1

and scCoVseq. *Left*: Performance metrics for each method across k values from 2 to 5.

Right: performance metrics for each method with k = 2. **B.** Visualization of infection

classification by different methods. **C.** Viral gene expression of cells by infection status,

determined by pam clustering method. **D.** Percent of infected cells per sample as

measured by flow cytometry, immunofluorescence, and infection classification with

unsupervised (pam) method or supervised infection classification by classifying infected

cells as those with at least 375 total viral UMIs. Because the same sample was

sequenced with 10X 5′ and 10X 5′ Extended R1, flow cytometry and

immunofluorescence results are duplicated for ease of visualization. Error bars for

immunofluorescence indicate mean ± one standard deviation of percent infected cells

based on three fields per sample.

Supplemental Figure 3: Detection of junction sites in SARS-CoV-2 reads with 10X 5′

Extended R1. Junction sites are represented by the 5′ start site and 3′ end site on the y

and x-axis, respectively. Below each axis, the number of UMIs supporting a position as

a junction start or end site is indicated with a density plot.