Spatial metabolomics reveals localized impact of influenza virus infection on the lung

tissue metabolome

Danya A. Dean,[a,b] London Klechka,[c] Ekram Hossain,[a,b] Adwaita R. Parab,[d] Krystin Eaton,[e]

Myron Hinsdale,[e] Laura-Isobel McCall[a,b,d#]

[a]Department of Chemistry and Biochemistry, University of Oklahoma, Norman, Oklahoma

[b]Laboratories of Molecular Anthropology and Microbiome Research, University of Oklahoma,

Norman, Oklahoma

[c]Department of Biology, University of Oklahoma, Norman, Oklahoma

[d]Department of Microbiology and Plant Biology, University of Oklahoma, Norman, Oklahoma

[e]Department of Physiological Sciences, Oklahoma State University, Stillwater, Oklahoma

#Address correspondence to Laura-Isobel McCall, lmccall@ou.edu

Running Head title: Spatial metabolomics of IAV infection

Abstract

The influenza virus (IAV) is a major cause of respiratory disease, with significant

infection increases in pandemic years. Vaccines are a mainstay of IAV prevention, but are

complicated by consideration of IAV's vast strain diversity, manufacturing and vaccine uptake

limitations. While antivirals may be used for treatment of IAV, they are most effective in early

stages of the infection and several virus strains have become drug resistant. Therefore, there is

a need for advances in IAV treatment, especially host-directed, personalized therapeutics.

Given the spatial dynamics of IAV infection and the relationship between viral spatial distribution

and disease severity, a spatial approach is necessary to expand our understanding of IAV

pathogenesis. We used spatial metabolomics to address this issue. Spatial metabolomics

combines liquid chromatography-tandem mass spectrometry of metabolites extracted from

systematic organ sections, 3D models and computational techniques, to develop spatial models

of metabolite location and their role in organ function and disease pathogenesis. In this project,

we analyzed plasma and systematically sectioned lung tissue samples from uninfected or

infected mice. Spatial mapping of sites of metabolic perturbations revealed significantly lower

metabolic perturbation in the trachea compared to other lung tissue sites. Using random forest

machine learning, we identified metabolites that responded differently in each lung position

based on infection, including specific amino acids, lipids and lipid-like molecules, and

nucleosides. These results support the implementation of spatial metabolomics to understand

metabolic changes upon IAV infection and to identify candidate pathways to be targeted for IAV

therapeutics.

Importance

The influenza virus is a major health concern. Over 1 billion people become infected

annually despite the wide distribution of vaccines, and antiviral agents are insufficient to address

current clinical needs. In this study, we used spatial metabolomics to understand changes in the

lung and plasma metabolome of mice infected with influenza A virus, compared to uninfected

controls. We determined metabolites altered by infection in specific lung tissue sites and

distinguished metabolites perturbed by infection between lung tissue and plasma samples. Our

findings highlight the importance of a spatial approach to understanding the intersection

between lung metabolome, viral infection and disease severity. Ultimately, this approach will

expand our understanding of respiratory disease pathogenesis and guide the development of

novel host-directed therapeutics.

Introduction

Influenza virus outbreaks are a continuous public health issue. Seasonal global

epidemics caused by both influenza A viruses (IAV) and influenza B viruses cause 300,000-

500,000 deaths each year (1). Vaccinations are the current method of prevention, but they fail to

account for every possible viral strain. Antiviral drugs are used for treatment of IAV but are most

effective within a short window during early infection. Additionally, it is believed that some

strains have developed resistance to these drugs (2). One study indicated that while influenza

A(H1N1)pdm09-infected intensive care unit patients treated with neuraminidase inhibitors have

greater survival rates than untreated patients, one in four treated patients still die (3). These

findings indicate a strong need for new treatments for IAV infection, and the potential for host-

targeted therapeutics to supplement antiviral agents. Their development, however, necessitates

an understanding of disease pathogenesis, which remains incompletely elucidated for IAV.

We used metabolomics to identify and analyze metabolites affected by IAV infection.

Metabolomics is a method of analysis that focuses on small molecules involved in biological

processes. This technique allows us to gain insight on the host chemical response to viral

infection. A comprehensive understanding of the relationship between host and virus could aid

in the development of more effective prevention and treatment options. Previous studies applied

metabolomic methods to lung tissue and serum during IAV infection. These studies found that

nucleosides such as uridine, lipids such as sphingosine, sphinganine, and amino acid

metabolites such as kynurenine are increased during infection in lung tissue (4). Additionally,

carbohydrates such as mannitol, myo-inositol and glyceric acid are decreased during active

infection. However, location of IAV within the respiratory tract is dynamic. Viral localization and

location of tissue damage within the respiratory tract also influences disease symptoms, disease

severity and transmissibility of the infection (5, 6, 7, 8). Thus, a spatial perspective is necessary

with regards to IAV and tissue metabolism. Chemical cartography is an approach that combines

liquid chromatography-mass spectrometry (LC-MS) with 3D visualizations, leading to detailed

spatial maps of metabolite distribution compared to pathogen load, tissue damage, or immune

responses (9, 10, 11). This approach, when applied to other infectious diseases enabled the

discovery of new treatments for these conditions (10). This method has been used to study the

impact of cystic fibrosis on the local lung metabolome (12, 13), but had not previously been

applied to IAV infection.

We therefore hypothesized that spatial metabolomics could provide new insights

regarding IAV infection. Using this method, we analyzed the distribution of small (*m/z* 100-

1,500) metabolites within the infected trachea and lung, in comparison to plasma samples and

to uninfected animals. We identified changes in the lung metabolome and determined limited

overlap in metabolites perturbed by IAV infection between lung tissue and plasma. Additionally,

we identified several metabolites altered by infection such as amino acids, lipids and lipid-like

molecules, and nucleosides. Interestingly, these metabolites were differentially affected at each

lung position. Ultimately, our study highlights the application of spatial metabolomics to

understand IAV infection and further our understanding of respiratory disease pathogenesis, to

guide the development of novel host-directed therapeutics.

Results

Viral distribution influences IAV transmissibility, viral reassortment and disease severity (5, 14, 15). While a few studies have investigated the changes in the metabolome during IAV infection, a spatial perspective of the metabolic disturbances is lacking (4, 16–18). We therefore used spatial metabolomics to identify candidate pathways and metabolites altered by infection in specific lung locations. Plasma, trachea and lungs were collected from IAV-infected mice at 3 days post-infection. Lungs were systematically sectioned into 11 segments (Fig. 1A), and all samples analyzed by liquid chromatography-tandem mass spectrometry (LC-MS/MS), followed by 3D reconstruction of metabolomics data. In addition, tissue homogenate bioluminescence was measured as an indicator of local viral burden in each tissue segment.

Using principal coordinate analysis (PCoA), we first compared lung and plasma metabolomes and found that the local lung tissue metabolome does not reflect the circulating metabolome (PERMANOVA <0.001, Fig 1D). Furthermore, lung tissue overall was found to be greatly impacted by infection (PERMANOVA <0.001, R^2 0.093931) (Fig 1E). A similar but non-significant, trend was seen between infected and uninfected plasma samples (Fig 1F).

Spatial analyses of lung tissue showed that viral load was largely localized to the lung tissue (positions 1-11) with minimal viral load in the trachea (position 12) (Dunn's test $p<0.05$, FDR-corrected, for comparisons between trachea and all positions except left lung middle (position 3, Fig 1B)). Overall local impact of infection on the metabolome was quantified using PERMANOVA R^2 at each sampling site. Magnitude of metabolic perturbation was variable between tissue segments, with the highest degree of metabolic perturbation in two segments of the left lung (position 1 and position 4), and the right lung middle lobe (position 8), whereas the trachea metabolome was least affected (R^2 range from 0.05 in trachea to 0.18 at position 4; Fig 1C and Table S15). Thus, sites of highest and lowest metabolic perturbation match sites with

high vs low viral load, respectively, enabling insight into the specific metabolites most affected

by infection.

We next sought to determine whether the specific metabolites perturbed by infection

differed between lung and plasma (Fig 1G-I). We used a random forest classifier, applied to

plasma on the one hand, and to lung tissue on the other hand (all positions combined). After

applying significance cutoffs (see Materials and Methods), this approach yielded a total list of 9

plasma metabolites and 374 lung tissue metabolites significantly perturbed by infection. There

was strikingly limited overlap of infection-perturbed metabolites between lung tissue samples

and plasma samples, indicating that both sites respond differentially to infection (Fig 1H) and

concurring with overall PCoA analysis findings (Fig 1D). We then sought to assess whether the

metabolites perturbed by infection were uniquely elicited by infection, or found under both

conditions but at differential levels. The majority of infection-perturbed metabolites were

common to infected and uninfected samples but found at different levels, with a minority

uniquely detected in uninfected samples only (Fig 1I). A similar trend was seen in infected and

uninfected plasma samples, where most infection-perturbed metabolites were present in both

infected and uninfected samples, albeit at different levels (Fig 1G). This signifies that the effect

of IAV infection on the metabolome is primarily on metabolite levels rather than induction of

novel metabolites.

Figure 1. Localized impact of IAV infection in lung tissue and plasma. A: 3D model of lung tissue showing sampling positions. **B:** Median viral burden distribution of IAV. Lower infection levels were observed in the trachea (Dunn's test, FDR-corrected, $p < 0.05$ for comparisons between trachea and all positions except left lung middle (position 3)). **C:** Magnitude of metabolic perturbation compared to uninfected (PERMANOVA R^2). **D:** Principal coordinate analysis (PCoA) plot showing differences in overall metabolome by sample type (PERMANOVA p-value <0.001). **E:** Impact of infection on overall lung tissue metabolome (PERMANOVA p-value <0.001). **F:** No significant impact of infection on the overall plasma metabolome by PCoA. **G-I:** Unique and common metabolites perturbed by infection **G:** Most metabolites perturbed by infection in plasma samples are found in both infected and uninfected samples, albeit at different levels. 4/8 metabolites were increased by infection and 4/8 were decreased by infection, while only one of the statistically significant metabolites was uniquely observed only in uninfected plasma samples. **H:** Overlap of metabolites perturbed by infection in plasma (orange) or lung tissue all positions combined (green). **I:** Most metabolites perturbed by infection in lung samples are found in both infected

and uninfected samples, albeit at different levels (181/382 metabolites increased by infection and 181/362

decreased by infection, with 11 of the statistically significant metabolites uniquely detected in uninfected

lung tissue).

Random forest classifier was also used to analyze the common and unique responses of

each lung and tracheal tissue site to infection. A random forest model was built for each lung

position and for plasma, classifying infected versus uninfected samples. Strikingly, most

infection-impacted metabolites were only affected at one or a few tissue sites (Fig 2A). This is

not due to divergence in overall metabolome between sites, as analysis of all metabolites,

irrespective of abundance, revealed large commonality across lung tissue sites and plasma (Fig

2B). Likewise, infection-perturbed metabolites do not overlap appreciably with metabolites

differing in abundance between lung and plasma (Fig S3). Very few perturbed metabolites were

identified for position 12 (trachea), perhaps as a consequence of the low viral load at that site

(Fig 1B).

Figure 2: Location-specific impact of IAV infection on metabolism. A: UpSet plot (Number of intersections to show: 60) showing uniqueness of statistically significant infection-impacted metabolites in lung tissue and plasma perturbed, based on 13 random forest models analyzing each lung segment and plasma separately. B: UpSet plot (Number of intersections to show: 50) of total metabolites found in lung tissue and plasma showing large commonality across tissue sites and plasma. Position numbers as in Fig. 1A.

Metabolites perturbed by infection were annotated using molecular networking (19).

Among these, carnitine, glutamine, kynurenine, and cytosine were found to be significantly and

markedly perturbed by IAV infection and in a spatial manner (Fig 3, Fig S1). Glutamine,

cytosine, and kynurenine were all significantly increased by IAV infection at all tissue sites

except the trachea (Wilcoxon FDR-corrected p-value <0.05 at all sites except trachea) (Fig 3,

Fig S1). The opposite trend was seen for carnitine which was decreased by infection at all

tissue sites except the trachea (Wilcoxon FDR-corrected p-value <0.05 at all sites except

trachea) (Fig 3, Fig S1). In contrast, these metabolites were not significantly affected by

infection in the plasma.

Figure 3: Representative metabolites perturbed by IAV infection. Molecular networks display summed peak area of metabolites in infected (red) and uninfected (blue) samples. Each connected node represents structurally-related metabolites, as determined by molecular networking. 3D lung 'ili plots show median peak area of each displayed metabolite across tissue sites. Wilcoxon FDR-corrected p-values comparing matched infected and uninfected lung tissue sites *<0.05; **<0.01; *** <0.001.

Additional metabolites perturbed by infection in plasma and lung tissue include amino acids, acylcarnitines and nucleobases (Fig 4, Table S1, Table S2), with contrasting effects between lung tissue positions and sample types. For example, amino acids had dissimilar effects following IAV infection in plasma and lung tissue samples (Fig 4): methionine was decreased in infected plasma samples (Wilcoxon FDR-corrected p-value <0.001) while phenylalanine and L-citrulline were increased in infected lung tissue samples (Wilcoxon FDR-corrected p-value <0.05) (Fig 4 A, E-F). However, the dipeptide methionine-phenylalanine was decreased in infected lung tissue (Wilcoxon FDR-corrected p-value <0.05) (Fig 4K). Acylcarnitines (oleylcarnitine and palmitoylcarnitine) were significantly increased in infected plasma samples (Wilcoxon FDR-corrected p-value <0.001) compared to uninfected samples (Fig 4B-C). Guanine, S-adenosy-homocyteine, spermidine, 4-hydroxynonenal, and hemin cation were all significantly increased upon infection (Wilcoxon FDR-corrected p-value <0.05) at select lung tissue sites (Fig 4F-H, G and L). Infection had the opposite effect on palmitamide, which was significantly decreased in infected tissue samples (Fig 4I). In contrast, none of these metabolites apart from phenylalanine were significantly affected by infection in the trachea.

Fig 4: Representative metabolites perturbed by IAV infection in plasma and lung tissue. A-C:

Statistically significant metabolites altered by infection in plasma (Wilcoxon FDR-corrected p-value

<0.001). **A:** Methionine (*m/z* 150.058, RT 0.323 min). **B:** Oleyl L-carnitine (*m/z* 426.357, RT 2.982 min).

C: Palmitoylcarnitine (*m/z* 400.341, RT 2.952 min). **D-L:** Statistically significant metabolites altered by

infection in lung tissue. **D:** Phenylalanine (*m/z* 166.042, RT 2.246 min) statistically different at all positions

(Wilcoxon FDR-Corrected p-value <0.05) except positions 2, 5, 10 and 12. **E:** Citrulline (*m/z* 198.085, RT

0.309 min) statistically significant at all positions (Wilcoxon FDR-Corrected p-value <0.05) except position

12. **F:** Guanine (*m/z* 150.977, RT 0.291 min) statistically different at all positions (Wilcoxon FDR-

Corrected p-value <0.05) except position 12. **G:** S-Adenosyl-homocysteine (*m/z* 385.212, RT 2.166 min)

statistically significant at all positions (Wilcoxon FDR-Corrected p-value <0.05) except position 12. **H:**

Spermidine (*m/z* 146.165, RT 0.287 min) statistically significant at all positions (Wilcoxon FDR-Corrected

p-value <0.05) except positions 3, 10, and 12. **I:** Palmitamide (*m/z* 256.073 RT 2.563 min) statistically

significant at all positions (Wilcoxon FDR-Corrected p-value <0.05) except positions 2, 5, 6, 7, 11, and 12.

J: 4-Hydroxynonenal (*m/z* 139.112, RT 2.644 min) statistically significant at all positions (Wilcoxon FDR-

Corrected p-value <0.05) except positions 1, 2, 3, 5, 11, and 12. **K:** Met-Phe (*m/z* 297.123, RT 2.444 min)

statistically significant at all positions (Wilcoxon FDR-Corrected p-value <0.05) except positions 3, 4, 6,

and 7. L: Hemin cation (*m/z* 616.358, RT 2.994 min) statistically significant at all positions (Wilcoxon FDR-

Corrected p-value <0.05) except positions 3, 4, 9, and 10.

Discussion

Here, we generated spatial maps of the metabolic impact of IAV infection on the mouse

lung and trachea. This approach revealed differential effects of infection across tissue sites and

between lung and plasma, as well as differential viral burden between trachea and lung tissue.

Based on random forest analysis and molecular networking, infection-impacted metabolites of

biological significance were annotated as acylcarnitines, amino acids, phospholipids, and

nucleotides, amongst others.

L-carnitine was significantly decreased by IAV infection at all lung positions except the

trachea (Fig 3 top left). Likewise, short chain acylcarnitines such as CAR 2:0 and CAR 6:0 were

also decreased in lung tissue (Table S3-14). In contrast, in both plasma and lung tissue,

different long chain acylcarnitines were increased by infection, including oleyl carnitine (CAR

18:1) and palmitoylcarnitine (CAR 16:0) in plasma and CAR 14:1 and CAR 20:1 in lung tissue

(Fig 4B-C, Table S1-14). Carnitines and acylcarnitines are key intermediates in energy

production via fatty acid beta oxidation, and influenza virus replication is sensitive to fatty acid

beta oxidation activity (20),(21). Likewise, these metabolic alterations may also be contributing

to the differential responses to vaccination in obese vs lean animals (22).

Our study indicated that glutamine is increased upon IAV infection (Fig 3 top right). T cell

proliferation and cytokine secretion relies heavily on glutamine presence (23). Thus, elevated

glutamine levels may enable anti-viral immunity IAV-infected cells are also more dependent on

glutamine availability than uninfected cells for survival, suggesting a pro-survival effect of our

observed elevated glutamine levels (24). Phenylalanine and citrulline were increased in infected

lung tissue, in contrast with methionine, which was decreased in plasma (Fig 4D-E). Both

phenylalanine and citrulline are involved in immune response by aiding in T-cell function (25).

Phenylalanine aids in activation of T-cells while citrulline has downstream effects through the

production of arginine which is used for T-cell growth and response (25–27). Upregulation of

both amino acids suggest their role in overall immune response to IAV.

Nucleotides, cytosine and guanine were elevated by infection in all lung sites except

trachea (Fig 3 bottom left and Fig 4F). Pyrimidine nucleotide biosynthesis is elevated upon IAV

infection, and indeed its replication was dependent on pyrimidine biosynthesis (28–30).

Interestingly, cytosine was found discriminatory in the plasma between RT-PCR positive COVID

patients and RT-PCR COVID negative patients (29).

Several of the metabolites annotated and identified as infection-impacted in the lung

tissue and plasma in this study are congruent with prior studies of respiratory infection (4).

Amino acids and nucleotides in particular were upregulated in lung tissue in other IAV studies

as well as in *Mycobacterium tuberculosis* infection (TB) and respiratory syncytial virus (RSV)

studies (4, 31–33). Phenylalanine was increased in mouse lung tissue in another IAV study and

in the serum and lung of mice infected with TB, while L-citrulline is elevated in mouse lung

tissue infected with RSV and in mice infected with TB (4, 17, 31, 32, 34). This coincides with our

findings showing elevations of these amino acids in IAV-infected lung tissue (Fig 4D and E).

Glutamine was identified in mouse lung tissue infected with TB and was elevated alongside

other immune response amino acids, corresponding with our study (Fig 3 top right) (32, 34).

Kynurenine, an infection-induced anti-inflammatory molecule, is consistently upregulated in

mouse lung tissue of several respiratory infection studies including TB, RSV, and our IAV study

(Fig 3 bottom left) (4, 31, 33, 34). Upregulation in cytosine in the lung tissue, oleylcarnitine in

plasma, and several phospholipids in the lung tissue are also consistent with current literature

(Fig 3 bottom right, Fig 4B and Table S1) (4, 17, 33).

Analysis of circulating metabolites has been performed in multiple studies on IAV

vaccinology (35), (36). Our findings of differential impact of infection on lung and plasma

metabolites (Fig. 1H, Table S1, Table S2) indicate that the metabolic changes observed in those

studies may not be directly linked to lung metabolic patterns, and this discrepancy was not due

purely to differences between plasma and lung overall (Fig. 2B). Likewise, the majority of

COVID-19 metabolomic studies have relied on serum or plasma samples (37). By extension,

based on this study's results, they may not be relevant to the pathogenesis of COVID-19 in the

lung, hampering translatability for drug development purposes. Thus, studies seeking to build on

metabolomics to design new treatments should rely on analysis of the affected organ, rather

than biofluids, even if the latter are more readily available.

We also observed differential impact of infection across lung sections, with the majority

of infection-perturbed metabolites only significantly perturbed at one lung segment (Fig. 2A).

The trachea in particular was especially divergent from the lung lobes in terms of viral burden

(Fig. 1B), overall magnitude of infection-induced metabolic perturbations (Fig. 1C) and specific

metabolic changes (Fig. 3, Fig. 4). Jointly, these results highlight the strength of our spatial

perspective.

As with any untargeted metabolomics studies, a significant fraction of infection-impacted

metabolites could not be annotated (~65%). The most commonly observed metabolite

subclasses in our dataset overall were amino acids, peptides, and analogues,

glycerophosphocholines, amines and fatty amides. Although this represents a broad diversity of

metabolite classes, nevertheless complementary metabolite extraction or data acquisition

methods could further expand this list. We further acknowledge that the mouse model of

influenza virus infection and mouse-adapted influenza viral strain may not be the most

representative of the functional lung alterations that would occur during human infection (38).

Indeed, the predominance of lower respiratory tract metabolic alterations over changes in the

trachea observed in this study. may be a consequence of this model. A further limitation is that

we focused on a single time point, in a lethal infection model. However, our findings serve as

proof-of-concept of the applicability of our approach to study respiratory viruses, with future

work applying this method in other biological systems.

We anticipate our findings to serve as a resource upon which the research community

can build to study the impact of different disease modifiers on the relationship between spatial

changes in the lung metabolome and disease severity, for example the impact of age,

comorbidities or treatment. Our findings and our approach also serve as a framework to study

how the metabolome is restored in a spatially-dependent fashion during recovery from

respiratory infection, or fails to recover during chronic disease in additional disease models, and

to identify markers of treatment response and infection outcome. Overall, we anticipate our

approach to be broadly applicable to many other respiratory infections, helping expand our

understanding of respiratory disease pathogenesis and to drive the development of host-

targeted therapeutic regimens.

Materials and Methods

In vivo infection

All vertebrate animal studies were performed under a protocol approved by Oklahoma State University Institutional Animal Care and Use Committee (protocol number VM20-36), in accordance with the USDA Animal Welfare Act and the Guide for the Care and Use of Laboratory Animals of the National Institutes of Health.

Female C57BL/6J mice were anesthetized with ketamine and xylazine intraperitoneally and then intranasally inoculated with PR8-Glu (39) (a generous gift from Dr. Peter Palese, Icahn School of Medicine at Mount Sinai, Department of Microbiology, New York, New York, USA) at 2×10^3 pfu/mouse in 25-50 µl PBS. Controls were inoculated with PBS alone. Body weights were monitored daily. Three days later, mice were anesthetized with ketamine and xylazine and injected IV with a working solution of coelenterazine (GoldBio, St. Louis, Mo.) made according to standard protocol (40). Briefly, using a stock solution of 7.5 mg/ml in acidified alcohol, each mouse was administered 98 µg substrate in PBS (13 µl stock in 137 µl PBS) into the retroorbital sinus. After substrate administration, the thoracic cavity was opened followed by exsanguination by cardiac blood collection for serum and by lung tissue harvesting. Upper trachea, cranial to tracheal bifurcation, was also collected. Subsequently, samples were submerged in coelenterazine 0.3 mg/ml in a 96 well plate. Once collected the tissue pieces were blotted off and flash frozen in liquid nitrogen for later analyses.

Metabolite extraction and UHPLC-MS/MS

A two-step metabolite extraction procedure was performed according to Want *et al.*(41) for both tissue and plasma samples. Samples were homogenized in LC-MS grade water with steel beads utilizing a Qiagen TissueLyzer at 25 Hz for 3 min, and 1 µl removed for luminescence analysis. Methanol was added for a final concentration of 50%, samples

homogenized again for 3 mins and centrifuged at 16,000xg for 10 mins at 4ºC. The supernatant

(aqueous extract) was collected, dried overnight in a Speedvac and frozen at -80ºC until LC-MS

analysis. The pellet produced from centrifugation was collected for organic extraction via

addition of 3:1 dichloromethane:methanol, homogenized for 5 mins and centrifuged for 10 mins

at 4ºC. The organic extract was air dried overnight and then frozen at -80ºC until LC-MS

analysis. Dried aqueous and organic extracts were resuspended in 1:1 methanol and water

spiked with the internal standard sulfadimethoxine and combined. Samples were then

sonicated, centrifuged, and the supernatant collected for analysis. A Thermo Scientific Vanquish

UHPLC system was used for tissue and plasma analysis using a Kinetex 1.7µm C8 100 Å LC

column (50 x 2.1mm). Chromatography was done with water + 0.1% formic acid (mobile phase

A) and acetonitrile + 0.1% formic acid (mobile phase B), at a 0.5 mL/min flow rate (7.5 mins)

with a 40°C column temperature. LC gradient can be found in Table 1. Data acquisition was

performed in random sample order, with a blank and pooled quality control every 12 samples.

To monitor instrumental drift, a 6-mix solution with 6 known molecules was run at the beginning

and end of LC/MS/MS analysis. Calibration of the instrument was also done immediately prior to

instrument analysis using Pierce LTQ Velos ESI positive ion calibration solution. MS/MS

detection was conducted on a Q Exactive Plus (Thermo Scientific) high resolution mass

spectrometer (Table 2). Ions were generated for MS/MS analysis in positive mode.

Table 1: LC Gradient

Time(min)	Flow (ml/min)	%B	Curve
0.000	Run		
0.000	0.500	2.0	5
1.000	0.500	2.0	5
2.500	0.500	98.0	5

4.500	0.500	98.0	5
5.500	0.500	2.0	5
7.500	0.500	2.0	5
7.500	Stop Run		

348   Table 2: Q Exactive Plus (Thermo Scientific) instrument parameters

Runtime	0 to 7.5 min
Polarity	Positive
Default charge state	1
Full MS	
Resolution	70,000
AGC target	3e6
Maximum IT	246 ms
Scan range	100 to 1500 *m/z*
dd-MS2/dd-SIM	
Resolution	17,500
AGC target	1e5
Maximum IT	54 ms
Loop count	5
TopN	5
Isolation window	1.0 *m/z*
(N)ce/stepped (N)CE	NCE: 20, 40, 60
dd Settings	
Minimum AGC target	8.00e3
Intensity Threshold	1.5e5
Peptide match	preferred

Exclude isotopes	on
Dynamic exclusion	10.0s
Tune Data	
Spray Voltage (+)	3800.00
Capillary Temperature (+ or +-)	320.00
Sheath Gas (+ or +-)	35.00
Aux Gas (+ or +-)	10.00
Spare Gas (+ or +-)	0.00
Max Spray Current (+)	100.00
Probe Heater Temp (+ or +-)	0.00
S-Lens RF Level	50.00
Ion Source	HESI

Luminescence analysis

Coelenterazine was prepared according to GoldBio standard protocols. Briefly, 1 mg

coelenterazine was added to 1 mL of acidified methanol to make a stock solution. The stock

solution was made to a final concentration of 1.5 µM. A 1:10 solution of sample homogenate to

colentrazine was analyzed on a GloMax Explorer (Promega).

LC-MS Data analysis

Data analysis was performed using MZmine version 2.53, according to Table 3

parameters, to develop the feature table for further analysis. Blank removal with a 3-fold

threshold was performed and Jupyter notebooks were used to perform total ion current (TIC)

normalization. Principal coordinate analysis (PCoA) was performed on the TIC-normalized MS1

feature table using the Bray-Curtis dissimilarity metric in QIIME2 (42). Three-dimensional PCoA

plots were developed and visualized using EMPeror (43). Lung three-dimensional model

developed using Sketchup and Mesh lab, and modelling was completed using 'ili

(http://ili.embl.de/) (44).

Random forest analysis was conducted using R in Jupyter notebooks. Number of trees

were restricted to 500 and random forest classifier cutoff was based on variable importance

score of mean decrease accuracy >1. Lists were further restricted to FDR-corrected Mann-

Whitney p-value less than 0.05 and fold change <0.05 or >2.0. Venn diagrams and UpSet plots

were developed to quantify metabolite overlap within lung tissue positions and between lung

tissue and plasma using Intervene Shiny App (45).

Global Natural Products Social Molecular Networking (GNPS) was used to perform

feature-based molecular networking and MolNetEnhancer according to the parameters in Table

4. Cytoscape 3.8.2. was used to visualize all molecular networks. All reported annotations are at

Metabolomics Standards Initiative confidence level 2 (specific metabolite name provided) or

level 3 (metabolite family name provided only) (46). Lipids were annotated based on GNPS

library or analog matches, and using standard LipidMAPS nomenclature (47).

Table 3: MzMine 2.53 parameters

MS^1	
Retention Time (min)	0-7.5
Noise Level	5E5
MS^2	
Retention Time (min)	0-7.5
Noise Level	1E3
Chromatogram Builder	
Mass list	Masses
Minimum Time span (min)	0.01
Minimum Height	1.5E6

m/z Tolerance (ppm)	10.0
Algorithm	Baseline cut-off
Chromatogram Deconvolution	
m/z Range for MS² scan pairing (Da)	0.001
RT Range for MS² Scan pairing (min)	0.2
Minimum peak height	1.5E6
Peak duration range (min)	0-3.5
Baseline level	5E5
Deisotoping	
m/z tolerance (ppm)	10.0
Retention time tolerance (min)	0.5
Monotonic shape	Yes
Maximum Charge	3
Representative Isotope	Most intense
Alignment	
m/z Tolerance (ppm)	10.0
Weigh for *m/z*	1
Weight for Retention time	1
Retention time tolerance (min)	0.5
Row filtering	
Minimum peaks in a row	3
Retention time (min)	0.2-6.5
Keep only peaks with MS² scans	YES
Reset Peak No. ID	YES

380   Table 4: GNPS Parameters

Feature-Based Molecular Network	
Precursor Ion Mass Tolerance	0.02
Fragment Ion Mass Tolerance	0.02
Minimum Matched Fragment Ions	4
Maximum Connected Component Size (Beta)	100
Maximum shift between precursors	500
Library Search Min Matched Peaks	4
Search analogs	Do Search
Top results to report per query	1
Score Threshold	0.7
Maximum analog difference	100.0 Da
Minimum Peak Intensity	0.0
Filter Precursor Window	Filter
Filter peaks in 50 Da Window	Filter
Filter Library	Filter Library
Normalization per file	Row Sum Normalization (Per file Sum to 1,000,000)
Aggregation Method for peak abundances per group	Mean
PCoA	braycurtis
Run Dereplicator	Run

Data availability: All metabolomics data is publicly available in MassIVE under

accession number MSV000085389 (ftp://massive.ucsd.edu/MSV000085389).

MOLNet Enhancer link:

https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=f9f194c9b723409f8927c84470f9a0c5

Original GNPS link:

https://gnps.ucsd.edu/ProteoSAFe/status.jsp?task=bb7a5f7fb32045208b6040908a1453f7

Acknowledgments

This project was supported by a pilot grant from the Oklahoma Center for Respiratory

and Infectious Diseases (OCRID) under NIH award number P20GM103648 and used the

OCRID Animal Model Core, supported by the National Institute of General Medical Sciences

under award number P20GM103648. The authors further wish to acknowledge partial support

from NIH award number R21AI148886 and to thank Dr. Peter Palese, Icahn School of Medicine

at Mount Sinai, Department of Microbiology, New York, New York, USA, for providing the PR8-

Glu viral strain. Laura-Isobel McCall, Ph.D. holds an Investigators in the Pathogenesis of

Infectious Disease Award from the Burroughs Wellcome Fund. The content is solely the

responsibility of the authors and does not necessarily represent the official views of the National

Institutes of Health or any of the other funders.

References

1. Erbelding EJ, Post DJ, Stemmy EJ, Roberts PC, Augustine AD, Ferguson S, Paules CI,

Graham BS, Fauci AS. 2018. A Universal Influenza Vaccine: The Strategic Plan for the

National Institute of Allergy and Infectious Diseases. J Infect Dis 218:347–354.

2. 2012. Suppressing production of reactive oxygen species (ROS) for influenza A virus

therapy. Trends Pharmacol Sci 33:3–8.

3. Louie JK, Yang S, Acosta M, Yen C, Samuel MC, Schechter R, Guevara H, Uyeki TM.

2012. Treatment with neuraminidase inhibitors for critically ill patients with influenza A

(H1N1)pdm09. Clin Infect Dis 55:1198–1204.

4. Bernatchez JA, McCall L-I. 2020. Insights gained into respiratory infection pathogenesis

using lung tissue metabolomics. PLoS Pathog 16:e1008662.

5.   Richard M, van den Brand JMA, Bestebroer TM, Lexmond P, de Meulder D, Fouchier RAM,

Lowen AC, Herfst S. 2020. Influenza A viruses are transmitted via the air from the nasal

respiratory epithelium of ferrets. Nat Commun 11:766.

6.   van Riel D, Munster VJ, de Wit E, Rimmelzwaan GF, Fouchier RAM, Osterhaus ADME,

Kuiken T. 2007. Human and avian influenza viruses target different cells in the lower

respiratory tract of humans and other mammals. Am J Pathol 171:1215–1223.

7.   Karlsson EA, Meliopoulos VA, Savage C, Livingston B, Mehle A, Schultz-Cherry S. 2015.

Visualizing real-time influenza virus infection, transmission and protection in ferrets. Nat

Commun 6:6378.

8.   Tran V, Moser LA, Poole DS, Mehle A. 2013. Highly sensitive real-time in vivo imaging of

an influenza reporter virus reveals dynamics of replication and spread. J Virol 87:13321–

13329.

9.   Dean DA, Gautham, Siqueira-Neto JL, McKerrow JH, Dorrestein PC, McCall L-I. Spatial

metabolomics identifies localized chemical changes in heart tissue during chronic cardiac

Chagas disease.

10. Hossain E, Khanam S, Dean DA, Wu C, Lostracco-Johnson S, Thomas D, Kane SS, Parab

AR, Flores K, Katemauswa M, Gosmanov C, Hayes SE, Zhang Y, Li D, Woelfel-Monsivais

C, Sankaranarayanan K, McCall L-I. 2020. Mapping of host-parasite-microbiome

interactions reveals metabolic determinants of tropism and tolerance in Chagas disease.

Sci Adv 6:eaaz2015.

11. McCall L-I, Morton JT, Bernatchez JA, de Siqueira-Neto JL, Knight R, Dorrestein PC,

McKerrow JH. 2017. Mass Spectrometry-Based Chemical Cartography of a Cardiac

Parasitic Infection. Anal Chem 89:10414–10421.

12. Garg N, Wang M, Hyde E, da Silva RR, Melnik AV, Protsyuk I, Bouslimani A, Lim YW,

Wong R, Humphrey G, Ackermann G, Spivey T, Brouha SS, Bandeira N, Lin GY, Rohwer

F, Conrad DJ, Alexandrov T, Knight R, Dorrestein PC. 2017. Three-Dimensional

Microbiome and Metabolome Cartography of a Diseased Human Lung. Cell Host Microbe

22:705–716.e4.

13. Melnik AV, Vázquez-Baeza Y, Aksenov AA, Hyde E, McAvoy AC, Wang M, da Silva RR,

Protsyuk I, Wu JV, Bouslimani A, Lim YW, Luzzatto-Knaan T, Comstock W, Quinn RA,

Wong R, Humphrey G, Ackermann G, Spivey T, Brouha SS, Bandeira N, Lin GY, Rohwer

F, Conrad DJ, Alexandrov T, Knight R, Dorrestein PC, Garg N. 2019. Molecular and

Microbial Microenvironments in Chronically Diseased Lungs Associated with Cystic

Fibrosis. mSystems 4.

14. Richard M, Herfst S, Tao H, Jacobs NT, Lowen AC. 2018. Influenza A Virus Reassortment

Is Limited by Anatomical Compartmentalization following Coinfection via Distinct Routes.

Journal of Virology.

15. Yan J, Grantham M, Pantelic J, Bueno de Mesquita PJ, Albert B, Liu F, Ehrman S, Milton

DK, EMIT Consortium. 2018. Infectious virus in exhaled breath of symptomatic seasonal

influenza cases from a college community. Proc Natl Acad Sci U S A 115:1081–1086.

16. Tisoncik-Go J, Gasper DJ, Kyle JE, Eisfeld AJ, Selinger C, Hatta M, Morrison J, Korth MJ,

Zink EM, Kim Y-M, Schepmoes AA, Nicora CD, Purvine SO, Weitz KK, Peng X, Green RR,

Tilton SC, Webb-Robertson B-J, Waters KM, Metz TO, Smith RD, Kawaoka Y, Suresh M,

Josset L, Katze MG. 2016. Integrated Omics Analysis of Pathogenic Host Responses

during Pandemic H1N1 Influenza Virus Infection: The Crucial Role of Lipid Metabolism. Cell

Host Microbe 19:254–266.

17. Cui L, Zheng D, Lee YH, Chan TK, Kumar Y, Ho WE, Chen JZ, Tannenbaum SR, Ong CN.

2016. Metabolomics Investigation Reveals Metabolite Mediators Associated with Acute

Lung Injury and Repair in a Murine Model of Influenza Pneumonia. Scientific Reports.

18. Milner JJ, Wang J, Sheridan PA, Ebbels T, Beck MA, Saric J. 2014. 1H NMR-based

profiling reveals differential immune-metabolic networks during influenza virus infection in

obese mice. PLoS One 9:e97238.

19. Wang M, Carver JJ, Phelan VV, Sanchez LM, Garg N, Peng Y, Nguyen DD, Watrous J,

Kapono CA, Luzzatto-Knaan T, Porto C, Bouslimani A, Melnik AV, Meehan MJ, Liu W-T,

Crüsemann M, Boudreau PD, Esquenazi E, Sandoval-Calderón M, Kersten RD, Pace LA,

Quinn RA, Duncan KR, Hsu C-C, Floros DJ, Gavilan RG, Kleigrewe K, Northen T, Dutton

RJ, Parrot D, Carlson EE, Aigle B, Michelsen CF, Jelsbak L, Sohlenkamp C, Pevzner P,

Edlund A, McLean J, Piel J, Murphy BT, Gerwick L, Liaw C-C, Yang Y-L, Humpf H-U,

Maansson M, Keyzers RA, Sims AC, Johnson AR, Sidebottom AM, Sedio BE, Klitgaard A,

Larson CB, P CAB, Torres-Mendoza D, Gonzalez DJ, Silva DB, Marques LM, Demarque

DP, Pociute E, O'Neill EC, Briand E, Helfrich EJN, Granatosky EA, Glukhov E, Ryffel F,

Houson H, Mohimani H, Kharbush JJ, Zeng Y, Vorholt JA, Kurita KL, Charusanti P, McPhail

KL, Nielsen KF, Vuong L, Elfeki M, Traxler MF, Engene N, Koyama N, Vining OB, Baric R,

Silva RR, Mascuch SJ, Tomasi S, Jenkins S, Macherla V, Hoffman T, Agarwal V, Williams

PG, Dai J, Neupane R, Gurr J, Rodríguez AMC, Lamsa A, Zhang C, Dorrestein K, Duggan

BM, Almaliti J, Allard P-M, Phapale P, Nothias L-F, Alexandrov T, Litaudon M, Wolfender J-

L, Kyle JE, Metz TO, Peryea T, Nguyen D-T, VanLeer D, Shinn P, Jadhav A, Müller R,

Waters KM, Shi W, Liu X, Zhang L, Knight R, Jensen PR, Palsson BO, Pogliano K,

Linington RG, Gutiérrez M, Lopes NP, Gerwick WH, Moore BS, Dorrestein PC, Bandeira N.

2016. Sharing and community curation of mass spectrometry data with Global Natural

Products Social Molecular Networking. Nat Biotechnol 34:828–837.

20. Longo N, Frigeni M, Pasquali M. 2016. Carnitine transport and fatty acid oxidation. Biochim

Biophys Acta 1863:2422–2435.

21. Munger J, Bennett BD, Parikh A, Feng X-J, McArdle J, Rabitz HA, Shenk T, Rabinowitz JD.

2008. Systems-level metabolic flux profiling identifies fatty acid synthesis as a target for

antiviral therapy. Nat Biotechnol 26:1179–1186.

22. Karlsson EA, Hertz T, Johnson C, Mehle A, Krammer F, Schultz-Cherry S. 2016. Obesity

Outweighs Protection Conferred by Adjuvanted Influenza Vaccination. MBio 7.

23. Carr EL, Kelman A, Wu GS, Gopaul R, Senkevitch E, Aghvanyan A, Turay AM, Frauwirth

KA. 2010. Glutamine uptake and metabolism are coordinately regulated by ERK/MAPK

during T lymphocyte activation. J Immunol 185:1037–1044.

24. Smallwood HS, Duan S, Morfouace M, Rezinciuc S, Shulkin BL, Shelat A, Zink EE, Milasta

S, Bajracharya R, Oluwaseum AJ, Roussel MF, Green DR, Pasa-Tolic L, Thomas PG.

2017. Targeting Metabolic Reprogramming by Influenza Infection for Therapeutic

Intervention. Cell Rep 19:1640–1653.

25. Sikalidis AK. 2015. Amino acids and immune response: a role for cysteine, glutamine,

phenylalanine, tryptophan and arginine in T-cell function and cancer? Pathol Oncol Res

21:9–17.

26. Lee Y-C, Su Y-T, Liu T-Y, Tsai C-M, Chang C-H, Yu H-R. 2018. L-Arginine and L-Citrulline

Supplementation Have Different Programming Effect on Regulatory T-Cells Function of

Infantile Rats. Front Immunol 9:2911.

27. Wijnands KAP, Castermans TMR, Hommen MPJ, Meesters DM, Poeze M. 2015. Arginine

and citrulline and the immune response in sepsis. Nutrients 7:1426–1463.

28. DeVito SR, Ortiz-Riaño E, Martínez-Sobrido L, Munger J. 2014. Cytomegalovirus-mediated

activation of pyrimidine biosynthesis drives UDP-sugar synthesis to support viral protein

glycosylation. Proc Natl Acad Sci U S A 111:18019–18024.

29. Blasco H, Bessy C, Plantier L, Lefevre A, Piver E, Bernard L, Marlet J, Stefic K, Benz-de

Bretagne I, Cannet P, Lumbu H, Morel T, Boulard P, Andres CR, Vourc'h P, Hérault O,

Guillon A, Emond P. 2020. The specific metabolome profiling of patients infected by SARS-

COV-2 supports the key role of tryptophan-nicotinamide pathway and cytosine metabolism.

Sci Rep 10:16824.

30. Hoffmann H-H, Kunz A, Simon VA, Palese P, Shaw ML. 2011. Broad-spectrum antiviral

that interferes with de novo pyrimidine biosynthesis. Proc Natl Acad Sci U S A 108:5777–

5782.

31. Sarkar I, Zardini Buzatto A, Garg R, Li L, van Drunen Littel-van den Hurk S. 2019.

Metabolomic and Immunological Profiling of Respiratory Syncytial Virus Infection after

Intranasal Immunization with a Subunit Vaccine Candidate. J Proteome Res 18:1145–1161.

32. Shin J-H, Yang J-Y, Jeon B-Y, Yoon YJ, Cho S-N, Kang Y-H, Ryu DH, Hwang G-S. 2011.

(1)H NMR-based metabolomic profiling in mice infected with Mycobacterium tuberculosis. J

Proteome Res 10:2238–2247.

33. Chandler JD, Hu X, Ko E-J, Park S, Lee Y-T, Orr M, Fernandes J, Uppal K, Kang S-M,

Jones DP, Go Y-M. 2016. Metabolic pathways of lung inflammation revealed by high-

resolution metabolomics (HRM) of H1N1 influenza virus infection in mice. Am J Physiol

Regul Integr Comp Physiol 311:R906–R916.

34. Fernández-García M, Rey-Stolle F, Boccard J, Reddy VP, García A, Cumming BM, Steyn
AJC, Rudaz S, Barbas C. 2020. Comprehensive Examination of the Mouse Lung
Metabolome Following Infection Using a Multiplatform Mass Spectrometry Approach. J
Proteome Res 19:2053–2070.

35. Zhou W, Sailani MR, Contrepois K, Zhou Y, Ahadi S, Leopold SR, Zhang MJ, Rao V, Avina
M, Mishra T, Johnson J, Lee-McMullen B, Chen S, Metwally AA, Tran TDB, Nguyen H,
Zhou X, Albright B, Hong B-Y, Petersen L, Bautista E, Hanson B, Chen L, Spakowicz D,
Bahmani A, Salins D, Leopold B, Ashland M, Dagan-Rosenfeld O, Rego S, Limcaoco P,
Colbert E, Allister C, Perelman D, Craig C, Wei E, Chaib H, Hornburg D, Dunn J, Liang L,
Rose SMS-F, Kukurba K, Piening B, Rost H, Tse D, McLaughlin T, Sodergren E, Weinstock
GM, Snyder M. 2019. Longitudinal multi-omics of host-microbe dynamics in prediabetes.
Nature 569:663–671.

36. Hagan T, Cortese M, Rouphael N, Boudreau C, Linde C, Maddur MS, Das J, Wang H,
Guthmiller J, Zheng N-Y, Huang M, Uphadhyay AA, Gardinassi L, Petitdemange C,
McCullough MP, Johnson SJ, Gill K, Cervasi B, Zou J, Bretin A, Hahn M, Gewirtz AT,
Bosinger SE, Wilson PC, Li S, Alter G, Khurana S, Golding H, Pulendran B. 2019.
Antibiotics-Driven Gut Microbiome Perturbation Alters Immunity to Vaccines in Humans.
Cell 178:1313–1328.e13.

37. Mussap M, Fanos V. 2021. Could metabolomics drive the fate of COVID-19 pandemic? A
narrative review on lights and shadows. Clin Chem Lab Med https://doi.org/10.1515/cclm-
2021-0414.

38. Thangavel RR, Bouvier NM. 2014. Animal models for influenza virus pathogenesis,
transmission, and immunology. Journal of Immunological Methods.

39. Heaton NS, Leyva-Grado VH, Tan GS, Eggink D, Hai R, Palese P. 2013. In vivo

bioluminescent imaging of influenza a virus infection and characterization of novel cross-

protective monoclonal antibodies. J Virol 87:8272–8281.

40. Tannous BA. 2009. Gaussia luciferase reporter assay for monitoring biological processes in

culture and in vivo. Nat Protoc 4:582–591.

41. Want EJ, Masson P, Michopoulos F, Wilson ID, Theodoridis G, Plumb RS, Shockcor J,

Loftus N, Holmes E, Nicholson JK. 2013. Global metabolic profiling of animal and human

tissues via UPLC-MS. Nat Protoc 8:17–32.

42. Caporaso JG, Kuczynski J, Stombaugh J, Bittinger K, Bushman FD, Costello EK, Fierer N,

Peña AG, Goodrich JK, Gordon JI, Huttley GA, Kelley ST, Knights D, Koenig JE, Ley RE,

Lozupone CA, McDonald D, Muegge BD, Pirrung M, Reeder J, Sevinsky JR, Turnbaugh

PJ, Walters WA, Widmann J, Yatsunenko T, Zaneveld J, Knight R. 2010. QIIME allows

analysis of high-throughput community sequencing data. Nat Methods 7:335–336.

43. Vázquez-Baeza Y, Pirrung M, Gonzalez A, Knight R. 2013. EMPeror: a tool for visualizing

high-throughput microbial community data. Gigascience 2:16.

44. Protsyuk I, Melnik AV, Nothias L-F, Rappez L, Phapale P, Aksenov AA, Bouslimani A,

Ryazanov S, Dorrestein PC, Alexandrov T. 2018. 3D molecular cartography using LC-MS

facilitated by Optimus and 'ili software. Nat Protoc 13:134–154.

45. Khan A, Mathelier A. 2017. Intervene: a tool for intersection and visualization of multiple

gene or genomic region sets. BMC Bioinformatics.

46. Sumner LW, Amberg A, Barrett D, Beale MH, Beger R, Daykin CA, Fan TW-M, Fiehn O,

Goodacre R, Griffin JL, Hankemeier T, Hardy N, Harnly J, Higashi R, Kopka J, Lane AN,

Lindon JC, Marriott P, Nicholls AW, Reily MD, Thaden JJ, Viant MR. 2007. Proposed

minimum reporting standards for chemical analysis. Metabolomics.

47. Fahy E, Sud M, Cotter D, Subramaniam S. 2007. LIPID MAPS online tools for lipid

research. Nucleic Acids Res 35:W606–12.

Carnitine

m/z 162.113
RT 0.297

Infected

Uninfected

Glutamine

m/z 147.0765
RT 0.32

Infected

Uninfected

Kynurenine

m/z 209.0921
RT 0.631

Infected

Uninfected

Cytosine

m/z 112.0508
RT 0.31

Infected

Uninfected

Infected

Uninfected

Plasma

Lung tissue

Lung tissue
&
Plasma