Genomic variation across a clinical *Cryptococcus* population linked to disease

outcome

Running title: Variants and virulence in *C. neoformans*

Poppy Sephton-Clark[1], Jennifer L. Tenor[2], Dena L. Toffaletti[2], Nancy Meyers[2], Charles

Giamberardino[2], Síle F. Molloy[3,4], Adrienne Chan[5], Tarsizio Chikaonda[8], Robert

Heyderman[6], Mina Hosseinipour[7], Newton Kalata[8], Cecilia Kanyama[7], Christopher

Kukacha[8], Duncan Lupiya[9], Henry C. Mwandumba[8], Thomas Harrison[3,4], Tihana

Bicanic[3,4], John R. Perfect[2], and Christina A. Cuomo*[1]

[1]Infectious Disease and Microbiome Program, Broad Institute of MIT and Harvard,

Cambridge MA 02142 USA. [2]Division of Infectious Diseases, Department of Medicine,

Duke University School of Medicine, Durham, NC 27710, USA. [3]Centre for Global

Health, Institute of Infection and Immunity, St George's University of London, London,

United Kingdom. [4]Clinical Academic Group in Infection, St George's University Hospital,

London, United Kingdom. [5]Sunnybrook Health Sciences Centre, Toronto, Canada.

[6]Division of Infection and Immunity, UCL, London, United Kingdom. [7]UNC Project

Malawi, University of North Carolina, USA. [8]Malawi-Liverpool-Wellcome Trust Clinical

Research Programme, Blantyre, Malawi. [9]Dignitas International, Zomba, Malawi.

*Address correspondence to: Christina A. Cuomo (cuomo@broadinstitute.org)

Abstract

Cryptococcus neoformans is the causative agent of cryptococcosis, a disease with poor

patient outcomes, accounting for approximately 180,000 deaths each year. Patient

outcomes may be impacted by the underlying genetics of the infecting isolate, however,

our current understanding of how genetic diversity contributes to clinical outcomes is

limited. Here, we leverage clinical, in vitro growth and genomic data for 284 *C.*

neoformans isolates to identify clinically relevant pathogen variants within a population

of clinical isolates from patients with HIV-associated cryptococcosis in Malawi. Through

a genome-wide association study (GWAS) approach, we identify variants associated

with fungal burden and growth rate. We also find both small and large-scale variation,

including aneuploidy, associated with alternate growth phenotypes, which may impact

the course of infection. Genes impacted by these variants are involved in transcriptional

regulation, signal transduction, glycolysis, sugar transport, and glycosylation. When

combined with clinical data, we show that growth within the CNS is reliant upon

glycolysis in an animal model, and likely impacts patient mortality, as CNS burden

modulates patient outcome. Additionally, we find genes with roles in sugar transport are

under selection in the majority of these clinical isolates. Further, we demonstrate that

two hypothetical proteins identified by GWAS impact virulence in animal models. Our

approach illustrates links between genetic variation and clinically relevant phenotypes,

shedding light on survival mechanisms within the CNS and pathways involved in this

persistence.

Importance

Infection outcomes for cryptococcosis, most commonly caused by *C. neoformans*, are

influenced by host immune responses, as well as host and pathogen genetics. Infecting

yeast isolates are genetically diverse, however, we lack a deep understanding of how

this diversity impacts patient outcomes. To better understand both clinical isolate

diversity and how diversity contributes to infection outcome, we utilize a large collection

of clinical *C. neoformans* samples, isolated from patients enrolled in a clinical trial

across 3 hospitals in Malawi. By combining whole-genome sequence data, clinical data,

and in vitro growth data, we utilize genome-wide association approaches to examine the

genetic basis of virulence. Genes with significant associations show virulence

phenotypes in both murine and rabbit models, demonstrating that our approach

successfully identifies links between genetic variation and biologically significant

phenotypes.

Introduction

Cryptococcus neoformans is a pathogenic yeast that most commonly affects

immunocompromised individuals, causing an estimated 180,000 deaths annually, with

75% of these occurring in sub-Saharan Africa. One of the leading causes of death in
adults living with HIV/AIDS, cryptococcal infections are especially problematic in low-
income countries where, despite a widespread roll-out of antiretroviral therapy, deaths
due to opportunistic infections such as cryptococcal meningitis remain high (1). The
infecting propagules of this pathogen generally enter human hosts via inhalation. From
infections within the lung, *C. neoformans* may disseminate throughout the bloodstream
and central nervous system of susceptible patients, causing meningitis (2). In a sample
of healthcare systems across low-income countries, the 1-year mortality rate for
individuals who develop cryptococcal meningitis is estimated to be 70% for those in
care (uncertainty interval 59–81%) (1). A better understanding of *C. neoformans* strain
virulence and fitness within the host is necessary to improve patient outcomes and
develop new treatment options.

Whilst the majority of cryptococcosis cases are caused by *Cryptococcus neoformans*
var grubii (3), there are often high levels of genetic diversity within clinical populations of
C. neoformans (4–7). Furthermore, isolates of the same multilocus sequence type
(MLST) have been shown to cause infections that range in severity from mild to extreme
(8). To examine how genetic variation contributes to virulence phenotypes, a recent
study carried out logistic regression analysis with 38 clinical *C. neoformans* isolates of
the same sequence type to identify single nucleotide polymorphisms (SNPs) associated
with patient survival, clinical parameters including cytokine response, immune cell
counts and infection clearance, as well as in vitro data on absolute yeast growth and
macrophage interactions (9). This study identified 40 candidate genes based on these
association parameters, 6 of which (out of 17 genes tested) were important for survival
in a murine model of *C. neoformans* infection. In a larger cohort of 230 *C. neoformans*
samples from patients in South Africa, isolate sequence type was associated with
patient outcome, in vitro cerebrospinal fluid (CSF) survival, and phagocytosis response
(10). Full scale genome-wide association studies (GWAS) have also examined how
natural variation within a *C. neoformans* population differentiates clinical and
environmental isolates, identifying loss-of-function variants present in clinical *C.*

neoformans (VNB) populations that impact a transcription factor important for
melanization, a well-studied virulence factor (11).

Furthermore, copy number variation, such as aneuploidy, has also been frequently
identified within clinical populations of *C. neoformans*. Disomy of chromosome 1 is
commonly reported for isolates exposed to azoles, and the higher copy number of two
key genes, the *AFR1* transporter and the *ERG11* drug target, confer increased
resistance to antifungals such as fluconazole (12–14). Chromosome duplication as a
result of in vivo passage has also been noted in clinical isolates (15–17), and the
emergence of aneuploidy in this setting has been proposed as a mechanism by which
both *Cryptococcus* and *Candida* species might rapidly adapt to high-stress
environments (18, 19). In *C. neoformans* aneuploidy is often transient and passage
under non-selective conditions allows for reversion to euploidy (14, 17). In total,
aneuploidy of chromosomes 1, 2, 4, 6, 8, 9, 10, 12, 13, and 14 have been reported in *C.*
neoformans (14, 16, 17, 20–23). Despite appearing consistently throughout clinical
populations, the impact of these other chromosomal aneuploidies is not yet well
understood.

To better understand how genetic variation among *C. neoformans* isolates contributes
to infection outcomes in patients, we carried out genome-wide association studies
(GWAS) with 266 *C. neoformans* clinical isolates from the VNI lineage, selected to
reduce the confounding effects of population structure between lineages. In addition to
comparing selected clinical data, all isolates were also measured for in vitro growth
under diverse conditions. Through our GWAS approach, we identify two hypothetical
proteins associated with fungal burden in patients which also contribute to virulence in a
murine model. Additionally, we show that growth in a rabbit model of CNS infection is
dependent on glycolytic genes identified by GWAS, and corroborate findings that patient
outcome is highly correlated with fungal burden in the CNS. Partial and full
chromosomal duplications are commonly detected within this clinical population, yet
these aneuploidies reduce *C. neoformans* fitness under in vitro growth conditions.

Materials and Methods

Sample Preparation and Sequencing

Clinical cryptococcal isolates derived from patient CSF subculture were procured through the Antifungal Combinations for Treatment of Cryptococcal Meningitis in Africa Trial (ACTA) (24); repeat cultures and duplicates were excluded. Collected strains were grown overnight in 10 ml of YPD at 30°C and 225 rpm. Genomic DNA was then extracted for sequencing with the MasterPure Yeast DNA Purification Kit, as described by Desjardins et al. (11). DNA was sheared to 250bp using a Covaris LE instrument, and adapted for Illumina sequencing as described by Fisher et al. (25). Libraries were sequenced on a HiSeq X10, generating 150bp paired reads (minimum average coverage 100x).

Data Processing and Variant Calling

To determine sample species, reads were first aligned to a composite pan-*Cryptococcus* genome, consisting of reference genomes for *Cryptococcus neoformans var. grubii* H99, *Cryptococcus neoformans var. neoformans* JEC21, and representative genomes for lineages VGI, VGII, VGIIIa, VGIIIb, VGIV, and VGV of *Cryptococcus gattii* (26–29). To identify variants for *C. neoformans* species, reads were aligned to the *Cryptococcus neoformans var grubii* H99 reference genome (GCA_000149245.3) (27) with BWA-MEM version 0.7.17 (30). GATK v4 variant calling was carried out as documented in our publicly available cloud-based pipeline (31) (https://github.com/broadinstitute/fungal-wdl/tree/master/gatk4). Post calling, variants were filtered on the following parameters: QD < 2.0, QUAL < 30.0, SOR > 3.0, FS > 60.0 (indels > 200), MQ < 40.0, GQ < 50, alternate allele percentage = 0.8, DP < 10. Variants were annotated with SNPeff, version 4.3t (32).

Population Genomic Analysis

A maximum likelihood phylogeny was estimated using 72,258 segregating SNP sites present in one or more isolates, allowing ambiguity in a maximum of 10% of samples, with RAxML version 8.2.12 GTRCAT rapid bootstrapping (33), and visualized with ggtree (R 3.6.0) rooted to VNII isolates. To estimate linkage disequilibrium (LD) decay,

vcftools version 0.1.16 was used to calculate LD for 1000bp windows, with a minimum

minor allele frequency of 0.1, and the --hap-r2 option. Region deletions and duplications

were identified using CNVnator v0.3 (significant instance e value < 0.01) (34). To

identify regions under selection, composite likelihood ratio analysis was performed with

PopGenome (R 3.5.0, PopGenome 2.7.5) per chromosome, by 5kb windows (35). The

top 5% scoring regions (centromeric regions excluded) were tested for enrichment using

a hypergeometric test with FDR correction. Large duplications and aneuploidies were

visualized   using   funpipe   (coverage   analysis)   version   0.1.0

(https://github.com/broadinstitute/funpipe).

Genome-Wide Association Studies

Association analysis between clinical data, in vitro phenotypes, and variants was carried

out using PLINK v1.08p formatted files and Gemma version 0.94.1 (36) (options:

centered relatedness matrix gk 1, linear mixed model), as previously described (11).

Variants were considered in two scenarios, one in which rare variants (present in < 5%

of the population) were collapsed by gene, and another in which loss-of-function

variants (SNPeff impact high) were considered independently. Significant variants were

considered to have a test score < 1.00E-6.

Clinical Data Analysis

De-identified clinical metadata detailing CSF fungal burden (CFU/ml), fungal clearance

(EFA), patient mortality, and Glasgow coma score were provided by investigators, with

these parameters determined as previously described (24). Correlation between clinical

parameters was determined in R 3.6.0 with Pearson correlation coefficient, Spearman's

rank correlation coefficient, point-biserial correlation, or Phi coefficient. Survival curves

were generated using Prism v9.1.0 and statistics were carried out in R 3.6.0.

In vitro Phenotyping

Strains were grown at 30°C for two to five days. For each strain, a single colony was

selected and added to 96 well microtiter plates containing 200 µL of YPD broth. Each 96

well plate contained six control strains (H99, *LAC1*, *MPK1*, *CNA1*, *RAS1*, and *HOG1*)

and a YPD control. The 96 well plates were incubated for one to two days at 30°C.

Colonies were pin replicated into 384 well microtiter plates containing 80 µL of YPD

broth in each well. The 384 well plates were incubated for one to two days at 30°C.

They were then pinned onto one well solid agar plates in duplicate using the BM5-BC

Colony Processing Robot (S & P Robotics) in 1536 array format. In three separate

biological replicates, isolates were grown at 30°C, 37°C, and 39°C on YPD agar.

Isolates were also pinned onto YPD+10 µg/ml fluconazole and YPD+64 µg/ml

fluconazole. Images were captured after approximately 24, 48, and 72 h. Colony size at

48h was determined using gitter (37), and used to assess growth.

Gene Deletion Strains

Strains used for the animal studies and the primer sequences used are listed in

Supplemental Table 1. KN99alpha (CM026) was used as the reference wild-type strain

for deletions obtained from a genome-wide *Cryptococcus* deletion library (38). Four

deletion strains were generated in wild-type *C. neoformans* strain H99 (*cnag_00544Δ,*

cnag_04102Δ, cnag_05324Δ, cnag_06033Δ) in this study. Three DNA fragments were

amplified by PCR: approximately 0.7-1 kb of 5' flank sequence, the nourseothricin

(NAT) drug selection cassette amplified from pAI3 (39), and 0.7-1 kb of 3' flank

sequence were prepared for each gene. The PCR products were gel extracted using

the NucleoSpin Gel and PCR Clean-up Kit (Macherey-Nagel). Next, the PCR products

were cloned into pUC19 using NEBuilder HiFi DNA Assembly, transformed into

Escherichia coli, and positive plasmids were confirmed by PCR. For biolistic

transformation, 2 µg of plasmid was transformed into *C. neoformans* strain H99 as

previously described (40) with a slight modification that the cells were recovered on

YPD containing 0.5 M sorbitol and 0.5 M mannitol. The cells were allowed to recover for

2.5 h before transferring to the selective medium, YPD+100 µg/ml NAT. Positive

transformants were confirmed by PCR.

Capsule production

To evaluate the capsule size, capsule inducing medium (10% Sabouraud broth in 50

mM MOPS pH 7.3) was used as previously described (41). Five milliliters of capsule

inducing medium was inoculated with a single freshly streaked yeast colony and grown

in an incubator shaker (225 rpm) for approximately 24 and 48 h. India ink was used as a

counterstain at a 1:5 ratio (ink:cell suspension). Images were captured by microscopy

(Zeis Axio Imager 1). Cell body and capsule size were measured in ImageJ V1.53a.

Murine model of infection

C. neoformans strains were grown in YPD broth at 30°C in a shaking incubator (220

rpm) for 24 h, centrifuged, and washed twice in phosphate-buffered saline (PBS). Cell

counts were determined using a T4 cell counter (Nexcelom). Five male CD-1 mice

(Charles River Laboratories) were infected with approximately 5×10^4 yeast cells per

mouse via oropharyngeal aspiration while under isoflurane anesthesia. Mice were

monitored and weighed daily. Mice with a total body weight loss of ≥ 20% or that

exhibited behavioral, neurological, or respiratory symptoms were sacrificed following

IACUC guidelines. Kaplan-Meier survival plots and analysis (the log-rank test) were

completed using Prism software v9.1.0.; GraphPad Software. A p-value of ≤ 0.05 was

considered statistically significant.

Rabbit model of infection

To assess the fitness and virulence of deletion strains in rabbit CSF, 3 New Zealand

White male rabbits were inoculated intracisternally with 300 µl of approximately $1x10^8$

cells. Animals were sedated with ketamine and xylazine for inoculation and serial CSF

cisternal taps. The rabbits were treated with hydrocortisone acetate (2.5 mg/kg) by

intramuscular injections daily starting one day prior to yeast inoculation. Cisternal taps

were performed on days 3, 7, and 10 followed by serial dilution of the CSF and

enumeration of colonies. The time series fungal burden data were then assessed by

using repeated measures analysis in R v3.6.1.

Animal studies

Animal experiments were performed in compliance with the Animal Welfare Act, the

*Guide for the Care and Use of Laboratory Animal*s (42), and the Duke Institutional

Animal Care and Use Committee (IACUC).

Results

The VNI lineage dominates clinical isolates and shows selection for sugar transporters

To examine variation within clinical populations, *C. neoformans* samples were isolated from HIV-infected patients as part of the ACTA trial (24), which evaluated the efficacy of fluconazole partnered with flucytosine, compared to amphotericin B combined with either fluconazole or flucytosine, as induction therapy for cryptococcal meningitis. Baseline (pre-antifungal exposure) isolates were collected from three hospitals in Malawi between 2013 and 2016. We performed whole-genome sequencing on 344 isolates and called variants against the *C. neoformans var grubii* H99 reference genome (27). Isolates identified as *Cryptococcus gattii* (45), hybrid AD *C. neoformans* (4), diploid (2), or with low coverage (9), based on evaluating read alignments to a pan-*Cryptococcus* collection of reference genomes, were removed from the analysis. To examine the population structure, a maximum likelihood phylogeny was built using 72,258 segregating SNP sites (Figure 1). Isolates can be clearly identified as VNI (266), VNII (9), and VNB (9) based on phylogenetic comparison with previously typed isolates reported by Desjardins et al (11). VNI isolates split into VNIa (217) and VNIb (49) with 100% bootstrap support; these recently described sub-lineages show strong evidence of separation (11). Of the two mating type loci found in *Cryptococcus*, mating type α predominated among these isolates, with only one VNB isolate (ACTA3525-D) possessing the alternate *MAT***a**. To assess recombination within the large VNI population, we calculated linkage disequilibrium (LD) decay and found levels of decay for lineage VNI (LD50 30kb) (Supplemental Figure 1) similar to those reported by Desjardins et al. (LD50 for VNI, VNBI, and VNBII being < 50kb) (11). There is increased decay in the VNI population as a whole when compared to individual VNIa and VNIb sub-groups (LD50 of 110kb for VNIb, LD50 not reached within 250kb for VNIa), suggesting that VNIa and VNIb isolates do not recombine exclusively within their groups.

To identify genomic regions under positive selection, we performed composite likelihood ratio analysis (43). We found that regions with scores in the top 10% in more than one

lineage (examining VNIa, VNIb, VNII, and VNB) include subtelomeric regions,
centromeres, *ERG11*, and *AFR1* (Supplemental Figure 2). To examine if genes within
these regions are associated with shared functions that appear under selection, we
performed gene ontology (GO) enrichment analysis (Hypergeometric test, FDR
correction) on regions with selection scores in the top 5% (excluding centromeres). For
VNII isolates we found nucleotide excision repair enriched in these regions (corrected p-
value = 6.77E-3). Sugar transport, including inositol transport, appeared robustly
enriched in both VNIa and VNB lineages (corrected p-value's = 1.07E-3 and 8.60E-3),
supporting previous work that identified these functions as under selection
(Supplemental Table 2) (11). Sugar transport and utilization have been identified as key
to success in nutrient scarce environments such as the CNS (23, 44, 45), during
interactions with amoebae, and are required for virulence and resistance to external
stress (46–49).

We also identified regions that were duplicated or deleted in these *C. neoformans*
isolates via copy number variation analysis (Supplemental Table 3). An 8kb region was
found to be duplicated in 43 VNIa isolates, containing 4 genes including a sugar
transporter (TCDB: 2.A.1.1, glycerol transport), a predicted non-coding RNA, a fungal
specific transcription factor (Zn2Cys6, *SIP402*), and a short-chain dehydrogenase. A
separate 34kb region was found duplicated across 48 VNIa isolates encoding 11 genes
including 3 dehydrogenase enzymes and 2 hydrolase enzymes. Duplicated regions
unique to VNIb included an un-annotated 19kb region specific to 20 isolates and an 8kb
region that encodes 2 hypothetical proteins duplicated across 21 VNIb isolates.
Duplications of genes involved in resistance to azoles such as *ERG11* were found
exclusively in VNII isolates, however, this did not correlate with an enhanced ability to
grow in the presence of fluconazole at 64ug/ml. While these duplicated regions are not
directly associated with tested phenotypes, the duplication of regions containing genes
such as *ERG11* and sugar transporters may still contribute to phenotypic variation that
is relevant to clinical outcomes through modulation of growth and virulence phenotypes
when grown in alternate conditions.

GWAS identifies multiple variants associated with fungal burden

We next used clinical data associated with these isolates to investigate the relationship between clinical factors, across lineages. We confirmed previous findings that mortality correlates strongly with high baseline fungal burden (CSF CFU/ml taken at diagnosis of cryptococcal meningitis) (Figure 2a, p=7.70E-7) as shown in prior studies (50–52), and observed similar outcome ratios across lineages (log-rank test, p=0.916) (Figure 2b-c), suggesting no major lineage-specific differences in virulence, though low numbers of VNB and VNII infecting isolates may limit our power to detect significant differences here. Additionally, we noted similar levels of baseline fungal burden and rates of clearance between VNIa and VNIb infecting isolates (Figure 2d-e). The data suggest a reduction in baseline fungal burden of 1.29E+06 CSF CFU/ml on average for VNII when compared to VNI isolates (Wilcoxon test, p=0.024), however, due to the limited number of VNII isolates included, this finding should be confirmed with additional cases.

To determine whether the variation in baseline fungal burden, which appears well distributed throughout this population (Figure 1), is linked to a specific genetic component, we performed genome-wide association studies (GWAS) to identify variants associated with higher fungal load, when taken as a continuous phenotype. We selected VNI isolates for this analysis as they represent the major genetic group present, and to avoid confounding factors of population structure between lineages. This analysis revealed 53 variants that were significantly (GEMMA score test, p < 1.00E-6) associated with CSF fungal burden levels (Figure 3a), 16 of which were predicted to result in a loss-of-function mutation. These variants impacted genes encoding 15 hypothetical proteins and 6 ncRNAs; an additional 4 variants fell into non-coding centromeric regions (Supplemental Table 4). Of the annotated genes impacted, 4 have been previously identified to modulate virulence phenotypes, these include the protein S-acyl transferase *PFA4* (CNAG_03981), the calcineurin catalytic subunit *CNA1* (CNAG_04796), and the mitochondrial co-chaperone *MRJ1* (CNAG_00938) that are required for virulence in the murine model, as well as the iron permease *FTR1* (CNAG_06242) that is required for capsule regulation (53–56). Additionally, 2 genes with variants are known to impact titan cell formation, these include the multidrug

transporter CNAG_04546 and the adenylate cyclase *CAC1* (CNAG_03202) (57, 58). A

high proportion (28%) of variants with high GWAS scores (GEMMA score test, p <

1.00E-6) appeared in genes annotated as hypothetical proteins. For phenotype

characterization, we decided to focus on genes impacted by variants associated with

higher fungal burden, that were predicted to result in a loss-of-function within coding

regions.

345   Two hypothetical proteins impact virulence in a murine model

To determine if the genes identified through our GWAS analysis of fungal burden impact

virulence, we tested a total of 10 gene deletion strains across murine and rabbit models.

Previous work has shown that infection outcomes from human infections are well

recapitulated in murine models (8), while rabbit models have proven useful in evaluating

CNS infections, as fungal burden within the CNS can be determined through

longitudinal tapping of CSF (59). The most striking result from GWAS analysis was a

pileup of 5 variants in the hypothetical protein CNAG_04102, with the highest scoring

variant within this gene being the third most significant overall (GEMMA score test,

p=1.30E-11). Deletion of CNAG_04102 in a H99 background led to reduced virulence in

a murine model (5 male CD-1 mice), compared to a H99 isolate control (log-rank test,

p=0.0427) (Figure 3b). This gene contains a Kyakuja-Dileera-Zisupton (KDZ)

superfamily motif (pfam18758), which has been found within species from

basidiomycota, mucoromycotina, *rhizophagus* and *allomyces* (60), with CNAG_04102

homologs containing this motif found in *Cryptococcus gattii* and *Cryptococcus floricola*.

The KDZ motif is also commonly located within TET/JBP genes which are involved in

genomic organization and epigenetic regulation (61), suggesting a role for gene

expression regulation. Deletion of a second hypothetical protein, CNAG_05608, (in a

CMO26 KN99 background) also resulted in reduced virulence within a murine model (5

male CD-1 mice), with isolates lacking a functional CNAG_05608 exhibiting reduced

virulence, when compared to wild type (log-rank test, p=0.0154) (Figure 3c). While

CNAG_05608 is annotated as a hypothetical protein, this gene is predicted to contain a

single transmembrane domain and has homologs in *Cryptococcus gattii*, *Cryptococcus*

amylolentus, *Kwoniella* species, and *Wallemia* species. Furthermore, this gene is

upregulated during growth in both murine and monkey lungs (62), and slightly
downregulated when grown in the presence of glucose (63), suggesting a role during
infection.

Sugar transport and metabolism impacts persistence in a rabbit CNS infection
model

When loss-of-function variants were considered independently, the most highly
significant variant was a frameshift in the phosphofructokinase gene, CNAG_06033
(pfkB) (GEMMA score test, p=1.69E-09). Deletion of CNAG_06033 in a H99
background resulted in significantly decreased CSF burden within the rabbit model (3
New Zealand white male rabbits) when compared to a H99 isolate, with CSF CFU/ml
counts dependent on both the infecting strain and the number of days post-infection.
The CSF loads were comparable across three rabbits for H99 and the CNAG_06033
deletion at 3 days post-infection, but decreased for the CNAG_06033 deletion at days 7
and 10, in contrast to the H99 isolate which showed increased CSF load over time
(repeated measures analysis, p=0.0225) (Figure 3d), highlighting the need for efficient
glycolysis within the mammalian CSF (44). The overall virulence within a murine model
(5 male CD-1 mice) for the CNAG_06033 deletion strain appeared similar to wild type
(log-rank test, p=0.1198) (Figure 3e), demonstrating that defects in glycolytic
metabolism do not appear to restrict persistence and dissemination but critically may be
important at specific body sites. Loss-of-function variants within this gene have been
shown to emerge over the course of in vivo passage in CSF during human infection and
relapse (15), suggesting a role for the loss of CNAG_06033 in adaptation to the host at
specific body sites. This result is consistent with prior work that tested deletions of other
genes involved in glycolysis; loss of pyruvate kinase (*pyk1Δ*) resulted in decreased
persistence in the rabbit CSF, but unperturbed dissemination in a murine model (44).
Further supporting the role of glycolysis and sugar transport in the survival of *C.*
neoformans in CSF, we identified significant variants in additional genes involved in
these pathways. A predicted xylose transporter, CNAG_05324, contained a frameshift
variant present in 33 isolates (GEMMA score test, p=4.00E-07). In preliminary
experiments deletion of CNAG_05324 in a H99 background led to an increase in CSF

load in one rabbit when compared to its H99 control. However, additional experiments
are required to confirm these results (Supplemental Figure 3a). Given the predicted role
of this gene as a xylose transporter, and the presence of xylose in cryptococcal capsule,
we undertook preliminary phenotypic capsule screening of the CNAG_05324 deletion
strain. Capsule analysis of this deletion mutant via cultivation in capsule-inducing media
and India ink staining revealed a significant increase in capsule thickness (Welch's T-
test, p=3.9E-14) (Supplemental Figure 3b), suggesting a role for CNAG_05324 in
capsule size modulation. Previous work has shown that modulation of xylose transport
and xylosylation can drastically alter virulence, capsule size, and immune evasion (64,
65), highlighting this capsular mechanism as an area for further exploration.

Aneuploidy is common and slows growth
To determine how natural variation might affect growth and other clinically relevant
phenotypes, we performed in vitro phenotyping of isolates. Isolates displayed a range of
growth levels on rich media (YPD) at 30°C, 37°C, and 39°C (Figure 4a-c), with colony
size across conditions and replicates showing strong correlation (Figure 4d,e, replicate
per condition R^2 > 0.8). To determine whether this variation is linked to a specific
genetic component, we performed GWAS to identify variants associated with increased
and decreased colony size. Significantly associated with the rapid growth of large
colonies on YPD were loss-of-function variants in CNAG_06637 (UBP8 Ubiquitin-
specific protease, component of the SAGA complex), CNAG_03818 (sensory
transduction histidine kinase), and CNAG_10082 (tRNA Threonine) (GEMMA score
test, p < 1.00E-6). A single loss-of-function variant was found significantly associated
with decreased colony size, a frameshift in the dolichyldiphosphatase encoding gene,
CNAG_03014 (GEMMA score test, p=9.90E-12). Naturally occurring loss-of-function
variants such as these may play a role in the fitness variation we see between clinical
isolates.

In addition to SNP and indel mutations, we evaluated the level of chromosome copy
number variation across these clinical isolates. We observed both fully and partially
duplicated chromosomes, with the most commonly duplicated chromosomes being 12,

9, 14, and 1; overall, duplication of an entire chromosome occurs in 8.5% of clinical
isolates. Partial duplications, where at least 25% of the chromosome shows continuous
duplication, occur most frequently in chromosomes 2 and 6 (15 instances, Figure 5a).
Aneuploid isolates appear well distributed throughout this clinical dataset (Supplemental
Figure 4a), suggesting frequent and independent origins for these events occurring in
vivo.

To evaluate the impact of these large aneuploidies, we next compared the growth of
aneuploid and euploid isolates. On rich media (YPD), at 30°C, 37°C, and 39°C, isolates
harboring a fully duplicated chromosome showed significantly poorer growth than
euploid isolates (Wilcoxon test, $p < 5.00E-07$), or isolates featuring only a partial
chromosomal duplication (Wilcoxon test, $p < 0.01$) (Figure 5b-d). To determine whether
this fitness reduction occurs in both clinical and environmental populations, we carried
out a metanalysis of the data for these isolates, and data previously generated using the
same assay for a diverse set of clinical and environmental isolates (Supplemental
Figure 4b) (11). Isolates with aneuploidies present in both datasets displayed a
significant reduction in growth on YPD at 37°C ($p=2.00E-07$), demonstrating that this
reduction in fitness holds true for both clinical and environmental isolates, across
lineages VNI and VNB.

Aneuploidy can be advantageous under certain stressors such as antifungal treatment,
however, with optimal nutrients at a range of temperatures we found that aneuploidy
significantly reduces fitness. Modulation of chromosome 1 ploidy has been linked to
apoptosis-inducing factor 1 (*AIF1*) (13), however, variants within *AIF1* were not present
in this population, suggesting an alternate mechanism may be responsible for
modulation of chromosome copy number here.

Discussion
Phylogenetic analysis of 284 *C. neoformans* samples from patients with cryptococcosis
revealed a mixed population of VNI, VNB, and VNII isolates, dominated by the VNI
lineage. We leverage clinical metadata to identify multiple variants associated with
fungal burden, a factor known to impact patient mortality, in combination with additional

factors such as host immune response, host genetic background, and antifungal
treatment. We identify both SNPs and indels significantly associated with fungal burden,
but not with other clinical factors such as mortality and rate of yeast clearance. We
utilize these association results to identify three genes involved in virulence. Of these,
two are annotated as hypothetical proteins and when deleted show reduced virulence in
a murine model, and one encodes for a phosphofructokinase that when deleted shows
reduced persistence in a rabbit CNS model. We employ in vitro phenotyping to identify
multiple variants associated with both poor and prolific growth phenotypes, and through
copy number analysis, we reveal multiple and varied aneuploidies within this clinical
population, as well as a previously reported environmental population. These
aneuploidies appear to reduce fitness levels in isolates from both clinical and
environmental sources, across both VNI and VNB lineages.

Isolates collected from patients in Malawi as part of the ACTA study were dominated by
the VNI genetic group, the most commonly observed global lineage of *C. neoformans*.
This population consists of two of the three previously identified VNI lineages (VNIa,
and VNIb) (66, 67). VNI isolates are found around the globe and appear relatively clonal
when compared to the highly diverse VNB lineage, often isolated from rural niches,
such as mopane trees, in Africa and South America (11, 68–70).

Sugar transporters have previously been identified as under selection in both VNI and
VNB isolates from Botswana (11), and we find they also appear under selection in both
VNI and VNB isolates in this population from Malawi. Specifically, we find inositol,
xylose, glucose, lactose, and glucoside transporters under selection. The expansion of
inositol transporters in *C. neoformans* may offer an advantage in both woodland areas
and the CNS, as these environments have abundant inositol (46). Xylose transport is
important for *C. neoformans* capsule production, the variable xylosylation of which can
enable immune evasion (65). The signaling molecule and preferred carbon source
glucose, the precursor of glucoside, is known to regulate hexose transporters that are
required for virulence (47) and is a key glycolytic metabolite, a pathway required for
growth in the CNS (44).

We identified multiple variants significantly associated with clinical and growth
phenotypes by taking a GWAS approach, however, clinical phenotypes such as
mortality and mental-status did not show a strong association with any variants
identified, perhaps due to the complex nature of these characteristics. Additionally, we
used a culture-based method to select for isolates from patients, and culture-negative
individuals were excluded. As a result, we were unable to detect variants associated
with very low levels of fungal burden in the CNS. While extensively applied to human
data, genome-wide association studies have also been applied to study fungal human
pathogens (9–11) and plant-pathogens (71, 72). A major challenge is adapting these
association approaches for the population structure of each species; for *C. neoformans*,
while there is recombination within the population, LD50 values for VNI, VNBI, and
VNBII populations are < 50kb (11). Expanding the sample size for such associations or
focusing narrowly on particular genetic groups can help increase the power to detect
variants, however, the choice of GWAS approach also needs to be optimized for the
population under study through consideration of population structure and size.

Through analysis of the clinical metadata, we found CSF fungal burden, a measure of
the quantity of live yeasts at the site of infection, to be the strongest factor in carrying
out GWAS. We found that high fungal burden within the CSF of an individual strongly
correlated with patient mortality, in accordance with prior work showing that high fungal
load is a predictor of mortality (50–52). We found that isolates lacking a functional
phosphofructokinase B (CNAG_06033) exhibited a reduced CSF load within a rabbit
model; phosphofructokinase plays a key regulatory role in the glycolytic pathway.
Patient isolates containing naturally occurring loss-of-function variants in CNAG_06033
showed no growth defects when grown on YPD at 37°C, and our phosphofructokinase
mutant showed a CSF load reduction similar to that observed for the pyruvate kinase
(*PYK1*) deletion strain within a rabbit model, likely due to the similar regulatory effects of
both enzymes in glycolysis (44). Additionally, loss-of-function variants have previously
been identified in CNAG_06033 (PfkB) after in vivo human passage (15). Under stress
conditions, metabolically heterogeneous populations may emerge (73, 74), this

metabolic diversity might explain the emergence of isolates less reliant upon glycolysis
through growth within the host. Additionally, we identified two hypothetical proteins that
appear to play a role in virulence within the murine model; this highlights the power of
our unbiased GWAS approach to systematically identify gene candidates implicated in
virulence, in the absence of additional functional or pathway information. Other
systematic studies utilizing RNA-Seq have also identified genes encoding hypothetical
proteins that are strong candidates for further study, due to their high expression in CSF
(23, 45). While the large proportion of *C. neoformans* genes annotated as hypothetical
proteins are more challenging to study, it is critical that we more widely characterize the
roles of all genes involved in the pathogenesis of *C. neoformans*.

We also found evidence that ploidy directly impacts the fitness of both clinical and
environmental *C. neoformans* isolates. Aneuploidy has been linked to broad-spectrum
stress resistance in *Candida* species (75), and in *C. neoformans*, disomy of
chromosome 1 is known to arise in isolates treated with azoles both in vitro and in vivo
and confers resistance to azoles such as fluconazole, through the increase in copy
number of *AFR1* and *ERG11* (12, 14). Suggested mechanisms for modulation of
chromosome 1 ploidy include regulation via the apoptosis-inducing factor Aif1 (13),
however, we did not find evidence for Aif1 disruption in these isolates. Specific impacts
of disomy have also been noted for chromosome 13, disomy of which results in reduced
melanization (20). In *S. cerevisiae*, disomy of select chromosomes also results in
reduced proliferation rates (76). Whilst the reduction in fitness we observed does not
seem specific to any particular chromosome, the questions of how and why ploidy
appears subject to change in stressful conditions and whether the most frequently
observed aneuploidies confer a specific advantage are intriguing and will require further
study.

By combining genetic, in vitro, and clinical data, we glean insights into the impact of
naturally-occurring genetic variation and the implications for infection outcomes. As
whole genome sequencing on an ever-larger scale becomes more accessible, and
association techniques for microbial communities grow in sophistication, so too will our

power to detect functionally relevant genetic variation across cryptococcal populations. Combined with data from large pan-African clinical trials, and approaches that leverage both fungal and human variant identification, we can further dissect the interaction between pathogen and host genetics. Together, this will enable a better understanding of how these variations impact the ability of *Cryptococcus* to adapt to, and thrive in, the wide range of environments it finds itself within.

References

1. Rajasingham R, Smith RM, Park BJ, Jarvis JN, Govender NP, Chiller TM, Denning DW, Loyse A, Boulware DR. 2017. Global burden of disease of HIV-associated cryptococcal meningitis: an updated analysis. Lancet Infect Dis 17:873–881.

2. Maziarz EK, Perfect JR. 2016. Cryptococcosis. Infect Dis Clin North Am 30:179–206.

3. Chayakulkeeree M, Perfect JR. 2006. Cryptococcosis. Infect Dis Clin North Am 20:507–544, v–vi.

4. Litvintseva AP, Thakur R, Vilgalys R, Mitchell TG. 2006. Multilocus sequence typing reveals three genetic subpopulations of *Cryptococcus neoformans* var. *grubii* (serotype A), including a unique population in Botswana. Genetics 172:2223–2238.

5. Chen Y, Litvintseva AP, Frazzitta AE, Haverkamp MR, Wang L, Fang C, Muthoga C, Mitchell TG, Perfect JR. 2015. Comparative analyses of clinical and environmental populations of *Cryptococcus neoformans* in Botswana. Mol Ecol 24:3559–3571.

6. Nyazika TK, Hagen F, Machiridza T, Kutepa M, Masanganise F, Hendrickx M, Boekhout T, Magombei-Majinjiwa T, Siziba N, Chin'ombe N, Mateveke K, Meis JF, Robertson VJ. 2016. *Cryptococcus neoformans* population diversity and clinical outcomes of HIV-associated cryptococcal meningitis patients in Zimbabwe. J Med Microbiol 65:1281–1288.

7.   Fernandes KE, Brockway A, Haverkamp M, Cuomo CA, van Ogtrop F, Perfect JR,
Carter DA. 2018. Phenotypic variability correlates with clinical outcome in
Cryptococcus isolates obtained from Botswanan HIV/AIDS patients. mBio
9(5):e02016-18.

8.   Mukaremera L, McDonald TR, Nielsen JN, Molenaar CJ, Akampurira A, Schutz C,
Taseera K, Muzoora C, Meintjes G, Meya DB, Boulware DR, Nielsen K. 2019. The
mouse inhalation model of *Cryptococcus neoformans* infection recapitulates strain
virulence in humans and shows that closely related strains can possess differential
virulence. Infection and Immunity 87(5):e00046-19.

9.   Gerstein AC, Jackson KM, McDonald TR, Wang Y, Lueck BD, Bohjanen S, Smith
KD, Akampurira A, Meya DB, Xue C, Boulware DR, Nielsen K. 2019. Identification
of pathogen genomic differences that impact human immune response and
disease during *Cryptococcus neoformans* infection. mBio 10(4):e01440-19.

10. Beale MA, Sabiiti W, Robertson EJ, Fuentes-Cabrejo KM, O'Hanlon SJ, Jarvis JN,
Loyse A, Meintjes G, Harrison TS, May RC, Fisher MC, Bicanic T. 2015. Genotypic
diversity Is associated with clinical outcome and phenotype in Cryptococcal
meningitis across southern Africa. PLOS Neglected Tropical Diseases
9(6):e0003847.

11. Desjardins CA, Giamberardino C, Sykes SM, Yu C-H, Tenor JL, Chen Y, Yang T,
Jones AM, Sun S, Haverkamp MR, Heitman J, Litvintseva AP, Perfect JR, Cuomo
CA. 2017. Population genomics and the evolution of virulence in the fungal
pathogen *Cryptococcus neoformans*. Genome Research 27:1207–1219.

12. Sionov E, Lee H, Chang YC, Kwon-Chung KJ. 2010. *Cryptococcus neoformans*
overcomes stress of azole drugs by formation of disomy in specific multiple
chromosomes. PLOS Pathogens 6(4):e1000848.

13. Semighini CP, Averette AF, Perfect JR, Heitman J. 2011. Deletion of *Cryptococcus*
neoformans AIF ortholog promotes chromosome aneuploidy and fluconazole-

resistance in a metacaspase-independent manner. PLOS Pathogens
7(11):e1002364.

14. Stone NRH, Rhodes J, Fisher MC, Mfinanga S, Kivuyo S, Rugemalila J, Segal ES,
Needleman L, Molloy SF, Kwon-Chung J, Harrison TS, Hope W, Berman J, Bicanic
T. 2019. Dynamic ploidy changes drive fluconazole resistance in human
cryptococcal meningitis. J Clin Invest 129:999–1014.

15. Chen Y, Farrer RA, Giamberardino C, Sakthikumar S, Jones A, Yang T, Tenor JL,
Wagih O, Van Wyk M, Govender NP, Mitchell TG, Litvintseva AP, Cuomo CA,
Perfect JR. 2017. Microevolution of serial clinical isolates of *Cryptococcus*
neoformans var. *grubii* and *C. gattii*. mBio 8(2):e00166-17.

16. Rhodes J, Beale MA, Vanhove M, Jarvis JN, Kannambath S, Simpson JA, Ryan A,
Meintjes G, Harrison TS, Fisher MC, Bicanic T. 2017. A population genomics
approach to assessing the genetic basis of within-host microevolution underlying
recurrent cryptococcal meningitis infection. G3: Genes, Genomes, Genetics
7:1165–1176.

17. Fu MS, Liporagi-Lopes LC, Dos SR, Júnior S, Tenor JL, Perfect JR, Cuomo CA,
Casadevall A. 2021. Amoeba predation of *Cryptococcus neoformans* results in
pleiotropic changes to traits associated with virulence. mBio 12(2):e00567-21.

18. Gerstein AC, Fu MS, Mukaremera L, Li Z, Ormerod KL, Fraser JA, Berman J,
Nielsen K. 2015. Polyploid titan cells produce haploid and aneuploid progeny to
promote stress adaptation. mBio 6(5):e01340-15.

19. Berman J. 2016. Ploidy plasticity: a rapid and reversible strategy for adaptation to
stress. FEMS Yeast Research 16(3):fow020.

20. Hu G, Wang J, Choi J, Jung WH, Liu I, Litvintseva AP, Bicanic T, Aurora R, Mitchell
TG, Perfect JR, Kronstad JW. 2011. Variation in chromosome copy number
influences the virulence of *Cryptococcus neoformans* and occurs in isolates from
AIDS patients. BMC Genomics 12:526.

21.   Ormerod KL, Morrow CA, Chow EWL, Lee IR, Arras SDM, Schirra HJ, Cox GM,
637   Fries BC, Fraser JA. 2013. Comparative genomics of serial isolates of
638   *Cryptococcus neoformans* reveals gene associated with carbon utilization and
639   virulence. G3: Genes, Genomes, Genetics 3:675–686.

640   22.   Rhodes J, Desjardins CA, Sykes SM, Beale MA, Vanhove M, Sakthikumar S, Chen
641   Y, Gujja S, Saif S, Chowdhary A, Lawson DJ, Ponzio V, Colombo AL, Meyer W,
642   Engelthaler DM, Hagen F, Illnait-Zaragozi MT, Alanio A, Vreulink JM, Heitman J,
643   Perfect JR, Litvintseva AP, Bicanic T, Harrison TS, Fisher MC, Cuomo CA. 2017.
644   Tracing genetic exchange and biogeography of *Cryptococcus neoformans* var.
645   *grubii* at the global population level. Genetics 207:327–346.

646   23.   Yu C-H, Sephton-Clark P, Tenor JL, Toffaletti D, Giamberardino C, Haverkamp M,
647   Cuomo C, Perfect J. 2021. Gene expression of diverse *Cryptococcus* isolates
648   during infection of the human central nervous system. mBio 12(6):e0231321.

649   24.   Molloy SF, Kanyama C, Heyderman RS, Loyse A, Kouanfack C, Chanda D,
650   Mfinanga S, Temfack E, Lakhi S, Lesikari S, Chan AK, Stone N, Kalata N,
651   Karunaharan N, Gaskell K, Peirse M, Ellis J, Chawinga C, Lontsi S, Ndong J-G,
652   Bright P, Lupiya D, Chen T, Bradley J, Adams J, van der Horst C, van Oosterhout
653   JJ, Sini V, Mapoure YN, Mwaba P, Bicanic T, Lalloo DG, Wang D, Hosseinipour
654   MC, Lortholary O, Jaffar S, Harrison TS, ACTA Trial Study Team. 2018. Antifungal
655   combinations for treatment of cryptococcal meningitis in Africa. N Engl J Med
656   378:1004–1017.

657   25.   Fisher S, Barry A, Abreu J, Minie B, Nolan J, Delorey TM, Young G, Fennell TJ,
658   Allen A, Ambrogio L, Berlin AM, Blumenstiel B, Cibulskis K, Friedrich D, Johnson
659   R, Juhn F, Reilly B, Shammas R, Stalker J, Sykes SM, Thompson J, Walsh J,
660   Zimmer A, Zwirko Z, Gabriel S, Nicol R, Nusbaum C. 2011. A scalable, fully
661   automated process for construction of sequence-ready human exome targeted
662   capture libraries. Genome Biology 12(1):R1.

26. Loftus BJ, Fung E, Roncaglia P, Rowley D, Amedeo P, Bruno D, Vamathevan J,
Miranda M, Anderson IJ, Fraser JA, Allen JE, Bosdet IE, Brent MR, Chiu R,
Doering TL, Donlin MJ, D'Souza CA, Fox DS, Grinberg V, Fu J, Fukushima M,
Haas BJ, Huang JC, Janbon G, Jones SJM, Koo HL, Krzywinski MI, Kwon-Chung
JK, Lengeler KB, Maiti R, Marra MA, Marra RE, Mathewson CA, Mitchell TG,
Pertea M, Riggs FR, Salzberg SL, Schein JE, Shvartsbeyn A, Shin H, Shumway M,
Specht CA, Suh BB, Tenney A, Utterback TR, Wickes BL, Wortman JR, Wye NH,
Kronstad JW, Lodge JK, Heitman J, Davis RW, Fraser CM, Hyman RW. 2005. The
genome of the basidiomycetous yeast and human pathogen *Cryptococcus*
neoformans. Science 307:1321–1324.

27. Janbon G, Ormerod KL, Paulet D, Byrnes EJ, Yadav V, Chatterjee G, Mullapudi N,
Hon CC, Billmyre RB, Brunel F, Bahn YS, Chen W, Chen Y, Chow EWL, Coppée
JY, Floyd-Averette A, Gaillardin C, Gerik KJ, Goldberg J, Gonzalez-Hilarion S,
Gujja S, Hamlin JL, Hsueh YP, Ianiri G, Jones S, Kodira CD, Kozubowski L, Lam
W, Marra M, Mesner LD, Mieczkowski PA, Moyrand F, Nielsen K, Proux C,
Rossignol T, Schein JE, Sun S, Wollschlaeger C, Wood IA, Zeng Q, Neuvéglise C,
Newlon CS, Perfect JR, Lodge JK, Idnurm A, Stajich JE, Kronstad JW, Sanyal K,
Heitman J, Fraser JA, Cuomo CA, Dietrich FS. 2014. Analysis of the Genome and
Transcriptome of *Cryptococcus neoformans* var. *grubii* reveals complex RNA
expression and microevolution leading to virulence attenuation. PLoS Genetics
10(4):e1004261.

28. Farrer RA, Desjardins CA, Sakthikumar S, Gujja S, Saif S, Zeng Q, Chen Y, Voelz
K, Heitman J, May RC, Fisher MC, Cuomo CA. 2015. Genome evolution and
innovation across the four major lineages of *Cryptococcus gattii*. mBio
6(5):e00868-15.

29. Farrer RA, Chang M, Davis MJ, van Dorp L, Yang DH, Shea T, Sewell TR, Meyer
W, Balloux F, Edwards HM, Chanda D, Kwenda G, Vanhove M, Chang YC, Cuomo
CA, Fisher MC, Kwon-Chung KJ. 2019. A new lineage of *Cryptococcus gattii* (VGV)
discovered in the central zambezian miombo woodlands. mBio 10(6):e02306-19.

30. Li H. 2013. Aligning sequence reads, clone sequences and assembly contigs with
BWA-MEM. arXiv:13033997 [q-bio].

31. Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-
Moonshine A, Jordan T, Shakir K, Roazen D, Thibault J, Banks E, Garimella KV,
Altshuler D, Gabriel S, DePristo MA. 2013. From FastQ data to high confidence
variant calls: the Genome Analysis Toolkit best practices pipeline. Curr Protoc
Bioinformatics 43:11.10.1-11.10.33.

32. Cingolani P, Platts A, Wang LL, Coon M, Nguyen T, Wang L, Land SJ, Lu X,
Ruden DM. 2012. A program for annotating and predicting the effects of single
nucleotide polymorphisms, SnpEff: SNPs in the genome of *Drosophila*
melanogaster strain w [1118]; iso-2; iso-3. Fly 6:80–92.

33. Stamatakis A. 2014. RAxML version 8: a tool for phylogenetic analysis and post-
analysis of large phylogenies. Bioinformatics 30:1312–1313.

34. Abyzov A, Urban AE, Snyder M, Gerstein M. 2011. CNVnator: An approach to
discover, genotype, and characterize typical and atypical CNVs from family and
population genome sequencing. Genome Res 21:974–984.

35. Pfeifer B, Wittelsbürger U, Ramos-Onsins SE, Lercher MJ. 2014. PopGenome: an
efficient Swiss army knife for population genomic analyses in R. Mol Biol Evol
31:1929–1936.

36. Zhou X, Stephens M. 2012. Genome-wide efficient mixed-model analysis for
association studies. 7. Nature Genetics 44:821–824.

37. Wagih O, Parts L. 2014. gitter: A robust and accurate method for quantification of
colony sizes from plate images. G3 (Bethesda) 4:547–552.

38. Chun CD, Madhani HD. 2010. Applying genetics and molecular biology to the study
of the human pathogen *Cryptococcus neoformans*. Methods Enzymol 470:797–
831.

39. Idnurm A, Reedy JL, Nussbaum JC, Heitman J. 2004. *Cryptococcus neoformans*
virulence gene discovery through insertional mutagenesis. Eukaryot Cell 3:420–
429.

40. Toffaletti DL, Rude TH, Johnston SA, Durack DT, Perfect JR. 1993. Gene transfer
in *Cryptococcus neoformans* by use of biolistic delivery of DNA. J Bacteriol
175:1405–1411.

41. Zaragoza O, Casadevall A. 2004. Experimental modulation of capsule size in
Cryptococcus neoformans. Biol Proced Online 6:10–15.

42. National Research Council (US) Committee for the Update of the Guide for the
Care and Use of Laboratory Animals. 2011. Guide for the Care and Use of
Laboratory Animals, 8th ed. National Academies Press (US), Washington (DC).

43. Nielsen R, Williamson S, Kim Y, Hubisz MJ, Clark AG, Bustamante C. 2005.
Genomic scans for selective sweeps using SNP data. Genome Res 15:1566–1575.

44. Price MS, Betancourt-Quiroz M, Price JL, Toffaletti DL, Vora H, Hu G, Kronstad
JW, Perfect JR. 2011. *Cryptococcus neoformans* requires a functional glycolytic
pathway for disease but not persistence in the host. mBio 2(3):e00103-e111.

45. Chen Y, Toffaletti DL, Tenor JL, Litvintseva AP, Fang C, Mitchell TG, McDonald
TR, Nielsen K, Boulware DR, Bicanic T, Perfect JR. 2014. The *Cryptococcus*
neoformans transcriptome at the site of human meningitis. mBio 5(1):e01087-13.

46. Xue C, Liu T, Chen L, Li W, Liu I, Kronstad JW, Seyfang A, Heitman J. 2010. Role
of an expanded inositol transporter repertoire in *Cryptococcus neoformans* sexual
reproduction and virulence. mBio 1(1):e00084-10.

47. Liu T-B, Wang Y, Baker GM, Fahmy H, Jiang L, Xue C. 2013. The glucose sensor-
like protein Hxs1 Is a high-affinity glucose transporter and required for virulence in
Cryptococcus neoformans. PLOS ONE 8(5):e64239.

48. Li LX, Rautengarten C, Heazlewood JL, Doering TL. 2018. UDP-Glucuronic acid
transport is required for virulence of *Cryptococcus neoformans*. mBio 9(1):e02319-
17.

49. Li LX, Rautengarten C, Heazlewood JL, Doering TL. 2018. Xylose donor transport
is critical for fungal virulence. PLOS Pathogens 14(1):e1006765.

50. Brouwer AE, Rajanuwong A, Chierakul W, Griffin GE, Larsen RA, White NJ,
Harrison TS. 2004. Combination antifungal therapies for HIV-associated
cryptococcal meningitis: a randomised trial. Lancet 363:1764–1767.

51. Bicanic T, Muzoora C, Brouwer AE, Meintjes G, Longley N, Taseera K, Rebe K,
Loyse A, Jarvis J, Bekker L-G, Wood R, Limmathurotsakul D, Chierakul W,
Stepniewska K, White NJ, Jaffar S, Harrison TS. 2009. Independent association
between rate of clearance of infection and clinical outcome of HIV-associated
cryptococcal meningitis: analysis of a combined cohort of 262 patients. Clin Infect
Dis 49:702–709.

52. Jarvis JN, Bicanic T, Loyse A, Namarika D, Jackson A, Nussbaum JC, Longley N,
Muzoora C, Phulusa J, Taseera K, Kanyembe C, Wilson D, Hosseinipour MC,
Brouwer AE, Limmathurotsakul D, White N, van der Horst C, Wood R, Meintjes G,
Bradley J, Jaffar S, Harrison T. 2014. Determinants of mortality in a combined
cohort of 501 patients with HIV-associated Cryptococcal meningitis: implications for
improving outcomes. Clin Infect Dis 58:736–745.

53. Jung WH, Sham A, Lian T, Singh A, Kosman DJ, Kronstad JW. 2008. Iron source
preference and regulation of iron uptake in *Cryptococcus neoformans*. PLOS
Pathogens 4(2):e45.

54. Nichols CB, Ost KS, Grogan DP, Pianalto K, Hasan S, Alspaugh JA. 2015. Impact
of protein palmitoylation on the virulence potential of *Cryptococcus neoformans*.
Eukaryotic Cell 14:626–635.

55. Fu C, Donadio N, Cardenas ME, Heitman J. 2018. Dissecting the roles of the
calcineurin pathway in unisexual reproduction, stress responses, and virulence in
Cryptococcus deneoformans. Genetics 208:639–653.

56. Horianopoulos LC, Hu G, Caza M, Schmitt K, Overby P, Johnson JD, Valerius O,
Braus GH, Kronstad JW. 2020. The novel J-Domain protein Mrj1 is required for
mitochondrial respiration and virulence in *Cryptococcus neoformans*. mBio
11(3):e01127-20.

57. Okagaki LH, Wang Y, Ballou ER, O'Meara TR, Bahn Y-S, Alspaugh JA, Xue C,
Nielsen K. 2011. Cryptococcal titan cell formation is regulated by G-Protein
signaling in response to multiple stimuli. Eukaryot Cell 10:1306–1316.

58. Trevijano-Contador N, Oliveira HC de, García-Rodas R, Rossi SA, Llorente I,
Zaballos Á, Janbon G, Ariño J, Zaragoza Ó. 2018. *Cryptococcus neoformans* can
form titan-like cells in vitro in response to multiple signals. PLOS Pathogens
14(5):e1007007.

59. Perfect JR, Lang SD, Durack DT. 1980. Chronic cryptococcal meningitis: a new
experimental model in rabbits. Am J Pathol 101:177–194.

60. Muszewska A, Steczkiewicz K, Stepniewska-Dziubinska M, Ginalski K. 2017. Cut-
and-paste transposons in fungi with diverse lifestyles. Genome Biology and
Evolution 9:3463–3477.

61. Iyer LM, Zhang D, de Souza RF, Pukkila PJ, Rao A, Aravind L. 2014. Lineage-
specific expansions of TET/JBP genes and a new class of DNA transposons shape
fungal genomic and epigenetic landscapes. Proc Natl Acad Sci USA 111:1676–
1683.

62. Li H, Li Y, Sun T, Du W, Li C, Suo C, Meng Y, Liang Q, Lan T, Zhong M, Yang S,
Niu C, Li D, Ding C. 2019. Unveil the transcriptional landscape at the
Cryptococcus-host axis in mice and nonhuman primates. PLOS Neglected Tropical
Diseases 13(7):e0007566.

63. Burke JE, Longhurst AD, Merkurjev D, Sales-Lee J, Rao B, Moresco JJ, Yates JR,
Li JJ, Madhani HD. 2018. Spliceosome profiling visualizes operations of a dynamic
RNP at nucleotide resolution. Cell 173:1014-1030.e17.

64. Moyrand F, Klaproth B, Himmelreich U, Dromer F, Janbon G. 2002. Isolation and
characterization of capsule structure mutant strains of *Cryptococcus neoformans*.
Mol Microbiol 45:837–849.

65. Li LX, Hole CR, Rangel-Moreno J, Khader SA, Doering TL. 2020. *Cryptococcus*
neoformans evades pulmonary immunity by modulating xylose precursor transport.
Infect Immun 88(8):e00288-20.

66. Vanhove M, Beale MA, Rhodes J, Chanda D, Lakhi S, Kwenda G, Molloy S,
Karunaharan N, Stone N, Harrison TS, Bicanic T, Fisher MC. 2017. Genomic
epidemiology of *Cryptococcus* yeasts identifies adaptation to environmental niches
underpinning infection across an African HIV/AIDS cohort. Molecular Ecology
26:1991–2005.

67. Ashton PM, Thanh LT, Trieu PH, Van Anh D, Trinh NM, Beardsley J, Kibengo F,
Chierakul W, Dance DAB, Rattanavong S, Davong V, Hung LQ, Chau NVV, Tung
NLN, Chan AK, Thwaites GE, Lalloo DG, Anscombe C, Nhat LTH, Perfect J,
Dougan G, Baker S, Harris S, Day JN. 2019. Three phylogenetic groups have
driven the recent population expansion of *Cryptococcus neoformans*. Nature
Communications 10(1):2035.

68. Litvintseva AP, Carbone I, Rossouw J, Thakur R, Govender NP, Mitchell TG. 2011.
Evidence that the human pathogenic fungus *Cryptococcus neoformans* var. *grubii*
may have evolved in Africa. PLoS ONE 6(5):e19688.

69. Khayhan K, Hagen F, Pan W, Simwami S, Fisher MC, Wahyuningsih R,
Chakrabarti A, Chowdhary A, Ikeda R, Taj-Aldeen SJ, Khan Z, Ip M, Imran D, Sjam
R, Sriburee P, Liao W, Chaicumpar K, Vuddhakul V, Meyer W, Trilles L, Iersel LJJ
van, Meis JF, Klaassen CHW, Boekhout T. 2013. Geographically structured

populations of *Cryptococcus neoformans* variety *grubii* in Asia correlate with HIV
status and show a clonal population structure. PLOS ONE 8:e72222.

70. Andrade-Silva LE, Ferreira-Paim K, Ferreira TB, Vilas-Boas A, Mora DJ, Manzato
VM, Fonseca FM, Buosi K, Andrade-Silva J, Prudente B da S, Araujo NE, Sales-
Campos H, Silva MV da, Júnior VR, Meyer W, Silva-Vergara ML. 2018. Genotypic
analysis of clinical and environmental *Cryptococcus neoformans* isolates from
Brazil reveals the presence of VNB isolates and a correlation with biological
factors. PLOS ONE 13(3):e0193237.

71. Palma-Guerrero J, Hall CR, Kowbel D, Welch J, Taylor JW, Brem RB, Glass NL.
2013. Genome wide association identifies novel loci involved in fungal
communication. PLoS Genetics 9(8):e1003669.

72. Gao Y, Liu Z, Faris JD, Richards J, Brueggeman RS, Li X, Oliver RP, McDonald
BA, Friesen TL. 2016. Validation of genome-wide association studies as a tool to
identify virulence factors in *Parastagonospora nodorum*. Phytopathology
106:1177–1185.

73. Alanio A, Vernel-Pauillac F, Sturny-Leclère A, Dromer F. 2015. *Cryptococcus*
neoformans host adaptation: toward biological evidence of dormancy. mBio
6(2):e02580-14.

74. Hommel B, Sturny-Leclère A, Volant S, Veluppillai N, Duchateau M, Yu C-H,
Hourdel V, Varet H, Matondo M, Perfect JR, Casadevall A, Dromer F, Alanio A.
2019. *Cryptococcus neoformans* resists to drastic conditions by switching to viable
but non-culturable cell phenotype. PLOS Pathogens 15(7):e1007945.

75. Yang F, Teoh F, Tan ASM, Cao Y, Pavelka N, Berman J. 2019. Aneuploidy
enables cross-adaptation to unrelated drugs. Molecular Biology and Evolution
36:1768–1782.

76. Torres EM, Sokolsky T, Tucker CM, Chan LY, Boselli M, Dunham MJ, Amon A.
2007. Effects of aneuploidy on cellular physiology and cell division in haploid yeast.
Science 317:916–924.

Data Availability

Isolate sequence data can be accessed in NCBI via accession PRJNA764746.

Ethics

The ACTA trial from which the isolates described here were collected had ethical approval from the London School of Hygiene and Tropical Medicine Research Ethics Committee and by all the site national research ethics committees and regulatory bodies. De-identified clinical metadata (fungal burden, fungal clearance (EFA), patient outcome) was provided by investigators for analysis performed here.

Acknowledgments

We thank the Broad Institute Genomics Platform for generating the sequence data for this study. We thank all of the patients and their families, as well as the staff at all the sites not directly involved in the ACTA trial; Andrew Nunn, Halima Dawood, Andrew Kitua, and William Powderly for serving on the data and safety monitoring committee; Graeme Meintjes, Calice Talom, Newton Kumwenda, and Maryline Bonnet for serving on the trial steering committee; and the ANRS Staff in Paris (Brigitte Bazin, Claire Rekacewicz, and Paula Garcia) for constant support.

Funding

This project has been funded in part with Federal funds from the National Institute of Allergy and Infectious Diseases, National Institutes of Health, Department of Health and Human Services, under award U19AI110818 to the Broad Institute and Public Health Service Grants AI73896 and AI93257 to JRP. CAC is a CIFAR fellow in the Fungal Kingdom Program. RSH is a NIHR Senior Investigator. The views expressed in this

874 publication are those of the author(s) and not necessarily those of the NIHR or the
875 Department of Health and Social Care.

Figure 1. Maximum likelihood phylogeny of patient isolates, estimated from 72,258 segregating SNP sites, rooted to VNII. Isolates separate distinctly into VNI, VNB, and VNII, with all lineages having 100% bootstrap support. All isolates possess MATα except ACTA3523 (highlighted). Colored circles correspond to patient survival. Greyscale squares indicate patient fungal burden of cerebrospinal fluid prior to treatment, Log10(CFU/ml).

Figure 2. A) Correlation coefficient between clinical phenotypes, p < 0.0001 in orange. Asterisk indicates phi coefficient. Days survived indicates the total number of days survived for all individuals. Days survived (Died) indicates the number of days survived only for individuals that died during the ACTA clinical trial. B) Deaths (top, grey) and survival (bottom) of patients by infecting isolate lineage. C) Probability of survival, by lineage of infecting isolate. D) Log_{10} CSF CFU/ml (fungal burden) by infecting isolate lineage, asterisk indicates p < 0.05, Wilcoxon test. E) Rate of clearance (EFA) by infecting isolate lineage. Displayed as -1(gradient), asterisk indicates p < 0.05 and 0.01, Wilcoxon test.

Figure 3. A) Manhattan plot displaying variants associated with high fungal burden. Variants with an association score < 0.000005 (score test) are labeled in red. B) Survival of mice infected with parental strain (H99) and a CNAG_04102 mutant strain. Five CD-1 mice were infected with approximately 5×10^4 CFU by oropharyngeal aspiration. C) Survival of mice infected with parental strain (CM026) and a CNAG_05608 mutant strain. Five CD-1 mice were infected with approximately 5×10^4 CFU by oropharyngeal aspiration. One mouse infected with CNAG_05608 was excluded from analysis as an outlier. At Day 45 of infection, this mouse was assessed for fungal burden in the lung and brain and both tissues were sterile. D) Rabbit CSF CFU's for the parental strain (H99) and a CNAG_06033 mutant strain extracted on days 3, 7, and 10 post-infection. Three rabbits were infected per strain. E) Survival of mice infected with parental strain (H99) and a CNAG_06033 mutant strain. Five CD-1 mice were infected with approximately 5×10^4 CFU by oropharyngeal aspiration.

Figure 4. Colony size (area, px) of isolates grown on YPD at A) 30°C, B) 37°C, C) 39°C. Correlation of isolate growth (area, px) on YPD, with axes corresponding to colony size when grown at D) 30°C and 37°C, E) 37°C and 39°C. Colors correspond to isolate lineage: VNIa (dark blue), VNIb (light blue), VNII (purple), VNB (orange).

Figure 5. A) Whole (aneuploid) and partial chromosomal duplication frequency throughout the population, by chromosome. Colony size (growth) on YPD, by ploidy state at B) 30°C, C) 37°C. D) 39°C.