A chromosome-level reference genome of *Ensete glaucum* gives insight into diversity, chromosomal and repetitive sequence evolution in the Musaceae

Ziwei Wang[1,2], Mathieu Rouard[3,4], Manosh Kumar Biswas[5], Gaetan Droc[4,6,7], Dongli Cui[1,2], Nicolas Roux[3], Franc-Christophe Baurens[4,6], Xue-Jun Ge[1], Trude Schwarzacher[1,5], Pat (J.S.) Heslop-Harrison[1,5*] and Qing Liu[1,2*]

[1]Key Laboratory of Plant Resources Conservation and Sustainable Utilization / Guangdong Provincial Key Laboratory of Applied Botany, South China Botanical Garden, Chinese Academy of Sciences, Guangzhou, 510650, China

[2]College of Life Sciences, University of the Chinese Academy of Sciences, Beijing 100049, China

[3]Bioversity International, Parc Scientifique Agropolis II, 34397 Montpellier Cedex 5, France

[4]French Institute of Bioinformatics (IFB) - South Green Bioinformatics Platform, Alliance Bioversity and CIAT, CIRAD, INRAE, IRD, F-34398 Montpellier, France

[5]Department of Genetics and Genome Biology, University of Leicester, Leicester, LE1 7RH, UK

[6]CIRAD, UMR AGAP Institut, F-34398 Montpellier, France

[7]UMR AGAP Institut, Univ Montpellier, CIRAD, INRAE, Institut Agro, F-34398 Montpellier, France

*Correspondence: liuqing@scib.ac.cn (QL); phh4@le.ac.uk (JSHH)

Running title: A chromosome scale reference genome of *Ensete glaucum*

Abstract

Background: *Ensete glaucum* ($2n = 2x = 18$) is a giant herbaceous monocotyledonous plant in the small Musaceae family along with banana (*Musa*). A high-quality reference genome sequence of *E. glaucum* offers a vital genomic resource for functional and evolutionary studies of *Ensete*, the Musaceae, and more widely in the Zingiberales. **Findings:** Using a

combination of Illumina and Oxford Nanopore Technologies (ONT) sequencing, genome-

wide chromosome conformation capture (Hi-C), and RNA survey sequence, we report a high-

quality assembly of the 481.5Mb genome with 9 pseudochromosomes and 36,836 genes

(BUSCO 94.7%). A total of 55% of the genome is composed of repetitive sequences with

LTR-retroelements (37%) and DNA transposons (7%) predominant. The 5S and 45S rDNA

were each present at one locus, and the 5S rDNA had an exceptionally long monomer length

of c.1,056 bp, contrasting with the c. 450 bp monomer at multiple loci in *Musa*. A tandemly

repeated c. 134 bp satellite, 1.1% of the genome (with no similar sequence in *Musa*), was

present around all nine centromeres, with a LINE retroelement also found at *Musa*

centromeres. The assembly, including centromeric positions, enabled us to characterize in

detail the chromosomal rearrangements occurring between the $x = 9$ species and $x = 11$

species of *Musa*. Only one chromosome has the same gene content as *M. acuminata* (ma).

Three ma chromosomes represent part of only one *E. glaucum* (eg) chromosome, while the

remaining seven ma chromosomes are fusions of parts of two, three, or four eg chromosomes,

demonstrating complex and multiple evolutionary rearrangements in the change between $x =$

9 and $x = 11$. **Conclusions:** The advance towards a Musaceae pangenome including *E.*

glaucum, tolerant of extreme environments, makes a complete set of gene alleles available for

crop breeding and understanding environmental responses**.** The chromosome-scale genome

assembly show how chromosome number evolves, and features of the rapid evolution of

repetitive sequences.

Keywords: Chromosome scale assembly; *Ensete glaucum* genome; Nanopore; Hi-C

sequencing; Musaceae evolution; repetitive DNA; centromeres; retrotransposons;

translocations; synteny

Background

The genus *Ensete* Bruce ex Horaninow includes 10 species of giant, herbaceous monocotyledonous plants in the family Musaceae, native to tropical Africa and Asia [1]. Among them, the African species *E. ventricosum* (Welw.) Cheesman is an important food crop for more than 20 million people in Ethiopia [2]. Its sister genus *Musa,* grown throughout the tropics for food and fibre, includes diploid species, triploids and hybrids of *M. acuminata* and *M. balbisiana,* with banana cultivars. Sister to the grasses (Poales) and palms (Arecales) in monocots, both *Ensete* and *Musa,* along with a third genus *Musella,* belong to Musaceae in the order Zingiberales (gingers and bananas) [3]. Following rapid diversification of the Zingiberales at the Cretaceous/ Tertiary boundary (>65 Mya) the crown node age of the Musaceae family is soon after, with the *Musa* genus diverging from *Ensete* and *Musella* about 40 Mya [4,5].

Ensete glaucum (Roxb.) Cheesman (NCBI:txid482298), like other species in *Ensete*, is monocarpic with a dilated and characteristically glaucous basal pseudostem, with a small number of large seeds (10 mm in diameter) in elongated, banana-like fruits, borne in hands with a terminal flower (Fig. 1A-E), and is diploid with $2n = 2x = 18$ chromosomes [1,6–10]. *E. glaucum* is widely distributed in Asia and has records from Burma, China, India, Indonesia, India, Indonesia, Laos, Myanmar, Vietnam, Philippine, Papua New Guinea, Thailand, Solomon Islands [11].

Originating in the tropics and subtropics at lower elevations, most species in Musaceae lack cold acclimation. Cold stress is one of the key limitations in extending banana planting and production to higher altitudes and beyond the tropics [12]. In contrast to other Musaceae species which would not survive, *Ensete glaucum* can be found above 1000m in the mountains of Yunnan in China, where the temperature drops lower than 0°C, with limited

rainfall in winter. As one of the most cold-resistant and perhaps the most drought-tolerant

species in Musaceae, *E. glaucum* is a potential gene and germplasm resource for abiotic stress

tolerance in banana breeding, likely to be required for the adaptation to a more variable and

extreme climate in the future.

Whole genome sequences (genome assemblies) are published for some species in

Musaceae, with pseudochromosome-level data only available in *Musa* [13–15]. Draft genome

assemblies in *Ensete* species are limited to only *Ensete ventricosum,* and these are with tens

of thousands of contigs with an N50 lengths mostly between 10,000 and 21,000 bp [16,17].

As such, effective introduction and utilization of genetic resources present in wild species of

Ensete are a pressing need for banana improvement.

We applied Illumina, Oxford Nanopore Technologies (ONT), and chromosome

conformation capture (Hi-C) sequencing to generate a high-quality chromosome-level

assembly of the *Ensete glaucum* genome. We aimed to use the chromosome sequence to

show the genome structure and gene composition, as well as revealing the repetitive DNA

organization. The structural variations of *Ensete glaucum* ($x = 9$) were studied in a

comparative context with *Musa* (x = 11) species, showing the evolutionary history of the

family. The study aims to be useful in expanding the genepool available not only to banana

and enset breeders, but also for plant conservation of biodiversity in ecologically sensitive or

threatened areas, and for fundamental research on chromosome and genome evolution.

Analyses, Results and Discussion

De novo chromosome-scale genome assembly

A *de novo* chromosome level assembly of *Ensete glaucum* was made by combining high-

coverage ONT long-read sequencing, Illumina 150bp paired-end sequences, and Hi-C

chromosome conformation capture sequence data (Table 1). From the first assembly (with

124 contigs with an N50 of 10.256 Mb, Table 2 and Supplementary Table S1), we assembled

nine pseudomolecules, eg01 to eg09 (Fig. 2 and Supplementary Table S2), corresponding to

the chromosome number (2n=18) and observed chromosome morphology.

Table 1: Statistics of whole-genome sequencing and transcriptome analysis of *Ensete*

***glaucum* using Illumina, ONT, and Hi-C.**

Type	Method	Number of reads	Clean data (Gb)	Length (bp)	Coverage (×)
Genome	Illumina	245,852,534	37	2 × 150	74
	ONT	4,357,035	109	38,885 (N50)	220
	Hi-C	319,793,734	48	2 × 150	
Transcriptome	Illumina (Leaf)	62,410,840	94	2 × 150	
	Illumina (Root)	13,712,542	21	2 × 150	

A Hi-C/ONT-only assembly was constructed first, by using an OLC (overlap layout-

consensus)/string graph method with corrected reads. Contigs were refined using Illumina

short reads, and after discarding redundant contigs, the final genome assembly was 481Mb

long, with 9 chromosomes between 42,457,113 and 67,484,389bp long. Our BUSCO analysis

[18] was used to assess the assembly in 'genome' mode showing 98.3% complete single and

duplicated Embryophyta core gene sets from the embryophyta_odb10 database (consisting of

1614 genes) (Table 2, Supplementary Table S3a); few genes were fragmented or missing. *E.*

glaucum chromosome designations were chosen to follow major regions of synteny with

Musa acuminata chromosomes [19,20].

118 Genome size and heterozygosity estimation

The final genome assembly after Hi-C scaffolding (481,507,213 bp) anchored 97.2%

sequences of the contig-level assembly (495,175,598 bp; Table 2). Some arrays of tandem

repeats, including the rDNA (see below), were collapsed and chromosome termini were not

fully assembled. Around 55% of the assembled genome was estimated to be repeat sequences

(RepeatMasker; Table 2). The genome size was estimated as 563,295,571bp (highest 17-mer

peak frequency), with slightly higher estimates made by findGSE software (588,939,614 bp).

The genome size of *E. glaucum* is similar to that of the *x*=11 *Musa* species (see [20]) using

sequencing methods, and to estimates of both genera by flow cytometry [21].

127   The heterozygosity rate of *E. glaucum* was 0.164% (Supplementary Fig. S1 estimated

with k=21 using GenomeScope). Heterozygosity in plants is influenced by mating-systems

and pollination [22], life span, habitat fragmentation and cultivation [23]. There is relatively

little known about the breeding system and pollination of *Ensete* species (see [24]) although

we observed insects (including the hornet *Vespa bicolor,* a widespread pollinator in southern

China) visiting flowers (Fig. 1C). Our low level of heterozygosity is within the range found

in individual plants in populations of *M. acuminata* ssp. *banksii* (0.02% to 0.34% in 24

individuals, [25]; and 0.13% to 0.23%, [26]), and in other wild monocotyledonous species

including two (most likely self-pollinating) diploid oat species (0.07% heterozygosity in

Avena atlantica and 0.12% *A. eriantha* [27]); it is however, low compared to other species

(e.g. walnut, *Juglans nigra* 1.0%, [28]; *Nyssa sinensis* 0.87%, [29]) and in particular many

Musa species, some with known hybrid genome composition [26]. The low value seen in

species including *E. glaucum* here, is consistent with frequent self-pollination and inbreeding,

or a population bottleneck of this monocarpic tropical plant [30].

Table 2: Statistics of *Ensete glaucum* genome assembly and annotation. RepeatMasker did
not identify satellite sequences. 5S, 45S and Egcen were identified manually in assemblies
and the abundance measured in raw read data; microsatellite abundance was calculated from
the assemblies. (See Supplementary Tables S1, S2, S3, S10, S12, S14)

Genome assembly	Value
k-mer estimation of genome size (17-mer)	563,295,571 bp
Total contig length	495,175,598 bp
% of estimated genome	87.9 %
Anchored into chromosomes	481,507,213 bp
GC content	38.21 %
Contigs	
Number	124
N50 length	10,255,891 bp
Longest	31,226,749 bp
Pseudo-chromosomes	
Number	9
Shortest	42,457,113 bp
Longest	67,484,389 bp
N50 length	55,030,165 bp
L50 number	5
Complete BUSCOs (C) of genome	98.3 %
RepeatMasker repetitive DNA	
Transposable elements	
LTR retroelements	37.20 %
Copia	17.64 %
Gypsy	19.25 %
LINEs	0.77 %
Class II DNA transposons	7.18 %
Unclassified dispersed repeats	8.74 %
Simple repeats & low complexity	1.13 %
Total repeats (RepeatMasker)	55.02 %
Tandem repeat content	
45S rDNA	1.21 %
5S rDNA	0.08 %
Egcen satellite	1.32 %
Microsatellites (<8bp motif)	0.59%
Protein-coding genes	
Number	36836
Average number of exons per gene	4.86
Average exons length per gene	1114 bp
Average intron length	2816 bp
Average length of predicted proteins	371 aa
Complete BUSCOs (C) of predicted genes	94.7%
Functional annotation	
NR	31599 (85.78%)
InterPro	30160 (81.88%)
GO	24436 (66.34%)
KO	11192 (30.38%)
Total	31804 (86.34%)

Gene distribution and whole-genome duplication analysis

Genes were unevenly distributed along chromosomes (Fig. 2 circle b), and generally depleted in broad centromeric regions; few genes were found on the short arm of the more acrocentric chromosome eg04 and the Nucleolus Organizing Regions (NOR) bearing chromosome arm of eg06 (see rDNA below). The centromeric, gene poor regions, are rich in repeats (Fig. 2 circle c), transposable elements (*Copia* and *Gypsy* LTR retroelements, Fig. 2 circles d, e, and, less markedly, DNA transposons, Fig. 2 circle f), as observed in many species (including *Musa*,[14,15]).

The Ks (the synonymous rates of substitution) between genes in paired collinearity gene groups were calculated between *E. glaucum* and *M. acuminata* to see whether they share the same three WGD events [14]. The two genomes have a nearly identical Ks density distribution (Fig. 3A), both having two peaks at around 0.55 and 0.9. This result indicates that Musaceae share the same WGD events. The more recent peak at 0.55 most likely represents the α and β duplications, while the peaks at 0.9 may represent the more ancient γ duplication event [14]. Fig. 2 (center) links the genomic locations of paralogous gene clusters: most chromosome regions show shared relationships with three other chromosome regions, reflecting the α and β whole genome duplications, as shown by D'Hont et al. [14] (their Supplementary Figure 12).

Gene identification

Genes and gene ontology

In total, 36,836 genes were predicted (BUSCO score: C:94.7%, Supplementary Table S3B) with 31,804 (86.34%) functionally annotated with protein domain signatures and 24,436 (66.34%) associated with GO terms (Table 2; Supplementary Table S4). *E. glaucum* has a similar gene space (Supplementary Table S5) to the sequenced *Musa* species *M. acuminata*

(35,264), *M. balbisiana* (35,148), *M. itinerans* (32,456) and *M. schizocarpa* (32,809). In the

Musaceae (i.e., *M. acuminata, M. balbisiana, M. schizocarpa, E. ventricosum* and *E.*

glaucum), we identified a total of 29,639 orthogroups including 173,025 (88.1%) assigned

genes and 23,355 (11.9%) unassigned genes (Fig. 3B and Supplementary Table S5). Between

all species, the analysis showed 48% (n=14,523) of assigned genes were shared (core or

pseudo-core genes; rising to 66%, n=19,583 if genes missing in only one species are

discounted as possible annotation artefacts). The analyses highlighted 5% (1471) of

orthogroups that are *Ensete* genus specific and not found in *Musa*. A total of 162 orthogroups

were found only in *E. glaucum* (Fig. 3B; lower than the value for *E. ventricosum* but the latter

is a draft genome status without RNA support and with fragmented contigs with likelihood of

a large number of redundant predicted genes). The predicted genes of *Ensete glaucum* were

compared to their orthologous genes in *M. acuminata* and Ka/Ks values between orthologous

pairs were calculated. Genes with Ka/Ks > 1 were under positive selection (Supplementary

Table S6).

Gene family expansion and contraction

Using *Musa* species and two other monocotyledonous species (in the same clade of the

Commelinids), *Phoenix dactylifera* (Arecaceae) and *Oryza sativa* (Poaceae), we explored

gene family expansion and contractions in *E. glaucum* (Fig. 3C and Supplementary Table S7).

Among 12,384 gene families shared by the MRCA (Most Recent Common Ancestor) of

these monocotyledons, there were large numbers of gene families expanding (1498 to 2184)

or contracting (817 to 3444) between the genomes of Musaceae, *Phoenix* and *Oryza* (Fig.

3C), presumably reflecting substantial differences in plant form between them. Similar,

although slightly lower, figures were reported between, for example, dicotyledons as diverse

as *Arabidopsis* (Brassicaceae), *Solanum* (Solanaceae) and *Cuscuta* (Convolvulaceae) [31].

Notably, though, our results show the largest expansion of gene families in the Musaceae

(2184), likely reflecting the whole genome duplication events not shared with the Poaceae or

Arecaceae (see also [32] in pineapple), and we find additional expansion in *E. glaucum*.

Large gene family losses were noted in *Oryza sativa*, *Phoenix dactylifera* and *Musa*

balbisiana (Supplementary Table S7).

Overall, *E. glaucum* showed enrichment of several GO biological processes

(Supplementary Fig. S2, Supplementary Table S8) compared to *Musa*. Among them,

"monosaccharide transmembrane transporter" (equal top hit), "carbohydrate transmembrane

transport" and "carbohydrate transport"; and among molecular functions, "monosaccharide

transmembrane transporter activity", "sugar transmembrane transporter activity", and

"carbohydrate transmembrane transporter activity" were all included in the top 20

enrichments. The genus *Ensete* is notable for its accumulation of starch in the pseudostem

and leaf bases with *E. ventricosum* cultivated as a staple starchy food in East Africa [2], and

perhaps this is reflected in the enrichment of certain carbohydrate transport GO terms.

Transcription Factors (TFs)

In total, 2637 putative TF genes were identified in the *E. glaucum* assembly, representing 7%

of all genes (Supplementary Table S9), which were classified by their signature DNA

Binding Domain (DBD) into 58 TF families (Fig. 3D). Similar to *M. acuminata*, the MYB

(myeloblastosis) superfamily of transcription factors (including 260 MYB TFs plus 109

MYB-related) was the largest family, with between 140 and 210 copies of each of the bHLH,

AP2/ERF, NAC, C2H2, WRKY and bZIP families. The identification and classification of

the TFs here provides a framework to explore regulatory networks in plants [33] with their

target genes, and to identify specific factors involved in important responses. Cenci et al. [34]

analyzed transcription factors involved in the regulation of tissue development and responses

to biotic and abiotic stresses and, particularly, the NAC plant-specific gene family, while

Xiao et al [35] discuss the importance of a HLH factor involved in starch degradation during

fruit ripening. *Ensete* and *Musa* differ in these characteristics so it will be interesting to

analyze differences in transcription factors responsible.

Repetitive DNA analysis

Repeat identification

A range of different programs were applied for repeat analysis, and, as has been considered

previously [36], there were differences in the repeats identified between approaches, and

small changes in parameters and reference sequences give substantial changes. Repeated

elements in the genome assembly were identified by RepeatMasker (Table 2 and

Supplementary Table S10) and amounted to 55% of the genome, the same range as other

plant species with similar DNA amount and, particularly, the genus *Musa* [14,15]. For

assembly-free identification of repeats, we used RepeatExplorer [37] to generate graph-based

clusters of similar sequence fragments: Illumina sequence reads are available from six

Musaceae species, allowing assembly-free comparisons (Supplementary Table S11 and

Supplementary Fig. S3); while there was a little more variation in proportion of reads in the

most abundant clusters, all had between 33% and 46% in the top clusters (>0.01% genomic

abundance, as defined in [35]). Transposable elements including LTR and non-LTR

retroelements, and class II DNA transposons, were found (Fig 4 and Supplementary Table

S11). Microsatellites and other repeats were further characterized by mining and dotplot

analysis as well as fluorescent *in situ* hybridization to chromosomes (Supplementary Tables

S12-S14 and Figs 5-7; see below). The organization of repetitive regions in the assembly was

sometimes verified by mapping individual ONT long reads to assembled repeat regions (eg.

Fig. 5A), and organization was generally confirmed, except for some long tandem arrays

which seem to be collapsed in the assembly due to high homology between repeat units. A

few ONT reads were found which included reversals of tandem arrays (head-to-head or tail-

to-tail junctions), potentially artefacts from both strands of the DNA molecule passing

sequentially through one pore, and these junctions need further investigation.

Fig. 4A compares the abundance and species-distributions of major repeat classes in

the Musaceae using the comparative genome analysis function of RepeatExplorer. All species

shared many transposons and rDNA sequences (Fig 4A, central region). However, genus-

specific retroelement variants were identified in *Musa* (Fig. 4A, left) and *Ensete*-with-

Musella (Fig. 4A, right), showing the separation of the two phylogenetic branches, supported

by extensive divergence of the repetitive sequence sub-families, and evolution in copy

number. Notably, satellite sequences (Fig. 4A center-right) were much more abundant and

some sequences (see centromere sequence below) were present exclusively in *Ensete*.

Transposable Elements

The most abundant class of repetitive elements were transposable elements, in particular LTR

retroelements. The distributions of *Copia* and *Gypsy* LTR retroelements along assembled

pseudo-chromosomes (Fig. 2 circles d and e) show greater abundance in proximal

chromosome regions. Approximately equal numbers of *Copia* and *Gypsy* elements (18 and 19%

of the genome respectively, Table 2) were found. This result contrasts with *M. acuminata,*

where *Copia* elements were considerably more frequent (29%) compared to *Gypsy* elements

(11%;[14]; Supplementary Table S11). The relative change in proportions of the two element

families, while the overall abundance remains the same, has implications for genome

evolution and the expansion or contraction of retrotransposon families which can be explored

in detail using the high-quality genome sequences where the elements are neither truncated

nor collapsed.

Analysis of RT domains identified subfamilies of LTR retroelements, with the families

showing different abundances in *E. glaucum* and *M. acuminata* (Supplementary Fig. S4).

Insertion times of LTR retroelement subfamilies (Figs 4B, C and Supplementary Fig. S5)

270 were calculated based on LTR divergence for *E. glaucum* and recalculated for *Musa* to allow

271 for identical software settings (see Material and Methods). In *E. glaucum,* both *Copia* and

272 *Gypsy* families show relatively constant activity over the last 2.5 My, with the major a peak

273 of insertion activity 3.5 to 5.5 Mya, (Fig. 4B, C) corresponding to the half-life of LTR-

274 elements [14]. The dynamic amplification of these elements is emphasized by individual sub-

275 families having bursts of amplification (Fig. 4B for *E. glaucum* and Supplementary Fig. S4

276 for *Musa*), with rounds of expansion of different elements. As shown by Wang et al. [15], *M.*

277 *balbisiana* has the most extensive LTR activity in the last 500,000 years, and *M. acuminata*

278 activity peaks around 1.5 Mya (Fig. 4C), in both cases with greater activity of *Copia* elements

279 (Supplementary Fig. S5), contrasting with *E. glaucum* with equal activity of both *Gypsy* and

280 *Copia* elements leading to a higher proportion of *Gypsy* elements within the genome of *E.*

281 *glaucum* compared to *Musa* (see above, and Supplementary Tables S10 and S11). This is also

282 evidenced by the larger number of *Musa* specific clusters identified as *Copia* Angela or Sire

283 elements while *Ensete* with *Musella* specific LTRs include more *Gypsy* Reina and Retand

284 elements (Fig. 4A). Wu et al. [38] discuss the rounds of amplification in *M. itinerans* with an

285 amplification burst after separation from *M. acuminata* about 5.8 Mya suggesting high

286 turnover of the elements. The results suggest a burst of retroelement amplification (the older

287 ones), sometime after the split of *Musa* and *Ensete,* and again more recently, perhaps after *E.*

288 *glaucum* split from other *Ensete* species.

289 **Tandem (satellite) repeats and centromeric sequences**

290 The repeat analysis revealed the presence of an abundant tandemly repeated sequence with a

291 monomer length of c. 134bp (Fig. 5A). The sequence, named Egcen, represents about 1.3%

292 of the *E. glaucum* genome (45,000 copies) (Table 2, Supplementary Table S12; GenBank:

293 OL310717), and forms arrays that are at places interspersed by the LINE element Nanica

294 (described in *M. acuminata;* [14]) and other sequences (Fig. 5B, see below). One or two

major arrays of Egcen repeats were found in the assemblies of all nine chromosomes (Fig.

5C). *In situ* hybridization of Egcen showed it was located around the primary, centromeric,

constrictions as seen by DAPI staining (Fig 5D, see also Figs. 7A and C). The hybridization

pattern of the FISH signal on all chromosomes showed variable strength and several sites

grouped closely together, corresponding to the pattern seen in the assembly. The location of

the Egcen arrays was therefore used to infer the centromere mid-point position in the *E.*

glaucum chromosome assemblies (Fig 2 outer circle, Fig. 8A and Supplementary Table S13).

Egcen was also detected at the centromeres of all *E. ventricosum* and *Musella*

lasiocarpa chromosomes (Fig. 5E, F), showing similar distribution patterns as in *E. glaucum*

(Fig. 5D), but it was not seen on *Musa* chromosomes by *in situ* hybridization (example of *M.*

balbisiana, Supplementary Fig. S6) nor found in analysis of assemblies of *M. acuminata, M.*

balbisiana or *M. schizocarpa* (Supplementary Fig. S7A). The comparative RepeatExplorer

clustering shows multiple satellite sequences found only in the *Ensete* and *Musella* genomes

that are not present in the three *Musa* species tested (Fig. 4A) and supports Egcen being part

of the tandem repeat birth and amplification that has occurred in *Ensete* and *Musella* after the

split from *Musa,* and contrasts with the younger insertion times found for *Musa* retroelements

(Supplementary Table S12 and Fig. 4C).

Tandem repeats or satellite DNA sequences are found around the centromeres of

many plant (and animal) species [39,40] and may be 'centromeric' or 'pericentromeric'. No

equivalent tandem repeats were found in *Musa* [14,20,41] and the *E. glaucum* Egcen is not

present either in *Musa* (Supplementary Figs S6 and, S7A). However, the centromeric regions

of all *M. acuminata* chromosomes have been shown to include multiple copies of a LINE

non-LTR retroelement, Nanica, both by *in situ* hybridization and bioinformatic analysis

[14,20]. Nanica-related sequences were also identified in the *E. glaucum* assembly, but with

less abundance than in *Musa* (Supplementary Fig. S7B); about 350 copies were mapped to

chromosomes, mostly (but not exclusively) present interspersed within, or adjacent, to Egcen

arrays (Fig. 5B, C and Supplementary Fig. S7B).

Assembly across centromeric regions including abundant repeats is difficult and

normally the tandem repeat elements are collapsed. The ONT long-molecule sequences

allowed detailed examination of parts of the centromere region of chromosomes. A dotplot of

an ONT read (coded 9c7e99b5, 96,300bp long) aligned to the assembly of eg04 shows the

complex organization of the Egcen array (Fig. 5B): this 100kb region includes a total of six

Egcen tandem blocks with between 3 and 126 repeats (a total of 385), five copies of Nanica

(some rearranged, pink boxes), and three diverse retroelements flanked by LTRs (green

boxes). Further copies of the Egcen tandem repeat occur in larger blocks over the following

450,000bp of the assembly, and no genes were identified in the region.

A characteristic 17bp long sequence, the canonical CENP-B box, is found within a

monomer of a tandem repeat at centromeres of many species including human [42] and has

been postulated to be necessary for binding of the centromeric CENP-B proteins regulating

formation of centromere-specific chromatin. Within the Egcen sequence, there was a CENP-

B related motif:

```
Egcen        ggctaaTACGTTGGTAATGGccagt
             |  ||||||| || ||
CENP-B       TNNNNTTCGTTGGAAACGGGA
             |  ||||| |   |  |||
WheatCCS1      TCGGTTGCATGCAGGA
             |  ||   ||   | |
ATcon     ctttgtCTTTGTATCTTCTAACAACAA
```

Although, similar CENP-B motifs have been found from wheat and *Brachypodium*

(CCS1, [43]) to *Arabidopsis* (ATcon, [44]; see also review [45]), the relevance of the CENP-

B related box to centromere function remains uncertain, particularly when no similar tandem

repeat is present in other species or the related genera such as *Musa* (see above). As is the

case in many other species, the relative roles of retroelements, tandem repeats, and

interspersed centromeric sequences, leading to recruitment of the centromeric proteins, are

uncertain: the exact sequence or sequences that mark the functional centromeres, remain

enigmatic. The identification of a centromeric tandem repeat and the assembly across all

centromere regions in *E. glaucum* together with data from *Musa* will allow protein binding

studies (with Chromatin Immunoprecipitation ChIP analysis) to resolve the functional

centromere.

Microsatellites (Simple sequence repeats, SSRs)

Microsatellites were searched using Phobos and an SSR mining pipeline ([46]; perfect SSRs

from mono to hexa-nucleotide repeats with 11 to 3 repeat numbers respectively). SSR

abundances, array lengths, and nucleotide base composition (70% were AT-rich) are shown

in Fig. 6A and Supplementary Table S14. The overall nature and abundances of

microsatellites were generally similar to *M. acuminata*, *M. balbisiana, M. itinerans* and *E.*

ventricosum (Table S14). An average of one SSR was found per 4000bp, with the density

lowest around the centromere and higher at the telomeres (Fig. 6B); they are excluded from

the 45S NOR chromosome arm of eg06, and their overall distribution is similar to the

distribution of genes but contrasts the more proximal distribution of LTR retroelements and

DNA transposons (Fig. 2). Individual microsatellite motifs, however, showed characteristic

and different distributions. The abundant microsatellites, CT and AAG, were synthesized as

labelled oligonucleotides probes and used as probes for FISH on chromosomes. Both the

bioinformatic analysis of the assembly (Fig. 6B) and FISH to chromosomes (Fig. 6C, D)

showed (AAG/CTT) has a relatively uniform distribution along chromosomes, while

(AG/CT) shows depletion in centromeric regions and greater abundance in distal parts of

chromosomes that are gene-rich (Fig.2). The constraints on microsatellite spread in the

genome are motif-specific, and, if SSR markers were to be used for genetic mapping, those

associated with genes (such as AG/CT) would potentially be more useful.

5S and 45S rDNA and rRNA genes

Tandem repeats of the rDNA were predominantly located within extended, complex, loci on

chromosomes eg05 (5S rDNA) and eg06 (45S rDNA) (Figs 2, 5C, 7, and 8). The 45S rDNA

monomer containing the 18S rRNA gene - ITS1 - 5.8S rRNA gene - ITS2 - 26S rRNA gene -

NTS (GenBank: OL310719) was 9984 bp long, typical but slightly longer than other plant

species. The NTS region includes in most cases 16 copies of a degenerate 180bp tandem

repeat. Based on occurrence in the unassembled Illumina reads, there were 587 copies of the

45S rDNA monomer (1.21% of the genome, Tables 2 and Supplementary Table S12).

Although in the whole genome assembly the rDNA array was collapsed, the strength of the *in*

situ hybridization signal using the rDNA sequence from wheat (Figs 7A,B) is consistent with

representing 1% of the genome. The long chromosome arm carrying the 45S NOR locus was

depleted in genes by 10-fold (average of 12.6 genes/Mb compared to 127.6/Mb on the short

arm; (Fig. 7D). The single site of 45S rDNA at the NOR (Nucleolar Organizing Region) per

chromosome set is similar to *Musa acuminata* and other *Musa* species [47], although not *E.*

gilletii (2n=18), where there are 4 pairs of sites [21].

The 5S rDNA (GenBank: OL310718) comprised the 5S rRNA gene (119bp long,

typical for all plants) and intergenic spacer (937bp), representing 0.078% of the genome or

approximately 366 copies, with a complete motif length of 1056bp (Supplementary Table

S12). The 5S rDNA locus lies in the middle of the short arm of chromosome eg05, in three

parts around 34.5M, 37.5M and 45.5M with multiple interruptions. An example of insertion

of a 4.7kb LTR retrotransposon-related sequences in the 5S rDNA array of tandem repeats is

shown in Fig 7E. In other regions of the *E. glaucum* ONT reads or assembly, the

retroelement-related sequences named Brep, reported in *Musa* [48], was also found in the 5S

rDNA arrays. Garcia et al. [49] show the rather unusual and highly complex structure of 5S

rDNA in *M. acuminata* using graph-based clusters of reads with multiple IGS and

retroelement components, supporting the complexity reported here in the *E. glaucum*

assembly. The multiple hybridization sites evident from the *in situ* hybridization site on one

pair of chromosomes (Fig. 7B,C), with several, non-continuous, signals visible in the

extended prometaphase chromosomes (Fig. 7C, insets) support the non-continuous nature of

the 5S rDNA array.

The 5S rDNA monomer length of 1056bp was exceptionally long in comparison to

any other plant species (typically 400-500 bp long). The first 400bp of the intergenic spacer

had no significant BLAST hits in GenBank, while the second part showed only short regions

with weak homology largely to chromosome assemblies of *Musa* species in GenBank. There

were no motifs characteristic of retroelements in the 937bp intergenic spacer. It is unclear

why the monomer length for the 5S rDNA in *E. glaucum* should be twice that typical in other

species, including *Musa*, and was to be relatively homogeneous over all copies (Fig. 7E).

Furthermore, in contrast to the single locus on eg04 of *E. glaucum*, all species of *Musa*

examined so far have multiple 5S rDNA sites (2, 3 or 4 per genome), and *E. gilletii* had 3

pairs of sites [21,47].

Different species in the Triticeae show wide variation in numbers and locations of both

45S and 5S rDNA sites, suggesting multiple and complex evolutionary rearrangements of the

chromosome arms [50] even in the absence of chromosomal rearrangements including

translocations and inversions. Dubcovsky and Dvorák [51] have considered the 45S rDNA

loci as the "nomads of the Triticeae genomes" given their repeated evolutionary changes in

position during species radiation without rearrangements of the genes of the linkage groups.

The depletion of genes in chromosomal regions extending over most of a chromosome arm

around the 45S rDNA genes is notable in *Musa* and *Ensete*, so chromosome rearrangements

can lead to loci moving, although other recombination, duplication, deletion or translocation

events must occur to alter the numbers of both 5S and 45S loci observed.

Synteny and chromosome rearrangements to *Musa*

Structural comparisons of the *E. glaucum* genome sequence (x = 9, chromosomes eg01 to

eg09) were performed with the high-quality assembled genomes of *M. acuminata* (x = 11,

ma01 to ma11; v4, [20]) using translated proteins. SynVisio [52] was used for syntenic block

visualization (Fig. 8A) and the comparison was extended to *M. balbisiana* (mb01 to

mb11;[15]; Supplementary Fig. S8). Sequence dotplots (Fig. 8B) and comparative karyotypes

(Fig. 8C and Supplementary Table S15) of the *E. glaucum* genome against the *M. acuminata*

genome were also analyzed.

Overall, the genome assemblies of *M. acuminata* and *E. glaucum* are very similar in

length and gene content (Supplementary Table S5). We observed high identity between

segments of the 9 chromosomes of *E. glaucum* and of the 11 chromosomes of the *Musa* (Fig

8A). Broad centromeric regions with few protein-coding genomes (Fig. 2) cannot show

syntenic domains. The number of collinear genes was 48,956 between *E. glaucum* and *M.*

acuminata, and 39,604 between *E. glaucum* and *M. balbisiana* (by comparison, the A and B

genome of *Musa* show 42,854 collinear genes). Chromosomes show rearrangements,

inversions, expansions or contractions by crossed, converging or spreading lines in the

Synvisio plots. The dotplot (Fig. 8B; single chromosome comparison in Supplementary Fig.

S9) shows that there are some syntenic regions where the genes are distributed over the same

length of chromosomes in both species (diagonal lines showing synteny at 45°, eg08/ma10),

while in other cases there is expansion in one genome and not in the other (lines of synteny

more vertical, eg04/ma04, or nearer horizontal, eg03/ma04). Many syntenic segments showed

curved lines (ma03/eg03), showing relative expansion of one genome at one end of the

conserved syntenic block, and expansion of the other genome at the other end.

One complete chromosome, ma05/eg05/mb05, was similar with the same gene content

in all three species (Fig. 8A and Supplementary Fig. S8), but it showed multiple internal

inversions and expansions/contractions. A nested pair of inversions was evident covering

10.4Mb near the start of the chromosome in *E. glaucum* with respect to *M. acuminata*

(8.84Mb region) in dotplots (Supplementary Fig. S9A) and by comparison of locations of

orthologous genes (Supplementary Fig. S10). In the context of the *E. glaucum* inversions, we

could also examine the ancestral structure of *M. acuminata* and *M. balbisiana* reported by

Wang et al. [15]. Notably, a major rearrangement involving an inversion between *M.*

acuminata ma05 and *M. balbisiana* mb05 [15] was the same inverted region as found one

eg05, with an additional nested inversion of 3.1Mb in eg05 (Supplementary Figs S8, S10).

Using positions of orthologous genes at the boundaries of syntenic regions, the inversion

structure between chromosomes eg05, ma05 and mb05 was clear. Regardless of the ancestral

condition, the result indicates closely similar inversion breakpoints were involved (at the ends

of the segment, Supplementary Fig. S10) twice during evolution.

Apart from chromosome 5, three further whole chromosomes of *M. acuminata* are

represented largely by a single, whole chromosome regions/arms of *E. glaucum,* with some

rearrangements occurring within the chromosomes (Fig. 8): ma01 is mainly the second (right)

arm of eg01; and ma02 is mainly the second (right) arm of eg02 (Fig. 8A); ma11 is entirely

the first (left) arm of eg09 (see details in the dot blot of Supplementary Fig. S9B). The other

arms of these three *E. glaucum* chromosomes (eg01, eg02 and eg09) and the remaining six

chromosomes are related to blocks of the remaining seven *Musa* chromosomes. Four *Musa*

chromosomes have translocated fusions of segments of two *Ensete* chromosomes. ma09 has

the intercalary region of eg07, with an intercalary segment of eg06 inserted within the eg07

region. ma10 includes parts of three *Ensete* chromosomes, while ma04 has four segments

from *Ensete* chromosomes (Figs 8A, C). The 45S rDNA on eg06 and ma10 are surrounded

by syntenic regions but are both depleted in genes (Fig. 2). In contrast, the 5S rDNA sites are

not surrounded by other orthologous genes (see above). The non-reciprocal translocation

noted by Wang et al. [15] of a terminal segment between ma03 and mb01 lies within a larger

syntenic block shared between ma03 and eg03. This suggests that the translocation occurred

in the *M. balbisiana* lineage (Supplementary Fig. S8).

In several chromosomes of both *E. glaucum* and *M. acuminata*, the breakpoints occur

in the centromeric regions (e.g., in eg02, eg03, eg07, eg08 and eg09; ma01, ma06, and ma10;

Fig 8A). While some breakpoints occur at or adjacent to centromeres, the exact relationship

of any breakpoint to the centromere and Egcen or Nanica sequences is diverse. Notably, eg03

is spanning the centromere of three *Musa* chromosomes, ma03, ma04 and ma08, and in other

cases centromere regions are different despite surrounding synteny (e.g., eg07, ma07 and

ma09). Telomeric or sub-telomeric regions are conserved between the two species in 7 of the

18 *E. glaucum* chromosome arms (e.g., eg01/ma01, eg06/ma07, eg08/ma10; the dotplot

homology lines end in the corners of the chromosomes, Figs 8B and Supplementary Fig.

S9B). In other chromosomes, telomeres in *E. glaucum* are in intercalary regions of *Musa* (see

eg03/ma03 with inversion, eg08/ma06 in Fig 8B; and in detail eg08 and ma10 in

Supplementary Fig. 9C). The homology of the whole chromosome ma11 and the left arm of

eg09 (see above) indicates a fusion/fission event with loss/gain of centromere and telomere

function, but they are also predicted for the other rearrangements discussed above.

Song et al. [10] review the data on the basic chromosome number of the Zingiberales,

concluding that x = 11 is most reasonable original basic number, with x = 9 as a derived basic

number. With chromosome numbers of x = 9, 10 and 11 predominant in Musaceae, this

family is particularly suitable to explore the nature and locations of chromosomal fusions and

fissions that are predicted to often occur in similar position during karyotype evolution (e.g.

in wheat [53]). The availability of high-quality genome sequences, based on the ONT, Hi-C

and in *Musa* BioNano and PacBio technologies, will allow the nature of breakpoints in

chromosome fission events to be investigated at the sequence level between the *Musa* x=11

and *Ensete* x=9 species, as well as being able to shed light to centromere and telomere

function.

Conclusions

We provide a chromosome-scale assembly of *Ensete glaucum,* a sister genus to *Musa*. This

assembly is valuable to infer *Musa* genome evolution, enabling comparison with putative last

common ancestors of *M. acuminata* (A genome) and *M. balbisiana* (B genome) at protein

and chromosomal levels. Most striking was the multiple rearrangements of chromosome

structures between *E. glaucum* and the *Musa* A and B genomes, with only 4 of the 11 *M.*

acuminata chromosomes (and only 3 of the 11 in *M. balbisiana*) showing synteny with only

one or part of one *E. glaucum* chromosome. Despite new insight to chromosome evolution,

the sampling still makes difficult to conclude between ascending or descending dysploidy,

but it should be clarified once additional chromosome scale genomes are released in

Musaceae (in particular the Callimusa section with n=7, 9 and 10) and in sister clades of the

Zingiberales.

As well as the complex chromosome rearrangements, repetitive sequences differ

extensively between the *Musa* and *Ensete* genera. There is a major tandem repeat at the

centromeres of only the *Ensete* species, showing lack of conservation of this key structural

element of chromosomes, although both genera have multiple copies of the Nanica

retroelement in centromeric regions. *E. glaucum* has only one 5S rDNA locus, with an

unusually long monomer of 1056bp.

The complete sequence provides an accurate reference for the genus for gene

identification, marker development, Genotyping-By-Sequencing and Genome Wide

Association Studies (GWAS) and will accelerate our understanding of the molecular bases of

traits such as cold tolerance and starch accumulation, and allow identification of relevant

genes. The work builds towards a complete pangenome of the Musaceae family, defining

structural, gene and genetic diversity which can be used for genetic improvement across the

Musaceae and potentially more widely.

Material, Methods and Validation

Sample collection and distribution

The individual *Ensete glaucum* plant used for genome sequencing and analysis was collected

from Puer city, Yunnan province, China and maintained in the South China Botanical

Garden, Guangdong province, China (accession no. 19990288; Fig. 1A-E). The distribution

of *Ensete* and *Musa* species were identified in databases of Flora of China, South China

Botanical Garden, iNaturalist, GBIF [54] (excluding cultivation sites) and regional

distributions maps. Fig. 1F was then made from POWO [55] (overlaid and color-adjusted in

Adobe Photoshop CC2018).

DNA extraction and sequencing

Young leaves of *Ensete glaucum* were collected and ground into powder in liquid nitrogen.

High molecular weight genomic DNA was extracted using the DNeasy Plant Mini Kit

(Qiagen, Hilden, Germany). DNA quality was assessed by agarose gel electrophoresis and

NanoDrop 2000c spectrophotometry, followed by Thermo Fisher Scientific Qubit

fluorometry.

Illumina sequencing

A genomic DNA library with 400bp fragments was constructed using Truseq Nano DNA HT

Sample preparation Kit (Illumina USA Preparation Kit (Illumina, USA), and 150bp paired

ends were sequenced with Illumina Novaseq by Grandomics Research Services (previously

known as Nextomics, Wuhan, PR China). After applying Trimmomatic v0.36 [56] to trim

adaptors, filtering out low quality reads and further quality control with fastQC v0.11.9 [57],

246 million paired reads and 36.88Gb of data resulted (Table 1).

Oxford Nanopore (ONT) sequencing

ONT (Oxford Nanopore, Oxford, UK) sequencing was performed by Grandomics

Biosciences Co., Ltd. (Wuhan, China): long fragments longer than 12 kb were selected with

BluePippin (Sage Sciences), and the SQK-LSK109 kit (Oxford Nanopore) was used to build

a library that was sequenced using PromethION. The base calling was performed with Guppy

and reads mean_q score_template (Phred) > 7 (base call accuracy >80%) were selected. A

total of 129 Gb ONT reads (~ 250X coverage) was generated. fastp v0.19.7 [58] was used for

quality control including adaptor-trimming, filtering reads with too many Ns or mean q score

lower than 7 and resulted in remaining clean data of 109 Gb (Table 1). The mean read length

was c. 20kb, with the longest >120kb (Supplementary Fig. S11).

Hi-C chromatin interaction data

The Hi-C library was prepared followed by a procedure with an improved modification [59].

Nuclear DNA from fresh leaves was first fixed and cross-linked in situ with formaldehyde

[60], extracted, and digested by the restriction enzyme *Dpn*II. The resulting sticky ends were

biotinylated, diluted, and ligated. The biotinylated fragments which contained DNA of two

physically interacting regions were used to make a paired-end library and sequenced with

Illumina HiSeq platform (performed by Grandomix, loc. cit.). Quality control of Hi-C raw

data was performed using Hi-C-Pro v2.8.1 (HiC-Pro, RRID:SCR_017643): low-quality

sequences (quality scores<20), adaptor sequences, and sequences shorter than 30 bp were

filtered out using fastp v0.19.7 [58].

Genome and chromosome assembly

ONT data were corrected by Nextdenovo v2.0-beta.1 (Nextomics, 2019) and the 109 Gb

filtered data were assembled by SMARTdenovo (SMARTdenovo, RRID:SCR_017622) [61].

To polish the assembly, contigs were refined with Racon (Racon, RRID:SCR_017642) [62],

BWA v0.7.17 (BWA, RRID:SCR_010910) [63] was used to map the filtered Oxford

Nanopore reads to the assembly and NextPolish v1.3.1 [64] was used to discard possibly

redundant contigs and generate a final assembly; similarity searches were performed with the

parameters "identity 0.8 – overlap 0.8". Finally, BWA and Pilon v1.21 [65] were used to

further correct the assembly using the Illumina Novaseq reads, and two rounds of mapping

back to the assembly each time with further correction were undertaken. A 494Mb assembly

with 124 contigs was achieved (Table 2 and Supplementary Tables S1, S2).

Read pairs from the Hi-C data were mapped to the draft assembly using bowtie2 v2.3.2

(bowtie2; RRID:SCR_016368) [66] with the settings -end-to-end, -very-sensitive and -L 30

to select unique mapped paired-end reads. Valid interaction paired-end reads were identified

by HiC-Pro v2.8.1 (HiC-Pro, RRID:SCR_017643) [67] and retained for further analysis

while invalid read pairs, including dangling-end, self-cycle, re-ligation, and dumped products

were discarded. The scaffolds were further clustered, ordered, and oriented onto pseudo-

chromosomes by LACHESIS (RRID:SCR_017644) [68],with parameters as follows:

cluster_min_re_sites = 100, cluster_max_link_density = 2.5, cluster noninformative ratio =

1.4, order min n res in trunk = 60, order min n res in shreds=60. Finally, placement and

orientation errors exhibiting obvious discrete chromatin interaction patterns were manually

adjusted. A Hi-C contact matrix plot (Supplementary Fig. S12) showed no discontinuities in

the assembly.

Validation of the assembly was performed using Benchmarking Universal Single-Copy

Orthologs with (BUSCO, RRID:SCR_015008) v5 [18] to assess the completeness and

presence of 1614 genes in the embryophyta_odb10 database in "genome" mode

(Supplementary Table S3a).

RNA extraction, sequencing and transcriptome assembly

Total RNA was extracted from fresh leaves and roots of the same individual of *Ensete*

glaucum that was used for genomic sequencing using RNeasy Plant Mini Kit (Qiagen).

Illumina libraries were built from 1 µg total RNA of each sample with TruSeq RNA Library

Preparation Kit (Illumina, USA) and were then sequenced using Illumina Novaseq platform

to generate paired-end reads. A transcriptome assembly was produced using RNAseq data

using Trinity (Trinity, RRID:SCR_013048) [69] and mapped on the genome with PASA

(PASA, RRID:SCR_014656) [70].

Genome size estimation

Using the Illumina DNA sequence, genome size was estimated from the 17-mer frequency

using Jellyfish v2.0 (Jellyfish, RRID:SCR_005491) [71] with the formula k-num/k-depth

(where k-num is the total number of 17-mers, 30,417,960,841; and k-depth the highest k-mer

depth, 54; Table 2). 21-mer data were used in findGSE [72] and Genomescope 2.0

(Genomescope R, RID:SCR_017014) [73] to estimate the genome size and heterozygosity

(Supplementary Fig. S1).

Gene annotation

We adopted a combination of ab initio gene prediction, homology-based gene prediction and transcriptome-based gene prediction strategy. RepeatMasker v4.0.9 (RepeatMasker, RRID:SCR_012954) with option "-no_is -xsmall" was used to generate a repeat softmasked genome file. RNAseq data from leaf and root tissues were mapped to the masked genome assembly with STAR v2.7 (STAR, RRID:SCR_004463) [74]. The RNA alignment was input into BRAKER2 v2.1.5 (BRAKER, RRID:SCR_018964) [75], a combination of GeneMark (GENEMARK, RRID:SCR_011930) [76] and AUGUSTUS (RRID:SCR_008417) [77], to perform ab initio gene predictions. The gene models from BRAKER2 were inputted into MAKER v2.31.10 (MAKER, RRID:SCR_005309) [78] as model, and the RNA alignment of *E. glaucum* and proteins form *M. acuminata* v2 were used as EST and protein evidence, respectively. We also utilized GeMoMa v2.3 (GeMoMa, RRID:SCR_017646) [79] to perform homology-based gene prediction using Musa acuminata v2 [19] as reference annotated genome. EvidenceModeler (EvidenceModeler, RRID:SCR_014659) [80] was used to combine de novo and homology-based predictions and our transcriptome evidence to produce the final structural gene annotation.

To annotate the function of predicted genes, we performed BLASTP (e-value = 1e−10) (BLASTP, RRID:SCR_001010) from the BLAST+ package [81] for each predicted coding sequence against the databases: UniProtKB/Swiss-Prot, UniProtKB/TrEMBL [82] and NR (non-redundant protein database at NCBI). These sequences are then processed to produce a non-identical (often referred to as pseudo non-redundant) prediction. To assign a putative function to a polypeptide we kept only the best hit based on three parameters: (1) Qcov (Query coverage = length high-scoring segment pair (HSP)/length query), (2) Scov (Subject coverage = length HSP/length subject) and (3) identity. Additional functional information

was added by scanning sequences with InterProScan v5.46 (InterProScan,

RRID:SCR_005829) [83]. Blast2GO v6.0.1 (Blast2GO, RRID:SCR_005828) [84] was used

to integrate the results of blast and InterProScan, and to link the GO (Gene ontology) terms to

genes accordingly (Supplementary Table S4) The functional annotation procedure is given in

greater detail at [85].

BUSCO (BUSCO, RRID:SCR_015008; v5.0.0) was run in mode "transcriptome" using

the embryophyta_odb10 database to assess the gene annotation results and found 1529

(94.7%) complete BUSCOs (Supplementary Table S3b).

Gene family analyses

Orthogroups (OGs) identification in Musaceae

Protein-coding genes from *M. acuminata* [19], *M. balbisiana* v1.1 [15] and *M. schizocarpa*

*v*1 [13] were retrieved from the Banana Genome Hub [86]. Protein-coding genes predicted

from *E. ventricosum* was downloaded at NCBI Genome (GCA_000818735.3) to allow

discrimination of *Ensete* specific OGs *E. glaucum* specific OGs. Combined with *E. glaucum*

protein-coding genes, we used OrthoFinder v2.5.2 (RRID:SCR_017118) [87] and Diamond

[88] with default parameters (summary in Supplementary Fig. S5). Visualization (Fig. 3B)

was produced with UpsetR [89]. Gene ontology (GO) enrichments were calculated using

TopGO [90] with Fisher's exact test (Supplementary Fig. S2 and Supplementary Table S8).

Gene family expansion and contraction

To identify gene family expansion and contraction, we expanded previous analyses with

OrthoFinder by adding a representative of Musaceae sister clades in Palms (*Phoenix*

dactylifera, date palm, [91]) and Poales (*Oryza sativa v7*, rice, [92]; data downloaded from

Phytozome [93]) but omitting *E. ventricosum* due to gene redundancy. The longest transcripts

were kept if alternative splicing occurred. Divergence time estimation with Approximate

Likelihood Calculation used MCMCTREE in PAML v4.9j (PAML, RRID:SCR_014932).

Computational Analysis of gene Family Evolution (CAFE v4.2.1, RRID:SCR_018924) [94]

was used to model the evolution of gene family sizes and stochastic birth and death processes

and summarized in the phylogenetic tree (Fig. 3C).

Transcription factors

Protein coding gene sequences for *E. glaucum* and *M. acuminata* v2 were searched in

PlantTFDB v5.0 (PLANTTFDB, RRID:SCR_003362) and iTAK online v1.6 [95]. Predicted

transcription factors (TFs) were verified through a Hidden Markov Model (HMM) with

PFAM searching tools using the cutoff E-value of 0.01 (Fig. 3D and Supplementary Table

S9). Genes were verified by PFAM (Pfam, RRID:SCR_004726), CDD (Conserved Domain

Database, RRID:SCR_002077, and SMART (SMART, RRID:SCR_005026) databases.

671   Whole-genome duplication (WGD)

To identify the whole-genome duplication events, we applied WGDI pipeline (whole-genome

duplication identification v0.4.7 [96]). The predicted proteins of *E. glaucum* were blasted

against themselves and then a collinearity analysis was conducted. The *K*s (the synonymous

rates of substitution) between genes in paired collinearity gene groups were calculated and

the *K*s peak was detected. For comparison, the same processes were also applied to *Musa*

acuminata v2 [19].

678   Synteny analyses

Structural comparisons of the *E. glaucum* genome were performed with *M. acuminata* v4

(designated the A genome) and *M. balbisiana* (B genome). The very recent release of *M.*

acuminata v4 assembly was preferred for this analysis as it improved pericentromeric regions

and provided telomere-to-telomere gapless chromosomes [20]. Assemblies were aligned with

minimap2 (Minimap2, RRID:SCR_018550) [97] and visualized results using D-Genies (D-

GENIES, RRID:SCR_018967) v1.2.0 [98]. Protein-coding genes were processed to identify

reciprocal best hits (RBH) with BLASTP (e-value 1e-10) followed by MCScanX (e-value 1e-

05, max gaps 25) [99] and results imported in SynVisio [52] for syntenic block visualization.

Scale bars and coloring of the chromosome bars was adjusted using Adobe Photoshop

CC2018.

The karyotype of *E. glaucum* in Fig. 8C, was prepared from lengths of each pseudo

chromosome (Supplementary Table S2) with the estimated centromere position

(Supplementary Table S13) to estimate the left (darker colored) and right (lighter colored)

chromosome arms. Chromosome lengths and centromere positions for *M. acuminata* were

taken from [20] (Fig. 2a); syntenic blocks were calculated using the SynVisio diagram (Fig.

8A).

Repetitive DNA identification and annotation

For repetitive DNA analysis, publicly available programs (see below and [36]) as well as

manual searches and sequence comparisons were applied. Geneious v.10.2.6 (Geneious,

RRID:SCR_010519) (Biomatters Ltd., Auckland, New Zealand) was used to produce the

dotplots of Figs 5A, B, 7E and 8B.

In the assembly, repeated sequences were first searched with REPET v2.5 pipeline

[100]. The top 100 repeated sequences were plotted on the four reference *Musa* genome

assemblies (i.e. *M. acuminata*, *M. balbisiana, M. schizocarpa* and *E. glaucum*) using

BlastAndDrawDensity.py script described in [20] and available on the Github repository

[101],

Transposable Elements

Transposable elements were annotated by EDTA pipeline [102], which integrates various

software to discovers TE including: long terminal repeat (LTR) retrotransposons [103–105],

terminal inverted repeat (TIR) transposons [106], short TIR transposons or miniature inverted

transposable elements (MITEs) [107], and Helitrons [108]. According to suggestions in

[109], we also adopt RepeatModeler2 (RepeatModeler, RRID:SCR_015027) v2.0.1 [109] to

find remaining TEs.

The TE-denovo procedure was used on masked assembly to produce a batch of 4229

TE consensus sequences. From these 2800 consensus sequences, only those with full length

fragments present in the assembly were kept for further analysis, quantification and

annotation with the TE-ANNOT procedure. A first annotation was performed using public

Repbase (Repbase, RRID:SCR_021169) release 20.05, followed by *Gypsy/Copia*

retroelement family identification using Hidden Markov Models (hmmsearch version 3) to

search consensus for corresponding retrotransposase PFAM domains PF04195 and PF14244

respectively. The above results were then combined and CD-HIT v4.1.8 (CD-HIT,

RRID:SCR_007105) [110] was used to reduce redundancy. The LTR retrotransposons were

sent to TEsorter [111] to classify into lineage level and RT domain amino acid sequences

were extracted. Phylogenetic trees of *Copia* and *Gypsy* were inferred by RT domain

alignment results (Supplementary Figure S4). The proportions of transposons in the assembly

are given in Table 2 (Supplementary Tables S11 and S12) and chromosomal distributions in

Fig. 2.

To estimate ages of LTR transposons and the time of insertion (Fig 4B, C and

Supplementary Figure S5), complete elements were found by LTRharvest v1.6.1

(LTRharvest, RRID:SCR_018970) [103] and LTR_retriever (LTR_retriever,

RRID:SCR_017623) [104] and then classified by TEsorter v1.3. The estimation of time was

based on the divergence of of the 5' and 3' end LTRs, and these two LTRs of every LTR

retrotransposons were extracted into separate files with a custom script. The 5' and 3' LTRs

were aligned by MUSCLE v3.8.1551 [112]. The divergence distances under K2P

evolutionary model were calculated by R package ape v5.4-1 (ape, RRID:SCR_017343). The

average base substitution rate was selected to be 11.3E-8 [113]. The insertion time T was

calculated as T = K/(2r), with r as the rate of nucleotide substitution and K as the divergence

distance between LTR pairs. Our script to perform the analysis is on github (2021) [114].

Graph based clustering of reads using RepeatExplorer

A sample of 2Gb of the Illumina HiSeq raw reads were used for assembly-free analysis by

RepeatExplorer2 [37]. Graph-based clusters of similar sequence fragments were generated

under default parameters. Clusters were assigned to repeat classes and retroelement lineages

using the automated Repeat Masker and Domain hits provided by the program

(Supplementary Table S11). Comparative analysis with sample Illumina sequence reads from

other five Musaceae species, namely *M. acuminata* v2 [19], *M. balbisiana* v1.1 [15] and *M.*

schizocarpa v1 [13], *E. ventricosum* [17] and *Musella lasiocarpa* (Ziwei Wang, Qing Liu and

Pat Heslop-Harrison pers communication) were also analyzed (Supplementary Figure S3) and

compared with RepeatExplorer2 following "comparative repeat analysis" protocol. The

results were visualized by R script "plot_comparative_clustering_summary.R" (Fig, 4A).

SSR Tandem Repeats

The genome assembly was searched for SSR (microsatellite) motifs using the SSR mining

pipeline developed by Biswas et al [46]. Searches were standardized for mining perfect SSRs

from mono to hexa-nucleotide repeats (minimum repeat number of 12 for mononucleotides, 8

for di-, 5 for tri-, tetra- and penta-, and 4 repeats for penta- and hexa-nucleotides). SSRs

abundance and nature was analyzed based on density in the genome (about 1 per 4000bp),

array length (Figure 7B and Supplementary Table S14; SSR search parameters minimum

lengths mono=1*12=12nt, di=8*2=16nt, tri=3*5=15nt, tetra=4*5=15nt, penta=5*4=20nt and

hexa=6*4=24nt; total SSR count 123884; Class I>20nt and Class II≤20nt), nucleotide base

composition of the SSR loci (70% were AT-rich) and abundance of each motif).

758 Fluorescent *in situ* hybridization (FISH)

Chromosome preparation and FISH was performed as described by Schwarzacher and

Heslop-Harrison [115] with minor modifications. Plants of *E. glaucum, E. ventricosum,*

Musella lasiocarpa (purchased commercially) and *M. balbisiana* 'Butuhan' (ITC1074) [116]

were grown in the glasshouse at the University of Leicester, UK. Actively growing root tips

were treated with 2 mM 8-hydroxyquinoline and fixed with 96% ethanol:glacial acetic acid

(3:1). For chromosome preparations, roots were digested with a modified enzyme solution

(32U/ml cellulose, Sigma-Aldrich C1184; 20U/ml 'Onozuka' RS cellulose; 35U/ml pectinase

from *Aspergillus niger,* Sigma-Aldrich P4716; 20U/ml Viscozyme, Sigma-Alderich V2010)

in 10mM citric acid/sodium citrate buffer (pH4.6) for 3-5h at 37°C and then kept in buffer for

12-30h at 4°C. Meristems were dissected in 60% acetic acid and routinely 2-6 slide

preparations were made from each root. Slides were stored at -20°C until FISH.

The 45S rDNA probe was labelled by random priming (Invitrogen) with digoxigenin

dUTP (Roche) using the linearized clone pTa71 (Gerlach and Bedbrook 1979) containing the

45S rDNA repeat unit of *Triticum aestivum*. 50-100ng of labelled probe was used per slides

and detection of hybridization sites was carried out with Fluorescein-conjugated anti-

digoxigenin (Roche). The remaining probes were designed from the consensus sequence of

the centromeric repeat Egcen (Fig. 5) and the 5S rDNA (Fig. 7) or as simple sequence repeats

(Fig. 6); as directly labelled oligonucleotides (200-500ng per slide) they needed no further

detection and were as follows:

CenCy3: EGL_2640R: [Cyanine3]GAC CGT CGC ATT TTT TGG CGA AAC CAT GCT
CGT ACG ACT TCC CAT GGG CTA AAA CGT TAG GA

CenFAM: EGL_G2640L: [6FAM]GGC CTA TAT TTT GAA ATT CCG AGA CGG TGC
ATG AAA AAC CGA TCG AAA CGA AAC ATT GCG

5S_4M_Cy3: [Cyanine3]TCA GAA CTC CGA AGT TAA GCG TGC TTG GGC GAG
AGT AGT AC

5S_3R_Cy3: [Cyanine3]AGT ACT AGG ATG GGT GAC CCC CTG GGA AGT CCT CGT
GTT GC

5S_6L_Cy3: [Cyanine3]GCG ATC ATA CCA GCA CTA AAG CAC CGG ATC CCA TCA
GAA CTC C

(AAG)15_FAM: [6FAM]AAG AAG AAG AAG AAG AAG AAG AAG AAG AAG AAG
AAG AAG AAG AAG

(CT)23_TAMRA: [TAMRA]CTC TCT CTC TCT CTC TCT CTC TCT CTC TCT CTC
TCT CTC TCT CTC T

For hybridization, probes were prepared in 40% (v/v) formamide, 20% (w/v) dextran

sulphate, 2x SSC (sodium chloride sodium citrate), 0.03μg of salmon sperm DNA, 0.12%

SDS (sodium dodecyl sulphate) and 0.12mM EDTA (ethylenediamine-tetra acetic acid).

Chromosomes and 40-50μl of probe mixture were denatured together at 72oC for 8 mins,

cooled down slowly and allowed to hybridize overnight at 37°C. Post-hybridization washes

were at 42°C in 0.1xSSC, giving a stringency of 80-85% for the short oligo probes, and 70-

75% for the 45SrDNA probe. Chromosomes were counterstained with 4μg/ml DAPI (4´,6-

diamidino-2-phenylindole) and mounted in CitifluorAF. Slides were examined using Nikon

Eclipse 80i microscope and images were captured with a DS-QiMc monochrome camera, and

NIS-Elements v2.34 (Nikon, Tokyo, Japan). Overlays of hybridization signal and DAPI

images were viewed enhanced with Adobe Photoshop CC2018 using only cropping and

functions that treat all pixels equally. Seven FISH runs with different combinations of probes

and replicates were performed, and between 5 and 15 metaphases per slide (99 metaphases in

total from 15 slides) were analyzed in detail.

Availability of Supporting Data and Materials

Raw sequence reads (RNA-seq, Illumina HiSeq, the Oxford Nanopore and Hi-C) were

deposited in the NCBI under accession number: PRJNA736572. In specifically, the Oxford

Nanopore raw reads: SRR15039768 and SRR15039769; RNA-sequencing raw reads, as

follows: leaf: SRR15039767, root: SRR15039766; genomic Illumina short-read data:

SRR15039770; raw reads of the Hi-C library: SRR15039764 and SRR15039765.The raw

reads data were also deposited in Genome Sequence Archive (GSA) of the China National

Center for Bioinformation (accession code: CRA004283).

The assembled genome was also deposited to GenBank in NCBI under the accession

number: JAHSUZ000000000. Genome Assembly, gene and TE annotation data,

transcriptomic data are also available on the Banana Genome Hub (http://banana-genome-

hub.southgreen.fr/) for download or exploration via a dedicated Genome Browser (Jbrowse)

and syntenic browser (SynVisio).

Additional Files

Supplementary Figures

Pdf -word document containing Supplementary Figures S1-S12

Supplementary Figure S1. Genomescope analysis of heterozygosity.

Supplementary Figure S2. GO enrichment terms.

Supplementary Figure S3. RepeatExplorer clustering summary in Musaceae species.

Supplementary Figure S4. LTR retroelement trees EGL and MAC.

Supplementary Figure S5. Gypsy and Copia insertion times in *Musa.*

Supplementary Figure S6: Egcen FISH to *Musa* chromosomes.

Supplementary Figure S7. EgCen and Nanica in Assemblies of *E. glaucum* and *Musa.*

Supplementary Figure S8 Synteny of *E. glaucum* with *Musa* A and B genome.

Supplementary Figure S9. Dotplots of individual chromosomes.

Supplementary Figure S10. Inversions on chromosome 5 in three Musaceae species.

Supplementary Figure S11. Length distribution of ONT reads.

Supplementary Figure S12. Hi-C interaction contact map.

Supplementary Tables

Excel Spreadsheet containing Supplementary Tables S1-S14

Supplementary Table S1. Contig statistics based on assembly of ONT sequencing data.

Supplementary Table S2. Chromosome lengths and number of contigs anchored in *Ensete*
glaucum assembly.

Supplementary Table S3. Quality assessment of the gene annotation of *Ensete glaucum* using
BUSCOs v5.

Supplementary Table S4. Complete gene list. homology and GO.

Supplementary Table S5. Statistics for shared orthogroups (OG) and gene clustering among
E. glaucum, E. ventricosum, Musa acuminata, M. balbisiana, and *M. schizocarpa*
genomes.

Supplementary Table S6. Positively selected genes and their annotation.

Supplementary Table S7: Result of gene family size change analysis.

Supplementary Table S8 a) Top 20 GO molecular function enrichments for *E. glaucum* and
shared *E. glaucum/E. ventricosum* gene families; b) Top 20 GO biological pathways
enrichments for *E. glaucum* and shared *E. glaucum/E. ventricosum* gene families.

Supplementary Table S9. Comparison of transcriptional factor between *Ensete glaucum* and
Musa acuminata.

Supplementary Table S10. Transposable elements and other repeat proportions comparison in
assembly (RepeatMasker).

Supplementary Table S11. Repeat content (RepeatExplorer) comparison between different
Musaceae genomes.

Supplementary Table S12. Abundance of major tandemly repeated DNA repeats in Illumina
raw reads.

Supplementary Table S13. Inferred centromere positions from locations of interrupted
tandem arrays of the Egcen centromeric sequence on the chromosome assemblies.

Supplementary Table S14. Comparative survey of microsatellite sequences in *Ensete*
glaucum genome with other sister species.

Declarations

List of abbreviations

BLAST: Basic Local Alignment Search Tool; bp: base pairs; BUSCO: Benchmarking

Universal Single-Copy Orthologs; BWA: Burrows-Wheeler Aligner; FISH: fluorescence in

situ hybridization; GeMoMa: Gene Model Mapper; Gb: gigabase pairs; GC: guanine-

cytosine; GO: gene ontogeny; CTAB: cetyl trimethylammonium bromide; GWAS: Genome

Wide Association Studies; Hi-C: High-throughput chromosome conformation capture; ITS:

internal transcribed spacer of rDNA; kb: kilobase pairs; KEGG: Kyoto Encyclopedia of

Genes and Genomes; GeMoMa: Gene Model Mapper; LACHESIS: Ligating Adjacent

Chromatin Enables Scaffolding In Situ; LINE: long interspersed nucleotide elements; LTR:

long terminal repeat; Mb: megabase pairs; ML: maximum likelihood; miRNA: microRNA;

Mya: million years ago; NCBI: National Center for Biotechnology Information; NR: RefSeq

non-redundant proteins; NOR: Nucleolar Organizing Region; NTS: non transcribed spacer of

rDNA; ONT: Oxford Nanopore Technologies; PAML: Phylogenetic Analysis by Maximum

Likelihood; PacBio: Pacific Biosciences; PASA: Program to Assemble Spliced Alignments;

RAxML: Randomized Accelerated Maximum Likelihood; RNA-seq: RNA sequencing;

rDNA: ribosomal DNA; SRA: Sequence Read Archive; SSR: Simple sequence repeat; TE:

transposable element; TF: Transcription Factor; tRNA: transfer RNA; WGD: whole genome

duplication.

Consent for publication

The origin of *E.glaucum* plants is given in Materials and Method. They were collected in

China and conserved in the South China Botanical Garden, Chinese Academy of Sciences,

with appropriate agreements. No live material was exported out of the country. Other plants

for chromosome preparations was obtained from the International Transit Centre, ITC

genebank, with official Standard Material Transfer Agreement (SMTA), acknowledged in the

manuscript.

Competing Interests

892 The authors declare that they have no competing interests.

Funding

This work was supported by grants from National Science Foundation of China (32070359),

Guangdong Basic and Applied Basic Research Foundation (2021A1515012410), Overseas

Distinguished Scholar Project of SCBG (Y861041001) and Undergraduate Innovation

Training Program of Chinese Academy of Sciences (KCJH-80107-2020-004-97). M.R.

acknowledges the support of the CGIAR Research Program on Roots, Tubers and Bananas

(RTB).

Author contributions

Q.L. and J.S.H.H. designed the project and with M.R. and Z.W. contributed to project

coordination. Z.W. and Q.H. collected samples and conducted DNA and RNA extraction.

T.S. and J.S.H.H. conducted FISH experiments. Z.W. carried out genome assemblies; Z.W.,

M.R. G.D., and M.K.B. conducted gene annotation and comparative genomic analyses; Z.W.,

T.S., J.S..H.H. and F.C.B. conducted repetitive sequence analysis. All authors contributed to

writing and editing the manuscript.

Acknowledgments

This work was technically supported by the high-performance cluster of the UMR AGAP -

CIRAD of the South Green Bioinformatics Platform (http://www.southgreen.fr). Assistance

and discussion with Celia Hansen and Paulina Tomaszewska, University of Leicester, are

acknowledged. We thank Ye Yushi for the assistance with growing and maintaining the

plants. We thank the International Musa Germplasm Transit Centre

(https://www.bioversityinternational.org/banana-genebank/) for samples of *M. balbisiana.*

References

1. Wu Z, Raven PH, Hong D, Missouri Botanical Garden. Musaceae. 1. *Ensete*. *Flora of*

China. Science Press and Missouri Botanical Garden Press. Beijing and St. Louis; 297–8,

2000.

2. Borrell JS, Biswas MK, Goodwin M, Blomme G, Schwarzacher T, Heslop-Harrison JS

(Pat), et al. Enset in Ethiopia: a poorly characterized but resilient starch staple. *Ann of Bot*.

2019; doi: 10.1093/aob/mcy214.

3. Zhao T, Zwaenepoel A, Xue J-Y, Kao S-M, Li Z, Schranz ME, et al. Whole-genome

microsynteny-based phylogeny of angiosperms. *Nat Commun*. 2021; doi: 10.1038/s41467-

021-23665-0.

4. Christelová P, Valárik M, Hřibová E, De Langhe E, Doležel J. A multi gene sequence-

based phylogeny of the Musaceae (banana) family. *BMC Evol Biol*. 2011; doi: 10.1186/1471-

2148-11-103.

5. Janssens SB, Vandelook F, De Langhe E, Verstraete B, Smets E, Vandenhouwe I, et al.

Evolutionary dynamics and biogeography of Musaceae reveal a correlation between the

diversification of the banana family and the geological and climatic history of Southeast Asia.

New Phytol. 2016; doi: 10.1111/nph.13856.

6. Cheesman EE. Classification of the Bananas: The Genus *Ensete* Horan. *Kew Bull*. 1947;

doi: 10.2307/4109206.

7. Simmonds NW. Notes on Banana Taxonomy. *Kew Bull*. 1960; doi: 10.2307/4114778.

8. Li HW. Musaceae of Yunnan. *Acta phytotaxonomica sinica*. 1978;

9. Ochiai Y. From forests to homegardens: A case study of *Ensete glaucum* in Myanmar and

Laos. *Tropics*. 2012; doi: 10.3759/tropics.21.59.

10. Song J-J, Liao J-P, Tang Y-J, Chen Z-Y. Chromosome numbers in Orchidantha

(Lowiaceae) and their biogeographic and systematic implications. *Annales Botanici Fennici*.

Finnish Zoological and Botanical Publishing Board; 41:429–332004;

11. Majumdar K, Sarkar A, Deb D, Majumder J, Datta B. Distribution record of *Ensete*

glaucum (Roxb.) Cheesm.(Musaceae) in Tripura, Northeast India: a rare wild primitive

banana. *Asian J Conserv Biol.* 2:164–72013;

12. Yang Q-S, Gao J, He W-D, Dou T-X, Ding L-J, Wu J-H, et al. Comparative

transcriptomics analysis reveals difference of key gene expression between banana and

plantain in response to cold stress. *BMC Genomics*. 2015; doi: 10.1186/s12864-015-1551-z.

13. Belser C, Istace B, Denis E, Dubarry M, Baurens F-C, Falentin C, et al. Chromosome-

scale assemblies of plant genomes using nanopore long reads and optical maps. *Nature*

Plants. 2018; doi: 10.1038/s41477-018-0289-4.

14. D'Hont A, Denoeud F, Aury J-M, Baurens F-C, Carreel F, Garsmeur O, et al. The banana

(*Musa acuminata*) genome and the evolution of monocotyledonous plants. *Nature*.

488:2132012;

15. Wang Z, Miao H, Liu J, Xu B, Yao X, Xu C, et al. *Musa balbisiana* genome reveals

subgenome evolution and functional divergence. *Nature Plants*. 2019; doi: 10.1038/s41477-

019-0452-6.

16. Yemataw Z, Muzemil S, Ambachew D, Tripathi L, Tesfaye K, Chala A, et al. Genome

sequence data from 17 accessions of *Ensete ventricosum*, a staple food crop for millions in

Ethiopia. *Data in Brief*. 2018; doi: 10.1016/j.dib.2018.03.026.

17. Harrison J, Moore KA, Paszkiewicz K, Jones T, Grant MR, Ambacheew D, et al. A Draft

Genome Sequence for *Ensete ventricosum*, the Drought-Tolerant "Tree Against Hunger."

Agronomy.   Multidisciplinary   Digital   Publishing   Institute;   2014;   doi:

10.3390/agronomy4010013.

18. Simão FA, Waterhouse RM, Ioannidis P, Kriventseva EV, Zdobnov EM. BUSCO:

assessing genome assembly and annotation completeness with single-copy orthologs.

Bioinformatics. 2015; doi: 10.1093/bioinformatics/btv351.

19. Martin G, Baurens F-C, Droc G, Rouard M, Cenci A, Kilian A, et al. Improvement of the

banana "*Musa acuminata*" reference sequence using NGS data and semi-automated

bioinformatics methods. *BMC Genomics*. 2016; doi: 10.1186/s12864-016-2579-4.

20. Belser C, Baurens F-C, Noel B, Martin G, Cruaud C, Istace B, et al. Telomere-to-

telomere gapless chromosomes of banana using nanopore sequencing. *Commun Biol*. 2021;

doi: 10.1038/s42003-021-02559-3.

21. Bartos J, Alkhimova O, Dolezelová M, De Langhe E, Dolezel J. Nuclear genome size and

genomic distribution of ribosomal DNA in *Musa* and *Ensete* (Musaceae): taxonomic

implications. *Cytogenet Genome Res*. 2005; doi: 10.1159/000082381.

22. Wang R, Yang Y, Jing Y, Segar ST, Zhang Y, Wang G, et al. Molecular mechanisms of

mutualistic and antagonistic interactions in a plant-pollinator association. *Nat Ecol Evol*.

2021; doi: 10.1038/s41559-021-01469-1.

23. González AV, Gómez-Silva V, Ramírez MJ, Fontúrbel FE. Meta-analysis of the

differential effects of habitat fragmentation and degradation on plant genetic diversity.

Conserv Biol. 2020; doi: 10.1111/cobi.13422.

24. Liu A-Z, Kress WJ, Wang HF, Li D. Insect pollination of *Musella* (Musaceae), a

monotypic genus endemic to Yunnan, China. *Plant Syst and Evol*. 235:135–462002;

25. Sardos J, Breton C, Perrier X, Houwe IVD, Paofa J, Rouard M, et al. Wild to

domesticates: genomes of edible diploid bananas hold traces of several undefined genepools.

bioRxiv. 2021; doi: 10.1101/2021.01.29.428762.

26. Martin G, Cardi C, Sarah G, Ricci S, Jenny C, Fondi E, et al. Genome ancestry mosaics

reveal multiple and cryptic contributors to cultivated banana. *The Plant J*. 2020; doi:

10.1111/tpj.14683.

27. Maughan PJ, Lee R, Walstead R, Vickerstaff RJ, Fogarty MC, Brouwer CR, et al.

Genomic insights from the first chromosome-scale assemblies of oat (*Avena* spp.) diploid

species. *BMC Biol*. 2019; doi: 10.1186/s12915-019-0712-y.

28. Marrano A, Britton M, Zaini PA, Zimin AV, Workman RE, Puiu D, et al. High-quality

chromosome-scale assembly of the walnut (*Juglans regia* L.) reference genome. *Gigascience*.

2020; doi: 10.1093/gigascience/giaa050.

29. Yang X, Kang M, Yang Y, Xiong H, Wang M, Zhang Z, et al. A chromosome-level

genome assembly of the Chinese tupelo *Nyssa sinensis*. *Sci Data*. 2019; doi: 10.1038/s41597-

019-0296-y.

30. Voillemot M, Pannell JR. Inbreeding depression is high in a self-incompatible perennial

herb population but absent in a self-compatible population showing mixed mating. *Ecol Evol*.

2017; doi: 10.1002/ece3.3354.

31. Sun G, Xu Y, Liu H, Sun T, Zhang J, Hettenhausen C, et al. Large-scale gene losses

underlie the genome evolution of parasitic plant *Cuscuta australis*. *Nat Commun*. 2018; doi:

10.1038/s41467-018-04721-8.

32. Redwan RM, Saidin A, Kumar SV. The draft genome of MD-2 pineapple using hybrid

error correction of long reads. *DNA Res*. 2016; doi: 10.1093/dnares/dsw026.

33. Franco-Zorrilla JM, López-Vidriero I, Carrasco JL, Godoy M, Vera P, Solano R. DNA-

binding specificities of plant transcription factors and their potential to define target genes.

Proc Natl Acad Sci U S A. 2014; doi: 10.1073/pnas.1316278111.

34. Cenci A, Guignon V, Roux N, Rouard M. Genomic analysis of NAC transcription factors

in banana (*Musa acuminata*) and definition of NAC orthologous groups for monocots and

dicots. *Plant Mol Biol*. 2014; doi: 10.1007/s11103-013-0169-2.

35. Xiao Y-Y, Kuang J-F, Qi X-N, Ye Y-J, Wu Z-X, Chen J-Y, et al. A comprehensive

investigation of starch degradation process and identification of a transcriptional activator

MabHLH6 during banana fruit ripening. *Plant Biotechnol J*. 2018; doi: 10.1111/pbi.12756.

36. Lerat E. Identifying repeats and transposable elements in sequenced genomes: how to

find your way through the dense forest of programs. *Heredity (Edinb)*. 2010; doi:

10.1038/hdy.2009.165.

37. Novák P, Neumann P, Macas J. Global analysis of repetitive DNA from unassembled

sequence reads using RepeatExplorer2. *Nat Protoc*. 2020; doi: 10.1038/s41596-020-0400-y.

38. Wu W, Yang Y-L, He W-M, Rouard M, Li W-M, Xu M, et al. Whole genome sequencing

of a banana wild relative *Musa itinerans* provides insights into lineage-specific

diversification of the Musa genus. *Sci Rep*. 2016; doi: 10.1038/srep31586.

39. Biscotti MA, Olmo E, Heslop-Harrison JSP. Repetitive DNA in eukaryotic genomes.

Chromosome Res. 2015; doi: 10.1007/s10577-015-9499-z.

40. Heslop-Harrison JSP, Schwarzacher T. Organisation of the plant genome in

chromosomes. *Plant J*. 2011; doi: 10.1111/j.1365-313X.2011.04544.x.

41. Čížková J, Hřibová E, Humplíková L, Christelová P, Suchánková P, Doležel J. Molecular

Analysis and Genomic Organization of Major DNA Satellites in Banana (*Musa* spp.). *PLoS*

ONE. 2013; doi: 10.1371/journal.pone.0054808.

42. Suntronpong A, Kugou K, Masumoto H, Srikulnath K, Ohshima K, Hirai H, et al. CENP-

B box, a nucleotide motif involved in centromere formation, occurs in a New World monkey.

Biol Lett. 2016; doi: 10.1098/rsbl.2015.0817.

43. Aragón-Alcaide L, Miller T, Schwarzacher T, Reader S, Moore G. A cereal centromeric

sequence. *Chromosoma*. 1996; doi: 10.1007/BF02524643.

44. Heslop-Harrison JS, Murata M, Ogura Y, Schwarzacher T, Motoyoshi F. Polymorphisms

and genomic organization of repetitive DNA from centromeric regions of *Arabidopsis*

chromosomes. *Plant Cell*. 1999; doi: 10.1105/tpc.11.1.31.

45. Lermontova I, Sandmann M, Mascher M, Schmit A-C, Chabouté M-E. Centromeric

chromatin and its dynamics in plants. *Plant J*. 2015; doi: 10.1111/tpj.12875.

46. Biswas MK, Natarajan S, Biswas D, Nath UK, Park J-I, Nou I-S. LSAT: Liliaceae Simple

Sequences Analysis Tool, a web server. *Bioinformation*. 2018; doi:

10.6026/97320630014181.

47. Osuji JO, Crouch J, Harrison G, Heslop-Harrison JS. Molecular Cytogenetics of *Musa*

Species, Cultivars and Hybrids: Location of 18S-5.8S-25S and 5S rDNA and Telomere-like

Sequences. *Ann Bot*. 1998; doi: 10.1006/anbo.1998.0674.

48. Baurens F-C, Noyer J-L, Lanaud C, Lagoda PJL. Assessment of a species-specific

element (Brep 1) in banana. *Theor Appl Genet*. 1997; doi: 10.1007/s001220050643.

49. Garcia S, Wendel JF, Borowska-Zuchowska N, Aïnouche M, Kuderova A, Kovarik A.

The Utility of Graph Clustering of 5S Ribosomal DNA Homoeologs in Plant Allopolyploids,

Homoploid Hybrids, and Cryptic Introgressants. *Front Plant Sci*. 2020; doi:

10.3389/fpls.2020.00041.

50. Castilho A, Heslop-Harrison JS. Physical mapping of 5S and 18S-25S rDNA and

repetitive DNA sequences in *Aegilops umbellulata*. *Genome*. 1995; doi: 10.1139/g95-011.

51. Dubcovsky J, Dvorák J. Ribosomal RNA multigene loci: nomads of the Triticeae

genomes. *Genetics*. 1995; doi: 10.1093/genetics/140.4.1367.

52. Bandi V, Gutwin C. Interactive Exploration of Genomic Conservation. *46th Graphics*

Interface Conference on Proceedings of Graphics Interface 2020. Waterloo, Canada:

Canadian Human-Computer Communications Society.

53. Li W, Challa GS, Zhu H, Wei W. Recurrence of Chromosome Rearrangements and Reuse

of DNA Breakpoints in the Evolution of the Triticeae Genomes. *G3*. 2016; doi:

10.1534/g3.116.035089.

54. GBIF.org: GBIF. https://doi.org/10.15468/dl.f9meez (2021). Accessed 2021 Apr 25.

55. POWO: Plants of the World Online | Kew Science. Plants of the World Online.

http://www.plantsoftheworldonline.org/ Accessed 2021 Sep 3.

56. Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence

data. *Bioinformatics*. 2014; doi: 10.1093/bioinformatics/btu170.

57. Andrews S. Babraham Bioinformatics—FastQC A Quality Control Tool for High

Throughput Sequence Data. https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

58. Chen S, Zhou Y, Chen Y, Gu J. fastp: an ultra-fast all-in-one FASTQ preprocessor.

Bioinformatics. 2018; doi: 10.1093/bioinformatics/bty560.

59. Belton J-M, McCord RP, Gibcus JH, Naumova N, Zhan Y, Dekker J. Hi-C: a

comprehensive technique to capture the conformation of genomes. *Methods*. 2012; doi:

10.1016/j.ymeth.2012.05.001.

60. Belaghzal H, Dekker J, Gibcus JH. Hi-C 2.0: An optimized Hi-C procedure for high-

resolution genome-wide mapping of chromosome conformation. *Methods*. 2017; doi:

10.1016/j.ymeth.2017.04.004.

61. Liu H, Wu S, Li A, Ruan J, Wu S, Li A, et al. SMARTdenovo: a de novo assembler using

long noisy reads. *Gigabyte*. 2021; doi: 10.46471/gigabyte.15.

62. Vaser R, Sović I, Nagarajan N, Šikić M. Fast and accurate de novo genome assembly

from long uncorrected reads. *Genome Res*. 2017; doi: 10.1101/gr.214270.116.

63. Li H, Durbin R. Fast and accurate long-read alignment with Burrows-Wheeler transform.

Bioinformatics. 2010; doi: 10.1093/bioinformatics/btp698.

64. Hu J, Fan J, Sun Z, Liu S. NextPolish: a fast and efficient genome polishing tool for long-

read assembly. *Bioinformatics*. 2020; doi: 10.1093/bioinformatics/btz891.

65. Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: an

integrated tool for comprehensive microbial variant detection and genome assembly

improvement. *PLoS One*. 2014; doi: 10.1371/journal.pone.0112963.

66. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. *Nat Methods*.

2012; doi: 10.1038/nmeth.1923.

67. Servant N, Varoquaux N, Lajoie BR, Viara E, Chen C-J, Vert J-P, et al. HiC-Pro: an

optimized and flexible pipeline for Hi-C data processing. *Genome Biol*. 2015; doi:

10.1186/s13059-015-0831-x.

68. Burton JN, Adey A, Patwardhan RP, Qiu R, Kitzman JO, Shendure J. Chromosome-scale

scaffolding of de novo genome assemblies based on chromatin interactions. *Nat Biotechnol*.

2013; doi: 10.1038/nbt.2727.

69. Grabherr MG, Haas BJ, Yassour M, Levin JZ, Thompson DA, Amit I, et al. Trinity:

reconstructing a full-length transcriptome without a genome from RNA-Seq data. *Nat*

Biotechnol. 2011; doi: 10.1038/nbt.1883.

70. Haas BJ, Delcher AL, Mount SM, Wortman JR, Smith RK, Hannick LI, et al. Improving

the Arabidopsis genome annotation using maximal transcript alignment assemblies. *Nucleic*

Acids Res. 2003; doi: 10.1093/nar/gkg770.

71. Marçais G, Kingsford C. A fast, lock-free approach for efficient parallel counting of

occurrences of k-mers. *Bioinformatics*. 2011; doi: 10.1093/bioinformatics/btr011.

72. Sun H, Ding J, Piednoël M, Schneeberger K. findGSE: estimating genome size variation

within human and Arabidopsis using k-mer frequencies. *Bioinformatics*. 2018; doi:

10.1093/bioinformatics/btx637.

73. Ranallo-Benavidez TR, Jaron KS, Schatz MC. GenomeScope 2.0 and Smudgeplot for

reference-free profiling of polyploid genomes. *Nat Commun*. 2020; doi: 10.1038/s41467-020-

14998-3.

74. Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR: ultrafast

universal RNA-seq aligner. *Bioinformatics*. 2013; doi: 10.1093/bioinformatics/bts635.

75. Brůna T, Hoff KJ, Lomsadze A, Stanke M, Borodovsky M. BRAKER2: Automatic

Eukaryotic Genome Annotation with GeneMark-EP+ and AUGUSTUS Supported by a

Protein Database. *bioRxiv*. 2020; doi: 10.1101/2020.08.10.245134.

76. Lomsadze A, Burns PD, Borodovsky M. Integration of mapped RNA-Seq reads into

automatic training of eukaryotic gene finding algorithm. *Nucleic Acids Res*. 2014; doi:

10.1093/nar/gku557.

77. Stanke M, Diekhans M, Baertsch R, Haussler D. Using native and syntenically mapped

cDNA alignments to improve de novo gene finding. *Bioinformatics*. 2008; doi:

10.1093/bioinformatics/btn013.

78. Campbell MS, Law M, Holt C, Stein JC, Moghe GD, Hufnagel DE, et al. MAKER-P: a

tool kit for the rapid creation, management, and quality control of plant genome annotations.

Plant Physiol. 2014; doi: 10.1104/pp.113.230144.

79. Keilwagen J, Hartung F, Grau J. GeMoMa: Homology-Based Gene Prediction Utilizing

Intron Position Conservation and RNA-seq Data. *Methods Mol Biol*. 2019; doi: 10.1007/978-

1-4939-9173-0_9.

80. Haas BJ, Salzberg SL, Zhu W, Pertea M, Allen JE, Orvis J, et al. Automated eukaryotic

gene structure annotation using EVidenceModeler and the Program to Assemble Spliced

Alignments. *Genome Biol*. 2008; doi: 10.1186/gb-2008-9-1-r7.

81. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:

architecture and applications. *BMC Bioinformatics*. 2009; doi: 10.1186/1471-2105-10-421.

82. Magrane M, Consortium U. UniProt Knowledgebase: a hub of integrated protein data.

Database (Oxford). 2011; doi: 10.1093/database/bar009.

83. Zdobnov EM, Apweiler R. InterProScan - an integration platform for the signature-

recognition methods in InterPro. *Bioinformatics*. 2001; doi: 10.1093/bioinformatics/17.9.847.

84. Conesa A, Götz S, García-Gómez JM, Terol J, Talón M, Robles M. Blast2GO: a

universal tool for annotation, visualization and analysis in functional genomics research.

Bioinformatics. 2005; doi: 10.1093/bioinformatics/bti610.

85. Droc G: ensete_annotation. https://github.com/gdroc/ensete_annotation (2021). Accessed

2021 Oct 25.

86. Droc G, Lariviere D, Guignon V, Yahiaoui N, This D, Garsmeur O, et al. The Banana

Genome Hub. *Database*. 2013; doi: 10.1093/database/bat035.

87. Emms DM, Kelly S. OrthoFinder: solving fundamental biases in whole genome

comparisons dramatically improves orthogroup inference accuracy. *Genome Biol*. 2015; doi:

10.1186/s13059-015-0721-2.

88. Buchfink B, Xie C, Huson DH. Fast and sensitive protein alignment using DIAMOND.

Nat Methods. 2015; doi: 10.1038/nmeth.3176.

89. Lex A, Gehlenborg N, Strobelt H, Vuillemot R, Pfister H. UpSet: Visualization of

Intersecting Sets. *IEEE Trans Vis Comput Graph*. 2014; doi: 10.1109/TVCG.2014.2346248.

90. Alexa A, Rahnenfuhrer J. topGO: enrichment analysis for gene ontology. R package

version 2.24.0. 2010.

91. Hazzouri KM, Gros-Balthazard M, Flowers JM, Copetti D, Lemansour A, Lebrun M, et

al. Genome-wide association mapping of date palm fruit traits. *Nat Commun*. 2019; doi:

10.1038/s41467-019-12604-9.

92. Ouyang S, Zhu W, Hamilton J, Lin H, Campbell M, Childs K, et al. The TIGR Rice

Genome Annotation Resource: improvements and new features. *Nucleic Acids Res*. 2007;

doi: 10.1093/nar/gkl976.

93. Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, et al. Phytozome: a

comparative platform for green plant genomics. *Nucleic Acids Res*. 2012; doi:

10.1093/nar/gkr944.

94. Han MV, Thomas GWC, Lugo-Martinez J, Hahn MW. Estimating gene gain and loss

rates in the presence of error in genome assembly and annotation using CAFE 3. *Mol Biol*

Evol. 2013; doi: 10.1093/molbev/mst100.

95. Zheng Y, Jiao C, Sun H, Rosli HG, Pombo MA, Zhang P, et al. iTAK: A Program for

Genome-wide Prediction and Classification of Plant Transcription Factors, Transcriptional

Regulators, and Protein Kinases. *Mol Plant*. 2016; doi: 10.1016/j.molp.2016.09.014.

96. Sun P, Jiao B, Yang Y, Shan L, Li T, Li X, et al. WGDI: A user-friendly toolkit for

evolutionary analyses of whole-genome duplications and ancestral karyotypes. 2021; doi:

10.1101/2021.04.29.441969.

97. Li H. Minimap2: pairwise alignment for nucleotide sequences. *Bioinformatics*. 2018; doi:

10.1093/bioinformatics/bty191.

98. Cabanettes F, Klopp C. D-GENIES: dot plot large genomes in an interactive, efficient and

simple way. *PeerJ*. 2018; doi: 10.7717/peerj.4958.

99. Wang Y, Tang H, DeBarry JD, Tan X, Li J, Wang X, et al. MCScanX: a toolkit for

detection and evolutionary analysis of gene synteny and collinearity. *Nucleic Acids Res*.

2012; doi: 10.1093/nar/gkr1293.

100. Amselem J, Cornut G, Choisne N, Alaux M, Alfama-Depauw F, Jamilloux V, et al.

RepetDB: a unified resource for transposable element references. *Mob DNA*. 2019; doi:

10.1186/s13100-019-0150-y.

101.   Belser   C: Pahang-associated-data.   GitHub.   https://github.com/institut-de-
genomique/Pahang-associated-data (2021). Accessed 2021 Oct 25.

102. Ou S, Su W, Liao Y, Chougule K, Agda JRA, Hellinga AJ, et al. Benchmarking
transposable element annotation methods for creation of a streamlined, comprehensive
pipeline. *Genome Biol*. 2019; doi: 10.1186/s13059-019-1905-y.

103. Ellinghaus D, Kurtz S, Willhoeft U. LTRharvest, an efficient and flexible software for
de novo detection of LTR retrotransposons. *BMC Bioinformatics*. 2008; doi: 10.1186/1471-
2105-9-18.

104. Ou S, Jiang N. LTR_retriever: A Highly Accurate and Sensitive Program for
Identification of Long Terminal Repeat Retrotransposons. *Plant Physiol*. 2018; doi:
10.1104/pp.17.01310.

105. Xu Z, Wang H. LTR_FINDER: an efficient tool for the prediction of full-length LTR
retrotransposons. *Nucleic Acids Res*. 2007; doi: 10.1093/nar/gkm286.

106. Su W, Gu X, Peterson T. TIR-Learner, a New Ensemble Method for TIR Transposable
Element Annotation, Provides Evidence for Abundant New Transposable Elements in the
Maize Genome. *Mol Plant*. 2019; doi: 10.1016/j.molp.2019.02.008.

107. Shi J, Liang C. Generic Repeat Finder: A High-Sensitivity Tool for Genome-Wide De
Novo Repeat Detection. *Plant Physiol*. 2019; doi: 10.1104/pp.19.00386.

108. Xiong W, He L, Lai J, Dooner HK, Du C. HelitronScanner uncovers a large overlooked
cache of Helitron transposons in many plant genomes. *Proc Natl Acad Sci U S A*. 2014; doi:
10.1073/pnas.1410068111.

109. Flynn JM, Hubley R, Goubert C, Rosen J, Clark AG, Feschotte C, et al. RepeatModeler2

for automated genomic discovery of transposable element families. *Proc Natl Acad Sci U S*

A. 2020; doi: 10.1073/pnas.1921046117.

110. Fu L, Niu B, Zhu Z, Wu S, Li W. CD-HIT: accelerated for clustering the next-

generation sequencing data. *Bioinformatics*. 2012; doi: 10.1093/bioinformatics/bts565.

111. Zhang R-G, Wang Z-X, Ou S, Li G-Y. TEsorter: lineage-level classification of

transposable elements using conserved protein domains. Accessed 2019 Oct 15.

112. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high

throughput. *Nucleic Acids Res*. 2004; doi: 10.1093/nar/gkh340.

113. Ma J, Bennetzen JL. Rapid recent growth and divergence of rice nuclear genomes. *Proc*

Natl Acad Sci U S A. 2004; doi: 10.1073/pnas.0403715101.

114. Wang Z: LTR-insertion-time-estimation. GitHub. https://github.com/wangziwei08/LTR-

insertion-time-estimation (2021). Accessed 2021 Oct 25.

115. Schwarzacher T, Heslop-Harrison JS. Practical in situ Hybridization. Oxford, UK: BIOS

Scientific Publishers Ltd; 2000

116. Ruas M, Guignon V, Sempere G, Sardos J, Hueber Y, Duvergey H, et al. MGIS:

managing banana (*Musa* spp.) genetic resources information and high-throughput genotyping

data. *Database*. 2017; doi: 10.1093/database/bax046.

1218

Figure legends

Figure 1. *Ensete glaucum* plant morphology and distribution map.

(A) *E. glaucum* in South China Botanical Garden, Chinese Academy of Sciences. The pseudostem of this plant is about 3.5m tall. (B) Inflorescence with male and female flowers showing bracts and flowers alternately arranged along the main axis. (C) Staminate flowers, and visiting black shield wasp (*Vespa bicolor,* Vespidae, Hymenoptera) (D) Female flowers. (E) Fruits. Bars in B-E represent 2 cm. (F) Native distribution of *Ensete glaucum, E. superbum, Musa* and *Ensete* species by countries or provinces (for China, India-Assam, and Australia). Musaceae are not currently native in the Americas, although *Ensete* is present in the fossil record (Manchester et al., 1993). *E. glaucum* always occurs in the same provinces as *Musa* and sometimes with other Asian *Ensete* species. Map adapted from POWO [52].

Figure 2. *Ensete glaucum* chromosome assembly and genome features.

Circos plot of (a) The nine pseudomolecules of the EGL assembly corresponding to 9 chromosomes. Black dots indicate centromere positions and 5S and 45S the positions of rDNA loci; scale in Mbp; (b) Gene density; (c) Repeat density; (d) *Copia* LTR retroelement density; (e) *Gypsy* LTR retroelement density; (f) DNA transposon density; (g) Short tandem repeat (microsatellite) density; (h) Syntenic genomic blocks, linked by curved lines (arbitrary color) in middle of the plot.

Figure 3. Gene family evolution and conservation.

(A) Gene family expansion and contraction with a phylogenetic tree showing timeline of divergence of monocot species. MRCA: Most Recent Common Ancestor. Numbers denote the gene family expansion (orange) and contraction (green). (B) The synonymous substitutions (Ks) frequency density distributions of orthologs within EGL or MAC, whose

peaks indicate whole-genome duplications (WGDs). (C) Intersection diagram showing the

distribution of shared orthogroups (OGs) (at least two sequences per OG) among *Musa* and

Ensete genomes. codes: E, *E. glaucum*; V, *E. ventricosum;* A, *M. acuminata*; B, *M.*

balbisiana; S, *M. schizocarpa.* (D) Histogram of the comparative abundance (number of

genes) of transcriptions factors between *M. acuminata* and *E. glaucum.*

Figure 4. Comparative analysis of repetitive DNA in Musaceae using RepeatExplorer.

(A) Bar plot showing the sizes (numbers of reads) of the most abundant individual graph-

based read clusters with (lower part) distribution among six Musaceae species (coloured

rectangle size is proportional to the number of reads in a cluster for each species. Clusters and

species were sorted by using hierarchical clustering. Individual rectangles are colored based

on the annotation of the clusters. Figure was adapted from the output of the visualization

script (plot_comparative_clustering_summary.R). Species codes: EGL, *Ensete glaucum*;

EVE, *E. ventricosum*; MAC, *Musa acuminata*; MBA, *M. balbisiana*; MLA, *Musella*

lasiocarpa; MSC, *M. schizocarpa.* (B) The nature and ages of LTR retroelements in *E.*

glaucum. (B) The ages of retroelement insertions in *E. glaucum, M. acuminata* and *M.*

balbisiana. Mya = Million years ago.

Figure 5. *Ensete glaucum* centromeric repeat structure.

(A) Dot plot (self-comparison of sequences) showing start of a 134bp Egcen tandem array.

(B) Dot plot showing part of a chromosomes assembly (eg04) plotted against part of a single

ONT read with blocks of the Egcen tandem repeat (appearing as dense rectangles at this

scale) interspersed with Nanica elements (red; five homologous copies in both orientations)

and LTR retroelements (green; two non-homologous sub-families), (C) Bar chart showing

frequency distribution of the EGcen centromeric tandem repeat, Nanica transposable

elements (x10 on axis), and locations of 45S and 5S rDNA along the assemblies for each

chromosome. Long Egcen arrays occur at one of more sites at the centromeric regions of all

chromosomes. (D, E, F) *In situ* hybridization of Egcen probes detected by red fluorescence to

cyan-fluorescing DAPI-stained chromosomes of (D) EGL, *Ensete glaucum*; (E) EVE, *E.*

ventricosum; and (F) MLA, *Musella lasiocarpa*. The red Egcen locations collocate with the

primary centromeric constriction on all 9 pairs of chromosomes. Bar=5µm.

Figure 6. Microsatellite (SSR) distribution in *Ensete glaucum*.

(A) Abundance (count) and total number of monomers microsatellites (SSR) with motifs

between 1 and 6 bp long. (B, C) *In situ* hybridization of synthetic microsatellite probes to

DAPI-stained (blue) chromosomes, showing (B) AAG is relatively uniformly distributed

along chromosomes compared to (C) where the greater abundance of AG/CT in distal

chromosome regions is seen. (D) Abundance of AAG, AG, all microsatellites, and genes

along the chromosome assemblies. In agreement with the *in situ* hybridization result, AAG is

more uniformly distributed, while AG (along with genes and all the microsatellites pooled)

show greater abundance in distal chromosome regions except for the arm of chromosome

eg06 carrying the 45S rDNA (NOR). Bar=5µm.

Figure 7. rDNA in *Ensete glaucum*

(A-C) *In situ* hybridization to chromosomes (stained blue with DAPI) showing locations of

(A) the 45S rDNA (green) on one pair of chromosomes while Egcen (red) locates all 9 pairs

of chromosomes at the centromeres. (B) Chromosomes showing location of 45S (green on

eg06; the two sites are on a pair of chromosomes which are adjacent to each other) and 5S

(red) rDNA loci. (C) the 5S rDNA (red) in a more dispersed pattern on one pair of

chromosomes with Egcen (green) at all 9 pairs of centromeres; inset shows 5S rDNA

chromosome pair at higher magnification. The 5S rDNA signal is dispersed over a longer

region of a chromosome, while the 45S locus occupies much of the chromosome arm. Bar

5µm. X=stain precipitate. (D) Histogram showing density of genes (light green), Egcen (blue)

and 45S rDNA copies on chromosome eg06. The arm carrying the 45S rDNA is depleted in

genes. (E) Part of a single ONT read covering 24kb spanning part of the 5S rDNA array. The

unusually long 1056bp tandemly repeated 5S rDNA monomers (14 copies) are interrupted by

an LTR retroelement. LTRs, with no homology to the 5S rDNA, are seen (bottom left and top

right) in the red box.

Figure 8. Synteny of *Ensete glaucum* and *Musa acuminata*

(A) Synteny plot (Synvisio) connecting syntenic genes in the 9 chromosomes of *E. glaucum*

(egxx) and 11 chromosomes of *M. acuminata* (maxx). syntenic blocks of high homology are

indicated by uniformly coloured areas in the graphs. Only eg05 and ma05 maintain synteny

over the full chromosome length, although there are some rearrangements. Three ma

chromosomes are represented by part of one eg chromosome, while other ma chromosomes

are fusions of more than one eg chromosome. (B) Dot plot comparing DNA sequences of *E.*

glaucum and *M. acuminata*. (C) representation of the syntenic blocks in the karyotypes of *E.*

glaucum and *M. acuminata*. Chromosome rearrangements are shown, complementing the

Synteny plot, while inversions and relative expansions and contractions of genome regions

are clear.

Figure 1: *Ensete glaucum* plant morphology and distribution map

Figure 2 : *Ensete glaucum* chromosome assembly and genome features

a) Nine pseudomolecules of EGL assembly; black circles indicate centromere positions
b) Genes
c) Repeats
d) *Copia* LTR retroelements
e) *Gypsy* LTR retroelements
f) DNA transposons
g) Microsatellites
h) Syntenic genomic blocks

Figure 3: Gene families

A

E. glaucum
M. acuminata

Median of Ks

C Gene family **expansions** and **contractions**

O. sativa (+1498/-2793)

MRCA
(12384)

+234
-137

M. balbisiana (+550/-2670)

M. acuminata (+872/-400)

+394
-177

+2184
-871

M. schizocarpa (+573/-1139)

+0
-11

E. glaucum (+1229/-837)

P. dactyliferava (+2009/-3444)

80   60   40   20   0   Mya

B

Number of OGs

14523

Core and
pseudo core

Dispensable

Species
specific

Musa
specific

Ensete
specific

93   1597   418   1146   1806   16   92   235   75   601   263   410   135   368   346   108   189   37   525   260   1471   281   1429   121   224   219   599   162   1866   260

BS   GS   GS   BV   SV   AB   AS   GA   AV

M. balbisiana (B)
M. schizocarpa (S)
E. glaucum (G)
E. ventricosum (V)
M. acuminata (A)

D

Count

Transcription Factor families

Ensete glaucum   *Musa acuminata*

AP2/ERF-AP2, AP2/ERF-ERF, AP2/ERF-RAV, B3, B3-ARF, BBR-BPC, BES1, bHLH, bZIP, CO-like, C2C2-Dof, GATA, C2C2-LSD, YABBY, C2H2, C3H, CAMTA, CPP, DBB, E2F-DP, EIL, FAR1, GARP-ARR-B, G2-like, GeBP, GRAS, GRF, HB-HD-ZIP, HB-KNOX, HB-other, HB-PHD, HB-WOX, HRT, HSF, LFY, MADS-MIKC, MADS-M-type, MYB, MYB-related, NAC, NF-X1, NF-YA, NF-YB, NF-YC, S1Fa-like, SAP, SBP, SRS, STAT, TCP, Trihelix, VOZ, Whirly, WRKY, zf-HD

Figure 4: Repeat classes and retroelement insertion times

Figure 5: Centromere structure

A

B Chromosome eg04 (nt 6,890,000- 6,990,400)

Single Nanopore read (94kb)
9c7e99b5-17a8-4bba-8955-81006158d07b

C Chromosome maps

eg01 (67.48Mb)

eg02 (55.95Mb)

eg03 (55.03Mb)

eg04 (44.62Mb)

5S

eg05 (50.34Mb)

45S

eg06 (42.46Mb)

eg07 (55.60Mb)

eg08 (55.56Mb)

eg09 (54.47Mb)

Egcen
Nanica

D EGL

E EVE

F MLA

Figure 6: Microsatellites

Figure 7: rDNA

A eg06 — 45S FITC / Cen Cy3 — X

B eg05, eg06, eg06, eg05 — 45S green / 5S Cy3

C eg05, eg05 — Cen FAM / 5S Cy3

D Chromosome eg06

Legend:
- Egcen
- 45S rDNA
- Genes

x-axis: Mb (1, 4, 7, 10, 13, 16, 19, 22, 25, 28, 31, 34, 37, 40, 43)
y-axis: 0, 50, 100, 150

E

ONT read (1,000–24,000nt)

070d5196-0757-44da-8860-ee90B5F28bb9

ONT read (3,000–14,500nt)

5S rDNA monomers

5S rDNA monomers

LTR

retroelement

LTR

5S rDNA monomers

5S rDNA monomers

Figure 8: Synteny of *Ensete glaucum* (eg) chromosomes and *Musa acuminata* Pahang V4 (ma)