SmdA is a novel cell morphology determinant in

Staphylococcus aureus

Ine Storaker Myrbråten[1], Gro Anita Stamsås[1], Helena Chan[2,3], Danae Morales Angeles[1], Tiril Mathiesen Knutsen[1], Zhian Salehian[1], Volha Shapaval[4], Daniel Straume[1] and Morten Kjos[1].

[1] Faculty of Chemistry, Biotechnology and Food Science, Norwegian University of Life Sciences, Ås, Norway.

[2] Structural Cellular Biology Unit, Okinawa Institute of Science and Technology, Okinawa, Japan.

[3] The ithree institute, University of Technology Sydney, Broadway, NSW, Australia.

[4] Faculty of Science and Technology, Norwegian University of Life Sciences, Ås, Norway.

Abstract

Cell division and cell wall synthesis in staphylococci need to be precisely coordinated and controlled to allow the cell to multiply while maintaining their nearly spherical shape. The mechanisms ensuring correct placement of the division plane and synthesis of new cell wall have been studied intensively, however, hitherto unknown factors and proteins are likely to play key roles in this complex interplay. We here identified and investigated a protein with major influence on cell morphology in *Staphylococcus aureus*. The protein, named SmdA (for staphylococcal morphology determinant A), is a membrane-protein with septum-enriched localization. By CRISPRi knockdown and overexpression combined with different microscopy techniques, we demonstrate that proper levels of SmdA is necessary for cell division, including septum formation and cell splitting. We also identified conserved residues in SmdA that are critical for its functionality. Pulldown- and bacterial two-hybrid interaction experiments showed that SmdA interacts with several known cell division- and cell wall synthesis proteins, including penicillin binding proteins (PBPs) and EzrA. Notably, SmdA also affects susceptibility to cell wall targeting antibiotics, particularly in methicillin-resistant *S. aureus* (MRSA). Together, our results show that *S. aureus* is dependent on balanced amounts of membrane-attached SmdA in order to carry out proper cell division.

Importance

Staphylococcus aureus is an important human and animal pathogen. Antibiotic resistance is a

major problem in treatment of staphylococcal infections, and cell division and cell wall

synthesis factors have previously been shown to modulate susceptibility to antibiotics in this

species. In the current work we investigated the function of an essential protein named SmdA,

which was identified based on its septal localization and knockdown phenotype resulting in

defective cellular morphologies. We demonstrate that this protein is critical for normal cell

division in *S. aureus*. Depletion of SmdA sensitize resistant staphylococci to β-lactam

antibiotics. This work thus reveals a new staphylococcal cell division factor and a potential

future target for narrow spectrum antimicrobials or compounds to resensitize antibiotic

resistant staphylococcal strains.

Introduction

Most bacteria are surrounded by a shape-determining cell envelope which protects against lysis and interacts with the extracellular milieu. The cell envelope of the opportunistic, Gram-positive pathogen *Staphylococcus aureus* consists of a thick layer of peptidoglycan (PG) along with teichoic acids (TA) and cell wall-associated surface proteins. During a bacterial cell cycle, synthesis of PG and TA needs to be precisely regulated and coordinated with cell division, DNA replication and chromosome segregation. Tight control of these processes is critical for staphylococcal cells to maintain their integrity and nearly spherical shape as they multiply, and they are therefore attractive targets for antimicrobials (1). Exactly how such control is mediated in *S. aureus* is still not fully established, and hitherto unknown factors may be involved. In this work we describe a new staphylococcal cell morphology determinant.

Staphylococcal cell division is initiated by assembly of the Z-ring, consisting of polymerized FtsZ-proteins, that localizes to the future division septa (2). The Z-ring functions as a scaffold for cell division- and cell wall synthesis proteins which together constitute the divisome (3). Cell division in *S. aureus* occurs in alternating orthogonal planes, meaning that the new cell division plane is always perpendicular to the previous (4). Timely and spatial control of localization of the Z-ring assembly is most likely linked with chromosome segregation and DNA replication, involving proteins such as the nucleoid occlusion factor Noc, which ensures that the cells do not establish new septa across the chromosomes (5), and CcrZ, which connects initiation of DNA replication to cell division (6). The chromosomes and chromosome segregation also contribute to establish a physical barrier allowing the Z-ring only to be formed in an angle perpendicular to the previous division plane (4).

The Z-ring directs the synthesis of new PG in *S. aureus* to the septum. Synthesis of PG starts in the cytoplasm, where UDP-MurNAc-pentapeptide is first synthesized and then attached to the membrane by the enzyme MraY to form the PG precursor lipid I (7, 8). A

GlcNAc residue and a pentaglycine side chain is attached to produce lipid II-Gly$_5$ (9), which

is flipped to the outer leaflet of the membrane by MurJ (10, 11) where it is incorporated into

the existing PG mesh by transpeptidation (TP) and transglycosylation (TG) reactions.

Specifically, the shape, elongation, division and sporulation (SEDS) proteins, FtsW and RodA

with TG activity, work in pairs with monofunctional transpeptidases, the penicillin binding

proteins PBP1 and PBP3, respectively (12, 13). While the PBP1-FtsW pair is essential and

performs the septal cross wall synthesis, the non-essential PBP3-RodA pair is responsible for

the slight elongation taking place in *S. aureus*. Additionally, *S. aureus* possesses two other

PBPs; the bifunctional PBP2 with both TG and TP activity, whose role is essential in *S. aureus*,

and the low-molecular weight PBP4, which controls the degree of PG crosslinks (14-16).

Finally, MRSA strains have an additional PBP, PBP2A, a transpeptidase with low-affinity for

β-lactam antibiotics (17, 18). In the final step of division, PG hydrolases break covalent bonds

in PG for cell wall remodeling and daughter cell splitting. The major, bifunctional autolysin

Atl, together with Sle1, for which expression are regulated by the two-component system

WalKR, are the primary enzymes responsible for hydrolyzing the septal PG to allow splitting

of daughter cells (19-22). The actual cross wall splitting is a mechanical process occurring

within milliseconds (23, 24).

The spatiotemporal control of cell division and PG synthesis is directly and indirectly

influenced by several factors. One of these is the anionic TA polymers, the second major

component of the cell wall, which are either covalently linked to the PG (wall teichoic acids,

WTA), or linked via a lipid-anchor to the plasma membrane (lipoteichoic acids, LTA).

Mutations in enzymes involved in either WTA or LTA biosynthesis result in cells of abnormal

shape and lack of septum synthesis control, probably via different mechanisms (15, 25, 26).

Furthermore, proteases, chaperones and secretion proteins, involved in production, folding

and/or secretion of cell cycle proteins, may also directly or indirectly affect coordination of cell

division and septum formation in *S. aureus*. For example, Clp-protease complexes can target

both FtsZ and Sle1, and thereby have major effects on these processes (27-29).

Evidently, cell shape maintenance and control of cell division and septum formation

are a complex interplay between many cellular processes where unknown key factors are yet

to be discovered. We here identified a hitherto novel protein named SmdA (for staphylococcal

morphology determinant A). We show that the level of the membrane-attached SmdA protein

is critical for maintaining normal division progression and morphology in *S. aureus*, as

silencing or overexpression of *smdA* resulted in defective cell division and septal cross wall

synthesis, as well as increased sensitivity towards cell wall targeting antibiotics.

Results

SmdA is a conserved staphylococcal membrane protein

To identify novel proteins potentially involved in cell cycle and morphology control in *S.*

aureus, we performed a combined depletion- and subcellular localization analysis of essential

staphylococcal proteins with no annotated functions (30-32). From this, we identified a septum-

enriched protein (SAOUHSC_01908, named SmdA), whose knockdown resulted in cells of

variable sizes that formed large clusters (see below). The function of SmdA in *S. aureus* was

investigated further here.

SmdA is a protein of 302 amino acids which is fully conserved in species within the

Staphylococcaceae family (Fig. S1). The *smdA* gene is monocistronic and located >100 bp

away from the neighboring genes (SAOUHSC_01907; unknown function, and *metK*; S-

adenosylmethionine synthase). The protein has a predicted N-terminal transmembrane helix

and a C-terminal cytoplasmic part with partial homology to a so-called nuclease related domain

(NERD) (PF08375, E = $1.08 \cdot 10^{-5}$) (Fig. 1A). The domain was named based on distant

similarities to endonucleases, and is found in bacterial, archaeal, as well as plant proteins (33).

However, the functional role of NERD in bacteria has to our knowledge never been studied.

SmdA has been found to be essential in transposon mutagenesis studies in *S. aureus* (30, 32).

Growth analysis of SmdA CRISPRi knockdown (hereafter SmdA^{down}) in *S. aureus* SH1000 in

rich medium at 37°C resulted in a reduction in growth compared to the control strain when

spotted on agar plates (Fig. 1B). The growth rate in liquid medium was only slightly affected

when CFU and OD$_{600}$ was determined at different time points in the same culture during growth

(Fig. S2A). By RT-PCR we verified that the expression of *smdA* was indeed fully knocked

down by the CRISPRi system in *S. aureus* SH1000 (Fig. S2B). Since the SmdA^{down} strains

were still viable, we attempted to construct deletion mutants of *smdA* by allelic replacement

with a spectinomycin resistance cassette using the pMAD-vector (34). However, we were not

able to obtain the deletion mutant in *S. aureus* SH1000. Similarly, for *S. aureus* strains

NCTC8325-4, HG001 and the MRSA strain COL, SmdA^{down} resulted in reduced growth on

agar plates, with HG001 being most affected (Fig. 1B), but we were not able to obtain any

deletion mutants in these strains. We therefore used the CRISPRi system in the different *S.*

aureus strains to study the phenotypes of SmdA further.

Depletion and overexpression of SmdA result in cells with highly aberrant cell shapes

During the initial analysis, we observed that SmdA^{down} in *S. aureus* SH1000 resulted in clusters

containing cells of variable sizes. The knockdown experiment was repeated, and exponentially

growing cells were stained with fluorescent vancomycin (VanFL, binds to non-crosslinked

stem peptides throughout the cell wall). SmdA^{down} indeed resulted in severe phenotypic defects

(Fig. 1C); increased cell clustering and a large fraction of cells with multiple septa (20.1 % for

SmdA^{down} as opposed to 1.6 % for the control strain, Fig. 1D) and abnormal, non-spherical

morphology (Fig. 1E). The SmdA^{down} cells were also significantly larger than the control strain

(Fig. 1F).

Transmission- and scanning electron microscopy (TEM and SEM) were used to obtain more detailed images of the defects in morphology and septal placement found in SmdA[down] cells. Strikingly, SmdA depleted *S. aureus* SH1000 displayed highly aberrant septum formation (Fig. 2A). In addition to lysis, cells with several non-perpendicular- or parallel septa were frequently observed, resulting in cells, or small cell clusters, with aberrant morphologies (Fig. 2A). This was also evident from the SEM micrographs, which showed clustered cells with various morphologies (Fig. 2B). Similar phenotypes from TEM and SEM analyses were observed for the NCTC8325-4, HG001 and COL strains, with the HG001 strain being more affected by SmdA[down] than the other two (Fig. S3 and Fig. S4).

Since reduced levels of SmdA led to defects in cell division and morphology of *S. aureus*, we next wondered whether overexpression of SmdA would affect the cells. An ectopic copy of *smdA* under control of an IPTG-inducible promotor in the plasmid pLOW (35) was expressed in *S. aureus* NCTC8325-4. As for SmdA[down], the cells were stained with VanFL (Fig. S5A). Although less evident than for SmdA[down], overexpression of SmdA also resulted in clusters of cells with several septa per cell as defined by VanFL staining (Fig. S5A) (3.9 %, n = 181 for SmdA overexpression, compared to 20.1 % for SmdA[down], and 1.6 % for the control). Together, these knockdown- and overexpression experiments clearly demonstrate that septum formation and splitting in *S. aureus* are dependent on proper levels of SmdA.

Depletion of SmdA results in increased sensitivity towards antimicrobials targeting cell wall synthesis

Given the negative effect SmdA depletion had on cell morphology and -division, we reasoned that reduced expression of SmdA could influence the sensitivity of *S. aureus* to cell wall targeting antibiotics. To test this, SmdA[down] strains were treated with PBP-targeting β-lactams (oxacillin, cefotaxime, cefoxitin and imipenem), the glycopeptide vancomycin (blocking cell

wall synthesis by targeting the terminal D-Ala-D-Ala on the stem peptides of nascent PG (36)),

tunicamycin (targeting TarO and MraY, enzymes involved in the early stages of WTA and PG

synthesis, respectively (37)), targocil (targeting the WTA exporter TarG (38)) and Congo Red

(inhibitor of the LTA biosynthesis enzyme LtaS (39)) (Table 1). Two antibiotics with

alternative targets; tetracycline (targeting protein synthesis) and ciprofloxacin (a quinolone

targeting DNA synthesis), were also included.

For the methicillin-susceptible *S. aureus* (MSSA) strain SH1000, a two-fold reduction

in minimum inhibitory concentration (MIC), compared to the control, was observed for

oxacillin and cefoxitin in SmdA[down]. More notably, in the MRSA strain COL, SmdA[down]

sensitized the COL strain towards all β-lactams with a 2- to 8-fold reduction in MIC compared

to the control (Table 1). SmdA depletion did not seem to significantly influence vancomycin

susceptibility. Strikingly, however, we observed that SmdA[down] cells became highly sensitive

towards tunicamycin, with a 64-fold reduction in MIC compared to the control for *S. aureus*

SH1000 and >4-fold reduction for COL (40, 41). We therefore also tested the tunicamycin

sensitivity of the NCTC8325-4 and HG001 SmdA[down] strains, which also showed increased

susceptibility towards tunicamycin (HG001; 125-fold reduced MIC and NCTC8325-4; 4-fold

reduced). While no difference in MIC was observed for targocil, the SmdA[down] strains also

displayed increased sensitivity towards Congo Red. Finally, SmdA depletion in *S. aureus*

SH1000 led to a 2- to 4-fold reduction in MIC against tetracycline, compared to the control,

however, depletion of SmdA in COL did not change its sensitivity towards tetracycline or

ciprofloxacin (Table 1).

SmdA has no major effects on the TA biosynthetic pathways

The increased sensitivity to tunicamycin and Congo Red in SmdA[down] strains, prompted us to

study whether there were any major alterations in the TA in these cells, although it was also

noted that no changes in sensitivity was observed for the WTA export inhibitor targocil (Table

1). Notably, the SmdA[down] strain displayed morphologies reminiscent of what has previously

been reported for cells depleted of TA, i.e., with larger cell sizes, cells with irregular septum

formations and reduced splitting (26, 37, 38, 42). It has previously been shown that there is a

synthetic lethal relationship between the WTA and LTA biosynthetic pathways (42, 43), and it

could therefore be hypothesized that hypersensitivity to tunicamycin could result from deficient

LTA biosynthesis in the *smdA[down]* mutants, Using an anti-LTA antibody, we compared the

quantity and lengths of LTA in the SmdA[down] - and control strains for SH1000, NCTC8325-4,

HG001 and COL (Fig. S6A). For HG001, the strain with most severe division defects, we

observed a reduction in LTA amounts in SmdA[down] and this can possibly contribute to the

severe phenotype of this strain. However, no consistent changes in LTA amounts or lengths

were observed between the SmdA depletions and the controls in the four strains. Furthermore,

we could not detect any LTA release into the growth medium in the depletion strains, indicating

that the stability of LTA (44) was intact (Fig. S6B). We therefore conclude that SmdA does

not have any consistent effect on LTA synthesis across strains.

WTA has been shown to protect cells from the LTA-inhibitor Congo Red. Without

WTA, cells became hypersensitive towards Congo Red (MIC of <4 μg/ml for *tarO* deletion

mutants and >1024 μg/ml for wild-type cells) (45). Although to a much lesser degree, SmdA

depletion strains were also more sensitive to Congo Red compared to the controls (Table 1),

and we therefore looked into whether WTA could be disturbed in a SmdA[down] strain. Cells

without WTA has previously been shown to lack the dark, electron-dense layer observed in

TEM images of crosswalls of *S. aureus* wild type cells (Fig. S6C) (37, 46). TEM images of

TarO[down] and SmdA[down] showed that TarO[down], as expected (37, 46), lacked this dark, high-

density layer, while it was still present in the SmdA[down] strain, suggesting that WTA was still

produced (Fig. S6C). We also performed Fourier transform infrared spectroscopy (FTIR),

which has been used before to detect differences in the composition of WTA due to variable

glycosylation patterns (44, 47). As expected, changes in the FTIR spectra were evident in the

polysaccharide region (1200 cm^{-1} - 800 cm^{-1}) for the TarO^{down} strain compared to the control,

with the most significant differences recorded for the peaks at 1076 cm^{-1}, 1048 cm^{-1}, 1033 cm^{-1}

^{1} and 1000 - 970 cm^{-1}, representing α- and β-glycosidic bonds in WTA (47). However, no

changes were observed between SmdA^{down} and the control (Fig. S6D). Together, these results

suggests that SmdA has no major effect on the WTA biosynthetic pathway.

SmdA is important in several stages of staphylococcal cell division

To identify potential protein interaction partners of SmdA, we next performed a protein

pulldown experiment using GFP-trapping with a chromosomal *smdA-m(sf)gfp* fusion strain.

Interestingly, the major staphylococcal autolysin Atl, as well as the bifunctional PBP2 were

identified, along with 12 other proteins (Table S1). The experiment was repeated in a strain

with plasmid-based expression of SmdA-m(sf)GFP, and this setup resulted in an extended list

of proteins that were pulled down, probably due to the elevated expression of SmdA-m(sf)GFP.

Selecting the proteins for which at least 10 unique peptides were detected and with a fold

change of >2 compared to the control, resulted in a list of 57 proteins (Table S1). Several

proteins with activity involved in protein folding, secretion and/or degradation (e.g., FtsH,

PrsA, SpsB, ClpB, ClpC, SecD) were identified in this assay, in addition to the penicillin-

binding proteins PBP1, PBP2, PBP3 and the early division protein EzrA. All the 14 proteins

identified in the initial experiment were also pulled down in the second experiment.

Bacterial two-hybrid analyses of the SmdA-PBP1-3 and SmdA-EzrA were performed

to see whether these interactions could be reproduced in a heterologous system. The proteins

were fused either N- or C-terminally to the domains of adenylate cyclase, an enzyme which

catalyzes the production of cyclic adenosine monophosphate (cAMP) and eventually induction

of β-galactosidase production when brought in proximity by interaction between the target proteins. Indeed, SmdA interacted with PBP2, as well as PBP1, PBP3 and EzrA in the two-hybrid assays (Fig. 3A). By expressing a version of SmdA without its N-terminal membrane domain (SmdAΔTMH), we also show that the observed interactions between SmdA and the PBPs in this assay occur via the transmembrane segment, while the interaction with EzrA was retained for SmdAΔTMH. The latter suggests that EzrA interacts with the intracellular part of SmdA (Fig. 1A). Furthermore, we also show that SmdA is able to self-interact, and also this interaction is dependent on the transmembrane helix (Fig. S7).

The observed interactions between SmdA and PBPs may suggest that SmdA is somehow important for proper regulation and/or localization of PBPs in *S. aureus* (Fig. 1 and Fig. 2). The VanFL approach used above (Fig. 1C) labels non-crosslinked PG throughout the cell surface. Therefore, to determine more precisely the sites of active PG synthesis in *S. aureus* NCTC8325-4 SmdA^{down} and SmdA overexpression cells, we pulse-labelled the cells for 90 sec with the fluorescent D-amino acid 7-hydroxycoumarincarbonylamino-D-alanine (HADA), a molecule which is integrated into PG by the specific activity of transpeptidases (48). In the control cells, the expected midcell-localized band was observed (Fig. 3B). However, in the SmdA^{down} strain, we observed a more diffuse HADA signal with PG synthesis often occurring at several sites within the cell, forming patterns visible as crosses or Y-shapes (Fig. 3C). Similar observations were made in the cells overexpressing SmdA, although with a lower frequency (Fig. 3D-E). Thus, the site of active transpeptidation (and thus probably at least one of the PBPs) is mis-localized when the levels of SmdA is altered.

Moreover, we also studied whether SmdA could affect the localization of EzrA by knocking down *smdA* in a strain expressing a chromosomal *ezrA-gfp* fusion (Fig. 3F). Indeed, abnormal localization patterns for EzrA-GFP was observed with similar frequency as the abnormal HADA labelling (Fig. 3G). These results thus suggest that depletion of SmdA,

directly or indirectly, influences the localization of both early and late divisome proteins in *S.*

aureus.

Furthermore, the major autolysin Atl was also pulled down with SmdA. Atl is a secreted

multidomain enzyme, which is processed to an acetylmuramyl-L-alanine amidase and β-N-

acetylglucosaminidase, involved in septal cross wall splitting resulting in daughter cell

separation (19). SmdA[down] cells frequently displayed increased clustering (Fig. 2, Fig. S4). This

indicates reduced cross wall splitting, a phenotype also observed in Δ*atl* mutants and WalKR

depleted cells (22, 49). WalKR is the two-component regulatory system controlling the

expression of *atl* and other cell wall hydrolase encoding genes (22). To assess the reduced cross

wall splitting phenotype in more detail, we performed Triton X-100-induced autolysis assays

on the cultures. Indeed, reduced autolysis was observed in SmdA[down] cells, demonstrating

reduced autolytic activity (Fig. 4A). The lysostaphin sensitivity was also reduced (Fig. 4B),

suggesting alterations in the cell wall affecting the lytic properties of this enzyme. It should be

noted that the resistance towards Triton X-100- and lysostaphin-induced autolysis in the

SmdA[down] strain was clearly less compared to the control strain where *walR*, and thus all the

regulated autolysins, was knocked down. It should also be noted that Triton-X induced

autolysis was not severely altered upon overexpression of SmdA (Fig. S5B).

SmdA localizes at the septal region after FtsZ

The SmdA-m(sf)GFP fusion protein displayed a septum-enriched signal when expressed

ectopically from the low copy-number plasmid pLOW (Fig. 5A). In order to analyze the SmdA

localization with native expression levels, a chromosomally integrated version of *smdA-*

m(sf)gfp was made, in which SmdA was expressed with m(sf)GFP fused to its C-terminus.

Localization analyses in these cells further confirmed the septum-enriched localization of the

fusion protein (Fig. 5A), with an average septum/periphery fluorescence signal ratio of 3.4 (n

= 52, see Materials and Methods). To visualize the localization of SmdA relative to the cell

division process, we used FtsZ as a marker. Expression of *ftsZ*-fusion genes in the *smdA-*

m(sf)gfp chromosomal fusion strain, however, only resulted in strains with extremely poor

growth, suggesting that the cells did not tolerate such double-labelling. Instead, we therefore

created a double-labelled strain in which SmdA-mYFP and FtsZ-mKate2 were co-expressed

from plasmids, while the native *smdA* and *ftsZ* genes were still present on the chromosome.

Structured illumination microscopy (SIM) analysis showed that SmdA-mYFP is localized

around in the membrane when the Z-ring is formed (Fig. 5B), and that a septum-enriched

localization occurs as the septal cross wall is being synthesized. This was further confirmed by

stimulated emission depletion (STED) microscopy analysis (Fig. 5C). STED imaging also

revealed that there was no apparent co-localization of SmdA-mYFP and FtsZ-mKate2 in

newborn cells before FtsZ-constriction initiates. Combined, the localization of SmdA is

reminiscent of the localization of PBPs in *S. aureus* (11), but appears to localize to the septal

area after the early divisome proteins such as EzrA (50, 51).

The SmdA protein is anchored to the membrane by a single transmembrane helix (Fig.

1A). To verify its significance for the septal localization, we ectopically expressed a version of

SmdA without the transmembrane helix (SmdAΔTMH). The N-terminally truncated

SmdAΔTMH-m(sf)GFP localized to the cytoplasm of the cells (Fig. 5A). This shows that while

the transmembrane helix is critical for protein-protein interactions and subcellular localization

of SmdA, the interaction observed between EzrA and SmdAΔTMH (and full-length SmdA)

(Fig. 3A) is probably not involved in determining the localization of this protein.

During these latter experiments, we noticed that SmdAΔTMH-m(sf)GFP

overexpression led to cells with obvious morphology defects. In fact, overexpression of

SmdAΔTMH (without the GFP-tag) (Fig. 6A) strikingly resulted in a more extreme phenotype

than overexpression of full length SmdA (Fig. 3D, Fig. S5). The cells were often inflated or

with a bean-shaped appearance (Fig. 6A). HADA pulse labelling of these cells revealed a

highly abnormal PG incorporation pattern, with the fluorescent signal forming both condensed

clumps and Y-shapes in different directions in almost 50 % of the cells (Fig. 6A and B). This

demonstrates that membrane attachment is critical for the localization and function of SmdA.

Conserved amino acids in the predicted NERD domain are critical for the function of SmdA

The functional importance of the NERD domain (Fig. 1A) has, to our knowledge, neither been

studied nor verified previously in any bacterial protein. As mentioned, we observed that

overexpression of SmdAΔTMH led to severe phenotypic changes in the staphylococcal

morphology (Fig. 6A). We decided to use this as a tool to gain insight into whether the

predicted NERD domain is important for the function of SmdA. Multiple sequence alignment

of SmdA proteins from different staphylococcal species (Fig. S1) revealed conserved residues

in this domain, and site directed mutagenesis was used to create two versions of SmdAΔTMH

containing mutations in the NERD domain: one in which H145 was changed to Ala (mut1),

and a second containing the mutations R150A and T151A (mut2). Furthermore, we also

observed from the alignment that the C-terminus of SmdA is highly conserved (Fig. S1), and

another C-terminally mutated variant was created (mut3; F280A, H281A). The three-

dimensional structure of SmdA, predicted by AlphaFold (52), suggests that the N-terminal part

of the protein folds into a long α-helix and that the NERD domain is located within the C-

terminal structured part of the protein (Fig. 6C-II). The mutated SmdAΔTMH versions were

overexpressed in a wild-type background. Expression of these mutated variants indeed

partially, or fully, abolished the phenotypic defects observed when overexpressing the non-

mutated SmdAΔTMH version. HADA staining of mut1 (H145A) showed septum placement

and PG synthesis comparable to wild-type cells, while mut2 (R150A, T151A) and mut3

(F280A, H281A) also clearly reduced the functionality of the protein (Fig. 6A and B).

Furthermore, the growth inhibition upon overexpression of SmdAΔTMH (Fig. 6D) was also

abolished when the mutated variants were expressed under the same conditions. This strongly

suggests that the given residues in both the predicted NERD and the conserved C-terminus are

important for the function of SmdA.

Discussion

The results presented here provide the first functional insights into the conserved

staphylococcal protein SAOUHSC_01908, here named SmdA. SmdA, a protein conserved

within the *Staphylococcaceae* family, is critical for maintaining cell morphology and for proper

progression of cell division in different *S. aureus* strains (including MRSA and MSSA).

Unbalanced levels, as well as loss of proper subcellular localization, of SmdA, result in severe

cell morphology defects due to mis-localized cell division, uncontrolled cell wall synthesis,

and lack of proper cross wall splitting.

We observed that knockdown of SmdA increased the sensitivity of *S. aureus* to several

cell wall targeting antibiotics, including β-lactams and tunicamycin (Table 1). Many of the

frequently used antibiotics, including β-lactams, target cell wall synthesis, but the rise of both

methicillin- resistant- and vancomycin-resistant *S. aureus* (MRSA and VRSA) have made *S.*

aureus infections more difficult to treat and novel anti-staphylococcal targets and strategies are

needed. Targeting cell cycle proteins have been shown to re-sensitize resistant staphylococci

towards existing antibiotics due to synergistic effects (53-55). For example, inactivation of

some proteins involved in WTA synthesis sensitizes MRSA to β-lactams (37, 56). The same is

also the case for a diversity of factors contributing to cell division and cell wall biogenesis,

including FtsZ, FtsA, PBP4 and most proteins involved in the PG synthesis pathway (55). The

secretion-associated proteins SecDF and the chaperones PrsA and HtrA1 have also been tightly

linked to β-lactam sensitivity (57, 58), and recently, inactivation of the autolytic cell wall

amidase Sle1 (21), or the membrane proteins AuxA and AuxB (44), were shown to result in

increased sensitivity to β-lactams. Mechanisms of re-sensitization in these cases vary and may

be a result of weakened cell wall, as well as inactivation or mis-localization of PG synthesis,

including the key resistance determinant PBP2A. The large number of factors affecting β-

lactam sensitivity reflects the tight links between different processes involved in cell wall

biosynthesis, and most likely SmdA affects several steps in the cell division process (Fig. 1-3).

The exact mechanism by which SmdA affects these processes remains to be determined.

Our results suggest that cell division and PG synthesis, but not teichoic acid biosynthesis, is

affected. This is supported by observations showing mis-localization of cell division and PG

synthesis in SmdA^{down} cells (Fig. 1-3). Furthermore, SmdA^{down} cells displayed hypersensitivity

towards tunicamycin (targeting enzymes involved both in PG and WTA synthesis), but not

towards targocil (targeting the WTA exporter only) (38). Possibly, SmdA influences the

localization of cell division and cell wall synthesis by direct protein-protein interactions. This

is supported by the SmdA-EzrA and SmdA-PBP1-3 interactions, combined with the abnormal

localization of EzrA-GFP and HADA incorporation in SmdA^{down} cells (Fig. 3). In fact, the

division defects observed in SmdA^{down} cells (Fig. 1-3) are reminiscent of previous studies of

ezrA null mutants, strains with point mutations in FtsZ or cells exposed to Z-ring inhibitors (50,

51, 59, 60). However, the contribution of the identified interactions to the observed phenotypes

must be investigated further, particularly since the subcellular localization of SmdA-m(sf)GFP

does not appear to overlap with the Z-ring throughout the cell cycle. The SmdA-localization,

which depends on the transmembrane helix, is instead more similar to the reported localization

of PBPs (Fig. 5) (11, 50, 51).

Lack of cross wall splitting is another prominent feature of the SmdA depleted cells

(Fig. 2 and 4). After septum formation, cells remain attached, forming clusters of mis-shaped

cells. Indeed, the autolytic activity was reduced in the SmdA^{down} cells (Fig. 4). The major

autolysin Atl was also pulled down together with SmdA-m(sf)GFP as one of the major hits

(Table S1). Atl is an extracellular processed protein whose amidase and glucosaminidase

domains together process PG in the septum to allow splitting. It is possible that SmdA

somehow influences the export or processing of Atl. In this context, it is interesting to note that

proteases, foldases and chaperones (FtsH, PrsA, SpsB, ClpC, ClpB, HtrA1) were also pulled

down with SmdA (Table S1). It remains to be determined whether these interactions have any

functional relevance, however, perturbation of such pathways may have major consequences

for cell division and septal placement as they are important for proper folding and secretion of

key proteins involved in such processes. This has, for example, previously been demonstrated

for the chaperone ClpX, which is critical for coordination of autolysins and cell division

proteins (29). Involvement in such a mechanism could also explain the pleiotropic phenotypes

observed in SmdA^{down} cells, affecting several stages in the cell division process.

A part of SmdA displays limited similarity to the so-called nuclease related domain

(NERD). This domain, which is found in a broad range of bacterial species and in some archaea

and plants, was initially identified on a virulence-plasmid from *Bacillus anthracis* (33), and

has later been suggested to belong to a superfamily of phosphodiesterases (61). The majority

of NERD proteins are single-domain proteins, but they are also occasionally found together

with a kinase-, pseudokinase- or helicase domain (62). The actual function of the NERD has,

to our knowledge, however, not been studied experimentally. The results in this paper show

that conserved residues in the predicted NERD are important for the functionality of SmdA,

thus demonstrating for the first time a functional role of this domain (Fig. 6). Although highly

speculative at this point, it would be interesting if a cell division factor such as SmdA had

DNA-interacting capabilities and thus forming a potential link between the membrane-

associated cell division proteins and the nucleoid. It should also be noted that several of the

most conserved residues of the NERD domain (33) are not found in SmdA, suggesting a functional diversity among proteins harboring homology to this domain and implying that the NERD of SmdA may have functions unrelated to the nucleoid.

It is also interesting to note that the effect of SmdA knockdown was different between strains. For example, SmdA[down] resulted in a more severe phenotype in HG001 compared to NCTC8325-4 and SH1000 (Fig. 1, Fig. S3-S4), although these strains derive from the same parent (63). While HG001 contains prophages, NCTC8325-4 and SH1000 have been cured of these, and it could for example be speculated that the stress imposed by knockdown of SmdA somehow results in prophage induction in HG001 and thereby a more severe phenotype. Understanding of these and other strain differences could also help to further pinpoint the exact function of SmdA.

The staphylococcal genome encodes hundreds of essential proteins, which all represent potential target sites for antimicrobials. To be able to fully exploit the antibiotic target repertoire, it is critical to understand how essential proteins are involved and linked between different cell cycle processes. In this work, we identified and characterized SmdA as a novel factor essential for cell morphology and cell division in *S. aureus*. Based on the results presented here, future research should aim at pinpointing the molecular mechanism by which SmdA affect different stages of the cell division process. Finally, since SmdA depletion results in increased sensitivity to β-lactams, it may as such also be a possible future target for combatting β-lactam resistance by re-sensitizing MRSA to these antibiotics.

Materials and Methods

Bacterial strains, growth conditions and transformations

Escherichia coli IM08B (64), *E. coli* XL1-Blue and *E. coli* BTH101 were grown in lysogeny broth (LB) at 30-37°C with shaking or on lysogeny agar (LA) plates at 30-37°C, with 100

µg/ml ampicillin and/or 50 µg/ml kanamycin for selection. *S. aureus* SH1000, NCTC8325-4,

HG001 and COL were grown in brain-heart infusion (BHI) broth or tryptic soy broth (TSB) at

37°C with shaking or on BHI/TSA plates at 37°C. For selection, 10 µg/ml of chloramphenicol,

5 µg/ml of erythromycin, 15 µg/ml neomycin or 100 µg/ml spectinomycin were added. For

induction of gene expression, 50 or 300 µM isopropyl β-D-1-thiogalactopyranoside (IPTG) or

2.5 ng/ml anhydrotetracycline (ATc) were added to the bacterial cultures.

A standard heat shock protocol was used for transformation of *E. coli* IM08B. Isolated

plasmids from *E. coli* were transformed into *S. aureus* by electroporation. Preparation of

electrocompetent *S. aureus* cells and electroporation were performed according to Löfblom et

al (65). Bacterial strains used in this study are listed in Table S2.

Genetic modifications

For all cloning, DNA fragments were amplified from *S. aureus* SH1000 genomic DNA, and

cloning was performed with restriction digestion (New England Biolabs [NEB]) and

subsequent ligation using T4 DNA Ligase (NEB), unless otherwise stated. Ligation mixtures

were transformed into *E. coli* IM08B, and sequence-verified plasmids were transformed into

S. aureus. All strains and primers used in this study are listed in Table S2 and Table S3,

respectively.

Fluorescent fusion constructs

Construction of pLOW-smdA-m(sf)gfp. Monomeric superfolder GFP, *m(sf)gfp*, was initially

fused to *ftsZ* in the plasmid pLOW (35). *m(sf)gfp* was amplified from the plasmid pMK17 (66)

using forward primer im1_linker_FP_F_BamHI (annealing to the linker sequence) and reverse

primer im2_m(sf)gfp_R_NotI_EcoRI. The amplified fragment encodes a linker sequence N-

terminally of the *m(sf)gfp* gene. The fragment was digested with BamHI and EcoRI and

subsequently ligated into the corresponding sites of the plasmid pLOW-*ftsZ-gfp* to replace the

original *gfp*-gene with the linker_m(sf)gfp sequence. SAOUSHSC_01908 was then amplified

with primers im77_SA1908_F_SalI_RBS and im78_SA1908_R_BamHI, the product was

digested with SalI and BamHI and ligated into the corresponding sites of pLOW-*ftsZ-m(sf)gfp*

to create pLOW-*smdA-m(sf)gfp*.

Construction of pMAD-smdA-m(sf)gfp_aad9 for chromosomal integration. To tag the

chromosomal *smdA*, the plasmid pMAD-*smdA-flag_aad9* was constructed initially. The insert

in this plasmid (*smdA-flag_aad9*) was assembled by overlap extension PCR (primers im147-

im152), with the flag-tag sequence inserted by primer overhangs, and cloned into pMAD (34)

using restriction enzymes MluI and BamHI and T4 DNA Ligase (NEB). The insert was

designed so that the flag-tag could be removed using NotI and SpeI. The *m(sf)gfp* sequence

was amplified from template pMK17 (66) using the primers im153 and im154. The PCR

product was digested with NotI and SpeI and subsequently ligated with NotI- and SpeI-digested

pMAD-*smdA-flag_aad9*. The ligation mix was transformed to *E. coli* IM08B, and the resulting

plasmid, pMAD-*smdA-m(sf)gfp_aad9,* was verified by sequencing and transformed into

electrocompetent *S. aureus* SH1000. Integration of *smdA-m(sf)gfp* in the native locus of *smdA*

using the temperature-sensitive pMAD-system was carried out as previously described (34).

Construction of pHC-ftsZ-mKate2. *ftsZ-mKate2* was amplified from the plasmid pLOW-*ftsZ-*

mKate2. pLOW-*ftsZ-mKate2* was made in a similar manner as described for pLOW-*ftsZ-*

m(sf)gfp, but with reverse primer im5_mKate_R_NotI_EcoRI and genomic DNA derived from

strain MK119 (67) as template for amplification of *mKate2*. Then, *ftsZ-mKate2* was amplified

with primers USHC109 and USHC148, generating a product with SalI and MluI as overhang.

The PCR product and the plasmid pSK9065 (68) were digested with SalI and MluI and thereafter ligated.

Construction of pLOW-smdA-mYFP. This plasmid was constructed by using pLOW-*ftsZ-mYFP* as starting point, made similarly as described for pLOW-*ftsZ-m(sf)gfp*, but with reverse primer im3_cfp_myfp_R_NotI_EcoRI and plasmid pMK20 (lab collection) as template for amplifying *mYFP*. The plasmids pLOW-*smdA-m(sf)gfp* (described above) and pLOW-*ftsZ-mYFP* were digested with SalI and BamHI and subsequently ligated, resulting in pLOW-*smdA-mYFP*.

CRISPRi constructs

Construction of pCG248-sgRNA(smdA). For gene knockdowns, the CRISPR interference (CRISPRi) system developed by Stamsås et al. (69) was used. In this system, *dcas9* is placed downstream of an IPTG-inducible promoter in the plasmid pLOW-*dcas9*. A second plasmid is carrying the single guide RNA (pCG248-sgRNA(*x*), where *x* represents the targeted gene), which constitutively expresses the sgRNA, including the 20 nt base-pairing region specific for the gene to be knocked down. The pLOW-*dcas9* plasmid contains an erythromycin resistance gene, and the pCG248-sgRNA(*x*) plasmid a chloramphenicol resistance gene. The gene-specific 20 nt sequences were replaced in the pCG248-sgRNA(*x*) plasmids using an inverse PCR approach as described earlier (69) using primers mk299 and mk323.

Construction of pLOW-dCas9_aad9. In order to make the CRISPRi system compatible with the MRSA strain COL (which is intrinsically erythromycin resistant), the erythromycin resistance gene *ermC* in pLOW-*dcas9* was replaced with *aad9*, encoding a spectinomycin resistance cassette. The primers im183 and im184 were used to amplify the entire pLOW-*dcas9*

plasmid, except the *ermC* gene. The *aad9* gene was amplified with primers im185 and im186

using pCN55 (70) as template, where im185 contained the sequence of im183 as overhang, and

im186 the sequence of im184 as overhang. Thus, the two fragments had overlapping sequences

and were fused using NEBuilder® HiFi DNA Assembly (NEB). The construct was transformed

to *E. coli* IM08B, and the plasmid verified by PCR and sequencing. For CRISPRi in *S. aureus*

COL, the pLOW-*dcas9_aad9* plasmid was used along with pCG248-sgRNA(*x*).

Construction of plasmids used for overexpression studies

Construction of pLOW-smdA and mutated versions. *smdA* was amplified from *S. aureus*

genomic DNA using primers im77_SA1908_F_SalI_RBS and mk517_1908_R_NotI. The

fragment was digested with SalI and NotI and ligated into the corresponding sites of plasmid

pLOW-*dcas9* (69) to produce the plasmid pLOW-*smdA*, with IPTG-inducible overexpression

of *smdA*. pLOW-*smdAΔTMH* was constructed in a similar manner, except that primer

mk518_1908_F_RBS_SalI was used instead of im77 to remove the 29 N-terminal amino acids

of SmdA, predicted to encode the TMH and extracellular part. Site-directed mutagenesis in the

two plasmids was performed by a two-step overlap extension PCR approach, where the

mutations were introduced in the primers. The primers mk519 and mk520 were used for

introducing mutation H145A (mut1), mut2 (R150A, R151A) was made with primers mk521

and mk522, and mut3 (F280A, H281A) with primers mk529 and mk530.

Phase contrast- and fluorescence microscopy analysis

For induction of plasmid-encoded fluorescent fusions, exponentially growing cultures were

diluted to OD_{600} 0.05 in medium with 50 μM IPTG an incubated for 2 hours prior to

microscopy. For CRISPRi-knockdown and overexpression experiments, cultures of $OD_{600} =$

0.4 were diluted 250-fold in medium with 300 μM IPTG and grown until OD_{600} was

approximately 0.4. For labelling of cell wall, BODIPY^{TM} FL vancomycin (VanFL) (Invitrogen) was added. To label newly synthesized peptidoglycan, HADA (van Nieuwenhze group, Indiana University) was added actively growing cultures (OD$_{600}$ approximately 0.4) at a final concentration of 250 μM. The cells were incubated with HADA at 37°C for 90 sec and then put on ice. The cultures were pelleted by centrifugation at 10 000 × g at 4°C for 1 min, washed with cold 1 × PBS, pH 7.4 and finally resuspended in 25 μl PBS. Cells were subsequently added to agarose pads and microscopy was performed on a Zeiss AxioObserver with ZEN Blue software. An ORCAFlash4.0 V2 Digital complementary metal-oxide semiconductor (CMOS) camera (Hamamatsu Photonics) was used to capture images through a 100 × PC objective. HPX 120 Illuminator (Zeiss) was used as a light source for fluorescence microscopy.

All microscopy analyses were repeated at least two times. Images were processed and prepared for publication using Fiji (71). For analysis of cell roundness, cell area and fluorescent labelling patterns, MicrobeJ (72) was used to determine cell outlines. Outlines were corrected manually, when necessary. Cell area and roundness as a measure of morphology were calculated and plotted using MicrobeJ. After cell outline detection, categorization of cells into normal or abnormal labelling patterns were done manually using images from independent experiments. The septum/periphery fluorescence signal ratio was measured as described before (73) in cells with full septal signal of SmdA-GFP. A fluoresce ratio between 2.5 and 3.5 is typical for septum-enriched proteins (11).

Structured illumination microscopy (SIM) and stimulated emission depletion (STED) microscopy

Super-resolution SIM imaging was performed using a Zeiss ELYRA PS.1 microscope equipped with a 100 × 1.46 NA alpha plan apochromat oil immersion objective and a pco.edge sCMOS camera. Fluorescence images were acquired sequentially using 200–300 ms exposure

times per image, for a total of 15 images per SIM reconstruction. All imaging was performed

at room temperature (~23°C). Raw data were reconstructed using the SIM algorithms in ZEN

2011 SP7 software (black edition, Carl Zeiss). Brightfield images were captured using

widefield imaging mode. Images had a final pixel size of 25 nm.

Gated STED (gSTED) images were acquired on a Leica TCS SP8 STED 3× system,

using an HC PL Apo 100 × oil immersion objective with NA 1.40. Fluorophores were excited

using a white excitation laser operated at 509 nm for mYFP and 563 nm for mKate2. A STED

depletion laser line was operated at 592 nm and 775 nm for mYFP and mKate2, respectively,

using a detection time-delay of 0.8–1.6 ns for both fluorophores. The total depletion laser

intensity was in the order of 20–40 MW cm^{-2} for all STED imaging. The final pixel size was

13 nm and scanning speed was 400 Hz. The pinhole size was set to 0.9 AU.

Images were processed and analyzed using Fiji (71). Line scans were analyzed using

the Plot Profile function in Fiji, using a line width of 1.5. Fluorescence intensities were

normalized to the highest value for each channel.

Scanning- and transmission electron microscopy analysis

Overnight cultures were diluted to approximately $OD_{600} = 0.1$. When OD_{600} reached 0.4, the

cultures were diluted 1:250. Antibiotic and IPTG were added when appropriate. The cultures

(10 ml) were grown to $OD_{600} = 0.3$ and 1 volume of fixation solution, containing 5 % (w/v)

glutaraldehyde and 4 % (w/v) paraformaldehyde in 1×PBS, pH 7.4, was added. The tubes were

carefully inverted a few times and incubated for one hour at room temperature before being

placed at 4°C overnight. The following day, the cultures were centrifugated at 5000 × g, and

the pellets washed three times with PBS. Further preparations of samples to be analyzed with

TEM were performed as described before (69).

Samples for SEM were, after washing with PBS, dehydrated with EtOH, essentially in

the same manner as for sample preparations for TEM (69). The samples were subjected to

critical point drying by exchanging the EtOH with CO_2. Then, the samples were coated with a

conductive layer of Au-Pd before being analyzed in a ZEISS EVO50 EP Scanning electron

microscope. Images were analyzed and prepared using Fiji (71).

Growth assays

For examining growth on solid media, overnight cultures were diluted 1:250 in BHI containing

300 µM IPTG, unless otherwise specified. When reaching exponential phase, OD_{600} were

adjusted to 0.3 for all samples. A 10-fold dilution series were made for all strains, and 2 µl of

each dilution were spotted on BHI agar containing proper antibiotics and 300 µM IPTG. The

plates were incubated at 37°C for approximately 16 h, and pictures of the plates were captured

in a Gel Doc^{TM} XR + Imager (Bio-Rad).

For measurement of growth in liquid cultures, cells were at OD_{600} 0.4, were diluted

1:250 in medium containing 300 µM IPTG. Every hours for five hours, OD_{600} was measured

spectrophotometrically using Genesys 30 (Thermo Scientific) and dilutions of the cultures were

plated for CFU-counting.

Minimum inhibitory concentration (MIC) assays

The experiments were set up in 96-well microtiter plates with a total volume of 300 µl. A two-

fold dilution series of the antibiotics were prepared in BHI containing selective antibiotics and

IPTG when appropriate. The overnight cultures were diluted 1:1000 in BHI containing 300

µg/ml IPTG for induction. The cells were grown at 37°C, and the plate was shaken for 5 sec

before measurements of OD_{600} were taken every 10th min throughout the experiment, using

either a Synergy^{TM} H1 Hybrid Multi-Mode Reader (BioTek Instruments) or a Hidex Sense

(Hidex Oy). The experiments were repeated at least two times with the same results.

RNA isolation and RT-PCR

To verify that *smdA* expression was knocked down by CRISPRi, RNA was isolated from

exponentially growing cultures of IM284 (SH1000, CRISPRi(control)), IM165 (SH000

CRISPRi(empty)) and IM269 (SH1000 CRISPRi(*smdA*)). Isolation of total RNA and cDNA

synthesis were performed as previously described (69). A PCR reaction (30 cycles) was run

with Phusion® High-Fidelity DNA polymerase (NEB). The primer pairs im126/im127 and

im137/im138 were used to target the reference gene *pta* (74) and *smdA*, respectively.

Detection of lipoteichoic acid (LTA) by Western blotting

Detection of LTA was performed by western blotting, and sample preparations were done

according to descriptions found in Hesser et al (26). The samples were separated on a 4-20 %

gradient Mini Protean TGX acrylamide gel (BioRad), and subsequently transferred to a

polyvinylidene difluoride (PVDF) membrane by semi-dry electroblotting. The membrane was

blocked for 1 h in 5 % (w/v) skim milk in PBST and placed overnight at 4°C. Next, the

membrane was incubated for 1 h with α-LTA (Hycult) 1:4000 in PBST, washed three times

(10 minutes each) with PBST before incubation for 1 h with α-Mouse IgG HRP Conjugate

(Promega) secondary antibody (1:10 000 in PBST). The membrane was again washed three

times and LTA bands were visualized by using SuperSignal^{TM} West Pico PLUS

Chemiluminescent substrate (Thermo Fisher Scientific) in an Azure Imager c400 (Azure

Biosystems).

Same procedure as in (44) was carried out for possible detection of LTA released to the

medium. The supernatants, after harvesting cells during sample preparations, were kept.

Supernatant samples were centrifugated for 16 000 × *g* for 10 min, and 75 µl was mixed with

25 µl 4 × SDS-PAGE sample buffer. These samples were boiled for 30 min and applied on the

4-20 % Mini Protean TGX acrylamide gel. Thereafter, the same immunodetection procedure

as described above was followed.

Fourier-transform infrared spectroscopy (FTIR) analysis

Cultures of the strains IM313 (HG001, CRISPRi(control)), IM312 (HG001, CRISPRi(*smdA*))

and IM357 (HG001, CRISPRi(*tarO*)) were initially pre-grown to exponential phase, back

diluted to OD_{600} = 0.05 and induced with 300 µM IPTG. The bacterial cells (1 ml) were

harvested at OD_{600} = 0.4 by centrifugation at 5000 × *g*, 4°C, for 3 min. The pelleted cells were

kept at -20°C prior to further processing. Pellets were resuspended in 40 µl 0.1 % (w/v) NaCl,

and 10 µl of the suspensions were added to an IR-light-transparent silicon 384-well microplate

(Bruker Optic, Germany), with three technical replicates for each sample. The plates were left

to dry at room temperature for approximately 2 h. FTIR spectra were recorded in transmission

mode using a high-throughput screening extension (HTS-XT) unit coupled to a Vertex 70 FTIR

spectrometer (Bruker Optik GmbH, Leipzig, Germany). Spectra were recorded in the region

4000-500 cm^{-1}, with a spectral resolution of 6 cm^{-1}, a digital spacing of 1.928 cm^{-1}, and an

aperture of 5 mm. For each spectrum, 64 scans were averaged. The OPUS software (Bruker

Optik GmbH, Leipzig,Germany) was used for data acquisition and instrument control. The

obtained spectra were processed by taking second derivatives and extended multiplicative

signal correction (EMSC) preprocessing in Unscrambler X version 11 (CAMO Analytics,

Oslo, Norway). Results presented are averaged spectra from 3 biological replicates (each with

3 technical replicates) for the region with wavelengths between 1200 cm^{-1} and 800 cm^{-1}.

GFP-trap and liquid chromatography with tandem mass spectrometry (LC-MS/MS)

Cultures (*S. aureus* SH1000 wild-type, IM308 SH1000 *smdA-m(sf)gfp*, IM104 SH1000 pLOW-*smdA-m(sf)gfp* and IM164 SH1000 pLOW-*smdA-flag*) were pre-grown to exponential phase, back diluted to $OD_{600} = 0.05$ and induced if necessary. When reaching OD_{600} at 0.4, 80 ml of each culture were harvested by centrifugation at $4000 \times g$, 4°C for 3 min. Supernatants were decanted and pellets resuspended in cold TBS prior to transfer to 1.5 ml microcentrifuge tubes. Centrifugation was repeated for 1 min, and pelleted cells were stored at -80°C prior to further use.

For GFP-trap, cells were resuspended in 1 ml cold buffer containing 10 mM Tris pH 7.5, 150 mM NaCl, 1 mM PMSF, 6 µg/ml RNase and 6 µg/ml DNase. Suspensions were transferred to 2 ml lysing matrix B tubes (MP Biomedicals) containing 0.8 g ≤ 106 µm glass beads (Sigma-Aldrich) and subjected for mechanical lysis by agitation in a FastPrep-24™ (MP Biomedicals) for 3×30 sec at 6.5 m/s, with 1 min pause on ice between the runs. Tubes were centrifugated at $5000 \times g$, 4°C for 10 min, and supernatants transferred to new tubes. Concentrations were determined measuring Abs280 using NanoDrop™ 2000 (Thermo Fisher Scientific), where a small amount of the samples were added a final concentration of 1 % (w/v) SDS prior to measurements. GFP-Trap beads (25 µl per sample) (Chromotek) were washed three times with 500 µl ice cold Dilution/Wash buffer (10 mM Tris pH 7.5, 150 mM NaCl, 0.5 mM EDTA) and centrifugated at $2500 \times g$, 4°C for 5 min. Lysates were diluted in Dilution/Wash buffer to a final concentration of 1 mg in a total volume of 500 µl, before being transferred to GFP-Trap beads. Samples were placed in a Bio RS-24 Multi rotor (Biosan) at 4°C for 1 h. Then, samples were centrifugated at $2500 \times g$, 4°C for 5 min, supernatant removed, and beads washed three times with Dilution/Wash buffer. During last washing step, solutions were transferred to new tubes, and after centrifugation and removal of supernatant, beads were resuspended in 50 µl 5 % (w/v) SDS, 50 mM Tris pH 7.6. Tubes were incubated at 95°C for 5

min, and centrifuged at maximum speed for 30 sec. After standing at the bench a few minutes,

30-50 µl were transferred into new tubes. Samples were kept at -20°C and heated for 2 min at

95°C prior to the sample preparation method Suspension trapping (STrap), conducted as

described by A. Zougman et al. (75).

The peptide samples were analyzed by coupling a nano UPLC (nanoElute, Bruker) to a

trapped ion mobility spectrometry/quadrupole time of flight mass spectrometer (timsTOF Pro,

Bruker). The peptides were separated by an Aurora Series 1.6 µm C18 reverse-phase 25 cm ×

75 µm analytical column with nanoZero and CaptiveSpray Insert (IonOpticks, Australia). The

flow rate was set to 400 nl/min and the peptides were separated using a gradient from 2 % to

95 % acetonitrile solution (in 0.1 % (v/v) formic acid) over 120 minutes. The timsTOF Pro was

ran in positive ion data dependent acquisition PASEF mode, with a mass range at 100-1700

m/z. The acquired spectra were analyzed against a *S. aureus* NCTC8325 proteome database.

Bacterial two-hybrid (BACTH) assays

Plasmid construction, and procedure for the BACTH assays, were conducted in a same manner

as previously described (69), and primers used are listed in Table S3. Briefly, gene fusions of

selected genes, to the T18 or T25 domains of adenylate cyclase form *Bordetella pertussis,* were

made by restriction cutting and ligation in the plasmid vectors pKT25, pKNT25, pUT18 or

pUT18C (Euromedex). *E. coli* XL1-Blue cells were used for transformation, and plasmids

verified by sequencing before BACTH assays (76) were set up according to the manufacturer

(Euromedex). Co-transformation of plasmids containing fusion-genes of opposite domains,

that is T25 in one plasmid and T18 in the other, were done in *E. coli* BTH101 with 50 µg/ml

kanamycin and 100 µg/ml ampicillin as selection markers. Five random colonies were picked,

grown in liquid LB to visible growth, and spotted on LA plates containing 40 µg/ml X-gal and

0.5 mM IPTG, in addition to the selection markers. Plates were incubated dark at 30°C for 20-

48 h before being inspected, and blue colonies are an indication of positive interaction between

tested genes. Presented results are representative for at least six independent replicates.

Acknowledgements

We would like to acknowledge the NMBU Imaging Center for help with electron microscopy,

Maria Victoria Heggenhougen and Marita Torrissen Mårli (both NMBU) for access to

unpublished strains and Henriette Olsen (NMBU) for help with FTIR. Mass spectrometry-

based proteomic analyses were performed by The MS and Proteomics Core Facility,

Norwegian University of Life Sciences (NMBU). This facility is a member of the National

Network of Advanced Proteomics Infrastructure (NAPI), which is funded by the Research

Council of Norway INFRASTRUKTUR-program (project number: 295910). We acknowledge

the van Nieuwenhze group, Indiana University, for providing HADA. The work is supported

by grants from the Research Council of Norway (project number 250976) and JPI-AMR

(project number 296906). Work in the Structural Cell Biology Unit (OIST) is supported by

OIST core subsidy. Ine S. Myrbråten acknowledge support from "Pasteurlegatet".

References

1. Sass P, Brötz-Oesterhelt H. 2013. Bacterial cell division as a target for new antibiotics. Curr Opin Microbiol 16:522-530.

2. Pinho MG, Kjos M, Veening J-W. 2013. How to get (a) round: mechanisms controlling growth and division of coccoid bacteria. Nat Rev Microbiol 11:601-614.

3. Errington J, Daniel RA, Scheffers D-J. 2003. Cytokinesis in bacteria. Microbiol Mol Biol Rev 67:52-65.

4. Saraiva BM, Sorg M, Pereira AR, Ferreira MJ, Caulat LC, Reichmann NT, Pinho MG. 2020. Reassessment of the distinctive geometry of *Staphylococcus aureus* cell division. Nat Comm 11:1-7.

5. Veiga H, Jorge AM, Pinho MG. 2011. Absence of nucleoid occlusion effector Noc impairs formation of orthogonal FtsZ rings during *Staphylococcus aureus* cell division. Mol Microbiol 80:1366-1380.

6. Gallay C, Sanselicio S, Anderson ME, Soh YM, Liu X, Stamsås GA, Pelliciari S, van Raaphorst R, Dénéréaz J, Kjos M, Murray H, Gruber S, Grossman AD, Veening JW. 2021. CcrZ is a pneumococcal spatiotemporal cell cycle regulator that interacts with FtsZ and controls DNA replication by modulating the activity of DnaA. Nat Microbiol 6:1175-1187.

7. Barreteau H, Kovač A, Boniface A, Sova M, Gobec S, Blanot D. 2008. Cytoplasmic steps of peptidoglycan biosynthesis. FEMS Microbiol Rev 32:168-207.

8. Bouhss A, Trunkfield AE, Bugg TD, Mengin-Lecreulx D. 2007. The biosynthesis of peptidoglycan lipid-linked intermediates. FEMS Microbiol Rev 32:208-233.

9. Rohrer S, Berger-Bächi B. 2003. FemABX peptidyl transferases: a link between branched-chain cell wall peptide formation and β-lactam resistance in gram-positive cocci. Antimicrob Agents Chemother 47:837-846.

10. Sham L-T, Butler EK, Lebar MD, Kahne D, Bernhardt TG, Ruiz N. 2014. Bacterial cell wall. MurJ is the flippase of lipid-linked precursors for peptidoglycan biogenesis. Science 345:220-222.

11. Monteiro JM, Pereira AR, Reichmann NT, Saraiva BM, Fernandes PB, Veiga H, Tavares AC, Santos M, Ferreira MT, Macario V, VanNieuwenhze MS, Filipe SR, Pinho MG. 2018. Peptidoglycan synthesis drives an FtsZ-treadmilling-independent step of cytokinesis. Nature 554:528-532.

760   12.   Reichmann NT, Tavares AC, Saraiva BM, Jousselin A, Reed P, Pereira AR, Monteiro
761   JM, VanNieuwenhze MS, Fernandes F, Pinho MG. 2019. SEDS-bPBP pairs direct
762   lateral and septal peptidoglycan synthesis in *Staphylococcus aureus*. Nat Microbiol
763   4:1368-1377.

764   13.   Meeske AJ, Riley EP, Robins WP, Uehara T, Mekalanos JJ, Kahne D, Walker S, Kruse
765   AC, Bernhardt TG, Rudner DZ. 2016. SEDS proteins are a widespread family of
766   bacterial cell wall polymerases. Nature 537:634-638.

767   14.   Wyke AW, Ward JB, Hayes MV, Curtis NA. 1981. A role *in vivo* for penicillin-binding
768   protein-4 of *Staphylococcus aureus*. Eur J Biochem 119:389-393.

769   15.   Atilano ML, Pereira PM, Yates J, Reed P, Veiga H, Pinho MG, Filipe SR. 2010.
770   Teichoic acids are temporal and spatial regulators of peptidoglycan cross-linking in
771   *Staphylococcus aureus*. Proc Natl Acad Sci U S A 107:18991-18996.

772   16.   Pinho MG, Errington J. 2005. Recruitment of penicillin-binding protein PBP2 to the
773   division site of *Staphylococcus aureus* is dependent on its transpeptidation substrates.
774   Mol Microbiol 55:799-807.

775   17.   Hartman BJ, Tomasz A. 1984. Low-affinity penicillin-binding protein associated with
776   beta-lactam resistance in *Staphylococcus aureus*. J Bacteriol 158:513-516.

777   18.   Ubukata K, Yamashita N, Konno M. 1985. Occurrence of a beta-lactam-inducible
778   penicillin-binding protein in methicillin-resistant staphylococci. Antimicrob Agents
779   Chemother 27:851-857.

780   19.   Oshida T, Sugai M, Komatsuzawa H, Hong Y-M, Suginaka H, Tomasz A. 1995. A
781   *Staphylococcus aureus* autolysin that has an N-acetylmuramoyl-L-alanine amidase
782   domain and an endo-beta-N-acetylglucosaminidase domain: cloning, sequence
783   analysis, and characterization. Proc Natl Acad Sci U S A 92:285-289.

784   20.   Nega M, Tribelli PM, Hipp K, Stahl M, Götz F. 2020. New insights in the coordinated
785   amidase and glucosaminidase activity of the major autolysin (Atl) in *Staphylococcus*
786   *aureus*. Commun Biol 3:1-10.

787   21.   Thalsø-Madsen I, Torrubia FR, Xu L, Petersen A, Jensen C, Frees D. 2019. The Sle1
788   cell wall amidase is essential for β-lactam resistance in community-acquired
789   methicillin-resistant *Staphylococcus aureus* USA300. Antimicrob Agents Chemother
790   64:e01931-19.

791   22.   Dubrac S, Boneca IG, Poupel O, Msadek T. 2007. New insights into the WalK/WalR
792   (YycG/YycF) essential signal transduction pathway reveal a major role in controlling

cell wall metabolism and biofilm formation in *Staphylococcus aureus*. J Bacteriol
189:8257-69.

23. Zhou X, Halladin DK, Rojas ER, Koslover EF, Lee TK, Huang KC, Theriot JA. 2015.
Mechanical crack propagation drives millisecond daughter cell separation in
Staphylococcus aureus. Science 348:574-578.

24. Monteiro JM, Fernandes PB, Vaz F, Pereira AR, Tavares AC, Ferreira MT, Pereira PM,
Veiga H, Kuru E, VanNieuwenhze MS. 2015. Cell shape dynamics during the
staphylococcal cell cycle. Nat Comm 6:1-12.

25. Brown S, Santa Maria Jr JP, Walker S. 2013. Wall teichoic acids of Gram-positive
bacteria. Ann Rev Microbiol 67:313-336.

26. Hesser AR, Matano LM, Vickery CR, Wood BM, Santiago AG, Morris HG, Do T,
Losick R, Walker S. 2020. The length of lipoteichoic acid polymers controls
Staphylococcus aureus cell size and envelope integrity. J Bacteriol 202:e00149-20.

27. Silber N, Pan S, Schäkermann S, Mayer C, Brötz-Oesterhelt H, Sass P. 2020. Cell
division protein FtsZ is unfolded for N-terminal degradation by antibiotic-activated
ClpP. mBio 11:e01006-20.

28. Feng J, Michalik S, Varming AN, Andersen JH, Albrecht D, Jelsbak L, Krieger S,
Ohlsen K, Hecker M, Gerth U, Ingmer H, Frees D. 2013. Trapping and proteomic
identification of cellular substrates of the ClpP protease in *Staphylococcus aureus*. J
Proteome Res 12:547-58.

29. Jensen C, Bæk KT, Gallay C, Thalsø-Madsen I, Xu L, Jousselin A, Torrubia FR,
Paulander W, Pereira AR, Veening J-W. 2019. The ClpX chaperone controls autolytic
splitting of *Staphylococcus aureus* daughter cells, but is bypassed by β-lactam
antibiotics or inhibitors of WTA biosynthesis. PLoS Pathog 15:e1008044.

30. Chaudhuri RR, Allen AG, Owen PJ, Shalom G, Stone K, Harrison M, Burgis TA,
Lockyer M, Garcia-Lara J, Foster SJ, Pleasance SJ, Peters SE, Makell DJ, Charles IG.
2009. Comprehensive identification of essential *Staphylococcus aureus* genes using
Transposon-Mediated Differential Hybridisation (TMDH). BMC Genomics 10:291.

31. Santiago M, Matano LM, Moussa SH, Gilmore MS, Walker S, Meredith TC. 2015. A
new platform for ultra-high density *Staphylococcus aureus* transposon libraries. BMC
Genomics 16:1-18.

32. Valentino MD, Foulston L, Sadaka A, Kos VN, Villet RA, Santa Maria J, Lazinski DW,
Camilli A, Walker S, Hooper DC. 2014. Genes contributing to *Staphylococcus aureus*
fitness in abscess-and infection-related ecologies. mBio 5:e01729-14.

33.   Grynberg M, Godzik A. 2004. NERD: a DNA processing-related domain present in the
anthrax virulence plasmid, pXO1. Trends Biochem Sci 29:106-110.

34.   Arnaud M, Chastanet A, Débarbouillé M. 2004. New vector for efficient allelic
replacement in naturally nontransformable, low-GC-content, Gram-positive bacteria.
Appl Environ Microbiol 70:6887-6891.

35.   Liew AT, Theis T, Jensen SO, Garcia-Lara J, Foster SJ, Firth N, Lewis PJ, Harry EJ.
2011. A simple plasmid-based system that allows rapid generation of tightly controlled
gene expression in *Staphylococcus aureus*. Microbiology 157:666-676.

36.   Reynolds PE. 1989. Structure, biochemistry and mechanism of action of glycopeptide
antibiotics. Eur J Clin Microbiol Infect Dis 8:943-950.

37.   Campbell J, Singh AK, Santa Maria Jr JP, Kim Y, Brown S, Swoboda JG, Mylonakis
E, Wilkinson BJ, Walker S. 2011. Synthetic lethal compound combinations reveal a
fundamental connection between wall teichoic acid and peptidoglycan biosyntheses in
Staphylococcus aureus. ACS Chem Biol 6:106-116.

38.   Campbell J, Singh AK, Swoboda JG, Gilmore MS, Wilkinson BJ, Walker S. 2012. An
antibiotic that inhibits a late step in wall teichoic acid biosynthesis induces the cell wall
stress stimulon in *Staphylococcus aureus*. Antimicrob Agents Chemother 56:1810-20.

39.   Vickery CR, Wood BM, Morris HG, Losick R, Walker S. 2018. Reconstitution of
Staphylococcus aureus lipoteichoic acid synthase activity identifies Congo red as a
selective inhibitor. J Am Chem Soc 140:876-879.

40.   Price NP, Tsvetanova B. 2007. Biosynthesis of the tunicamycins: a review. J Antibiot
60:485-491.

41.   Swoboda JG, Campbell J, Meredith TC, Walker S. 2010. Wall teichoic acid function,
biosynthesis, and inhibition. Chembiochem 11:35-45.

42.   Santa Maria JP, Jr., Sadaka A, Moussa SH, Brown S, Zhang YJ, Rubin EJ, Gilmore
MS, Walker S. 2014. Compound-gene interaction mapping reveals distinct roles for
Staphylococcus aureus teichoic acids. Proc Natl Acad Sci U S A 111:12510-12515.

43.   Oku Y, Kurokawa K, Matsuo M, Yamada S, Lee B-L, Sekimizu K. 2009. Pleiotropic
roles of polyglycerolphosphate synthase of lipoteichoic acid in growth of
Staphylococcus aureus cells. J Bacteriol 191:141-151.

44.   Mikkelsen K, Sirisarn W, Alharbi O, Alharbi M, Liu H, Nøhr-Meldgaard K, Mayer K,
Vestergaard M, Gallagher LA, Derrick JP. 2021. The novel membrane-associated
auxiliary factors AuxA and AuxB modulate β-lactam resistance in MRSA by stabilizing
lipoteichoic acids. Int J Antimicrob Agents 57:106283.

45. Suzuki T, Campbell J, Kim Y, Swoboda JG, Mylonakis E, Walker S, Gilmore MS.
2012. Wall teichoic acid protects *Staphylococcus aureus* from inhibition by Congo red
and other dyes. J Antimicrob Chemother 67:2143-2151.

46. Chan YG-Y, Kim HK, Schneewind O, Missiakas D. 2014. The capsular polysaccharide
of *Staphylococcus aureus* is attached to peptidoglycan by the LytR-CpsA-Psr (LCP)
family of enzymes. J Biol Chem 289:15680-15690.

47. Grunert T, Jovanovic D, Sirisarn W, Johler S, Weidenmaier C, Ehling-Schulz M, Xia
G. 2018. Analysis of *Staphylococcus aureus* wall teichoic acid glycoepitopes by
Fourier Transform Infrared Spectroscopy provides novel insights into the
staphylococcal glycocode. Sci Rep 8:1-9.

48. Radkov AD, Hsu Y-P, Booher G, VanNieuwenhze MS. 2018. Imaging bacterial cell
wall biosynthesis. Annu Rev Biochem 87:991-1014.

49. Biswas R, Voggu L, Simon UK, Hentschel P, Thumm G, Götz F. 2006. Activity of the
major staphylococcal autolysin Atl. FEMS Microbiol Letters 259:260-268.

50. Jorge AM, Hoiczyk E, Gomes JP, Pinho MG. 2011. EzrA contributes to the regulation
of cell size in *Staphylococcus aureus*. PLoS One 6:e27542.

51. Lund VA, Wacnik K, Turner RD, Cotterell BE, Walther CG, Fenn SJ, Grein F,
Wollman AJ, Leake MC, Olivier N, Cadby A, Mesnage S, Jones S, Foster SJ. 2018.
Molecular coordination of *Staphylococcus aureus* cell division. eLife 7:e32057.

52. Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool
K, Bates R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ,
Cowie A, Romera-Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman
D, Clancy E, Zielinski M, Steinegger M, Pacholska M, Berghammer T, Bodenstein S,
Silver D, Vinyals O, Senior AW, Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly
accurate protein structure prediction with AlphaFold. Nature 596:583-589.

53. Roemer T, Schneider T, Pinho MG. 2013. Auxiliary factors: a chink in the armor of
MRSA resistance to β-lactam antibiotics. Curr Opin Microbiol 16:538-548.

54. Tan CM, Therien AG, Lu J, Lee SH, Caron A, Gill CJ, Lebeau-Jacob C, Benton-
Perdomo L, Monteiro JM, Pereira PM, Elsen NL, Wu J, Deschamps K, Petcu M, Wong
S, Daigneault E, Kramer S, Liang L, Maxwell E, Claveau D, Vaillancourt J, Skorey K,
Tam J, Wang H, Meredith TC, Sillaots S, Wang-Jarantow L, Ramtohul Y, Langlois E,
Landry F, Reid JC, Parthasarathy G, Sharma S, Baryshnikova A, Lumb KJ, Pinho MG,
Soisson SM, Roemer T. 2012. Restoring methicillin-resistant *Staphylococcus aureus*
susceptibility to β-lactam antibiotics. Sci Transl Med 4:126ra35.

55. Lee SH, Jarantow LW, Wang H, Sillaots S, Cheng H, Meredith TC, Thompson J,
Roemer T. 2011. Antagonism of chemical genetic interaction networks resensitize
MRSA to β-lactam antibiotics. Chem Biol 18:1379-1389.

56. Brown S, Xia G, Luhachack LG, Campbell J, Meredith TC, Chen C, Winstel V, Gekeler
C, Irazoqui JE, Peschel A. 2012. Methicillin resistance in *Staphylococcus aureus*
requires glycosylated wall teichoic acids. Proc Natl Acad Sci U S A 109:18909-18914.

57. Quiblier C, Zinkernagel AS, Schuepbach RA, Berger-Bächi B, Senn MM. 2011.
Contribution of SecDF to *Staphylococcus aureus* resistance and expression of virulence
factors. BMC Microbiol 11:72.

58. Roch M, Lelong E, Panasenko OO, Sierra R, Renzoni A, Kelley WL. 2019.
Thermosensitive PBP2a requires extracellular folding factors PrsA and HtrA1 for
Staphylococcus aureus MRSA β-lactam resistance. Commun Biol 2:417.

59. Pereira AR, Hsin J, Król E, Tavares AC, Flores P, Hoiczyk E, Ng N, Dajkovic A, Brun
YV, VanNieuwenhze MS, Roemer T, Carballido-Lopez R, Scheffers DJ, Huang KC,
Pinho MG. 2016. FtsZ-dependent elongation of a coccoid bacterium. mBio 7:e00908-
16.

60. Steele VR, Bottomley AL, Garcia-Lara J, Kasturiarachchi J, Foster SJ. 2011. Multiple
essential roles for EzrA in cell division of *Staphylococcus aureus*. Mol Microbiol
80:542-55.

61. Steczkiewicz K, Muszewska A, Knizewski L, Rychlewski L, Ginalski K. 2012.
Sequence, structure and functional diversity of PD-(D/E)XK phosphodiesterase
superfamily. Nucleic Acids Res 40:7016-7045.

62. Kwon A, Scott S, Taujale R, Yeung W, Kochut KJ, Eyers PA, Kannan N. 2019. Tracing
the origin and evolution of pseudokinases across the tree of life. Sci Signal
12:eaav3810.

63. Herbert S, Ziebandt AK, Ohlsen K, Schafer T, Hecker M, Albrecht D, Novick R, Gotz
F. 2010. Repair of global regulators in *Staphylococcus aureus* 8325 and comparative
analysis with other clinical isolates. Infect Immun 78:2877-89.

64. Monk IR, Tree JJ, Howden BP, Stinear TP, Foster TJ. 2015. Complete bypass of
restriction systems for major *Staphylococcus aureus* lineages. mBio 6:e00308-15.

65. Löfblom J, Kronqvist N, Uhlén M, Ståhl S, Wernérus H. 2007. Optimization of
electroporation-mediated transformation: *Staphylococcus carnosus* as model organism.
J Appl Microbiol 102:736-747.

66. van Raaphorst R, Kjos M, Veening J-W. 2017. Chromosome segregation drives
division site selection in *Streptococcus pneumoniae*. Proc Natl Acad Sci U S A
114:E5959-E5968.

67. Kjos M, Veening JW. 2014. Tracking of chromosome dynamics in live *Streptococcus*
pneumoniae reveals that transcription promotes chromosome segregation. Mol
Microbiol 91:1088-105.

68. Brzoska AJ, Firth N. 2013. Two-plasmid vector system for independently controlled
expression of green and red fluorescent fusion proteins in *Staphylococcus aureus*. Appl
Environ Microbiol 79:3133-3136.

69. Stamsås GA, Myrbråten IS, Straume D, Salehian Z, Veening JW, Håvarstein LS, Kjos
M. 2018. CozEa and CozEb play overlapping and essential roles in controlling cell
division in *Staphylococcus aureus*. Mol Microbiol 109:615-632.

70. Charpentier E, Anton AI, Barry P, Alfonso B, Fang Y, Novick RP. 2004. Novel
cassette-based shuttle vector system for Gram-positive bacteria. Appl Environ
Microbiol 70:6076-6085.

71. Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch
S, Rueden C, Saalfeld S, Schmid B. 2012. Fiji: an open-source platform for biological-
image analysis. Nat Methods 9:676-682.

72. Ducret A, Quardokus EM, Brun YV. 2016. MicrobeJ, a tool for high throughput
bacterial cell detection and quantitative analysis. Nat Microbiol 1:16077.

73. Pereira PM, Filipe SR, Tomasz A, Pinho MG. 2007. Fluorescence ratio imaging
microscopy shows decreased access of vancomycin to cell wall synthetic sites in
vancomycin-resistant Staphylococcus aureus. Antimicrob Agents Chemother 51:3627-
33.

74. Valihrach L, Demnerova K. 2012. Impact of normalization method on experimental
outcome using RT-qPCR in *Staphylococcus aureus*. J Microbiol Methods 90:214-216.

75. Zougman A, Selby PJ, Banks RE. 2014. Suspension trapping (STrap) sample
preparation method for bottom-up proteomics analysis. Proteomics 14:1006-1000.

76. Karimova G, Pidoux J, Ullmann A, Ladant D. 1998. A bacterial two-hybrid system
based on a reconstituted signal transduction pathway. Proc Natl Acad Sci U S A
95:5752-6.

77. Omasits U, Ahrens CH, Müller S, Wollscheid B. 2014. Protter: interactive protein
feature visualization and integration with experimental proteomic data. Bioinformatics
30:884-886.

78.   Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li W, Lopez R, McWilliam H,

963   Remmert M, Söding J, Thompson JD, Higgins DG. 2011. Fast, scalable generation of

964   high-quality protein multiple sequence alignments using Clustal Omega. Mol Syst Biol

965   7:539.

Figures and figure legends

Fig. 1. Phenotypes resulting from depletion of SmdA. (A) Predicted topology of SmdA using

Protter (77). SmdA is predicted to have one transmembrane (TM) helix with a short

extracellular N-terminus and a large intracellular domain. The sequence with predicted

similarity to the NERD domain is highlighted in yellow. **(B)** Growth on solid media of SmdA

knockdown strains (SmdA^down) in *S. aureus* SH1000 (IM269), NCTC8325-4 (IM311), HG001

(IM312) and COL (IM294). Strains carrying a non-targeting sgRNA were used as controls

(IM284, IM307, IM313 and IM295 for the respective strains). From non-induced overnight

cultures, 10-fold dilution series were made and spotted onto plates with 300 µM isopropyl-β-

D-thiogalactopyranoside (IPTG). **(C)** SmdA^down (IM269) and control strain (IM284) analyzed

by phase contrast- and fluorescence microscopy of cells stained with the cell wall label VanFL.

White arrows point at mis-shaped cells and cells with perturbed septum formation. Magnified

insets of representative cells are shown for the VanFL micrographs. Scale bars, 5 µm. **(D)**

Fraction of cells with multiple septa per cell for the SmdA^down strain IM269 (n = 225) and the

non-target control strain IM284 (n = 242) are plotted. The asterisks indicate significant

difference (Fisher's exact test, P<0.001). **(E)** Cell roundness, as determined using MicrobeJ,

was used as a measure of the morphology of the cells. Spherical cells will have values close to

1. Cell roundness measures for the control strain IM284 (n = 198) and the SmdA^down strain

IM269 (n = 191) are plotted. **(F)** Cell area (in μm^2) as determined using MicrobeJ of the control

strain IM284 (n = 198) and the SmdA^down strain IM269 (n = 191). In E and F, significant

differences between the distributions are indicted by asterisks (**, P < 0.001). P-values were

derived from a Mann-Whitney test.

Fig. 2. SmdA^{down} in *S. aureus* SH1000 visualized by electron microscopy. (A) Transmission electron- and **(B)** scanning electron micrographs of SH1000 CRISPRi control strain (IM284) and SH1000 SmdA^{down} (IM269). In (A), red arrows in the TEM micrographs point at lysed cells. Representative examples of cells with parallel septa or multiple septa are shown. Green arrows point to initiation of septum synthesis at multiple sites within the same cell. Different magnifications are shown, indicated by the scale bars.

Fig. 3. SmdA interacts with important cell division proteins and is necessary for proper localization of EzrA and peptidoglycan synthesis. (A) Protein-protein interactions tested with bacterial two-hybrid assays, where SmdA and SmdAΔTMH were tested against PBP1, PBP2, PBP3 and EzrA. The proteins were fused to the T18 or T25 domains as indicated. Blue bacterial spots and plus symbols indicate positive interactions and white spots and minus symbols indicate no interaction. **(B-D).** Micrographs of HADA labelled *S. aureus* NCTC8325-4 with depletion and overexpression of SmdA. Arrows point at cells with misplaced septum synthesis. Scale bars, 2 μm. **(B)** CRISPRi control strain, IM307 **(C)** SmdA^{down} strain, IM311 and **(D)** SmdA overexpression strain, MK1866. **(E)** Frequency plot of cells with normal or abnormal HADA labelling pattern. Categorization of normal or abnormal labelling patterns are indicated. The number of cells analyzed were 259, 179 and 189 for B, C and D, respectively. The asterisks indicate significant difference from the control (Fisher's exact test, $P<0.001$). **(F)** Micrographs showing co-localization of EzrA-GFP and HADA incorporation in *S. aureus* SH1000 strains with (MK1952) or without (MK1953) knockdown of SmdA. Phase contrast- and fluorescence images of HADA labelling and GFP (EzrA-GFP) are shown. Scale bars, 2 μm. Arrows point to cells with abnormal localization of both HADA and EzrA-GFP. **(G)** Frequencies of cells (from F) with normal or abnormal localization of HADA and EzrA-GFP are plotted. The number of cells analyzed were 285 (for MK1953) and 298 (for MK1952). The

1016 asterisks indicate significant difference form the respective controls (Fisher's exact test,

1017 P<0.001).

1018

1019 **Fig. 4**. Autolysis of *S. aureus* SH1000 depletion strains (control = IM284; WalR^{down} = IM293;

1020 SmdA^{down} = IM269) measured in presence of (**A**) 0.5 % Triton X-100 and (**B**) 100 ng/ml

1021 lysostaphin. Results presented as % of initial OD$_{600}$. Error bars represent standard error

1022 calculated from four technical replicates.

1023

Fig. 5. Subcellular localization analysis of SmdA. (**A**) Micrographs of cells with induced

expression of SmdA-m(sf)GFP from plasmid (top panel, IM104), native chromosomal

expression of the SmdA-m(sf)GFP fusion protein (middle panel, IM308), and expression of

SmdAΔTMH-m(sf)GFP from plasmid (bottom panel, IM373). Scale bars, 2 μm. (**B**) SIM

images of fixed *S. aureus* SH1000 with plasmid-expressed SmdA-mYFP and FtsZ-mKate2

(HC060). (**B-I**) Side-view of a cell showing that FtsZ localizes at septum before arrival of

SmdA. As cell division progresses, (**B-II**) SmdA concentrates at sites where septum formation

is initiated, and (**B-III**) displays a septal localization at the two septal membranes as FtsZ

constricts and septum formation proceeds. (**B-IV**) Top-view of a cell showing the FtsZ-ring

inside of the SmdA-ring. All scale bars, 0.5 μm. (**C**) STED images of fixed *S. aureus* SH1000

with plasmid-expressed SmdA-mYFP and FtsZ-mKate2 (HC060), where line scans show

fluorescence intensity of selected areas. (**C-I**) At an early stage of cell division, the rings of

both SmdA and FtsZ have a similar diameter, but do not overlap in their distribution, and a

heterogenous distribution with a patchy signal is observed. (**C-II**) Top-view of a cell as cell

division progresses, with the FtsZ-ring laying inside of SmdA, approximately 100-150 nm

apart from each other. (**C-III**) Side-view of a cell showing how FtsZ is located innermost, and

how SmdA is located at the two septal membranes. Scale bars, 0.5 μm.

Fig. 6. Membrane attachment and the NERD domain are important for SmdA function.

(**A**) HADA stained *S. aureus* NCTC8325-4 control cells not expressing any SmdA-variant (IM307) and cells overexpressing SmdAΔTMH (MK1911), SmdAΔTMH_mut1 (IM377), _mut2 (IM378) and _mut3 (IM379). Scale bars, 2 µm. (**B**) The frequency of cells from panel A with normal or abnormal HADA staining is plotted. The number of cells plotted were 259 the control, 82 for SmdAΔTMH, 101 for _mut1, 113 for _mut2 and 122 for _mut3. The asterisks indicate significant difference form the control (Fisher's exact test, $P<0.001$). ns indicates that there is not significant difference from the control. (**C**) Structure of SmdA, predicted by AlphaFold (52), where (**C-I**) shows the structure colored by a "per-residue confidence score" (pLDDT) with dark blue pLDDT > 90, light blue 90 > pLDDT > 70, and yellow 70 > pLDDT > 50, (**C-II**) shows the structured domain magnified with the predicted NERD colored in green and residues that have been mutated in red. (**C-III**) A magnified inset of (II) with annotated residues. The positions of the mutated residues are indicated in red. (**D**) Growth assay on solid media of *S. aureus* NCTC8325-4 expressing SmdAΔTMH, SmdAΔTMH_mut1, _mut2 and _mut3.

1057

Figures and figure legends

 Table 1. Minimum inhibitory concentration (MIC, in µg/ml) of different antimicrobials
 when SmdA was depleted in *S. aureus* SH1000 and COL.

Antibiotics	*S. aureus* SH1000 (MSSA)			*S. aureus* COL (MRSA)		
	SmdA[down]	Control	Fold change	SmdA[down]	Control	Fold change
Oxacillin	0.12	0.24	2	64-128	256	2-4
Cefotaxime	2	2	1	50	400	8
Cefoxitin	1	2	2	47-94	188	2-4
Imipenem	0.016	0.016	1	150	>300	>2
Vancomycin	2.5	2.5	1	2.5	2.5	1
Tetracycline	0.05	0.1-0.2	2-4	≥64	≥64	1
Ciprofloxacin	0.32	0.32	1	0.32	0.32	1
Tunicamycin	0.094	6	64	24	>96	>4
Targocil	1	1	1	1	1	1
Congo Red	256	>1024	>4	512-1024	>1024	≥2

Legends to supplementary material

Fig. S1. Multiple sequence alignment of SmdA from different staphylococcal species. Protein sequences were aligned with Clustal Omega (78). The blue shaded residues are predicted to be extracellular, the transmembrane domain is shaded in grey, and the predicted NERD domain is shaded in yellow. *S. aureus* NCTC8325-4 is highlighted in bold, and residues that were mutated are marked in green. The accession numbers of the sequences are indicated, and the first five letter in the sequence tags indicate the genus or species corresponding the to the sequences (for example, Nosoc; *Nococomiicoccus*, Aliic; *Aliicoccus*, Salin; *Salinicoccus*, Jeotg; *Jeotgalicoccus*, Auric; *Auricoccus*, Abyss; *Abyssicoccus*, Mepid; *Macrococcus epidermidis*, Mcase; *Macrococcus caseolyticus*, Ssciu; *Staphylococcus sciuri*, Sinte; *Staphylococcus intermedius*).

Fig. S2. Growth of SmdA knockdown strains in liquid cultures and verification of *smdA* silencing. (A) Growth of SmdA^{down} in *S. aureus* SH1000 (IM269) compared to the CRISRPi-control strain with a non-targeting sgRNA (IM284). IPTG (300 µg/ml) was added to induce expression of the CRISPRi-system and CFU/ml and OD$_{600}$ measured every hour for five hours. (**B**) Verification of *smdA* silencing by PCR with RT-PCR. cDNA was synthesized from RNA isolated from induced and un-induced cultures of SH1000 SmdA^{down} (IM269) and the CRISPRi control strains (IM284; non-targeting sgRNA and IM165; empty plasmid without sgRNA). Primers targeting either *smdA* or the housekeeping gene *pta*.

Fig. S3. Transmission electron microscopy (TEM) of different *S. aureus* strains with SmdA knockdown. SmdA^{down} and control cells analyzed with TEM in the *S. aureus* strains

1088 (**A**) NCTC8325-4 (IM311 and IM307), (**B**) HG001 (IM312 and IM313) and (**C**) COL (IM294

1089 and IM295). Th e sizes of the scale bars are indicated in the images.

1090

1091 **Fig. S4. Scanning electron microscopy (SEM) of cells depleted of SmdA.** SEM micrographs

1092 of SmdA^{down} and control cells in *S. aureus* (**A**) HG001 (IM312 and IM313) and (**B**) COL

1093 (IM294 and IM295). All scale bars, 1 μm.

1094

Fig. S5. Overexpression of SmdA. (**A**) Induced expression of an ectopic copy of *smdA* in the

plasmid pLOW in *S. aureus* NCTC8325-4 (MK1866). Cells were labelled with the cell wall

marker fluorescent vancomycin (VanFL). Scale bar, 2 μm. (**B**) Autolysis of *S. aureus* SmdA

overexpression strain (MK1866) compared to plasmid control strain (MK1465) monitored in

presence of 0.5 % Triton X-100. Results presented as % of initial OD_{600}. Error bars represent

standard error calculated from four technical replicates.

1101

Fig. S6. Analysis of teichoic acids. (**A-B**) Lipoteichoic acid (LTA) detection by

immunoblotting with α-LTA antibody was performed with (**A**) cell extract samples from

SmdA^{down} and CRISPRi control cells from *S. aureus* SH1000 (IM269 and IM284),

NCTC8325-4 (IM311 and IM307), HG001 (IM312 and IM313) and COL (IM294 and IM295),

and (**B**) cell extract (CE)- and supernatant (SN) samples from SmdA^{down} and CRISPRi control

cells from *S. aureus* SH1000 and NCTC8325-4 (IM269, IM284, IM311 and IM307,

respectively). In (A), the mean intensities in the bands (background subtracted) were

determined using Fiji (71). The mean intensities of the LTA bands in the *smdA* depletions

relative to their controls are plotted. All control strains express a non-targeting sgRNA. (**C-D**)

SmdA does not have major impact on the synthesis of wall teichoic acids. (**C**) TEM

micrographs of *S. aureus* NCTC8325-4 control strain (IM307) compared to SmdA and TarO

knockdown strains IM311 and IM358, respectively. The black arrows indicate presence of a

high-density layer in the septum, which is missing in the TarO depletion strain (white arrow).

(**D**) Fourier transform infrared spectroscopy (FTIR) of *S. aureus* HG001 control strain (IM313)

and knockdown strain of SmdA and TarO (IM312 and IM357, respectively). The

polysaccharide region of the spectrum is shown, and the indicated peaks represents α- and β-

glycosidic bonds.

1119

1120   **Fig. S7.** Bacterial two-hybrid analysis demonstrating self-interaction between SmdA proteins.

1121   Blue bacterial spots indicate positive interactions and white spots indicate no interaction.

1122

1123   **Table S1. Proteins pulled down with SmdA-GFP.**

1124

1125   **Table S2. Strains used in this study.**

1126

1127   **Table S3. Primers used in this study.**

1128   **Fig. 1**

1129

1130   **Fig. 2**

1131

1132

Fig. 3

1136   **Fig. 4**

1137

1138   **Fig. 5**

1139

1140

1141   **Fig. 6**