A bioactive phlebovirus-like envelope protein in a hookworm endogenous retrovirus

Monique Merchant[1,2,‡], Carlos P. Mata[1,2 ,‡ ,#], Yangci Liu[1,2], Haoming Zhai[1,2], Anna V. Protasio[3] and Yorgo Modis[1,2,*]

5   [1]Molecular Immunity Unit, Department of Medicine, University of Cambridge, MRC Laboratory of Molecular Biology, Francis Crick Avenue, Cambridge, CB2 0QH, United Kingdom

[2]Cambridge Institute of Therapeutic Immunology & Infectious Disease (CITIID), University of Cambridge School of Clinical Medicine, Cambridge CB2 0AW, UK

[3]Department of Pathology, University of Cambridge, Tennis Court Road, Cambridge

10   CB2 1QP, United Kingdom

[‡]These authors contributed equally

[#]Current address: Astbury Centre for Structural Molecular Biology, School of Molecular and Cellular Biology, Faculty of Biological Sciences, University of Leeds, Leeds LS2 9JT, United Kingdom

15   *Corresponding author: ymodis@mrc-lmb.cam.ac.uk

Keywords

Transposable element; retrotransposon; endogenous retrovirus (ERV); envelope protein; membrane fusion; horizontal gene transfer (HGT); virus evolution; gene exaptation; nematode

Short title

A bioactive phlebovirus-like Env in a hookworm ERV

Abstract

Endogenous retroviruses (ERVs), accounting for 14% of our genome, serve as a genetic reservoir from which new genes can emerge. Nematode ERVs are particularly diverse and informative of retrovirus evolution. We identify Atlas virus – a novel, intact ERV in the human hookworm *Ancylostoma ceylanicum* with an envelope protein genetically related to G_N-G_C glycoproteins from phleboviruses. A cryo-EM structure of Atlas G_C reveals a class II viral membrane fusion protein fold not previously seen in retroviruses. Atlas G_C has the structural hallmarks of an active fusogen. Atlas G_C trimers insert into membranes with endosomal lipid compositions and low pH. When expressed on the plasma membrane, Atlas G_C has cell-cell fusion activity. Atlas virus is transcriptionally active at specific stages of hookworm development. The preserved activities and expression pattern of Atlas G_C suggest it has acquired a cellular function. Our work reveals an unexpected degree of structural and genetic plasticity in retroviruses.

Introduction

Retroviruses can integrate their genome, reverse-transcribed from RNA into DNA, into the host-cell genome. Viral genomes integrated into germline cells are inherited by future generations as endogenous retroviruses (ERVs). ERVs account for approximately 14% of the human genome, seven times more than protein-coding genes. ERVs and other transposons were initially viewed as parasitic DNA. It is now evident that they serve as a genetic reservoir, from which new genes and regulatory elements can emerge. Sequences of retroviral origin help control gene expression by serving as promoters, enhancers and other regulatory elements (*1, 2*). Genes coopted from ERVs have evolved to fulfill vital cellular functions (*3*). For example, syncytins, which drive cell-cell fusion of trophoblasts during placental development, are encoded by ERV envelope glycoprotein (*env*) genes (*1, 4*). Another recent example is the Gag capsid protein encoded by the *Caenorhabditis elegans* Cer1 retrotransposon, which encapsidates small non-

coding RNAs into nucleocapsids that can transfer RNAs conferring learned pathogen avoidance

50   behavior from infected parents to naïve progeny (*5*).

The reduced mutation rate of host versus retrovirus genomes (10^{-9} vs. 10^{-3} mutations

per site per year (*6*)) means that ERVs are windows to ancient retroviral sequences –

evolutionary fossils preserved from the time of integration. Some ERV genes are expressed in

human tissues and retain their biological activities, such as membrane fusion activity in the case

55   of Env proteins (Envs) (*1, 4, 7*). Aberrant expression of Envs is associated with disease (*8, 9*).

With the biology of ERVs still largely uncharted, it is likely that many cellular functions of ERVs

in health and disease remain undiscovered. Studying ERV genes with novel properties could

therefore provide insights on the evolutionary history of retroviruses and identify fundamental

principles in host-virus coevolution.

60   Nematode ERVs are particularly diverse and informative of retrovirus evolution. ERVs

from the *Belpaoviridae* (BEL/Pao) family (*10*), widespread across metazoa, have revealing

genetic features in nematodes (*11*). The presence in *C. elegans* ERVs of overlapping ORFs,

otherwise unique to complex vertebrate retroviruses, suggests retroviruses originated in early

metazoa with a common ancestor resembling belpaoviruses (*12*). Furthermore, nematode

65   belpaovirus ERVs encode Envs that are genetically unrelated to all other retrovirus Envs(*13*).

Instead of a class I viral membrane fusion protein with a core fold of three bundled α-helices

(*14-17*) – a defining feature of modern retroviruses (*18*) – belpaovirus Envs have sequence

similarity to G_C (G2) envelope glycoproteins from phleboviruses (family *Phenuiviridae*) (*13*). G_C

proteins are class II membrane fusion proteins, with a three-domain β-strand architecture (*19,*

70   *20*) also found in alphaviruses (*21*), flaviviruses (*22, 23*) and Rubella virus (*24*) but structurally

unrelated to class I fusion proteins. A series of conformational changes in class II fusion

proteins, triggered by endosomal acidification, catalyses fusion of the viral and endosomal

membranes to deliver the viral genome into the cytosol (*25-27*). A hydrophobic fusion loop first

inserts into the endosomal membrane. The proteins then form trimers and fold back on

themselves, pulling the cell membrane (held by the fusion loop) and the viral membrane (held

by a transmembrane anchor) together so they fuse (*18, 20*).

Class II fusion proteins are not limited to viruses: they also drive cell-cell fusion events of

fundamental importance, including syncytial epithelia formation in *C. elegans* (*28*), and gamete

fusion in protozoa, plants, algae and invertebrates (*29-31*). The identical topology and overall

arrangement of the three domains of viral and eukaryotic class II fusion proteins, along with

similarities in their membrane fusion mechanisms, makes it all but certain they evolved from a

common ancestor (*29*). Although the evolutionary origin of the ancestral class II fusion protein

remains unknown, the presence of class II fusion proteins in ERVs raises the provocative

prospect that a gene transfer from a virus to a cell led to the advent of sexual reproduction (*29,

32*).

Here, we identify a novel, intact belpaovirus ERV in the human hookworm *Ancylostoma

ceylanicum* (a parasitic nematode) with an Env more similar than any other eukaryotic sequence

to phlebovirus G_C protein sequences. We expressed and purified the G_C-homologous fragment

from this ERV, henceforth Atlas virus. A cryo-electron microscopy (cryo-EM) structure of Atlas

G_C reveals a class II viral fusion protein fold remarkably similar to phlebovirus G_C proteins and

not previously seen in retroviruses. We show that Atlas G_C has all the hallmarks of an active

class II membrane fusion protein. It undergoes a monomer-to-trimer transition and inserts into

lipid membranes with a specific lipid composition in response to a low pH trigger. Our work

provides biochemical validation for the hypothesis that acquisition of a fusion protein from an

infectious virus, as exemplified by Atlas virus, represents a general paradigm for how

retrotransposons can become retroviruses (*13*), and indeed how ancestral reverse-transcribing

viruses may have originated (*10*). The preserved biological activities of Atlas G_C, including

membrane fusion activity, raise the question of whether these activities, and those of ERV gene products more broadly, have cellular functions or cause disease.

100 **Results**

An intact ERV with a phlebovirus-like Env in the hookworm *A. ceylanicum*

The bioinformatic discovery of nematode ERVs with phlebovirus G_C-like Env sequences not previously seen in retroviruses (*11-13*), requires biochemical validation. To identify phlebovirus-like ERV Envs suitable for biochemical analysis, we performed PSI-BLAST searches (*33*) for

105 protein sequences similar to biochemically characterized phlebovirus G_C proteins. A search with Rift Valley Fever virus (RVFV) G_C as the query identified the gene *Acey_s0020.g108* (UniProt A0A016UZK2) in the human hookworm *A. ceylanicum* as containing the most similar sequence outside infectious virus taxa (Expected value 10^{-20}). The homologous sequence lies within a 9,204-nucleotide element possessing all the features of an intact ERV, including 100% identical

110 271-nt LTRs and a coding sequence encoding a single 2,828-residue Gag-Pol-Env polyprotein without any stop codons or introns (**Fig. 1a; Supplementary Fig. 1a**). We refer to this element as *A. ceylanicum* Atlas virus. It is one of nine *A. ceylanicum* ERVs that encode complete Gag-Pol-Env polyproteins (**Fig. 1b,c**). These ERVs have the distinguishing genomic features of belpaoviruses from other nematode species, including an atypical Env and an aspartate to

115 asparagine substitution (Y[X]D<u>D</u> → YVD<u>N</u>) in the most conserved reverse transcriptase motif, the polymerase site (*12, 34, 35*) (**Supplementary Fig. 1b**). With a Gag sequence less than 20% identical to its closest homolog (*Acey_s0020.g1106*), Atlas virus is a candidate for classification as a member of the *Belpaoviridae* family (which contains a single genus, *Semotivirus*) (**Supplementary Fig. 1c**).

120 The phlebovirus G_C-like sequence spans the last 498 residues of the Atlas virus polyprotein (residues 2330-2828). It contains a single predicted C-terminal transmembrane helix, like phlebovirus G_C proteins. Phleboviruses and other *Phenuiviridae* family members

express a glycoprotein precursor that is cleaved by cellular proteases into two envelope

glycoproteins, G_N and G_C (or G1 and G2). G_N is the receptor-binding protein required for cellular

125   attachment and G_C is the membrane fusion protein required for cell entry. G_N is highly antigenic

and more variable in sequence than G_C. Our analysis of the Atlas virus Env sequence detected

a slight but statistically significant similarity in the 421 residues preceding the G_C-like sequence

(residues 1909-2329) to G_N glycoproteins from *Phenuiviridae* family viruses (*E*-value $\geq 10^{-7}$).

Moreover, the distribution of predicted transmembrane helices and proteolytic cleavage sites in

130   and adjacent to the Atlas G_N- and G_C-like sequences is the same as in phlebovirus

glycoproteins (**Fig. 1a**). Together, these sequence features suggest the Atlas virus Env contains

tandem phlebovirus-like G_N and G_C glycoproteins instead of a retrovirus-like glycoprotein. With

all the features of a recently active ERV and an apparently intact set of phlebovirus-like

glycoproteins, Atlas virus is an excellent candidate for biochemical analysis.

Figure 1. Phylogenetic analysis of class II-like Env genes and their closest relatives from modern infectious viruses

(a) Gene architecture of *Acey_s0020.g108*, an endogenous retrovirus with a phlebovirus-like Env from the *Ancylostoma ceylanicum* human hookworm parasite. We refer to this element here as *A. ceylanicum* Atlas virus. Inset: annotated closeup of the Atlas Env region encoding phlebovirus G_N- and G_C-like glycoproteins. Red residue number refer to the G_C sequence alone.

(b) List of *A. ceylanicum* ERVs encoding complete Gag-Pol-Env polyproteins with phlebovirus-like Envs. The length and sequence identity of LTRs is listed. LTRs sequence identities for two of the genes could not be calculated: *Acey_s0206.g1994* had degenerate LTRs, and one of the

LTRs of *Acey_s0025.g1106* was truncated in the genome assembly. Amino acid sequence

identities of Env and Gag sequences are listed.

(c) Phylogenetic tree of G_C proteins from *Phenuiviridae* and G_C-like (G2-like) sequences from

retrotransposons. Ace, *A.* ceylanicum; Cel, *C. elegans*; Ath, *Arabidopsis thaliana*; Cre,

150   *Chlamydomonas reinhardtii*; Ddi, *Dictyostelium discoideum*; CCHFV, Crimean Congo

hemorrhagic fever virus; SFV, Semliki Forest virus; DabieV, Dabie bandavirus; HRTV,

Heartland virus; JEV, Japanese encephalitis virus; WNV, West Nile virus; ZIKV, Zika virus;

DENV, dengue virus. Drawn with Interactive Tree Of Life (iTOL) (*36*).

Atlas G_C has a class II membrane fusion protein fold not previously seen in retroviruses

155   As the molecular structure of phlebovirus G_C proteins, and how they drive fusion of the viral and

cellular membranes, are well established from previous studies (*19, 27*), we focused our

biochemical analyses on the G_C-like sequence from Atlas virus. A recombinant ectodomain

fragment of Atlas G_C (polyprotein residues 2330-2772) was expressed in *Drosophila*

melanogaster D.mel-2 cells as a secreted protein. The purified protein was a soluble, folded

160   homotrimer (**Supplementary Fig. 2**). The structure of the G_C trimer was determined by single-

particle cryo-EM image reconstruction at an overall resolution of 3.76 Å (**Fig. 2a,b;**

Supplementary Table 1; Supplementary Fig. 3). The map was sufficiently detailed for an

atomic model to be built and refined for Atlas G_C residues 2330-2769 using the crystal structure

of RVFV G_C (*27*) as a starting model (see Methods; **Supplementary Fig. 4**). The structure

165   reveals a three-domain class II membrane fusion protein fold (**Fig. 2c**). All previously described

retroviral Env structures have a helical coiled-coil class I fusion protein fold (*14-17*), exemplified

by the hemagglutinin of influenza virus (*18*). The structure of the Atlas G_C ectodomain fragment

is remarkably similar to phlebovirus G_C structures, specifically the trimeric postfusion G_C

structures from RVFV (*27*), Dabie bandavirus (DABV, formerly SFTS phlebovirus) (*37*) and

170   Heartland virus (HRTV) (*38*) (**Fig. 2d-e**). Domain I, a 10-stranded β-barrel augmented by a 3-

stranded sheet, organizes the structure. Two insertions in domain I form the elongated, mostly β-stranded domain II. Domain III has the 7-stranded β-sandwich topology of fibronectin type III (FN3) domains also found in macroglobulin domains (*31, 39*), but the hydrophobic core and disulfide bonding pattern of domain III differ from these and other annotated domains from non-

175   viral species. A 16-amino acid portion of the stem region, which links domain III to the C-terminal transmembrane anchor in class II fusion proteins, could be modeled, spanning 5 nm from the end of domain III to within approximately 1 nm of the tip of domain II. The stem forms trimer contacts, adding a β-strand to domain II of a different subunit, as seen in RVFV $G_{C(27)}$. The overall configuration bears strong similarity to other viral and cellular class II fusion proteins

180   including, in order of decreasing similarity: alphavirus E1 proteins (*21*); EFF-1/AFF-1 cell-cell fusion proteins from *C. elegans (28)*; HAP2 gamete fusion proteins from protozoa (*29-31, 40*) and plants (*30*); and flavivirus E proteins (*41*) (**Fig. 2e**). Despite these structural similarities, the only proteins or domains of known structure with detectable amino acid sequence similarity to Atlas G_C (*E*-value <1 in PSI-BLAST (*33*)) are the phlebovirus G_C proteins (22-24% sequence

185   identity; **Supplementary Fig. 5**).

       Atlas G_C has the same structural features that distinguish phlebovirus glycoproteins from other class II fusion proteins: a larger number of disulfide bonds, ten of which are conserved in phlebovirus G_C sequences but not in other class II proteins; N-linked glycosylation in domain III; and a more extensive and rigid interface between domains I and II (**Fig. 2d-e; Supplementary**

190   **Fig. 5**). The most notable differences between Atlas virus and phlebovirus G_C structures are differences in the disulfide bonding pattern and in the composition of side chains lining the glycerophospholipid (GPL) headgroup binding pocket conserved in arboviral class II fusion proteins (*27*). We discuss these differences and their potential functional implications below. Atlas G_C also has a different glycosylation pattern, with a single predicted N-linked glycosylation

195   site at Asn414 in domain III with a weak corresponding feature in the density map

(**Supplementary Fig. 6**). In contrast, phlebovirus G_C proteins contain two N-linked glycans in domain III, at two different sites. One of these, Asn1035 in RVFV, covers the fusion loop in the prefusion conformation of RVFV G_C and stabilizes the prefusion dimer by forming contacts across the dimer interface (*19*), as also seen in flavivirus E proteins (*22, 23*). This glycosylation site is conserved in DABV and HRTV G_C but absent in Atlas G_C. Despite these minor differences, the striking overall structural similarity of Atlas G_C to phlebovirus G_C proteins in the postfusion conformation reveals a surprisingly strong evolutionary link between nematode ERVs from the *Belpaoviridae* family and the fusion proteins of phleboviruses.

Figure 2. Cryo-EM structure of a phlebovirus G$_C$-like Env fragment from *A. ceylanicum* Atlas virus.

(a) Domain organization of Atlas virus G$_C$. Domains are colored as in **Fig. 1a**.

(b) 3D density map of the cryo-EM image reconstruction of a soluble ectodomain fragment of Atlas virus G$_C$ at 3.76 Å overall resolution. The purified protein was a homotrimer (**Supplementary Fig. 2**) and threefold (C3) symmetry was imposed. The map is colored by domain as in **(a).** A representative cryo-EM micrograph is shown in **Supplementary Fig. 4a.**

(c) Overview of the refined atomic model of the Atlas virus G_C trimer. Atlas G_C has the fold of a class II fusion protein in its postfusion conformation. The protomer in the foreground is shown as a cartoon and the two other protomers in surface representation. Disulfide bonds (green) and an N-linked glycan (blue) are shown as sticks.

(d) Protein fold of single Atlas G_C protomer.

(e) The structures most similar to Atlas G_C are G_C glycoproteins from Rift Valley Fever virus (RVFV, Rmsd 2.6 Å, Z-score (*42*) 29, PDB:6EGU (*27*)), Dabie bandavirus (DABV, formerly SFTS phlebovirus; Rmsd 2.6 Å, Z-score 28, PDB:5G47 (*37*)) and Heartland virus (HRTV, Rmsd 2.6 Å, Z-score 28, PDB:5YOW (*38*)). Other representative class II fusion proteins are shown for comparison: Semliki Forest virus (SFV) E1 (Z-score 19, PDB:1RER (*26*)), *C. elegans* EFF-1 (Z-score 19, PDB:4OJD (*28*)), *Chlamydomonas reinhardtii* HAP2 (Z-score 16; PDB:5MF1 (*29*)) and dengue virus (DENV) E (Z-score 15, PDB:4GSX (*43*)).

Structure of the putative lipid membrane anchor of Atlas virus G_C

Viral fusion proteins insert a membrane anchor – the fusion loop, in class II proteins – into the host cell membrane to initiate virus-cell membrane fusion. The putative fusion loop of Atlas G_C can be clearly identified by analogy to phlebovirus G_C proteins as spanning residues 127-140. The local resolution of the cryo-EM density for this region is lower than for the rest of the map but the large number of structural constraints imposed by the positions of disulfide-bonded cysteines and other residues conserved in phleboviruses G_C proteins allowed an atomic model to be built unambiguously (**Fig. 3a; Supplementary Fig. 6b,c**). Specifically, the structure of the fusion loop is constrained by the positions of four disulfide bonds conserved in phleboviruses, a fifth disulfide specific to Atlas G_C (Cys129-Cys138), a phenylalanine (Phe136) required in phleboviruses at the apex of the fusion loop for membrane binding and fusion (*27, 37, 44*), and two conserved glycines (Gly128 and Gly134) that provide the torsional flexibility necessary for the fusion loop's tightly folded conformation (**Fig. 3a-c**).

The structure and chemical properties of the Atlas G_C fusion loop are similar to phlebovirus fusion loops and consistent with a membrane anchoring function. The side chains of Ile130, Phe136, Phe137 and the Atlas-specific disulfide (Cys129-Cys138) form a hydrophobic

240   surface at the narrow end of the trimer similar to phlebovirus and flavivirus fusion proteins (*25, 27*) (**Fig. 3c**). The area of this surface, extended by the side chain of Arg89 on an adjacent loop, analogous to Leu779 in RVFV G_C, is greater than in most other viral class II fusion proteins. By analogy with other viral class II fusion proteins, the location and extent of the hydrophobic surface formed by the Atlas G_C fusion loop suggest it could function as a membrane anchor.

245   In addition to inserting nonpolar side chains into the hydrophobic region of the membrane, viral class II fusion proteins form polar contacts with lipid headgroups via the fusion loop and an adjacent glycerophospholipid (GPL) binding pocket (*27*). By selecting for headgroups with complementary electrostatic potential, polar contacts confer a degree of specificity to lipid binding. In phleboviruses, a set of conserved polar residues in the GPL

250   binding pocket bind selectively to zwitterionic GPLs (*27*). The Atlas G_C structure reveals a putative GPL binding pocket with both conserved and novel features (**Fig. 3**). The arginine that forms bidentate hydrogen bonds with the GPL phosphate moiety in phleboviruses (*27*) is conserved in Atlas virus (Arg86). The disulfide bond and short-chain hydrophobic residue that bind the GPL glycerol moiety are also conserved (Cys87-Cys135, Ala90). However, an

255   aspartate-arginine pair that binds choline and ethanolamine GPL moieties in phleboviruses is not conserved in Atlas G_C, which instead has two histidines at the corresponding positions (His271 and His85). Moreover, Atlas virus has an extra residue in the fusion loop, Asp133, compared to phlebovirus G_C proteins. The side chain of Asp133 points into the GPL binding pocket and is located near the position of the choline GPL moiety in the superimposed structure

260   of RVFV G_C bound to a phosphatidylcholine ligand (*27*), suggesting that Asp133 could compensate for the lack of a conserved aspartate at position 271 (**Fig. 3a-c**). Hence the GPL binding pocket of Atlas G_C appears to have the necessary physicochemical attributes to support

GPL binding, with Arg86 binding the phosphate moiety, Cys87/Cys135/Ala90 binding the glycerol moiety, and His85/Asp133/His271 coordinating the end of the headgroup. We note the presence in the cryo-EM reconstruction of a bulge in the density around the GPL binding pocket that is unaccounted for by the atomic model (**Supplementary Fig. 6c**). Additionally, the absorbance at 260 nm of purified Atlas G_C was higher than expected despite treatment with nucleases during purification (**Supplementary Fig. 2a**). These two observations would be consistent with lipid molecules with unsaturated acyl chains copurifying with Atlas G_C, but the local resolution of the map is insufficient to ascertain whether the GPL binding pocket contains a ligand.

Figure 3. Structure of the putative fusion loop and lipid binding pocket of Atlas G$_C$.

(a) Local alignment of the sequences surrounding the fusion loops of Atlas G$_C$, RVFV G$_C$, and DENV E. Fusion loop sequences are boxed in orange. Disulfides are marked with green dumbbells. Disulfides shown below the RVFV sequence are conserved in Atlas and RVFV.

(b) Residues contributing to binding of the phosphate (Phos), glycerol and choline (Chol) moieties of glycerophospholipid (GPL) molecules by RVFV G$_C$ (*27*). The aligned sequences of G$_C$ from Atlas, Dabie bandavirus (DABV, formerly SFTSV) and heartland virus (HRTV) are

280   shown. Colors indicate side chain properties: blue, positive charge; pink, negative charge;

magenta, hydrophobic; green, sulfhydryl; light blue, positive charge at endosomal pH.

(c) Closeup of the Atlas G_C fusion loop and GPL binding pocket structure. The corresponding

structures of RVFV G_C and dengue E are shown for comparison. A single protomer from the

postfusion trimer is shown, with key residues shown in stick representation. Label colors

285   indicate side chain properties: blue positive charge; red, negative charge; orange, hydrophobic.

(d) Closeup of the fusion loop and GPL binding pocket region of the Atlas G_C trimer showing

histidine residues in or near the pocket. Prime symbols following residue numbers denote the

protomer to which the residue belongs. **(e)** Surface representation of the same view as in (**d**),

colored by side chain hydrophobicity, except for histidine residues shown in light blue.

290   **Atlas G_C binds membranes with endosome-like lipid composition at low pH**

A key step in viral membrane fusion is insertion of the fusion protein into the host cell

membrane. To determine whether Atlas G_C has this activity, we assessed binding to liposomes

in density gradient centrifugation experiments, supported by dynamic light scattering (DLS)

measurements. Viruses containing class II fusion proteins, like many retroviruses, undergo

295   membrane insertion and fusion in endosomal compartments where the pH is acidic. We

therefore assayed liposome binding at a range of pH values. In contrast to RVFV G_C, Atlas G_C

did not bind liposomes containing phosphatidylcholine (PC), phosphatidylethanolamine (PE),

cholesterol and sphingomyelin (SM) at neutral or acidic pH (pH 4-8; **Fig. 4a,b**). At neutral pH

(pH 7.8), Atlas G_C also failed to bind liposomes containing anionic lipids enriched in early or late

300   endosomes: phosphatidylcholine (PS) or bis(monoacylglycerol)phosphate (BMP, also known as

lysobisphosphatidic acid or LBPA), respectively. At pH 4, however, Atlas G_C bound tightly to

liposomes containing PS or BMP, with weaker binding observed at pH 4.6 (**Fig. 4a,b;**

Supplementary Fig. 7). Remarkably, Atlas G_C bound only weakly to liposomes containing

phosphatidylglycerol (PG) instead of BMP even though PG and BMP are regioisomers with

305   identical chemical composition and electrostatic charge (of -1). PG and BMP differ only in the

position of the second acylglycerol linkage, resulting in a linear configuration for BMP instead of

the usual branched configuration for PG. Our liposome binding data show that Atlas G_C binds to

membranes containing specific GPLs that are enriched in the endosomal pathway in a pH-

dependent manner. No other class II fusion protein has been reported to require low pH, PS or

310   BMP for membrane insertion. However, phleboviruses, which require only PE or PC and

cholesterol for membrane insertion (*27*), subsequently require BMP and low pH for fusion (*45*).

Similarly, flaviviruses require BMP, PS or other anionic lipids and low pH for efficient fusion (*46-*

48).

        The optimal pH for membrane insertion of Atlas G_C (pH 4-4.5) is similar to the optimal pH

315   of hemifusion of Uukuniemi virus (*45*), a model phlebovirus, and would be consistent with

membrane insertion in late endosomes or endolysosomes, as is the case for phleboviruses (*49*).

Since the imidazole side chain of histidine has a pK_a value in the range of 6-6.4, the side chains

of His85 and His271, in the GPL binding pocket of Atlas G_C, would be fully protonated at pH 4-

4.5. The resulting net positive charge of the His85/Asp133/His271 triad (+1/-1/+1), analogous to

320   the Arg/Asp pair that coordinates the end of GPL headgroups in phleboviruses, would precisely

mirror the charge of the phosphoserine headgroup of PS (-1/+1/-1), and to a lesser extent the

glycerol-phospho-glycerol headgroup of BMP (-1). Moreover, Atlas G_C contains four additional

histidines (residues 88, 94, 99 and 100) exposed to the solvent in the immediate vicinity of the

GPL binding pocket and fusion loop (**Fig. 3d,e**), none of which are conserved in phleboviruses.

325   The protonation of these histidines at low pH may promote further interactions with anionic

moieties of lipid headgroups, complementing the function of His85/His271. The presence of six

histidines in and around the GPL binding pocket provides a possible explanation for the

observed pH-dependent insertion of Atlas G_C into membranes containing PS and BMP.

Consistent with a conserved role for the GPL binding pocket in determining lipid specificity of

330   class II fusion proteins, mutations in alphaviruses at a position equivalent to His271 in Atlas G_C,

in the *ij* loop, determine the extent to which alphaviruses depend on cholesterol for membrane

binding (*50*).

Figure 4. Atlas G$_C$ binds liposomes in a lipid- and pH-dependent manner.

(a) Liposome coflotation lipid binding assay. A liposome-G$_C$ mixture in 40% OptiPrep density

gradient medium was overlaid with a 30% OptiPrep cushion and centrifuged at 100,000 g.

Flotation was defined as the amount of Atlas G_C co-floating with liposomes in the top-half

fraction (T) divided by the total amount of Atlas G_C in the top- and bottom-half (B) fractions.

Atlas G_C was quantified by Coomassie-stained SDS-PAGE.

340 **(b)** Binding of Atlas G_C to liposomes measured by dynamic light scattering (DLS). Liposome

diameter was measured by DLS in the presence and absence of Atlas G_C.

(c) Binding of Atlas G_C F136A/F137A to liposomes measured by flotation and DLS as above.

(d) Binding of Atlas G_C R86A to liposomes measured by flotation and DLS as above. See

Supplementary Dataset 1 for source data and **Supplementary Fig. 8** for uncropped gels for all

345 replicates. Error bars for flotation data show the standard deviation with three replicates except

PC:PE:SM:CH:PS + WT (two replicates) and PC:PE:SM:BMP + WT (four replicates); error bars

for DLS data show the standard deviation of three to seven replicates. Significance was

determined by 2-way ANOVA analysis of the mean change in liposome diameter, using Sidak's

multiple comparisons test with a 95% confidence interval in GraphPad Prism 8 (see

350 **Supplementary Fig. 7**). **, $0.001 < p < 0.01$; *, $p < 0.05$; ns, not significant.

Atlas G_C membrane insertion does not strictly require cholesterol and occurs via the

fusion loop

In addition to GPLs, phleboviruses and alphaviruses (but not flaviviruses) require cholesterol for

efficient membrane binding and subsequent fusion (*27, 45, 50*). Alphaviruses additionally

355 require sphingolipids (such as SM) for efficient fusion (*51, 52*). We found that neither cholesterol

nor SM were required for Atlas G_C to bind liposomes (**Fig. 4a,b**). Removal of cholesterol

reduced the fraction of G_C bound by 50% in the liposome flotation assay (**Fig. 4a**) though

binding was still detected in the DLS assay (**Fig. 4b**). Hence, although cholesterol and SM are

not strictly required for binding, they enhance binding, possibly by increasing membrane fluidity.

360 Notably, the concentration of cholesterol in nematode cell membranes is approximately 20 times

lower than in vertebrates (*53, 54*). This is insufficient for cholesterol to regulate the structure or

fluidity of nematode membranes, in which cholesterol is thought to be instead a precursor for

low-abundance metabolites (*53-55*). Likewise, *Drosophila* can grow indefinitely with only trace

amounts of exogenous sterols (*53*), suggesting that arthropods, which are obligate vectors of

365   the vast majority of viruses containing class II fusion proteins, rely on lipids other than

cholesterol to regulate membrane fluidity.

In the structure of Atlas G_C, the hydrophobic residues Ile130 and Phe137 in the fusion

loop occupy the positions of more hydrophilic side chains (Trp821 and Asn827 in RVFV G_C),

which are conserved across phleboviruses and flaviviruses and have been implicated in

370   cholesterol binding. Indeed, molecular dynamics simulations suggested that the side chains of

Trp821 and Asn827 in the RVFV fusion loop form hydrogen bonds with cholesterol upon

membrane insertion (*27*). Moreover, the N827A RVFV G_C mutant lacks fusion activity (*44*), and

the N827A and W821H mutants both fail to bind membranes (*27*). We speculate that the

observed ability of Atlas G_C to bind GPL membranes lacking cholesterol – a functionality

375   consistent with the low cholesterol abundance in nematodes – may be due to its more

hydrophobic fusion loop surface, with Ile130, Phe137 and the Cys129-Cys138 disulfide

complementing the conserved Phe136.

To determine whether Atlas G_C binds membranes in a manner analogous to other viral

class II fusion proteins – via the fusion loop and GPL binding pocket – we generated Atlas G_C

380   variants with mutations in the two phenylalanine residues in the fusion loop (F136A/F137A), or

in the arginine predicted to bind the GPL phosphate moiety (R86A). The F136A/F137A mutant

failed to bind liposomes containing PS and coflotation with liposomes containing BMP at pH 4

and pH 4.6 was reduced to approximately one third of wild-type (**Fig. 4c**). The R86A mutation

reduced binding to liposomes containing PS or BMP at pH 4, to one third and 75% of wild-type,

385   respectively (**Fig. 4d**). For both variants, preparations contained trimers as the major species

but a small monomeric fraction was also present (**Supplementary Fig. 2**), suggesting that the

mutated lipid binding residues are required for efficient trimer assembly. We conclude that Atlas

G_C binds to lipid membranes through insertion of hydrophobic fusion-loop residues and

coordination of lipid headgroups in the GPL binding pocket, as in other viral class II fusion

390   proteins.

Evidence for monomeric and trimeric states of Atlas G_C

Class II fusion protein ectodomains can be monomeric, dimeric or form icosahedral shells in

their prefusion conformation, but the fusogenic conformational change is always accompanied

by reorganization into trimers (*25, 45, 56, 57*). Fusion proteins from classes I and III, including

395   retrovirus fusogens, remain trimeric throughout the fusion reaction, but no class II fusion

proteins are known to be trimeric in their prefusion conformation. Having established that Atlas

G_C can insert into membranes as a trimer with a postfusion-like conformation, we set out to

determine whether it could undergo a conformational change as seen in the fusion reaction of

class II proteins from infectious viruses. The Atlas G_C construct described above was expressed

400   as a trimer with no trace of monomers or dimers (**Fig. 5a, Supplementary Fig. 2a**). However,

we found that a construct with the stem region truncated, G_C(DI-III), spanning residues 2330-

2751, was expressed as a mixture of monomers, trimers and higher order oligomers (**Fig. 5b,**

Supplementary Fig. 2b). In contrast to G_C trimers, which were stable at different protein

concentrations and pH values, G_C(DI-III) monomers were unstable over time. As noted above,

405   monomeric fractions were also present in preparations of the fusion loop mutant (F136A/F137A)

and GPL binding pocket mutant (R86A) (**Supplementary Fig. 2c,d**). Whether these monomeric

species are in a prefusion conformation remains to be determined, but the presence of

metastable monomers and stable trimers recapitulates a key property of class II fusogens from

infectious viruses.

410   **pH-dependent stabilization of the Atlas G_C trimer by protonated histidine residues**

The increase in positive charge resulting from histidine protonation is an important part of the

pH sensing mechanism of viral class II fusion proteins. Protonation of conserved histidines at

the domain I-domain III interface of alphavirus, flavivirus and phlebovirus glycoproteins promotes the fusogenic conformational change by destabilizing the prefusion conformation and

415   stabilizing the postfusion conformation (*19, 27, 58-61*). For example, histidines in domain III of phlebovirus G_C proteins form interprotomer salt bridges with negatively charged side chains in the postfusion trimer and mutation of one such histidine in RVFV, His1087, renders the virus uninfectious (*62*). Similarly, in Atlas G_C His258, His333 and His336 form interprotomer or interdomain polar contacts (**Fig. 5c**). By stabilizing the trimeric postfusion-like conformation of

420   Atlas G_C specifically at low pH these histidine-dependent contacts would reduce the reversibility of any preceding conformational change in the protein in acidic endosomal compartments. The striking parallels in how Atlas G_C and class II fusogens from infectious viruses respond at the ultrastructural level to environmental cues support the hypothesis that Atlas G_C would have membrane fusion activity in late endosomes, like phleboviruses and many retroviruses.

425   **Fifteen disulfide bonds stabilize Atlas G_C in its postfusion-like conformation**
With 30 cysteine residues forming 15 disulfide bonds, Atlas G_C contains twice the average abundance of cysteines, more than has been found in any other class II protein. 20 of these cysteines form disulfides that are structurally conserved in phleboviruses (including RVFV, DABV and HRTV) but not in other class II proteins (**Supplementary Fig. 5**). An eleventh

430   disulfide, in domain III, is conserved in Atlas virus, DABV and HRTV but not RVFV. However, Atlas G_C contains four additional disulfides not conserved in phleboviruses: one in the fusion loop, two in domain II in the β-hairpin containing the *ij* loop (one of the cysteines forming these disulfides is conserved in phleboviruses but forms a disulfide with a cysteine in a different β-strand in domain II), and one in domain III (**Figs. 2c, 3a; Supplementary Fig. 5**). The Atlas-

435   specific disulfide in the fusion loop, Cys129-Cys138, appears likely to confer novel biochemical properties. As discussed above, the side chains of Cys129 and Cys138 extend the hydrophobic surface formed by conserved residues in the fusion loop that are required for membrane

insertion. Furthermore, due to the location of Cys129 and Cys138 close to the threefold

symmetry axis of the trimer, the side chains of the two residues can be rearranged by torsional

440 rotation, without changes to the polypeptide backbone, to form intermolecular disulfides across

the trimer interface, thereby crosslinking all three protomers in the trimer (**Fig. 5d,e**). SDS-

PAGE under non-reducing conditions showed that approximately 15% of protein purified without

reducing agents contained one or more intermolecular disulfide bonds. However, under the

reducing conditions used to purify Atlas G_C for cryo-EM imaging (0.5 mM TCEP), no crosslinked

445 protein was detected (**Fig. 5f**). Exchange of the Cys129-Cys138 bonds from intramolecular to

intermolecular disulfides could provide a novel mechanism to stabilize the Atlas G_C trimer in its

postfusion-like conformation and reduce the reversibility of any preceding conformational

change. Disulfide-mediated crosslinking of the postfusion conformation has not been described

in any viral or non-viral membrane fusion proteins but could potentially function as a mechanism

450 to promote a fusogenic conformational change in Atlas G_C or in infectious viruses with fusion

proteins related to Atlas G_C.

Figure 5. Oligomeric states and disulfide bonding pattern of Atlas G$_C$.

(a) SEC-MALS of Atlas G$_C$ ectodomain (residues 2330-2772). The protein formed stable trimers.

(b) SEC of Atlas G$_C$(DI-III) (residues 2330-2751) is expressed in a mixture of oligomeric states, including monomers, trimers and larger aggregates (void peak). G$_C$(DI-III) monomers were unstable.

(c) His258, His333 and His336 form interprotomer or interdomain polar contacts predicted to stabilize the G$_C$ trimer specifically at pH <6, when the histidine side chains are charged. Residues from different protomers are denoted with a prime symbol.

(d-e) Views along the threefold axis of the G_C trimer without, (**d**), and with, (**e**), intermolecular

disulfides modeled between the cysteines most proximal to the axis, Cys129 and Cys138.

(f) Coomassie-stained SDS-PAGE gels of Atlas G_C purified under non-reducing conditions (NR)

or with 0.5 mM TCEP reducing agent (R). Gel samples were prepared with either non-reducing

465   (NR) or reducing (R) SDS-PAGE loading buffer. Approximately 15% of the Atlas G_C sample

purified under non-reducing conditions contained intermolecular disulfides. Arrowheads indicate

bands corresponding to monomers (blue), dimers (red) and trimers (black).

Atlas G_C retains membrane fusion activity

Having identified the hallmarks of a fusion protein in Atlas G_C, we measured its membrane

470   fusion activity in a cell-cell fusion assay. CHO cells were transfected with plasmids encoding

Atlas G_C or Atlas G_N-G_C. To promote plasma membrane localization and minimize ER retention,

we replaced the predicted transmembrane domain and cytosolic tail of G_C with the C-terminal

transmembrane anchor and cytosolic tail from HLA-A2, known to localize to the plasma

membrane (*63*). Plasmids encoding vesicular stomatitis virus (VSV) G, or no protein, were used

475   as positive and negative controls, respectively. Since Atlas G_C trimers require low pH and BMP

or PS to efficiently bind membranes, we treated transfected cells with exogenous BMP and then

transferred them to pH 4.5 buffered medium to trigger fusion. Similar treatments have been

used previously to measure cell-cell fusion activity of flavi- and alphaviruses (*46*). Confocal light

microscopy with nuclear and plasma membrane stains showed that cells with three or more

480   nuclei were common in cells expressing Atlas G_C following treatment with BMP and pH 4.5,

though less abundant than in cells expressing VSV G, with or without treatment (**Fig. 6a**). Cells

with more than three nuclei were rare in cells expressing Atlas G_N-G_C. To quantify cell-cell

fusion we counted nuclei and multinuclear cells in micrographs. We found that in cells

expressing Atlas G_C, the fraction of multinuclear cells, defined as cells containing two or more

485   nuclei, was 25 ± 3 % following treatment with BMP and pH 4.5, versus 10 ± 0.2 % without

treatment (**Fig. 6b**). Among cells expressing Atlas G_N-G_C, 13 ± 1 % were multinuclear with

treatment, versus 10 ± 1 % without treatment. By comparison, in cells expressing VSV G, 73 ± 8

% of treated cells and 41 ± 8 % of untreated cells were multinuclear. 8 ± 1 % of cells transfected

with empty vector were binuclear with or without treatment, but none contained more than two

490   nuclei (**Supplementary Fig. 9**). We conclude that Atlas G_C has approximately one third of the

membrane fusion activity of VSV G under the treatment conditions tested, which is significant

given that VSV G is considered highly fusogenic and widely as a model fusogen. The Atlas G_N-

G_C construct tested had little fusion activity. The reason for this remains unclear – possible

explanations include incomplete proteolytic processing at the G_N-G_C junction, lower expression

495   level, or absence of a priming factor in CHO cells.

Figure 6. Cell-cell fusion assay with cells transfected with Atlas G$_C$ or G$_N$-G$_C$.

(a) Confocal micrographs of CHO cells expressing Atlas G$_C$ fused to the C-terminal

transmembrane anchor from HLA-A2, or VSV G. Cells were treated with pH 4.5 buffer

containing BMP, or pH 7.4 containing no exogenous lipid. Blue, Hoechst 33342 nuclear stain;

Magenta, CellBrite Red plasma membrane dye. Micrographs were selected to show fusion

events – see **Supplementary Fig. 9** for representative raw micrographs.

(b) Confocal micrograph of a multinucleated CHO cell expressing Atlas G_C following treatment

with pH 4.5 and BMP. Blue, Hoechst 33342; magenta, CellBrite Red; green, α-HA antibody for

505   Atlas G_C detection. All scale bars in (**a**) and (**b**) are 10 µm.

(c) The fraction of multinuclear cells transfected with plasmids encoding Atlas G_C, Atlas G_N-G_C,

VSV G or no protein (pHLsec empty vector control) was calculated by counting mono- and

multinucleated cells in confocal micrographs, using the Hoechst and CellBrite stains. Error bars

represent standard deviation between measurements, n = 3. See **Supplementary Dataset 2** for

510   source data.

Transcriptional activity of Atlas virus

The transcription of endogenous retroviruses is tightly regulated to prevent harmful gene

expression and genome damage from transposition events. However, some ERVs escape

repression and are highly transcribed, including a subset of belpaoviruses from the parasitic

515   trematode *Schistosoma mansoni* (blood fluke) (*64*). As a first step to assess the expression

level of Atlas virus in *A. ceylanicum*, we analyzed transcript levels of Atlas and other intact

belpaoviruses in previously published transcriptional profiling (RNA-Seq) data (*65*). Transcript

abundance of the nine belpaovirus ERVs encoding complete and intact polyproteins (**Fig. 1a**)

varied, ranging from undetectable to above the median value for all annotated genes (**Fig. 7,**

520   **Supplementary Fig. 10**). For Atlas virus, transcript levels varied across developmental stages:

expression was above the median for annotated genes at the larval L3 stage, below the first

quartile at stage L4, and slightly below the median in adult hookworms (**Fig. 7a**). More precisely,

Atlas virus transcript expression increased 15-fold from the L4 to adult (mixed sex) stages

(adjusted *p*-value <0.05, see **Supplementary Dataset 3** for differential expression calculations).

525   Because the L3 sample was not replicated, statistical significance could not be established for

the differences in expression between L3 and other stages. RNA-Seq reads coverage over the Atlas virus gene was greater in mixed sex adults than in L4 larvae (**Fig. 7b**). In contrast, the difference in Atlas transcription from L4 to male adults was not statistically significant. This suggests the increase in expression from L4 to mixed sex adults was primarily driven by

530 increased expression in female tissues or unlaid eggs. In conclusion, Atlas virus is transcriptionally active at specific stages of hookworm development.

Figure 7. Transcriptional activity of Atlas virus in *A. ceylanicum* hookworm.

(a) Boxplot representation of RNA-Seq expression in Reads Per Kilobase of transcript per

535 Million mapped reads (RPKM) for all annotated genes. The RPKM values for Atlas virus and two other representative intact *A. ceylanicum* belpaovirus ERVs at different developmental stages (L3 and L4 larva, adult male and adult mixed sex) are highlighted. RNA-Seq data are from Bernot *et al. (65).* Trendlines were added to aid interpretation.

(b) Distribution of raw RNA-Seq reads over the Atlas virus (*Acey_s0020.g108*) coding sequence

540 for the Adult (mixed sex) and L4 developmental stages. A snapshot of the same region from the Integrative Genomics Viewer (IGV) with individual unmerged BAM files is shown in **Supplementary Fig. 10**.

Discussion

Here, we identify Atlas virus as an intact ERV with novel features in the human hookworm *A. ceylanicum*. The cryo-EM structure of the Atlas Env reveals a class II viral fusion protein fold remarkably similar to that of the G_C glycoprotein from Rift Valley Fever virus. Atlas G_C has the hallmarks of an active class II membrane fusion protein: a stable trimeric assembly, a putative

550 fusion loop, membrane insertion triggered by low pH with specificity for late endosomal lipid composition, and membrane fusion activity. Moreover, Atlas virus mRNA is expressed at specific stages of hookworm development.

The envelope proteins from retroviruses, including ERVs, that have been biochemically characterized were all found to be class I fusion proteins with an α-helical coiled-coil as the core

555 fold. Viral class II fusion proteins have so far been found only in non-integrating RNA viruses. Our discovery of an ERV with a functional, phlebovirus-like class II fusion protein that is structurally unrelated to other retrovirus Envs reveals an unexpected degree of structural and genetic plasticity in retroviruses. Our work supports the model first proposed based on phylogenetic studies that the nematode belpaoviruses acquired their Env by horizontal gene

560 transfer from a virus from the *Phenuiviridae* family or a phlebovirus-like ancestor(*13*). It remains unclear whether RNA encoding the phlebovirus-like glycoprotein integrated into the belpaovirus ancestor as mRNA in a splicing event, or by first becoming a substrate for the reverse transcriptase with subsequent genomic integration as dsDNA.

While rare, horizontal gene transfer of atypical fusogens into retroelements is not unique

565 to the belpaoviruses. The Tas element from the nematode *Ascaris lumbricoides*, the most common parasitic worm in humans, has an *env* gene with weak genetic similarity to herpesvirus gB proteins (*13*), which have a class III fusion protein fold (also found in *Rhabdoviridae* and *Baculoviridae*). Taken together, these findings lead us to hypothesize that acquisition of a fusion protein from an infectious virus represents a general paradigm for how retrotransposons can

570   become retroviruses, and indeed how ancestral reverse-transcribing viruses may have

originated.

The *env* gene is often first element to be lost in ERVs, as it is not required for

intracellular proliferation, so it is notable that the G_N-G_C *env* module is intact in Atlas virus. With

its identical LTRs, and no stop codons or frameshift mutations, Atlas virus shows all the signs of

575   being intact and recently active. This supports the notion that the envelope may be functional

(*11*). The preserved biological activities of Atlas G_C suggest these activities could have cellular

functions in health and disease, as reported for a small but increasing number of other ERV *env*

and *gag* gene products (*1, 3-5*). Further studies are required to determine the full extent to

which protein expression from transposable elements – and its dysregulation – contribute to

580   basic cellular functions, embryonic development and disease outcomes. This work provides a

blueprint for such efforts.

Materials and Methods

Sequence analysis of *A. ceylanicum* Atlas virus. A PSI-BLAST search (*33*) for protein

585   sequences similar to biochemically characterized phlebovirus fusion proteins identified the gene

Acey_s0020.g108 (UniProt A0A016UZK2; Genbank JARK01001356.1, genomic translation

EYC20859.1) in the human hookworm *Ancylostoma ceylanicum* as the most similar sequence

outside infectious virus taxa, with an Expected value (E-value) of 10^{-20} against the RVFV G_C

sequence. A second iteration performed using position-specific scoring matrix based on an

590   alignment of sequences identified in the first iteration gave an E-value of 10^{-144}.

Protein expression and purification. Synthetic genes encoding soluble ectodomain fragments

of the Env of endogenous retrovirus Y032_0020g108 from *A. ceylanicum* were subcloned into

the pMT/BiP/V5-His vector (ThermoFisher) in frame with the BiP signal sequence and the C-

terminal V5 and six-histidine tags. The constructs referred to here as Atlas G_C and Atlas G_C(DI-

595   III) span amino acids 2330-2772 and 2330-2751 from UniProt A0A016UZK2, respectively. Atlas

G_C mutants were generated by Dpn I-based site-directed mutagenesis. D.mel-2 insect cells

(ThermoFisher) were co-transfected with the expression construct and blasticidin resistance

marker pCoBlast (ThermoFisher) at a 20:1 molar ration and cultured for 6 weeks in 0.5 µg ml^{-1}

blasticidin to obtain a population of expressor cells. Expression was induced in a shaking cell

600   suspension at 27˚C with 0.5 mM $CuSO_4$ at a cell density of 5 x 10^6 cells ml^{-1}. The cell culture

medium was harvested 4 to 5 days after induction, centrifuged to remove cells (2,000 g) and cell

debris (17,000 g), filtered with a 0.2 µm filter, concentrated by tangential-flow filtration, and

buffer exchanged into 20 mM Tris pH 7.8, 0.3 M NaCl, 5% glycerol, 20 mM imidazole, 0.5 mM

TCEP (tris(2-carboxyethyl)phosphine). Atlas G_C was purified by nickel-affinity chromatography

605   with a HisTrap Excel column (Cytiva), followed by anion-exchange chromatography with a

MonoQ or Resource Q column (Cytiva) using 20 mM Tris pH 8.0, 50 mM NaCl, 5% glycerol, 0.5

mM TCEP as the binding buffer and binding buffer plus 1 M NaCl as the elution buffer. Peak

fractions were concentrated and further purified by size-exclusion chromatography (SEC) with a

Superdex 200 Increase (10/300) column (Cytiva) in 20 mM Tris pH 7.8-8.0, 0.15 M NaCl, 5%

610   glycerol and 0.5 mM TCEP. The C-terminal V5 and histidine tags were optionally cleaved by

incubation with carboxypeptidase A (CPA) for 3 h at 4˚C (1:500 CPA:G_C molar ratio).

Liposome binding assay. 1-palmitoyl-2-oleoyl-sn-glycero-3-phosphocholine (PC), 1-palmitoyl-

2-oleoyl-snglycero-3-phosphoethanolamine (PE), egg sphingomyelin (SM), 1-palmitoyl-2-oleoyl-

sn-glycero-3-phospho-L-serine (PS), 1-palmitoyl-2-oleoyl-sn-glycero-3-phospho-(1'-rac-glycerol)

615   (PG), (S,R) Bis(Monoacylglycero)Phosphate (BMP) (Avanti Polar Lipids), and 1-cholesterol

(Sigma-Aldrich) were dissolved in chloroform. 25 mM lipid solutions were mixed at various molar

ratios and dried under nitrogen gas for over 4 h. The lipid film was resuspended in Liposome

Buffer (20 mM Tris pH 7.8, 0.15 M NaCl, 5% glycerol, 0.5 mM TCEP, 2 mM $MgCl_2$, 2 mM $CaCl_2$,

2 mM KCl) and subjected to five cycles of freeze-thawing in liquid nitrogen, followed by 25

620   cycles of extrusion through two 0.2 µm polycarbonate filter membranes (Whatman). Purified

Atlas G_C was added in a 1:771 protein:lipid molar ratio and incubated at 37˚C for 5 min. The pH

was reduced by adding a 2 M stock solution of sodium acetate pH 4.6 or 4.0 to a final

concentration of 0.2 M. Following a 2 h incubation at 37˚C the pH of the suspension was

neutralized with 1 M Tris pH 8. OptiPrep density gradient medium (Sigma-Aldrich) was added to

625   a concentration of 40%, maintaining 0.15 M NaCl throughout. Approximately 0.5 ml of the

liposome suspension was placed in a centrifuge tube, overlayed with a 2.5 ml cushion of 30%

OptiPrep solution, and centrifuged at 100,000 g for 1 h at 4˚C in a TLA100.3 rotor (Beckman

Coulter). Top (T) and bottom (B) fractions (approximately 1.5 ml each) were collected from the

top meniscus with a micropipette. Atlas G_C was quantified by densitometry of the absorbance at

630   700 nm of bands in Coomassie-stained SDS-PAGE gels with an Odyssey scanner (LI-COR).

Flotation was defined as the amount of Atlas G_C in the top fraction divided by the total amount of

Atlas G_C in both fractions.

For measurement of liposome diameter by dynamic light scattering (DLS), the liposome

suspensions were diluted ten-fold in Liposome Buffer prior to the addition of 40% OptiPrep

635   solution. Following centrifugation liposome diameters were measured in 384-well clear-

bottomed optical imaging plates (Corning) with a DynaPro Plate Reader III (Wyatt

Technologies). The mean diameter was calculated as the average of three independent

measurements, each consisting of fifteen two-second acquisitions. Protein-free acidified

liposome controls were treated and measured in parallel, with Liposome Buffer instead of Atlas

640   G_C solution.

Size-exclusion chromatography and multi-angle scattering (SEC-MALS) analysis. 100 µl

samples containing 1.6 – 2.5 mg ml^{-1} Atlas G_C were analysed by size-exclusion chromatography

(SEC) at 293 K on a Superdex 200 (10/300) column (Cytiva) in 20 mM Tris pH 7.8, 0.15 M

NaCl, 5% glycerol and 0.5 mM TCEP with a flow rate of 0.5 ml min^{-1}. The SEC system was

645　coupled to both multi-angle light scattering (MALS) and quasi-elastic light scattering (QELS)

modules (DAWN-8+, Wyatt Technology). The protein was also detected as it eluted from the

column with a differential refractometer (Optilab T-rEX, Wyatt Technology) and a UV detector at

280 nm (Agilent 1260 UV, Agilent Technology). Molar masses of peaks in the elution profile

were calculated from the light scattering and protein concentration, quantified using the

650　differential refractive index of the peak assuming a dn/dc of 0.1860, with ASTRA6 (Wyatt

Technology).

Cryo-EM sample preparation and data collection. Purified Atlas G_C trimer (3 µl at a

concentration of 0.025 mg ml^{-1}) in 20 mM Tris pH 7.8, 0.15 M NaCl, 5% glycerol and 0.5 mM

TCEP, was applied onto glow-discharged R1.2/1.3 400 mesh copper grids (Quantifoil Micro

655　Tools, Germany). The grids were blotted for 4 s and plunge-frozen in liquid ethane with a

Vitrobot Mark IV (ThermoFisher) at 4˚C and 100% humidity. Preliminary sample screening and

initial datasets were acquired on a FEI Tecnai F20 microscope operated at 200 kV equipped

with Falcon II direct electron detector (ThermoFisher) at -4 µm defocus. High-resolution cryo-EM

dataset collection was performed on a Titan Krios microscope (ThermoFisher) operated at

660　300 kV equipped with a 20 eV slit-width GIF quantum energy-filtered Gatan K2-Summit direct

electron detector in counting mode. A total of 3,027 movies were recorded at a calibrated

magnification of 130,000x leading to a magnified pixel size of 1.047 Å on the specimen. Each

movie comprised 36 frames with an exposure rate of 1.28 e$^-$ Å^{-2} per frame, with a total exposure

time of 8 s and an accumulated exposure of 46.18 e$^-$ Å^{-2}. Data acquisition was performed with

665　EPU Automated Data Acquisition Software for Single Particle Analysis (ThermoFisher) with

three shots per hole at -1.3 µm to -3.5 µm defocus.

Image processing. Micrographs from initial datasets allowed us to obtain a consistent model at

~19 Å resolution from 3,790 particles selected after 2D and 3D classification, and consequent

auto-refinement. All movies from high-resolution datasets were motion-corrected and dose-

670   weighted with MOTIONCOR2 (*66*). Aligned, non-dose-weighted micrographs were then used to

estimate the contrast transfer function (CTF) with GCTF (*67*). All subsequent image processing

steps were performed using RELION 3.0 (*68, 69*). 2D references from initial datasets were used

to auto-pick the micrographs. One round of reference-free 2D classification was performed to

produce templates for better reference-dependent auto-picking, resulting in a total of 987,570

675   particles. After a first round of 2D classification, 595,011 particles were selected to perform a

second 2D classification, resulting in a final number of 320,041 selected particles. Then, a 3D

classification imposing C3 symmetry was performed using the model from the initial datasets

filtered at 40 Å resolution as initial model. The best class, containing 197,145 particles, was

selected and subjected to 3D auto-refinement imposing C3 symmetry, yielding a map with an

680   overall resolution at 4.11 Å based on the gold-standard (FSC = 0.143) criterion. After

refinement, the CTF refinement (per-particle defocus fitting and beam tilt estimation) and

Bayesian polishing (*70*) routines implemented in RELION 3.0 were performed, yielding a final

map with an overall resolution at 3.76 Å. Local resolution was estimated with RELION.

Model building and refinement. The most similar sequence to Atlas G_C with a structure

685   available was glycoprotein G_C from Rift Valley Fever virus (RVFV). The crystal structure of

RVFV G_C in the postfusion conformation (PDB:6EGU (*27*)) was used as template to build a

homology model with the sequence of Atlas G_C using the Swiss-Model server

(https://swissmodel.expasy.org). The output model was docked as a rigid body into the density

with UCSF Chimera (*71*). Initial docking was performed manually and was followed by real

690   space fitting with the Fit in Map routine. A preliminary step of real space refinement was

performed on the three-subunit model, with Phenix 1.13 (*72*), with global minimization, atomic

displacement parameter (ADP), simulated annealing and morphing options selected. The model

was then rebuilt in Coot (*73*) to optimize the fit to the density. Due to low resolution information

in the fusion loop region, the density was converted to .mtz file using CCP-EM software

package tools and blurring of the density allowed us to localize bulky residues and disulphide bonds, and thus use them as a guide to build the entire fusion loop. A final step of real space refinement was performed with Phenix 1.15, with global minimization and ADP options selected. The following restraints were used in the real space refinement steps: secondary structure restraints, non-crystallographic symmetry (NCS) restraints between the protein subunits, side

700   chain rotamer restraints, and Ramachandran restraints. Key refinement statistics are listed in **Supplementary Table 1**.

Model validation and analysis. The FSC curve between the final model and full map after post-processing in RELION, Model vs Map, is shown in **Supplementary Fig. 3**. Cross-validation against overfitting was performed as described (*74*). The atoms in the final atomic model were

705   displaced by 0.5 Å in random directions with Phenix. The shifted coordinates were then refined against one of the half-maps generated in RELION, the "work set". This test refinement was performed in Phenix using the same procedure as for the refinement of the final model (see above). The other half-map, the "test set" was not used in refinement for cross-validation. FSC curves of the refined shifted model against the work set, FSCwork, and against the test set,

710   FSCtest, are shown in **Supplementary Fig. 3**. The FSCwork and FSCtest curves are not significantly different, consistent with the absence of overfitting in our final models.

The quality of the atomic models, including basic protein geometry, Ramachandran plots, clash analysis, was assessed and validated with Coot, MolProbity (*75*) as implemented in Phenix 1.15, and with the Worldwide PDB (wwPDB) OneDep System (https://deposit-

715   pdbe.wwpdb.org/deposition).

Cell-cell fusion assay. Chinese hamster ovary (CHO) Lec3.2.8.1 cells were transfected with pHL-Sec plasmids encoding ectodomain fragments of Atlas G_N-G_C (polyprotein residues 1909-2795) or Atlas G_C (residues 2330-2795) fused to the C-terminal transmembrane domain from

HLA-A2 (residues 288-345) and cloned in frame with the vector's secretion signal and a C-

720   terminal HA tag. Empty pHL-Sec plasmid, and pcDNA encoding vesicular stomatitis virus (VSV)

G, were used as negative and positive controls, respectively. 16-20 h post-transfection, cells

were transferred to PBS supplemented with 2.5 mM BMP (18:1(S,S)

Bis(Monoacylglycero)Phosphate (BMP); Avanti Polar Lipids). To obtain a homogeneous BMP

suspension, the mixture was freeze-thawed five times using liquid nitrogen and a water bath,

725   followed by a 3 min incubation in a sonicating water bath. Cells were incubated in the BMP

suspension (or PBS for the untreated control) at 37˚C for 5 min, shown previously to be

sufficient for anionic lipid incorporation into the plasma membrane (46). Cell were transferred to

pH 4.5 complete medium (DMEM adjusted to pH 4.5 with HCl supplemented with 10% foetal

bovine serum), or pH 7.4 complete medium for the untreated control, and centrifuged at 2,500 g

730   at 37˚C for 2 min. Cells were immediately resuspended in complete media and plated out.

Following reattachment (4-6 h after treatment), cells were stained with Hoechst 33342 (Bio-Rad)

and CellBrite Red cytoplasmic membrane dye (Biotium, cat. no. 30023), and imaged on a Nikon

iSIM Swept Field inverted confocal microscope with a 60x/1.2NA water objective. Cluster

analysis of the Hoechst channel with Fiji (76) was used to count single nuclei and identify

735   polynuclear clusters. Nuclei within polynuclear clusters were counted by visual inspection. The

plasma membrane stain was used to confirm polynuclear clusters and count the number of

multinuclear cells, defined as cells with two or more nuclei, by visual inspection. The fraction of

multinucleated cells (F) was calculated using the formula:

$$F \; = \; 1 \; - \; (n.\,mononucleated\;cells) \; / \; (total\;n.\,nuclei)$$

740   **RNA-Seq analysis.** We analysed data published by Hawdon and colleagues (65).

ArrayExpress run accession numbers and corresponding life cycle stages were: SRR6359160,

L3; SRR6359161 and SRR6359163, L4; SRR6359164 and SRR6359165, Adult mixed pooled

worms; SRR6359162 and SRR6359166, Adult male pooled worms. Our mapping and counting

strategy was modified as recommended for repetitive genomic features (*77*). Reads were

745   mapped against the reference *A. ceylanicum* genome (*78*) with STAR v2.7.5a (*79*) with

parameters: "--outSAMtype BAM SortedByCoordinate --runMode alignReads --

outFilterMultimapNmax 1000 --outSAMmultNmax 1 --outFilterMismatchNmax 3 --

outMultimapperOrder Random --winAnchorMultimapNmax 1000 --alignEndsType EndToEnd --

alignIntronMax 1 --alignMatesGapMax 350". FeatureCounts v2.0.1 (*80*) with parameters "-M -F

750   GFF -s 0 -p -t exon -g gene_id" was used to count reads over a modified annotation file based

on the original annotation (*78*) and downloaded from Wormbase Parasite (*81*). The original

annotation file in GFF format was slightly modified to contain a "gene_id" feature in the ninth

column and hence facilitate the calculation of aggregate reads by gene (for more details see

header of the GFF file, **Supplementary Dataset 4**). The resulting table of counts was imported

755   into R (*82*) and processed for differential expression analysis using DESeq2 v1.26.0 (*83*). Plots

were generated using ggplot2 (*84*) and karyoploteR (*85*). All scripts and instructions for how to

use them for RNA-Seq analysis are available from Github

[https://github.com/annaprotasio/Merchant_et_al_2020].

Statistics. Error bars represent the standard deviation or standard error –as indicated in the

760   respective figure legend– of 2-7 replicates conducted across at least two independent

experiments. SDS-PAGE gels and DLS data shown are representative of at least two

independent experiments. Significance was determined by 2-way ANOVA analysis using

Sidak's multiple comparisons test with a 95% confidence interval, in Prism 8 (GraphPad). **,

$0.001 < p < 0.01$; *, $p < 0.05$; ns, not significant. Source data are provided in **Supplementary**

765   **Datasets 1 and 2.** No statistical methods were used to predetermine sample size, experiments

were not randomized, and the investigators were not blinded to experimental outcomes.

Data availability

The atomic coordinates were deposited in the Protein Data Bank with code 7A4A

[https://doi.org/10.2210/pdb7A4A/pdb]. The cryo-EM density was deposited in the EM Data

Bank with code EMD-11630 [https://www.ebi.ac.uk/pdbe/entry/emdb/EMD-11630]. The raw

electron micrographs were deposited in EMPIAR with code EMPIAR-10266

[https://dx.doi.org/10.6019/EMPIAR-10266]. RNA-Seq analysis scripts and instruction are

available from Github [https://github.com/annaprotasio/Merchant_et_al_2020]. Other data are

available from the corresponding author upon reasonable request.

References

1. M. Friedli, D. Trono, The developmental control of transposable elements and the evolution of higher species. *Annu Rev Cell Dev Biol* **31**, 429-451 (2015).

2. E. B. Chuong, N. C. Elde, C. Feschotte, Regulatory evolution of innate immunity through co-option of endogenous retroviruses. *Science* **351**, 1083-1087 (2016).

3. J. A. Frank, C. Feschotte, Co-option of endogenous viral sequences for host cell function. *Curr Opin Virol* **25**, 81-89 (2017).

4. A. Dupressoir, C. Lavialle, T. Heidmann, From ancestral infectious retroviruses to bona fide cellular genes: role of the captured syncytins in placentation. *Placenta* **33**, 663-671 (2012).

5. R. S. Moore *et al.*, Horizontal and vertical transmission of transgenerational memories via the *Cer1* transposon. *bioRxiv*, 2020.2012.2028.424563 (2020).

6. T. Gojobori, S. Yokoyama, Rates of evolution of the retroviral oncogene of Moloney murine sarcoma virus and of its cellular homologues. *Proc Natl Acad Sci U S A* **82**, 4198-4201 (1985).

7. K. R. McCarthy *et al.*, The ancient fusogen EnvP(b)1 is expressed in human tissues and its structure informs the evolution of gammaretrovirus envelope proteins. *bioRxiv*, 2020.2004.2022.056234 (2020).

8. D. Kremer *et al.*, Human endogenous retrovirus type W envelope protein inhibits oligodendroglial precursor cell differentiation. *Ann Neurol* **74**, 721-732 (2013).

9. W. Li *et al.*, Human endogenous retrovirus-K contributes to motor neuron disease. *Sci Transl Med* **7**, 307ra153 (2015).

10. M. Krupovic *et al.*, Ortervirales: New Virus Order Unifying Five Families of Reverse-Transcribing Viruses. *J Virol* **92**, (2018).

11. I. G. Frame, J. F. Cutfield, R. T. Poulter, New BEL-like LTR-retrotransposons in Fugu rubripes, Caenorhabditis elegans, and Drosophila melanogaster. *Gene* **263**, 219-230 (2001).

12. N. J. Bowen, J. F. McDonald, Genomic analysis of Caenorhabditis elegans reveals ancient families of retroviral-like elements. *Genome Res* **9**, 924-935 (1999).

13. H. S. Malik, S. Henikoff, T. H. Eickbush, Poised for contagion: evolutionary origins of the infectious abilities of invertebrate retroviruses. *Genome Res* **10**, 1307-1318 (2000).

14. D. Fass, S. C. Harrison, P. S. Kim, Retrovirus envelope domain at 1.7 angstrom resolution. *Nat Struct Biol* **3**, 465-469 (1996).

15. D. C. Chan, D. Fass, J. M. Berger, P. S. Kim, Core structure of gp41 from the HIV envelope glycoprotein. *Cell* **89**, 263-273 (1997).

16. W. Weissenhorn, A. Dessen, S. C. Harrison, J. J. Skehel, D. C. Wiley, Atomic structure of the ectodomain from HIV-1 gp41. *Nature* **387**, 426-430 (1997).

17. B. Kobe, R. J. Center, B. E. Kemp, P. Poumbourios, Crystal structure of human T cell leukemia virus type 1 gp21 ectodomain crystallized as a maltose-binding protein chimera reveals structural evolution of retroviral transmembrane proteins. *Proc Natl Acad Sci U S A* **96**, 4319-4324 (1999).

18. S. C. Harrison, Viral membrane fusion. *Virology* **479-480**, 498-507 (2015).

19. M. Dessau, Y. Modis, Crystal structure of glycoprotein C from Rift Valley fever virus. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 1696-1701 (2013).

20. Y. Modis, Relating Structure to Evolution in Class II Viral Membrane Fusion Proteins. *Curr Opin Virol*, DOI:10.1016/j.coviro.2014.1001.1009 (2014).

820   21. J. Lescar *et al.*, The Fusion glycoprotein shell of Semliki Forest virus: an icosahedral assembly primed for fusogenic activation at endosomal pH. *Cell* **105**, 137-148 (2001).

22. F. A. Rey, F. X. Heinz, C. Mandl, C. Kunz, S. C. Harrison, The envelope glycoprotein from tick-borne encephalitis virus at 2 A resolution. *Nature* **375**, 291-298 (1995).

23. Y. Modis, S. Ogata, D. Clements, S. C. Harrison, A ligand-binding pocket in the dengue

825   virus envelope glycoprotein. *Proc Natl Acad Sci U S A* **100**, 6986-6991 (2003).

24. R. M. DuBois *et al.*, Functional and evolutionary insight from the crystal structure of rubella virus protein E1. *Nature* **493**, 552-556 (2013).

25. Y. Modis, S. Ogata, D. Clements, S. C. Harrison, Structure of the dengue virus envelope protein after membrane fusion. *Nature* **427**, 313-319 (2004).

830   26. D. L. Gibbons *et al.*, Conformational change and protein-protein interactions of the fusion protein of Semliki Forest virus. *Nature* **427**, 320-325 (2004).

27. P. Guardado-Calvo *et al.*, A glycerophospholipid-specific pocket in the RVFV class II fusion protein drives target membrane insertion. *Science* **358**, 663-667 (2017).

28. J. Perez-Vargas *et al.*, Structural basis of eukaryotic cell-cell fusion. *Cell* **157**, 407-419

835   (2014).

29. J. Fedry *et al.*, The Ancient Gamete Fusogen HAP2 Is a Eukaryotic Class II Fusion Protein. *Cell* **168**, 904-915 e910 (2017).

30. J. Fedry *et al.*, Evolutionary diversification of the HAP2 membrane insertion motifs to drive gamete fusion across eukaryotes. *PLoS Biol* **16**, e2006357 (2018).

840   31. J. Feng *et al.*, Fusion surface structure, function, and dynamics of gamete fusogen HAP2. *eLife* **7**, (2018).

32. T. Clark, HAP2/GCS1: Mounting evidence of our true biological EVE? *PLoS Biol* **16**, e3000007 (2018).

33. S. F. Altschul *et al.*, Gapped BLAST and PSI-BLAST: a new generation of protein database

        search programs. *Nucleic Acids Res* **25**, 3389-3402 (1997).

34. O. Poch, I. Sauvaget, M. Delarue, N. Tordo, Identification of four conserved motifs among

        the RNA-dependent polymerase encoding elements. *EMBO J* **8**, 3867-3874 (1989).

35. H. Felder *et al.*, Tas, a retrotransposon from the parasitic nematode Ascaris lumbricoides.

        Gene **149**, 219-225 (1994).

36. I. Letunic, P. Bork, Interactive Tree Of Life (iTOL) v4: recent updates and new

        developments. *Nucleic Acids Res* **47**, W256-W259 (2019).

37. S. Halldorsson *et al.*, Structure of a phleboviral envelope glycoprotein reveals a

        consolidated model of membrane fusion. *Proc Natl Acad Sci U S A* **113**, 7154-7159 (2016).

38. Y. Zhu *et al.*, The Postfusion Structure of the Heartland Virus Gc Glycoprotein Supports

        Taxonomic Separation of the Bunyaviral Families Phenuiviridae and Hantaviridae. *J Virol*

        92, (2018).

39. D. J. Leahy, W. A. Hendrickson, I. Aukhil, H. P. Erickson, Structure of a fibronectin type III

        domain from tenascin phased by MAD analysis of the selenomethionyl protein. *Science*

        258, 987-991 (1992).

40. J. F. Pinello *et al.*, Structure-Function Studies Link Class II Viral Fusogens with the

        Ancestral Gamete Fusion Protein HAP2. *Curr Biol* **27**, 651-660 (2017).

41. M. Kielian, F. A. Rey, Virus membrane-fusion proteins: more than one way to make a

        hairpin. *Nat. Rev. Microbiol.* **4**, 67-76 (2006).

42. L. Holm, P. Rosenstrom, Dali server: conservation mapping in 3D. *Nucleic Acids Res* **38**,

        W545-549 (2010).

43. D. E. Klein, J. L. Choi, S. C. Harrison, Structure of a dengue virus envelope protein late-

        stage fusion intermediate. *J Virol* **87**, 2287-2293 (2013).

44. S. Murakami, K. Terasaki, S. I. Ramirez, J. C. Morrill, S. Makino, Development of a novel, single-cycle replicable rift valley Fever vaccine. *PLoS neglected tropical diseases* **8**, e2746 (2014).

45. D. Bitto, S. Halldorsson, A. Caputo, J. T. Huiskonen, Low pH and Anionic Lipid-dependent Fusion of Uukuniemi Phlebovirus to Liposomes. *J Biol Chem* **291**, 6412-6422 (2016).

46. E. Zaitseva, S. T. Yang, K. Melikov, S. Pourmal, L. V. Chernomordik, Dengue virus ensures its fusion in late endosomes using compartment-specific lipids. *PLOS Pathog.* **6**, e1001131 (2010).

47. D. L. Esposito, J. B. Nguyen, D. C. DeWitt, E. Rhoades, Y. Modis, Physico-chemical requirements and kinetics of membrane fusion of flavivirus-like particles. *J Gen Virol* **96**, 1702-1711 (2015).

48. A. M. Nour, Y. Li, J. Wolenski, Y. Modis, Viral membrane fusion and nucleocapsid delivery into the cytoplasm are distinct events in some flaviviruses. *PLoS Pathog* **9**, e1003585 (2013).

49. P. Y. Lozach *et al.*, Entry of bunyaviruses into mammalian cells. *Cell Host Microbe* **7**, 488-499 (2010).

50. M. Umashankar *et al.*, Differential cholesterol binding by class II fusion proteins determines membrane fusion properties. *J Virol* **82**, 9245-9253 (2008).

51. P. K. Chatterjee, C. H. Eng, M. Kielian, Novel mutations that control the sphingolipid and cholesterol dependence of the Semliki Forest virus fusion protein. *J Virol* **76**, 12712-12722 (2002).

52. J. M. Smit, R. Bittman, J. Wilschut, Low-pH-dependent fusion of Sindbis virus with receptor-free cholesterol- and sphingolipid-containing liposomes. *J Virol* **73**, 8476-8484 (1999).

53. T. V. Kurzchalia, S. Ward, Why do worms need cholesterol? *Nat Cell Biol* **5**, 684-688 (2003).

54. M. Merris, J. Kraeft, G. S. Tint, J. Lenard, Long-term effects of sterol depletion in C. elegans: sterol content of synchronized wild-type and mutant populations. *J Lipid Res* **45**, 2044-2051 (2004).

55. M. Ruiz *et al.*, Membrane fluidity is regulated by the C. elegans transmembrane protein FLD-1 and its human homologs TLCD1/2. *eLife* **7**, (2018).

56. K. Stiasny, S. L. Allison, J. Schalich, F. X. Heinz, Membrane interactions of the tick-borne encephalitis virus fusion protein E at low pH. *J. Virol.* **76**, 3784-3790 (2002).

57. D. L. Gibbons, A. Ahn, P. K. Chatterjee, M. Kielian, Formation and characterization of the trimeric form of the fusion protein of Semliki Forest Virus. *J Virol* **74**, 7772-7780 (2000).

58. R. Fritz, K. Stiasny, F. X. Heinz, Identification of specific histidines as pH sensors in flavivirus membrane fusion. *J Cell Biol* **183**, 353-361 (2008).

59. V. Nayak *et al.*, Crystal structure of dengue virus type 1 envelope protein in the postfusion conformation and its implications for membrane fusion. *J. Virol.* **83**, 4338-4344 (2009).

60. Z. L. Qin, Y. Zheng, M. Kielian, Role of conserved histidine residues in the low-pH dependence of the Semliki Forest virus fusion protein. *J Virol* **83**, 4670-4677 (2009).

61. Y. Zheng, C. Sanchez-San Martin, Z. L. Qin, M. Kielian, The domain I-domain III linker plays an important role in the fusogenic conformational change of the alphavirus membrane fusion protein. *J Virol* **85**, 6334-6342 (2011).

62. S. M. de Boer *et al.*, Acid-activated structural reorganization of the Rift Valley fever virus Gc fusion protein. *J Virol* **86**, 13642-13652 (2012).

63. M. A. Vega, J. L. Strominger, Constitutive endocytosis of HLA class I antigens requires a specific portion of the intracytoplasmic tail that shares structural features with other endocytosed molecules. *Proc Natl Acad Sci U S A* **86**, 2688-2692 (1989).

64. R. DeMarco *et al.*, Saci-1, -2, and -3 and Perere, four novel retrotransposons with high transcriptional activities from the human parasite Schistosoma mansoni. *J Virol* **78**, 2967-2978 (2004).

65. J. P. Bernot *et al.*, Transcriptomic analysis of hookworm Ancylostoma ceylanicum life cycle

920   stages reveals changes in G-protein coupled receptor diversity associated with the onset of

parasitism. *Int J Parasitol* **50**, 603-610 (2020).

66. S. Q. Zheng *et al.*, MotionCor2: anisotropic correction of beam-induced motion for improved

cryo-electron microscopy. *Nat Methods* **14**, 331-332 (2017).

67. K. Zhang, Gctf: Real-time CTF determination and correction. *J Struct Biol* **193**, 1-12 (2016).

925 68. S. He, S. H. W. Scheres, Helical reconstruction in RELION. *J Struct Biol* **198**, 163-176

(2017).

69. S. H. Scheres, RELION: implementation of a Bayesian approach to cryo-EM structure

determination. *J Struct Biol* **180**, 519-530 (2012).

70. J. Zivanov, T. Nakane, S. H. W. Scheres, A Bayesian approach to beam-induced motion

930   correction in cryo-EM single-particle analysis. *IUCrJ* **6**, 5-17 (2019).

71. E. F. Pettersen *et al.*, UCSF Chimera--a visualization system for exploratory research and

analysis. *J Comput Chem* **25**, 1605-1612 (2004).

72. P. D. Adams *et al.*, PHENIX: a comprehensive Python-based system for macromolecular

structure solution. *Acta Crystallogr D Biol Crystallogr* **66**, 213-221 (2010).

935 73. P. Emsley, K. Cowtan, Coot: model-building tools for molecular graphics. *Acta Crystallogr D

Biol Crystallogr* **60**, 2126-2132 (2004).

74. A. Amunts *et al.*, Structure of the yeast mitochondrial large ribosomal subunit. *Science* **343**,

1485-1489 (2014).

75. V. B. Chen *et al.*, MolProbity: all-atom structure validation for macromolecular

940   crystallography. *Acta Crystallogr D Biol Crystallogr* **66**, 12-21 (2010).

76. J. Schindelin *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat Methods*

9, 676-682 (2012).

77. A. Teissandier, N. Servant, E. Barillot, D. Bourc'his, Tools and best practices for

retrotransposon analysis using high-throughput sequencing data. *Mob DNA* **10**, 52 (2019).

945   78. E. M. Schwarz *et al.*, The genome and transcriptome of the zoonotic hookworm

Ancylostoma ceylanicum identify infection-specific gene families. *Nat Genet* **47**, 416-422

(2015).

79. A. Dobin, T. R. Gingeras, Mapping RNA-seq Reads with STAR. *Curr Protoc Bioinformatics*

51, 11 14 11-11 14 19 (2015).

950   80. Y. Liao, G. K. Smyth, W. Shi, featureCounts: an efficient general purpose program for

assigning sequence reads to genomic features. *Bioinformatics* **30**, 923-930 (2014).

81. K. L. Howe, B. J. Bolt, M. Shafie, P. Kersey, M. Berriman, WormBase ParaSite - a

comprehensive resource for helminth genomics. *Molecular and biochemical parasitology*

215, 2-10 (2017).

955   82. R. C. Team. (R Foundation for Statistical Computing, Vienna, Austria, 2019).

83. M. I. Love, W. Huber, S. Anders, Moderated estimation of fold change and dispersion for

RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

84. H. Wickham, *ggplot2: Elegant Graphics for Data Analysis*. (Springer-Verlag, New York,

2016).

960   85. B. Gel, E. Serra, karyoploteR: an R/Bioconductor package to plot customizable genomes

displaying arbitrary data. *Bioinformatics* **33**, 3088-3090 (2017).

86. S. Kurtz *et al.*, REPuter: the manifold applications of repeat analysis on a genomic scale.

Nucleic Acids Res **29**, 4633-4642 (2001).

87. M. Krupovic, E. V. Koonin, Homologous Capsid Proteins Testify to the Common Ancestry of

965   Retroviruses, Caulimoviruses, Pseudoviruses, and Metaviruses. *J Virol* **91**, (2017).

88. B. M. Muhire, A. Varsani, D. P. Martin, SDT: a virus classification tool based on pairwise

sequence alignment and identity calculation. *PLoS One* **9**, e108277 (2014).

Acknowledgements

We acknowledge the MRC Laboratory of Molecular Biology Electron Microscopy (EM) and UK national electron Bio-Imaging Centre (eBIC) at Diamond Light Source (DLS) for access and support of EM sample preparation and data collection. We acknowledge DLS for access and support of eBIC under proposals EM17434 and EM20121 funded by the Wellcome Trust, MRC and BBSRC. We are grateful to Kyle Dent for providing assistance in using the microscopes at eBIC. We are grateful for access to the MRC-LMB Scientific Computing facilities. This work was supported by Senior Research Fellowships 101908/Z/13/Z and 217191/Z/19/Z from the Wellcome Trust and NIH grant R01 GM102869-01 to Y.M.

Author contributions

M.M. and Y.M. conceived the experiments. M.M. purified the proteins and performed the biochemical assays. C.P.M collected and processed the cryo-EM data, and performed the image reconstruction. C.P.M and Y.M. built and refined the atomic model. Y.L. and H.Z. carried out the cell-cell fusion assay. A.V.P performed the RNA-Seq analysis. All authors contributed to the figures. Y.M. wrote the manuscript. Project supervision, administration and funding acquisition, Y.M.

Competing interests

Y.M. is a consultant for Related Sciences LLC and has profits interests in Danger Bio LLC.

Materials & Correspondence

Correspondence and material requests should be addressed to Y.M. (ymodis@mrc-lmb.cam.ac.uk).

A bioactive phlebovirus-like envelope protein in a hookworm endogenous retrovirus

Merchant, Mata *et al.*

Supplementary Information

Supplementary Figures 1-10

Supplementary Table 1

Supplementary Fig. 1. Atlas is an intact belpaovirus with a phlebovirus-like Env in the hookworm *A. ceylanicum*

(a) Nucleotide sequence of the Atlas virus LTRs, predicted by REPuter(*86*). The sequences of the two LTRs is 100% identical. **(b)** Aligned protein sequences of reverse transcriptase (RT)

polymerase site from Atlas and other representative belpaoviruses and retroviruses. Endogenous belpaoviruses in *A. ceylanicum* have an aspartate to asparagine substitution
1015 (Y[X]DD to YVDN) in the most conserved reverse transcriptase motif, the polymerase site (Motif V or Motif C). Nucleotide accession numbers are listed after virus names. MoMLV, Murine Moloney leukemia; HERVK, human ERV-K; *Dme*, *D. melanogaster*; *Sce, Saccharomyces cerevisiae*; *Bmo, Bombyx mori*; *Cel, C. elegans*; *Ace, A. ceylanicum*; *Alu, Ascaris lumbricoides*; *Ncr, Neurospora crassa*; *Cre, C. reinhardtii*; *Tbr, Trypanosoma brucei*. **(c)** Matrix of sequence
1020 identities for pairs of Gag protein sequences from representative viruses from the *Belpaoviridae* family and the belpaovirus ERVs from *A. ceylanicum* with intact protein coding regions. *Amy, Antheraea mylitta; Dsi, Drosophila simulans; Aga, Anopheles gambiae; Fru, Fugu rubripes.* With less than 20% Gag protein sequence identity to its closest homolog (*87*) (*Acey_s0020.g1106*), Atlas virus is a candidate for classification as new member of the *Belpaoviridae* family (which
1025 contains a single genus, *Semotivirus*). **(d)** Sequence identity matrix for phlebovirus-like Env protein sequences from nematode belpaoviruses. Identity matrices were calculated with SDT (*88*). **(e)** Phylogenetic tree of phlebovirus-like Env proteins from nematode belpaoviruses. Drawn with Interactive Tree Of Life (iTOL) (*36*).

Supplementary Fig. 2. Size-exclusion chromatography of Atlas G$_C$ following ion-exchange chromatography

Samples were analysed on a Superdex 200 Increase (10/300) column (Cytiva). The A$_{260}$:A$_{280}$ ratios for selected peaks are shown in bold. **(a)** The Atlas G$_C$ fragment used for cryo-EM image reconstruction (residues 2330-2772). The protein had higher absorbance at 260 nm (A$_{260}$) than expected for pure protein. The elution volume was consistent with a homotrimer. **(b)** Atlas G$_C$(DI-III), maintained at pH > 7 throughout expression and purification. Multiple oligomeric states were present including peaks with elution volumes consistent with monomers, trimers and higher order oligomers. **(c)** Atlas G$_C$ fusion loop mutant F136A/F137A. **(d)** Atlas G$_C$ GPL binding pocket mutant R86A.

Supplementary Fig. 3. Image reconstruction quality assessment and workflow
(**a**) Fourier Shell Correlations (FSC) of the reconstruction of Atlas G_C from two independently refined half-maps (map2map, in blue); and of the reconstruction from the whole dataset versus a map calculated from the refined atomic model (model2map, in red). The gold-standard cutoff (FSC = 0.143) and the FSC = 0.5 level are marked with dashed lines. The resolution values of each curve at these levels are indicated. (**b**) FSC plots for cross-validation as described by Amunts *et al.* (*74*). FSCwork (magenta), FSC of refined test model versus work set (half-map used in test refinement). FSCfree (green), FSC of refined test model versus test set (half-map not used in test refinement). The FSC = 0.5 level is indicated by a dashed line. (**c**) Flow chart showing the workflow pipeline for cryo-EM image processing, classification and model refinement, as described in the Materials and Methods.

Supplementary Fig. 4. Representative Cryo-EM micrograph and density with local resolution estimation

(**a**) Representative cryo-EM micrograph of Atlas G_C trimers. (**b**) Local resolution estimation for the cryo-EM volume, calculated in RELION 3.0 (*69*). (**c**) Representative samples of local cryo-EM density from the structure with fitted and refined atomic models. The deposited density map was contoured at 3 σ in Pymol (Schrodinger, LLC).

Supplementary Fig. 5. Protein sequence alignment of Atlas G$_C$ and its closest orthologs
Amino acid sequence alignment of Atlas G$_C$ and the most similar sequences from infectious
viruses: Rift Valley Fever virus (RVFV), Dabie bandavirus (DABV, formerly SFTSV phlebovirus)
and Heartland virus (HRTV). Domains and structural features are marked with lines above the
alignment. GPL, residues in or near the glycerophospholipid headgroup binding pocket. N-glyc.,
N-glycosylation site. Cyt. tail, cytoplasmic tail. Conserved disulfide bonds are numbered in
green. Disulfide bonds specific to Atlas virus are denoted with an "A". An RVFV-specific
disulfide is denoted with an "R". A phlebovirus-specific disulfide is denoted with a "P". The
disulfide marked 11' is conserved in Atlas, DABV and HRTV but not RVFV.

1070

Supplementary Fig. 6. Cryo-EM density for N-linked glycan and GPL pocket of Atlas G$_C$
(a) Cryo-EM density and atomic model of the N-linked glycan at Asn414. The deposited density map was contoured at 2.2 σ in Pymol (Schrodinger, LLC). **(b)** Cryo-EM density and atomic model for the fusion loop region of the Atlas G$_C$ trimer. A B-factor blurring correction of +150 Å^2

1075 was applied to the deposited density map. The resulting map was contoured at 1 σ in Pymol. **(c)** Cryo-EM density in the glycerophospholipid (GPL) headgroup binding pocket unaccounted for by the atomic model. Shown in grey stick representation is the phosphatidylcholine (PC) headgroup bound to RVFV G$_C$ structure (*27*) superimposed on the Atlas G$_C$ structure. A B-factor blurring correction of +163 Å^2 was applied to the deposited density map. The resulting map was

1080 contoured at 1.2 σ in Pymol.

Supplementary Fig. 7. Dynamic light scattering of Atlas G$_C$-liposome complexes
Liposome diameter dynamic light scattering (DLS) data from **Fig. 4** are shown as the mean change in liposome diameter. **(a)** Change in liposome diameter upon binding Atlas G$_C$ ectodomain. **(b)** Change in liposome diameter upon binding Atlas G$_C$ F136A/F137A. **(c)** Change in liposome diameter upon binding Atlas G$_C$ R86A. Error bars show the standard error of the mean (s.e.m.) of three to seven replicates (see **Supplementary Dataset 1** for source data). Significance was determined by 2-way ANOVA analysis using Sidak's multiple comparisons test with a 95% confidence interval, in Prism 8 (GraphPad). **, $0.001 < p < 0.01$; *, $p < 0.05$; ns, not significant.

Supplementary Fig. 8. Uncropped gels from Figures 4 and 5
Gels are shown for all liposome flotation assay replicates (Fig. 4) and non-reducing gels (Fig. 5).

Supplementary Fig. 9. Cell-cell fusion assay with CHO cells
(a) Representative confocal micrographs of CHO cells transfected with plasmid encoding Atlas G_C, Atlas G_N-G_C, VSV G, or no protein (pHLsec empty vector). Cells were treated with pH 4.5 buffer containing BMP, or pH 7.4 buffer without exogenous lipid. Blue, Hoechst 33342 nuclear stain; Magenta, CellBrite Red plasma membrane dye. **(b)** Permeabilized and fixed cells transfected with plasmid encoding G_C stained with anti-HA antibody, to detect the C-terminal HA tag on G_C. Blue, Hoechst 33342; green, anti-HA antibody. **(c)** Left, confocal micrograph of cells transfected with pHLsec plasmid stained with Hoechst 33342. Right, masks from cluster analysis of the Hoechst channel with Fiji (*76*). All scale bars in this figure are 10 μm.

Supplementary Fig. 10. RNA-Seq analysis of intact *A. ceylanicum* belpaovirus ERVs
(a) Boxplot representation of RNA-Seq expression in Reads Per Kilobase of transcript per Million mapped reads (RPKM) for all annotated genes, for Atlas virus, and for the eight other intact *A. ceylanicum* belpaovirus ERVs at the L3, L4 and adult (male and mixed sex) developmental stages. RNA-Seq data are from Bernot *et al. (65).* **(b)** Integrative Genomics Viewer (IGV) snapshot showing the distribution of raw RNA-Seq reads over the Atlas virus (*Acey_s0020.g108*) coding sequence in mixed sex adult and L4 worms with individual unmerged BAM files.

Supplementary Table 1. Cryo-EM data collection, structure determination, model building and refinement parameters and statistics.

	Atlas G$_C$
Data Collection and Processing	
Microscope	FEI Titan Krios
Voltage (kV)	300
Electron exposure (electrons Å^{-2})	46.12
Exposure per frame (electrons Å^{-2})	1.28
Defocus range (μm)	-1.3 to -3.5
Pixel size (Å)	1.047
N. initial particles	987,570
N. final particles	197,145
Map resolution (Å)	3.76
Map resolution range (Å)	3.50 - 5.20
FSC threshold for resolution limit	0.143
Model refinement	
Map sharpening B factor (Å^2)	-180
Symmetry imposed	C3
Mask correlation coefficient	0.78
Model composition	
N. non-hydrogen atoms	10,159
Protein residues (chains A, B, C)	1,320
R.m.s. deviations	
Bond lengths (Å)	0.005
Bond angles (˚)	0.694
Planarity (Å)	0.004
B-factors/ADPs*	
Minimum	35
Maximum	176
Mean	82
Validation (Phenix 1.15)	
MolProbity overall score	2.23
MolProbity all-atom clashscore	5.86
Rotamer outliers (%)	2.28
Ramachandran plot	
% favored	86.2
% allowed	13.8
% outliers	0.0
PDB code	7A4A
EMDB code	EMD-11630
EMPIAR code	10266

ADPs, atomic displacement parameters.