1
2
3
The RNA-binding protein Orb2 is associated with microcephaly and supports centrosome asymmetry in neural stem cells

Beverly V. Robinson[1,2], Junnan Fang[1,3], Dipen S. Mehta[3,4], Joseph Buehler[5], and Dorothy A. Lerit[1]*

Author Affiliations:
1 Department of Cell Biology, Emory University School of Medicine, Atlanta, GA 30322
2 Genetics and Molecular Biology Graduate Program, Emory University, Atlanta, GA 30322
3 These authors contributed equally
4 College of Science and Mathematics, Augusta University, Augusta, GA 30912
5 Biochemistry, Cell, and Developmental Biology Graduate Program, Emory University, Atlanta, GA 30322

* Correspondence: dlerit@emory.edu

Running title: CPEB ortholog Orb2 modulates centrosome activity

Keywords: centrosome, asymmetry, neural stem cell, Orb2, CPEB, RNA-binding protein, post-transcriptional regulation, MTOC

Funding information: This work was supported by NIH grant T32GM008490 (BVR), American Heart Association postdoctoral fellowship 20POST35210023 (to JF) and NIH grant R01GM138544 (to DAL).

Abstract

To maintain a balance of self-renewal versus neurogenesis, neural stem cells (NSCs) undergo

repeated cycles of asymmetric cell division along an invariant polarity axis instructed by

centrosomes. During interphase, the NSC centrosomes are defined by marked asymmetries in

protein composition and functional activity as microtubule-organizing centers. Here, we show a

conserved RNA-binding protein, Orb2, supports centrosome asymmetry in interphase NSCs.

While Orb2 localizes to the active apical centrosome, it promotes the transient inactivation of the

basal centrosome required for centrosome segregation and spindle morphogenesis. Orb2 is

required cell autonomously within NSCs to support centrosome asymmetry and maintenance of

the stem cell pool. Conversely, loss of *orb2* manifests in microcephaly independent of Orb2

function in NSCs. We suggest Orb2 plays opposing roles in centrosome activation and

inactivation, possibly through the translational regulation of multiple mRNA substrates.

Bioinformatics uncovers a significant overlap among RNA targets between *Drosophila* Orb2 and

human CPEB4, consistent with a conserved role for CPEB proteins in centrosome regulation

and neurodevelopment.

53
Introduction

Neural stem cells (NSCs) undergo asymmetric cell division (ACD) along an invariant

apical-basal polarity axis to segregate cell fate determinants, giving rise to two differentially

fated progeny: a self-renewing stem cell and a ganglion mother cell (GMC) fated for neural

differentiation (Doe et al., 1991; Knoblich et al., 1995; Kraut et al., 1996; Broadus and Doe,

1997). This balance in NSC self-renewal is critical for neurogenesis, as its deregulation can lead

to brain tumors or neurodevelopmental disorders, such as microcephaly (Bond et al., 2002;

Cabernard and Doe, 2009). Key to NSC homeostasis are centrosomes, which instruct the

division axis and organize the bipolar mitotic spindle required to segregate the pro-stem and

pro-differentiation cell fate determinants (Cabernard and Doe, 2009; Januschke and Gonzalez,

2010; Wang et al., 2011).

Centrosomes are microtubule (MT)-organizing centers (MTOC) consisting of a central

pair of centrioles surrounded by pericentriolar material (PCM), which recruits the γ-Tubulin

(γTub) ring complex required for MT nucleation (Conduit et al., 2015). Normally, centrosomes

recruit the robust levels of PCM necessary for microtubule-nucleating activity just before mitotic

onset, a process called centrosome maturation (Gould and Borisy, 1977; Khodjakov and Rieder,

1999). Following mitotic exit, centrosomes shed PCM.

In NSCs, however, centrosomes are subject to an asymmetric centrosome maturation

cycle, wherein the apical (daughter) centrosome recruits PCM and organizes MTs, while the

basal-fated (mother) centrosome is transiently inactivated until mitotic onset (Rebollo et al.,

2007; Rusan and Peifer, 2007; Conduit and Raff, 2010; Januschke et al., 2011). NSC

centrosome asymmetry is implicated in apical-basal spindle pole alignment and centrosome

segregation (Januschke and Gonzalez, 2010; Januschke et al., 2013; Lerit and Rusan, 2013;

Ramdas Nair et al., 2016). A basic molecular framework required for NSC centrosome

asymmetry involves asymmetric localization of Centrobin (Cnb) and Polo kinase to the daughter

centrosome in a mechanism also requiring Wdr62 to promote centrosome maturation

(Januschke et al., 2013; Ramdas Nair et al., 2016; Gallaud et al., 2020). Conversely, transient

inactivation of the basal centrosome requires Bld10/Cep135, Pericentrin-like protein (PLP), and

Polo-like kinase 4 (PLK4/SAK) (Lerit and Rusan, 2013; Singh et al., 2014; Gambarotto et al.,

2019). Nevertheless, how centrosome asymmetry is regulated remains incompletely

understood.

Intriguingly, *Cnb*, *Cep135*, *plp*, *polo*, and *Wdr62* mRNAs were identified as putative

mRNA targets for the RNA-binding protein (RBP) Orb2 through an unbiased transcriptomics

study, raising the possibility that Orb2 might regulate centrosome asymmetry in NSCs (Stepien

et al., 2016). Orb2 is a member of the cytoplasmic polyadenylation element binding (CPEB)

proteins orthologous to mammalian CPEB2–4 and implicated in mRNA localization and

translational control (Huang et al., 2006; Keleman et al., 2007; Hafer et al., 2011). Although prior

work supports a role for *orb2* in NSC spindle orientation and neuronal specification, whether

Orb2 contributes to centrosome regulation is unknown (Hafer et al., 2011).

Here, we identify an NSC-autonomous role for Orb2 in establishing centrosome

asymmetry associated with misaligned spindles and NSC loss. We also identify an NSC-

independent role for Orb2 in regulating brain size, as *orb2* loss leads to microcephaly. Finally,

we examine potential targets of Orb2 and propose a revised model of asymmetric centrosome

maturation.

99
Results and Discussion

Orb2 disrupts centrosome activity in interphase NSCs

To determine if Orb2 contributes to centrosome asymmetry, we examined γTub distributions at

apical and basal centrosomes in wild-type (WT) vs. *orb2* null mutant NSCs during late

interphase, when centrosomes are normally asymmetric (Rebollo et al., 2007; Rusan and

Peifer, 2007). To directionally measure protein localization to centrosomes, we calculated an

asymmetry index (AI; *Methods*), wherein a value near 0 indicates symmetric distribution, while

values near 1 or -1 indicate asymmetric enrichment at the apical or basal centrosome,

respectively. Centrosomes visualized with the centriole marker Asterless (Asl; (Varmark et al.,

2007) displayed the expected symmetric distribution among apical and basal centrosomes in

both genotypes (mean AI $\pm$ S.D.= 0.0$\pm$0.2 for WT and 0.1$\pm$0.3 for *orb2*; Fig. 1A–C). In

comparison, γTub was significantly enriched on the apical centrosome in WT NSCs (Fig. 1A, C).

In contrast to WT, *orb2* NSCs showed impaired centrosome asymmetry, evident by increased

γTub localization to the basal centrosome and decreased AI values (Fig. 1B, C). γTub AI was

reduced by over 20% within *orb2* NSCs, as compared to WT (Fig 1C; *p*=0.05 by Kolmogorov-

Smirnov test). Consistently, ~30% of *orb2* NSCs (N= 12/40) had γTub AI values >2 S.D. from

the WT mean (Fig 1D; *p*< 0.0001 by chi-square test). Measuring the levels of γTub localized at

the apical and basal centrosomes revealed a 2.7-fold increase in γTub recruitment to the basal

centrosomes of symmetrized *orb2* NSCs relative to WT (Fig. 1E; *p*<0.0001 by Mann-Whitney

test). We conclude Orb2 primarily promotes centrosome asymmetry during interphase by

blocking the precocious activation of the basal centrosome, although diminished recruitment of

γTub to the apical centrosome may compound this effect.

Orb2-dependent centrosome regulation is cell autonomous

ACD is regulated through intrinsic and extrinsic cellular pathways (Siegrist and Doe, 2006; Doe,

2008). Prior work indicates Orb2 is broadly expressed throughout the larval central nervous

system (Keleman et al., 2007; Hafer et al., 2011). To elucidate if the reduction of centrosome

asymmetry observed in *orb2* mutants arose from a requirement for Orb2 within NSCs, we

depleted *orb2* specifically in NSCs using an *orb2* dsRNA transgene (*UAS-orb2^{RNAi}*) driven by the

NSC-specific *worniu* (*wor*)-*GAL4* (Albertson et al., 2004)*.* Both the no *GAL4* control and the *wor-*

GAL4>orb2^{RNAi} (hereafter, *orb2^{RNAi}*) interphase NSCs showed equal distributions of Asl at apical

and basal centrosomes (Fig. 1F–H). While controls appeared WT with an enrichment of γTub on

the apical centrosome, a subset of *orb2^{RNAi}* NSCs (~30%; N=16/51) recruited γTub to the basal

centrosome precociously and showed reduced γTub at the apical centrosome (Fig. 1G–J).

These data indicate Orb2 promotes centrosome asymmetry cell autonomously within NSCs

through opposing activities at the apical and basal centrosomes.

Loss of *orb2* is associated with supernumerary centrosomes

The precocious activation of the basal centrosome is associated with errant centrosome

segregation during ACD, resulting in both centrosomes retained within the self-renewing stem

cell (Lerit and Rusan, 2013). To assay whether *orb2* loss similarly impairs centrosome

segregation, we quantified the frequency of supernumerary centrosomes in WT and *orb2* NSCs.

While >90% WT NSCs (N=25/30) had the expected 2 centrosomes, ~35% *orb2* mutant NSCs

(N=15/40) had extra centrosomes (Fig 2A–C; $p<0.05$ by chi-square test). A similar frequency of

supernumerary centrosomes was observed in *orb2^{RNAi}* NSCs (~25%; N=13/51; $p<0.05$ by chi-

square test), demonstrating Orb2 functions within NSC to regulate centrosome number, likely

due to failed segregation of the basal centrosome to the GMC (Fig. 2D–F).

Orb2 is required for mitotic spindle morphogenesis

Given that Orb2 helps regulate centrosome activity and segregation, we next examined its role

in spindle orientation. During ACD, the NSC mitotic spindle normally orients along an invariant

apical-basal polarity axis entrained by the concerted action of multiple protein complexes

(Siegrist and Doe, 2005; Siller et al., 2006; Cabernard and Doe, 2009). Apical polarization is

initiated by the Par-complex, comprising Bazooka (Baz)/Par-3, Par-6, and atypical protein

kinase C (αPKC; (Kuchinke et al., 1998; Cai et al., 2003; Goldstein and Macara, 2007)).

Inhibitory phosphorylation events by αPKC restrict basal fate determinants and the adapter

protein Miranda (Mira) to the basal cortex (Betschinger et al., 2003; Atwood and Prehoda,

2009).

We examined polarity establishment in mitotic WT and *orb2* NSCs by monitoring the

localization of Baz and Mira. Localization of Baz to the apical cortex in late interphase initiates

NSC polarization (Wodarz and Huttner, 2003; Knoblich, 2008), while localization of Mira to the

basal cortex during mitosis represents a late polarization step (Shen et al., 1998; Rolls et al.,

2003; Atwood and Prehoda, 2009). WT and *orb2* mutant NSCs showed similar distributions of

Baz and Mira to the apical and basal cortices, respectively (Fig 2G–I; n.s. by chi-square test).

These data argue that polarization is not significantly disrupted in *orb2* larval NSCs.

We next used multiple coordinate analysis to define spindle orientation in WT and *orb2*

NSCs stained for β-Tubulin to label MTs, Baz, and Asl (see *Methods*; Fig. 2J–L). First, we

measured spindle orientation (θ_1),, the angle between the apical cortical polarity and spindle

alignment axes (Fig. 2J, M–N). In WT, most spindles aligned within 30° of the polarity axis, yet

~20% of *orb2* NSC spindles were misoriented >30°, indicating *orb2* spindles are more

randomized than controls (Fig 2M, N; *p*<0.0001 by chi-square test).

To ascertain how defects in spindle orientation arise, we next quantified apical polarity

alignment, θ_2, the angle between the apical centrosome and polarity axes (Fig. 2J, O–P). The

anchoring of the apical centrosome to the apical cortex happens during early interphase and

influences polarization in NSCs and other stem cells (Yu et al., 2003; Januschke and Gonzalez,

2010; Inaba et al., 2015). Both WT and *orb2* NSCs showed alignment of the apical centrosome

to the apical polarity axis, consistent with our observations that apical polarity is unaffected in

orb2 mutants (Fig. 2G-I, O-P).

Finally, we examined centrosome alignment, θ_3, the angle between the apical and basal

centrosome axes (Fig 2J, Q-R). About 90% of WT mitotic NSCs aligned the basal centrosome

~180° away from the apical centrosome (Fig 2K, Q-R). In contrast, θ_3 is reduced in *orb2* NSCs

(Fig. 2L, Q). While the average reduction in θ_3 is not statistically significant, a subpopulation of

orb2 NSCs (~20%; N=7/32 cells) show defective θ_3 <120° (Fig. 2L, Q). Consequently, spindle

morphogenesis is significantly impaired, resulting in a higher frequency (~25%; N=9/34 cells;

p<0.0001 by chi-square test) of bent spindles in *orb2* NSCs relative to WT (Fig 2R). Taken

together, these data argue the precocious activation of the basal centrosome in *orb2* NSCs

during interphase impairs centrosome migration to the basal cortex, resulting in aberrant spindle

alignment and morphology. Alternatively, *orb2* may have a distinct functions in interphase

centrosome asymmetry and mitotic spindle orientation.

Loss of *orb2* results in microcephaly

Defects in centrosome regulation contribute to neurodevelopmental disorders, including

microcephaly and intellectual disability (Robinson et al., 2020). Previous work illustrates *orb2* is

crucial for learning and memory in the adult *Drosophila* brain (Keleman et al., 2007; Mastushita-

Sakai et al., 2010; Kruttner et al., 2012; Kacsoh et al., 2015; Hervas et al., 2016; Sanguanini

and Cattaneo, 2018). Moreover, reduced brain volumes were noted from serial sectioning adult

orb2 brains (Kruttner et al., 2012). Thus, to assess larval neurodevelopment in *orb2* mutants, we

measured the volume of single optic lobes from age-matched third instar larva (Link et al.,

2019). Compared to WT brains, *orb2* brains were significantly smaller (Fig 3A,B,E; *p*<0.0001 by

t-test). About 75% (N=10/13 cells) of *orb2* brains had volumes < 2-S.D. from WT, consistent

with microcephaly.

To determine if NSC loss contributes to *orb2*-dependent microcephaly, we counted the

number of Mira+ NSCs in control and *orb2* brains. While WT had ~80 NSCs, *orb2* mutants

showed a ~20% reduction with ~60 NSCs per lobe (Fig 3C,D,F; *p*<0.001 by t-test). These data

indicate Orb2 is required to maintain the NSC pool.

Are the centrosome and spindle defects observed in *orb2* mutants correlated with NSC

loss and/or microcephaly? To begin to address this question, we depleted *orb2* specifically in

NSCs and compared the brain volume of age matched *orb2^RNAi^* brains relative to no *GAL4*

controls and noted no difference (Fig. 3 G,H,K), indicating *orb2*-dependent microcephaly is

nonautonomous to NSCs – Orb2 is required in other cell types to affect brain size. In contrast,

NSC loss is a cell autonomous response to *orb2* depletion. Similar to *orb2* mutants, we detected

~30% fewer NSC per optic lobe in *orb2^RNAi^* brains relative to no *GAL4* controls (Fig 3I,J,L;

p<0.0001 by t-test). We conclude the frequency of NSC loss in *orb2* mutants is similar to the

incidence of centrosome defects, arguing these responses are correlated. Moreover, NSC loss

may be uncoupled from *orb2*-dependent microcephaly, consistent with the idea that Orb2

supports neurodevelopment in multiple cellular lineages.

To elucidate mechanisms underlying NSC loss in *orb2* mutants, we first tested the

hypothesis that NSCs are eliminated by cell death. We quantified the coincidence of pro-

apoptotic cleaved Caspase 3 (CC3; (Fan and Bergmann, 2010)) in cells marked with the NSC-

specific Deadpan (Dpn) antibody (Bier et al., 1992). Similar rates of apoptosis were observed in

WT and *orb2* NSCs, indicating NSC loss occurs by other mechanisms (Fig S1 A–C; n.s. by t-

test). Another mechanism whereby NSCs may be depleted is via premature differentiation

(Cabernard and Doe, 2009; Lai and Doe, 2014; Abdel-Salam et al., 2020). Normally, the pro-

differentiation marker Prospero (Pros) is confined to the nucleus of the differentiating GMCs and

absent from NSCs (Doe et al., 1991; Vaessin et al., 1991), while retention of Pros in the NSC

nucleus promotes premature differentiation (Lai and Doe, 2014). We quantified the coincidence

of Pros with Dpn and detected no significant difference relative to WT, indicating premature

differentiation does not precipitate NSC loss in *orb2* mutants (Fig S1D–F; n.s. by t-test).

Finally, we examined whether *orb2* affects mitotic progression, reasoning impaired NSC

self-renewal might contribute to NSC loss. However, the mitotic indices in WT and *orb2* brains

were not significantly different ($33.6\pm4.9\%$ per lobe in N=6 WT vs. $36.3\pm10.1\%$ in N=10 *orb2*

brains; Fig S1G–I; n.s. by t-test). These data suggest NSC loss is not due to increased

quiescence. Loss of NSCs observed in *orb2* mutants may be actuated by non-apoptotic cell

death pathways. Alternatively, the altered neuronal specification observed in *orb2* embryos may

impinge upon larval neurodevelopment, resulting in fewer NSCs (Hafer et al., 2011).

Orb2 localizes to centrosomes within cycling NSCs

Our data indicate Orb2 functions autonomously within NSCs to regulate centrosome

asymmetry. Previous work indicates CPEB proteins localize to centrosomes in *Xenopus*

embryos and cultured mammalian cells (Groisman et al., 2000; Pascual et al., 2020a; Pascual

et al., 2020b). Although Orb2 localizes to neuronal lineages, subcellular localization of Orb2 is

not well defined (Keleman et al., 2007; Hafer et al., 2011). Therefore, to inform mechanisms of

how Orb2 functions within NSCs to support centrosome asymmetry and spindle morphogenesis,

we examined endogenous Orb2 localization in NSCs using monoclonal antibodies (Hafer et al.,

2011). In interphase NSCs, Orb2 appeared dispersed throughout the cytoplasm with a notable

enrichment on the apical centrosome (Fig 4A, *interphase*). Apical centrosome enrichment of

Orb2 was observed in N=21/24 interphase NSCs, indicated by positive Orb2 AI values (Fig. 4B).

On average, interphase apical centrosomes contained 1.7-fold more Orb2 than basal

centrosomes (Fig. 4C; *p*<0.01 by t-test). Upon mitotic entry, Orb2 localization increased at both

centrosomes, demonstrating Orb2 is recruited to active centrosomes during centrosome

maturation (Fig 4A, *prophase* and *metaphase*). Our localization analysis reveals two populations

of Orb2 in interphase NSCs, a cytoplasmic versus centrosomal pool, raising the possibility that

Orb2 may normally function to regulate centrosome activity locally at centrosomes, from a

distance within the cytoplasm, or both.

Orb2 regulates PLP protein levels in larval brains

Because Orb2 is an RNA-binding protein known to promote or repress translation of its target

mRNAs, we reasoned it may contribute to centrosome regulation indirectly through regulation of

an mRNA associated with centrosome asymmetry (Mastushita-Sakai et al., 2010; Xu et al.,

2012; Khan et al., 2015). A genome-wide analysis for mRNA targets bound by Orb2 within

Drosophila S2 cell extracts uncovered thousands of putative targets, including mRNAs of genes

required for NSC centrosome asymmetry, e.g., *Cep135*, *Cnb*, *plp*, *polo*, and *Wdr62* mRNAs

(Januschke et al., 2013; Lerit and Rusan, 2013; Singh et al., 2014; Ramdas Nair et al., 2016;

Stepien et al., 2016; Gallaud et al., 2020). While these transcripts were identified as potential

Orb2 targets, they varied in their biologic complexity (BC), a metric of reproducibility across

biological replicates (Licatalosi et al., 2008).

Aspects of the *orb2* null phenotype in NSCs resemble *plp* loss, as both mutants display

precocious activation of the basal centrosome in interphase NSCs coincident with

supernumerary centrosome and spindle defects (Lerit and Rusan, 2013). Given these

similarities, we tested if Orb2 regulates PLP protein expression by comparing PLP levels in WT

vs. *orb2* larval brain extracts. Semi-quantitative western blotting uncovered an average 30%

reduction in PLP in *orb2* relative to WT, suggesting Orb2 promotes PLP translation (Fig. 5A, B).

To assay a requirement for Orb2 in regulating PLP in NSCs, we examined PLP localization to

the apical versus basal centrosomes in interphase NSCs. In WT NSCs, PLP was enriched on

the inactive, basal centrosome, consistent with prior work (Fig. 5C; (Lerit and Rusan, 2013;

Singh et al., 2014). Despite a reduction in PLP levels in whole brain extracts, robust localization

of PLP to centrosomes was detected in *orb2* NSCs, comparable to WT (Fig. 5C, D; n.s. by

Kolmogorov-Smirnov test). Further, levels of PLP localized at apical vs. basal centrosomes

were not significantly different from WT (Fig. 5E). Taken together, these data imply Orb2-

dependent regulation of PLP may occur outside of NSCs and raise the possibility that other

Orb2 targets contribute to NSC centrosome regulation.

To refine the list of targets that may be involved in NSC centrosome regulation, we

compared the Orb2 mRNA targets identified in *Drosophila* S2 cells (Stepien et al., 2016) to a

recent list of mRNA targets bound by CPEB4 in HeLa cells (Pascual et al., 2020b) and identified

1083 overlapping genes (Fig S2A; Table S1). Gene ontology (GO) analysis for cellular

components uncovered a significant enrichment of organelle terms (Fig S2B). In contrast,

centrosome-related ontologies were not significantly enriched among the overlapping genes

(Fig S2C). However, because Orb2 has multiple orthologs, CPEB2-4, other putative Orb2

targets absent from the CPEB4 dataset are omitted from these pairwise analyses. Therefore,

we examined centrosome-related ontologies of Orb2 targets with a BC $\geq$ 3. This analysis

identified 2150 mRNAs, among which 53 mRNAs were annotated with centrosome ontologies,

which may be subject to Orb2 regulation (Fig. S2D; Table S1).

Model Summary

Our data indicate Orb2 is required for robust inactivation of the basal centrosome and may also

enhance apical centrosome activity in interphase NSCs. Orb2 also functions in other cell types

for normal brain size (Fig 5F). Given its local enrichment, we propose Orb2 promotes apical

centrosome maturation (Fig 5F, *top box*), while cytoplasmic Orb2 inactivates the basal

centrosome, perhaps by repressing pro-maturation factors (Fig 5F, *bottom box*). Consistent with

opposing activities at the apical vs. basal centrosomes, Orb2 activates or represses the

translation of its targets depending on its oligomerization status ((Khan et al., 2015)). This model

predicts cytoplasmic Orb2 is predominantly monomeric, thereby functioning as a translational

repressor, while centrosome-localized Orb2 is oligomerized and functions as a translational

activator. Whether on-site translation occurs at the apical centrosome remains to be tested and

specific Orb2 targets require identification.

307   Microcephaly is quite heterogenous; however, a significant number of genes associated

308   with heritable microcephaly and intellectual disability are also associated with centrosome

309   biogenesis and regulation (Thornton and Woods, 2009; Jayaraman et al., 2018; Robinson et al.,

310   2020). Our study implicates Orb2 at the intersection of these pathways.

Materials and Methods

Fly Stocks

The following strains and transgenic lines were used: y^1w^{1118} (Bloomington *Drosophila* Stock

Center (BDSC) #1495) was used as the WT control unless otherwise noted. *orb2* brains were

isolated from the homozygous null allele, $orb2^{\Delta36}$ (gift from P. Schedl, Princeton University (Xu

et al., 2012)). NSC-specific depletion of *orb2* by $orb2^{RNAi}$ ($P\{TRiP.HMJ22715\}^{attP40}$; BDSC

#60424) was driven by $(P\{wor.GAL4.A\}^2$; BDSC #56555). All strains were maintained on

Bloomington formula cornmeal-agar media (Lab-Express, Inc.; Ann Arbor, MI) at 25°C in a light

and temperature-controlled incubator.

Immunofluorescence

Crawling third instar larva were used for dissections. Larval brains were prepared for

immunofluorescence as previously described (Lerit et al., 2014). Briefly, brains were dissected

in Schneider's *Drosophila* Medium (ThermoFisher Scientific, #21720024), fixed in 9%

paraformaldehyde for 15 min, blocked in PBT buffer (Phosphate Buffered Saline (PBS)

supplemented with 1% BSA and 0.1% Tween-20) for one hour at room temperature prior to

overnight incubation in primary antibodies in PBT with nutation at 4°C. Samples were further

blocked with modified PBT (2% BSA, 0.1% Tween-20, and 4% normal goat serum in PBS) for 1

hour before incubation for two hours at room temperature with secondary antibody and DAPI.

Brains were oriented and mounted in Aqua-Poly/Mount (Polysciences, Inc) prior to imaging.

The following primary antibodies were used: guinea pig anti-Asl (1:4000, gift from G.

Rogers, University of Arizona), mouse anti-GTU88 (γTub; 1:250–350, Sigma T6557), rabbit anti-

Cnn (1:4000 gift from Tim Megraw, Florida State University), guinea pig anti-PLP (1:4000, gift

from Nasser Rusan, NIH), rabbit anti-phospho-Histone H3 Ser10 (pH3; 1:2000, Sigma-Millipore,

05-570), mouse anti-β-Tub (clone E7, 1:250; Developmental Studies Hybridoma Bank (DSHB)),

mouse anti-Orb2 (undiluted 1:1 mix of clones 2D11 and 4G8 (DSHB); (Xu et al., 2012)), rat anti-

Mira (1:500, Abcam, ab197788), rabbit anti-Baz (1:2000, gift from A. Harris, University of

Toronto), rat anti-Dpn (1:500, Abcam, ab195173), rabbit anti-cleaved Caspase 3 (CC3, 1:75;

Cell Signaling Technology, 9661s), and mouse anti-Pros (clone MR1A, 1:500; DSHB).

Secondary antibodies: Alexa Fluor 488, 568, and 647 (1:500, Molecular Probes). DAPI

(ThermoFisher Scientific) was used at 10 ng/mL.

Microscopy

Images were acquired on a Nikon Ti-E inverted microscope fitted with a Yokogawa CSU-X1

spinning disk head (Yokogawa Corp. of America), Orca Flash 4.0 v2 CMOS camera

(Hamamatsu Corp.), Perfect Focus system (Nikon), and a Nikon LU-N4 solid-state laser launch

(15 mW; 405, 488, 561, and 647 nm) using the following Nikon objectives: 100x 1.49-NA Apo

Total Internal Reflection Fluorescence oil immersion, 40x 1.3-NA Plan Fluor oil immersion, and

2x× 0.75-NA Plan Apo. Images were acquired at 25°C through Nikon Elements AR software on

a 64-bit HP Z440 workstation (Hewlett-Packard).

Image analysis

Images were assembled using Fiji (National Institutes of Health;(Schindelin et al., 2012)), Adobe

Photoshop, and Adobe Illustrator software to separate or merge channels, crop regions of

interest, generate maximum-intensity projections, and adjust brightness and contrast.

Centrosome asymmetry

Interphase NSCs were identified by the absence of pH3, round nuclear morphology, and

presence of duplicated centrosomes in large ($\geq$10 μm) cells. To blind the experimenter to

genotype, maximum projected images were generated, randomized, and used to measure

background-subtracted integrated densities from regions of interest (ROIs) centered at the

apical or basal centrosomes. Apical centrosome integrated density (*A*) and basal centrosome

integrated density (*B*) were used to calculate the asymmetry index, (*A-B*)/(*A+B*) ((Lerit and

Rusan, 2013)). The data were then unblinded and mean $\pm$ S.D. were calculated per genotype.

Polarity, NSC number, and mitotic index

Mitotic NSCs were identified by the presence of pH3 in large, Mira+ cells. To score polarity,

maximum projected images were anonymized to blind the experimenter to genotype, and each

NSC was scored for the absence or presence of Baz or Mira crescents at the apical or basal

cortices, respectively. All Mira+ NSCs were counted to calculate the number of NSCs per optic

lobe. Mitotic index is defined as the number of Mira+, pH3+ NSCs per total Mira+ NSCs.

Spindle morphology

Z-stack images of mitotic NSCs labeled with anti-β-Tub to label the mitotic spindle, Asl to mark

the centrioles, and Baz to label the apical polarity axis were randomized to blind the

experimenter to genotype. The point tool in Fiji was used to record the X,Y, and Z coordinates of

four ROIs per cell: 1) center of the apical centrosome, 2) center of the basal centrosome, 3)

center of the Baz apical crescent, and 4) center of the DAPI+ condensed chromosomes. These

points were used to calculate four different vectors, and vector analysis was used to calculate

the angles between specified vectors. The following angles (θ) were calculated: θ_1= *spindle*

orientation, the angle between the vectors defined by points 1 and 2 (division axis) relative to

points 3 and 4 (polarity axis); θ_2= *apical polarity alignment* is the angle between the vectors

defined by points 3 and 4 (apical polarity axis) relative to points 1 and 4 (apical centrosome

axis); and θ_3= *centrosome alignment* is the angle between the vectors defined by points 1 and 4

(apical centrosome axis) relative to points 2 and 4 (basal centrosome axis). Angles that fell

outside $\pm$ 2 S.D. from the control mean were defined as defective. For spindle alignment, NSCs

with spindle angles >75° were classified as orthogonal.

Age-matched brain volume

To age match larva, 20 female virgins and 10 males of the appropriate genotype were allowed

to seed a vial for 24 hrs. After removal of the adults, vials were incubated at 25 $^{\circ}$C until crawling

third instar larva emerged. *orb2* null progeny showed a developmental delay of 24-48 hrs as

compared to controls; null larva took ~7-8 days to emerge. Age-matched crawling third instar

larva were dissected and prepared for immunofluorescence. The entire volume of the DAPI-

labeled brain was imaged. Imaris software (Oxford Instruments) was used to select an ROI of

the optic lobe and measure the volume using the 3D surfaces tool (Link et al., 2019).

NSC differentiation and death rate

NSCs were identified by the presence of Dpn. To score for premature differentiation, maximum

projected images of Dpn and Pros stained brains were randomized to blind the experimenter to

genotype, and an ROI of the central brain region was used to calculate the Pearson's

Correlation Coefficient on background-subtracted and automatic threshold–masked images

using the Coloc 2 plugin for Fiji (Schindelin et al., 2012). To score cell death, a similar analysis

was run on Dpn and CC3 stained brains. To assess specificity of overlapping signals for both

experiments, the red channel (Pros or CC3) was rotated 90° clockwise, and colocalization was

remeasured.

Immunoblotting

Larval brain extracts were prepared from 20 crawling third instar larva dissected in Schneider's

medium, removed of imaginal discs, transferred to fresh media, then rinsed once in cold PBST.

Samples were homogenized on ice in 30 μL of fresh PBST using a cordless motor and plastic,

disposable homogenizer, supplemented with 20 μL 5x SDS loading buffer, boiled for 10 min at

95 °C, then stored at -20 °C or resolved on a commercial 7.5% polyacrylamide gel (Bio-Rad,

#4568023. Proteins were transferred to a 0.2 μm nitrocellulose membrane (GE Healthcare) by

wet transfer in a buffer containing 25 mM Tris-HCl, pH 7.6, 192 mM glycine, 10% methanol, and

0.02% SDS at 4 °C. Membranes were blocked in 5% dry milk in TBST (Tris-buffered saline,

0.05% Tween-20), washed well with TBST, and incubated overnight at 4 °C with primary

antibodies. After washing with TBST, membranes were incubated for 1.5 hr in secondary

antibodies diluted 1:5000 in TBST. Bands were visualized with Clarity ECL substrate (Bio-Rad,

1705061) on a Bio-Rad ChemiDoc imaging system.

The following primary antibodies were used: rabbit anti-PLP (1:4000, gift from Nasser

Rusan, NIH), guinea pig anti-Asl (1:10,000, gift from Greg Rogers, University of Arizona), and

mouse anti-Orb2 (1:25 dilution each of 2D11 and 4G8; DSHB, Paul Schedl, Princeton University

(Hafer et al., 2011)). Secondary antibodies: goat anti-mouse HRP (1:5000, ThermoFisher

#31430), goat anti-rabbit HRP (1:5000, ThermoFisher #31460), and goat anti-guinea pig HRP

(1:5000, ThermoFisher #A18769). Densitometry was measured in Adobe Photoshop and

protein levels are normalized to the Asl loading control. Full-size, replicated blots are available

to view on FigShare: https://doi.org/10.6084/m9.figshare.17052722 . The experiment was

repeated with three biological replicates per genotype.

Bioinformatics

Gene names were converted to FlyBase identifiers using the Flybase.org tool 'Query by

symbols'(Larkin et al., 2021). Overlapping genes were identified by the 'COUNT IF' function in

Excel and Venn diagrams were plotted in R-Studio. GO cellular component analysis was done

using the Panther statistical overrepresentation test (http://www.pantherdb.org/), and Fisher's

exact test was used to generate an adjusted p-value, i.e., false discovery rate (FDR)(Mi et al.,

2021) .

Statistical analysis

Data were plotted and statistical analysis performed using Microsoft Excel, GraphPad Prism,

and RStudio software. Normality distributions were determined using a D'Agnostino-Pearson or

Shapiro-Wilk normality test. Goodness-of-Fit chi-squared analysis were performed using control

distributions as the expected distributions. Data were further analyzed using parametric two-

tailed t-test, non-parametric Mann-Whitney test or Kolmogorov-Smirnov (KS) test, or an

appropriate post-hoc test as noted in the figure legends. Data are plotted as mean $\pm$ S.D. and

are representative results from at 2 or more independent experiments.

Supplemental materials

Fig. S1 NSC differentiation and survival in WT vs. *orb2* mutants. Maximum intensity

projections of (A) WT and (B) *orb2* brains stained with Dpn (green; NSC nuclei) and CC3 (red;

pro-apototic. Insets (boxes) are enlarged to the right to highlight NSCs (dashed ovals). (C)

Quantification of the Pearson's correlation coefficient between Dpn and CC3 from N=10 WT and

15 *orb2* optic lobes pooled from two replicates. The CC3 channel was rotated clockwise (90°) to

test for specificity of overlapping signals. (D) Maximum intensity projections of WT and (E) *orb2*

brains stained with Dpn (green) and Pros (red; differentiated nuclei). (F) Quantification of the

Pearson's correlation coefficient between Dpn and Pros from N=16 WT and 18 *orb2* optic lobes

pooled from two replicates. The Pros channel was rotated clockwise (90°) to test for specificity

of overlapping signals. In (D) and (F), each data point is from one ROI per optic lobe. (G)

Projected images of WT and (H) *orb2* NSCs (dashed ovals) stained for Mira (green; NSCs) and

the mitotic marker pH3 (red). (I) Quantification of mitotic index from N=6 WT and 10 *orb2* brains.

n.s., not significant by Student's t-test. Bars: (A–E) 20 μm and 8 μm (insets); (G and H) 10 μm.

Fig. S2 Ontological analysis of Orb2 and human CPEB4. A) Venn diagram showing relative

transcript pool sizes and overlap detected from the Stepien et al. Orb2 CLIP and Pascual et al.

CPEB4 datasets (Stepien et al., 2016; Pascual et al., 2020b). B) Top 5 gene ontologies

overrepresented in the overlapping transcripts by lowest raw *p* value. C) Overrepresentation

tests for gene ontologies implicated in the centrosome were not significant. D)

Overrepresentation test of genes in the Orb2 CLIPseq with BC $\geq$3 yields significant centrosome

ontologies. -log *p*-values are plotted with a significance cut-off of *p*=0.05, noted by the dashed

line and asterisk. Genes are listed in Table S1.

Table S1. Orb2 and CPEB4 common RNA targets. Sheet 1 lists FlyBase IDs for all transcripts

detected in (Stepien et al., 2016; Pascual et al., 2020b). Sheet 2 lists overlapping genes;

column E is sortable to list genes with centrosome ontologies. Sheet 3 lists GO terms

overrepresented in the overlapping genes, column J is sortable to list genes with centrosome

ontologies. Sheet 4 lists Orb2 targets from (Stepien et al., 2016) with BC $\geq$ 3; column F is

sortable to list genes with centrosome ontologies. Sheet 5 shows an overrepresentation test for

centrosome ontologies from the Orb2 BC $\geq$ 3 gene list. Sheet 6 lists each gene included in the

indicated centrosome ontologies.

Competing interest statement

The authors have no competing interests to declare.

Acknowledgements

We thank Drs. Timothy Megraw, Nasser Rusan, Greg Rogers, and Paul Schedl for gifts of

reagents. We are grateful to members of the Lerit lab and Drs. Victor Faundez and Nasser

Rusan for constructive feedback on the manuscript.

This work was supported by NIH grant T32GM008490 (BVR), American Heart Association

grant 20POST35210023 (to JF), and R01GM138544 (to DAL).

Author contributions

Beverly V. Robinson– data curation, formal analysis, funding acquisition, investigation,

methodology, visualization, writing–original draft, and writing–review & editing.

Junnan Fang– formal analysis, funding acquisition, investigation, methodology, and writing–

review & editing.

Dipen S. Mehta– investigation

Joseph D. Buehler– formal analysis, software, visualization, and writing–review & editing.

Dorothy A. Lerit– conceptualization, formal analysis, funding acquisition, investigation,

methodology, project administration, supervision, visualization, writing–original draft, and

writing–review & editing.

References

Abdel-Salam, G.M.H., I.S.M. Sayed, H.H. Afifi, S.F. Abdel-Ghafar, M.R. Abouzaid, S.I. Ismail, M.S. Aglan, M.Y. Issa, H.T. El-Bassyouni, G. El-Kamah, L.K. Effat, M. Eid, M.S. Zaki, S.A. Temtamy, and M.S. Abdel-Hamid. 2020. Microcephalic osteodysplastic primordial dwarfism type II: Additional nine patients with implications on phenotype and genotype correlation. *Am. J. Med. Genet. A*. 182:1407-1420.

Albertson, R., C. Chabu, A. Sheehan, and C.Q. Doe. 2004. Scribble protein domain mapping reveals a multistep localization mechanism and domains necessary for establishing cortical polarity. *J. Cell Sci.* 117:6061-6070.

Atwood, S.X., and K.E. Prehoda. 2009. aPKC phosphorylates Miranda to polarize fate determinants during neuroblast asymmetric cell division. *Curr. Biol.* 19:723-729.

Betschinger, J., K. Mechtler, and J.A. Knoblich. 2003. The Par complex directs asymmetric cell division by phosphorylating the cytoskeletal protein Lgl. *Nature*. 422:326-330.

Bier, E., H. Vaessin, S. Younger-Shepherd, L.Y. Jan, and Y.N. Jan. 1992. deadpan, an essential pan-neural gene in Drosophila, encodes a helix-loop-helix protein similar to the hairy gene product. *Genes Dev.* 6:2137-2151.

Bond, J., E. Roberts, G.H. Mochida, D.J. Hampshire, S. Scott, J.M. Askham, K. Springell, M. Mahadevan, Y.J. Crow, A.F. Markham, C.A. Walsh, and C.G. Woods. 2002. ASPM is a major determinant of cerebral cortical size. *Nat. Genet.* 32:316-320.

Broadus, J., and C.Q. Doe. 1997. Extrinsic cues, intrinsic cues and microfilaments regulate asymmetric protein localization in Drosophila neuroblasts. *Curr. Biol.* 7:827-835.

Cabernard, C., and C.Q. Doe. 2009. Apical/basal spindle orientation is required for neuroblast homeostasis and neuronal differentiation in Drosophila. *Dev. Cell*. 17:134-141.

Cai, Y., F. Yu, S. Lin, W. Chia, and X. Yang. 2003. Apical complex genes control mitotic spindle geometry and relative size of daughter cells in Drosophila neuroblast and pI asymmetric divisions. *Cell*. 112:51-62.

Conduit, P.T., and J.W. Raff. 2010. Cnn dynamics drive centrosome size asymmetry to ensure daughter centriole retention in Drosophila neuroblasts. *Curr. Biol.* 20:2187-2192.

Conduit, P.T., A. Wainman, and J.W. Raff. 2015. Centrosome function and assembly in animal cells. *Nat. Rev. Mol. Cell Biol.* 16:611-624.

Doe, C.Q. 2008. Neural stem cells: balancing self-renewal with differentiation. *Development*. 135:1575-1587.

Doe, C.Q., Q. Chu-LaGraff, D.M. Wright, and M.P. Scott. 1991. The prospero gene specifies cell fates in the Drosophila central nervous system. *Cell*. 65:451-464.

Fan, Y., and A. Bergmann. 2010. The cleaved-Caspase-3 antibody is a marker of Caspase-9-like DRONC activity in Drosophila. *Cell Death Differ.* 17:534-539.

Gallaud, E., A. Ramdas Nair, N. Horsley, A. Monnard, P. Singh, T.T. Pham, D. Salvador Garcia, A. Ferrand, and C. Cabernard. 2020. Dynamic centriolar localization of Polo and Centrobin in early mitosis primes centrosome asymmetry. *PLoS Biol.* 18:e3000762.

Gambarotto, D., C. Pennetier, J.M. Ryniawec, D.W. Buster, D. Gogendeau, A. Goupil, M. Nano, A. Simon, D. Blanc, V. Racine, Y. Kimata, G.C. Rogers, and R. Basto. 2019. Plk4 Regulates Centriole Asymmetry and Spindle Orientation in Neural Stem Cells. *Dev. Cell*. 50:11-24 e10.

Goldstein, B., and I.G. Macara. 2007. The PAR proteins: fundamental players in animal cell polarization. *Dev. Cell*. 13:609-622.

Gould, R.R., and G.G. Borisy. 1977. The pericentriolar material in Chinese hamster ovary cells nucleates microtubule formation. *J Cell Biol*. 73:601-615.

Groisman, I., Y.S. Huang, R. Mendez, Q. Cao, W. Theurkauf, and J.D. Richter. 2000. CPEB, maskin, and cyclin B1 mRNA at the mitotic apparatus: implications for local translational control of cell division. *Cell*. 103:435-447.

Hafer, N., S. Xu, K.M. Bhat, and P. Schedl. 2011. The Drosophila CPEB protein Orb2 has a
novel expression pattern and is important for asymmetric cell division and nervous
system function. *Genetics*. 189:907-921.
Hervas, R., L. Li, A. Majumdar, C. Fernandez-Ramirez Mdel, J.R. Unruh, B.D. Slaughter, A.
Galera-Prat, E. Santana, M. Suzuki, Y. Nagai, M. Bruix, S. Casas-Tinto, M. Menendez,
D.V. Laurents, K. Si, and M. Carrion-Vazquez. 2016. Molecular Basis of Orb2
Amyloidogenesis and Blockade of Memory Consolidation. *PLoS Biol.* 14:e1002361.
Huang, Y.S., M.C. Kan, C.L. Lin, and J.D. Richter. 2006. CPEB3 and CPEB4 in neurons:
analysis of RNA-binding specificity and translational control of AMPA receptor GluR2
mRNA. *EMBO J.* 25:4865-4876.
Inaba, M., Z.G. Venkei, and Y.M. Yamashita. 2015. The polarity protein Baz forms a platform for
the centrosome orientation during asymmetric stem cell division in the Drosophila male
germline. *Elife*. 4.
Januschke, J., and C. Gonzalez. 2010. The interphase microtubule aster is a determinant of
asymmetric division orientation in Drosophila neuroblasts. *J Cell Biol*. 188:693-706.
Januschke, J., S. Llamazares, J. Reina, and C. Gonzalez. 2011. Drosophila neuroblasts retain
the daughter centrosome. *Nat Commun*. 2:243.
Januschke, J., J. Reina, S. Llamazares, T. Bertran, F. Rossi, J. Roig, and C. Gonzalez. 2013.
Centrobin controls mother-daughter centriole asymmetry in Drosophila neuroblasts. *Nat.*
Cell Biol. 15:241-248.
Jayaraman, D., B.I. Bae, and C.A. Walsh. 2018. The Genetics of Primary Microcephaly. *Annu*
Rev Genomics Hum Genet. 19:177-200.
Kacsoh, B.Z., J. Bozler, S. Hodge, M. Ramaswami, and G. Bosco. 2015. A novel paradigm for
nonassociative long-term memory in Drosophila: predator-induced changes in
oviposition behavior. *Genetics*. 199:1143-1157.
Keleman, K., S. Kruttner, M. Alenius, and B.J. Dickson. 2007. Function of the Drosophila CPEB
protein Orb2 in long-term courtship memory. *Nat. Neurosci.* 10:1587-1593.
Khan, M.R., L. Li, C. Perez-Sanchez, A. Saraf, L. Florens, B.D. Slaughter, J.R. Unruh, and K.
Si. 2015. Amyloidogenic Oligomerization Transforms Drosophila Orb2 from a Translation
Repressor to an Activator. *Cell*. 163:1468-1483.
Khodjakov, A., and C.L. Rieder. 1999. The sudden recruitment of gamma-tubulin to the
centrosome at the onset of mitosis and its dynamic exchange throughout the cell cycle,
do not require microtubules. *J Cell Biol*. 146:585-596.
Knoblich, J.A. 2008. Mechanisms of asymmetric stem cell division. *Cell*. 132:583-597.
Knoblich, J.A., L.Y. Jan, and Y.N. Jan. 1995. Asymmetric segregation of Numb and Prospero
during cell division. *Nature*. 377:624-627.
Kraut, R., W. Chia, L.Y. Jan, Y.N. Jan, and J.A. Knoblich. 1996. Role of inscuteable in orienting
asymmetric cell divisions in Drosophila. *Nature*. 383:50-55.
Kruttner, S., B. Stepien, J.N. Noordermeer, M.A. Mommaas, K. Mechtler, B.J. Dickson, and K.
Keleman. 2012. Drosophila CPEB Orb2A mediates memory independent of Its RNA-
binding domain. *Neuron*. 76:383-395.
Kuchinke, U., F. Grawe, and E. Knust. 1998. Control of spindle orientation in Drosophila by the
Par-3-related PDZ-domain protein Bazooka. *Curr. Biol.* 8:1357-1365.
Lai, S.L., and C.Q. Doe. 2014. Transient nuclear Prospero induces neural progenitor
quiescence. *Elife*. 3.
Larkin, A., S.J. Marygold, G. Antonazzo, H. Attrill, G. Dos Santos, P.V. Garapati, J.L. Goodman,
L.S. Gramates, G. Millburn, V.B. Strelets, C.J. Tabone, J. Thurmond, and C. FlyBase.
2021. FlyBase: updates to the Drosophila melanogaster knowledge base. *Nucleic Acids*
Res. 49:D899-D907.
Lerit, D.A., K.M. Plevock, and N.M. Rusan. 2014. Live imaging of Drosophila larval neuroblasts.
J Vis Exp.

Lerit, D.A., and N.M. Rusan. 2013. PLP inhibits the activity of interphase centrosomes to ensure their proper segregation in stem cells. *J Cell Biol*. 202:1013-1022.

Licatalosi, D.D., A. Mele, J.J. Fak, J. Ule, M. Kayikci, S.W. Chi, T.A. Clark, A.C. Schweitzer, J.E. Blume, X. Wang, J.C. Darnell, and R.B. Darnell. 2008. HITS-CLIP yields genome-wide insights into brain alternative RNA processing. *Nature*. 456:464-469.

Link, N., H. Chung, A. Jolly, M. Withers, B. Tepe, B.R. Arenkiel, P.S. Shah, N.J. Krogan, H. Aydin, B.B. Geckinli, T. Tos, S. Isikay, B. Tuysuz, G.H. Mochida, A.X. Thomas, R.D. Clark, G.M. Mirzaa, J.R. Lupski, and H.J. Bellen. 2019. Mutations in ANKLE2, a ZIKA Virus Target, Disrupt an Asymmetric Cell Division Pathway in Drosophila Neuroblasts to Cause Microcephaly. *Dev. Cell*. 51:713-729 e716.

Mastushita-Sakai, T., E. White-Grindley, J. Samuelson, C. Seidel, and K. Si. 2010. Drosophila Orb2 targets genes involved in neuronal growth, synapse formation, and protein turnover. *Proc. Natl. Acad. Sci. U. S. A.* 107:11987-11992.

Mi, H., D. Ebert, A. Muruganujan, C. Mills, L.P. Albou, T. Mushayamaha, and P.D. Thomas. 2021. PANTHER version 16: a revised family classification, tree-based classification tool, enhancer regions and extensive API. *Nucleic Acids Res.* 49:D394-D403.

Pascual, R., J. Martin, F. Salvador, O. Reina, V. Chanes, A. Millanes-Romero, C. Suner, G. Fernandez-Miranda, A. Bartomeu, Y.S. Huang, R.R. Gomis, and R. Mendez. 2020a. The RNA binding protein CPEB2 regulates hormone sensing in mammary gland development and luminal breast cancer. *Sci Adv*. 6:eaax3868.

Pascual, R., C. Segura-Morales, M. Omerzu, N. Bellora, E. Belloc, C.L. Castellazzi, O. Reina, E. Eyras, M.M. Maurice, A. Millanes-Romero, and R. Mendez. 2020b. mRNA spindle localization and mitotic translational regulation by CPEB1 and CPEB4. *RNA*.

Ramdas Nair, A., P. Singh, D. Salvador Garcia, D. Rodriguez-Crespo, B. Egger, and C. Cabernard. 2016. The Microcephaly-Associated Protein Wdr62/CG7337 Is Required to Maintain Centrosome Asymmetry in Drosophila Neuroblasts. *Cell Rep.* 14:1100-1113.

Rebollo, E., P. Sampaio, J. Januschke, S. Llamazares, H. Varmark, and C. Gonzalez. 2007. Functionally unequal centrosomes drive spindle orientation in asymmetrically dividing Drosophila neural stem cells. *Dev. Cell*. 12:467-474.

Robinson, B.V., V. Faundez, and D.A. Lerit. 2020. Understanding microcephaly through the study of centrosome regulation in Drosophila neural stem cells. *Biochem. Soc. Trans.* 48:2101-2115.

Rolls, M.M., R. Albertson, H.P. Shih, C.Y. Lee, and C.Q. Doe. 2003. Drosophila aPKC regulates cell polarity and cell proliferation in neuroblasts and epithelia. *J Cell Biol*. 163:1089-1098.

Rusan, N.M., and M. Peifer. 2007. A role for a novel centrosome cycle in asymmetric cell division. *J Cell Biol*. 177:13-20.

Sanguanini, M., and A. Cattaneo. 2018. A continuous model of physiological prion aggregation suggests a role for Orb2 in gating long-term synaptic information. *R Soc Open Sci*. 5:180336.

Schindelin, J., I. Arganda-Carreras, E. Frise, V. Kaynig, M. Longair, T. Pietzsch, S. Preibisch, C. Rueden, S. Saalfeld, B. Schmid, J.Y. Tinevez, D.J. White, V. Hartenstein, K. Eliceiri, P. Tomancak, and A. Cardona. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods*. 9:676-682.

Shen, C.P., J.A. Knoblich, Y.M. Chan, M.M. Jiang, L.Y. Jan, and Y.N. Jan. 1998. Miranda as a multidomain adapter linking apically localized Inscuteable and basally localized Staufen and Prospero during asymmetric cell division in Drosophila. *Genes Dev.* 12:1837-1846.

Siegrist, S.E., and C.Q. Doe. 2005. Microtubule-induced Pins/Galphai cortical polarity in Drosophila neuroblasts. *Cell*. 123:1323-1335.

Siegrist, S.E., and C.Q. Doe. 2006. Extrinsic cues orient the cell division axis in Drosophila embryonic neuroblasts. *Development*. 133:529-536.

Siller, K.H., C. Cabernard, and C.Q. Doe. 2006. The NuMA-related Mud protein binds Pins and
regulates spindle orientation in Drosophila neuroblasts. *Nat. Cell Biol.* 8:594-600.
Singh, P., A. Ramdas Nair, and C. Cabernard. 2014. The centriolar protein Bld10/Cep135 is
required to establish centrosome asymmetry in Drosophila neuroblasts. *Curr. Biol.*
24:1548-1555.
Stepien, B.K., C. Oppitz, D. Gerlach, U. Dag, M. Novatchkova, S. Kruttner, A. Stark, and K.
Keleman. 2016. RNA-binding profiles of Drosophila CPEB proteins Orb and Orb2. *Proc.*
Natl. Acad. Sci. U. S. A. 113:E7030-E7038.
Thornton, G.K., and C.G. Woods. 2009. Primary microcephaly: do all roads lead to Rome?
Trends Genet. 25:501-510.
Vaessin, H., E. Grell, E. Wolff, E. Bier, L.Y. Jan, and Y.N. Jan. 1991. prospero is expressed in
neuronal precursors and encodes a nuclear protein that is involved in the control of
axonal outgrowth in Drosophila. *Cell*. 67:941-953.
Varmark, H., S. Llamazares, E. Rebollo, B. Lange, J. Reina, H. Schwarz, and C. Gonzalez.
2007. Asterless is a centriolar protein required for centrosome function and embryo
development in Drosophila. *Curr. Biol.* 17:1735-1745.
Wang, C., S. Li, J. Januschke, F. Rossi, Y. Izumi, G. Garcia-Alvarez, S.S. Gwee, S.B. Soon,
H.K. Sidhu, F. Yu, F. Matsuzaki, C. Gonzalez, and H. Wang. 2011. An ana2/ctp/mud
complex regulates spindle orientation in Drosophila neuroblasts. *Dev. Cell*. 21:520-533.
Wodarz, A., and W.B. Huttner. 2003. Asymmetric cell division during neurogenesis in
Drosophila and vertebrates. *Mech. Dev.* 120:1297-1309.
Xu, S., N. Hafer, B. Agunwamba, and P. Schedl. 2012. The CPEB protein Orb2 has multiple
functions during spermatogenesis in Drosophila melanogaster. *PLoS Genet*.
8:e1003079.
Yu, F., Y. Cai, R. Kaushik, X. Yang, and W. Chia. 2003. Distinct roles of Galphai and Gbeta13F
subunits of the heterotrimeric G protein complex in the mediation of Drosophila
neuroblast asymmetric divisions. *J Cell Biol*. 162:623-633.
690
691

692   **Abbreviations used:**
693
ACD, asymmetric cell division
AI, asymmetry index
Asl, Asterless
Baz, Bazooka/PAR-3
BC, biologic complexity
Cep135, Centrosomal protein 135kDa
Cnb, Centrobin
Cnn, Centrosomin
γTub, γ-Tubulin
GMC, ganglion mother cell
Mira, Miranda
MT, microtubule
MTOC, microtubule-organizing center
NSC, neural stem cell
PCM, pericentriolar material
PLP, Pericentrin-like protein
Polo, Polo kinase
Wdr62, WD repeat domain 62
wor, worniu
WT, wild-type
714

Figure Legends

Figure 1. Orb2 contributes to centrosome asymmetry. Maximum intensity projections of

interphase NSCs (dashed circles) stained for Asl (centrioles, magenta) and γTub (PCM, cyan).

Solid and dashed boxes mark apical vs. basal centrosomes; enlarged at right. (A) WT NSC with

γTub enriched at the apical centrosome. (B) *orb2* NSC with γTub at both centrosomes. (C) AIs

of Asl and γTub in N=30 NSCs from n=8 WT brains and N=40 NSCs from n=10 *orb2* brains.

Each dot is a measurement from one cell. (D) Frequency distribution of Asl and γTub AIs in WT

vs. *orb2* NSCs. Light grey (outlier) values are > 2 S.D. from the control mean. (E) Scatter plot of

γTub levels at apical vs. basal centrosomes in N=24 NSCs from n=8 WT brains and N=10

symmetrized NSCs from n=7 *orb2* brains (defined as AI >2 S.D. from WT mean). (F) no *GAL4*

control NSCs resemble WT. (G) *orb2^{RNAi}* NSC with γTub at both centrosomes. (H) AIs from

N=39 NSCs from n=6 no GAL4 brains and N=51 NSC from n=6 *orb2^{RNAi}* brains. (I) Frequency

distribution of Asl and γTub AIs in control vs. *orb2^{RNAi}* brains. (J) Scatter plot of γTub at

centrosomes from N=39 NSCs from n=6 control brains and N=16 NSCs from n=6

worGAL4>orb2^{RNAi} brains. Mean ± SD displayed. The experiments were repeated $\geq$3

independent replicates and significance determined by (C, H) Kolmogorov-Smirnov test, (D, I)

chi-square test, (E, J) Mann-Whitney test: n.s., not significant; *, $p \leq 0.05$; **$p \leq 0.01$ ***, $p < 0.001$;

and ****, $p < 0.0001$. Bars: 5 μm, 1 µm (insets).

Figure 2. Orb2 is required for centrosome segregation and spindle morphogenesis.

Maximum intensity projections of interphase NSCs (dashed circles) from (A) WT, (B) *orb2*, (D)

no *GAL4* control, and (E) *orb2^{RNAi}* stained for Asl (arrows). Solid and dashed boxes mark apical

vs. basal centrosomes; enlarged at right. (C and F) Frequency distributions of supernumerary

centrosomes in interphase NSCs. (G) WT and (H) *orb2* NSCs stained for Baz (apical cortex,

cyan), pH3 (mitotic, cyan), Mira (basal cortex, magenta), and Asl (yellow; arrowheads). (I)

Frequency distribution of apical Baz and/or basal Mira crescents in mitotic NSCs. (J) Cartoon

depicts points used to measure within mitotic NSCs: θ_1 spindle orientation, θ_2 apical polarity

alignment, and θ_3 centrosome alignment (*Methods*). (K) WT and (L) *orb2* NSCs stained for MTs

(β-Tub, magenta), Asl (yellow), Baz (cyan), and DAPI (DNA, grey). (M) Plot and (N) frequency

distribution of θ_1 spindle orientation. (O) Plot and (P) frequency distribution of θ_2 apical polarity

alignment. (Q) Plot and (R) frequency distribution of θ_3 centrosome alignment from N=33 WT

and N=32 *orb2* mitotic NSCs from a single experiment across two replicates. Light grey θ are >2

SD from the WT mean. Mean ± SD indicated; significance determined by (C, F, I, N, P, R) Chi-

squared test or (M, O, Q) Two-tailed t-test. n.s., not significant; *, $p < 0.05$; and ****, $p < 0.0001$.

Bars: 5 µm; 1 µm (insets).

Figure 3. NSC-autonomous and non-autonomous Orb2 activities support

neurodevelopment. Projected third instar larval optic lobes (dashed lines) stained for DAPI or

Mira. In age-matched brains, (A) WT are larger than (B) *orb2* and (C) WT have more NSCs than

(D) *orb2*. (E) Volume quantified from N=15 WT and 13 *orb2* brains. (F) Quantification of NSCs in

N=6 WT and 10 *orb2* brains. In age-matched brains, (G) no *GAL4* controls are sized as (H)

*orb2^RNAi^* brains. Yet, (I) no *GAL4* controls have more NSCs than (J) *orb2^RNAi^*. (K) Volume and (L)

NSC counts quantified from N=6 brains per genotype. Brain volumes were measured from one

lobe per brain and normalized to the control mean (Link et al., 2019). Mean ± SD displayed.

Significance determined by Student's t-test: n.s., not significant; *, $p<0.05$***; and $p<0.001$.

Bars: 40 µm.

Figure 4. Orb2 localizes to active NSC centrosomes. (A) Maximum-intensity projections of

WT NSCs (dashed circles) stained for Asl (magenta), Cnn (PCM; yellow), and Orb2 (cyan).

Solid and dashed boxes note apical vs. basal centrosomes; enlarged at right. (B) Orb2 AI and

(C) Orb2 levels on apical vs. basal centrosomes quantified from N=24 late interphase NSCs.

Mean ± SD displayed; **, $p=0.003$ by Welch's t-test. Bars: 5 µm, 1 µm (inset).

Figure 5. Orb2 regulates PLP protein levels. (A) Immunoblot of PLP, Asl, and Orb2 proteins

from WT and *orb2* third instar larval brain extracts. (B) Normalized (to Asl) relative PLP levels in

WT and *orb2* larval brains from 3 biological replicates. (C) Max projected WT and *orb2* NSCs

(dashed circles) stained for γTub (magenta) and PLP (cyan). Solid and dashed boxes note

apical vs. basal centrosomes; enlarged at right. (D) PLP AI calculated from N=24 WT and 32

orb2 interphase NSCs. (E) PLP localization to apical and basal centrosomes in N=24 WT vs.

N=43 *orb2* interphase NSCs. Data representative from a single experiment across 2 replicates.

(F) Model depicts NSC-autonomous function of Orb2 to support interphase centrosome

asymmetry and NSC-independent requirement for normal brain volume. Loss of *orb2* activity

(grey cells) impairs basal centrosome inactivation (PCM, green). Centrosome-localized Orb2

(WT apical; *top box*) enhances centrosome activation, perhaps by translational activation of one

or more targets (black, Orb2 target BC$\geq$3; see Table S1). Conversely, cytoplasmic Orb2

supports basal centrosome inactivation (*bottom box*), required for centrosome separation,

spindle orientation, and centrosome segregation to the daughter cells. Mean $\pm$ SD displayed; *,

p<0.05 by unpaired, one-tailed t-test; n.s., not significant. Uncropped, replicated blots are

available to view: https://doi.org/10.6084/m9.figshare.17052722 Bars: 5 µm; insets 1 µm.

Robinson and Lerit Figure 1. orb2 is required for centrosome asymmetry

Robinson et al.
Figure 2. Orb2 is required for centrosome segregation and spindle morphogenesis

Robinson et al.
Figure 3. NSC-autonomous and non-autonomous Orb2 activities support neurodevelopment

Robinson et al.
Figure 4. Orb2 localizes to active centrosomes

Robinson et al.
Figure 5. Orb2 regulates PLP protein levels

Robinson et al.
Figure S1. NSC differentiation and survival in WT vs. *orb2* mutants

A

Orb2

4206 1083 821

CPEB4

B

Intracellular membrane-bound organelle GO:0043231 — $p=6.7E-55$
Organelle | GO:0043226 — $p=7.5E-61$
Intracellular organelle | GO:0043229 — $p=6.5E-62$
Cellular anatomical entity | GO:0110165 — $p=5.8E-65$
Intracellular anatomical structure GO:0005622 — $p=3.5E-103$

-log(p-value)

C

Microtubule-organizing center GO:0005815 — $p=0.68$
Centrosome | GO:0005813 — $p=.52$
Pericentriloar material | GO:0000242 — $p=.09$
Spindle pole | GO:0000922 — $p=.08$

-log(p-value)

D

Spindle pole | GO:0000922 — $p=.015$
Microtubule-organizing center GO:0005815 — $p=.015$
Centrosome | GO:0005813 — $p=.0075$

-log(p-value)

Robinson et al.
Figure S2. Ontological analysis of Orb2 and human CPEB4