A newly-evolved chimeric lysin motif receptor-like kinase in *Medicago truncatula*

spp. *tricycla* R108 extends its Rhizobia symbiotic partnership

Thi-Bich Luu[1], Anna Ourth[1], Cécile Pouzet[2], Nicolas Pauly[1,3] and Julie Cullimore[1]

[1]Laboratory of Plant-Microbe Interactions and Environment (LIPME), University Toulouse

III, INRAE, CNRS, Castanet-Tolosan, France; [2]FRAIB-TRI Imaging Platform Facilities,

FR AIB, Université de Toulouse, CNRS, 31320 Castanet-Tolosan, France; [3]Institut

Sophia Agrobiotech, Université Côte d'Azur, INRAE, CNRS, Sophia Antipolis Cedex,

France;

Author for correspondence:

Julie Cullimore

Tel: +33 (0)5 61 28 55 13

Email: julie.cullimore @inrae.fr

Total word count (Excluding summary, references, and legends):	4911	No. of figures:	9 (Figs 1-9 in colour)
Summary:	192	No. of Tables:	0
Introduction:	1013	No of Supporting Information files:	2 (Table S1, Fig. S1, Table S2,S4; Table S3)
Materials and Methods:	872		
Results:	1426		
Discussion:	1600		
Acknowledgements:	127		

Summary

- Rhizobial lipochitooligosaccharidic Nod factors, specified by *nod* genes, are the primary determinants of host specificity in the legume-Rhizobia symbiosis. A *Sinorhizobium meliloti nodF/nodL* mutant produces Nod factors that differ from wild-type ones in lacking an *O*-acetate, and with a different acyl chain on the terminal non-reducing sugar. This mutant is defective in nodulation with various *Medicago* hosts.

- We examined the nodulation ability of *M. truncatula* cv Jemalong A17 and *M. truncatula* ssp. *tricycla* R108 with the *nodF/nodL* mutant. We then applied genetic and functional approaches to study the genetic basis and mechanism of nodulation of R108 by this mutant.

- We show that the *nodF/nodL* mutant can nodulate R108 but not A17. Using genomics and reverse genetics, we identified a newly-evolved gene in R108, *LYK2bis*, which is responsible for the phenotype. Transformation with *LYK2bis* allows A17 to gain nodulation with the *nodF/nodL* mutant. We found that LYK2bis is involved in specific NF signalling and interacts with the key receptor protein NFP.

- Our findings reveal that a newly-evolved gene in R108, *LYK2bis*, extends nodulation specificity to strains producing non-*O*-acetylated NFs. Interaction between LYK2bis and NFP provides a means of integrating the nodulation signalling pathways.

Key words: *Medicago truncatula*, legume-Rhizobia symbiosis, LysM receptor-like kinase, Nod factor perception, lipochitooligosaccharide signalling, nodulation specificity

Introduction

The importance of legumes in agriculture and the environment is largely due to their ability to form a N_2-fixing symbiosis with Rhizobia bacteria, allowing them to grow independently of an additional nitrogen source (Graham & Vance, 2003). This ability has stimulated research on the evolution of such a successful mutualism, and the mechanisms that allow it to develop (Oldroyd *et al.*, 2011; Griesmann *et al.*, 2018). Legumes represent a huge family and show considerable variation in their specificity for their bacterial symbiont (Andrews & Andrews, 2017). Indeed, Rhizobia are polyphyletic, defined primarily by their ability to induce root nodules on their legume host, in which they differentiate into nitrogen-fixing forms. Different genera/species/strains may show either a narrow or promiscuous host range (Masson-Boivin *et al.*, 2009). Similarly, some legumes are highly promiscuous whereas others show exquisite fidelity to a narrow range of rhizobia (Andrews & Andrews, 2017; Chen *et al.*, 2021)

Coevolution has ensured that Legume-Rhizobia couples are highly adapted to each other and has led to improvement of plant and bacterial fitness (Younginger & Friesen, 2019). However, as Rhizobia are horizontally transmitted symbionts, host-partner coevolution would not be possible if there were not signalling mechanisms to select the evolved partner from the external microbial milieu. Thus, partner choice signalling is a crucial step in the evolution of a successful symbiosis (Younginger & Friesen, 2019). Nod factors (NF) are key rhizobial signals for initiating and maintaining the interaction between most well-studied legume-Rhizobia couples (Krönauer & Radutoiu, 2021). NFs are lipochitooligosaccharides (LCO) with a basic structure of 3-5 N-acetyl glucosamine residues, *N*-acylated on the terminal non-reducing sugar. Various chemical substitutions on the basic structure lead to the primary mechanism of partner specificity (D'Haeze & Holsters, 2002; Oldroyd *et al.*, 2011; Krönauer & Radutoiu, 2021). Although other signals may be important for some legume-Rhizobia interactions, such as cell-surface polysaccharides or protein effectors (Wang *et al.*, 2018; Chen *et al.*, 2021), the genomic linkage of the NF (*nod*) genes to nitrogen fixation (*nif*) genes (generally on a symbiotic megaplasmid or a genomic island), provides a means by which NF signalling can be used

by the legume host to select its co-evolved and beneficial nitrogen-fixing partner
(Younginger & Friesen, 2019).

On the plant side, lysin motif receptor like kinase (LysM-RLK) genes have evolved as key
elements of NF perception and signalling. They encode plasma-membrane located
proteins with three lysin motifs in the external region connected through a single
transmembrane spanning domain to an internal kinase like domain. These genes are
involved in defence and mychorriza signalling in many plants, but the family is particularly
expanded in legumes, where two genes show neofunctionalization to play essential roles
in NF perception and nodulation (Buendia *et al.*, 2018; Chiu & Paszkowski, 2020). One
of these genes encodes a receptor with an inactive kinase domain and is known as
NFP/NFR5 (according to the species), whereas the other gene encodes a receptor with
an active kinase domain and is known as *LYK3/NFR1* (Krönauer & Radutoiu, 2021). This
latter gene evolved by tandem duplication, prior to the speciation of the legume family
and it has been suggested that its neofunctionalization was fundamental to the
evolutionary gain of a legume-specific NF receptor (De Mita *et al.*, 2014). Further, more
recent duplications of this and other genes in the cluster have led to a source of variation
in NF signalling (De Mita *et al.*, 2014; Sulima *et al.*, 2017).

Medicago truncatula Gaertn. (closely related to alfalfa and pea), together with *Lotus*
japonicus, is a widely used model for studying the molecular mechanisms leading to the
Rhizobial symbiosis and also, more generally, for traits important for the successful use
of legumes in agriculture (Garmier *et al.*, 2017). Initially most work used the genotype *M.*
truncatula ssp. *truncatula* cv Jemalong line A17 (referred to here as A17), but more
recently, genomic resources have been developed for many other genotypes (see the
Hapmap project – https://medicagohapmap2.org/) in order to exploit the considerable
genetic variation of this species for these studies (Garmier *et al.*, 2017). Of note, *M.*
truncatula ssp. *tricycla* line R108-1 (referred to as R108) has achieved considerable
success due to the development of a *Tnt1*-insertion population resource and its superior
genetic transformation properties (Garmier *et al.*, 2017; Kaur *et al.*, 2021). Studies on
A17 have shown that it has a narrow partner specificity, being nodulated primarily by
*Sinorhizobium (*also known as *Ensifer)* species, *S. meliloti* and *S. medicae* (Andrews &

Andrews, 2017; Kazmierczak *et al.*, 2017). Analysis of nodulation by *S. meliloti* strain
2011 has shown that it produces NFs that are 6-*O*-sulphated on the reducing sugar and
contain predominantly a C16:2 acyl chain on the terminal non-reducing sugar, which may
also be 6-*O*-acetylated. The presence of the sulphate (specified by the *nodH* gene) is
essential for nodulation (Roche *et al.*, 1991) and the modifications on the terminal non-
reducing sugar are also important. For example, a double *nodF/nodL* mutant which
couples mutations in *nodF* (leading to N-acylation mainly with a C18:1 fatty acid) and
nodL (leading to lack of 6 *O*-acetylation), is almost completely Nod⁻, whereas a *nodF* or
a *nodL* mutant induce similar or a reduced number of nodules (Ardourel *et al.*, 1994; Smit
et al., 2007). On the plant side, *NFP* is required for all NF responses and its role can be
partially substituted by *NFP* genes from other species (Bensmihen *et al.*, 2011; Girardin
et al., 2019). Studies on a weak *lyk3* mutant of A17 (Smit *et al.*, 2007) and recent structural
studies (Bozsoki *et al.*, 2020) suggest that *LYK3* is involved in the perception of specific
NF features.

Comparative studies between A17 and R108 have shown that they are highly divergent
members of the *M. truncatula* species (Zhou *et al.*, 2017), show differences in their
symbiotic effectiveness with different Rhizobial strains (Kazmierczak *et al.*, 2017) and
exhibit differences in partner choice when confronted with a mixed *Sinorhizobium*
(*Ensifer*) inoculum (Burghardt *et al.*, 2018). In this article, we report that R108 has an
extended NF-related strain specificity compared to A17 and have identified a newly
evolved LysM-RLK gene, unique to R108, which is required for this phenotype.

Materials and methods

Seed germination and growth conditions

Medicago seeds were scarified in 95% sulphuric acid for 5 min, washed with distilled
water two times and surface-sterilized in bleach (3.2% chlorine) for 3 min, then washed
three times. After that, the seeds were soaked in water for 1 h and placed on 1% agar,
supplemented with 1 µg ml⁻¹ GA3. The plates were kept upside-down at 4ºC for 5 d and
then put at 16ºC overnight for germination. Seedlings were transferred to pots and grown
in a growth chamber at 22ºC with a 16 h photoperiod.

cDNA cloning

R108 roots were collected and immediately frozen in liquid N_2. Total RNA was extracted using a Nucleospin plant RNA extraction kit (Marcherey-Nagel GmbH & Co. KG, Germany). cDNA was synthesized using SuperScript IV Reverse Transcriptase kit (Thermo Fisher Scientific, USA). Briefly, 300 ng of total RNA was incubated with 50 μM Oligo dT and 10 mM dNTPs in a reaction tube at 65ºC for 5 min, then put on ice for 1 min. After that, SuperScript IV Reverse Transcriptase with SSIV buffer, DTT and RNAase inhibitor were added and incubated in a thermocycler at 55ºC for 30 min, followed by an enzyme deactivation step at 80ºC for 10 min.

LYK2bis coding sequence (CDS) was amplified with Phusion High Fidelity DNA Polymerase (F530S, Thermo Fisher Scientific, USA) using R108 root cDNA as template and the following primers (5'-ATGAAACTAAAAAATGGCTTAC-3' and 5'-TCTAGTTGACAACAAATTTATG-3'). The PCR product was cloned into pJet1.2/blunt using CloneJET PCR Cloning Kit (K1231, Thermo Fisher Scientific, USA). The correct clone was selected by sequencing, then used for further cloning steps.

Constructs for in planta *protein expression*

For *M. truncatula* root transformation, full length CDS of *LYK2bis* and of *LYK3* from A17 (LYK3-A17) were fused at the C-terminus with 3xFLAG under the control of the Ubiquitin promoter from *L. japonicus* (ProLjUbi) by Golden gate cloning, using a vector based on pCAMBIA2200 expressing DsRED (Fliegmann *et al.*, 2016). For *Nicotiana benthamiana* leaf agro-infiltration, full length CDS of LYK2bis and NFP were fused at the C-terminus with mCherry and GFP tag proteins, respectively, under the control of ProLjUbi. The vector pCambia2200ΔDsRED was used (Fliegmann *et al.*, 2016).

Mutants and nodulation tests

Tnt1 insertional mutants, *lyk2-1R* (NF13076), *lyk2bis-1R* (NF15454) and *lyk3-1R* (NF2752) of R108, obtained from the Noble Research Institute (USA), were used for nodulation tests. R108 and A17 were used as wild-type (WT) controls. Note that "*R*" designates a R108 mutant.

Germinated seedlings were transferred into pots (8x8x8 cm^3 in size), filled with sterilized

attapulgite clay granules (Oil Dri, UK). To each pot, 80 ml of Fahraeus medium

supplemented with 1 mM $CaCl_2$ and 0.2 mM NH_4NO_3, was added. After 3 d, plants were

inoculated with approximately 10,000 bacteria/plant of *S. meliloti* 2011 WT or mutant

strains (Table **S1**). The number of nodules per plant was counted at 21 dpi. When

appropriate, bacterial infection was determined by LacZ straining with X-gal. All results

were confirmed in separate biological experiments.

Complementation and gain-of-function assays

Using *Agrobacterium rhizogenes*-mediated transformation (Boisson-Dernier *et al.*, 2001),

seedlings of *lyk2bis-1R* or A17 were transformed using strains containing either empty

vector (EV) or ProLjUb:LYK2bis-3xFLAG or ProLjUb:LYK3-A17-3xFLAG constructs, and

transformants were selected on medium containing 25 μg ml^{-1} kanamycin, and after two-

weeks growth, by expression of the DsRed marker. R108 seedlings transformed with EV

were used as control. Nodulation was analysed in pots as above after 4 wk. The number

of nodulated plants and the number of nodules/nodulated plant were analysed.

Agro-infiltration and Fluorescence Lifetime Imaging on Nicotiana benthamiana leaves

Agrobacterium tumefaciens LBA4404 strains containing ProLjUb:LYK2bis-mCherry or

NFP-GFP fusion constructs were used to agro-infiltrate the three to four oldest leaves of

each *N. benthamiana* plant in which the OD_{600} of LYK2bis-mCherry was five times higher

than that of NFP-GFP in order to have similar expression levels.

Two days post infiltration, the protein expression in leaves was assessed by confocal

microscopy. Forster resonance energy transfer (FRET) between the fluorophores was

analysed by Fluorescence Lifetime Imaging Measurements (FLIM) on a Leica TCS SP8

SMD which consists of an inverted LEICA DMi8 microscope equipped with a TCSPC

system from PicoQuant. The excitation of the FITC donor at 470 nm was carried out by a

picosecond pulsed diode laser at a repetition rate of 40 MHz, through an oil immersion

objective (63×, N.A. 1.4). The emitted light was detected by a Leica HyD detector in the

500-550 nm emission range. Images were acquired with acquisition photons of up to 1500

per pixel.

From the fluorescence lifetime images, the decay curves were calculated per pixel and

fitted (by Poissonian maximum likelihood estimation) with either a mono- or double-

exponential decay model using the SymphoTime 64 software (PicoQuant, Germany). The

mono-exponential model function was applied for donor samples with only GFP present.

The double-exponential model function was used for sample containing GFP and

mCherry. Experiments were repeated at least three times. The efficiency of energy

transfer (E) based on the fluorescence lifetime (τ) was calculated as $E = 1 - (\tau_{D+A}/\tau_{D-A})$,

where τ_{D+A} is donor fluorescence lifetime in the presence of acceptor while τ_{D-A} is the

donor fluorescence lifetime in the absence of acceptor.

Kinase assays

Glutathione-S-transferase (GST) tagged proteins of the predicted intracellular region of

LYK2bis (termed the KD) were expressed in *E. coli* DH5α and the proteins purified using

glutathione resin (GE Healthcare, USA) as described (Fliegmann *et al.*, 2016). LYK2bis-

KD was released from the resin using PreScission Protease (GE27-0843-01, Sigma

Aldrich, Germany). LYK2bis-KD was incubated with kinase buffer containing ^{32}P-ATP

alone or with purified GST/NFP-KD, GST/LYR2-KD, GST/LYR3-KD, GST/LYR4-KD,

GST/LYK3-deadKD (G334E mutation), Myelin Basic Protein (MyBP) or GST at 25ºC for

1 h and the proteins analysed by SDS-PAGE, followed by Coomassie staining and

Phosphor Imaging.

Mycorrhization tests

Mycorrhization tests were performed using the gridline intersect method as described in

Gibelin-Viala *et al.* (2019). Briefly, *lyk2bis-1R* and R108 seedlings were inoculated with

200 spores per plant of *Rhizophagus irregularis* DAOM197198 (Agronutrition, Toulouse,

France) and colonisation was assessed at 3- and 5-wk post inoculation (wpi) using 15

plants/genotype/time-point.

Bioinformatic analysis

The 1st exon encoding the whole LysM domain of *LYK2* (640 bases), *LYK3* (637 bases)

and *LYK2bis* (640 bases) from R108 were used with blastn to screen all *M. truncatula*

genomes available in the Medicago BLAST Service of the Hapmap2 database

(https://medicagohapmap2.org/). The first hit for each query was extracted. The

percentages of identity (%ID) between hits found with the LYK3R108 probe and the

LYK2bis sequence were calculated by EMBOSS Needle.

Results

R108 shows an extended nodulation specificity with a S. meliloti nodF/nodL mutant strain

To determine the nodulation specificity of the *M. truncatula* R108 genotype, seedlings of

R108 and A17 were inoculated with either *S. meliloti* 2011 WT or its *nodF/nodL* mutant.

The number of nodules was counted at 21 dpi. No nodules were found on A17 plants

inoculated with *nodF/nodL,* which is consistent with the results on *M. truncatula* cv.

Jemalong obtained previously (Ardourel *et al.*, 1994) (Fig. **1**). However, R108 showed a

similar nodulation capacity with both WT and *nodF/nodL* strains (Fig. **1**). Most (> 95%) of

the R108 nodules with both strains were pink, and LacZ staining showed that they were

well-infected with bacteria. This suggests that R108 has an extended nodulation

specificity, compared to A17, which includes the *nodF/nodL* mutant.

A newly-evolved gene in R108, LYK2bis, is required for nodulation with the nodF/nodL

strain

The perception of NF and nodulation in *M. truncatula* have been shown to require LysM-

RLKs, which are NFP and LYK3 in A17 (Arrighi *et al.*, 2006; Smit *et al.*, 2007). In the R108

genotype, NFP is reported to play essential roles for these processes (Feng *et al.*, 2019),

therefore, to understand the extension of nodulation in this ecotype, we focused on the

LYK gene cluster, close to *LYK3*.

Using the genome sequence of R108 v1.0 (Kaur *et al.*, 2021), we located the *LYK* gene

cluster on chromosome 5, using the sequences of *LYK2* (MtrunA17Chr5g0439701) and

LYK3 (MtrunA17Chr5g0439631) from A17. Between *LYK2* and *LYK3*, an additional

LysM-RLK gene was found, which is not present in A17, and is designated as *LYK2bis*

(Fig. **2a**). Notably, the exon/intron organization is conserved among these genes with 12

exons in each (Fig. **2a**). The predicted amino acid sequence alignment between LYK2,

LYK2bis and LYK3 proteins of A17 and R108 indicates that LYK2bis is a chimera with a

similar extracellular domain to LYK2 up to the middle of LysM3 and then shares a highly

conserved sequence with LYK3 (Fig. **2b**, Fig. **S1** and Table **S2**). In order to examine the

presence of *LYK2bis* in *M. truncatula* ecotypes, the nucleotide sequence of the first exon

of *LYK2*, *LYK3* and *LYK2bis* (containing the whole LysM region) were used to screen the

23 other *M. truncatula* genomes available for BLAST screening in the Hapmap project

(https://medicagohapmap2.org/). Separate sequences corresponding to *LYK2* and *LYK3*

were identified in all cases (Table **S3**). However, in genotype HM026, the available *LYK3*

sequence seems to be not complete. The *LYK2bis* probe identified primarily the *LYK2*

gene in all the genomes, with a lesser homology of this exon to the *LYK3* sequence. We

did not find any evidence for an additional *LYK2* or *LYK3*-like gene in these genomes.

These results suggest that *LYK2bis* is unique to R108 in these *M. truncatula* Hapmap

genomes.

To determine the roles of these genes in nodulation, *Tnt1* insertional mutants of *LYK2*,

LYK2bis and *LYK3* in the R108 background were identified from the Noble Foundation

resources (Fig. **2b**). The mutants were inoculated with either *S. meliloti* WT or *nodF/nodL*

strains and phenotyped at 21 dpi. All tested lines formed a similar number of nodules with

the WT strain (Fig. **2c**) indicating that, unlike the crucial role of *LYK3* in A17, none of those

genes plays an essential role in nodulation of R108 with the *S. meliloti* WT strain.

Similarly, there was no significant difference in the number of nodules formed in *lyk2-1R*

and *lyk3-1R* mutants with the *nodF/nodL* strain, compared to R108. In contrast, no

nodules formed on *lyk2bis-1R* plants inoculated with *nodF/nodL* suggesting that the

LYK2bis gene is responsible for nodulation of R108 with the *nodF/nodL* mutant strain.

To validate the involvement of *LYK2bis* in nodulation with the *nodF/nodL* strain, we

performed a complementation experiment of the *lyk2bis-1R* mutant. Using *A. rhizogenes*-

mediated transformation, constructs of *LYK2bis*, *LYK3-A17* and empty vector (EV) were

expressed in the *lyk2bis-R1* mutant. As shown in Fig. **3**, while neither *LYK3* nor EV

transformed plants formed nodules with the *nodF/nodL* strain, nodulation was restored in

most plants transformed with *LYK2bis*. This result confirms the essential role of *LYK2bis*

in nodulation with *nodF/nodL* in R108.

LYK2bis allows A17 to gain the ability to nodulate with the nodF/nodL strain

As A17 is not able to nodulate with the *nodF/nodL* strain and does not contain a *LYK2bis*

gene in the genome, we performed a gain-of-function assay in A17 to determine whether

introduction of *LYK2bis* is sufficient to gain nodulation with the *nodF/nodL* strain in this

genotype. The experiment was done using a similar approach to the *lyk2bis-1R*

complementation. Most of the A17 plants transformed with the *LYK2bis* construct

nodulated with *nodF/nodL* while none of the ones transformed with *LYK3* and EV formed

nodules (Fig. **4**). This evidence reveals that *LYK2bis* is central for interacting and

nodulating with the *nodF/nodL* strain and can transfer this ability to another genotype.

LYK2bis is involved in the perception of specific NF decorations leading to successful

nodulation in R108

S. meliloti nodF/nodL produces NFs that differ from those of the WT strain in lacking the

O-acetylation and containing C18:1 fatty acid chains rather than C16 on the terminal non-

reducing sugar (Ardourel *et al.*, 1994). To determine the specificity of NF decorations

recognized by *LYK2bis*, we tested the nodulation of *lyk2bis-1R* with two other *S. meliloti*

mutant strains: *nodFE* which produces NFs with the C18:1 acyl chain and *nodL* which

produces non-*O*-acetylated NFs (Ardourel *et al.*, 1994). All tested strains could form

nodules on R108 (Fig. **5**). While a similar number of nodules formed on *lyk2bis-1R* and

R108 with the *nodFE* strain, no nodules were found on *lyk2bis-1R* inoculated with *nodL*

(Fig. **5**). This result suggests that while other protein(s) can recognize rhizobia producing

C18:1 NFs in R108, LYK2bis is particularly involved in the perception of rhizobia

producing non-*O*-acetylated NFs.

LYK2bis does not play an important role in mycorrhization

As some Myc-factors have an identical structure to the major *nodF/nodL* NFs (Maillet *et*

al., 2011), we also tested the role of *LYK2bis* in forming arbuscular mycorrhiza (AM).

Seedlings of *lyk2bis-1R* and R108 were inoculated with spores of *Rhizophagus irregularis*

and phenotyped for AM colonization at 3 wpi and 5 wpi. Both genotypes showed good

mycorrhization at both time points (Fig**. 6**), indicating that *LYK2bis* is not essential for
mycorrhization.

LYK2bis physically interacts with NFP

NFP is required for nodulation in R108 (Feng *et al.*, 2019) as well as in A17 (Arrighi *et al.*,
2006), therefore, to test whether LYK2bis could be part of an NFP receptor complex, we
investigated whether the two proteins interact using FRET-FLIM technology.

A construct of the LYK2bis protein fused with mCherry was co-expressed with a construct
of a NFP-GFP fusion in *N. benthamiana* leaves using *A. tumefaciens*-mediated
expression. Interestingly, at 3 dpi, all leaves expressing both proteins showed strong cell-
death (Fig. **7a**) similar to that observed in the co-expression of NFP and LYK3-A17
(Pietraszewska-Bogiel *et al.*, 2013). This suggests that LYK2bis may functionally interact
with NFP in a similar mechanism to LYK3.

At 2 dpi, the cell death response was much reduced and confocal microscopy revealed
that the two proteins co-localise at the periphery of the cells (Fig. **7b**). As NFP was
previously reported to localise at the plasma membrane of *N. benthamiana* leaves
(Lefebvre *et al.*, 2012), this evidence indicates that LYK2bis co-localises with NFP at the
plasma membrane.

Similar leaf samples were used for FRET FLIM analysis in which NFP-GFP was used as
the donor and LYK2bis-mCherry was the acceptor. A highly significant decrease in
lifetime of NFP-GFP was obtained when co-expressed with LYK2bis-mCherry (Fig. **7c**).
This data not only supports the co-localization of LYK2bis and NFP at the plasma
membrane but also provides strong evidence for the physical interaction between the two
proteins.

LYK2bis possesses an active kinase and can transphosphorylate the pseudo kinase of
NFP

329 NFP has an inactive kinase domain (KD) which can be weakly trans-phosphorylated by
330 the active kinase domain of LYK3 (Arrighi *et al.*, 2006; Fliegmann *et al.*, 2016). To test
331 whether LYK2bis has an active kinase domain and can trans-phosphorylate NFP, we

performed *in vitro* phosphorylation assays on the intracellular regions of the two proteins (which includes the KD), purified after expression in *E. coli*. We also tested the ability of LYK2bis-KD to trans-phosphorylate related proteins, LYR2-KD, LYR3-KD, LYR4-KD and LYK3-deadKD (Fliegmann *et al.*, 2016), and the model kinase substrate myelin basic protein (MyBP). LYK2bis-KD had strong autophosphorylation activity and could trans-phosphorylate GST fusions of NFP-KD, LYR2-KD, LYR3-KD, LYR4-KD and LYK3-deadKD and MyBP in presence of [ɣ-^{32}P] ATP, but not GST (Fig. **8**). These results show that LYK2bis has an active kinase domain and can trans-phosphorylate NFP, some other kinase domains and the model kinase substrate, MyBP.

Discussion

The symbiotic interactions between legumes and rhizobia are highly specific and require proper recognition of bacterial signals by plant receptors. NFs are key determinants of host specificity due to their species-specific chemical substitutions on both the reducing and non-reducing ends of the NF, which are recognized by their compatible hosts. In this study, we demonstrate that two commonly used ecotypes of *M. truncatula*, A17 and R108, display a contrasting nodulation specificity with the *S. meliloti nodF/nodL* mutant; while A17 has a completely Nod$^-$ phenotype with this strain, R108 is able to form fully-infected nodules with both the *S. meliloti* WT and *nodF/nodL* strains (Fig. **1**). Using reverse genetics, we have identified *LYK2bis* as the genetic determinant underlying this extension of nodulation specificity. Complementation studies have clearly confirmed the essential role of *LYK2bis* in nodulation with the *nodF/nodL* strain of R108 (Fig. **3**). Moreover, this exceptional characteristic can be transferred to another genotype. As shown in Fig. **4**, the introduction of *LYK2bis* into A17 allows the plants to form nodules with the *nodF/nodL* mutant strain. These results provide conclusive evidence for the role of *LYK2bis* in nodulation with the *nodF/nodL* strain in R108.

The evolution of LYK2bis

LYK2bis is predicted to encode a LysM-RLK, which is located between *LYK2* and *LYK3* on chromosome 5 of the *M. truncatula* R108 genome (Fig. **2**). It is a chimeric gene with most of the LysM domains of *LYK2* and the rest, including the kinase domain, of *LYK3* (Fig. **S1**). Such a gene does not occur in other legumes such as *L. japonicus*, soybean

and pea (De Mita *et al.*, 2014; Sulima *et al.*, 2017). In *M. truncatula*, although all the 23

genotypes available for BLAST screening in the Hapmap collection

(https://medicagohapmap2.org/) contain sequences highly related to *LYK2* and *LYK3*

(Table **S3**), only R108 contains *LYK2bis*. R108 is the most distant of the *M. truncatula*

genomes (Zhou *et al.*, 2017), and the only one representing *M. truncatula* ssp. *tricycla*.

The most parsimonious explanation for the presence of *LYK2bis* in R108 is that it is a

newly evolved gene in this genotype/subspecies, although we cannot exclude the

possibility that it pre-existed in the most recent common ancestor of *M. truncatula* and

has been lost at an early stage of the divergence of the two *M. truncatula* subspecies. In

either case its phenotypic attributes suggest that it is a good example of recent adaptive

gene duplication (Kondrashov, 2012).

The specificity of nodulation directed by LYK2bis

The *nodF/nodL* strain is a double mutant of *nodF* and *nodL* genes, which in the WT strain

are responsible for C16:2 *N*-acylation (*nodF*) and *O*-acetylation (*nodL*) on the non-

reducing sugar of NFs: the *nodF/nodL* strain thus produces non-*O*-acetylated NFs, with

predominantly C18 acyl chains (Ardourel *et al.*, 1994). Using various *Medicago* species,

the authors reported that most of the tested hosts were unable to induce nodules with the

nodF/nodL strain, including *M. truncatula* cv Jemalong which has been confirmed in our

study using *M. truncatula* cv Jemalong A17. A17 shows only a slightly reduced nodulation

with a *nodF* mutant but a much reduced nodulation with a *nodL* mutant, and this has been

attributed to the *LYK3* gene (Smit *et al.*, 2007), which may thus show a preference for

recognizing *O*-acetylated NFs, but may tolerate C18 ones. In our study, *LYK2bis* was

found to play essential roles in nodulation of R108 with both *nodF/nodL* and *nodL* but not

with the *nodFE* mutant strain. This evidence indicates a specific role of *LYK2bis* in

nodulation with strains producing non-*O*-acetylated NFs whereas other gene(s) in R108

can enable nodulation with strains producing *O*-acetylated NFs, including ones without

the specific C16 acyl chain. This may be due to *LYK3* as suggested above for A17, and

it is relevant to note that in pea, nodulation with C18:1 NFs produced by a *Rhizobium*

leguminosarum nodE mutant is associated with certain haplotypes of *PsSYM37,* which is

orthologous and functionally equivalent to *LYK3/NFR1* (Li *et al.*, 2011).

In pea, an interesting phenotype linked to *O*-acetylation of the reducing sugar of NFs has

been identified in cv Afghanistan: nodulation by *R. leguminosarum* requires this particular

NF structural modification, which is produced by strains containing the *nodX* gene (Firmin

et al., 1993). On the plant side, *PsSym2* was identified as the determinant of this strain

selectivity, and incidentally, synteny with closely related *M. truncatula* led to the

identification of the *LYK* cluster, containing genes *LYK1 - LYK7* (Limpens *et al.*, 2003).

Recent studies have identified three *LysM-RLKs* in the corresponding pea *LYK* cluster:

PsSYM37, *PsK1* and *PsLYKX,* all of which play roles in nodulation (Zhukov *et al.*, 2008;

Li *et al.*, 2011; Kirienko *et al.*, 2019; Sulima *et al.*, 2017). By analysing different pea

cultivars, it has been shown that nodulation with *nodX*[+] strains is correlated with

haplotypes of the *PsLYKX* gene, but not *PsSYM37* or *PsK1*, suggesting that *PsLYKX*

corresponds to *Sym2* (Sulima *et al.*, 2017, 2019).

Nevertheless, there are also factors that regulate nodulation specificity beyond NF

signalling. In soybean, a plant resistance gene has been identified in which allelic variants

restrict nodulation with specific *Bradyrhizobium japonicum* and *Sinorhizobium fredii*

strains (Yang *et al.*, 2010). The involvement of the plant immune responses in this

interaction is dependent on the type 3 secretory system (T3SS) and the recognition of

rhizobial effectors (Wang *et al.*, 2018). In *M. truncatula* the natural variation in nodulation

with the *S. meliloti* Rm41 strain is reported to be due to the restriction of bacterial infection

but not recognition of NFs, but is independent of a T3SS (Liu *et al.*, 2014). The authors

identified the genetic basis of this phenotype as a single dominant gene, *M. truncatula*

NODULATION SPECIFICITY 1 (Mt-NS1), which may trigger plant defence responses

restricting root nodule symbiosis. In addition, Rhizobial extracellular polysaccharides of

various structures, are important factors of nodulation efficiency (Marczak *et al.*, 2017;

Maillet *et al.*, 2020). In *Lotus* a promiscuity locus permitting nodulation of *L. burttii* with

rhizobia from 5 different genera has been located near the *LYK* cluster but is not due to

NFR1 (Zarrabian *et al.*, 2021).

The importance of LYK2bis structure for NF recognition

Recently, Bozsoki *et al.* (2020) have published the crystal structure of the extracellular

domain of LYK3-A17. By using a domain swapping and point mutation approach, the

authors have identified the importance of two regions in the LysM1 domain of LYK3-A17

for the specificity of NF signalling. As shown in Fig. **S1**, the extracellular domain of

LYK2bis shows strong divergence to LYK3-A17, especially in LysM1 (Table **S3,** Table

S4). In particular, the two regions II and IV of LysM1 that are essential in LYK3 for

discrimination of NF and CO ligands and for specific nodulation are very poorly conserved

in LYK2bis. These regions could therefore be involved in the discrimination of non-*O*-

acetylated and *O*-acetylated NFs. In pea, nodulation with *nodX*[+] strains is associated with

LYKX haplotypes containing specific amino acids in regions II and IV of LysM1, as

described above (Sulima *et al.*, 2019; Solovev *et al.*, 2021). Together, this evidence

indicates the importance of specific regions in the LysM1 of these highly-related LYK

cluster proteins in the recognition of specific NF decorations.

Mechanism of action of LYK2bis

MtNFP, and its orthologs in *L. japonicus* (NFR1)*, P. sativum* (SYM10) and other legumes,

is the key LysM-RLK involved in NF perception and signalling (Krönauer & Radutoiu,

2021). Many studies have shown the involvement of this protein and one or two LYK

cluster proteins in perception of NFs and signalling for nodulation (Radutoiu *et al.*, 2003;

Kirienko *et al.*, 2019). In this study, we observed that the co-expression of LYK2bis and

NFP in *N. benthamiana* leaves leads to defence-like responses, which were also

observed in co-expression of LYK3-A17 and NFP, indicating a similar mechanism of

action between these two protein pairs. In addition, we have shown that LYK2bis

physically interacts with NFP in *N. benthamiana* leaves suggesting the formation of a

heteromer *in planta,* which is possibly responsible for recognition of non-*O*-acetylated

NFs. This hypothesis is supported by the work on molecular modelling of the interaction

between PsLYKX and PsSYM10 receptor complex with NFs (Solovev *et al.*, 2021). In this

study, using molecular modelling and ligand docking, the authors have shown the

potential for stable heterodimers between PsLYKX and PsSYM10, which may interact

with NFs at the heterodimer interface. Indeed in LYK3, region III of LysM1 is envisaged

to interact with NFP (Bozsoki *et al.*, 2020), and is quite highly conserved between LYK2bis

and LYK3 (7/12 residues – Fig. **S1**). Moreover, the LYK2bis intracellular domain is almost

identical to that of LYK3-A17 (three amino acids in difference) and both possess an active

kinase, which *in vitro* can trans-phosphorylate NFP-KD, albeit non-specifically (Fig. **8** and Fliegmann *et al.*, 2016). This similarity suggests a mechanism by which LYK2bis could integrate into the LYK3 signal transduction pathway. Based on data obtained in this study, we propose a model for the nodulation of R108 in which LYK2bis forms a receptor complex with NFP that perceives non-*O*-acetylated NFs and activates the kinase of LYK2bis, which then trans-phosphorylates NFP leading to downstream signalling and finally to nodule formation (Fig. **9**).

In conclusion, we have identified a newly-evolved gene in R108, *LYK2bis*, which is responsible for extending the nodulation specificity of R108 to include the *nodF/nodL* mutant strain. *LYK2bis* is a chimera formed from the *LYK2* and *LYK3* genes and is located between them in the *LYK* cluster on chromosome 5. Studies on the diversity of this locus in the *tricycla* subspecies coupled with analysis of rhizobial symbionts, would establish how the evolution of this gene in *M. truncatula* has led to host-range expansion.

Acknowledgements

We thank Clare Gough, Sandra Bensmihen, Benoit Lefebvre, Benjamin Gourion (LIPME) and Pierre-Marc Delaux and Maxime Bonhomme (Laboratoire de Recherche en Sciences Végétales, Toulouse) for discussions and for their comments on the manuscript. Funding for part of this work was gratefully received from the Fédération de Recherche Agrobiosciences, Interactions et Biodiversité (FR AIB – project CHAIN, coordinators J. Cullimore and C. Jacquet) and the Agence Nationale de la Recherche (ANR – project DUALITY, ANR-20-CE20-0017-01, coordinator C. Gough). This study is set within the framework of the "Laboratoires d'Excellences (LABEX)" TULIP (ANR 10 LABX 41) and of the "École Universitaire de Recherche (EUR)" TULIP GS (ANR 18 EURE 0019). TBL gratefully acknowledges receipt of a Bourse d'Excellence from the Ambassade de France au Vietnam, to fund her PhD in France.

References

Andrews M, Andrews ME. **2017**. Specificity in legume-Rhizobia symbioses. *International Journal of Molecular Sciences* **18**: 705.

Ardourel M, Demont N, Debelle F, Maillet F, de Billy F, Prome JC, Denarie J, Truchet G. **1994**. *Rhizobium meliloti* lipooligosaccharide nodulation factors: Different structural requirements for bacterial entry into target root hair cells and induction of plant symbiotic developmental responses. *Plant Cell* **6**: 1357–1374.

Arrighi JF, Barre A, Ben Amor B, Bersoult A, Soriano LC, Mirabella R, De Carvalho-Niebel F, Journet EP, Ghérardi M, Huguet T, *et al.* 2006. The *Medicago truncatula* lysin motif-receptor-like kinase gene family includes *NFP* and new nodule-expressed genes. *Plant Physiology* **142**: 265–279.

Bensmihen S, Billy F de, Gough C. **2011**. Contribution of NFP LysM domains to the recognition of Nod factors during the *Medicago truncatula/Sinorhizobium meliloti* symbiosis. *PLoS ONE* **6**: e26114.

Boisson-Dernier A, Chabaud M, Garcia F, Bécard G, Rosenberg C, Barker D-G. **2001**. *Agrobacterium rhizogenes*-transformed roots of *Medicago truncatula* for the study of nitrogen-fixing and endomycorrhizal symbiotic associations. *Molecular Plant-Microbe Interactions* **14**: 695–700.

Bozsoki Z, Gysel K, Hansen SB, Lironi D, Krönauer C, Feng F, de Jong N, Vinther M, Kamble M, Thygesen MB, *et al.* 2020. Ligand-recognizing motifs in plant LysM receptors are major determinants of specificity. *Science* **369**: 663–670.

Buendia L, Girardin A, Wang T, Cottret L, Lefebvre B. **2018**. LysM receptor-like kinase and LysM receptor-like protein families: An update on phylogeny and functional characterization. *Frontiers in Plant Science* **871**: 1531.

Burghardt LT, Epstein B, Guhlin J, Nelson MS, Taylor MR, Young ND, Sadowsky MJ, Tiffin P. **2018**. Select and resequence reveals relative fitness of bacteria in symbiotic and free-living environments. *Proceedings of the National Academy of Sciences of the United States of America* **115**: 2425-2430.

Chen WF, Wang ET, Ji ZJ, Zhang JJ. **2021**. Recent development and new insight of diversification and symbiosis specificity of legume rhizobia: mechanism and application. *Journal of Applied Microbiology* **131**: 553–563.

Chiu CH, Paszkowski U. **2020**. Receptor-like kinases sustain symbiotic scrutiny. *Plant*
Physiology **182**: 1597-1612.

De Mita S, Streng A, Bisseling T, Geurts R. **2014**. Evolution of a symbiotic receptor
through gene duplications in the legume-rhizobium mutualism. *New Phytologist* **201**:
961–972.

D'Haeze W, Holsters M. **2002**. Nod factor structures, responses, and perception during
initiation of nodule development. *Glycobiology* **12**: 79R-105R.

Feng F, Sun J, Radhakrishnan G V., Lee T, Bozsóki Z, Fort S, Gavrin A, Gysel K,
Thygesen MB, Andersen KR, *et al.* 2019. A combination of chitooligosaccharide and
lipochitooligosaccharide recognition promotes arbuscular mycorrhizal associations in
Medicago truncatula. *Nature Communications* **10**: 1–12.

Firmin JL, Wilson KE, Carlson RW, Davies AE, Downie JA. **1993**. Resistance to
nodulation of cv. Afghanistan peas is overcome by *nodX*, which mediates an *O*-
acetylation of the *Rhizobium leguminosarum* lipo-oligosaccharide nodulation factor.
Molecular Microbiology **10**: 351–360.

Fliegmann J, Jauneau A, Pichereaux C, Rosenberg C, Gasciolli V, Timmers ACJ,
Burlet-Schiltz O, Cullimore J, Bono JJ. **2016**. LYR3, a high-affinity LCO-binding
protein of *Medicago truncatula*, interacts with LYK3, a key symbiotic receptor. *FEBS*
Letters **590**: 1477–1487.

Garmier M, Gentzbittel L, Wen J, Mysore KS, Ratet P. **2017**. *Medicago truncatula*:
genetic and genomic resources. *Current Protocols in Plant Biology* **2**: 318–349.

Gibelin-Viala C, Amblard E, Puech-Pages V, Bonhomme M, Garcia M, Bascaules-
Bedin A, Fliegmann J, Wen J, Mysore KS, le Signor C, *et al.* 2019. The *Medicago*
truncatula LysM receptor-like kinase LYK9 plays a dual role in immunity and the
arbuscular mycorrhizal symbiosis. *New Phytologist* **223**: 1516–1529.

Girardin A, Wang T, Ding Y, Keller J, Buendia L, Gaston M, Ribeyre C, Gasciolli V,
Auriac M-C, Vernié T, *et al.* 2019. LCO receptors involved in arbuscular mycorrhiza
are functional for Rhizobia perception in legumes. *Current Biology* **29**: 4249–4259.

Graham PH, Vance CP. **2003**. Legumes: importance and constraints to greater use.
Plant Physiology **131**: 872–877.

Griesmann M, Chang Y, Liu X, Song Y, Haberer G, Crook MB, Billault-Penneteau B,

Lauressergues D, Keller J, Imanishi L, *et al.* 2018. Phylogenomics reveals multiple losses of nitrogen-fixing root nodule symbiosis. *Science* **361**: eaat1743.

Kaur P, Lui C, Dudchenko O, Nandety RS, Hurgobin B, Pham M, Aiden EL, Wen J, Mysore KS. **2021**. Delineating the *Tnt1* insertion landscape of the model legume *Medicago truncatula* cv. R108 at the Hi-C resolution using a chromosome-length genome assembly. *International Journal of Molecular Sciences* **22**: 4326.

Kazmierczak T, Nagymihály M, Lamouche F, Barrière Q, Guefrachi I, Alunni B, Ouadghiri M, Ibijbijen J, Kondorosi É, Mergaert P, *et al.* 2017. Specific host-responsive associations between *Medicago truncatula* accessions and Sinorhizobium strains. *Molecular Plant-Microbe Interactions* **30**: 399–409.

Kirienko AN, Vishnevskaya NA, Kitaeva AB, Shtark OY, Kozyulina PY, Thompson R, Dalmais M, Bendahmane A, Tikhonovich IA, Dolgikh EA. **2019**. Structural variations in LysM domains of LysM-RLK psK1 may result in a different effect on Pea–Rhizobial symbiosis development. *International Journal of Molecular Sciences* **20**: 1624.

Kondrashov FA. **2012**. Gene duplication as a mechanism of genomic adaptation to a changing environment. *Proceedings of the Royal Society B: Biological Sciences* **279**: 5048-5057.

Krönauer C, Radutoiu S. **2021**. Understanding Nod factor signalling paves the way for targeted engineering in legumes and non-legumes. *Current Opinion in Plant Biology* **62**: 102026.

Lefebvre B, Klaus-Heisen D, Pietraszewska-Bogiel A, Hervé C, Camut S, Auriac MC, Gasciolli V, Nurisso A, Gadella TWJ, Cullimore J. **2012**. Role of *N*-glycosylation sites and CXC motifs in trafficking of *Medicago truncatula* Nod factor perception protein to plasma membrane. *Journal of Biological Chemistry* **287**: 10812–10823.

Li R, Knox MR, Edwards A, Hogg B, Ellis THN, Wei G, Downie JA. **2011**. Natural variation in host-specific nodulation of pea is associated with a haplotype of the SYM37 LysM-type receptor-like kinase. *Molecular Plant-Microbe Interactions* **24**: 1396–1403.

Limpens E, Franken C, Smit P, Willemse J, Bisseling T, Geurts R. **2003**. LysM domain receptor kinases regulating rhizobial Nod factor-induced infection. *Science* **302**: 630–633.

Liu J, Yang S, Zheng Q, Zhu H. **2014**. Identification of a dominant gene in *Medicago*
truncatula that restricts nodulation by *Sinorhizobium meliloti* strain Rm41. *BMC Plant*
Biology **14**: 167.

Maillet F, Poinsot V, André O, Puech-Pagès V, Haouy A, Gueunier M, Cromer L,
Giraudet D, Formey D, Niebel A, *et al.* 2011. Fungal lipochitooligosaccharide
symbiotic signals in arbuscular mycorrhiza. *Nature* **469**: 58–63.

Maillet F, Fournier J, Mendis HC, Tadege M, Wen J, Ratet P, Mysore KS, Gough C,
Jones KM. **2020**. *Sinorhizobium meliloti* succinylated high-molecular-weight
succinoglycan and the *Medicago truncatula* LysM receptor-like kinase MtLYK10
participate independently in symbiotic infection. *The Plant Journal* **102**: 311–326.

Marczak M, Mazur A, Koper P, Żebracki K, Skorupska A. **2017**. Synthesis of rhizobial
exopolysaccharides and their importance for symbiosis with legume plants. *Genes* **8**:
360.

Masson-Boivin C, Giraud E, Perret X, Batut J. **2009**. Establishing nitrogen-fixing
symbiosis with legumes: how many rhizobium recipes? *Trends in Microbiology* **17**:
458–466.

Oldroyd GED, Murray JD, Poole PS, Downie JA. **2011**. The rules of engagement in the
legume-rhizobial symbiosis. *Annual Review of Genetics* **45**: 119–144.

Pietraszewska-Bogiel A, Lefebvre B, Koini MA, Klaus-Heisen D, Takken FLW,
Geurts R, Cullimore J V., Gadella TWJ. **2013**. Interaction of *Medicago truncatula*
Lysin motif receptor-like kinases, NFP and LYK3, produced in *Nicotiana benthamiana*
induces defence-like responses. *PLoS ONE* **8**: e650055.

Radutoiu S, Madsen LH, Madsen EB, Felle HH, Umehara Y, Grønlund M, Sato S,
Nakamura Y, Tabata S, Sandal N, *et al.* 2003. Plant recognition of symbiotic bacteria
requires two LysM receptor-like kinases. *Nature* **425**: 585–592.

Roche P, Debellé F, Maillet F, Lerouge P, Faucher C, Truchet G, Dénarié J, Promé
JC. **1991**. Molecular basis of symbiotic host specificity in *Rhizobium meliloti*: *nodH* and
nodPQ genes encode the sulfation of lipo-oligosaccharide signals. *Cell* **67**: 1131–1143.

Smit P, Limpens E, Geurts R, Fedorova E, Dolgikh E, Gough C, Bisseling T. **2007**.
Medicago LYK3, an entry receptor in rhizobial nodulation factor signaling. *Plant*
Physiology **145**: 183–191.

Sulima AS, Zhukov VA, Afonin AA, Zhernakov AI, Tikhonovich IA, Lutova LA. **2017**. Selection signatures in the first exon of paralogous receptor kinase genes from the *Sym2* region of the *Pisum sativum* L. genome. *Frontiers in Plant Science* **8**: 1957.

Sulima AS, Zhukov VA, Kulaeva OA, Vasileva EN, Borisov AY, Tikhonovich IA. **2019**. New sources of *Sym2^A* allele in the pea (*Pisum sativum* L.) carry the unique variant of candidate *LysM-RLK* gene *LykX*. *PeerJ* **7**: e8070.

Wang Q, Liu J, Zhu H. **2018**. Genetic and molecular mechanisms underlying symbiotic specificity in legume-rhizobium interactions. *Frontiers in Plant Science* **9**: 313.

Yang S, Tang F, Gao M, Krishnan HB, Zhu H. **2010**. *R* gene-controlled host specificity in the legume–rhizobia symbiosis. *Proceedings of the National Academy of Sciences of the United States of America* **107**: 18735-18740.

Solovev YV, Igolkina AA, Kuliaev PO, Sulima AS, Zhukov VA, Porozov YB, Pidko EA, Andronov EE. **2021**. Towards understanding Afghanistan pea symbiotic phenotype through the molecular modeling of the interaction between LYKX-SYM10 receptor heterodimer and Nod Factors. *Frontiers in Plant Science* **12**: 642591.

Younginger BS, Friesen ML. **2019**. Connecting signals and benefits through partner choice in plant–microbe interactions. *FEMS Microbiology Letters* **366**: fnz217.

Zarrabian M, Montiel J, Sandal N, Jin H, Lin Y-Y, Klingl V, Marín M, James E, Parniske M, Stougaard J, et al. **2021**. A promiscuity locus confers *Lotus burttii* nodulation with rhizobia from five different genera. *bioRxiv*. doi: 10.1101/2021.08.26.457880.

Zhou P, Silverstein KAT, Ramaraj T, Guhlin J, Denny R, Liu J, Farmer AD, Steele KP, Stupar RM, Miller JR, et al. **2017**. Exploring structural variation and gene family architecture with *de novo* assemblies of 15 Medicago genomes. *BMC Genomics* **18**: 261.

Zhukov V, Radutoiu S, Madsen LH, Rychagova T, Ovchinnikova E, Borisov A, Tikhonovich I, Stougaard J. **2008**. The pea *Sym37* receptor kinase gene controls infection-thread initiation and nodule development. *Molecular Plant-Microbe Interactions* **21**: 1600–1608.

Supporting Information

Additional supporting information may be found in the online version of this article.

Fig. S1 Amino acid alignment of LYK3-A17, LYK3-R108, LYK2bis, LYK2-A17 and LYK2-R108.

Table S1 *Sinorhizobium meliloti* strains used in this study

Table S2 Percentage identity/similarity of LYK2, LYK3 in A17 and LYK2, LYK2bis and LYK3 in R108.

Table S3 Presence of *LYK2bis* in *Medicago truncatula* genomes.

Table S4 Number of identical amino acids/total amino acids in each LysM between LYK2bis and LYK3-A17 and LYK2-A17.

 Figures and Tables

Fig. 1. *Medicago truncatula* R108 is able to nodulate with the *Sinorhizobium meliloti nodF/nodL* strain. The number of nodules in 20 plants/genotype/inoculation was analyzed at 21dpi. Statistical analyses were performed using Student's *t*-test (ns, not significant; ***, *P* < 0.001). White shading is for plants inoculated with the WT strain; green shading is for plants inoculated with the *nodF/nodL* strain

Fig. 2. *LYK2bis* is a newly-evolved gene in *Medicago truncatula* R108 and is responsible for nodulation with the *Sinorhizobium meliloti nodF/nodL* strain. (a) Schematic representation of the *LYK3-LYK2* region in *M. truncatula* A17 and R108. Coding sequences are shown in filled blocks; Non-coding sequences are shown in lines. kb = kilobases. (b) Schematic representation of LYK2, LYK2bis and LYK3 proteins of R108, showing the structure of the proteins and the *Tnt1* insertional sites in *lyk2-1R*, *lyk2bis-1R* and *lyk3-1R mutants*. The proteins are predicted to have three extracellular LysM domains (yellow/orange), a transmembrane domain (grey) and an intracellular kinase domain (blue/green). (c) *LYK2bis* is responsible for nodulation with the *nodF/nodL* mutant in R108. *lyk2-1R, lyk2bis-1R* and *lyk3-1R* mutants were phenotyped for nodulation with either *S. meliloti* WT or the *nodF/nodL* mutant strain. 20 plants/genotype/inoculation were analyzed at 21dpi. Statistical analyses were performed using One-way ANOVA ($P < 0.05$). Lowercase letters indicate significant difference. White shading is for plants inoculated with WT; Green shading is for plants inoculated with *nodF/nodL*.

Fig. 3. Complementation analysis confirms the role of *LYK2bis* in nodulation of *Medicago truncatula* R108 with the *Sinorhizobium meliloti nodF/nodL* strain. *lyk2bis-1R* roots were transformed with constructs of EV, LYK3-A17 and LYK2bis using *A. rhizogenes*. Transformed plants were analyzed at 4 wpi. Statistical analyses were performed using Student's *t*-test (*, $P < 0.05$). Number below indicates number of nodulated plants/total transformed plants.

Fig. 4. *LYK2bis* allows *Medicago truncatula* A17 to gain the ability to nodulate with the *Sinorhizobium meliloti nodF/nodL* strain. A17 roots were transformed with constructs of EV, LYK3-A17 and LYK2bis using *A. rhizogenes*. Transformed plants were analyzed at 4wpi. Statistical analyses were performed using Student's *t*-test (***, *P*-value< 0.001). Numbers below indicate number of nodulated plants/total transformed plants.

Fig. 5. Nodulation specificity of the *Medicago truncatula lyk2bis-1R* mutant with different *Sinorhizobium meliloti* mutant strains. The *lyk2bis-1R* mutant was tested for nodulation with *nodF/nodE* and *nodL* mutant strains. The number of nodules were counted at 21dpi. Statistical analyses were performed using Student's *t*-test (ns, not significant; ***, *P* < 0.001). White shading is for plants inoculated with WT; Light blue shading is for plants inoculated with *nodF/nodE;* Orange shading is for plants inoculated with *nodL.*

Fig. 6. *LYK2bis* does not play an essential role in mycorrhization. Seedlings were inoculated with *Rhizophagus irregularis* and examined at 3 and 5 wpi. 15 plants/genotype at each time point were analyzed. Statistical analyses were done using Students' t-test (*, P < 0.05). Dark blue shading is for the R108 genotype and dark green shading is for *lyk2bis-1R*.

(a)

LYK2bis
(0/36)

NFP
(0/27)

LYK2bis+NFP
(33/33)

(b)

NFP-GFP

LYK2bis-mCherry

Overlay

(c)

Fig. 7. LYK2bis physically interacts with NFP and this interaction can lead to cell-death at the plasma membrane of *Nicotiana benthamiana* leaves. (a) Co-expression of LYK2bis and NFP induces cell-death in *N. benthamiana* leaves at 3 dpi. Numbers in brackets indicate the number of leaves exhibiting cell death/total number of infiltrated leaves. Bars, 1 cm. (b) Co-localisation of NFP and LYK2bis at the plasma membrane of *N. benthamiana* leaves. Leaf discs were observed by confocal microscopy at 2 dpi. Bars, 25 μm. (c) LYK2bis physically interacts with NFP at the plasma membrane of *N. benthamiana* leaves by FRET FLIM technology. For each measurement, the difference to average GFP decay times (Δ) of the GFP-NFP in the absence of the LYK2bis-mCherry is given in ps; n: number of samples measured for each combination. Students' *t*-test was used for statistical analyses (***, *P* < 0.001).

Fig. 8. LYK2bis has an active kinase and can transphosphorylate NFP-KD *in vitro.* The kinase domains (KD) of LYK2bis and NFP were expressed and purified from *E. coli* as fusions with GST. GST-cleaved LYK2bis-KD was incubated individually or co-incubated with purified GST/NFP-KD (NFP), GST/LYR2-KD (LYR2), GST/LYR3-KD (LYR3), GST/LYR4-KD (LYR4), GST/LYK3-deadKD (LYK3dk), GST or Myelin Basic Protein (MyBP) in the presence of radioactive [γ-^{32}P] ATP. Assays were analyzed by SDS-PAGE, followed by coomassie staining (a) and phosphor imaging (b). Transphosphorylated proteins are marked by red arrowheads on the phosphor-image. The position of GST is marked on the coomassie gel (black arrowhead).

Fig 9. Model of NF recognition and nodulation in *Medicago truncatula* R108. Different rhizobia may secrete different proportions of WT-like and *nodFL*-like NFs. The nodulation of R108 with strains producing *nodFL*-like NFs is dependent on LYK2bis and possibly involves NFP while the WT-like NFs may involve other LYK(s) such as LYK3. Transphosphorylation (P) of NFP by the active LYK kinases may be important for signal transduction.