TFAP2 paralogs facilitate chromatin access for MITF at pigmentation genes but inhibit expression of cell-cell adhesion genes independently of MITF

Colin Kenny[1], Ramile Dilshat [2], Hannah Seberg[1+], Eric Van Otterloo[1*], Gregory Bonde[1], Annika Helverson[1], Christopher M. Franke[2], Eiríkur Steingrímsson[3], Robert A. Cornell[1, #]

1. Department of Anatomy and Cell Biology, College of Medicine, University of Iowa, Iowa City, Iowa

2. Department of Surgery, College of Medicine, University of Iowa, Iowa City, Iowa

3. Department of Biochemistry and Molecular Biology, BioMedical Center, Faculty of Medicine, University of Iceland, Sturlugata 8, 101 Reykjavik, Iceland

[+] Current address: University of Manchester

[*] Current address: Iowa Institute for Oral Health Research, Department of Anatomy and Cell Biology, Department of Periodontics, College of Dentistry, University of Iowa, Iowa City, Iowa

[#]Corresponding author:

cornellr@uw.edu

Conflict of interest statement: CK, RD, HS, EVO, GB, AH, CMF, ES and RAC declare no conflicts of interest.

Key words: MITF, TFAP2A, TFAP2C, TFAP2E, TFAP2B, melanocyte, melanoma, chromatin, nucleosome, pioneer factor, transcription, activator, repressor.

Abstract

In developing melanocytes and in melanoma cells, multiple paralogs of the Activating-enhancer-binding Protein 2 family of transcription factors (TFAP2) contribute to expression of genes encoding pigmentation regulators, but their interaction with Microphthalmia transcription factor (MITF), a master regulator of these cells, is unclear. Supporting the model that Tfap2 facilitates MITF's ability to activate expression of pigmentation genes, single-cell seq analysis of zebrafish embryos revealed that pigmentation genes are only expressed in the subset of *mitfa*-expressing cells that also express Tfap2 paralogs. To test this model in SK-MEL-28 melanoma cells we deleted the two *TFAP2* paralogs with highest expression, *TFAP2A* and *TFAP2C,* creating *TFAP2* knockout (*TFAP2*-KO) cells. We then assessed gene expression, chromatin accessibility, binding of TFAP2A and of MITF, and the chromatin marks H3K27Ac and H3K27Me3 which are characteristic of active enhancers and silenced chromatin, respectively. Integrated analyses of these datasets indicate TFAP2 paralogs directly activate enhancers near genes enriched for roles in pigmentation and proliferation, and directly repress enhancers near genes enriched for roles in cell adhesion. Consistently, compared to WT cells, *TFAP2*-KO cells proliferate less and adhere to one another more. TFAP2 paralogs and MITF co-operatively activate a subset of enhancers, with the latter necessary for MITF binding and chromatin accessibility. By contrast, TFAP2 paralogs and MITF do not appear to co-operatively inhibit enhancers. These studies reveal a mechanism by which TFAP2 profoundly influences the set of genes activated by MITF, and thereby the phenotype of pigment cells and melanoma cells.

Introduction

Gene expression in developing melanocytes and melanoma, a cancer derived from the melanocyte lineage, is regulated by transcription factors including Microphthalmia-associated transcription factor (MITF) and members of the SOXE, PAX and TFAP2 families [1-11]. MITF is required for differentiation of melanocytes during development, and its activity is regulated at both the transcriptional and post-translational levels [3]. In melanoma cells, high levels of MITF activity promote cell proliferation and pigmentation, while lower levels promote an invasive phenotype [12, 13]. Mass spectroscopy revealed that MITF interacts with components of both the PBAF chromatin remodeling complex, including BRG1 and CDH7, and the NURF

remodeling complex, including RBBP4 [14, 15]. Furthermore, chromatin

immunoprecipitation of BRG1 in cells depleted of *MITF* revealed that MITF recruits

BRG1 to the promoters of specific genes, including *TYR*, which encodes the rate-

limiting enzyme in melanin synthesis Tyrosinase [15]. Similar analysis suggested

that SOX10 also recruits BRG1 to chromatin, and at some loci it does so in co-

operation with MITF [15]. Conversely, there is evidence that PAX3 inhibits the

activity of MITF at the *DCT* promoter [16]. Furthermore, low MITF activity is

associated with an invasive phenotype, and deletion or knockdown of *MITF* results in

upregulation of genes that promote migration and invasion [17]. MITF CUT&RUN

peaks are found near some genes whose expression is upregulated in *MITF* mutant

cells, implying that MITF directly represses their expression [17]. This set of MITF

peaks is enriched for the binding site of FOXC1, a transcriptional repressor [18],

suggesting MITF has co-factors in its repressive function as well as its activating

one.

The activating enhancer-binding family of transcription factors, comprising five

members, TFAP2A-E, regulate development of many cell types and organs including

neural crest, placodes, epidermis, trophectoderm, heart, kidney, and brain [19-28]. In

several contexts, including melanocyte differentiation, TFAP2 paralogs function

redundantly [10, 29-31]. For instance, in zebrafish *tfap2a* loss-of-function mutant

embryos the number of melanocytes is lower than normal and pigmentation is

profoundly delayed relative to in wild-type embryos; this phenotype is exacerbated if

tfap2a mutant embryos are also depleted of *tfap2e* expression with antisense

morpholinos [10]. In zebrafish melanoma Tfap2a and Tfap2e also appear to act

redundantly to promote proliferation and, interestingly, to suppress cell adhesion and

cell migration [32]. Consistent with this, in the skin of mouse embryos with neural-

crest specific knockout of *Tfap2a* and *Tfap2b,* the two paralogs with highest

expression, fewer-than-normal cells express markers of melanocytes [33]. In

addition, evidence exists for sub-functionalization among Tfap2 paralogs [34]. Of

note, while *tfap2e* is the paralog with highest expression in zebrafish embryonic

melanophores, it's clear ortholog in mice, *Tfap2e,* is not expressed in embryonic

melanocytes [35]. This suggests that during evolution there has been some shuffling

of the function of individual TFAP2 paralogs among species.

TFAP2 paralogs and MITF appear to co-activate certain genes. For instance, in a

human melanoma cell line, the *in vitro* enhancer activity of an element within an *IRF4*

intron depended on the simultaneous binding of MITF and TFAP2 [36]. Further, in

zebrafish *tfap2a* and *mitfa* double mutant embryos there is a greater-than-additive

reduction in the number, and level of pigmentation, of melanocytes in comparison to

single mutants [33]. Supporting parallel function of Tfap2 paralogs and Mitfa, the

promoters of MITF target genes are enriched for TFAP2 consensus binding sites [15,

37], and ChIP-seq experiments in primary melanocytes indicate that TFAP2A and

MITF bind overlapping regions of chromatin near genes encoding regulators of

pigmentation [33]. Collectively, these observations indicate that TFAP2 paralogs co-

activate a subset of MITF target genes by binding at the same enhancers. Still

unclear, however, is whether TFAP2 paralogs and MITF act cooperatively or

independently at enhancers they co-regulate. It is also unclear whether they co-

repress enhancers.

TFAP2 paralogs may serve as pioneer factors for MITF, although not all evidence

supports this possibility. *Pioneer* or *initiating* TFs can bind and penetrate

nucleosome-bound DNA and recruit other TFs that lack this property called *settler*

TFs [reviewed in 38, 39, 40]. Evidence that TFAP2 paralogs are pioneer factors

includes, first, that TFAP2 binding site is over-represented within DNase1-protected

"footprints" in mouse embryonic stem cells induced to differentiate [41]. Second,

TFAP2A catalyzes assisted loading of androgen receptor (AR) in epididymis cells

[42] and estrogen receptor in MCF-7 cells [43]. Third, the TFAP2 binding site is

enriched for at the center of ATAC-seq peaks, implying it has a strong effect on

chromatin accessibility [44]. Fourth, ATAC-seq peaks in naïve-stated human ESC

showed reduced openness in TFAP2C-KO cells [45], and forcing expression of

TFAP2C in human ESC is sufficient to open chromatin at loci where it binds [46].

Finally, TFAP2A, TFAP2B and TFAP2C can bind nucleosomes [47]. Together these

findings support the possibility that TFAP2 displaces nucleosomes and thereby

facilitates chromatin binding by MITF. However, it is not clear that MITF needs a

pioneer factor to bind chromatin. In the *dynamic-assisted-loading model*, all classes

of TFs have short residency on chromatin [reviewed in 38]. In this model, *initiating*

TFs can recruit ATP-dependent chromatin remodelers (nBAF, SWI/SNF, INO80,

ISWI, NURD) and thereby make chromatin accessible to other TFs, i.e., the *assisted*

TFs [48]. Unlike the pioneer factor model, the assisted-loading-model predicts that

initiation factors are interchangeable, and that the recruitment of chromatin

remodeling complexes by initiation factors is dependent on the local chromatin

structure and accessibility [48]. As mentioned above, MITF binds various

components of the SWI/SNF complex [14, 49, 50] and the chromatin remodeler

CHD7 [15] and so meets the criteria for an initiating factor. If the dynamic-assisted-

loading model holds in this situation, MITF would have no need for a pioneer factor

like TFAP2 to assist its binding to chromatin.

To address these questions, we used single cell RNA-sequencing (scRNA-seq) to

investigate the temporal order of gene expression of *mitfa*, *tfap2* paralogs, and

pigmentation regulators in cells of the melanocyte lineage which were included

among GFP-expressing cells isolated from *Tg(mitfa-GFP)* zebrafish embryos at 28

hours post fertilization (hpf). Furthermore, to test how TFAP2 paralogs affect

enhancer activity and MITF binding, we deleted TFAP2 paralogs from SK-MEL-28

melanoma cells and assessed: nucleosome positioning, using the assay for

transposase-accessible chromatin with next generation sequencing (ATAC-seq);

enhancer activity, using cleavage under targets and release using nuclease

(CUT&RUN) with anti-H3K27Ac, anti-H3K4me3, and anti-H3K27me3; and binding of

MITF, using CUT&RUN with anti-MITF. We similarly assessed binding of TFAP2A in

cells harboring loss of function mutations in *MITF*. Our results support the notion that

TFAP2 factors behave like the canonical pioneer factor FOXA1: at many chromatin

elements bound by TFAP2A, loss of TFAP2 led to loss of enhancer activity, and in a

large subset, it also resulted in chromatin becoming condensed. At both subsets of

TFAP2-activated enhancers, MITF binding was TFAP2-dependent. In addition, we

find evidence that TFAP2 paralogs can also inhibit enhancers, and at a subset of

those, they exclude binding of MITF. Finally, the analyses suggest that TFAP2

directly inhibits many of the same genes that MITF inhibits, but we do not find

evidence that TFAP2 and MITF co-repress the same enhancers. Together these

findings illuminate the mechanisms by which TFAP2 and MITF coordinately regulate

differentiation of melanocytes and the phenotype of melanoma cells.

Results

Expression of *tfap2* paralogs in the melanocyte lineage precedes activation of Mitfa transcriptional programs *in-vivo*.

If Tfap2 paralogs cooperate with MITF to promote expression of melanocyte differentiation genes *in-vivo*, then high level expression of known Mitfa-target genes, including *dct, pmel, mreg (zgc:91968)* and *trpm1b*, should only occur in cells that express both *mitfa* and *tfap2* paralogs. To test this prediction, we performed scRNA-seq on GFP[+] cells sorted from *Tg(mitfa:GFP)* transgenic zebrafish embryos at 28 hours post fertilization (hpf) using the 10x Chromium system. We sequenced 11,217 cellular transcriptomes and visualized the data in two-dimensional space using Seurat [51] and uniform and manifold approximation and projection (UMAP) [52] (**Fig. 1A**). We assigned cell-type annotations using previously identified marker genes [53-56] and information from the Zebrafish Information Network (ZFIN) [57]. The remaining 28 clusters comprised 11 cell types (**Fig. 1A**); neural crest and basal cell lineages were the most strongly represented (**Supplemental Fig. S1A-C**). Interestingly, only a minority of the clusters expressed *mitfa.* Sorted cells that express GFP but not *mitfa* presumably reflect either non-specific activity of the *mitfa* promoter used in the transgene or earlier transient expression of *mitfa* revealed by the long half-life of GFP; importantly, for the upcoming analysis, such cells did not express pigmentation genes. We focused on *sox10*-positive clusters 6-12 as these comprised all stages of the melanocyte lineage, including all *mitfa*-expressing cells (**Supplemental Fig. S1D**). Re-clustering of *sox10*-positive cells identified ten clusters (**Fig. 1B**). Two of these were neural crest (NC) cell populations which shared expression of bona fide NC markers (*foxd3* and *sox10*), but were distinguished by expression in one cluster of members of the Notch signaling pathway (*her6, her4* and *her12*); this cluster may represent migrating cardiac or cranial NC populations [58] (**Fig. 1C**, **Supplemental Fig. S2A, S2D**). In addition, two clusters expressed markers of a tripotent precursor of melanoblasts, iridoblasts, and xanthoblasts (*cdkn1ca, slc15a2, ino80e, id3, mycn, tfec*), called MIX cells [59-62] (**Supplemental Fig. S2B**). Interestingly, while both MIX clusters express *tfap2a*, the clusters differed by one and not the other expressing *tfap2e* and *tfap2b* - indeed these genes were ranked first and third, respectively, among those differentially expressed between the two MIX clusters (**Fig. 1G; Supplemental Fig. S3A;**

Supplemental Data Table 1). Therefore, we will refer to the two MIX clusters as

tfap2-low and *tfap2*-high (**Fig. 1B**). An additional cluster expressed high levels of

melanoblast/ xanthoblast markers (*mitfa, impdh1b, gch2, id3*) suggesting it

corresponds to the previously described MX cluster [59-62]. Another cluster was

similar to the MX cluster but also expressed markers of differentiated melanophores

(*dct, pmela, tyrp1b, mreg, tyr*) (**Fig. 1F-G; Supplemental Fig. S4A-B**). Other cluster

pairs corresponding to xanthoblasts and xanthophores, and to iridoblasts and

iridophores, were identified (**Fig. 1B, Supplemental Fig. S2C-D**).

Pseudotime analysis supports a lineage trajectory for the melanophore lineage

leading from NC, to *tfap2*-low MIX cells, to *tfap2*-high MIX cells, to MX, to

melanophores (**Fig. 1B**). Within this trajectory, despite expression of *mitfa* in both

tfap2-low and *tfap2*-high MIX clusters, expression of select pigmentation genes (*dct,*

pmel, mreg and *trpm1b*) is first detected in *tfap2*-high MIX cells, and becomes higher

in MX cells concomitant with higher level expression of *mitfa* and of *tfap2e* (**Fig. 1C-**

G; Supplemental Fig. S4A-B). These data support a cooperative role for Mitfa and

Tfap2 paralogs in activation of pigmentation genes.

Tfap2a and Tfap2e redundantly promote the differentiation of zebrafish

embryonic melanocytes

The scRNA-seq analysis revealed expression of four *tfap2* paralogs at some point

within the embryonic melanophore lineage, i.e., in NC cells (*tfap2a*), in *tfap2*-low MIX

cells (*tfap2a*), *tfap2*-high MIX cells (*tfap2a, tfap2b, tfap2c*, and *tfap2e*), and in MX

(*tfap2a, tfap2c* and *tfap2e*) (**Fig. 1G; Supplemental Fig. S3A**); expression of all

tfap2 paralogs is reduced upon terminal differentiation of melanophores (cluster 10).

An overt phenotype of reduced melanocyte numbers and delayed melanization is

evident in *tfap2a* mutants [23, 63, 64] and morphants [65], but not in *tfap2b*

morphants [56], *tfap2c* mutants [29, 66], or *tfap2e* morphants [10]. However,

depletion of *tfap2e* from *tfap2a* mutants yields a more severe phenotype of reduced

melanocyte numbers and delayed melanization in comparison to control-MO-injected

tfap2a mutants [10]. Here we used zinc finger nucleases to generate a 157bp

deletion of exon 2 in *tfap2e*. Similar to the findings with morpholinos, homozygous

mutants were phenotypically normal (**Supplemental Fig. S5A-C**) while *tfap2a/tfap2e*

double mutants had fewer embryonic melanophores than *tfap2a* single mutants, and

pigmentation was further delayed (**Fig. 1H-L**). Nonetheless, by 48 hpf the intensity of

pigmentation within individual melanophores was not overtly different between wild-

type and *tfap2a/tfap2e* double mutants (**Supplemental Fig. S5G-J**).

Importantly, the phenotypes in single or double mutants thus far examined may be

suppressed by compensatory upregulation of other Tfap2 paralogs. Indeed, we

detected upregulated expression of *tfap2c* in *tfap2a* mutants and in *tfap2a/tfap2e*

double mutants, presumably driven by non-sense mediated decay (**Supplemental**

Fig. S6). In conclusion, the *in-vivo* consequence of loss of Tfap2 function upon

melanophore differentiation is difficult to assess because of the potential for

redundant function among Tfap2 paralogs and because of pleiotropy (see

Discussion), nonetheless mutant embryos with lower-than-normal expression of

tfap2 paralogs exhibit fewer melanophores than wild-types, and these melanophores

are delayed in pigmentation.

TFAP2A binds open and closed chromatin

To test how TFAP2 paralogs interact with MITF we switched to a cell line model. We

chose the SK-MEL-28 melanoma cell line because we have deleted all alleles of

MITF from this cell line and have evaluated MITF binding genome-wide using

CUT&RUN [17]. Here we carried out (1) CUT&RUN using antibodies to TFAP2A

(i.e., TFAP2A peaks), (2) CUT&RUN using antibodies to chromatin marks indicative

of active regulatory elements (H3K27Ac and H3K4Me3) [67, 68] and indicative of

inactive chromatin (H3K27Me3) [69], and (3) ATAC-seq to distinguish between open

and closed chromatin [70]. We used IgG as a background control and the MACS2

software to call peaks in each CUT&RUN dataset (**Supplemental Fig. S7A-B**).

Based on proximity to transcriptional start sites (TSS), about one-third of TFAP2A

peaks appeared proximal to promoters (within 3kb of a TSS). As expected, these

elements had strong H3K4Me3 signal (**Supplemental Fig. S7B**). At promoter-

proximal TFAP2A peaks, the H3K27Ac signal in WT cells was relatively consistent,

whereas at promoter-distal TFAP2A peaks (greater than 3 kb but within 100 kb of a

TSS) the H3K27Ac signal ranged from high to background level (**Supplemental Fig.**

S7B-C). About two-thirds of TFAP2A peaks overlapped ATAC-seq peaks, indicating

that they were in open chromatin (**Supplemental Fig. S7D-E**). Of note, the read

depth (height) of a peak approximates the number of chromosome molecules where

TFAP2A binds. The average read depth of the TFAP2A peaks in closed chromatin

was only about 50% of that in open chromatin but was nonetheless 80-fold higher

than the IgG background read depth (**Supplemental Fig. S7D-E, Supplemental**

Fig. S8A-B for example loci). Importantly, the TFAP2 binding site was strongly

enriched for in both TFAP2A-bound elements where the local ATAC-seq signal was

called as a peak and in counterparts where it was not ($p < 1 \times 10^{-1785}$ and $p < 1 \times 10^{-4375}$

, respectively), supporting the idea that TFAP2A binds DNA directly even when

the DNA is occupied by nucleosomes (**Supplemental Fig. S8C-D**). These results

indicate that TFAP2A binds at both open and closed chromatin, consistent with it

being a pioneer factor, and at enhancers and promoters with a range of activity

levels.

At a subset of TFAP2-activated enhancers, TFAP2 is necessary for chromatin

accessibility

Having shown that TFAP2A can bind closed chromatin, we next asked whether

TFAP2 factors play a role in opening chromatin [39]. We used CRISPR/Cas9

methods to introduce frame-shift mutations into *TFAP2A* and *TFAP2C*, the *TFAP2*

genes with high expression in SK-MEL-28 cells (**Supplemental Fig. S9A-D**). We

then carried out RNA-seq, ATAC-seq, and CUT&RUN with antibodies to H3K27Ac,

H3K4me3, and H3K27me3, in two independent knockout clones (hereafter, *TFAP2*-

KO cells). Western blot analysis showed an absence of immunoreactivity for both

proteins (**Supplemental Fig. 9E**). Control clones (hereafter WT cells) were derived

from the parental SK-MEL-28 line transiently transfected with Cas9 but not with

guide RNAs. RNA-seq revealed that expression of 532 genes was downregulated,

and expression of 609 genes was upregulated, in *TFAP2*-KO cells (i.e., in both

clones) versus in WT cells. We will refer to these sets as TFAP2-activated genes

and TFAP2-inhibited genes, respectively.

We defined TFAP2-activated enhancers as TFAP2A peaks a) that overlap H3K27Ac

and ATAC-seq peaks in WT cells (21,745/ 36,948, or 59% of all TFAP2A peaks), b)

that are greater than 3 kb from a transcription start site (11,005/ 21,745, or 51% of

TFAP2A peaks at H3K27Ac/ATAC peaks), and c) where the H3K27Ac signal was

significantly lower in *TFAP2*-KO cells relative to in WT cells (adj p < 0.05, log2FC <-

1; 3,858/11,005 or 35% of TFAP2A-bound enhancers). Interestingly, at about half

the TFAP2-activated enhancers (2002/3858), the ATAC-seq signal was significantly

lower in *TFAP2*-KO versus in WT cells (adj. p < 0.05, log2FC <-1) (**Fig. 3A, E-E'**).

The open status of these loci depends on TFAP2 paralogs, and we therefore infer

that TFAP2 paralogs directly or indirectly recruit the nucleosome-displacing

machinery. We further infer that TFAP2 paralogs bound these loci as closed

chromatin, but it is also possible they bound to another pioneer factor present at

these loci. In either case, it participates in a pioneering function, and we refer to this

subset of enhancers as TFAP2-pioneered-and-activated enhancers (**Fig. 2 Box 1A**).

We refer to the subset where ATAC-seq signal is unchanged between *TFAP2*-KO

and WT cells as non-pioneered TFAP2-activated enhancers **(Fig. 2 Box 1B; Fig.**

3B, F-F').

Both TFAP2-pioneered-and-activated enhancers and non-pioneered TFAP2A-

activated enhancers were associated with TFAP2-activated genes. Interestingly, the

association was stronger for the former subset (**Fig. 3 I, J**; **Table 1,** compare rows 4

and 5). Moreover, at both subsets, the H3K4me3 signal, which is associated with

enhancer activity [68], was reduced in *TFAP2*-KO cells relative to WT cells (**Fig.**

3E'', F''). While both subsets were strongly enriched for the TFAP2 binding site and

certain other binding sites (e.g., RUNX), the subset pioneered by TFAP2 was more

strongly enriched for the SOXE and MITF binding sites, while the non-pioneered

subset was more strongly enriched for the FRA1, TEAD and the ZFX binding sites

(**Fig. 3M, N**). Of note, FRA1 is a pioneer factor [71] which could explain why these

elements do not depend on TFAP2 to be free of nucleosomes.

Table 1: TFAP2-regulated enhancers and their association with TFAP2-regulated genes (Fisher's Exact Test, hypergeometric analysis):

Effect of TFAP2 on:				Association with **TFAP2**-activated genes	Association with **TFAP2**-inhibited genes
Enhancer	H3K27Ac	Chromatin	N (# of elements)	p-value |Log2FC| >1	p-value |Log2FC| >1
All TFAP2-activated	Increases	Any effect or none	3,838	1.12×10^{-24}	ns
TFAP2-pioneered-and-activated	Increases	Opens	2,002	3.7×10^{-27}	ns
Non-pioneered TFAP2-activated	Increases	None	1,836	2.1×10^{-12}	ns
All TFAP2-inhibited	Decreases	Any effect or none	1,304	ns	4.8×10^{-06}
TFAP2-pioneered-and-inhibited	Decreases	Condenses	864	ns	9.44×10^{-05}
Non-pioneered TFAP2-Inhibited	Decreases	None	440	ns	ns
All TFAP2A peaks	Any effect or none	Any effect or none	36,948	2.4×10^{-09}	ns
TFAP2A peaks	None	None	23,735	ns	ns

Column 1: groups of enhancers directly bound by TFAP2A, determined by CUT&RUN. Pioneered, refers to the set of enhancers where the chromatin is opened or condensed by TFAP2. Non-pioneered, refers to the set of enhancers where only the H3K27Ac signal is activated or repressed by TFAP2 (i.e. where the ATAC-Seq signal is independent of TFAP2 binding). Columns 2-3: TFAP2-dependent H3K27Ac signal (increased or decreased) and/or TFAP2-dependent nucleosome depleted regions (ATAC-Seq; opened or condensed) at TFAP2 bound enhancers. Column 4: the number of target elements regulated by TFAP2. Columns 5-6: the effect of TFAP2 on gene expression (two independent clones, 4 replicates each) and WT (4 replicates). Differential gene expression (adjusted p-value < 0.05) in *TFAP2*-KO cells and WT cells was determined by RNA-Seq. The p-value from the Fisher's Exact Test in which the null hypothesis is that no association exists. DEG: differentially expressed genes. Attention should be focused on columns 5-6 which indicate the strength of association between the regulation status of a gene and the TFAP2-regulated enhancer of interest; significant associations are highlighted yellow. ns; not statically significant.

TFAP2A inhibits enhancers by blocking the opening of chromatin

Because of evidence that TFAP2A directly represses gene expression [72-74] we next sought to identify enhancers directly inhibited by TFAP2 paralogs. To this end we filtered promoter-distal TFAP2A peaks for those where the local H3K27Ac signal was higher in *TFAP2*-KO cells than in WT cells (adj. p <0.05, log2FC>1). Analogously to TFAP2-activated enhancers, TFAP2-inhibited enhancers were split between a subset where the ATAC-seq signal was higher in *TFAP2*-KO cells than in WT cells (as illustrated in **Fig. 2 Box 2A; Fig. 3C, G-G'**), implying TFAP2 paralogs maintain condensed chromatin at these sites, and a subset where it was unchanged (**Fig. 2 Box 2B; Fig. 3D, H-H'**). The first but not the second subset was significantly associated with TFAP2-inhibited genes (**Fig. 3K**, **Table 1**), and the average H3K4me3 signal at the first but not the second subset was higher in *TFAP2*-KO cells than in WT cells (**Fig. 3G'', H''**). We define these enhancers as TFAP2-pioneered-and-inhibited. The binding site for TFAP2 was strongly enriched for in these enhancers, as were those for ETS1 and CTCF (**Fig. 3O**), both transcriptional repressors [75, 76]. By contrast, the subset of candidate TFAP2-inhibited enhancers where the ATAC- and average H3K4me3 signals were unchanged between TFAP2-KO and WT cells was not associated with TFAP2-inhibited genes. We infer these elements are not, in fact, TFAP2-inhibited enhancers. In conclusion, at TFAP2-inhibited enhancers TFAP2 recruits and/or retains a machinery that condenses chromatin and inhibits enhancer activity; the canonical pioneer factor FOXA1 also has this potential [77, 78]. Alternatively, TFAP2 itself may instead play a distinct role in keeping enhancers closed rather than actively closing them from a presumed prior open state.

We analyzed whether TFAP2 paralogs directly activate or directly inhibit promoters, and how they do so. Although TFAP2A peaks are frequently found at promoters (8277 genes have a TFAP2A peak within 3 kb of the TSS), it was uncommon for the underlying H3K27Ac and H3K4Me3 signals to be elevated (or reduced) in *TFAP2*-KO cells relative to WT cells. Based on this quality, we found 119 candidates for directly TFAP2-activated promoters, and 31 candidates for directly TFAP2-inhibited promoters. Nonetheless, similar to the trends for TFAP2 regulated enhancers, the pioneered subset of directly TFAP2-activated promoters was more strongly

associated with TFAP2-activated genes than the non-pioneered subset, and only the

pioneered subset of TFAP2-inhibited promoters was associated with TFAP2-

inhibited genes (**Supplemental Fig. S11A-L; additional examples in**

Supplemental Fig. S12A-B).

Interestingly, at the majority of TFAP2A peaks, whether at open or closed chromatin,

neither the ATAC-seq nor H3K27Ac signals were TFAP2-dependent. This large

subset of quiescent TFAP2A peaks was associated with neither TFAP2-activated nor

TFAP2-inhibited genes (**Table 1, row 10**). In contrast to the quiescent subset of

peaks, the set of all TFAP2A peaks was modestly associated with TFAP2-activated

genes (**Table 1, row 9**). Nonetheless, we infer that the common practice of using

ChIP-seq or CUT&RUN-seq data to identify genes directly regulated by a

transcription factor is subject to false positives.

At a subset of MITF/TFAP2 co-bound peaks, TFAP2 paralogs facilitate MITF

binding

A prediction of the TFAP2-as-pioneer-factor model is that binding of transcription

factors, like MITF, will depend on TFAP2. Among 36,621 MITF peaks in WT SK-

MEL-28 cells that we previously identified by CUT&RUN [17], we found that 15,752

(43%) overlap a TFAP2A peak. Of these, 9,413 (60%) were within open and active

(i.e., ATAC+ and H3K27Ac+) chromatin (**Supplemental Fig. S12A**). To assess MITF

binding in the absence of TFAP2, we carried out anti-MITF CUT&RUN in *TFAP2*-KO

cell lines. Of note, as *MITF* RNA levels in *TFAP2*-KO cells are only about 40-50% of

those in WT cells, an across-the-board decrease in the average height (read depth)

of MITF peaks was possible. Instead, we observed that the average height of MITF

peaks not overlapping TFAP2A peaks was equivalent in *TFAP2*-KO cells and in WT

(**Supplemental Fig. S12B** and **Supplemental Fig. S12D** for example at *MLANA*).

By contrast, the height of about 35% (5,443 /15,752) of MITF peaks overlapping

TFAP2A peaks was significantly lower in *TFAP2*-KO cells than in WT cells (adj. p

<0.05, log2FC < -1). (**Fig. 4A-D; Supplemental Fig. S12C-D** and all replicates

shown in **Supplemental Fig. S13**). We refer to these as directly TFAP2-activated

MITF peaks.

We reasoned that TFAP2 paralogs could facilitate MITF binding by displacing

nucleosomes (i.e., in pioneer factor mode) or alternatively by elevating MITF's affinity

for open DNA. Consistent with both models, we observed that directly TFAP2-

activated MITF peaks fall in three subsets with respect to the TFAP2-dependence of

the underlying ATAC-seq signal. Thus, at about 57% (3,083/5,443) of directly

TFAP2-activated MITF peaks the ATAC-seq signal was significantly lower (**Fig. 4B-**

C, 4F-F'), at 37% (2,022/5,443) it was unchanged (**Fig. 4D, 4H-H'**), and at 6% it was

higher (**Supplemental Fig. S14A-A'**) in *TFAP2*-KO cells compared to WT cells. The

first two subsets were strongly associated with TFAP2-activated genes

(Hypergeometric test; p-value $= 8.4$ x 10^{-26} and p-value $= 1.07$ x 10^{-13} respectively)

and with MITF-activated genes (Hypergeometric test; p-value $= 1.16$ x 10^{-21} and p-

value $= 4.3$ x 10^{-11} respectively) (**Fig. 4F', H', Table 2**)**.** We infer that at the first

subset of TFAP2-dependent MITF peaks, TFAP2 is a pioneer factor (or participates

in pioneering function), facilitating access to chromatin for MITF and other

transcription factors (illustrated in **Fig. 4F''**). Supporting this prediction, the

transcription factor binding sites for MITF, SOX10, RUNX and FRA1 were strongly

enriched at such elements (**Fig. 4F'''**). At the second subset, TFAP2 is a

transcriptional activator that recruits MITF, also functioning as a transcriptional

activator; we presume another protein serves as a pioneer factor at this subset

(illustrated in **Fig. 4H''**). Consistent with this notion, the binding site for JUN, a widely

deployed pioneer factor [79], site is strongly enriched in these elements (**Fig. 4H'''**).

Examples are shown of TFAP2-activated MITF peaks near *FRMD4B, CYP7B1,*

TRPM1, SOX9, EDNRB, MREG, GPR143, SNAI2, MEF2C, MYO5A, PAX3, EN1

and *FOXI3* genes (**Fig. 4B-D and Supplemental Fig. S12D**). At the third subset of

TFAP2-dependent MITF peaks, where ATAC-seq signal was higher in *TFAP2*-KO

cells than in WT cells (**Supplemental Fig. S14A-A'**), TFAP2 may serve as a pioneer

factor for MITF in MITF's proposed role as transcriptional repressor [17] (illustrated in

Supplemental Fig. S14A''). However, this category of element was not enriched

near genes inhibited by either TFAP2 or MITF (Hypergeometric test; p-value $= 6.02$ x

10^{-02} and p-value $= 9.12$ x 10^{-02} respectively). In summary, these results are

consistent with TFAP2 facilitating access for MITF as a transcriptional activator to

enhancers in both pioneer-factor and non-pioneer factor modes.

Table 2: TFAP2-dependent MITF peaks and gene expression (Fisher's Exact

Test; hypergeometric analysis):

Effect of TFAP2 on			Association with activated genes	Association with inhibited genes
MITF binding	Chromatin	N (# of elements)	p-value |Log2FC| >1	p-value |Log2FC| >1
All TFAP2 - activated	Any effect or none	5,443	8.4 x 10^{-26}	ns
Pioneered	Opens	3,083	1.6 x 10^{-26}	ns
Non-pioneered	None	2,358	1.07 x 10^{-13}	ns
Mutually - activated	Opens	717	3.47 x 10^{-05}	ns
All TFAP2 - inhibited	Any effect or none	1,605	ns	5.0 x 10^{-03}
Pioneered	Condenses	924	ns	3.5 x 10^{-04}
Non-pioneered	None	681	ns	ns
None	Any effect or none	10,418	ns	ns

Column 1: groups of directly TFAP2-regulated MITF peaks (activated or inhibited),
determined by anti-MITF CUT&RUN in *TFAP2*-KO and WT SK-MEL-28 cell lines.
Column 2: TFAP2-dependent MITF peaks at TFAP2-regulated nucleosome depleted
regions (ATAC-Seq; opened or condensed by TFAP2). Column 3: the number of
directly TFAP2-regulated MITF peaks. Column 4-7: The effect of TFAP2-regulated
MITF peaks on gene expression (two independent clones, 4 replicates each) and
WT (4 replicates). Differentially expressed genes (adjusted p-value <0.05) between
TFAP2-KO cells and WT cells was determined by RNA-Seq. The p-value from the
Fisher's Exact Test in which the null hypothesis is that no association exists. DEG:
differentially expressed genes. Attention should be focused on columns 4-5, which
indicate the strength of association between the regulation status of a gene and the
TFAP2-regulated MITF peaks of interest; significant associations are highlighted
yellow. ns; not statically significant.

To test whether depended on MITF, we carried out anti-TFAP2A CUT&RUN in

MITF-KO cells. *TFAP2A* mRNA and protein levels were equivalent in *MITF*-KO and

WT cells (**Supplemental Fig. S9E**), and the average TFAP2A peak height was

globally equivalent as determined by CUT&RUN. At 13% (717/ 5334) of TFAP2-

dependent MITF peaks, the TFAP2A peak was, reciprocally, significantly reduced in

MITF-KO cells (**Fig. 4A and 4C**). At such loci, the average ATAC-seq signal was

reduced in *TFAP2*-KO cells as compared to WT cells (**Fig. 4C, 4G-G'**). We termed

these peaks mutually-dependent (i.e. mutually-activated peaks) (**illustrated in Fig.**

4G'')**.** Interestingly, mutually-dependent MITF/ TFAP2 peaks were enriched in

binding motifs for TFAP2, MITF, BRM2 and TEAD4 but, unlike the other subsets of TFAP2-activated MITF peaks, not for SOXE (**Fig. 4H'''**). SOX10 co-binds many loci together with MITF [15] and if SOX10 is absent from mutually-dependent peaks this may explain the dependence of TFAP2A binding on MITF at these sites. At ~40% (288/ 717) of the mutually dependent peaks, and in the gene body surrounding these peaks, the repressive histone mark H3K27Me3, known to be applied by the polycomb complex [69, 80], was significantly higher in *MITF*-KO cells relative to WT cells but, unexpectedly, not in *TFAP2*-KO cells relative to WT cells, even though MITF binding was lower in *TFAP2*-KO cells (illustrated in **Fig 4G''**, **Supplemental Fig. 12E-G**).

In summary, the binding of MITF depends on TFAP2 at about one third of MITF peaks that overlap TFAP2A peaks. Such TFAP2-activated MITF binding occurs both at loci where nucleosome packing depends on TFAP2 (pioneer factor mode) and where it does not (non-pioneer factor mode). At a subset of the former, but none of the latter, TFAP2A binding is, reciprocally, MITF-dependent.

At a subset of MITF/TFAP2A co-bound peaks, TFAP2 paralogs inhibit chromatin access for MITF

In Figure 3 we established that at some TFAP2A peaks, TFAP2 paralogs maintain closed chromatin, and presumably inhibit binding of transcription factors. Consistent with this prediction, among MITF peaks overlapping TFAP2A peaks, 10% (1,605) of the MITF peaks were higher in *TFAP2*-KO cells than in WT cells (third cluster **Fig. 4A, E; Supplemental Fig. S13**), implying TFAP2 paralogs inhibit binding of MITF at this subset. At most peaks within this subset (58%, 924/1,605), the ATAC-seq signal was also significantly higher in *TFAP2*-KO cells versus WT cells (violin plots, **Fig. 4I, I',** illustrated in **Fig. 4I''**), implying TFAP2 maintains condensed chromatin at this subset. DNA elements underlying this subset were enriched for transcription factor binding sites including those for SP1, NFY, JUN and TFE3 (**Fig. 4I'''**). Moreover, these elements were modestly associated with TFAP2-inhibited genes (**Table 2;** row 8)**.**

Of note, at the majority of MITF peaks that overlap TFAP2A peaks (65%, 10,418/

15,752), the height of the MITF peak was equivalent in *TFAP2*-KO and WT cells

(TFAP2-independent MITF peaks) (**Supplemental Fig. S13, Supplemental Fig.**

S15A;), implying TFAP2 paralogs neither recruit nor repel MITF at these sites.

Interestingly, such TFAP2-independent MITF peaks were not strongly enriched for

the TFAP2 binding site (**Supplemental Fig. S15A**), implying that TFAP2 is attracted

to many of these sites via other proteins rather than binding directly to the DNA.

Such indirect binding may be less avid, as the average height of TFAP2-independent

MITF peaks was smaller than that of TFAP2-dependent MITF peaks (**Supplemental**

Fig. S13, compare MITF signal in first and fourth cluster WT cells). As expected,

TFAP2-independent MITF peaks were associated neither with TFAP2-activated nor

TFAP2-inhibited genes (**Table 2**; row 10).

TFAP2 and MITF co-regulate genes in the melanocyte gene regulatory network

The delayed pigmentation in zebrafish *tfap2a/tfap2e* double mutants, and the

reduced expression of many pigmentation genes in *TFAP2*-KO cells, is consistent

with two scenarios which are not exclusive of one another. In the first, TFAP2

paralogs directly activate *MITF* expression and thereby indirectly activate expression

of pigmentation genes. In the second, TFAP2 paralogs directly activate expression of

pigmentation genes. Supporting the first scenario, there is a TFAP2-pioneered-and-

activated enhancer in intron 2 of the *MITF* gene (**Supplemental Fig. S16**), and *MITF*

mRNA levels are about 40-50% lower in *TFAP2*-KO cells than in WT cells.

However, arguing against a strong role for this mechanism, many of the genes

whose expression was most strongly reduced in *MITF*-KO cells compared to WT

cells were completely TFAP2-independent, or were TFAP2-inhibited (**Supplemental**

Table 4). To assess the second scenario, we identified the set of genes activated

directly by MITF, defined as MITF-activated genes associated with an MITF peak,

and the set of genes directly activated by TFAP2, defined as TFAP2-activated genes

associated with an TFAP2-activated enhancer (i.e., of pioneered or non-pioneered

variety). Supporting the second mechanism, genes activated directly both by TFAP2

and by MITF were enriched for GO terms related to pigmentation and pigment cell

differentiation (**Fig. 5A-B**) [81]. Of note, genes related to pigmentation were also

among those apparently directly regulated solely by MITF or TFAP2 paralogs (**Fig.**

5C). We took a similar approach to identify genes directly inhibited by TFAP2 and/or

by MITF (**Fig. 5D**). Genes directly inhibited by both were strongly enriched for GO

terms related to cell motility and cell migration (**Fig. 5D-E**)**.** Additional categories of

genes strongly activated or inhibited by TFAP2 included genes associated with cell

proliferation and cell differentiation (**Fig. 6A; Supplemental Fig. S17A-B**) and with

cell-cell adhesion, respectively (**Fig. 6B; Supplemental Fig. S17C)**.

We next examined the association of TFAP2-activated and TFAP2-inhibited genes

with gene expression profiles from melanoma tumors and cell lines with distinct

phenotypes [82-88]. Enrichment analysis showed that melanoma profiles previously

found to be associated with high levels of MITF activity [89] were enriched for genes

directly activated by TFAP2, including the subset associated with TFAP2-dependent

MITF peaks (**Fig. 5F**). Moreover, melanoma profiles associated with low levels of

MITF activity were enriched for genes directly inhibited by TFAP2 (**Fig. 5F**). These

findings support other observations that the level of TFAP2 expression has a

profound effect on the melanoma phenotype [32, 90].

Finally, we performed cell proliferation, cell adhesion and cell migration assays on

wild-type and TFAP2-depleted SK-MEL-28 cells. Consistent with effects on gene

expression, *TFAP2*-KO cells showed reduced proliferation over 24, 48, 72 and 96

hours compared to WT cells (**Fig. 6E**), and in cluster formation assays, *TFAP2*-KO

cells showed increased cell-cell adhesion compared to WT cells (**Fig. 6E**). However,

in contrast to the effects on gene expression, while WT cells migrated to fill in a

scratch within 24 hours, neither *TFAP2*-KO clone filled it within that time

(**Supplemental Fig. S18A**). This finding also contrasts with the observation that the

expression of *tfap2e* correlates negatively with the migratory capacity of zebrafish

models of melanoma [32], but it is consistent with the accumulation of melanocytes

in the dorsum of zebrafish *tfap2a* knockout embryos [23, 63, 64] and *tfap2a/ tfap2e*

double mutant embryos. It is noteworthy that in *MITF*-KO cells, similar to in *TFAP2-*

KO cells, cell migration genes are upregulated, but cell migration *in-vitro* is inhibited

[17]. In summary, reduced expression of pigmentation, differentiation and cell growth

genes and elevated expression of cell adhesion and cell invasion genes in *TFAP2-*

KO cells as compared to WT cells is largely explained by the direct effects of TFAP2

paralogs on these genes.

Discussion

Here, single-cell seq analysis of zebrafish embryos indicated that expression of MITF-target genes encoding regulators of pigmentation occurs only in the subset of *mitfa*-expressing cells which also express *tfap2* paralogs. In embryos with loss-of-function mutations in *tfap2a* and *tfap2e*, the two *tfap2* paralogs with highest expression in melanophore precursors, there are fewer embryonic melanophores than in wild-type embryo's and pigmentation is delayed. The phenotype was similar, but less penetrant than the phenotype in *tfap2a* mutants injected with *tfap2e* MO [10]. We speculate that homeostatic upregulation of *tfap2c* in *tfap2a/tfap2e* double mutants, possibly mediated by transcriptional adaptation [91], suppresses the phenotype. Simultaneous deletion of *tfap2a* and *tfap2c* eliminates neural crest precluding the possibility of examining melanocyte development in such embryos [23, 29, 64]. It has been thus far impossible to evaluate the phenotypic consequence of loss, rather than just reduction, of Tfap2 activity in the melanophore lineage in zebrafish. Although RNA-less loss-of-function mutants of *tfap2a* and *tfap2e* should prevent transcriptional adaptation [91, 92], it is possible the homeostatic upregulation of *tfap2c* in the existing mutants documented here occurs by another mechanism. Of note, in mice with simultaneous tissue specific deletion of Tfap2a and Tfap2b in the neural crest lineage the number of embryonic melanocytes is reduced relative to wild-type [33]. Nonetheless, because a) MITF is known to directly activate expression of pigmentation genes and cell cycle genes, b) TFAP2A and MITF bind many of the same loci in melanocytes [33], and c) there is prior evidence that TFAP2 paralogs can serve as pioneer factors [41, 42, 45], we hypothesized that TFAP2 is a pioneer factor that facilitates chromatin access for MITF.

To test the TFAP2-pioneer-factor model, the two paralogs with highest expression, TFAP2A and TFAP2C, were deleted from the SK-MEL-28 cells. Creation and integration of several genomic datasets permitted us to identify genomic elements that were bound by TFAP2A in WT cells and that either lost or gained H3K27Ac signal in *TFAP2*-KO cells; we inferred that these elements were enhancers directly activated or inhibited by TFAP2. As expected by the TFAP2-as-pioneer-factor model, at a subset of TFAP2A-activated enhancers, TFAP2 paralogs were required to keep chromatin open, and at all TFAP2-inhibited enhancers it was required to keep it

closed. We also showed that TFAP2 paralogs facilitated binding of another

transcription factor (MITF), that this occurred at loci where TFAP2 facilitates

chromatin access, and that this dependency was not reciprocal (at most loci bound

by both transcription factors). These findings imply that TFAP2 paralogs recruit

distinct transcription factors to distinct categories of enhancer, that these in turn

recruit enzymes that open chromatin or that condense it (TFAP2 paralogs also may

directly recruit such enzymes) [93], and that the state of chromatin accessibility

determines access of other transcription factors. Therefore, TFAP2 paralogs

facilitate chromatin access by other transcription factors, either as a pioneer factor or

as a linker that connects pioneer factors to chromatin-remodeling machinery.

Notably, the role of MITF at the co-bound enhancers is still largely unexplored.

Of note, there are loci where the pioneer factor FOXA1 appears to recruit proteins

like GRG3 (groucho-related gene 3) that condense chromatin [77, 78]. At TFAP2-

inhibited enhancers, our data do not distinguish between whether TFAP2 binds open

loci and closes them or instead binds to closed loci and helps stabilize the closed

state. Of note, the majority of TFAP2A peaks did not coincide with enhancers either

activated or inhibited by TFAP2 paralogs and were thus quiescent. In some cases,

such apparently quiescent TFAP2A peaks may represent loci that TFAP2 paralogs

have stably pioneered; a precedent for this scenario is that at a subset of elements

pioneered by PAX7, chromatin remains open after the removal of PAX7 [94]. To

explore this possibility, TFAP2A and ATAC-seq will need to be monitored in a

developmental time series, or in TFAP2-KO cells before and after TFAP2A has been

re-expressed.

Like other pioneer factors, TFAP2 can activate enhancers in a non-pioneer factor

mode. At a subset of TFAP2A-activated enhancers the H3K27Ac signal, but not the

ATAC-seq signal, was lower in TFAP2-KO cells than in WT cells. Evidence that such

elements are indeed TFAP2-activated enhancers is that their average H3K4Me3

signal was also lower in TFAP2-KO cells than in WT cells. We infer that TFAP2

activates these enhancers, but not as a pioneer factor. At such enhancers the

continued presence of TFAP2 is necessary for continued acetylation of histone H3

lysine 27 (H3K27Ac), which fits with evidence that TFAP2 binds the histone acetyl

transferase p300/CBP [95] and inhibits the NURD histone-deacetylase complex [93].

TFAP2 may attract other transcription factors without affecting nucleosome

positioning; indeed, some TFAP2-dependent MITF peaks were found at non-

pioneered TFAP2-activated enhancers. The fact that the TFAP2 binding site is not

strongly enriched in non-pioneered TFAP2-activated enhancers implies TFAP2 binds

these elements indirectly. Although we refer to these elements as non-pioneered

TFAP2-activated enhancers, our experimental design could not rule out the

possibility that some such enhancers were stably pioneered by TFAP2 as discussed

above. However, the observation that these sites are less enriched for the TFAP2

binding site than the TFAP2-pioneered-and-activated enhancers, and that the

binding sites of pioneer factors FOS and JUN are enriched [96], supports the

alternative model that such elements are simply pioneered by different transcription

factors.

Interestingly, at a subset of MITF/TFAP2A overlapping peaks where MITF was lost in

TFAP2-KO cells, TFAP2A binding was lost in *MITF*-KO cells (MITF/TFAP2A

mutually dependent peaks). There is precedent for reciprocal dependence of the

binding of pioneer factors, in the cases of both FOXA1 and steroid hormone

receptors, at subsets of sites where they are co-bound [97]. Notably, at many

MITF/TFAP2A mutually dependent peaks, the repressive mark H3K27Me3

accumulated in MITF-KO cells. This is consistent with evidence that the SWI/SNF

complex, which MITF probably recruits to such loci, competes for access to

chromatin against the H3K27Me3-depositing Polycomb repressor complex and that

the binding of pioneer factors is impeded by condensed H3K27Me3-positive

chromatin [80, 98]. We propose that at the majority of TFAP2-dependent MITF

peaks, but not at MITF/TFAP2A mutually dependent peaks, a measure of BRG1

binding is retained in MITF-KO cells, possibly recruited by another activator like

SOX10, but this is not the case for MITF/TFAP2A mutually dependent peaks.

Finally, by identifying enhancers regulated by TFAP2, whether in pioneer mode or

not, we were able to identify with high-confidence genes directly activated and

directly inhibited by TFAP2. The genes directly activated by TFAP2, including by the

specific mechanism of facilitating MITF access at a nearby enhancer, were enriched

in GO terms related to pigmentation and proliferation, and overlapped significantly

with expression profiles previously associated with high levels of MITF activity [82-

88]. The MITF "rheostat model" predicts that high MITF activity promotes cell

proliferation [2, 12]. Proliferation of both *MITF*-KO [17] and *TFAP2*-KO cells (shown

here) is slower than of the wild-type counterparts. We suggest there are two

mechanisms to explain the shared expression profiles and phenotypes between

TFAP2-KO cells and *MITF*-KO cells. First, the expression of *MITF* mRNA is about

40-50% lower in *TFAP2*-KO than WT cells. It is unclear how important this effect is

because the average read depth of MITF peaks was equivalent in WT and TFAP2-

KO cells, in particular at MITF-activated genes that are not also TFAP2-activated

(e.g., *MLANA*). Second, TFAP2 facilitates binding of MITF to enhancers and

promoters, leading to higher expression of associated MITF-target genes. The

"rheostat model" also predicts that low MITF activity promotes cell invasion [12], and

there was a large overlap of TFAP2-inhibited and MITF-inhibited genes and these

genes were enriched for those involved in cell-cell adhesion. We found evidence that

TFAP2 directly inhibits enhancers, but this did not involve TFAP2 and MITF

cooperativity. Directly-TFAP2-inhibited genes were associated with cell-cell adhesion

and *TFAP2*-KO cells formed large clusters when cultured on low adhesive plates.

Interestingly, in zebrafish melanoma, low *tfap2e* expression correlates with increased

metastases [32]. We have previously shown that MITF binds to enhancers near

MITF-inhibited genes [17]. However, mechanisms by which MITF inhibits gene

expression in melanoma cells and *in-vivo* remain to be elucidates. Given that an

MITF-low is a deadly status for melanoma, an interesting possibility for therapy

would be to effectively covert them to MITF-high by manipulating pioneer factors or

other MITF-cofactors. In summary, MITF activity in melanoma cells – and thus the

phenotypes of these cells – depends in part on the presence of transcription factors

that give MITF access to specific regulatory elements.

Materials and Methods

Zebrafish lines and maintenance

D. rerio were maintained in the University of Iowa Animal Care Facility according to a

standard protocol (protocol no. 6011616). All zebrafish experiments were performed

in compliance with the ethical regulations of the Institutional Animal Care and Use

Committee at the University of Iowa and in compliance with NIH guidelines.

Zebrafish embryos were maintained at 28.5 ◻C and staged by hours or days post-
fertilization (hpf or dpf).

Single cell suspension, fluorescence activated cell sorting and library
preparation

50 Tg(*mitfa*:GFP) transgeneic zebrafish embryos [99] were collected at 28 hours
post fertilization (hpf) and washed in PBS without Ca2+ or Mg2+ (Life technology).
Embryos were manually dechorionated under a dissecting scope (Leica KL300
LED), washed twice in 1X PBS and then cell dissociated cells using a pestle and
incubated in trypsin (0.25%)-EDTA (Life technology) at 33°C for 30 minutes
(ensuring to pipette mixed every 5 minutes). Reactions were quenched by adding
PBS supplemented with 5% fetal bovine serum (FBS, Life technologies). Dissociated
cells were centrifuged at 500 g for 5 minutes and cell pellets were washed with PBS-
5%FBS before passing thought a Bel Art™ SP Scienceware™ Flowmi™ 40 µm cell
strainers (ThermoFisher Scientific). Dissociated cells were re-suspended into single-
cell solution and analyzed at the University of Iowa Flow Cytometry Facility, using an
Aria Fusion instrument (Becton Dickinson, Franklin Lakes, NJ).

Single-cell RNA-sequencing (10x genomics) and data processing.

The sorted cellular suspensions were loaded on a 10x Genomics Chromium
instrument to generate single-cell gel beads in emulsion (GEMs). Approximately
10,000 cells were loaded per channel. Single-cell RNA-Seq libraries were prepared
using Single Cell 3′ Reagent Kits v2: Chromium™ Single Cell 3′ Library & Gel Bead
Kit v2, PN-120237; Single Cell 3′ Chip Kit v2 PN-120236 and i7 Multiplex Kit PN-
120262 (10x Genomics) following the Single Cell 3′ Reagent Kits v2 User Guide
(Manual Part # CG00052 Rev A). Libraries were run on an Illumina HiSeq 4000 as
2◻×◻150 paired-end reads. Sequencing results were demultiplexed and converted
to FASTQ format using Illumina bcl2fastq software.

A custom reference genome for the Zebrafish was constructed with GRCz11 primary
assembly (Ensembl) using Cell Ranger (Cell Ranger mkref function). 10X Genomics
scRNA-seq reads were then processed and aligned to this reference with Cell

Ranger (Cell Ranger Count function). Further analysis and visualization was

performed using Seurat (v4.1.0) [51], cells with fewer than 200 RNA feature counts

and greater than 5% mitochondrial contamination were removed with filtering. RNA

counts were normalized, and FindVariableFeatures was run with the following

parameters: selection.method = "vst", nfeatures = 2,000. The cells were originally

clustered in a UMAP by RunPCA then RunUMAP with dims 1:30. FindNeighbors

was run with dims 1:30 followed by FindClusters using a resolution of 1.2. Pigment

clusters were further analyzed by sub-setting out and re-clustering using dims 1:30,

which identified 10 clusters. Pseudotime was performed using Monocle3 (v0.2.3.0)

[100].

Generation of a zebrafish *tfap2e* loss-of-function allele

To generate the tfap2e loss-of-function allele, we designed paired (e.g., left and

right) zinc finger nucleases (ZFN) targeting exon 2 of the tfap2e locus resulting in

non-homologous end-joining and disruption of the open reading frame for Tfap2e.

Briefly, the online tool, ZiFiT [101], was used to identify an optimal ZFN target site

[utilizing the CoDa approach [102]. Once identified, a custom DNA fragment

encoding the entire left or right zinc finger array (ZFA) along with flanking XbaI and

BamHI restriction sites was synthesized (Integrated DNA Technologies, Coralville,

IA). Subsequently, the ZFA fragment was subcloned into pMLM290 (Addgene,

plasmid 21872), which includes a modified FokI nuclease domain [103]. Next, the

fully assembled ZFN was PCR amplified, directionally cloned into pENTR-D/TOPO

(ThermoFisher Scientific), and finally subcloned into pCS2+DEST using Gateway LR

Clonase II enzyme mix (ThermoFisher Scientific). Once assembled, the final pCS2+

plasmids were sequence verified, linearized, mRNA synthesized in vitro (mMessage

mMachine SP6 Kit, Ambion/ThermoFisher Scientific). Synthesized RNA was cleaned

using the Qiagen RNeasy Kit (Qiagen) and both left and right ZFN components were

co-injected into 1-cell stage zebrafish embryos. Following injections, embryos were

initially screened via PCR and restriction enzyme digest to confirm editing at the

target site. Upon confirmation, additional embryos from a similar clutch (F0's) were

allowed to develop into adulthood, 'mosaics' identified and out-crossed, and a stable

F1 generation isolated.

Cell lines, reagents, and antibodies

The cells referred to as WT throughout the document are the parent SK-MEL-28 (HTB-72) line. They and the derivative line, delta6-MITF knockout cells (referred to as MITF-KO cells in this work), were obtained from the laboratory of Dr. Eirikur Steingrimsson. The cells were grown in RPMI 1640 medium (Gibco #5240025) supplemented with 10% FBS (Gibco #10270106) at 5% CO_2 and 37°C. Cells were tested for, and determined to be free of, mycoplasma. SK-MEL-28 cells harbor the $BRAF^{V600E}$ and $p53^{L145R}$ mutations [104]. The following primary antibodies and their respective dilutions were used in western blotting (WB) and CUT&RUN experiments: anti-Tubulin (Sigma, #T6199), 1:5000 (WB); anti-MITF (Sigma, #HPA003259), 1:2000 (WB), 1:100 (CUT&RUN); anti-TFAP2A (Abcam, ab108311), 1:5000 (WB), 1:200 (CUT&RUN); anti-TFAP2C (Santa-Cruz #SC-12762 X), 1:1000 (WB); anti-H3K27Ac (EMD Millipore, #07-360), 1:100 (CUT&RUN); anti-H3K4Me3 (EMD Millipore, #05-745R), 1:100 (CUT&RUN); H3K27Me3 (EMD Millipore, #07-449), 1:100 (CUT&RUN); Rabbit IgG (EMD Millipore,#12-370), 1:100 (CUT&RUN); Mouse IgG (EMD Millipore, #12-371), 1:100 (CUT&RUN). The following secondary antibodies and their respective dilutions were used: Anti-mouse IgG(H+L) DyLight 800 conjugate (Cell Signaling Technologies, #5257), 1:20000; and anti-rabbit IgG(H+L) DyLight 680 conjugate Cell Signaling Technologies, #5366), 1:100. Images were captured using an Odyssey CLx Imager (LICOR Biosciences).

Purification of pA/G-MNase

E.coli strain BL21-DE3 was transformed with plasmid DNA pAG-MNase-6xHis (Addgene, plasmid #123461). Recombinant pAG-MNase was purified from cells grown in LB medium to OD600 0.6 at 37°C. Cells were induced with 0.5 mM IPTG and cultured for 16 hours at 20°C. Cell pellets were homogenized in lysis buffer (10 mM Tris, pH 7.5, 300 mM NaCl, 10 mM imidazole) containing lysozyme and protease inhibitors, then sonicated and the slurry was cleared by centrifugation (35K RPM, Ti70 rotor). The supernatant was subjected to IMAC chromatography (NI-NTA column) and to size-exclusion fractionation (Superdex 75) using a BioLogic DuoFlow QuadTec FPLC system (Bio-Rad). The purified pAG-MNase was concentrated by buffer exchange with ultrafiltration (Amicon Ultra-15, 10K). Finally, the purified pAG-

MNase was diluted in dilution buffer (10 mM Tris pH7.5, 120 mM NaCl, 0.01mM

EDTA, and 50% glycerol), and stored at -80°C.

Generation of *TFAP2A*; *TFAP2C* knockout cell lines (*TFAP2*-KO)

TFAP2-KO clones were generated using the Alt-R CRISPR-Cas9 technology from

Integrated DNA Technologies (IDT). Briefly, crRNAs targeting exon 2 of *TFAP2A* and

TFAP2C were designed using the Cas9 guide RNA design checker tool (crRNA

sequences below). Equimolar concentrations of crRNA and tracrRNA (IDT,

#1072532) were annealed to form gRNA complexes. The ribonucleoprotein (RNP)

complex was prepared by mixing gRNAs and Cas9 protein (IDT #1081058). SK-

MEL-28 cells were transfected with constructs encoding components of RNP

complexes using the Lipofectamine CRISPRmax Cas9 transfection reagent

(ThermoFisher #CMAX00015) following the manufacturer's protocol. Single-cell

colonies were screened by PCR and Sanger sequencing using primers flanking the

cut sites (primer sequences below). Mutant clones (clone 2.12 and clone 4.3) were

selected and further screened by western blotting, using anti-TFAP2A and anti-

TFAP2C antibodies. The control cell lines used in this study were generated

following this protocol but without adding gRNA duplexes.

crRNA	Sequence (5'-3')
TFAP2A_ex2_gRNA1	CGTCACGACGGCACCAGCAAGTTTTAGAGCTATGCT
TFAP2A_ex2_gRNA2	CTTACCTCACGCCATCGAGGGTTTTAGAGCTATGCT
TFAP2C_ex2_gRNA1	CGCCACGACGGGAGCAGCAAGTTTTAGAGCTATGCT
TFAP2C_ex2_gRNA2	CCACGACATGCCTCACCAGAGTTTTAGAGCTATGCT
Primers	**Sequence (5'-3')**
TFAP2A_geno_Fw	TCTCTTGTGCCCCCTCCATA
TFAP2A_geno_Rv	GCCCACCGACTGTATGTTCCA
TFAP2C_geno_Fw	CCGTGACCCCGATTTTGGAT
TFAP2C_geno_Rv	CGGCTTCACAGACATAGGCA

SDS-PAGE and Western blotting

TFAP2-KO and WT cells were washed in ice-cold PBS. RIPA buffer containing

protease inhibitors (Roche, cOmplete Mini) was added and cells were lysed on ice

for 20 minutes. Cell lysates were centrifuged at 14,000 g for 20 minutes and the

quantity of protein in the supernatants was quantified using Bradford assays (Bio-

Rad #5000002). Laemmli sample buffer (Bio-Rad #1610747, 5% 2-mercaptoethanol)

was added to 20 μg protein and samples were boiled at 95°C for 5 minutes before

being loaded onto a 10% SDS-polyacrylamide gel (Bio-Rad #4568034). Protein was

transferred to polyvinylidene fluoride (PVDF) membranes (Thermo Scientific

#88520), which were incubated overnight with primary antibody. Membranes were

washed 3 times with TBS-T and incubated with horseradish peroxidase-conjugated

anti-rabbit or anti-mouse for 1 hour at room temperature, washed, and imaged using

an Amersham Imager 600.

ATAC-seq

ATAC-seq was performed according to [105, 106] with minor alterations. Briefly,

70,000 *TFAP2*-KO cells (clone 2.12 and clone 4.3, four replicates each) and WT

cells (four replicates) were lysed in ice-cold lysis buffer (10 mM Tris-HCl, pH 7.4, 10

mM NaCl, 3 mM MgCl2, 0.1% NP-40: Sigma). Transposition was performed directly

on nuclei using 25 μl tagmentation reaction mix (Tagment DNA Buffer #15027866,

Tagment DNA Enzyme #15027865 from Illumina Tagment DNA kit #20034210).

Tagged DNA was subjected to PCR amplification and library indexing, using the

NEBNext High-Fidelity 2x PCR Master Mix (New England Biolabs #M0451S) with

Nextera DNA CD Indexes (Illumina #20015882), according to the following program:

72 □C for 5 minutes; 98 □C for 30 seconds; 12 cycles of 98 □C for 10 seconds, 63

□C for 30 seconds, and 72 □C for 1 minute. The PCR product was purified with 1.8

times the volume of Ampure XP beads (Beckman Coulter #A63881). Library quality

was assessed using a BioAnalyzer 2100 High Sensitivity DNA Chip (Agilent

Technologies). All DNA libraries that exhibited a nucleosome pattern were pooled

and processed for 150bp paired-end sequencing.

ATAC-seq peak calling and differential analysis

ATAC-seq was performed using 150 bp paired-end sequencing reads. Raw ATAC-seq reads were trimmed using Trim Galore Version 0.6.3 (Developed by Felix Krueger at the Babraham Institute) and aligned to human genome assembly hg19 (GRCh37) using Bowtie 2 [107, 108] with default parameters. Sorting, removal of PCR duplicates, and identification of fragments shorter than 100 bp as the nucleosome-depleted-regions (NDRs), was performed using BAM filter version 0.5.9. DeepTools version 3.3.0 [109] was used to check the reproducibility of the biological replicates and generate bigWig coverage files for visualization. Peaks were called using model-based analysis of ChIP-seq 2 (MACS2, version 2.1.1.20160309.6) [110]. NDRs for which accessibility differed between *TFAP2*-KO and WT cells were identified using DiffBind version 2.10 [111] with log2 fold-change threshold of >1 and a false discovery rate (FDR) < 0.05. NDRs that are directly regulated by TFAP2 were identified by overlapping differentially accessible NDRs with anti-TFAP2A CUT&RUN peaks; a 1-bp window was used to define overlap. Peaks were assigned to genes using GREAT with a peak-to-gene association rule of the nearest-gene-within-100 kb [112]. Both the raw ATAC-seq files and processed sequencing data presented in this manuscript have been deposited in the Gene Expression Omnibus (GEO) repository (GSE number pending).

CUT&RUN

CUT&RUN sequencing was performed in *TFAP2*-KO cells (clone 2.12 and clone 4.3, two replicates each) and WT cells (two replicates) as previously described [113, 114], but with minor modifications. Cells in log-phase culture (approximately 80% confluent) were harvested by cell scraping, centrifuged at 600 g (Eppendorf, centrifuge 5424) and washed twice in calcium-free wash-buffer (20 mM HEPES, pH7.5, 150 mM NaCl, 0.5 mM spermidine, and protease inhibitor cocktail cOmplete Mini, EDTA-free from Roche). Pre-activated concanavalin A-coated magnetic beads (Bangs Laboratories, Inc) were added to cell suspensions (2×10^5 cells) and incubated for 15 minutes at 4°C. Antibody buffer (wash-buffer with 2mM EDTA and 0.05% digitonin) containing anti-TFAP2A, anti-MITF, anti-H3K4Me3, anti-H3K27Me3,

anti-H3K27Ac or Rabbit IgG was added and cells were incubated overnight at 4ºC.

Cells were washed in dig-wash buffer (wash buffer containing 0.03% digitonin), and

pAG-MNase was added at a concentration of 500 µg/mL. The pAG-MNase reactions

were quenched with 2X Stop Buffer (340mM NaCl, 20mM EDTA, 4mM EGTA, 0.05%

digitonin, 100 µg/mL RNAse A (10 mg/mL, Thermo Fisher Scientific #EN0531), 50

µg/mL glycogen (20mg/mL, Thermo Fisher Scientific #R0561) and 2 pg/mL

sonicated yeast spike-in control). Released DNA fragments were treated with

1µL/mL phosphatase K (20mg/mL, Thermo Fisher Scientific #25530049) for 1 hour

at 50ºC and purified by phenol/chloroform-extraction and ethanol-precipitation.

Fragment sizes were analyzed using a 2100 Bioanalyzer (Agilent).

CUT&RUN library preparation and data analysis

CUT&RUN libraries were prepared using the KAPA Hyper Prep Kit (Roche). Quality

control post-library amplification consisted of fragment analysis using the 2100

Bioanalyzer (Agilent). Libraries were pooled, brought to equimolar concentrations,

and sequenced with 150 bp paired-end reads on an Illumina HiSeq X platform

(Novogene, Sacramento, CA). For quality control, paired-end FASTQ files were

processed using FastQC (Babraham Bioinformatics). Reads were trimmed using

Trim Galore Version 0.6.3 (Developed by Felix Krueger at the Babraham Institute)

and then mapped against the hg19 genome assembly using Bowtie2 version 2.1.0

[107, 108]. The mapping parameters and peak calling of MACS2 peaks [110] were

performed as previously described [113, 114] against their matching control IgG

samples. Differential analysis of H3K27Ac and of H3K27Me3 signal in WT and

TFAP2-KO cells was preformed using MACS2 with a Log2 fold-change threshold of

1, and p-value < 1×10^{-5}. Differential H3K4Me3, MITF and TFAP2A signal in WT,

TFAP2-KO and when mentioned *MITF*-KO cells was determined using DiffBind

version 2.10.0 [111] with a Log2 fold-change threshold of 1, and FDR < 0.05.

DeepTools version 3.3.0 was used to check the reproducibility of the biological

replicates, to generate bigwig normalized (RPKM) coverage files for visualization and

to plot average CUT&RUN-seq and ATAC-seq profiles (-plotProfile) and generate

heatmaps (-plotHeatmap) of normalized reads [109]. MultiBigwigSummary was used

to extract read counts (-outRawCounts) [109] and Prism was used to generate Violin

and Box plots. Peaks were assigned to genes using GREAT with a peak-to-gene

association rule of the nearest-gene-within-100 kb [112]

RNA-seq

Four replicate RNA-seq experiments were performed on *TFAP2*-KO cells (clone 2.12

and clone 4.3) and WT cells. Total RNA was extracted by direct cell lysis using the

RNeasy Plus Mini Kit with QiaShredder (Qiagen #47134). RNA samples with an

RNA integrity number (RIN) above nine were used for library generation and 150 bp

paired-end sequencing on the Illumina HiSeq2500 platform (Novogene, Sacramento,

CA). FASTQ sequence files were processed using FastQC (Babraham

Bioinformatics) for quality control, and reads were trimmed using Trim Galore

Version 0.6.3 (Developed by Felix Krueger at the Babraham Institute) and

subsequently aligned to human genome assembly hg19 (GRCh37) using STAR

[115]. The output of the --quantMode GeneCounts function of STAR was used for

the calculation of differential transcript expression using DESeq2 [116].The rlog

function was used to generate log2-transformed normalized counts. Adjusted p-value

< 0.05 was used as the threshold for statistically significant differences. Functional

enrichment analyses was performed using PANTHER [117]. A full list of genes

differentially expressed between *TFAP2*-KO and WT cells is provided in

Supplemental Table 1-3.

Motif analyses

Both de novo and known motifs were identified within 200 bp of TFAP2-activated

and TFAP2-inhibited enhancer and promoter peak summits using HOMER (-

findMotifsGenome).

Statistical analysis

Fisher's Exact Test was used to assess association between TFAP2-regulated

elements (enhancers and promoters) with TFAP2-regulated gene

expression. Fisher's Exact Test is a statistical significance test based on the

hypergeometric distribution and is used to analyze contingency tables to determine if

there is an association between the classes in the columns and rows of the

contingency table. Use of this test is relevant for the analysis of differential
expression because we classify enhancers as either TFAP2-activated
(downregulated in TFAP2-KO cells) or TFAP2-inhbited (upregulated in TFAP2-KO
cells) as well as near a target of interest (i.e., TFAP2-activated or TFAP2-inhibited)
or not. This test determines if there is an association between genes that are near an
TFAP2-regulated enhancer and genes whose expression varies between TFAP2-KO
cells and WT cells.
956
957   Data processing and analysis was performed in R and the code can be found at
958   https://GitHub.com/ahelv/Differential_Expression. GraphPad-Prism was used to
959   perform Student's t-test as indicated in the figure legends.
960
961   **Proliferation assay**
962
Cell proliferation was monitored at 24, 48, 72 and 96 hours using the CellTiter 96
Aqueous One Solution reagent kit according to the manufacturer's protocol. Cells
were serum depleted for 24 hours before transferring 5,000 cells per well of cell
culture treated 96-well plate (corning) containing RPMI media supplemented with
10% FBS and 1% PS. One the day of the experiment, CellTiter 96 Aqueous One
Solution was added to cells at a ratio of 6:1 (Media: solution). After 1 hour at 37^{o}C in
a humidified, 5% CO_2 atmosphere, the absorbance at 450nm was recorded using an
ELISA plate reader. Background absorbance was measured from cell free wells
containing media and CellTiter 96 Aqueous One Solution. The experiment was
performed in triplicate and repeated 3 times.
973
974   **Cluster formation assay**
975
Cluster formation assays were performed as previously described [32]. In brief, cells
were trypsinized and resuspended in RPMI media supplemented with 10% FBS and
1% PS. A flat bottom Low Attachment Surface-well plate (Corning) was seeded with
5,000 cells/well in a final volume of 100uL. Clusters were allowed to form over the
course of 3 days in a humidified 37°C incubator at which time the number of clusters
were counted per well using Image J software. The experiment was performed in
triplicate and repeated twelve times.

Wound scratch assay

A total of 500K cells were seeded per well of 6-well plate (Thermo Scientific, #

1483210) to reach confluent monolayer. Cells were incubated in serum free media

for 6 hours before wounding with a 200 µL pipette tip. Scratches were manually

imaged on an inverted light microscope (Leica #10445930) every 6 hours over a 24-

hour time period. The distance of scratch closure between WT and *TFAP2*-KO cells

were analyzed with Image J software.

Data availability

994 All raw sequencing files and processed data presented in this manuscript have been

995 deposited in the Gene Expression Omnibus (GEO) repositor. For ATAC-Seq and

996 CUT&RUN-seq data (GSE190610 and GSE153020) and for single cell RNA-Seq

997 (GSE198791).

Acknowledgements

1000 This work was supported by grants from the National Institutes of Health (NIH) to

1001 RAC (R01-AR062457), and a postdoctoral fellowship from the American Association

1002 for Anatomy (CK). The ES laboratory is supported by grants 207067 and 217768

1003 from the Research Fund of Iceland.

1004

Author Contributions

1006

Conceptualization: CK, RAC.

Formal analysis: CK, RD.

Funding acquisition: CK, ES, RAC.

Investigation: CK, RAC.

Methodology: CK, RD, HS, EVO, AH, GB.

Statistical analysis: CK, CMF, AH.

Supervision: ES, RAC.

Writing – original draft: CK, RAC.

Figure legends

1016

Figure 1: Expression of *tfap2* paralogs in the melanophore lineage precedes

activation of Mitfa-target genes *in-vivo*: (**A**) Uniform Manifold Approximation and

Projection (UMAP) obtained after clustering (dimensions, dims = 30, resolution = 1.2)

GFP+ cells (n = 11,217 cells) sorted from *Tg(mitfa:GFP)* zebrafish embryos at 28

hours post fertilization (hpf). Annotated cell clusters as labelled. (**B**) UMAP obtained

after re-clustering *sox10*-expressing clusters 6-12 (n = 1918 cells). Black line -

Monocle pseudotime trajectory analysis starting at *foxd3*+ neural crest cells showing

the progression through different pigment cell clusters as shown. (**C-F**) Violin plots

showing expression of select genes *foxd3, mitfa, tfap2e* and *dct* for each cell cluster

represented in B. *Blue arrows* point to the MIX-*tfap2*-low cluster, and show that *mitfa*,

but not pigmentation genes, are expressed in this cluster (*tfap2a* is also expressed in

this cluster). *Red arrows* point to the MIX-*tfap2*-high cluster, and show that *mitfa*,

tfap2e, and pigmentation genes are expressed in this cluster. (**G**) Dot plot

representing the expression of *sox10*, *tfap2* paralogs, *mitfa* and Mitfa-target genes,

and the percentage of cells expressing these genes in neural crest, MIX, MX and

melanophore clusters. Expression of *tfap2* paralogs are highlighted in MIX *tfap2*-low

and MIX *tfap2*-high clusters (red box). Size of dots represents percentage of cells

expressing the gene, and blue-red scale (low-high) represents relative average

expression among cells. (**H-K**) Lateral views of head and trunk of live embryos at 29

hpf, anterior to the left and dorsal to the top. Genotype as shown. Boxes, regions

magnified in accompanying panels H'-K.' (**H-H'**) A wild-type or heterozygous mutant

(sibling) embryo with normal melanophores (white arrowheads). (**I-I'**) A

tfap2e$^{ui157ui/157}$ embryo, with melanophores that are normal in terms of number,

differentiation and pigmentation (white arrowheads). (**J-J'**) A *tfap2a$^{low/low}$*

homozygous mutant embryo, with fewer melanophores than *tfap2e$^{ui157ui/157}$* and WT

sibling embryos. (**K-K'**) A *tfap2a$^{low/low}$; tfap2e $^{ui157ui/157}$* double-mutant embryo, with

fewer and paler melanophores than in *tfap2a$^{low/low}$* siblings. (**L**) Box plot illustrating

the number of pigmented melanophores in the dorsum of *tfap2a$^{low/low}$*, *tfap2a$^{low/low}$;*

tfap2e$^{+/ ui157}$, and *tfap2a$^{low/low}$; tfap2e$^{ui157ui/157}$* double mutant embryos at 36 hpf.

Center line, mean; box limits, upper and lower quartiles; whiskers, minimum and

maximum values; black dots, number of melanocytes per individual embryo

(*tfap2a$^{low/low}$*; n=9, *tfap2a$^{low/low}$; tfap2e$^{+/ui157}$;* n=32, *tfap2a$^{low/low}$; tfap2e$^{ui157/ui157}$*, n=10).

P-value according to the Student's t-test.

1050

Figure 2: Categories of TFAP2-regulated enhancers.

Box 1: TFAP2A peaks at open and active chromatin. (**1A**) TFAP2-pioneered-and-

activated enhancers show reduced nucleosome accessibility (ATAC-seq) and

reduced levels of active chromatin marks (H3K27Ac and H3K4Me3) in *TFAP2-KO*

cells compared to in WT cells. We infer that TFAP2 paralogs pioneer chromatin

access for transcriptional co-activators, like MITF and SOX10 (purple box), that in

turn recruit chromatin remodelling enzymes and histone modifying enzymes. (**1B**)

Non-pioneered TFAP2-activated enhancers show loss of active chromatin marks but

unchanged nucleosome accessibility in *TFAP2*-KO cells compared to in WT cells. At

these enhancers, we infer that TFAP2 paralogs recruit the binding of transcription

factors that, in turn, recruit histone modifying enzymes. TFAP2 paralogs also may

recruit such enzymes. It is possible that these elements are stably pioneered by

TFAP2 paralogs [94]. (**1C**) At TFAP2-independent elements, neither the nucleosome

accessibility nor active histone marks are altered in *TFAP2*-KO cells relative to in WT

cells. Both types of TFAP2-activated enhancer are significantly enriched near genes

whose expression is reduced in *TFAP2*-KO cells relative to in WT cells (i.e., TFAP2-

activated genes). Such genes are associated with the gene ontology (GO) terms cell

proliferation and pigmentation. TFAP2-independent elements are associated with

neither TFAP2-activated nor TFAP2-inhibited genes. **Box 2**: TFAP2A peaks at

closed and inactive chromatin. (**2A**) TFAP2-pioneered-and-inhibited enhancers show

increased nucleosome accessibility and increased levels active chromatin marks in

TFAP2-KO cells compared to in WT cells. At these sites we infer that TFAP2

paralogs recruit or stabilize the binding of enzymes that condense chromatin and

that inhibit the binding of transcriptional activators that are otherwise inclined to bind

at them. These elements are significantly enriched near genes whose expression is

elevated in *TFAP2*-KO cells relative to in WT cells (i.e., TFAP2-inhibited genes).

Such genes were associated with the GO terms cell-cell adhesion and cell migration.

(**2B**) At TFAP2-independent elements, neither the nucleosome accessibility nor

active histone marks are altered in *TFAP2*-KO cells relative to in WT cells. These

elements were associated with neither TFAP2-activated nor TFAP2-inhibited genes.

1081

Figure 3: TFAP2 paralogs facilitate gene expression by opening and condensing chromatin. (**A-D**) Screenshot of IGV genome browser (GRCH37/hg19), visualizing anti-TFAP2A CUT&RUN-seq (red), ATAC-seq (black), anti-H3K4Me3 CUT&RUN-seq (blue), anti-H3K27Ac CUT&RUN-seq (green) and RNA-seq (magenta) datasets at: (**A**) an TFAP2-pioneered-and-activated enhancer at the *PAX3* (+60 kb) locus; (**B**) a non-pioneered TFAP2A-activated enhancer at the *ENTPD6* (+26kb) locus; (**C**) an TFAP2-pioneered-and-inhibited enhancer at the *ADAM19* (+21 kb) locus; and (**D**) a non-pioneered TFAP2A-inhibited enhancer at the *FGF5* (+40kb) locus. Genotypes as labeled; y-axis in E applies to E-H, etc. (**E-H''**) Violin plots conveying normalized reads of (**E-H**) anti-H3K27Ac (two independent replicates), (**E'-H'**) ATAC-seq (four independent replicates) and (**E''-H''**) anti-H3K4Me3 (two independent replicates) CUT&RUN-seq at **E-E''**) TFAP2-pioneered-and-activated enhancers, (**F-F''**) non-pioneered TFAP2-activated enhancers, (**G-G''**) TFAP2-pioneered-and-inhibited enhancers, and (**H-H''**) non-pioneered TFAP2-inhibited enhancers. The number of peaks in each group is indicated. P-values shown were determined by Student's t-test. (**I-L**) Hypergeometric analysis of TFAP2 regulated enhancers at TFAP2-activated (*) and TFAP2-inhibited (**) genes in WT cells (FDR < 0.05, |log2FC| > 1). ns; not significant. (**M-O**) Enrichment of transcription factor binding motifs at (**M**) TFAP2-pioneered-and-activated enhancers, at (**N**) non-pioneered TFAP2-activated enhancers and at (**O**) TFAP2-pioneered-and-inhibited enhancers as determined using HOMER motif analysis. P values were calculated using ZOOPS scoring (zero or one occurrence per sequence) coupled with hypergeometric enrichment analysis. TF; transcription factor.

Figure 4: TFAP2 paralogs facilitate chromatin access by MITF. (**A**) Density heatmaps of anti-MITF CUT&RUN-seq in WT and *TFAP2*-KO SK-MEL-28 cells, and anti-TFAP2 CUT&RUN-seq in WT and *MITF*-KO SK-MEL-28 cells at TFAP2-dependent MITF peaks (first cluster), mutually dependent peaks (second cluster) and TFAP2-inhibited MITF peaks (third cluster). Number of peaks in each group as labelled. Regions shown are +/- 3 kb from peak center, normalized reads (RPKM). (**B-E**) Screenshot of IGV genome browser (GRCH37/hg19), showing anti-TFAP2A (red) CUT&RUN-seq in WT and *MITF*-KO cells, and anti-MITF (blue) CUT&RUN-seq, ATAC-seq (black) and anti-H3K27Ac (green) CUT&RUN-seq profiles in WT and *TFAP2*-KO cells. Examples of MITF binding at (**B**) a TFAP2-pioneered-and-activated

enhancer (**C**) a TFAP2-pioneered-and-activated promoter (**D**) a non-pioneered

TFAP2-activated enhancer and (**E**) a TFAP2-pioneered-and-inhibited enhancer.

Genotypes as labeled; y-axis in F applies to F-I, etc. (**F-I'**) Violin plots representing

(**F,G,H,I**) anti-MITF CUT&RUN-seq (two independent replicates), and (**F', G',H', I'**)

ATAC-seq (four independent replicates) at the indicated number of peaks. P-value

according to Student's t-test, ns; not statistically significant, normalized reads RPKM.

Association with gene expression; hypergeometric analysis of TFAP2-depednent

and TFAP2-inhibited MITF peaks are shown at TFAP2-activated and MITF-activated

genes (FDR < 0.05, log2FC > |1|). (**F''- I''**) Schematic representation of TFAP2-

pioneered-and-activated, TFAP2-activated and TFAP2-inhibited MITF peaks as

labelled; B; BAF complex (SWI/SNF), P; alternative pioneer factor. R; repressor

protein. Transcription factor binding sites indicated by small rectangles, TFAP2 (red),

MITF (blue) and alternative pioneer factor (yellow), example activator SOX10 (green)

. (**F''' – I'''**) Enrichment of transcription factor motifs using HOMER at (**F'''**) TFAP2-

pioneered -and-activated MITF peaks, (**G'''**) non-pioneered TFAP2-activated MITF

peaks, (**H'''**) mutually dependent peaks and (**I'''**) TFAP2-inhibited MITF peaks. TF;

transcription factor.

1133

Figure 5: TFAP2 and MITF co-regulate pigmentation and cell differentiation

genes in SK-MEL-28 cell lines. (**A**) Venn diagram representing the overlap of

genes apparently directly activated by MITF (i.e., expression lower in *MITF*-KO than

in WT cells, FDR < 0.05, and an MITF peak within 100kb of the TSS) and genes

directly activated by TFAP2 (i.e., expression lower in *TFAP2*-KO than in WT cells,

FDR < 0.05, and an TFAP2-activated enhancer of any category within 100kb of the

TSS). The number of overlapping genes with TFAP2-dependent MITF peaks are

also shown (*). (**B**) Gene ontology (GO) biological process analysis enriched among

MITF- and TFAP2-activated genes are shown (Top 5 hits). (**C**) A curated list of

pigment-associated genes [118] was intersected with directly MITF-activated,

overlapping genes of directly MITF- and TFAP2-activated genes, and directly

TFAP2-acitvated genes and represented by gene list. (**D**) Venn diagram

representing directly MITF inhibited genes (MITF peak within 100kb of a TSS),

based on RNA-seq, in *MITF*-KO versus WT cells (FDR < 0.05) and directly TFAP2

inhibited genes (TFAP2-inhibited enhancers, of any category, within 100kb of a

TSS), based on RNA-seq, in *TFAP2*-KO versus WT cells (FDR < 0.05). (**E**) Gene

ontology (GO) biological process analysis enriched among MITF- and TFAP2-

inhibited genes are shown (Top 5 most enriched GO terms). GO analysis was

performed using PANTHER. (**F**) Dot plot of enrichment analysis showing the

enrichment of melanoma gene signatures from the literature in directly TFAP2-

activated and TFAP2-inhibted genes. P value is red lowest to blue highest; gene

ratio is the fraction of all genes in the gene signature category that are included in

the set identified here. TFAP2-activated genes associated with TFAP2-dependent

MITF peaks are shown (*).

1158

Figure 6: TFAP2 paralogs promote cell proliferation and inhibit cell-cell

adhesion a melanoma cell line. (**A**) Heatmap representing the top 55 directly

TFAP2-activated genes and (**B**) the top 55 directly TFAP2-inhibited genes that are

associated with the GO terms cell pigmentation and cell-cell adhesion, respectively.

Four replicate RNA-Seq experiments are shown for WT cells and two clones of

TFAP2-KO cells (Clone 2.12 and Clone 4.3) (FDR < 0.05). (**C**) Growth curves (mean

± SE of mean) over 100 hours of cultivation for WT and *TFAP2*-KO SK-MEL-28 cells.

x-axis is time and y-axis is absorbance at 450nm which is directly proportional to

number of living. (**D**) Box plots representing the quantification of cluster formation on

low-bind plates after 72 hours of culture (n = 12 independent experiments, p < 0.001

by Student's t-test, plot shows mean ± SD). (**E**) Representative images of clusters

formed in WT and *TFAP2*-KO cells after 72 hours.

1171

Figure S1: UMAPs showing expression of select genes in *foxd3+* neural crest,

***twist+* cranial neural crest and basal cell clusters.** (**A-C**) Uniform Manifold

Approximation and Projection (UMAP) obtained after clustering (dimensions, dims =

30, resolution = 1.2) GFP+ cells (n = 11,217 cells) sorted from *Tg(mitfa:GFP)*

zebrafish embryos at 28 hours post fertilization (hpf). Feature plots showing

expression of select marker genes (**A**) *foxd3*+ neural crest clusters (**B**) *twist*+ cranial

neural crest clusters and (**C**) basal cell clusters. (**D**) Feature plots showing

expression of *mitfa* and *sox10*. Note that *mitfa* positive clusters are a subset of all

sox10 positive clusters. Dotted lines indicate clusters of interest.

1181

Figure S2: Expression of select marker genes in *foxd3/her+* neural crest, MIX,

xanthophore and iridophore cell clusters. (**A**) Violin plots showing expression of

Notch pathway genes in *foxd3/her*+ NC cluster 6. Note the lack of notch signaling

gene expression in *foxd3*+ NC cluster 7. (**B**) Violin plots showing expression of

select genes in MIX cell clusters, some of which are also expressed in MX,

xanthoblast and iridoblast cell clusters. (**C**) Violin plots showing expression of select

marker genes in xanthoblast and xanthophore cell clusters. (**D**) Violin plots showing

expression of select marker genes in iridoblast and iridophore cell clusters.

1190

1191   **Figure S3: Expression of TFAP2 paralogs in the pigment cell lineage.** (**A**) Violin

1192   plots showing expression of *tfap2* paralogs in NC, MIX *tfap2*-low*,* MIX *tfap2*-high,

1193   MX, melanophore, xanthoblast, xanthophore, iridoblast and iridophore cell clusters.

1194   Numeric labels below each violin plot refer cluster numbers as described in Figure

1195   1B.

1196

Figure S4: Expression of *mitfa* and *tfap2e* precede activation of Mitfa target

gene in the pigment cell lineage. (**A**) UMAP feature plots and **B**) violin plots

showing expression of *mitfa, tfap2e,* and genes in the melanin synthesis pathway,

whose human orthologs are known to be directly activated by MITF, in the pigment

cell lineage. Although robust *mitfa* expression is present in the MIX-*tfap2*-low cluster

(cluster 8a), expression of genes in the melanin synthesis pathway is absent in this

cluster. Expression of the latter begins concurrent with expression of *tfap2e* in the

MIX-*tfap2*-high cluster (cluster 8b).

Figure S5: Zebrafish homozygous for a frame-shift inducing mutation in *tfap2e*

do not display a pigmentation phenotype, and melanophores in *tfap2a/tfap2e*

double mutants are melanized by 48 hpf. (**A**) A 157 base pair mutation at the end

of *tfap2e* exon 2 disrupts splicing and results in a premature stop codon. (**B**) PCR

using primers in *tfap2e* exon 2 and intron 2 (e2-i2) amplifies a band of the expected

450 base pair size in *tfap2e* mutants but not wildtype (WT), whereas primers in exon

2 and exon 3 (e2-e3) amplify only in wild type. NTC: not template control. (**C**) qRT-

PCR analysis of *tfap2e* expression shows that the transcript is strongly decreased in

tfap2e$^{ui157/\ ui157}$ mutants, consistent with nonsense-mediated decay (Student's t-test,

**** $p<0.0001$). (**D-E**) *tfap2e* mutant zebrafish at 36 hpf, *tfap2e*$^{+/\ ui157}$ (**D**) and *tfap2e*

$^{ui157/\ ui157}$ (**E**) are phenotypically indistinguishable. (**F**) Histogram illustrating the

number of pigmented melanocytes in the dorsum of *tfap2e*$^{+/\ ui157}$ and *tfap2e*$^{ui157/\ ui157}$

mutant zebrafish embryos. **G-J)** Lateral views of 48 hpf embryos in the chorion. **G)**

A wildtype embryo shows normal melanocyte patterning. **(H)** A *tfap2a$^{-/-}$* mutant

embryo, with fewer and paler embryonic melanocytes than wildtypes**. (I)** In *tfap2a$^{-/-}$*

;tfap2e$^{+/-}$ and (**J**) in *tfap2a$^{-/-}$;tfap2e$^{-/-}$* mutants individual melanocytes are grossly as

melanized as they are in wild-types. These mutants are phenotypically

indistinguishable from *tfap2a$^{-/-}$* mutant embryos.

Figure S6: qRT-PCR expression analysis of *tfap2* paralogs in *tfap2* zebrafish

mutants. (**A**) Expression of *tfap2* paralogs in pooled embryos. Wildtype (n=20),

tfap2a$^{low/low}$ (n=20), and *tfap2a$^{low/low}$*; *tfap2e$^{ui157/ui157}$* (n=18). One-way ANOVA with

Bonferroni correction, * signifies adj. $p < 0.05$. RQ, relative quantification.

1227

Figure S7: TFAP2 binds to open and closed chromatin. (**A**) Screenshot of IGV

genome browser (GRCH37/hg19), visualizing anti-TFAP2A and IgG CUT&RUN-seq

profiles. Peaks were called using MACS2 software (two independent replicates) and

are illustrated by blue bars under the anti-TFAP2A track. (**B**) Density heatmap

centred on the 36,867 TFAP2A peaks identified by anti-TFAP2A CUT&RUN in WT

SK-MEL-28 cells. Regions shown are +/- 3 kb from peak center. Promoter peaks are

< 3kb and enhancer peaks >3kb from a transcriptional start site (TSS). Anti-TFAP2A,

anti-H3K4Me3 and anti-H3K27Ac CUT&RUN-seq, and ATAC-Seq profiles are

shown. Color code reflects normalized Reads Per Kilobase, per Million mapped

reads (RPKM). (**C**) Histogram representing H3K27Ac signal, binned from low to high

read-depth (normalized RPKM) on the x-axis and percentage of TFAP2A promoter

peaks (black) and TFAP2A enhancer peaks (red) on the y-axis. (**D**) Violin plots

illustrating TFAP2A and IgG normalized reads (RPKM) at open (ATAC-peaks) and at

closed chromatin (no ATAC-peak) (**E**) Density heatmap representing TFAP2A

CUT&RUN and ATAC-seq profiles at TFAP2A peaks that overlap nucleosome

depleted regions (ATAC-peaks) and at nucleosome bound DNA (no ATAC-peak),

the number of TFAP2A peaks in each group are as labeled.

1245

1246   **Figure S8: Example loci of TFAP2A peaks at open and closed chromatin.** (**A-B**)

1247   IGV screenshots of TFAP2A peaks that (**A**) closed chromatin and (**B**) open

1248   chromatin, based on ATAC-seq, in SK-MEL-28 cells. Genes names and distance to

1249   that gene's TSS are as labeled. Note that scale for ATAC-seq reads (RPKM) is

constant but for TFAP2A CUT&RUN is variable. *Schematics*, renderings of TFAP2 in

A) closed and **B**) open chromatin. (**C**) HOMER motif analysis at TFAP2A peaks at

closed chromatin and (**D**) at open chromatin. TF, transcription factors; the top

ranking transcription factor motif is shown, with P-values calculated with HOMER-

based hypergeometric enrichment analysis.

Figure S9: Generation of *TFAP2A*; *TFAP2C* double mutant SK-MEL-28 cell

lines. (**A**) RNA-seq showing transcript counts of TFAP2 paralogs in SK-MEL-28

cells. Transcript counts for WT cells (n=4) and two *TFAP2A;TFAP2C* double

knockout clones (4 replicates each) are shown. The expression of WT and mutant

alleles of *TFAP2A* is comparable between cell lines whereas mutant alleles of

TFAP2C are strongly reduced. (**B**) Two CRISPR guide RNAs (gRNA) each were

designed to target exon 2 of *TFAP2A* and *TFAP2C* (yellow boxes). (**C**) A 401 base

pair inversion and a 452 base pair deletion at exon 2 of *TFAP2A* and *TFAP2C*,

respectively, was identified in clone 4.3. *TFAP2A* and *TFAP2C* mutant alleles

resulted in a frame-shift and premature stop codon in alleles of both genes. (**D**) A

405 base pair deletion at exon 2 of *TFAP2A* resulted in a frame-shift and premature

stop codon in clone 2.12. A 455 base pair deletion, and a 70 base pair insertion, 1

base pair deletion (Indel) was identified in exon 2 of *TFAP2C*. Such mutations

resulted in a frame-shift and premature stop codon. Additional permutations were not

identified at exon 2 of *TFAP2A* or *TFAP2C* in clone 4.3 or clone 2.12 cells. Inv,

inversion; Del, deletion. (**E**) Western blot analysis confirming loss of TFAP2A and

TFAP2C immunoactivity in clone 4.3 and clone 2.12 cell lines.

Figure S10: TFAP2 paralogs directly activate and inhibit promoters as pioneer

factors. (**A-D**) Screenshot of IGV genome browser (GRCH37/hg19), visualizing anti-

TFAP2A CUT&RUN-seq (red), ATAC-seq (black), anti-H3K4Me3 CUT&RUN-seq

(blue), anti-H3K27Ac CUT&RUN-seq (green) and RNA-seq (magenta) datasets at

(**A**) TFAP2-pioneered-and-activated promoters, (**B**) non-pioneered TFAP2-activated

promoters, (**C**) TFAP2-pioneered-and-inhibited promoters and (**D**) non-pioneered

TFAP2-activated promoters . Genotypes as labeled; y-axes are grouped scaled per

dataset. (**E-H**) Violin plots representing anti-H3K27Ac (two independent replicates)

(**E'-H'**) ATAC-seq (four independent replicates), and (**E''-H''**) anti-H3K4Me3 (two

independent replicates) normalized reads at (**E-E''**) TFAP2-pioneered-and-activated

promoters, (**F'F''**) non-pioneered TFAP2-activated promoters, (**G-G''**) TFAP2-

pioneered-and-inhibited promoters, (**H-H''**) non-pioneered TFAP2-inhibited

promoters. P-values shown were determined by Student's t-test. (**I-L**)

Hypergeometric analysis of TFAP2 regulated enhancers at TFAP2-activated (*) and

TFAP2-inhibited (**) genes in *TFAP2*-KO cells (FDR < 0.05, |log2FC| > 1). The

number of promoters in each category of TFAP2-regulated promoters is shown.

1291

Figure S11: Examples of TFAP2-activated and -inhibited promoters. (**A-B**)

Screenshots of IGV genome browser (GRCH37/hg19) visualizing anti-TFAP2A, anti-

H3K4Me3, anti-H3K27Ac CUT&RUN-seq, ATAC-seq and RNA-seq profiles at (**A**)

the TFAP2-pioneered-and-activated *ZNF540* promoter and (**B**) the TFAP2-

pioneered-and-inhibited *S100A16* promoter. Genotypes as labeled; y-axes are

scaled per dataset.

1298

Figure S12: TFAP2 paralogs modulate the binding of MITF. (**A**) Density heatmap

centred on 9,413 peaks co-bound by TFAP2A and MITF, showing anti-TFAP2, anti-

MITF, ATAC-seq, and anti-H3K27Ac CUT&RUN profiles in SK-MEL-28 cells.

Regions shown are +/- 5 kb from peak center. (**B**) Violin plot showing anti-MITF

CUT&RUN signal (RPKM) in *TFAP2*-KO and SK-MEL-28 cells at MITF peaks that

are not bound by TFAP2A. There is no change in the average read-depth of such

MITF peaks in WT vs TFAP2-KO cells. ns; non-significant by Student's t-test. **(C)**

Scatterplot of TFAP2-dependent (i.e., TFAP2-activated and TFAP2-inhibited) MITF

peaks. X axis is log2 normalized MITF CUT&RUN read depth in peaks in WT cells. Y

axis is log2FC of MITF CUT&RUN read depth in *TFAP2*-KO versus WT cells. **(D)**

Screenshots of IGV genome browser (GRCH37/hg19); genotypes as labeled,

visualizing 5kb windows centered on TFAP2A peaks and showing indicated genomic

data at several examples of TFAP2-activated MITF peaks and mutually-dependent

MITF/TFAP2 peaks, and a single example of a non-overlapping, TFAP2-independent

MITF peak. **(E)** Violin plot showing increased H3K27Me3 CUT&RUN signal at

mutually-dependent TFAP2A/MITF peaks in *MITF*-KO cells versus WT cells. P-

values were determined by Student's t-test. Y axis is normalized reads (RPKM). **(F-**

G) Screenshots of IGV genome browser (GRCH37/hg19), visualizing anti-

H3K27me3 CUT&RUN-seq profiles in *MIT*F-KO and WT cells. Peaks were called

using MACS2 software (two independent replicates) and are illustrated by blue bars. Two examples loci of mutually-dependent TFAP2A/MITF peaks showing increased H3K27Me3 signals are shown **(F)** at the *TRPM1* promoter and **(G)** at two *FRMD4B* enhancers. Note that H3K27Me3 signal extends much farther than just the region of the TFAP2A/MITF peak. Mutually-dependent peaks are indicated by red arrows.

Figure S13: Density heatmap of TFAP2 regulated MITF peaks. Density heatmap centered on TFAP2-activated MITF peaks (first cluster), mutually dependent (mutually activated) peaks (second cluster), TFAP2-inhibited MITF peaks and (third cluster) and TFAP2-independent MITF peaks (forth cluster), showing two replicates of anti-MITF CUT&RUN in SK-MEL-28 and *TFAP2*-KO cells, and two replicates of anti-TFAP2A CUT&RUN in SK-MEL-28 and *MITF*-KO cells. Regions shown are +/- 5 kb from peak center.

Figure S14: TFAP2 and MITF do not co-inhibit enhancers at TFAP2-inhibited or MITF-inhibited genes in SK-MEL-28 cells. (**A-A'**) Violin plot of TFAP2-activated MITF peaks at TFAP2-pioneered-and-inhibited enhancers, which are potentially TFAP2/MITF co-inhibited enhancers, showing (**A**) anti-MITF CUT&RUN and (**A'**) ATAC-Seq profiles in *TFAP2*-KO and WT cells. Such loci were not significantly enriched at MITF-inhibited or TFAP2-inhibited genes. (**A''**) Schematic of theoretical TFAP2/ MITF co-inhibited enhancers. In this example, TFAP2 is a pioneer factor recruiting MITF in its repressor form to condense chromatin. Loss of TFAP2 in *TFAP2*-KO cells results in loss of MITF-repressor binding and opening of chromatin by an alternative pioneer factor.

Figure S15: MITF-activated genes associated with TFAP2-independent MITF peaks are enriched for cell cycle and DNA-repair. (**A**) Plot-profile showing MITF CUT&RUN peak signal at TFAP2-independent MITF peaks in *TFAP2*-KO and WT SK-MEL-28 cell lines. (**B**) Genes associated with TFAP2-independent anti-MITF peaks were analyzed for enriched gene ontology biological process using GREAT (single nearest gene +/- 100kb).

Figure S16: A possible TFAP2-pioneered-and-activated enhancer at intron 2 of

MITF. Screenshot of IGV genome browser (GRCH37/hg19) visualizing anti-TFAP2A,

ATAC-seq, anti-H3K27Ac and RNA-seq profiles at intron 2 of MITF. Dashed

rectangle indicates an TFAP2-activated ATAC-seq peak; like the ATAC-seq signal,

the H3K27Ac-signal is significantly lower in TFAP2-KO cells relative to in WT cells

implying this is a TFAP2-pioneered-and-activated enhancer, although the log2 fold

change of the H3K27Ac signal does not meet the cut-off we used to define this

category of enhancer elsewhere in the paper. MITF and downstream regions are

shown, blue arrows indicate strand orientation and horizontal rectangles the exons.

Genotypes are as labeled; y-axes are grouped scaled per dataset.

1361

Figure S17: GO analysis of TFAP2-activated and inhibited genes. (**A**) Heatmap

representing the top 55 directly TFAP2-activated genes that are included in the Gene

Ontology (GO) biological process categories "cell pigmentation" and "cell

differentiation." Four replicate RNA-Seq experiments are shown for WT cells and two

clones of *TFAP2*-KO cells (Clone 2.12 and Clone 4.3) (FDR < 0.05). (**B-C**) GO

biological process terms enriched among (**B**) directly TFAP2-activated genes and

(**C**) directly TFAP2-inhibited genes.

1369

1370 **Figure S18: Scratch assay.** Micrographs of (top) WT and (bottom) *TFAP2*-KO cells

1371 in petri dishes at the indicated time after a scratch through the cells was made with a

1372 pipette tip. After 24 hours, the WT but not the *TFAP2*-KO cells had filled in the

1373 scratch through migration and/or proliferation.

1374

1375

1376

1377

1378

1379

1380

1381

1382

1383

References cited:

1.   Seberg HE, Van Otterloo E, Cornell RA. Beyond MITF: Multiple transcription
factors directly regulate the cellular phenotype in melanocytes and melanoma. Pigment Cell
Melanoma Res. 2017;30(5):454-66. doi: 10.1111/pcmr.12611. PubMed PMID: 28649789.
2.   Goding CR. Mitf from neural crest to melanoma: signal transduction and transcription
in the melanocyte lineage. Genes Dev. 2000;14(14):1712-28. Epub 2000/07/18. PubMed
PMID: 10898786.
3.   Hartman ML, Czyz M. MITF in melanoma: mechanisms behind its expression and
activity. Cell Mol Life Sci. 2015;72(7):1249-60. doi: 10.1007/s00018-014-1791-0. PubMed
PMID: 25433395; PubMed Central PMCID: PMCPMC4363485.
4.   Mollaaghababa R, Pavan WJ. The importance of having your SOX on: role of SOX10
in the development of neural crest-derived melanocytes and glia. Oncogene.
2003;22(20):3024-34. PubMed PMID: 12789277.
5.   Atchison ML. Function of YY1 in Long-Distance DNA Interactions. Front Immunol.
2014;5:45. Epub 2014/02/28. doi: 10.3389/fimmu.2014.00045. PubMed PMID: 24575094;
PubMed Central PMCID: PMCPMC3918653.
6.   Eckert D, Buhl S, Weber S, Jager R, Schorle H. The AP-2 family of transcription
factors. Genome biology. 2005;6(13):246. doi: 10.1186/gb-2005-6-13-246. PubMed PMID:
16420676; PubMed Central PMCID: PMC1414101.
7.   Hoek KS, Schlegel NC, Eichhoff OM, Widmer DS, Praetorius C, Einarsson SO, et al.
Novel MITF targets identified using a two-step DNA microarray strategy. Pigment Cell
Melanoma Res. 2008;21(6):665-76. Epub 2008/12/11. doi: 10.1111/j.1755-
148X.2008.00505.x. PubMed PMID: 19067971.
8.   Strub T, Giuliano S, Ye T, Bonet C, Keime C, Kobi D, et al. Essential role of
microphthalmia transcription factor for DNA replication, mitosis and genomic stability in
melanoma. Oncogene. 2011;30(20):2319-32. Epub 2011/01/25. doi: onc2010612 [pii]
10.1038/onc.2010.612. PubMed PMID: 21258399.
9.   Betancur P, Bronner-Fraser M, Sauka-Spengler T. Assembling neural crest regulatory
circuits into a gene regulatory network. Annu Rev Cell Dev Biol. 2010;26:581-603. Epub
2009/07/07. doi: 10.1146/annurev.cellbio.042308.113245. PubMed PMID: 19575671.
10.   Van Otterloo E, Li W, Bonde G, Day KM, Hsu MY, Cornell RA. Differentiation of
zebrafish melanophores depends on transcription factors AP2 alpha and AP2 epsilon. PLoS
genetics. 2010;6(9). Epub 2010/09/24. doi: 10.1371/journal.pgen.1001122. PubMed PMID:
20862309; PubMed Central PMCID: PMC2940735.
11.   Van Otterloo E, Li W, Garnett A, Cattell M, Medeiros DM, Cornell RA. Novel
Tfap2-mediated control of soxE expression facilitated the evolutionary emergence of the
neural crest. Development. 2012;139(4):720-30. Epub 2012/01/14. doi: 10.1242/dev.071308.
PubMed PMID: 22241841; PubMed Central PMCID: PMC3265060.
12.   Carreira S, Goodall J, Denat L, Rodriguez M, Nuciforo P, Hoek KS, et al. Mitf
regulation of Dia1 controls melanoma proliferation and invasiveness. Genes Dev.
2006;20(24):3426-39. Epub 2006/12/22. doi: 10.1101/gad.406406. PubMed PMID:
17182868; PubMed Central PMCID: PMC1698449.
13.   Rambow F, Marine JC, Goding CR. Melanoma plasticity and phenotypic diversity:
therapeutic barriers and opportunities. Genes Dev. 2019;33(19-20):1295-318. Epub
2019/10/03. doi: 10.1101/gad.329771.119. PubMed PMID: 31575676; PubMed Central
PMCID: PMCPMC6771388.
14.   de la Serna IL, Ohkawa Y, Higashi C, Dutta C, Osias J, Kommajosyula N, et al. The
microphthalmia-associated transcription factor requires SWI/SNF enzymes to activate
melanocyte-specific genes. J Biol Chem. 2006;281(29):20233-41. Epub 2006/05/02. doi:
M512052200 [pii]
10.1074/jbc.M512052200. PubMed PMID: 16648630.

15.   Laurette P, Strub T, Koludrovic D, Keime C, Le Gras S, Seberg H, et al.
Transcription factor MITF and remodeller BRG1 define chromatin organisation at regulatory
elements in melanoma cells. eLife. 2015;4. doi: 10.7554/eLife.06857. PubMed PMID:
25803486; PubMed Central PMCID: PMC4407272.
16.   Lang D, Lu MM, Huang L, Engleka KA, Zhang M, Chu EY, et al. Pax3 functions at a
nodal point in melanocyte stem cell differentiation. Nature. 2005;433(7028):884-7. Epub
2005/02/25. doi: nature03292 [pii]
10.1038/nature03292. PubMed PMID: 15729346.
17.   Dilshat R, Fock V, Kenny C, Gerritsen I, Lasseur RMJ, Travnickova J, et al. MITF
reprograms the extracellular matrix and focal adhesion in melanoma. eLife. 2021;10. Epub
2021/01/14. doi: 10.7554/eLife.63093. PubMed PMID: 33438577; PubMed Central PMCID:
PMCPMC7857731.
18.   Du J, Li L, Ou Z, Kong C, Zhang Y, Dong Z, et al. FOXC1, a target of polycomb,
inhibits metastasis of breast cancer cells. Breast Cancer Res Treat. 2012;131(1):65-73. Epub
2011/04/06. doi: 10.1007/s10549-011-1396-3. PubMed PMID: 21465172.
19.   Mitchell PJ, Timmons PM, Hebert JM, Rigby PW, Tjian R. Transcription factor AP-2
is expressed in neural crest cell lineages during mouse embryogenesis. Genes Dev.
1991;5(1):105-19.
20.   Moser M, Ruschoff J, Buettner R. Comparative analysis of AP-2 alpha and AP-2 beta
gene expression during murine embryogenesis. Developmental Dynamics. 1997;208(1):115-
24.
21.   Bamforth SD, Braganca J, Eloranta JJ, Murdoch JN, Marques FI, Kranc KR, et al.
Cardiac malformations, adrenal agenesis, neural crest defects and exencephaly in mice
lacking Cited2, a new Tfap2 co-activator. Nat Genet. 2001;29(4):469-74. PubMed PMID:
11694877.
22.   Brewer S, Jiang X, Donaldson S, Williams T, Sucov HM. Requirement for AP-2alpha
in cardiac outflow tract morphogenesis. Mech Dev. 2002;110(1-2):139-49. PubMed PMID:
11744375.
23.   Knight RD, Nair S, Nelson SS, Afshar A, Javidan Y, Geisler R, et al. lockjaw encodes
a zebrafish tfap2a required for early neural crest development. Development.
2003;130(23):5755-68. PubMed PMID: 14534133.
24.   Tan CC, Sindhu KV, Li S, Nishio H, Stoller JZ, Oishi K, et al. Transcription factor
Ap2delta associates with Ash2l and ALR, a trithorax family histone methyltransferase, to
activate Hoxc8 transcription. Proc Natl Acad Sci U S A. 2008;105(21):7472-7. Epub
2008/05/23. doi: 10.1073/pnas.0711896105. PubMed PMID: 18495928; PubMed Central
PMCID: PMCPMC2396708.
25.   Schorle H, Meier P, Buchert M, Jaenisch R, Mitchell PJ. Transcription factor AP-2
essential for cranial closure and craniofacial development. Nature. 1996;381(6579):235-8.
26.   Luo T, Matsuo-Takasaki M, Thomas ML, Weeks DL, Sargent TD. Transcription
factor AP-2 is an essential and direct regulator of epidermal development in Xenopus. Dev
Biol. 2002;245(1):136-44. PubMed PMID: 11969261.
27.   Wang X, Bolotin D, Chu DH, Polak L, Williams T, Fuchs E. AP-2alpha: a regulator
of EGF receptor signaling and proliferation in skin epidermis. J Cell Biol. 2006;172(3):409-
21. Epub 2006/02/02. doi: jcb.200510002 [pii]
10.1083/jcb.200510002. PubMed PMID: 16449191; PubMed Central PMCID:
PMC2063650.
28.   Kuckenberg P, Kubaczka C, Schorle H. The role of transcription factor
Tcfap2c/TFAP2C in trophectoderm development. Reproductive biomedicine online.
2012;25(1):12-20. doi: 10.1016/j.rbmo.2012.02.015. PubMed PMID: 22560121.

29.   Li W, Cornell RA. Redundant activities of Tfap2a and Tfap2c are required for neural
crest induction and development of other non-neural ectoderm derivatives in zebrafish
embryos. Dev Biol. 2007;304(1):338-54. Epub 2007/01/30. doi: S0012-1606(06)01501-6
[pii]
10.1016/j.ydbio.2006.12.042. PubMed PMID: 17258188; PubMed Central PMCID:
PMC1904501.
30.   Wang X, Pasolli HA, Williams T, Fuchs E. AP-2 factors act in concert with Notch to
orchestrate terminal differentiation in skin epidermis. J Cell Biol. 2008;183(1):37-48. Epub
2008/10/01. doi: jcb.200804030 [pii]
10.1083/jcb.200804030. PubMed PMID: 18824566; PubMed Central PMCID:
PMC2557040.
31.   Kołat D, Kałuzińska Ż, Bednarek AK, Płuciennik E. WWOX Loses the Ability to
Regulate Oncogenic AP-2γ and Synergizes with Tumor Suppressor AP-2α in High-Grade
Bladder Cancer. Cancers (Basel). 2021;13(12). Epub 2021/07/03. doi:
10.3390/cancers13122957. PubMed PMID: 34204827; PubMed Central PMCID:
PMCPMC8231628.
32.   Campbell NR, Rao A, Zhang M, Baron M, Heilmann S, Deforet M, et al. Cell state
diversity promotes metastasis through heterotypic cluster formation in melanoma.
Developmental Cell. 2021:2020.08.24.265140. doi: 10.1101/2020.08.24.265140.
33.   Seberg HE, Van Otterloo E, Loftus SK, Liu H, Bonde G, Sompallae R, et al. TFAP2
paralogs regulate melanocyte differentiation in parallel with MITF. PLoS genetics.
2017;13(3):e1006636. doi: 10.1371/journal.pgen.1006636. PubMed PMID: 28249010;
PubMed Central PMCID: PMCPMC5352137.
34.   Rothstein M, Simoes-Costa M. Heterodimerization of TFAP2 pioneer factors drives
epigenomic remodeling during neural crest specification. Genome Res. 2020;30(1):35-48.
Epub 20191217. doi: 10.1101/gr.249680.119. PubMed PMID: 31848212; PubMed Central
PMCID: PMCPMC6961570.
35.   Sennett R, Wang Z, Rezza A, Grisanti L, Roitershtein N, Sicchio C, et al. An
Integrated Transcriptome Atlas of Embryonic Hair Follicle Progenitors, Their Niche, and the
Developing Skin. Dev Cell. 2015;34(5):577-91. Epub 20150806. doi:
10.1016/j.devcel.2015.06.023. PubMed PMID: 26256211; PubMed Central PMCID:
PMCPMC4573840.
36.   Praetorius C, Grill C, Stacey SN, Metcalf AM, Gorkin DU, Robinson KC, et al. A
polymorphism in IRF4 affects human pigmentation through a tyrosinase-dependent
MITF/TFAP2A pathway. Cell. 2013;155(5):1022-33. doi: 10.1016/j.cell.2013.10.022.
PubMed PMID: 24267888; PubMed Central PMCID: PMC3873608.
37.   Rambow F, Job B, Petit V, Gesbert F, Delmas V, Seberg H, et al. New Functional
Signatures for Understanding Melanoma Biology from Tumor Cell Lineage-Specific
Analysis. Cell Rep. 2015;13(4):840-53. Epub 2015/10/23. doi: 10.1016/j.celrep.2015.09.037.
PubMed PMID: 26489459; PubMed Central PMCID: PMCPMC5970542.
38.   Voss TC, Hager GL. Dynamic regulation of transcriptional states by chromatin and
transcription factors. Nat Rev Genet. 2014;15(2):69-81. Epub 2013/12/18. doi:
10.1038/nrg3623. PubMed PMID: 24342920; PubMed Central PMCID: PMCPMC6322398.
39.   Zaret KS, Carroll JS. Pioneer transcription factors: establishing competence for gene
expression. Genes Dev. 2011;25(21):2227-41. doi: 10.1101/gad.176826.111. PubMed PMID:
22056668; PubMed Central PMCID: PMCPMC3219227.
40.   Zaret KS. Pioneer Transcription Factors Initiating Gene Network Changes. Annu Rev
Genet. 2020;54:367-85. Epub 2020/09/05. doi: 10.1146/annurev-genet-030220-015007.
PubMed PMID: 32886547; PubMed Central PMCID: PMCPMC7900943.

41.   Sherwood RI, Hashimoto T, O'Donnell CW, Lewis S, Barkal AA, van Hoff JP, et al.
Discovery of directional and nondirectional pioneer transcription factors by modeling DNase
profile magnitude and shape. Nat Biotechnol. 2014;32(2):171-8. Epub 2014/01/21. doi:
10.1038/nbt.2798. PubMed PMID: 24441470; PubMed Central PMCID: PMCPMC3951735.
42.   Pihlajamaa P, Sahu B, Lyly L, Aittomaki V, Hautaniemi S, Janne OA. Tissue-specific
pioneer factors associate with androgen receptor cistromes and transcription programs.
EMBO J. 2014;33(4):312-26. Epub 2014/01/24. doi: 10.1002/embj.201385895. PubMed
PMID: 24451200; PubMed Central PMCID: PMCPMC3989639.
43.   Tan SK, Lin ZH, Chang CW, Varang V, Chng KR, Pan YF, et al. AP-2gamma
regulates oestrogen receptor-mediated long-range chromatin interaction and gene
transcription. EMBO J. 2011;30(13):2569-81. doi: 10.1038/emboj.2011.151. PubMed PMID:
21572391; PubMed Central PMCID: PMC3155293.
44.   Grossman SR, Engreitz J, Ray JP, Nguyen TH, Hacohen N, Lander ES. Positional
specificity of different transcription factor classes within enhancers. Proc Natl Acad Sci U S
A. 2018;115(30):E7222-E30. Epub 2018/07/11. doi: 10.1073/pnas.1804663115. PubMed
PMID: 29987030; PubMed Central PMCID: PMCPMC6065035.
45.   Pastor WA, Liu W, Chen D, Ho J, Kim R, Hunt TJ, et al. TFAP2C regulates
transcription in human naive pluripotency by opening enhancers. Nat Cell Biol.
2018;20(5):553-64. Epub 2018/04/27. doi: 10.1038/s41556-018-0089-0. PubMed PMID:
29695788; PubMed Central PMCID: PMCPMC5926822.
46.   Li L, Wang Y, Torkelson JL, Shankar G, Pattison JM, Zhen HH, et al. TFAP2C- and
p63-Dependent Networks Sequentially Rearrange Chromatin Landscapes to Drive Human
Epidermal Lineage Commitment. Cell Stem Cell. 2019;24(2):271-84 e8. Epub 2019/01/29.
doi: 10.1016/j.stem.2018.12.012. PubMed PMID: 30686763; PubMed Central PMCID:
PMCPMC7135956.
47.   Fernandez Garcia M, Moore CD, Schulz KN, Alberto O, Donague G, Harrison MM,
et al. Structural Features of Transcription Factors Associating with Nucleosome Binding. Mol
Cell. 2019;75(5):921-32 e6. Epub 2019/07/16. doi: 10.1016/j.molcel.2019.06.009. PubMed
PMID: 31303471; PubMed Central PMCID: PMCPMC6731145.
48.   Swinstead EE, Paakinaho V, Presman DM, Hager GL. Pioneer factors and ATP-
dependent chromatin remodeling factors interact dynamically: A new perspective: Multiple
transcription factors can effect chromatin pioneer functions through dynamic interactions
with ATP-dependent chromatin remodeling factors. Bioessays. 2016;38(11):1150-7. Epub
2016/10/27. doi: 10.1002/bies.201600137. PubMed PMID: 27633730; PubMed Central
PMCID: PMCPMC6319265.
49.   Aras S, Saladi SV, Basuroy T, Marathe HG, Lores P, de la Serna IL. BAF60A
mediates interactions between the microphthalmia-associated transcription factor and the
BRG1-containing SWI/SNF complex during melanocyte differentiation. J Cell Physiol.
2019;234(7):11780-91. Epub 2018/12/06. doi: 10.1002/jcp.27840. PubMed PMID:
30515787; PubMed Central PMCID: PMCPMC6426657.
50.   Keenen B, Qi H, Saladi SV, Yeung M, de la Serna IL. Heterogeneous SWI/SNF
chromatin remodeling complexes promote expression of microphthalmia-associated
transcription factor target genes in melanoma. Oncogene. 2010;29(1):81-92. Epub
2009/09/29. doi: onc2009304 [pii]
10.1038/onc.2009.304. PubMed PMID: 19784067; PubMed Central PMCID: PMC2803337.
51.   Butler A, Hoffman P, Smibert P, Papalexi E, Satija R. Integrating single-cell
transcriptomic data across different conditions, technologies, and species. Nat Biotechnol.
2018;36(5):411-20. Epub 20180402. doi: 10.1038/nbt.4096. PubMed PMID: 29608179;
PubMed Central PMCID: PMCPMC6700744.

52.   McInnes LH, John; Saul, Nathaniel; and Grossberger, Lukas. UMAP: Uniform
Manifold Approximation and Projection. The Journal of Open Source Software. 2018;3:861.
doi: doi.org/10.48550/arXiv.1802.03426.
53.   Farnsworth DR, Saunders LM, Miller AC. A single-cell transcriptome atlas for
zebrafish development. Dev Biol. 2020;459(2):100-8. Epub 20191127. doi:
10.1016/j.ydbio.2019.11.008. PubMed PMID: 31782996; PubMed Central PMCID:
PMCPMC7080588.
54.   Saunders LM, Mishra AK, Aman AJ, Lewis VM, Toomey MB, Packer JS, et al.
Thyroid hormone regulates distinct paths to maturation in pigment cell lineages. eLife.
2019;8. Epub 20190529. doi: 10.7554/eLife.45181. PubMed PMID: 31140974; PubMed
Central PMCID: PMCPMC6588384.
55.   Wagner DE, Weinreb C, Collins ZM, Briggs JA, Megason SG, Klein AM. Single-cell
mapping of gene expression landscapes and lineage in the zebrafish embryo. Science.
2018;360(6392):981-7. Epub 20180426. doi: 10.1126/science.aar4362. PubMed PMID:
29700229; PubMed Central PMCID: PMCPMC6083445.
56.   Brombin A, Simpson DJ, Travnickova J, Brunsdon H, Zeng Z, Lu Y, et al. Tfap2b
specifies an embryonic melanocyte stem cell that retains adult multifate potential. Cell Rep.
2022;38(2):110234. doi: 10.1016/j.celrep.2021.110234. PubMed PMID: 35021087; PubMed
Central PMCID: PMCPMC8764619.
57.   Howe DG, Bradford YM, Conlin T, Eagle AE, Fashena D, Frazer K, et al. ZFIN, the
Zebrafish Model Organism Database: increased support for mutants and transgenics. Nucleic
Acids Res. 2013;41(Database issue):D854-60. Epub 20121015. doi: 10.1093/nar/gks938.
PubMed PMID: 23074187; PubMed Central PMCID: PMCPMC3531097.
58.   Mead TJ, Yutzey KE. Notch pathway regulation of neural crest cell development in
vivo. Dev Dyn. 2012;241(2):376-89. Epub 20120103. doi: 10.1002/dvdy.23717. PubMed
PMID: 22275227; PubMed Central PMCID: PMCPMC3266628.
59.   McMenamin SK, Bain EJ, McCann AE, Patterson LB, Eom DS, Waller ZP, et al.
Thyroid hormone-dependent adult pigment cell lineage and pattern in zebrafish. Science.
2014;345(6202):1358-61. Epub 20140828. doi: 10.1126/science.1256251. PubMed PMID:
25170046; PubMed Central PMCID: PMCPMC4211621.
60.   Singh AP, Dinwiddie A, Mahalwar P, Schach U, Linker C, Irion U, et al. Pigment
Cell Progenitors in Zebrafish Remain Multipotent through Metamorphosis. Dev Cell.
2016;38(3):316-30. Epub 20160721. doi: 10.1016/j.devcel.2016.06.020. PubMed PMID:
27453500.
61.   Petratou K, Subkhankulova T, Lister JA, Rocco A, Schwetlick H, Kelsh RN. A
systems biology approach uncovers the core gene regulatory network governing iridophore
fate choice from the neural crest. PLoS genetics. 2018;14(10):e1007402. Epub 20181004.
doi: 10.1371/journal.pgen.1007402. PubMed PMID: 30286071; PubMed Central PMCID:
PMCPMC6191144.
62.   Bagnara JT, Matsumoto J, Ferris W, Frost SK, Turner WA, Jr., Tchen TT, et al.
Common origin of pigment cells. Science. 1979;203(4379):410-5. doi:
10.1126/science.760198. PubMed PMID: 760198.
63.   Knight RD, Javidan Y, Nelson S, Zhang T, Schilling T. Skeletal and pigment cell
defects in the lockjaw mutant reveal multiple roles for zebrafish tfap2a in neural crest
development. Dev Dyn. 2004;229(1):87-98. Epub 2003/12/31. doi: 10.1002/dvdy.10494.
PubMed PMID: 14699580.
64.   Barrallo-Gimeno A, Holzschuh J, Driever W, Knapik EW. Neural crest survival and
differentiation in zebrafish depends on mont blanc/tfap2a gene function. Development.
2004;131(7):1463-77. PubMed PMID: 14985255.

65. O'Brien EK, d'Alençon C, Bonde G, Li W, Schoenebeck J, Allende ML, et al.
Transcription factor Ap-2alpha is necessary for development of embryonic melanophores,
autonomic neurons and pharyngeal skeleton in zebrafish. Dev Biol. 2004;265(1):246-61. doi:
10.1016/j.ydbio.2003.09.029. PubMed PMID: 14697367.
66. Dooley CM, Wali N, Sealy IM, White RJ, Stemple DL, Collins JE, et al. The gene
regulatory basis of genetic compensation during neural crest induction. PLoS genetics.
2019;15(6):e1008213. Epub 20190614. doi: 10.1371/journal.pgen.1008213. PubMed PMID:
31199790; PubMed Central PMCID: PMCPMC6594659.
67. Creyghton MP, Cheng AW, Welstead GG, Kooistra T, Carey BW, Steine EJ, et al.
Histone H3K27ac separates active from poised enhancers and predicts developmental state.
Proc Natl Acad Sci U S A. 2010;107(50):21931-6. doi: 10.1073/pnas.1016071107. PubMed
PMID: 21106759; PubMed Central PMCID: PMCPMC3003124.
68. Pekowska A, Benoukraf T, Zacarias-Cabeza J, Belhocine M, Koch F, Holota H, et al.
H3K4 tri-methylation provides an epigenetic signature of active enhancers. EMBO J.
2011;30(20):4198-210. Epub 2011/08/19. doi: 10.1038/emboj.2011.295. PubMed PMID:
21847099; PubMed Central PMCID: PMCPMC3199384.
69. Ringrose L, Paro R. Epigenetic regulation of cellular memory by the Polycomb and
Trithorax group proteins. Annu Rev Genet. 2004;38:413-43. Epub 2004/12/01. doi:
10.1146/annurev.genet.38.072902.091907. PubMed PMID: 15568982.
70. Buenrostro JD, Giresi PG, Zaba LC, Chang HY, Greenleaf WJ. Transposition of
native chromatin for fast and sensitive epigenomic profiling of open chromatin, DNA-
binding proteins and nucleosome position. Nat Methods. 2013;10(12):1213-8. doi:
10.1038/nmeth.2688. PubMed PMID: 24097267; PubMed Central PMCID:
PMCPMC3959825.
71. Lee BK, Uprety N, Jang YJ, Tucker SK, Rhee C, LeBlanc L, et al. Fosl1
overexpression directly activates trophoblast-specific gene expression programs in embryonic
stem cells. Stem Cell Res. 2018;26:95-102. Epub 2017/12/23. doi: 10.1016/j.scr.2017.12.004.
PubMed PMID: 29272857; PubMed Central PMCID: PMCPMC5899959.
72. Liu H, Tan BC, Tseng KH, Chuang CP, Yeh CW, Chen KD, et al. Nucleophosmin
acts as a novel AP2alpha-binding transcriptional corepressor during cell differentiation.
EMBO Rep. 2007;8(4):394-400. Epub 2007/02/24. doi: 10.1038/sj.embor.7400909. PubMed
PMID: 17318229; PubMed Central PMCID: PMCPMC1852768.
73. Lin CY, Chao A, Wang TH, Lee LY, Yang LY, Tsai CL, et al. Nucleophosmin/B23 is
a negative regulator of estrogen receptor alpha expression via AP2gamma in endometrial
cancer cells. Oncotarget. 2016;7(37):60038-52. Epub 2016/08/17. doi:
10.18632/oncotarget.11048. PubMed PMID: 27527851; PubMed Central PMCID:
PMCPMC5312367.
74. Wong PP, Miranda F, Chan KV, Berlato C, Hurst HC, Scibetta AG. Histone
demethylase KDM5B collaborates with TFAP2C and Myc to repress the cell cycle inhibitor
p21(cip) (CDKN1A). Mol Cell Biol. 2012;32(9):1633-44. Epub 2012/03/01. doi:
10.1128/MCB.06373-11. PubMed PMID: 22371483; PubMed Central PMCID:
PMCPMC3347242.
75. Kim S, Yu NK, Kaang BK. CTCF as a multifunctional protein in genome regulation
and gene expression. Exp Mol Med. 2015;47:e166. Epub 2015/06/06. doi:
10.1038/emm.2015.33. PubMed PMID: 26045254; PubMed Central PMCID:
PMCPMC4491725.
76. Mavrothalassitis G, Ghysdael J. Proteins of the ETS family with transcriptional
repressor activity. Oncogene. 2000;19(55):6524-32. Epub 2001/02/15. doi:
10.1038/sj.onc.1204045. PubMed PMID: 11175368.

77.   Sekiya T, Zaret KS. Repression by Groucho/TLE/Grg proteins: genomic site
recruitment generates compacted chromatin in vitro and impairs activator binding in vivo.
Mol Cell. 2007;28(2):291-303. Epub 2007/10/30. doi: 10.1016/j.molcel.2007.10.002.
PubMed PMID: 17964267; PubMed Central PMCID: PMCPMC2083644.
78.   Watts JA, Zhang C, Klein-Szanto AJ, Kormish JD, Fu J, Zhang MQ, et al. Study of
FoxA pioneer factor at silent genes reveals Rfx-repressed enhancer at Cdx2 and a potential
indicator of esophageal adenocarcinoma development. PLoS genetics. 2011;7(9):e1002277.
Epub 2011/09/22. doi: 10.1371/journal.pgen.1002277. PubMed PMID: 21935353; PubMed
Central PMCID: PMCPMC3174211.
79.   Vierbuchen T, Ling E, Cowley CJ, Couch CH, Wang X, Harmin DA, et al. AP-1
Transcription Factors and the BAF Complex Mediate Signal-Dependent Enhancer Selection.
Mol Cell. 2017;68(6):1067-82.e12. Epub 2017/12/23. doi: 10.1016/j.molcel.2017.11.026.
PubMed PMID: 29272704; PubMed Central PMCID: PMCPMC5744881.
80.   Wilson BG, Wang X, Shen X, McKenna ES, Lemieux ME, Cho YJ, et al. Epigenetic
antagonism between polycomb and SWI/SNF complexes during oncogenic transformation.
Cancer Cell. 2010;18(4):316-28. Epub 2010/10/19. doi: 10.1016/j.ccr.2010.09.006. PubMed
PMID: 20951942; PubMed Central PMCID: PMCPMC2957473.
81.   Baxter LL, Watkins-Chow DE, Pavan WJ, Loftus SK. A curated gene list for
expanding the horizons of pigmentation biology. Pigment Cell Melanoma Res.
2019;32(3):348-58. doi: 10.1111/pcmr.12743. PubMed PMID: 30339321.
82.   Hoek KS, Eichhoff OM, Schlegel NC, Döbbeling U, Kobert N, Schaerer L, et al. In
vivo switching of human melanoma cells between proliferative and invasive states. Cancer
Res. 2008;68(3):650-6. Epub 2008/02/05. doi: 10.1158/0008-5472.Can-07-2491. PubMed
PMID: 18245463.
83.   Verfaillie A, Imrichova H, Atak ZK, Dewaele M, Rambow F, Hulselmans G, et al.
Decoding the regulatory landscape of melanoma reveals TEADS as regulators of the invasive
cell state. Nat Commun. 2015;6:6683. Epub 2015/04/14. doi: 10.1038/ncomms7683. PubMed
PMID: 25865119; PubMed Central PMCID: PMCPMC4403341.
84.   Hoek KS, Schlegel NC, Brafford P, Sucker A, Ugurel S, Kumar R, et al. Metastatic
potential of melanomas defined by specific gene expression profiles with no BRAF signature.
Pigment Cell Res. 2006;19(4):290-302. Epub 2006/07/11. doi: 10.1111/j.1600-
0749.2006.00322.x. PubMed PMID: 16827748.
85.   Tsoi J, Robert L, Paraiso K, Galvan C, Sheu KM, Lay J, et al. Multi-stage
Differentiation Defines Melanoma Subtypes with Differential Vulnerability to Drug-Induced
Iron-Dependent Oxidative Stress. Cancer Cell. 2018;33(5):890-904 e5. Epub 2018/04/17.
doi: 10.1016/j.ccell.2018.03.017. PubMed PMID: 29657129; PubMed Central PMCID:
PMCPMC5953834.
86.   Jonsson G, Busch C, Knappskog S, Geisler J, Miletic H, Ringner M, et al. Gene
expression profiling-based identification of molecular subtypes in stage IV melanomas with
different clinical outcome. Clin Cancer Res. 2010;16(13):3356-67. Epub 2010/05/13. doi:
10.1158/1078-0432.CCR-09-2509. PubMed PMID: 20460471.
87.   Tirosh I, Izar B, Prakadan SM, Wadsworth MH, 2nd, Treacy D, Trombetta JJ, et al.
Dissecting the multicellular ecosystem of metastatic melanoma by single-cell RNA-seq.
Science. 2016;352(6282):189-96. Epub 2016/04/29. doi: 10.1126/science.aad0501. PubMed
PMID: 27124452; PubMed Central PMCID: PMCPMC4944528.
88.   Rambow F, Rogiers A, Marin-Bejar O, Aibar S, Femel J, Dewaele M, et al. Toward
Minimal Residual Disease-Directed Therapy in Melanoma. Cell. 2018;174(4):843-55 e19.
Epub 2018/07/19. doi: 10.1016/j.cell.2018.06.025. PubMed PMID: 30017245.
89.   Yu G, Wang LG, Han Y, He QY. clusterProfiler: an R package for comparing
biological themes among gene clusters. Omics. 2012;16(5):284-7. Epub 2012/03/30. doi:

10.1089/omi.2011.0118. PubMed PMID: 22455463; PubMed Central PMCID:
PMCPMC3339379.
90.   Huang S, Jean D, Luca M, Tainsky MA, Bar-Eli M. Loss of AP-2 results in
downregulation of c-KIT and enhancement of melanoma tumorigenicity and metastasis.
EMBO J. 1998;17(15):4358-69. doi: 10.1093/emboj/17.15.4358. PubMed PMID: 9687504;
PubMed Central PMCID: PMC1170769.
91.   El-Brolosy MA, Kontarakis Z, Rossi A, Kuenne C, Günther S, Fukuda N, et al.
Genetic compensation triggered by mutant mRNA degradation. Nature. 2019;568(7751):193-
7. Epub 20190403. doi: 10.1038/s41586-019-1064-z. PubMed PMID: 30944477; PubMed
Central PMCID: PMCPMC6707827.
92.   Kumari P, Sturgeon M, Bonde G, Cornell RA. Generating Zebrafish RNA-Less
Mutant Alleles by Deleting Gene Promoters with CRISPR/Cas9. Methods Mol Biol.
2022;2403:91-106. doi: 10.1007/978-1-0716-1847-9_8. PubMed PMID: 34913119.
93.   White JR, Thompson DT, Koch KE, Kiriazov BS, Beck AC, van der Heide DM, et al.
AP-2alpha-mediated activation of E2F and EZH2 drives melanoma metastasis. Cancer Res.
2021. Epub 2021/07/03. doi: 10.1158/0008-5472.CAN-21-0772. PubMed PMID: 34210752.
94.   Mayran A, Khetchoumian K, Hariri F, Pastinen T, Gauthier Y, Balsalobre A, et al.
Pioneer factor Pax7 deploys a stable enhancer repertoire for specification of cell fate. Nat
Genet. 2018;50(2):259-69. Epub 2018/01/24. doi: 10.1038/s41588-017-0035-2. PubMed
PMID: 29358650.
95.   Braganca J, Eloranta JJ, Bamforth SD, Ibbitt JC, Hurst HC, Bhattacharya S. Physical
and functional interactions among AP-2 transcription factors, p300/CREB-binding protein,
and CITED2. The Journal of biological chemistry. 2003;278(18):16021-9. Epub 2003/02/15.
doi: 10.1074/jbc.M208144200. PubMed PMID: 12586840.
96.   Bejjani F, Evanno E, Zibara K, Piechaczyk M, Jariel-Encontre I. The AP-1
transcriptional complex: Local switch or remote command? Biochim Biophys Acta Rev
Cancer. 2019;1872(1):11-23. Epub 2019/04/30. doi: 10.1016/j.bbcan.2019.04.003. PubMed
PMID: 31034924.
97.   Swinstead EE, Miranda TB, Paakinaho V, Baek S, Goldstein I, Hawkins M, et al.
Steroid Receptors Reprogram FoxA1 Occupancy through Dynamic Chromatin Transitions.
Cell. 2016;165(3):593-605. Epub 2016/04/12. doi: 10.1016/j.cell.2016.02.067. PubMed
PMID: 27062924; PubMed Central PMCID: PMCPMC4842147.
98.   Petruk S, Cai J, Sussman R, Sun G, Kovermann SK, Mariani SA, et al. Delayed
Accumulation of H3K27me3 on Nascent DNA Is Essential for Recruitment of Transcription
Factors at Early Stages of Stem Cell Differentiation. Mol Cell. 2017;66(2):247-57 e5. Epub
2017/04/16. doi: 10.1016/j.molcel.2017.03.006. PubMed PMID: 28410996; PubMed Central
PMCID: PMCPMC5412717.
99.   Curran K, Raible DW, Lister JA. Foxd3 controls melanophore specification in the
zebrafish neural crest by regulation of Mitf. Dev Biol. 2009;332(2):408-17. Epub 20090613.
doi: 10.1016/j.ydbio.2009.06.010. PubMed PMID: 19527705; PubMed Central PMCID:
PMCPMC2716409.
100.   Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, et al. The single-cell
transcriptional landscape of mammalian organogenesis. Nature. 2019;566(7745):496-502.
Epub 20190220. doi: 10.1038/s41586-019-0969-x. PubMed PMID: 30787437; PubMed
Central PMCID: PMCPMC6434952.
101.   Sander JD, Maeder ML, Reyon D, Voytas DF, Joung JK, Dobbs D. ZiFiT (Zinc
Finger Targeter): an updated zinc finger engineering tool. Nucleic Acids Res. 2010;38(Web
Server issue):W462-8. Epub 2010/05/04. doi: 10.1093/nar/gkq319. PubMed PMID:
20435679; PubMed Central PMCID: PMCPMC2896148.

102. Sander JD, Dahlborg EJ, Goodwin MJ, Cade L, Zhang F, Cifuentes D, et al.
Selection-free zinc-finger-nuclease engineering by context-dependent assembly (CoDA). Nat
Methods. 2011;8(1):67-9. Epub 2010/12/15. doi: nmeth.1542 [pii]
10.1038/nmeth.1542. PubMed PMID: 21151135; PubMed Central PMCID: PMC3018472.
103. Miller JC, Holmes MC, Wang J, Guschin DY, Lee YL, Rupniewski I, et al. An
improved zinc-finger nuclease architecture for highly specific genome editing. Nat
Biotechnol. 2007;25(7):778-85. Epub 2007/07/03. doi: 10.1038/nbt1319. PubMed PMID:
17603475.
104. Leroy B, Girard L, Hollestelle A, Minna JD, Gazdar AF, Soussi T. Analysis of TP53
mutation status in human cancer cell lines: a reassessment. Hum Mutat. 2014;35(6):756-65.
Epub 2014/04/05. doi: 10.1002/humu.22556. PubMed PMID: 24700732; PubMed Central
PMCID: PMCPMC4451114.
105. Liu H, Duncan K, Helverson A, Kumari P, Mumm C, Xiao Y, et al. Analysis of
zebrafish periderm enhancers facilitates identification of a regulatory variant near human
KRT8/18. eLife. 2020;9. Epub 2020/02/08. doi: 10.7554/eLife.51325. PubMed PMID:
32031521; PubMed Central PMCID: PMCPMC7039683.
106. Buenrostro JD, Wu B, Chang HY, Greenleaf WJ. ATAC-seq: A Method for Assaying
Chromatin Accessibility Genome-Wide. Curr Protoc Mol Biol. 2015;109:21.9.1-.9.9. Epub
2015/01/07. doi: 10.1002/0471142727.mb2129s109. PubMed PMID: 25559105; PubMed
Central PMCID: PMCPMC4374986.
107. Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. Genome biology. 2009;10(3):R25.
Epub 2009/03/06. doi: 10.1186/gb-2009-10-3-r25. PubMed PMID: 19261174; PubMed
Central PMCID: PMCPMC2690996.
108. Langmead B, Salzberg SL. Fast gapped-read alignment with Bowtie 2. Nat Methods.
2012;9(4):357-9. Epub 2012/03/06. doi: 10.1038/nmeth.1923. PubMed PMID: 22388286;
PubMed Central PMCID: PMCPMC3322381.
109. Ramírez F, Ryan DP, Grüning B, Bhardwaj V, Kilpert F, Richter AS, et al.
deepTools2: a next generation web server for deep-sequencing data analysis. Nucleic Acids
Res. 2016;44(W1):W160-5. Epub 2016/04/16. doi: 10.1093/nar/gkw257. PubMed PMID:
27079975; PubMed Central PMCID: PMCPMC4987876.
110. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-
based analysis of ChIP-Seq (MACS). Genome biology. 2008;9(9):R137. Epub 2008/09/19.
doi: 10.1186/gb-2008-9-9-r137. PubMed PMID: 18798982; PubMed Central PMCID:
PMCPMC2592715.
111. Ross-Innes CS, Stark R, Teschendorff AE, Holmes KA, Ali HR, Dunning MJ, et al.
Differential oestrogen receptor binding is associated with clinical outcome in breast cancer.
Nature. 2012;481(7381):389-93. Epub 2012/01/06. doi: 10.1038/nature10730. PubMed
PMID: 22217937; PubMed Central PMCID: PMCPMC3272464.
112. McLean CY, Bristor D, Hiller M, Clarke SL, Schaar BT, Lowe CB, et al. GREAT
improves functional interpretation of cis-regulatory regions. Nat Biotechnol. 2010;28(5):495-
501. Epub 2010/05/04. doi: nbt.1630 [pii]
10.1038/nbt.1630. PubMed PMID: 20436461.
113. Skene PJ, Henikoff S. An efficient targeted nuclease strategy for high-resolution
mapping of DNA binding sites. eLife. 2017;6. Epub 2017/01/13. doi: 10.7554/eLife.21856.
PubMed PMID: 28079019; PubMed Central PMCID: PMCPMC5310842.
114. Meers MP, Bryson TD, Henikoff JG, Henikoff S. Improved CUT&RUN chromatin
profiling tools. eLife. 2019;8. Epub 2019/06/25. doi: 10.7554/eLife.46314. PubMed PMID:
31232687; PubMed Central PMCID: PMCPMC6598765.

115.   Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, et al. STAR:
ultrafast universal RNA-seq aligner. Bioinformatics. 2013;29(1):15-21. Epub 2012/10/30.
doi: 10.1093/bioinformatics/bts635. PubMed PMID: 23104886; PubMed Central PMCID:
PMCPMC3530905.
116.   Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion
for RNA-seq data with DESeq2. Genome biology. 2014;15(12):550. Epub 2014/12/18. doi:
10.1186/s13059-014-0550-8. PubMed PMID: 25516281; PubMed Central PMCID:
PMCPMC4302049.
117.   Mi H, Ebert D, Muruganujan A, Mills C, Albou LP, Mushayamaha T, et al.
PANTHER version 16: a revised family classification, tree-based classification tool,
enhancer regions and extensive API. Nucleic Acids Res. 2021;49(D1):D394-d403. Epub
2020/12/09. doi: 10.1093/nar/gkaa1106. PubMed PMID: 33290554; PubMed Central
PMCID: PMCPMC7778891.
118.   Baxter LL, Loftus SK, Pavan WJ. Networks and pathways in pigmentation, health,
and disease. Wiley interdisciplinary reviews Systems biology and medicine. 2009;1(3):359-
71. Epub 2010/02/18. doi: 10.1002/wsbm.20. PubMed PMID: 20161540; PubMed Central
PMCID: PMC2804986.
1845

1846

Fig. 2

Box 1: TFAP2A peaks at open and active chromatin

WT

Proliferation and pigmentation genes

H3K27Ac+ ATAC+

TFAP2-KO

TF (e.g., MITF)

H3K27Ac- ATAC-

TF (e.g., MITF)

H3K27Ac- ATAC+

H3K27Ac+ ATAC+

1A) TFAP2-pioneered-and-activated

1B) Non-pioneered TFAP2-activated

1C) TFAP2-independent

Box 2: TFAP2A peaks at closed and inactive chromatin

WT

TFAP2 R

Cell-cell adhesion genes

H3K27Ac- ATAC-

TFAP2-KO

TF (e.g., MITF) R

H3K27Ac+ ATAC+

H3K27Ac- ATAC-

2A) TFAP2 pioneered-and-inhibited

2B) TFAP2 independent

ATAC

TFAP2

Pioneer factor

Repressor

H3K27Ac

Transcription Factors

TFAP2-activated genes

TFAP2-inhibited genes

Fig. 3

A TFAP2-pioneered - and - activated enhancer

B Non-pioneered TFAP2 - activated enhancer

C TFAP2-pioneered - and - inhibited enhancer

D Non-pioneered TFAP2 - inhibited enhancer

PAX3 (+60 kb) *ENTPD6* (+26 kb) *ADAM19* (+21 kb) *FGF5* (+40 kb)

E TFAP2-pioneered - and - activated enhancers (2,002 peaks)

F Non-pioneered TFAP2-activated enhancers (1,836 peaks)

G TFAP2-pioneered – and - inhibited enhancers (864 peaks)

H Non-pioneered TFAP2-inhibited enhancers (440 peaks)

I * 3.7×10^{-27} ** 8.1×10^{-02}

J * 2.1×10^{-12} ** 2.8×10^{-01}

K * ns ** 9.4×10^{-05}

L * ns ** ns

* p-val. <u>enhancer</u> association with TFAP2-activated genes
** p-val. <u>enhancer</u> association with TFAP2-inhibited genes

M TFAP2-pioneered-and-activated enhancers

Motif	TF	Rank	P-value
	TFAP2	1,2	1×10^{-600}
	EBF	3	1×10^{-51}
	SOXE	4,6	1×10^{-36}
	RUNX	7	1×10^{-31}
	MITF	28	1×10^{-12}

N Non-pioneered TFAP2-activated enhancers

Motif	TF	Rank	P-value
	TFAP2	1	1×10^{-280}
	FRA1	2	1×10^{-46}
	RUNX	4	1×10^{-27}
	TEAD	6	1×10^{-24}
	ZFX	7	1×10^{-22}

O TFAP2-pioneered-and-inhibited enhancers

Motif	TF	Rank	P-value
	TFAP2	1,2	1×10^{-63}
	RUNX	3	1×10^{-37}
	CST6	4	1×10^{-21}
	ETS1	6	1×10^{-14}
	CTCF	12	1×10^{-12}

Fig. 4

A Effect of TFAP2 on MITF:

MITF — WT, *TFAP2*-KO
TFAP2A — WT, *MITF-KO*

Activated (4,726 peaks)
Mutual (717 peaks)
Inhibited (1,605 peaks)

B C D TFAP2-activated MITF peaks **E** TFAP2-inhibited MITF peaks

	B	C	D	E
TFAP2:	pioneer	pioneer*	activator	pioneer
MITF:	activator	activator*	activator	activator

FRMD4B (-100Kb), *TRPM1* (Promoter), *CYP7B1* (Intron 1), *SNX29* (Intron 19)

TFAP2A — WT: 250, 300, 250, 100
MITF-KO
MITF — WT: 250, 300, 250, 150
TFAP2-KO
ATAC — WT: 50, 50, 50, 50
TFAP2-KO
H3K27Ac — WT: 25, 25, 25, 25
TFAP2-KO

*Mutually dependent

F TFAP2-activated MITF peaks, pioneered
(3,083 peaks)
MITF — Log2 (Normalized reads) — WT, *TFAP2*-KO — *P < 0.0001*

G TFAP2-activated MITF peaks, pioneered, mutually dependent
(717 peaks)
MITF — WT, *TFAP2*-KO — *P < 0.0001*

H TFAP2-activated MITF peaks, non-pioneered
(2,022 peaks)
MITF — WT, *TFAP2*-KO — *P < 0.0001*

I TFAP2-inhibted MITF peaks, pioneered
(924 peaks)
MITF — WT, *TFAP2*-KO — *P < 0.0001*

F' ATAC — Log2 (Normalized reads) — WT, *TFAP2*-KO — *P < 0.0001*
G' ATAC — WT, *TFAP2*-KO — *P < 0.0001*
H' ATAC — WT, *TFAP2*-KO — *ns*
I' ATAC — WT, *TFAP2*-KO — *P < 0.0001*

Association with gene expression:
MITF-activated, EXP-21
TFAP2-activated, EXP-26

MITF-activated, EXP-09
TFAP2-activated, EXP-05

MITF-activated, EXP-11
TFAP2-activated, EXP-13

TFAP2-inhibited, EXP-04

F''
TFAP2: pioneer
MITF: activator
WT, *TFAP2*-KO, *MITF*-KO
B = BAF (SWI/SNF)

G''
TFAP2: pioneer*
MITF: activator*
WT, *TFAP2*-KO, *MITF*-KO
* Mutually dependent
H3K27Me3

H''
TFAP2: activator
MITF: activator
WT, *TFAP2*-KO, *MITF*-KO
P = Alternative Pioneer factor

I''
TFAP2: pioneer
MITF: activator
WT, *TFAP2*-KO, *MITF*-KO
R = Repressor

F'''

Motif	TF	Rank	P-value
	TFAP2	1,2	1 x 10^-641
	MITF	3	1 x 10^-134
	SOXE	4,5	1 x 10^-104
	RUNX	6-7, 9	1 x 10^-90
	FRA1	11	1 x 10^-69

G'''

Motif	TF	Rank	P-value
	MITF	1	1 x 10^-35
	TFAP2	2-3	1 x 10^-31
	BRM2	4, 7-9	1 x 10^-18
	TEAD4	6	1 x 10^-10

H'''

Motif	TF	Rank	P-value
	JUNB	1	1 x 10^-163
	TFAP2	2	1 x 10^-140
	SOXE	3	1 x 10^-125
	RUNX	4	1 x 10^-124
	CBF/TFE3	5	1 x 10^-117

I'''

Motif	TF	Rank	P-value
	SP1	1	1 x 10^-86
	NFY	2	1 x 10^-52
	JUN	3	1 x 10^-43
	TFE3	4	1 x 10^-41
	TFAP2	14	1 x 10^-10

Fig. 5

A

RNA-Sequencing (FDR < 0.05)

Genes directly activated by MITF — 527

180
112*

Genes directly activated by TFAP2 — 358

*Associated with a TFAP2-dependent MITF peak

B

Genes directly activated by MITF and by TFAP2 paralogs (Five most enriched GO terms)

GO biological process	Enrichment	No. of genes	FDR
Pigmentation	16.75	11	2.79×10^{-06}
Pigment differentiation	28.02	6	9.79×10^{-04}
Cell differentiation	44	77	8.7×10^{-03}
Cellular pigmentation	13.49	5	4.68×10^{-02}
Receptor signaling	3.51	15	3.78×10^{-02}

C

Pigmentation related genes directly activated by:

MITF	MITF and TFAP2	TFAP2
DKC1	RAB32	IGFBP7
MLANA	HPS3	KCTD15
EIF3B	CLCN7	ABCA12
KBTBD8	CTNNB1	CHD7
RAB27A	CTSD	EDNRB
STX3	DCT	EN1
AFG3L2	ETS1	HIPK2
BNC2	GPR143	HTRA1
CCT2	MAFB	MCM4
DTNBP1	MFSD12	MYO5A
IRF4	MITF	PARN
MLPH	MREG	PAX3
MYO10	NR4A3	RAD21
OSTM1	OCA2	SDC2
PARD3	PEPD	SLC2A11
PMEL	RAB38	SOX9
PPARGC1A	SH3BP4	SYTL2
PSEN2	SH3PXD2A	TFAP2C
RPS14	SNAI2	ALCAM
RPS7	TRPM1	BLM
SCUBE2	VDR	DISC1
SLC45A2		ERBB3
STXBP1		FANCC
SZT2		LYST
TYRP1		MIB1
USF2		NBN
VPS18		PMCH
		PRKDC
		SOX5

D

RNA-Sequencing (FDR < 0.05)

Genes directly inhibited by MITF — 725

Genes directly inhibited by TFAP2 — 249

E

Genes directly inhibited by MITF and by TFAP2 paralogs (Five most enriched GO terms)

GO biological process	Enrichment	No. of genes	FDR
Vasculature development	6.8	32	1.8×10^{-12}
Tube development	5.13	36	1.4×10^{-10}
Cell motility	4.81	34	2.2×10^{-08}
Cell migration	3.51	31	3.7×10^{-07}
Cell adhesion	3.5	20	7.0×10^{-06}

F

TFAP2-activated   TFAP2-inhibited

all direct direct* all direct

Verfaillie-proliferation
Jonsson-pigmentation
Hoek-proliferative
Tsoi-melanocytic
Rambow-pigmented
Triosh-ALX program
Riesenberg-TNF response in melanoma
Verfaillie-invation
Rambow-invasive cluster

Gene ratio
● 0.2
● 0.4
● 0.6

p-adjust
0.01 0.03

*Associated with TFAP2-activated MITF peak

Fig. 6