Title: Prevalence and correlates of phenazine resistance in culturable bacteria from a dryland wheat field

Running title: Phenazine resistance in wheat rhizosphere community

Authors: Elena K. Perry[1], Dianne K. Newman[1,2*]

Affiliations:

[1]Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA

91125, USA

[2]Division of Geological and Planetary Sciences, California Institute of Technology, Pasadena, CA 91125,

USA

Contact information:

E-mail: dkn@caltech.edu

Tel: 626-395-3543

Abstract

Phenazines are a class of bacterially-produced redox-active natural antibiotics that have

demonstrated potential as a sustainable alternative to traditional pesticides for the biocontrol of

fungal crop diseases. However, the prevalence of bacterial resistance to agriculturally-relevant

phenazines is poorly understood, limiting both the understanding of how these molecules might

shape rhizosphere bacterial communities and the ability to perform risk assessment for off-target

effects. Here, we describe profiles of susceptibility to the antifungal agent phenazine-1-carboxylic

acid (PCA) across more than 100 bacterial strains isolated from a wheat field where PCA producers

are indigenous and abundant. We find that Gram-positive bacteria are typically more sensitive to

PCA than Gram-negative bacteria, but that there is also significant variability in susceptibility both

within and across phyla. Phenazine-resistant strains are more likely to be isolated from the wheat

rhizosphere, where PCA producers are also more abundant, compared to bulk soil. Furthermore,

PCA toxicity is pH-dependent for most susceptible strains and broadly correlates with PCA

reduction rates, suggesting that uptake and redox-cycling are important determinants of phenazine

toxicity. Our results shed light on which classes of bacteria are most likely to be susceptible to

phenazine toxicity in acidic or neutral soils. In addition, the taxonomic and phenotypic diversity of

our strain collection represents a valuable resource for future studies on the role of natural antibiotics

in shaping wheat rhizosphere communities.

Importance

Microbial communities contribute to crop health in important ways. For example, phenazine

metabolites are a class of redox-active molecules made by diverse soil bacteria that underpin the

biocontrol of wheat and other crops. Their physiological functions are nuanced: in some contexts

they are toxic, in others, beneficial. While much is known about phenazine production and the effect

of phenazines on producing strains, our ability to predict how phenazines might shape the

composition of environmental microbial communities is poorly constrained; that phenazine

prevalence in the rhizosphere is predicted to increase in arid soils as the climate changes provides an

impetus for further study. As a step towards gaining a predictive understanding of phenazine-linked

microbial ecology, we document the effects of phenazines on diverse bacteria that were co-isolated

from a wheat rhizosphere and identify conditions and phenotypes that correlate with how a strain

will respond to phenazines.

Introduction

Diverse microorganisms produce natural antibiotics that can kill or inhibit the growth of other

microbes (Bérdy, 2012; Granato et al., 2019). Several such compounds have been commercialized

as antimicrobial drugs for the treatment of infections, beginning with penicillin in the 1940s

(Aminov, 2010; Hutchings et al., 2019). Unfortunately, the selective pressures exerted by the

widespread use of antibiotics in medicine and agriculture have led to worrisome increases in the

prevalence of antimicrobial resistance among human and animal pathogens over the past several

decades (Davies and Davies, 2010; Manyi-Loh et al., 2018). Yet while this repercussion of human

antibiotic use has been well documented, comparatively little is known about the ecological effects

of microbially-produced antibiotics in natural environments (Aminov, 2009; Sengupta *et al.*, 2013).

Recent studies have suggested that natural antibiotics may serve a variety of functions for their

producers beyond the suppression of competing microbes (Davies, 2006; Davies *et al.*, 2006),

including nutrient acquisition (Wang and Newman, 2008; McRose and Newman, 2021),

conservation of energy in the absence of oxygen (Glasser *et al.*, 2014; Glasser *et al.*, 2017), and cell-

cell signaling (Dietrich *et al.*, 2006; Linares *et al.*, 2006). At the same time, toxicity to one or more

microorganisms is by definition a feature of these molecules, but the extent to which this trait shapes

their influence on microbial communities is unclear (Demain and Fang, 2000; Davies and Ryan,

2012; Bérdy, 2012). In addition, for many if not most natural antibiotics, the determinants and

prevalence of susceptibility or resistance to their toxicity remain unknown or poorly characterized

(Handelsman and Stabb, 1996). These gaps in knowledge hinder our ability to understand and predict

the impacts of these metabolites on microbial communities of interest.

One environmental context in which natural antibiotics are thought to be particularly

abundant and ecologically relevant is the rhizosphere—the narrow plane of soil immediately

adjacent to plant roots (Haas and Défago, 2005; Mavrodi *et al.*, 2012; Tyc *et al.*, 2017). Natural

antibiotics such as phenazines and 2,4-diacetylphloroglucinol have been directly detected in the

rhizospheres of multiple crops, including wheat, potato, and sugar beet (Thomashow *et al.*, 1990;

Bergsma-Vlami *et al.*, 2005; Mavrodi *et al.*, 2012), and phenazines have been shown to increase the

fitness of their producers when competing with other microbes in the rhizosphere (Mazzola *et al.*,

1992; Yu *et al.*, 2018). Production of these molecules has also been demonstrated to underpin the

ability of certain bacteria to control fungal crop diseases (Thomashow and Weller, 1988;

Thomashow *et al.*, 1990; Haas and Keel, 2003; Mazurier *et al.*, 2009; Yu *et al.*, 2018), further

indicating that natural antibiotics can act as agents of microbe-microbe antagonism in the

rhizosphere. As a result of this activity, phenazine-producing *Pseudomonas* strains have received

attention as potential biocontrol agents that could serve as a more sustainable alternative to traditional

synthetic pesticides in agriculture (Handelsman and Stabb, 1996; Haas and Keel, 2003). However,

several challenges remain with respect to the practical application of these strains, including

inconsistent efficacy under field conditions (Haas and Keel, 2003; Haas and Défago, 2005), limited

understanding of the mechanisms and evolutionary dynamics of resistance to phenazines (Mazzola

et al., 1995; Handelsman and Stabb, 1996), and the possibility of off-target effects (Haas and Keel,

2003).

Importantly, with regard to the latter concern, phenazines are toxic not only to fungi, but also

to some bacteria (Baron and Rowe, 1981; Turner and Messenger, 1986; Costa *et al.*, 2015). Yet

while the utility of phenazine-producing pseudomonads for biocontrol of fungal crop diseases has

been extensively investigated, the potential impact of these strains on non-target bacterial residents

of the rhizosphere is less well understood. One study found that inoculation of *Pseudomonas*

biocontrol strains shifted the rhizosphere community of maize seedlings, pushing the ratio of Gram-

positive to Gram-negative bacteria in favor of the latter; however, this analysis was based on

profiling colony growth rates and whole-cell fatty acids from pooled cultured isolates, greatly

limiting the taxonomic resolution and making it difficult to rule out whether the Gram-negative

biocontrol strains might themselves have directly contributed to the shift (Kozdrój *et al.*, 2004). On

the other hand, at least three studies found no notable or consistent effects of introduced

Pseudomonas species on the rhizosphere bacterial communities of wheat or potato (Gagliardi *et al.*,

2001; Bankhead *et al.*, 2004; Roquigny *et al.*, 2018), albeit the studies in wheat employed methods

with limited discriminatory power (namely, carbon source utilization profiling and terminal

restriction fragment length polymorphism). Given these mixed results and the lack of fine-grained

spatial or taxonomic resolution in most studies on this topic, whether phenazine-producing bacteria

actively shape the surrounding rhizosphere bacterial community through antibiosis, potentially at the

micron or millimeter scale, remains an open question.

In addition to the lack of clarity regarding the effects of phenazines on rhizosphere bacterial

communities, the taxonomic and physiological correlates of phenazine resistance in bacteria remain

incompletely understood. The toxicity of phenazines is generally ascribed to the generation of

reactive oxygen species (ROS) and interference with respiration (Hassan and Fridovich, 1980; Baron

and Rowe, 1981; Voggu *et al.*, 2006; Perry and Newman, 2019). Previous studies have suggested

that efflux pump expression, cell permeability, oxidative stress responses, and the composition of

the respiratory electron transport chain can affect bacterial susceptibility to phenazines (Voggu *et*

al., 2006; Khare and Tavazoie, 2015; Noto *et al.*, 2017; Wolloscheck *et al.*, 2018; Perry and

Newman, 2019; Meirelles *et al.*, 2021). In addition, a comparison of 14 bacterial strains indicated

that Gram-negative bacteria as a group may be more resistant to phenazine toxicity than Gram-

positive bacteria (Baron and Rowe, 1981). However, all of these studies focused on a specific

phenazine, pyocyanin, that is particularly toxic (Meirelles and Newman, 2018) and best known for

its role as a virulence factor during infections of humans and animals (Lau *et al.*, 2004; Liu and

Nizet, 2009). Whether the same observations hold true for more agriculturally-relevant phenazines

such as phenazine-1-carboxylic acid (PCA) (Thomashow and Weller, 1988; Thomashow *et al.*,

1990; Mavrodi *et al.*, 2012; Dar *et al.*, 2020) is unknown.

In this study, we set out to lay a foundation for addressing unresolved questions about the

ecological impact of phenazine toxicity in the rhizosphere by determining the prevalence of

phenazine resistance among bacteria isolated from a wheat field in the Inland Pacific Northwest, a

region where phenazine production and the biocontrol potential of indigenous *Pseudomonas* species

have been studied for decades (Thomashow and Weller, 1988; Thomashow *et al.*, 1990; Mazzola *et*

al., 1992; Mavrodi *et al.*, 2012). We designed a culture-based assay to measure sensitivity to PCA,

which is the best-studied and one of the most abundant phenazines in this environment (Thomashow

et al., 1990; Mavrodi *et al.*, 2012; Dar *et al.*, 2020). We also performed full-length 16S rRNA gene

sequencing of our isolates in order to assess the relationship between taxonomy and PCA resistance.

Finally, to assess potential physiological correlates of PCA resistance, we measured PCA reduction

rates for a subset of phenotypically-diverse strains and investigated the effects of broad-spectrum

efflux pump inhibitors on growth in the presence of PCA.

Results

Taxonomic diversity of culturable bacteria from dryland wheat rhizospheres and bulk soil

A total of 166 strains of bacteria were isolated from 12 soil samples collected from a wheat

field at Washington State University's Lind Dryland Research Station in early August 2019, shortly

after the wheat harvest. The samples comprised 4 replicates each of wheat rhizosphere ("Wheat"),

bulk soil collected between planted rows ("Between"), and bulk soil from a virgin patch of

uncultivated soil adjacent to the field ("Virgin"). Full-length 16S rRNA gene sequencing revealed

that the isolates represented 24 genera across 4 phyla: Actinobacteria, Bacteroidetes, Firmicutes, and

Proteobacteria. The vast majority of isolates from the bulk soil samples (both Between and Virgin)

were Actinobacteria or Firmicutes. In Wheat samples, on the other hand, the combined proportions

of these two phyla were lower, accounting for 18-50% and 10-25% of isolates respectively, while

Proteobacteria accounted for approximately 25-60% of isolates depending on the replicate, and 1-3

strains of Bacteroidetes were also detected in each replicate (5-12% of isolates) (Fig. 1).

Taxonomic and spatial distribution of PCA resistance phenotypes

We screened our isolates for resistance to PCA at both circumneutral and acidic pH (7.3 and

5.1, respectively), because the toxicity of PCA to diverse organisms is known to vary depending on

pH (Brisbane *et al.*, 1987; Cezairliyan *et al.*, 2013), and because the bulk soil and rhizosphere pH of

wheat fields in the Inland Pacific Northwest can vary by >3 units (pH 4.3-8.0) depending on

geographic location and fertilizer treatment status (Smiley and Cook, 1973). The pH-dependency of

PCA toxicity has been attributed to the fact that the deprotonated form of PCA is negatively charged

(Fig. S1); the negative charge on bacterial cell walls and the outer membrane of Gram-negative

bacteria (or the negative membrane potential of eukaryotic cells) likely hinders uptake of this species.

The protonated form of PCA, on the other hand, is neutral and presumably can passively diffuse

across cell membranes given the small size and hydrophobic nature of the molecule (Price-Whelan

et al., 2006; Cezairliyan *et al.*, 2013). The pKa of PCA is 4.24 at 25 °C; thus, at pH 7, only 0.17%

of PCA in solution is protonated compared to 14.8% at pH 5, typically leading to greater toxicity at

the lower pH (Brisbane *et al.*, 1987). We chose 100 μM as the working concentration of PCA as this

is likely to be in a physiologically relevant range based on concentrations measured both in pure

cultures and in the field. In broth cultures of biocontrol strains of *Pseudomonas*, PCA accumulates

to concentrations ranging from dozens to hundreds of micromolar (Séveno *et al.*, 2001; Tagele *et al.*,

2019). In natural wheat rhizospheres, PCA has been detected at nanomolar concentrations (Mavrodi

et al., 2012), but these bulk measurements almost certainly underestimate local concentrations at the

micron scale given that bacteria colonize the rhizosphere in a patchy manner (Thomashow *et al.*,

1990). Notably, PCA can accumulate in biofilms to concentrations 360-fold greater on a per volume

basis compared to broth cultures (Séveno *et al.*, 2001), suggesting that local concentrations of PCA

in the rhizosphere, where biocontrol strains form robust biofilms (LeTourneau *et al.*, 2018), may be

orders of magnitude higher than the reported bulk values.

The screen was performed using 24-well plates and 0.1x tryptic soy agar (TSA) that was

either left unadjusted (pH 7.3-7.5) or adjusted to pH 5.1. Each isolate was spotted onto agar in

separate wells to prevent crosstalk and antagonism between the strains, and image analysis was used

to derive quantitative information about the growth of each strain over the course of 7 days (with

one timepoint per day). Because several strains among our 166 isolates appeared to be duplicates of

each other based on 16S sequence and colony morphology, we restricted the screen to 108 strains

that we judged likely to be unique (Table S1A); where multiple strains appeared identical, we chose

a representative strain. We also included 30 strains obtained from public culture collections that

represented species found among our isolates (Table S1B), in order to investigate the extent to which

PCA resistance phenotypes are consistent within species across different strains isolated from

different geographical locations. Accurate quantification of growth was not possible in this screen

for certain strains that formed transparent colonies, produced dark pigments, or had a tendency to

turn mucoid and spread (Fig. S2). Nevertheless, this assay enabled us to derive detailed profiles of

PCA sensitivity and resistance for the vast majority of our strains.

To compare PCA susceptibility across strains, we first focused on a single-timepoint

snapshot of each strain's phenotype taken at the equivalent of early stationary phase (i.e. around the

time that the spots on non-PCA control plates reached their maximal density). In accordance with

previous reports of the pH dependency of PCA toxicity, we found that more of our strains were

sensitive to PCA at pH 5.1 than at pH 7.3, and strains that were mildly inhibited by PCA at pH 7.3

were typically inhibited more strongly at pH 5.1 (Fig. 2A-B). At pH 5.1, strains of Actinobacteria

and Firmicutes were strongly inhibited by PCA, while most Proteobacteria were relatively resistant.

Members of Bacteroidetes exhibited variable responses, ranging from mild to severe growth

inhibition (Fig. 2A-B). At pH 7.3, there was considerably more phenotypic variation among the

Actinobacteria (Fig. 2A-B), particularly within the *Streptomyces* genus. In some cases, there was

even noticeable variation within the same putative species of *Streptomyces* (Fig. S3). In addition,

two strains of *Microbacterium* (W2I7 and W4I20), as well as one strain of *Arthrobacter* (W3I6),

grew slightly better in the presence of PCA at pH 7.3 compared to the control condition, perhaps

suggesting that they can use PCA as a carbon source or otherwise benefit from its presence at a pH

where toxicity is limited (Fig. 2B). On the other hand, in contrast to the phenotypic variation seen

among Actinobacteria, Firmicutes remained almost universally sensitive to PCA at pH 7.3 (albeit

somewhat less so than at pH 5.1), while all members of Bacteroidetes were resistant to PCA at pH

7.3, in contrast to their sensitivity to PCA at pH 5.1 (Fig. 2A-B). Proteobacteria remained generally

resistant (Fig. 2A-B), with the most notable exception being a strain of *Sphingomonas faeni*, isolate

W4I17, that was inhibited by PCA at both pH 7.3 and pH 5.1 (Fig. 2A-B).

Interestingly, while most strains yielded qualitatively consistent results across independent

biological replicates that were separated by months (Fig. S4-S7), a handful of strains displayed

variable growth and lag times at pH 5.1 either in the absence of PCA (mostly among Actinobacteria

and Firmicutes), or in the presence of PCA (a few strains among the Bacteroidetes). Growth at pH

7.3 was generally more consistent, with the exception being two strains of *Neobacillus niacini*

(V4I10 and V4I25) that initially failed to grow in the presence of PCA, but grew with minimal to no

inhibition in subsequent replicates. The reasons for these discrepancies are unclear, but may be

related to variations in how long the strains had been in stationary phase prior to inoculating the

experimental cultures, which was difficult to control precisely due to differing growth rates across

the large number of strains. The trace nutrient content may also have varied between different

batches of media, as the first replicate was performed using a different lot of tryptic soy broth

compared to subsequent replicates. Thus, the growth and PCA sensitivity of some strains may be

influenced by environmental factors beyond pH that remain to be elucidated.

We next examined whether there was any evidence of correlation between PCA resistance

phenotypes and the type of soil each strain was isolated from (Between, Virgin, or Wheat). A

previous study based on samples taken from the same wheat field found that the relative abundance

of phenazine producers was higher in the wheat rhizosphere compared to adjacent bulk soil (Dar *et*

al., 2020), a finding that we also confirmed using shotgun metagenomic sequencing of our soil

samples (Fig. 2C). We therefore hypothesized that if PCA-mediated antibiosis had shaped the

bacterial community composition of this field, the prevalence of PCA resistance would be higher

among isolates from the rhizosphere samples. Indeed, the Wheat samples clearly had the highest

proportion of PCA-resistant isolates, regardless of whether resistance was assessed at pH 5.1 (Fig.

2D) or pH 7.3 (Fig. 2E). By contrast, all strains from Between or Virgin samples were highly

sensitive to PCA at pH 5.1, though at pH 7.3, the resistance phenotypes of isolates from either type

of bulk soil spanned the full range from highly sensitive to completely resistant. However, these

findings are conflated with the fact that all of our Bacteroidetes strains, and all but one strain of

Proteobacteria, were exclusively isolated from the Wheat samples, and that the members of these

two phyla were generally relatively resistant to PCA, especially at pH 7.3. Further experiments will

be necessary in order to distinguish whether the enrichment of PCA-resistant phenotypes in the
rhizosphere samples is a consequence of PCA-mediated antibiosis or merely an indirect reflection
of other factors that favor the growth of Proteobacteria and Bacteroidetes in the rhizosphere.

Finally, visualizing the growth of each strain over the full 7-day time course revealed more
nuanced variations among the PCA resistance phenotypes at both pH 5.1 and pH 7.3 (Figs. 3 and
S4-S7). For example, while some strains of Bacteroidetes displayed increased lag and/or slower
growth rates in the presence of PCA at pH 5.1, the growth on PCA often eventually caught up to the
PCA-free controls. At pH 7.3, the effect of PCA on *Peribacillus* species was a combination of
increased lag and lower final cell density, but apparently not decreased maximal growth rate, as
compared to no growth at all on PCA at pH 5.1. This was distinct from the slower growth rates seen
for certain strains of *Bacillus*, *Neobacillus*, and *Paenibacillus* at pH 7.3. A few strains of the latter
three genera appeared to be still unable to grow on PCA at pH 7.3 based on the values derived from
image analysis (Fig. S5). However, examination of the plate images by eye revealed that this
sometimes reflected a limitation in the sensitivity of our imaging and quantification methods to low
levels of growth, rather than a true lack of growth (Fig. S2).

PCA reduction rates in relation to PCA susceptibility

We next sought to determine whether there are physiological correlates of PCA susceptibility
or resistance across taxonomically-diverse bacteria. Given that phenazines are redox-active
molecules whose toxicity is thought to be related to ROS generation as a consequence of redox
cycling in the cell (Hassan and Fridovich, 1980; Mavrodi *et al.*, 2006), we hypothesized that
susceptibility to PCA would be correlated with higher redox-cycling rates. PCA oxidation occurs
rapidly and abiotically in the presence of oxygen (Wang and Newman, 2008), suggesting that the
rate of PCA reduction would be the primary driver of differences in redox-cycling rates across strains
under oxic conditions. Therefore, to test our hypothesis, we measured the rate of PCA reduction
under anoxia as a proxy for the true redox-cycling rate, using a subset of strains that covered a range
of taxonomic groups and PCA susceptibility phenotypes.

For each tested strain, the PCA reduction rate was typically higher at pH 5.1 than at pH 7.3
(Fig. 4A), matching the expectation that PCA uptake would tend to be faster at a lower pH and the

observation that most susceptible strains were more sensitive to PCA at pH 5.1. The exceptions were:

1) strains belonging to the Firmicutes, which had lower reduction rates at pH 5.1 in our assay despite

being more susceptible to PCA under this condition, and 2) *Sphingomonas faeni* W4I17, which was

in fact more susceptible to PCA at pH 7.3 than at pH 5.1 (Fig. 2A-B), matching the pH dependency

of its rate of PCA reduction. Notably, some of the tested Firmicutes often struggled to grow at pH

5.1 even in the absence of PCA (Fig. S5). This observation might account for the inhibitory effect

of acidic pH on PCA reduction in these strains, given that a functional metabolism would be required

for the generation of cellular reductants, such as NADH, that can indirectly or directly reduce PCA.

For the strains of Firmicutes that were able to grow at acidic pH (namely B1I1 and B1I6), the reason

for their lower PCA reduction rate at pH 5.1 remains unclear.

We also examined whether there was any correlation between PCA reduction rates and

resistance levels across different strains at each pH. To do so, we plotted each strain's PCA reduction

rate against the early stationary phase snapshot ratio of growth on PCA versus no PCA (Fig. 4B).

We also calculated Spearman's correlation coefficient (r_s). Interestingly, there was a statistically

significant negative correlation between reduction rate and resistance at pH 7.3 (r_s = -0.70, $p <$

0.001), but not at pH 5.1 (r_s = -0.16, p = 0.4485). However, there were exceptions to the trend even

at pH 7.3; for example, while PCA severely and consistently inhibited the growth of the four fastest

PCA-reducing strains, the fifth-fastest strain, W2I1 (*Pantoea agglomerans*), appeared to be

completely resistant to PCA in one biological replicate, though its growth was mildly inhibited by

PCA in another replicate (Fig. S7). In addition, strains with reduction rates in the middle range of

0.2-0.4 nmol/OD_{600}/min spanned the full range of resistance phenotypes, from a growth ratio of 0.12

(*Peribacillus simplex* B2I2) to 0.96 (*Phyllobacterium ifrigiyense* W4I11). Taken together, these data

suggest that a low reduction rate may be helpful in mitigating PCA toxicity but is neither necessary

nor sufficient for resistance.

Effects of efflux pump inhibitors on PCA susceptibility

Given that PCA reduction rate alone was insufficient to account for the range of sensitivities

to PCA, we asked if efflux pumps might serve as an important contributor to PCA resistance in some

strains. To address this question, we used two efflux pump inhibitors (EPIs) in combination: 1)

reserpine, which is thought to target efflux pumps in the major facilitator superfamily (MFS) (Kumar

et al., 2013), and 2) phenylalanine-arginine β-naphthylamide (PAβN), which is thought to target

resistance-nodulation-division (RND) efflux pumps (Jamshidi *et al.*, 2017), although it may also

permeabilize the outer membrane of Gram-negative bacteria (Lamers *et al.*, 2013; Schuster *et al.*,

2019). We tested these EPIs on a subset of the strains that were used in the PCA reduction assay; we

excluded the *Bacillus* strains due to their tendency to settle on the bottom of the wells during growth

in the plate reader, which confounded the OD_{600} readings. Preliminary experiments revealed that

different strains were differentially sensitive to the EPIs themselves, with growth inhibition evident

even in PCA-free controls. We therefore determined for each strain the maximal EPI concentration

that, when the two EPIs were used separately, did not markedly reduce growth of PCA-free controls

(up to a maximum dose of 20 μg/mL for reserpine and 50 μg/mL for PAβN) (Table S2).

Interestingly, for many strains, the combination of both EPIs remained highly toxic even after this

optimization step, particularly at pH 7.3, possibly indicating synergistic toxicity or redundant roles

of different efflux pumps in preventing intracellular accumulation of toxic metabolic intermediates

(Fig. S9); the toxicity appeared to be pH-dependent, suggesting that the protonation state of the EPIs

might influence their efficacy. In these cases, it was not possible to assess the impact of efflux pump

inhibition on PCA sensitivity. However, for the remaining strains, three types of responses to the

EPIs emerged. For one group of strains, treatment with both EPIs in combination did not affect their

resistance to PCA (Fig. 5A-B); however, we cannot distinguish whether this is because efflux does

not contribute to PCA resistance in these strains, or because the tested EPIs did not effectively target

the relevant efflux pumps. For the second group of strains, the EPIs enhanced the toxicity of PCA,

as would be expected if efflux is an important mechanism of resistance (Fig. 5C). Finally, for the

third group of strains, the EPIs counterintuitively appeared to mitigate the toxicity of PCA, or vice

versa (Fig. 5D). Although unexpected, the latter phenomenon indirectly suggests that PCA interacts

with efflux systems in these strains. For example, treatment with either PCA or the EPIs might

upregulate the expression of certain efflux pumps, leading to decreased toxicity relative to treatment

with either type of toxin alone. Alternatively, given that some pumps in the MFS family are thought

to be bidirectional (Pao *et al.*, 1998), it is possible that MFS pumps promote uptake rather than export

of PCA in certain strains.

Discussion

In this study, we have characterized profiles of resistance to an agriculturally-relevant
phenazine across taxonomically diverse bacteria isolated from a wheat field where phenazine
producers are indigenous. We also examined potential physiological correlates of phenazine
resistance in a subset of these strains. Our findings establish a basis for inferring whether intrinsic
resistance is a factor that affects how phenazine production shapes bacterial communities in the
rhizosphere. In particular, it will now be possible to test data-driven predictions regarding which
strains, species, or even phyla are most likely to be affected by phenazine-mediated antibiosis, even
though the mechanistic underpinnings of phenazine susceptibility or resistance are likely
multifactorial and appear to differ across species.

One of our more notable findings is that the previously reported pH dependency of PCA
toxicity varies across bacterial taxonomic groups. For most Actinobacteria and Firmicutes,
sensitivity to PCA was clearly higher at pH 5.1 than at pH 7.3, indicating greater toxicity at the lower
pH as expected according to the pKa of PCA. On the other hand, nearly all tested Proteobacteria
were completely resistant to 100 µM PCA regardless of pH, at least down to pH 5.1. Consequently,
at pH 5.1, PCA resistance phenotypes largely correlated with phylum—and more broadly, a Gram-
positive versus Gram-negative divide, which has previously been reported for the toxicity of another
phenazine, pyocyanin (Baron and Rowe, 1981). The Gram-positive versus Gram-negative divide at
pH 5.1 is perhaps not surprising, as at this pH, a significant proportion (~14%) of PCA in solution is
protonated and therefore would not be repelled by the negatively charged cell wall. Under this
condition, the outer membrane of Gram-negative species presumably presents an additional barrier
to the entry of PCA, helping to limit the intracellular accumulation of the toxin in the same manner
as for numerous other antibiotics (Nikaido, 1989). Less expected, however, was the within-phylum
and even within-species variation in PCA resistance phenotypes at pH 7.3 among certain taxonomic
groups. Interestingly, several *Streptomyces* strains exhibited at least some PCA-dependent growth
inhibition at pH 7.3, even though many *Streptomyces* species are capable of producing their own
toxic phenazines (Turner and Messenger, 1986; Dar *et al.*, 2020). Given that the major limit on PCA
toxicity at circumneutral pH is thought to be its ability to enter cells, the phenotypic variability at pH
7.3 may indicate the presence of transporters or channels capable of phenazine uptake in some PCA-
sensitive Actinobacteria. Alternatively, it is possible that some PCA-sensitive strains locally
acidified the growth medium, which was only weakly buffered (1.4 mM K_2HPO_4).

348   Importantly, despite the general Gram-positive versus Gram-negative divide in sensitivity to
349 PCA at pH 5.1, possessing an outer membrane is evidently not a leakproof shield against PCA
350 toxicity. Proteobacteria as a group were more resistant to PCA at pH 5.1 compared to strains of
351 Bacteroidetes, most of which exhibited increased lag in the presence of PCA, even though both phyla
352 comprise Gram-negative bacteria. Intriguingly, one major difference between these clades is that
353 Proteobacteria generally utilize ubiquinone as an electron carrier during aerobic growth (Collins and
354 Jones, 1981), while the Bacteroidetes genera screened in this study (*Chitinophaga*,
355 *Chryseobacterium*, and *Pedobacter*) utilize menaquinone (Lin *et al.*, 2015; Singh *et al.*, 2017; Kong
356 *et al.*, 2019). Menaquinones have a lower reduction potential compared to ubiquinones (White *et al.*,
357 2012). Although the standard reduction potential of menaquinone is still higher than that of PCA (-
358 74 mV compared to -177 mV) (Price-Whelan *et al.*, 2006; White *et al.*, 2012), indicating that PCA
359 likely is not reduced by menaquinol, this difference with ubiquinone nevertheless raises the
360 possibility that PCA may interact differently, and perhaps more readily, with the aerobic respiratory
361 electron transport chain of menaquinone-utilizing Bacteroidetes strains compared to Proteobacteria,
362 thereby generating more ROS and/or interfering with the generation of ATP. Interestingly, another
363 study has shown that the reduced form of different phenazine with a low reduction potential, neutral
364 red (3-amino-7-dimethylamino-2-methylphenazine), can directly transfer electrons to menaquinone,
365 bypassing the proton-pumping NADH dehydrogenase complex that normally transfers electrons
366 from NADH to menaquinone and thereby "short-circuiting" the electron transport chain. We
367 hypothesize that a similar phenomenon may occur with PCA in strains that rely on menaquinone. In
368 future studies, this hypothesis could be tested by 1) performing *in vitro* experiments with purified
369 menaquinone, ubiquinone, and reduced PCA to directly test whether quinones oxidize the latter and
370 if so, whether the kinetics differ between menaquinone and ubiquinone, 2) measuring ROS
371 production and steady-state ATP pools in selected strains of Bacteroidetes and Proteobacteria both
372 in the presence and absence of PCA, and 3) forcing species of Proteobacteria (such as *Escherichia*
373 *coli*) that utilize both ubiquinone and menaquinone in different branches of their electron transport
374 chains to rely only on the latter (for example, by deleting the genes for ubiquinone biosynthesis),
375 followed by reassessing their sensitivity to PCA to see if there is any effect.

376   Beyond the specific strains screened in this study, the risk assessment of phenazine-
377 producing biocontrol strains, and the understanding of phenazine biology in general, would benefit

greatly from the development of a platform for the prediction of phenazine resistance phenotypes
from genomic or phylogenetic information. The results of this study already indicate that, depending
on the environmental pH, phylogenetic information may be of limited utility for prediction of PCA
resistance, given the variation in phenotypes for Actinobacteria at circumneutral pH. However, this
variability may also hold the key to identifying genome-based predictive markers of phenazine
resistance. Given the taxonomic and phenotypic diversity of our strain collection, whole-genome
sequencing of our isolates, followed by comparative genomics, could potentially reveal such
markers.

In summary, this work has laid the groundwork for rectifying a major gap in studies of how
introduction of a phenazine-producing biocontrol strain affects rhizosphere bacterial communities.
Previous studies have lacked information about the baseline prevalence of resistance in the native
communities (Gagliardi *et al.*, 2001; Bankhead *et al.*, 2004; Kozdrój *et al.*, 2004; Roquigny *et al.*,
2018), and in the absence of such information, it is impossible to determine whether a negative result
(lack of change in the rhizosphere community) reflects a high prevalence of resistance to PCA that
is particular to the studied community, versus a general lack of toxicity of PCA to most bacteria or
fundamental abiotic constraints on the antibacterial activity of PCA in the rhizosphere (e.g. limited
diffusion, adsorption to soil particles, etc.). Distinguishing between these scenarios is key to
assessing the risk of unwanted side effects in rhizosphere communities upon the application of
phenazine-producing biocontrol strains. In addition, recent studies have demonstrated that
phenazines produced by the opportunistic pathogen *Pseudomonas aeruginosa* can promote bacterial
tolerance and resistance to clinical antibiotics (Schiessl *et al.*, 2019; Zhu *et al.*, 2019; Meirelles *et
al.*, 2021; VanDrisse *et al.*, 2021), and that these effects can extend to other opportunistic pathogens
that are resistant to phenazines (Meirelles *et al.*, 2021). Thus, understanding the prevalence and
genetic determinants of resistance to phenazines may have implications not only for agriculture but
also for human medicine and beyond, as we continue to uncover new ecological roles for these
multifaceted bacterial metabolites.

Methods

Isolation of bacteria from wheat rhizosphere and bulk soil samples

Three types of samples were collected from a non-irrigated wheat field at Washington State

University's Lind Dryland Research Station on August 9, 2019: wheat plants and surrounding soil,

bulk soil from in between the planted rows, and bulk soil from a "virgin" hillside site that has never

been farmed. The wheat had been harvested a few weeks prior to sample collection. All samples

were immediately stored on ice in clean plastic bags, and subsequently at 4 °C for four days until

processing. Rhizosphere soil samples were obtained by shaking the wheat roots until only 1-2 mm

of tightly-adhering soil remained, followed by excising the roots at the crown with a sterile razor

blade. The roots of 2-3 plants per replicate were placed in 50 mL conical tubes with 30 mL of sterile

deionized water, vortexed at top speed for 1 min, and treated in an ultrasonic water bath for 1 min to

dislodge bacteria from the roots. Bulk soil samples (1 g per sample) were processed in the same

manner. Large soil particles were allowed to settle to the bottom of the tubes on the bench top, and

100 µL each of a 10-fold dilution series of the supernatants was spread onto 0.1x TSA plates

containing 50 µg/mL nystatin to inhibit fungal growth. The plates were incubated upside down at

room temperature in the dark and monitored for the appearance of new colonies over the course of

a week. Colonies that appeared morphologically distinct in each sample were picked and restreaked

on 0.1x TSA until visually pure cultures were obtained. Multiple representatives were also picked

for the most common colony types in an attempt to account for strain variations that might not be

apparent to the eye. Once the streaks yielded uniform single colonies, the isolates were inoculated

into 1.5 mL of 0.1x tryptic soy broth (TSB) in 5 mL polycarbonate culture tubes and incubated at 30

°C with shaking at 250 rpm. After 1-3 days of incubation, depending on when the cultures became

turbid, 0.5 mL of each culture was mixed with 0.5 mL of 50% glycerol and stored at -80 °C. Some

cultures never became turbid under these growth conditions, but nevertheless yielded viable frozen

stocks.

Species identification by 16S rRNA gene sequencing

Single colonies or patches of morphologically pure streaks were picked and resuspended in

20 µL sterile nuclease-free water. Colony PCR was performed using GoTaq Green Master Mix

(Promega) in 50 µL reactions (1 µL of cell suspension) according to the manufacturer's instructions.

For putative streptomycetes (isolates that formed hard colonies rooted in agar, often with aerial

hyphae), the thermocycling protocol was modified to include a 10 min initial heating step at 95 °C

(compared to 2 minutes for other samples). The primers used were 27F

(AGAGTTTGATCMTGGCTCAG) and 1492R (TACGGYTACCTTGTTACGACTT) (Lane,

1991). The PCR products were run on a 1% agarose gel to verify the presence of a single band at the

expected size (~1500 bp), followed by purification with the Monarch PCR and DNA Cleanup Kit

(New England Biolabs). The purified products were submitted for Sanger sequencing at Laragen,

Inc., using the same 27F and 1492R primers. The resulting forward and reverse sequences were

aligned using MAFFT (https://mafft.cbrc.jp/alignment/server/) and subjected to BLAST against the

NCBI 16S ribosomal RNA sequences database. For a few strains, full-length sequences could only

be obtained following PCR on genomic DNA extracted using the DNeasy Blood & Tissue kit

(Qiagen), following the manufacturer's instructions for Gram-positive bacteria. In addition, for two

strains (B1I5 and B2I8), the forward sequence consistently appeared to contain multiple products.

We presumed that this was due to multiple primer binding sites or another sequencing artifact rather

than mixed cultures as the corresponding sequences from the other direction were clean. In these

cases, only the clean sequence was submitted to BLAST.

PCA resistance screen

The optimized version of the PCA resistance screen was performed with four conditions:

0.1x TSA (15 g/L agar plus 3 g/L tryptic soy broth no. 2 from MilliporeSigma) with or without 100

μM PCA at pH 7.3 or pH 5.1 (adjusted with HCl). PCA was purchased from Princeton BioMolecular

Research and dissolved in filter-sterilized 14 mM NaOH to make 10 mM stock solutions. The PCA

stock solution or solvent control (14 mM NaOH) was added at 1% v/v to autoclaved molten 0.1x

TSA; we verified that this addition did not noticeably alter the pH of the medium by pipetting

aliquots onto pH test strips. Subsequently, 1 mL of the medium was pipetted into each well in 24-

well polystyrene Cellstar Cell Culture plates (Greiner Bio-One). The plates were allowed to set and

dry with the lids off in a biological safety cabinet for 20-30 min, followed by storage upside with the

lids on at room temperature in the dark for two days prior to use.

Cell suspensions for inoculation in the screen were prepared in one of two ways. First,

individual strains were inoculated into 5 mL TSB cultures in glass culture tubes and incubated at 25

°C with shaking at 250 rpm. Strains that grew overnight were then diluted to an OD_{600} of 0.05.

Certain strains did not grow well in this condition—in particular, strains of *Agromyces*,

Streptomyces, *Paenibacillus*, and *Neobacillus*. For these, some replicates were prepared by directly

scraping cells from streaks grown on 0.1x TSA plates and resuspended the cells in 200 µL TSB with

pipetting and brief vortexing at top speed. The OD_{600} of these cell suspensions was adjusted to 0.05,

except for *Streptomyces* due to their filamentous nature and tendency to clump; instead, visible cell

clumps were allowed to settle to the bottom and inocula were taken from the visually clear upper

portion. Subsequently, 10 µL of each cell suspension was pipetted onto the agar in a single well in

the 24-well plates. Three adjacent wells per condition were inoculated with each cell suspension,

representing technical replicates. After the spots dried, the plates were incubated at room temperature

upside in the dark for up to 7 days. Every 24 hrs, the plates were imaged in color at 600 dpi with an

Epson Perfection V550 Photo flatbed scanner. For strains that were prepared both from liquid

cultures and by resuspension from plates, we did not observe any consistent effect on the resulting

PCA susceptibility phenotypes.

Image analysis and quantification of growth

Images from the scanner were analyzed using Fiji (Schindelin *et al.*, 2012). Circular regions

of interest (ROIs) were drawn around each culture spot and the mean gray value was measured for

each ROI. We also measured the mean gray values of equivalent ROIs in the wells of blank,

uninoculated plates for each condition. The latter values were averaged across each 24-well blank

plate to give the "background" gray value, which was then subtracted from the mean gray values of

the culture spots. The resulting numbers were reported as the metric for growth. Importantly, while

this method generally worked well for comparisons across conditions within each strain, there are a

few caveats. First, this metric underestimated growth for strains that produced a dark pigment.

Second, growth was difficult to quantify for a few strains that grew as nearly transparent colonies.

Finally, this metric is not very sensitive to low levels of growth. Nevertheless, for the vast majority

of strains, this approach captured visible differences in growth across the four conditions in the

screen.

Metagenomics-based estimation of the relative abundance of PCA producers

DNA was extracted from 250 mg of each soil sample using the DNeasy PowerSoil kit

(Qiagen). Illumina metagenomic DNA libraries were prepared and sequenced (150 bp single-end

reads) at the Millard and Muriel Jacobs Genetics and Genomics Laboratory at the California Institute

of Technology. The resulting sequence data was processed according to the pipeline described by

Dar et al. (2020).

PCA reduction assay

Selected strains were grown in triplicate overnight in 5 mL TSB cultures at 25 °C with

shaking at 250 rpm. The cells were then concentrated by serially centrifuging 1 mL aliquots at 9000

g for 2 min in microcentrifuge tubes, washed with 0.1x TSB (pH 7.3), and standardized to OD_{600} =

8 in 0.1x TSB (pH 7.3). Subsequently, 100 µL of each cell suspension was transferred into a 96-well

plate, and the plate was moved into a Coy anaerobic chamber (5% hydrogen/95% nitrogen

headspace). Using plasticware and media that had been passively degassed in the chamber for at

least 3 days, electrochemically-reduced PCA was added to a final concentration of 100 µM in 190

µL of 0.1x TSB (pH 5.1 or pH 7.3) per well in a black clear-bottom 96-well plate; a standard curve

was also prepared using reduced PCA at 1 µM, 10 µM, 50 µM, and 100 µM. Cells were added to

each well (except the standard curve) to an OD_{600} of ~0.4, and the plate was immediately transferred

to a Biotek Synergy 5 plate reader maintained at 30 °C. OD_{600} and fluorescence readings (360 nm

ex/528 nm em) were taken every 5 minutes, with continuous shaking on the "medium" setting.

Reduction rates were derived from the slope of a linear regression fit to the visually linear range of

the data using the function lm in R (see Fig. S8 for an example). The rates were converted to

nmol/OD/min by relating the fluorescence units to the standard curve of reduced PCA concentrations

and then dividing by the initial OD_{600} reading for each replicate. No appreciable change in OD_{600}

was observed over the course of the experiment for any of the tested strains.

Efflux pump inhibition growth curves

Reserpine was purchased from Sigma-Aldrich and dissolved in dimethyl sulfoxide (DMSO)

to make a 20 mg/mL stock solution. PAβN was purchased from MedChemExpress and dissolved

in distilled water to make a 50 mg/mL stock solution. Selected strains were grown in triplicate

overnight in 5 mL TSB cultures at 25 °C with shaking at 250 rpm, washed once with 0.1x TSB (pH

7.3), and then inoculated to an initial OD_{600} of 0.05 in 160 µL of 0.1x TSB (pH 5.1 or pH 7.3) in a

96-well plate. Each well contained either a combination of reserpine and PAβN (see Table S2 for

the concentrations used for each strain) or a solvent control (DMSO at 0.1% final concentration),

along with either 0 or 100 µM PCA. The wells were topped with 50 µL of autoclaved mineral oil to

prevent evaporation, and growth was monitored with OD_{600} readings every 15 minutes in a Biotek

Synergy 5 plate reader set to 30 °C with continuous shaking on the "medium" setting.

Data availability

The metagenomic sequencing data generated in this study were deposited in the Sequence Read

Archive (SRA) under accession PRJNA521160.

Acknowledgements

We thank Linda Thomashow and David Weller of the USDA Agricultural Research

Service for providing the wheat rhizosphere and soil samples used in this study, and the members

of the Newman lab for helpful feedback throughout the process of designing and analyzing the

results of the PCA resistance screen. We also thank Zevin Condiotte for assistance with the

initial rounds of testing the design of the PCA resistance screen and Richard Horak for assistance

with extracting genomic DNA. This material is based upon work supported by the National

Science Foundation Graduate Research Fellowship under Grant No. DGE-1745301. This work

was also supported by grants to DKN from the ARO (W911NF-17-1-0024), NIH

(1R01AI127850-01A1) and the Resnick Sustainability Institute at Caltech.

References

Aminov, R.I. (2009) The role of antibiotics and antibiotic resistance in nature. *Environ Microbiol*
11: 2970–2988.
Aminov, R.I. (2010) A brief history of the antibiotic era: lessons learned and challenges for the
future. *Front Microbiol* **1**: 134.
Bankhead, S.B., Landa, B.B., Lutton, E., Weller, D.M., and Gardener, B.B.M. (2004) Minimal
changes in rhizobacterial population structure following root colonization by wild type and
transgenic biocontrol strains. *FEMS Microbiol Ecol* **49**: 307–318.
Baron, S.S., and Rowe, J.J. (1981) Antibiotic action of pyocyanin. *Antimicrob Agents Chemother*
20: 814–820.
Bérdy, J. (2012) Thoughts and facts about antibiotics: where we are now and where we are
heading. *J Antibiot* **65**: 385–395.
Bergsma-Vlami, M., Prins, M.E., and Raaijmakers, J.M. (2005) Influence of plant species on
population dynamics, genotypic diversity and antibiotic production in the rhizosphere by
indigenous Pseudomonas spp. *FEMS Microbiol Ecol* **52**: 59–69.
Brisbane, P.G., Janik, L.J., Tate, M.E., and Warren, R.F. (1987) Revised structure for the

phenazine antibiotic from Pseudomonas fluorescens 2-79 (NRRL B-15132). *Antimicrob Agents*
Chemother **31**: 1967–1971.
Cezairliyan, B., Vinayavekhin, N., Grenfell-Lee, D., Yuen, G.J., Saghatelian, A., and Ausubel,
F.M. (2013) Identification of Pseudomonas aeruginosa phenazines that kill Caenorhabditis
elegans. *PLoS Pathog* **9**: e1003101.
Collins, M.D., and Jones, D. (1981) Distribution of isoprenoid quinone structural types in
bacteria and their taxonomic implication. *Microbiol Rev* **45**: 316–354.
Costa, K.C., Bergkessel, M., Saunders, S., Korlach, J., and Newman, D.K. (2015) Enzymatic
degradation of phenazines can generate energy and protect sensitive organisms from toxicity.
MBio **6**: e01520-15.
Dar, D., Thomashow, L.S., Weller, D.M., and Newman, D.K. (2020) Global landscape of
phenazine biosynthesis and biodegradation reveals species-specific colonization patterns in
agricultural soils and crop microbiomes. *Elife* **9**.
Davies, J. (2006) Are antibiotics naturally antibiotics? *J Ind Microbiol Biotechnol* **33**: 496–499.
Davies, J., and Davies, D. (2010) Origins and evolution of antibiotic resistance. *Microbiol Mol*
Biol Rev **74**: 417–433.
Davies, J., and Ryan, K.S. (2012) Introducing the parvome: bioactive compounds in the
microbial world. *ACS Chem Biol* **7**: 252–259.
Davies, J., Spiegelman, G.B., and Yim, G. (2006) The world of subinhibitory antibiotic
concentrations. *Curr Opin Microbiol* **9**: 445–453.
Demain, A.L., and Fang, A. (2000) The natural functions of secondary metabolites. *Adv Biochem*
Eng Biotechnol **69**: 1–39.
Dietrich, L.E., Price-Whelan, A., Petersen, A., Whiteley, M., and Newman, D.K. (2006) The
phenazine pyocyanin is a terminal signalling factor in the quorum sensing network of
Pseudomonas aeruginosa. Mol Microbiol **61**: 1308–1321.
Gagliardi, J.V., Buyer, J.S., Angle, J.S., and Russek-Cohen, E. (2001) Structural and functional
analysis of whole-soil microbial communities for risk and efficacy testing following microbial
inoculation of wheat roots in diverse soils. *Soil Biol Biochem* **33**: 25–40.
Glasser, N.R., Kern, S.E., and Newman, D.K. (2014) Phenazine redox cycling enhances
anaerobic survival in *Pseudomonas aeruginosa* by facilitating generation of ATP and a proton-
motive force. *Mol Microbiol* **92**: 399–412.
Glasser, N.R., Saunders, S.H., and Newman, D.K. (2017) The colorful world of extracellular
electron shuttles. *Annu Rev Microbiol* **71**: 731–751.
Granato, E.T., Meiller-Legrand, T.A., and Foster, K.R. (2019) The evolution and ecology of
bacterial warfare. *Curr Biol* **29**: R521–R537.
Haas, D., and Défago, G. (2005) Biological control of soil-borne pathogens by fluorescent
pseudomonads. *Nat Rev Microbiol* **3**: 307–319.
Haas, D., and Keel, C. (2003) Regulation of antibiotic production in root-colonizing
Peudomonas spp. and relevance for biological control of plant disease. *Annu Rev Phytopathol*
41: 117–153.
Handelsman, J., and Stabb, E.V. (1996) Biocontrol of soilborne plant pathogens. *Plant Cell* **8**:
1855–1869.
Hassan, H.M., and Fridovich, I. (1980) Mechanism of the antibiotic action pyocyanine. *J*
Bacteriol **141**: 156–163.
Hutchings, M.I., Truman, A.W., and Wilkinson, B. (2019) Antibiotics: past, present and future.
Curr Opin Microbiol **51**: 72–80.

Jamshidi, S., Sutton, J.M., and Rahman, K.M. (2017) Computational Study Reveals the
Molecular Mechanism of the Interaction between the Efflux Inhibitor PAβN and the AdeB
Transporter from Acinetobacter baumannii. *ACS Omega* **2**: 3002–3016.
Khare, A., and Tavazoie, S. (2015) Multifactorial competition and resistance in a two-species
bacterial system. *PLoS Genet* **11**: e1005715.
Kong, X.-K., Chen, D., Huang, J.-W., Cheng, X.-K., and Jiang, J.-D. (2019) Chitinophaga
deserti sp. nov., isolated from desert soil. *Int J Syst Evol Microbiol* **69**: 1783–1788.
Kozdrój, J., Trevors, J.T., and Elsas, J.D. van (2004) Influence of introduced potential biocontrol
agents on maize seedling growth and bacterial community structure in the rhizosphere. *Soil Biol*
Biochem **36**: 1775–1784.
Kumar, S., Mukherjee, M.M., and Varela, M.F. (2013) Modulation of Bacterial Multidrug
Resistance Efflux Pumps of the Major Facilitator Superfamily. *International Journal of*
Bacteriology **2013**.
Lamers, R.P., Cavallari, J.F., and Burrows, L.L. (2013) The efflux inhibitor phenylalanine-
arginine beta-naphthylamide (PAβN) permeabilizes the outer membrane of gram-negative
bacteria. *PLoS One* **8**: e60666.
Lane, D.J. (1991) 16S/23S rRNA sequencing. In *Nucleic acid techniques in bacterial*
systematics. Stackebrandt, E., and Goodfellow, M. (eds). John Wiley & Sons, Inc., New York,
New York. pp. 115–176.
Lau, G.W., Ran, H., Kong, F., Hassett, D.J., and Mavrodi, D. (2004) *Pseudomonas aeruginosa*
pyocyanin is critical for lung infection in mice. *Infect Immun* **72**: 4275–4278.
LeTourneau, M.K., Marshall, M.J., Cliff, J.B., Bonsall, R.F., Dohnalkova, A.C., Mavrodi, D.V.,
et al. (2018) Phenazine-1-carboxylic acid and soil moisture influence biofilm development and
turnover of rhizobacterial biomass on wheat root surfaces. *Environ Microbiol* **20**: 2178–2194.
Lin, S.-Y., Hameed, A., Wen, C.-Z., Liu, Y.-C., Shen, F.-T., Hsu, Y.-H., *et al.* (2015)
Chryseobacterium echinoideorum sp. nov., isolated from sea urchins (Tripneustes gratilla). *Int J*
Syst Evol Microbiol **65**: 3985–3990.
Linares, J.F., Gustafsson, I., Baquero, F., and Martinez, J.L. (2006) Antibiotics as intermicrobial
signaling agents instead of weapons. *Proc Natl Acad Sci USA* **103**: 19484–19489.
Liu, G.Y., and Nizet, V. (2009) Color me bad: microbial pigments as virulence factors. *Trends*
Microbiol **17**: 406–413.
Manyi-Loh, C., Mamphweli, S., Meyer, E., and Okoh, A. (2018) Antibiotic use in agriculture
and its consequential resistance in environmental sources: potential public health implications.
Molecules **23**.
Mavrodi, D.V., Blankenfeldt, W., and Thomashow, L.S. (2006) Phenazine compounds in
fluorescent Pseudomonas spp. biosynthesis and regulation. *Annu Rev Phytopathol* **44**: 417–445.
Mavrodi, D.V., Mavrodi, O.V., Parejko, J.A., Bonsall, R.F., Kwak, Y.-S., Paulitz, T.C., *et al.*
(2012) Accumulation of the antibiotic phenazine-1-carboxylic acid in the rhizosphere of dryland
cereals. *Appl Environ Microbiol* **78**: 804–812.
Mazurier, S., Corberand, T., Lemanceau, P., and Raaijmakers, J.M. (2009) Phenazine antibiotics
produced by fluorescent pseudomonads contribute to natural soil suppressiveness to Fusarium
wilt. *ISME J* **3**: 977–991.
Mazzola, M., Cook, R.J., Thomashow, L.S., Weller, D.M., and Pierson, L.S. (1992) Contribution
of phenazine antibiotic biosynthesis to the ecological competence of fluorescent pseudomonads
in soil habitats. *Appl Environ Microbiol* **58**: 2616–2624.
Mazzola, M., Fujimoto, D.K., Thomashow, L.S., and Cook, R.J. (1995) Variation in Sensitivity

of Gaeumannomyces graminis to Antibiotics Produced by Fluorescent Pseudomonas spp. and
Effect on Biological Control of Take-All of Wheat. *Appl Environ Microbiol* **61**: 2554–2559.
McRose, D.L., and Newman, D.K. (2021) Redox-active antibiotics enhance phosphorus
bioavailability. *Science* **371**: 1033–1037.
Meirelles, L.A., and Newman, D.K. (2018) Both toxic and beneficial effects of pyocyanin
contribute to the lifecycle of *Pseudomonas aeruginosa. Mol Microbiol* **110**: 995–1010.
Meirelles, L.A., Perry, E.K., Bergkessel, M., and Newman, D.K. (2021) Bacterial defenses
against a natural antibiotic promote collateral resilience to clinical antibiotics. *PLoS Biol* **19**:
e3001093.
Nikaido, H. (1989) Role of the Outer Membrane of Gram-Negative Bacteria in Antimicrobial
Resistance. In *Microbial resistance to drugs*. Bryan, L.E. (ed.). Springer Berlin Heidelberg,
Berlin, Heidelberg. pp. 1–34.
Noto, M.J., Burns, W.J., Beavers, W.N., and Skaar, E.P. (2017) Mechanisms of Pyocyanin
Toxicity and Genetic Determinants of Resistance in Staphylococcus aureus. *J Bacteriol* **199**.
Pao, S.S., Paulsen, I.T., and Saier, M.H. (1998) Major facilitator superfamily. *Microbiol Mol*
Biol Rev **62**: 1–34.
Perry, E.K., and Newman, D.K. (2019) The transcription factors ActR and SoxR differentially
affect the phenazine tolerance of *Agrobacterium tumefaciens. Mol Microbiol* **112**: 199–218.
Price-Whelan, A., Dietrich, L.E.P., and Newman, D.K. (2006) Rethinking "secondary"
metabolism: physiological roles for phenazine antibiotics. *Nat Chem Biol* **2**: 71–78.
Roquigny, R., Novinscak, A., Léger, G., Marcoux, N., Joly, D.L., and Filion, M. (2018)
Deciphering the Rhizosphere and Geocaulosphere Microbiomes of Potato Following Inoculation
with the Biocontrol Agent *Pseudomonas fluorescens* Strain LBUM223. *Phytobiomes* **2**: 92–99.
Schiessl, K.T., Hu, F., Jo, J., Nazia, S.Z., Wang, B., Price-Whelan, A., *et al.* (2019) Phenazine
production promotes antibiotic tolerance and metabolic heterogeneity in *Pseudomonas*
aeruginosa biofilms. *Nat Commun* **10**: 762.
Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., *et al.* (2012)
Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**: 676–682.
Schuster, S., Bohnert, J.A., Vavra, M., Rossen, J.W., and Kern, W.V. (2019) Proof of an Outer
Membrane Target of the Efflux Inhibitor Phe-Arg-β-Naphthylamide from Random Mutagenesis.
Molecules **24**.
Sengupta, S., Chattopadhyay, M.K., and Grossart, H.-P. (2013) The multifaceted roles of
antibiotics and antibiotic resistance in nature. *Front Microbiol* **4**: 47.
Séveno, N.A., Morgan, J.A.W., and Wellington, E.M.H. (2001) Growth of *Pseudomonas*
aureofaciens PGS12 and the dynamics of HHL and phenazine production in liquid culture, on
nutrient agar, and on plant roots. *Microb Ecol* **41**: 314–324.
Singh, P., Singh, H., Kim, Y.-J., and Yang, D.-C. (2017) Pedobacter panacis sp. nov., isolated
from Panax ginseng soil. *Antonie Van Leeuwenhoek* **110**: 235–244.
Smiley, R.W., and Cook, R.J. (1973) Relationship between take-all of wheat and rhizosphere pH
in soils fertilized with ammonium vs. nitrate-nitrogen. *Phytopathology* **63**: 882–890.
Tagele, S.B., Lee, H.G., Kim, S.W., and Lee, Y.S. (2019) Phenazine and 1-Undecene Producing
Pseudomonas chlororaphis subsp. aurantiaca Strain KNU17Pc1 for Growth Promotion and
Disease Suppression in Korean Maize Cultivars. *J Microbiol Biotechnol* **29**: 66–78.
Thomashow, L.S., and Weller, D.M. (1988) Role of a phenazine antibiotic from Pseudomonas
fluorescens in biological control of Gaeumannomyces graminis var. tritici. *J Bacteriol* **170**:
3499–3508.

Thomashow, L.S., Weller, D.M., Bonsall, R.F., and Pierson, L.S. (1990) Production of the antibiotic phenazine-1-carboxylic acid by fluorescent *Pseudomonas* species in the rhizosphere of wheat. *Appl Environ Microbiol* **56**: 908–912.

Turner, J.M., and Messenger, A.J. (1986) Occurrence, biochemistry and physiology of phenazine pigment production. *Adv Microb Physiol* **27**: 211–275.

Tyc, O., Song, C., Dickschat, J.S., Vos, M., and Garbeva, P. (2017) The ecological role of volatile and soluble secondary metabolites produced by soil bacteria. *Trends Microbiol* **25**: 280–292.

VanDrisse, C.M., Lipsh-Sokolik, R., Khersonsky, O., Fleishman, S.J., and Newman, D.K. (2021) Computationally designed pyocyanin demethylase acts synergistically with tobramycin to kill recalcitrant Pseudomonas aeruginosa biofilms. *Proc Natl Acad Sci USA* **118**.

Voggu, L., Schlag, S., Biswas, R., Rosenstein, R., Rausch, C., and Götz, F. (2006) Microevolution of cytochrome bd oxidase in Staphylococci and its implication in resistance to respiratory toxins released by Pseudomonas. *J Bacteriol* **188**: 8079–8086.

Wang, Y., and Newman, D.K. (2008) Redox reactions of phenazine antibiotics with ferric (hydr)oxides and molecular oxygen. *Environ Sci Technol* **42**: 2380–2386.

White, D., Drummond, J., and Fuqua, C. (2012) Electron Transport. In *The Physiology and Biochemistry of Prokaryotes*. Oxford University Press, New York, New York. pp. 146–174.

Wolloscheck, D., Krishnamoorthy, G., Nguyen, J., and Zgurskaya, H.I. (2018) Kinetic Control of Quorum Sensing in Pseudomonas aeruginosa by Multidrug Efflux Pumps. *ACS Infect Dis* **4**: 185–195.

Yu, J.M., Wang, D., Pierson, L.S., and Pierson, E.A. (2018) Effect of Producing Different Phenazines on Bacterial Fitness and Biological Control in Pseudomonas chlororaphis 30-84. *Plant Pathol J* **34**: 44–58.

Zhu, K., Chen, S., Sysoeva, T.A., and You, L. (2019) Universal antibiotic tolerance arising from antibiotic-triggered accumulation of pyocyanin in *Pseudomonas aeruginosa. PLoS Biol* **17**: e3000573.

Figure 1: Taxonomic distribution of bacterial isolates from wheat rhizosphere and bulk soil samples.

This plot depicts the proportion of isolates from each sample that belonged to the 4 represented phyla. Each column represents one soil sample, and each box within the columns represents an individual isolate colored by the phylum to which it belongs (e.g. 6 boxes comprising one column indicates that 6 strains were isolated from that sample). B = Between (bulk soil), V = Virgin (bulk soil), W = Wheat (rhizosphere).

Figure 2: Distribution of PCA resistance phenotypes across phyla and soil sample type.
A-B. Heat maps depicting the growth of the strains at pH 5.1 (A) and pH 7.3 (B). Each row represents a strain. The leftmost columns are colored according to the ratio of growth on PCA-containing agar versus PCA-free agar; magenta indicates sensitivity to PCA while green indicates resistance to PCA. The right two columns in each heat map are colored according to the separate values for growth on PCA-containing agar (+ PCA) or solvent control agar (- PCA), with darker green indicating more growth. Values are the mean of two to four biological replicates that each comprised three technical replicates. Asterisks indicate strains that displayed markedly variable susceptibility to PCA across biological replicates. Strains for which fewer than two biological replicates grew to stationary phase in the PCA-free control at the given pH, or for which color interfered with growth quantification, were omitted from this analysis. See Methods for a description of how growth was quantified. **C.** Relative abundance of putative phenazine producers (*phz*+ bacteria) across the three types of soil samples. Error bars represent the standard deviation. **D-E.** Density plots (i.e. smoothed histograms) representing the distribution of PCA resistance phenotypes at pH 5.1 (D) and pH 7.3 (E) among strains isolated from Between (bulk), Virgin (bulk), or Wheat (rhizosphere) soil**.** Higher values along the x-axis indicate greater resistance to PCA. If multiple identical strains were isolated from the same soil type, only one representative was counted; thus, 26 strains were isolated from Between, 39 from Virgin, and 45 from Wheat. Colored tick marks above the x-axis represent where individual isolates fall along the range of PCA resistance phenotypes.

Figure 3: Growth of representative strains over time with and without PCA.
Growth was quantified as described in the Methods. Solid lines represent the growth of spotted cultures on PCA-free agar, and dashed lines represent the growth of spotted cultures on agar containing 100 µM PCA. Blue represents growth at pH 7.3 and pink represents growth at pH 5.1. Data points are the mean of three technical replicates from a representative biological replicate for each strain, and the shaded ribbon represents the standard deviation. The selected strains depicted here cover the range of PCA susceptibility phenotypes observed in each genus. The full set of biological replicates for all tested isolates is displayed in supplementary Figs. S4-S7.

Figure 4: PCA reduction rates of selected strains at pH 5.1 and pH 7.3.
A. PCA reduction rates of representative strains from each phylum at pH 5.1 and pH 7.3. The values represent the mean of three biological replicates and error bars represent the standard deviation. Reduction rates were normalized to OD_{600} values. **B.** PCA reduction rates versus each strain's growth ratio (+PCA/-PCA) at early stationary phase. The growth ratios are the same values calculated and used in Fig. 2.

Figure 5: Effects of efflux pump inhibitors on PCA susceptibility.
A-B. Growth curves of strains for which treatment with a combination of the efflux pump inhibitors (EPI) reserpine and PAβN did not affect susceptibility to PCA at pH 5.1 (A) or pH 7.3 (B). **C.** Growth curves of strains for which treatment with reserpine and PAβN potentiated the toxicity of PCA. **D.** Growth curves of strains for which treatment with reserpine and PAβN improved growth in the presence of PCA (strain W4I17) or vice versa (strains W1I16 and W4I9-1). In all panels, data points are the mean of three biological replicates and the shaded ribbon represents the standard deviation. The concentrations of reserpine and PAβN used for each strain can be found in Table S2.