Native American Ancestry and Pigmentation Allele Contributions to Skin Color in a Caribbean

Population

[1,2,*]Khai C Ang, [1,2]Victor A Canfield, [1,2]Tiffany C Foster, [1,2]Katherine P Reid, [3]Shou L Leong, [4,5,6]Yuka I

Kawasawa, [4,7]Dajiang Liu, [8]John Hawley, [1,2,4,5,*]Keith C Cheng

[1]Department of Pathology, Penn State College of Medicine, Hershey, PA, USA [2]Jake Gittlen Laboratories

for Cancer Research, Penn State College of Medicine, Hershey, PA, USA [3]Department of Family &

Community Medicine, Penn State College of Medicine, Hershey, PA, USA [4]Department of Biochemistry

and Molecular Biology, Penn State College of Medicine, Hershey, PA, USA [5]Department of Pharmacology,

Penn State College of Medicine, Hershey, PA, USA [6]Institute of Personalized Medicine, Penn State College

of Medicine, Hershey, PA, USA [7]Department of Public Health Sciences, Penn State College of Medicine,

Hershey, PA, USA; [8]Salybia Mission Project, Dominica.

*Corresponding authors

Khai C Ang, kca2@psu.edu; Keith C Cheng, kcheng76@gmail.com

Abstract

Interest in the genetic basis of variation in skin pigmentation in Native American populations led us to

seek indigenous populations of the Western Hemisphere with African and minimal European admixture

to study the effect of Native American ancestry on skin color. Admixture analysis from DNA collected from

458 individuals in the Kalinago territory of the Commonwealth of Dominica showed shared ancestry with

East Asians at K=3 and 55% Native American, 32% African, and 11% European ancestry at K=6, the highest

Native American ancestry of Caribbean populations. Skin pigmentation was 20 to 80 melanin units,

averaging 46. Three albino individuals were homozygous for multi-nucleotide polymorphism $OCA2^{NW273KV}$

of African origin, whose population allele frequency was 0.03 and single allele effect size was -8 melanin

units. Hypopigmenting allele frequencies for $SLC24A5^{A111T}$ and $SLC45A2^{L374F}$ were 0.14 and 0.05, whose

single allele effect sizes were -6 and -3, respectively. Skin color plots of individuals lacking known

hypopigmenting alleles suggests that Native American Ancestry reduced pigmentation by more than 20

melanin units (low and high estimates 21.8 and 28.5). Shared ancestry with East Asians at K=3 suggests

potential sharing of one or more pigmentation alleles.

Introduction

Skin pigmentation is a highly heritable polygenic trait influenced by health and environment (Barsh, 2003). European variants that lighten pigmentation include non-ancestral coding polymorphisms in *SLC24A5* (Basu Mallick et al., 2013; Lamason et al., 2005; Soejima and Koda, 2007) and *SLC45A2* (Lucotte and Yuasa, 2013; Soejima and Koda, 2007; Yuasa et al., 2006) that are nearly fixed. However, the genetic basis for lighter skin pigmentation in Native American and East Asian populations, who carry the same ancestral alleles as Africans at these loci, has yet to be established. Dark skin was the ancestral phenotype for anatomically modern humans (Jablonski and Chaplin, 2000; Lamason et al., 2005), whose evolution towards lighter skin at higher latitudes occurred independently in eastern and western Eurasia (Adhikari et al., 2019; Basu Mallick et al., 2013; Norton et al., 2007; Yang et al., 2016), potentially driven by a UV-dependent photoactivation step in the formation of vitamin D (Engelsen, 2010; Hanel and Carlberg, 2020; Holick, 1981; Loomis, 1967).

Native Americans share common ancestry with East Asians (Derenko et al., 2010; Tamm et al., 2007), diverging before 15 kya (Gravel et al., 2013; Moreno-Mayar et al., 2018; Reich et al., 2012), but the extent to which these populations share pigmentation variants remains to be determined. High European admixture is characteristic of most Native American populations (Brown et al., 2017; Gravel et al., 2013; Klimentidis et al., 2009; Reich et al., 2012), complicating the characterization of pigmentation variants specific to Native Americans.

Prior to European contact, the Caribbean islands were inhabited by populations who migrated from the northern coast of South America (Benn-Torres et al., 2008; HARVEY et al., 1969; Honychurch, 2012; "Island Caribs," 2016; Torres et al., 2015, 2013). During the Colonial period, large numbers of Africans were introduced into the Caribbean as slave labor (Honychurch, 2012; Torres et al., 2013). As a consequence African and European admixture, and high mortality among the indigenous populations, Native American ancestry now contributes only a minor portion of the ancestry of most Caribbean islanders (1000 Genomes Project Consortium, 2010; The 1000 Genomes Project Consortium, 2015; Torres et al., 2015, 2013). The islands of Dominica and St. Vincent were the last colonized by Europeans, in the late 1700s (Honychurch, 2012, 1998; Rogoziński, 2000). In 1903, the British granted 15 km^2 (3,700 acres) on the eastern coast of Dominica as a reservation for the Kalinago, who were then called "Carib". When Dominica gained Independence in 1978, legal rights and a degree of protection from assimilation were gained by the inhabitants of the Carib Reserve (redesignated *Kalinago Territory* in 2015) (Honychurch, 2012). The

Kalinago, numbering about 3,000 living within the Territory ("Kalinago Territory," 2021), consider
themselves to be of primarily Native American and African ancestry.

Early in our genetic and phenotypic survey of the Kalinago, we noted one individual with albinism, and
discovered that two others were known to reside in the Territory. Oculocutaneous albinism (OCA) is a
recessive trait characterized by visual system abnormalities and hypopigmentation of skin, hair, and eyes
(Gargiulo et al., 2011; Gronskov et al., 2007; Grønskov et al., 2014; Hong et al., 2006; Vogel et al., 2008)
that is caused by mutations in any of several autosomal pigmentation genes (Carrasco et al., 2009;
Edwards et al., 2010; Gao et al., 2017; Grønskov et al., 2013; Kausar et al., 2013; King et al., 2003; Spritz
et al., 1995; Stevens et al., 1997, 1995; Vogel et al., 2008; Woolf, 2005; Yi et al., 2003). Here, we report on
ancestry, distribution of measured skin color, identification of an albinism allele, and the hypopigmenting
effects of this allele and the European $SLC24A5^{A111T}$ and $SLC45A2^{L374}$ alleles in a sample representing 15%
of the population (Figure S1).

Results & Discussion

Our search for a population admixed for Native American/African ancestries led us to the "Carib"
population in the Commonwealth of Dominica. Observations from an initial trip to Dominica suggested
wide variation in Kalinago skin color. Pursuit of the genetic studies described here required learning about
oral and written histories, detailed discussion with community leadership, and IRB approval from Ross
University (until Hurricane Maria in 2017, the largest medical school in Dominica), the Department of
Health of the Commonwealth of Dominica, and relationship-building with three administrations the
Kalinago Council over 15 years.

Population Sample

Our DNA and skin-color sampling program encompassed 458 individuals, representing 15% of the
population of the territory and including the three known albino individuals. Ages ranged from 6 to 93
(Table S1 and Figure S2). We were able to obtain genealogical information for about half of the parents
(243 mothers and 194 fathers); community-defined ancestry (described as 'black,' 'Kalinago,' or 'Mixed')
for both parents was obtained for 426 individuals (92% of sample), including 221 from which DNA samples
were obtained.

Kalinago Ancestry

The earliest western mention of the Kalinago (originally as "Caribs") was in Christopher Columbus's journal
dated 26th November 1492 (Honychurch, 2012). Little is known about the detailed cultural and genetic
similarities and differences between them and other pre-contact groups such as the Taino in the
Caribbean. African admixture in the present Kalinago population derived from the African slave trade, but
we were unable to identify written historical record that includes specific regional origin or well-defined
contributions from other groups. The population's linguistics is uninformative, as they speak the same
French-based Antillean Creole spoken on the neighboring islands of Guadeloupe and Martinique (in
addition to English).

Figure 1: Admixture analysis of Kalinago compared with Human Genome Diversity Project populations.
Results are depicted using stacked bar plots, with one column per individual. At K=3, the Kalinago, Native
Americans, Oceanians, and East Asians fall into the same cluster. At K=4, the Kalinago and the Native
Americans are separated from the East Asians.

To study Kalinago population structure, we combined our Kalinago SNP genotype data with HGDP (Li et
al., 2008) data and analyzed using ADMIXTURE (Figures 1 and S3) as described in Methods. Shared
ancestry of the Kalinago with East Asians at K=3 suggests the potential sharing of one or more lighter skin
alleles between those populations. At K=4 and higher, a Native American component (that predominates
in Kalinago) separates from the East Asian component. Consistent with prior work, an Oceanian
component appears at K=5, and a Central & South Asian component appears at K=6. These are minor
components in our Kalinago sample (average <1%) (Table S3). On average, the Kalinago show 55% Native
American, 32% African, and 11% European ancestry. The individual with least admixture has 94% Native
American and 6% African. Principal component analysis (Figure S4) provides additional insight into the
relationships between Kalinago and potential source populations.

Figure 2. Comparison of Kalinago ancestry with that of other populations in the Western Hemisphere.
Ternary plots show estimated proportions of African (AFR), European (EUR) and Native American (NAM)
ancestry. **A,** Comparison of individuals (n=452, omitting 6 individuals with EAS > 0.1) genotyped in this
study to individuals (n=38) from southern Dominica sampled by Torres et al., 2013, **B,** Comparison of
population averages. Kalinago, this study (n=458); Islands (BT) indicates Caribbean islanders reported in
Torres et al., 2013, with Dominica labeled; admixed (adm) AFR (1KGP) and admixed NAM (1KGP) represent
admixed populations from The 1000 Genomes Project Consortium, 2015, with Caribbean samples PUR
(Puerto Rico) and ACB (Barbados) labeled; and AMR (Reich) indicates mainland Native American samples
reported in Reich et al., 2012. Inset shows ancestries at vertices.

Our analysis of Kalinago ancestry revealed considerably more Native American and less European ancestry

than the Dominican samples of Torres et al. (2013) from outside of the Kalinago Territory (Figure 2A) and

those admixed populations from the 1000 Genomes Project (The 1000 Genomes Project Consortium,

2015). Some Western Hemisphere Native Americans reported in Reich et al. (2012) have varying

proportions of European but very little African admixture (Figure 2B) . Overall, the Kalinago have more

Native American and less European ancestry than any other Caribbean population.

Specifically, the 55% Native American ancestry observed for the Kalinago is far greater than the reported

13% in Puerto Rico (Gravel et al., 2013), 10-15% for Tainos across the Caribbean (Schroeder et al., 2018),

and 8% for Cubans (Marcheco-Teruel et al., 2014). Samples from Jamaica and the Lesser Antilles (Torres

et al., 2015, 2013) yielded an average of 7.7% Native American ancestry (range 5.6% to 16.2%), with the

highest value from a population of Kalinago ancestry outside the reservation in Dominica. Relevant to the

potential mapping of Native American light skin color alleles, the Kalinago has the lowest European

ancestry compared to other reported Caribbean Native Americans in St. Kitts (8.2%), Barbados (11.5%)

and Puerto Rico (71%) (Torres et al., 2013). Potential reasons for the high percentage of Native American

ancestry in the Kalinago likely include their segregation within the 3,700-acre Kalinago Territory in

Dominica granted by the British in 1903, and the Kalinago tradition of women marring non-Kalinago being

required to leave the Territory; non-Kalinago spouses of Kalinago men are allowed to move to the

Territory (KCC, KCA, Personal Communication with Kalinago Council, 2014). These factors help explain why

Kalinago samples collected outside the Kalinago territory (Torres et al., 2013), show lower fractional

Native American ancestry.

During our fieldwork, it was noted that members of the Kalinago community characterized themselves

and others in terms of perceived ancestry as "black," "Kalinago," or "mixed," based primarily on

phenotype. These folk categorizations were broadly supported by differences in admixture (Figures S5,

S6). Compared to individuals identified as "Mixed," those identified as "Kalinago" have on average more

Native American ancestry (67% vs 51%), less European ancestry (10% vs 14%), and less African ancestry

(23% vs 34%). Individuals described as "Kalinago" were slightly lighter and had a narrower MI distribution

(42.5 ± 5.6, mean ± SD) than those described as "Mixed" (45.8 ± 9.6).

Kalinago Skin Color Variation

Melanin index unit (MI) calculated from skin reflectance (see Methods) was used as a quantitative

measure of melanin pigmentation (Ang et al., 2012; Diffey et al., 1984). The MI in the Kalinago ranged

from 20.7 to 79.7 (Figure S7), averaging 45.7. The three Kalinago albino individuals sampled had the lowest

values (20.7, 22.4 and 23.8). Excluding these, the MI ranged between 28.7 to 79.7 and averaged 45.9. For

comparison, the MI averaged 25 and 21 for people of East Asian and European ancestry, respectively, as
measured with the same equipment in our laboratory (Ang et al., 2012; Tsetskhladze et al., 2012). This
range is similar to that of another indigenous population related to East Asians, the Senoi of Peninsular
Malaysia (MI 24 to 78; mean = 45.7) (Ang et al., 2012). The Senoi are believed include admixture from
Malaysian Negritos whose pigmentation is darker (mean = 55) (Ang et al., 2012) than that of the average
Kalinago. In comparison, the average MI was 53.4 for Africans in Cape Verde (Beleza et al., 2012) and 59
for African-Americans (Shriver et al., 2003).

An OCA2 albinism allele in the Kalinago

Oculocutaneous albinism (OCA) is a genetically determined condition characterized by nystagmus,
reduced visual acuity, foveal hypoplasia and strabismus as well as hypopigmentation of the skin, hair and
eye (Dessinioti et al., 2009; van Geel et al., 2013). The three sampled albino individuals had pale skin (MI
20.7, 22.4 and 23.8 vs. 29-80 for non-albinos), showed nystagmus, and reported photophobia and high
susceptibility to sunburn. In contrast to the brown irides and black hair of most Kalinago, including their
parents, the albino individuals had golden blonde hair and grey irides with varying amounts of green and
blue.

Whole exome sequencing of one albino individual and one parent (obligate carrier) identified 12 variant
alleles in 7 oculocutaneous albinism genes (or genomic regions) that were heterozygous in the parent and
homozygous for a non-reference allele in the albino individual (summarized in Table S4A); none was a
nonsense or splice site variant. Five of the twelve potential candidate mutations found by this approach
were intronic, one was synonymous, one was located in the 5'UTR, and three were in 3'UTR (Table S4B).
Two missense variants were found in *OCA2*: SNP rs1800401 (c.913C>T or p.Arg305Trp in exon 9), *R305W*,
and multi-nucleotide polymorphism rs797044784 in exon 8 (c.819_822delCTGGinsGGTC;
p.Asn273_Trp274delinsLysVal), *NW273KV*.

Of 458 Kalinago *OCA2* genotypes, 26 carried *NW273KV* and 60 carried *R305W* (Table 1). Only *NW273KV*
homozygotes were albino, and neither of the two *R305W* homozygotes (who were either heterozygous
or homozygous ancestral for *NW273KV*) were albino. Notably, one Kalinago individual who is homozygous
derived for *R305W* mutation but homozygous ancestral for *NW273* has an MI of 72, among the darkest in
the entire population. Therefore, *R305W* is not an albinism allele, as expected from its high population
frequencies (> 0.10 in some African, South Asian, and European populations)(The 1000 Genomes Project
Consortium, 2015). Notably, the black hair and dark eyes of Kalinago *R305W* homozygotes without the

NW273KV indicate that the *in silico* predictions by SIFT, Polyphen 2.0 and PANTHER that *R305W* is a likely

pathogenic variant (Kamaraj and Purohit, 2013) are incorrect. The observed patterns of zygosity suggest

that *NW273KV* arose on the background of a haplotype carrying the widespread *R305W* variant.

Table 1. Albinism among *NW273KV* and *R305W* genotypes.

Allele/Genotype		NW273KV genotype			
		Homozygous Ancestral[a]	Heterozygous	Homozygous Derived	Total
***R305W* genotype**	Homozygous Ancestral	398	0	0	398
	Heterozygous	33	22	0	55
	Homozygous Derived	1	1	3*	5
	Total	432	23	3*	458

193 [a] Ancestral=reference allele and derived=alternate allele for both variants.
194 * Albino phenotype.
195

Figure 3. Haplotype analysis for three albino individuals. Inner two lines indicate NAM (red) or AFR
(black) ancestry; no EUR ancestry was found in this genomic region. For this local ancestry analysis, the
region shown here consisted of 110 non-overlapping segments with 7 to 346 SNPs each (mean 65). The
deduced extent of shared albino haplotype (dotted light blue lines) is indicated on each
chromosome. Minimum homozygous region (determined by albino individual 1) shared by all three
albino individuals is shown at expanded scale below. Genes in this region are labeled, and the position of
the *NW273KV* polymorphism in *OCA2* is indicated by the red arrowhead.

The sole prior reported instance of albinism that includes the *NW273KV* variant involves a compound
heterozygote of African-American descent (Garrison et al., 2004; Lee et al., 1994). The conservation of the
NW sequence among vertebrates and its inclusion in a potential N-linked glycosylation site (Rinchik et al.,
1993), that is eliminated by the mutation, supports the pathogenicity of this mutation. The *NW273KV*
frequency in our sample (0.03) translates into a Hardy-Weinberg albinism frequency (p^2 = 0.0009) of ~1
per 1000, as observed (3 in a population of about 3000). Examination of publicly available data reveals
three $OCA2^{NW273KV}$ heterozygotes in the 1000 Genome Project, a pair of siblings from Barbados (ACB) and
one individual from Sierra Leone (MSL) (The 1000 Genomes Project Consortium, 2015). These three
individuals share a haplotype of 1.5 Mb but no more than 1.04 Mb are shared with the Kalinago albinos.

One of the three albinos was also heterozygous for $SLC24A5^{A111T}$, but his skin and hair color was not lighter
than that of the other two albinos, who were homozygous for the ancestral allele at $SLC24A5^{A111}$; this
observation is consistent with epistasis of *OCA2* albinism over $SLC24A5^{A111T}$. Eight sampled non-albino
individuals had combinations of lighter hair, fair skin, and lighter irides, among these, seven were
heterozygous for $SLC24A5^{A111T}$, and four were heterozygous for the albino variant. A precise
understanding of the phenotypic effects of the combinations of these hypopigmenting alleles will require
further study.

Genetic Contribution of Native American Ancestry to Kalinago Skin Color Variation

Among the primary goals of this work is quantification of the contribution of Native American ancestry to
skin pigmentation. Answering this question required identifying the Kalinago as a Native American
population admixed for African Ancestry. To minimize interference of pigmentation lightening alleles
specific to Europeans, we confirmed the relatively small 15% European ancestry of this population, and
now need to identify individuals lacking either of the two known hypopigmenting variants fixed in
Europeans, $SLC24A5^{A111T}$ and $SLC45A2^{L374F}$. In addition, we needed to identify individuals that are also
ancestral for the $OCA2^{NW273KV}$ albinism allele in this population.

Figure 4. Skin color distribution of Kalinago samples according to genotype. Triple ancestral for three pigmentation alleles ($SLC24A5^{111A}$, $SLC45A2^{374L}$ and $OCA2^{273NW}$) and derived (heterozygous or homozygous) for the indicated variant: $OCA2^{NW273KV}$; $SLC24A5^{A111T}$; and $SLC45A2^{L374F}$.

To control for the effects of the known European pigmentation loci, all Kalinago samples were genotyped

for two known pigmentation polymorphisms of European origin, $SLC24A5^{A111T}$ and $SLC45A2^{L374F}$. The

phenotypic effects of these variants and $OCA2^{NW273KV}$ are shown in the histograms of Figure 4. Each variant

decreases melanin pigmentation, and homozygotes are lighter than heterozygotes. The greatest effect is

seen in the $OCA2^{NW273KV}$ homozygotes (the albino individuals), as previously noted. The frequencies of the

derived alleles of $SLC24A5^{A111T}$ and $SLC45A2^{L374F}$ in the Kalinago sample are 0.14 and 0.05, respectively.

The higher frequency of $SLC24A5^{A111T}$ compared to $SLC45A2^{L374F}$ frequencies is not explained solely by

European admixture, given that most Europeans are fixed for both alleles. This deviation can, however,

be accounted for by the involvement of source populations that have a lower frequency of $SLC45A2^{L374F}$

than $SLC24A5^{A111T}$. The 0.03 excess of $SLC24A5^{A111T}$ frequency over EUR ancestry indicates a non-negligible

frequency of $SLC24A5^{A111T}$ in one or more African source populations. At 32% AFR ancestry, this

corresponds to an average $SLC24A5^{A111T}$ frequency of about 0.09 for the AFR source populations. Although

many sub-Saharan West African populations (the likeliest source of AFR ancestry in the

Kalinago)(Micheletti et al., 2020) have lower $SLC24A5^{A111T}$ frequencies, a similar frequency is observed in

the Mende of Sierra Leone (MSL) (Micheletti et al., 2020; The 1000 Genomes Project Consortium, 2015),

while some West African populations such as Hausa and Mandinka have frequencies exceeding 0.10

(Cheung et al., 2000; Rajeevan et al., 2012). The 0.06 deficit of $SLC45A2^{L374F}$ frequency compared to EUR

ancestry (11%) corresponds to an average $SLC45A2^{L374F}$ frequency in the European source population close

to 0.5. This is far below the frequency of 0.82 observed the 2015 Genomes Project Spanish population

sample (IBS) (The 1000 Genomes Project Consortium, 2015). It should be noted that the major component

in North African and Middle Eastern populations is not distinguished from Europeans in our analysis; these

populations (and also inhabitants of Andalusia in Spain) have a wide range of $SLC45A2^{L374F}$ frequencies

dropping considerably below that of IBS (Cheung et al., 2000; Rajeevan et al., 2012). We are unable with

existing information to definitively account for the higher frequency of $SLC24A5^{A111T}$ over that

of $SLC45A2^{L374F}$.

Table 2. Effect sizes for covariates in full model with 10 Principal Components

covariate	Effect size[a]	Adjusted *p*-value
rs1426654 ($SLC24A5^{A111T}$)	-5.8	1.5E-12
rs16891982 ($SLC45A2^{L374F}$)	-2.8	0.015
albino allele ($OCA2^{NW273KV}$)	-7.8	2.5E-05
sex (female vs male)	-2.4	0.0013

[a] per allele effect size, in melanin units, for *A111T* and *L374F*; effect of first allele for albino variant

To investigate the relative contributions of genetic variation to skin color, we performed genome wide

association analyses using an additive model for Melanin Index, conditioning on sex, ancestry, and

genotypes for $SLC24A5^{A111T}$, $SLC45A2^{L374F}$ and $OCA2^{NW273KV}$. These analyses omitted the three albino
individuals. We found that sex, all three genotyped polymorphisms, and the first principal component
were statistically significant (effect sizes shown in Table 2). Effect sizes were about -6 units (per allele)
for $SLC24A5^{A111T}$, -3 units for $SLC45A2^{L374F}$ and -8 units for the first $OCA2^{NW273KV}$ allele. When controlling for
$OCA2^{NW273KV}$ status, $OCA2^{R305W}$ had no detectable effect on skin color (not shown).

The effect size for $SLC24A5^{A111T}$ is consistent with previously reported results of -5 melanin units for
African-Americans (Lamason et al., 2005; Norton et al., 2007) and -5.5 for the admixed inhabitants of the
Cape Verde islands (Beleza et al., 2013). Reported effect sizes for continental Africans are both higher and
lower -7.7 in Crawford et al. (2017) and -3.6 Martin et al. (2017b), while the estimated effect size in the
CANDELA study (GWAS of combined admixed populations from Mexico, Brazil, Columbia, Chile and Peru)
(Adhikari et al., 2019) is about -3 melanin units. For $SLC45A2^{L374F}$, significance was found in Beleza et al.
(2013) and Adhikari et al. (2019) and for African Americans but not the African Caribbean subsample in
Norton et al (2007) (Norton et al., 2007)

Our estimate that a single $OCA2^{NW273KV}$ allele causes about -8 melanin units of skin lightening is the first
reported population-based effect size measurement for any albinism allele. To study the effect of
homozygosity, we applied the estimated parameters to the three albinos, who were lighter by an average
of 10 units than predicted by the additive model ($p< 0.0033$, 1-tailed t-test). An additive model for skin
color in albinos is rejected; the nature of the non-linearity or epistasis remains to be investigated.

The strong dependence of pigmentation on ancestry for individuals lacking hypopigmenting alleles
$SLC24A5^{A111T}$, $SLC45A2^{L374F}$ and $OCA2^{NW273KV}$ is depicted in Figure 5. Positive deviations from the best fit
are apparent at both high and low NAM ancestry, but these do not change the conclusions that AFR
ancestry contributes to darker skin compared to NAM ancestry, and more importantly, that there are skin-
lightening variants of Native American origin.

Figure 5. Dependence of Melanin Unit on ancestry for Kalinago. Only individuals who are ancestral for
SLC24A5[111A]*, SLC45A2*[374L], and *OCA2*[273NW] alleles are shown (n=276). The dotted red line represents the
best fit (linear regression).

To estimate the contribution of Native American ancestry to skin hypopigmenting alleles, we analyzed the

276 samples without the *SLC24A5*[A111T], *SLC45A2*[L374F] and *OCA2*[NW273KV] polymorphisms plotted in Figure 5.

One conservative estimate of the effect of Native American Ancestry on skin color, -21.8 melanin units, is

provided by the difference in pigmentation between Kalinago with less than 20% Native American

ancestry (MI= 65.6), and Kalinago of more than 80% Native American ancestry (MI= 43.8). An alternative

and larger estimate, -28.5 melanin units, derives if we estimate the pigmentation of Kalinago individuals

with >80% African ancestry (MI = 72.3). In light of the shared ancestry between Native Americans and East

Asians shown by admixture analysis at K=3 (Figure 1), it can be expected that individual hypopigmenting

alleles of significant affect size remain to be identified.

Material and Methods

Ethics Statement

The study was reviewed and approved by the Kalinago council and institutional review boards of Penn
State University (29269EP), Ross University, and the Dominica Ministry of Health (H125). Informed
consent was obtained from each participant enrolled in the study, and in the case of minors, consent was
also obtained from a parent or guardian.

Recruitment

Participants from among the Kalinago populations were recruited with the help of nurses from the
Kalinago Territory in 2014. Recruitment took place throughout the territory's 8 hamlets. Place and date
of birth, reported ancestry of parents and grandparents, number of siblings, and response to sun exposure
(tanning ability, burning susceptibility) were obtained by interview. Hair color and texture and eye color
(characterized as black, brown, gray, blue, green, hazel, no pigment) were noted visually but not measured
quantitatively.

Skin Reflectometry

Skin reflectance was measured using a Datacolor CHECK[PLUS] spectrophotometer and converted to melanin
unit as we have previously described (Ang et al., 2012; Diffey et al., 1984). To minimize the confounding
effects of sun exposure and body hair, skin color measurements were measured on each participant's
inner arm, and the average of triplicate measurements was generated. Measurements at this location are
generally used as an approximation for constitutive skin pigmentation (Choe et al., 2006; Park and Lee,
2005). Before skin color measurements were taken, alcohol wipes were used to minimize the effect of dirt
and/or oil. In order to minimize blanching due to occlusion of blood from the region being measured, care
was taken not to apply only sufficient pressure to the skin to prevent ambient light from entering the
scanned area (Fullerton et al., 1996).

DNA Collection

Saliva samples were collected using the Oragene Saliva kit, and DNA was extracted using the prepIT.L2P
kit, both from DNA Genotek (Ottawa, Canada). DNA integrity was checked by agarose gel electrophoresis
and quantitated using a NanoDrop spectrophotometer (Thermo Fisher Scientific, Waltham, MA). Further
quantification was done using Qubit Fluorometer (Thermo Fisher Scientific, Waltham, MA) as needed,
following manufacturer instructions.

Genotyping

Oculocutaneous albinism variants previously identified in African and Native Americans (Carrasco et al.,

2009; King et al., 2003; Stevens et al., 1997; Yi et al., 2003) were amplified by PCR in all albino individuals

as well as control samples using published conditions. Selected alleles of *SLC24A5, SLC45A2*, and *OCA2*

were amplified in all sampled individuals as described in Table S2. Amplicons generated by 30 cycles of

PCR using an Eppendorf thermocycler were sequenced (GeneWiz, South Plainfield, NJ) and the

chromatograms viewed using Geneious software.

Illumina SNP genotyping using the Infinium Omni2.5-8 BeadChip was performed for all the individuals

sampled. This was performed in three cohorts, using slightly different versions of the array, and the results

combined. Due to ascertainment differences between the cohorts, analysis is presented here only for the

combined sample. After quality control to eliminate duplicates and monomorphic variants, and to remove

variants and individuals with genotype failure rates > 0.05, 358 Kalinago individuals and 1 638 140 unique

autosomal SNPs remained.

Whole exome sequencing of albino and obligate carriers

In order to identify the causative variant for albinism in the Kalinago, 2 samples (one albino and one

parent) were selected for whole exome sequencing. Following shearing of input DNA (1 microgram) using

a Covaris E220 Foucsed-ultrasonicator (Woburn, MA), exome enrichment and library preparation was

done using the Agilent SureSelect V5+UTR kit (Santa Clara, CA). The samples were sequenced at 50x

coverage using a HiSeq 2500 sequencer (Illumina, San Diego, CA).

The *fastq* files were aligned back to Human Reference Genome GRCh37 (HG19) using BWA(Li and Durbin,

2009) and bowtie (Langmead et al., 2009). Candidate SNP polymorphisms were identified using GATK's

UnifiedGenotyper (McKenna et al., 2010), while the IGV browser was used to examine the exons of

interest for indels (Thorvaldsdottir et al., 2013). Variants with low sequence depth (< 10) in either sample

were excluded from further consideration.

Computational analysis

Association analysis, basic statistics, and merges with other datasets were performed using plink 1.9

(Chang et al., 2015; Purcell et al., 2007). Phasing and analysis of regions of homozygosity by descent and

identity by descent were performed with Beagle 4.1 (Browning and Browning, 2013, 2007), using 1000

Genomes Project (1KGP) phased data (The 1000 Genomes Project Consortium, 2015) as reference.

The genotyped individuals were randomly partitioned into nine subsets of 50 or 51 individuals (n=50

subsets) in which no pair exhibited greater than second-order relationship (PI_HAT > 0.25 using --genome

command in plink). Using the same criteria, a maximal subset of 184 individuals was also generated (n=184

subset).

Principle components analysis (PCA) was performed using the smartpca program (version 13050) in the

eigensoft package (Price et al., 2006). For comparison to HGDP populations, Kalinago samples were

projected onto principal components calculated for the HGDP samples alone. For use as covariates in

association analyses, the n=184 subset was used to generate the PCA, and the remaining individuals were

projected onto the same axes.

Admixture analysis was performed using the ADMIXTURE program (Alexander et al., 2009; Zhou et al.,

2011). Each of the nine n=50 subsets was merged with the N=940 subset of HGDP data (Li et al., 2008;

Rosenberg, 2006) for analysis and the outputs combined.

For association analysis, we removed the three albino individuals from the analysis. In addition to the

entire remaining sample, we also analyzed the n=184 subset and each n=50 subset; the latter results were

combined using METAL (Willer et al., 2010). P-values were adjusted for statistic inflation by genomic

control (median statistic method).

Statistical analysis of pigmentary effect of albinism involved fitting parameters to an additive model for

the sample containing carriers but lacking albinos, applying the same model to the albino individuals, and

comparing residuals for the albinos and the other individuals.

Local ancestry analysis of the region containing the albinism allele was performed using the PopPhased

version of rfmix (v1.5.4) with the default window size of 0.2 cM (Maples et al., 2013). A subset of 1KGP

data served as reference haplotypes for European and African populations, and the Native American

ancestry segments of the admixed samples as determined by (Martin et al., 2017a) were combined to

generate synthetic Native American reference haplotypes.

Data Availability Statement

The whole exome sequencing and whole genome SNP genotyping data underlying this article cannot be

shared publicly due to the privacy of individuals and stipulation by the Kalinago community. The data will

be shared on reasonable request to the corresponding author.

Acknowledgments

We would like to thank the Kalinago Council, Dominica Ministry of Health, nurses at the Kalinago Territory, Salybia Mission Project and the Kalinago community for their assistance and participation in this study. We would also like to acknowledge faculty of Ross University, Portsmouth, Dominica (now Bridgetown, Barbados), especially Drs. Gerhard Meisenberg (retired) and Liris Benjamin of Ross University in helping us to obtain the necessary IRB approval for fieldwork. This work was supported by the Hershey Rotary Club, Jake Gittlen Laboratories for Cancer Research and Department of Pathology for funding portions of this project. We would also like to acknowledge members of the Cheng Lab for their constructive comments and input.

References

1000 Genomes Project Consortium. 2010. A map of human genome variation from population-
scale sequencing. *Nature* **467**:1061–73. doi:10.1038/nature09534
Adhikari K, Mendoza-Revilla J, Sohail A, Fuentes-Guajardo M, Lampert J, Chacón-Duque JC,
Hurtado M, Villegas V, Granja V, Acuña-Alonzo V, Jaramillo C, Arias W, Lozano RB, Everardo
P, Gómez-Valdés J, Villamil-Ramírez H, Silva de Cerqueira CC, Hunemeier T, Ramallo V,
Schuler-Faccini L, Salzano FM, Gonzalez-José R, Bortolini M-C, Canizales-Quinteros S, Gallo C,
Poletti G, Bedoya G, Rothhammer F, Tobin DJ, Fumagalli M, Balding D, Ruiz-Linares A. 2019.
A GWAS in Latin Americans highlights the convergent evolution of lighter skin pigmentation
in Eurasia. *Nature Communications* **10**:1–16. doi:10.1038/s41467-018-08147-0
Alexander DH, Novembre J, Lange K. 2009. Fast model-based estimation of ancestry in unrelated
individuals. *Genome Res* **19**:1655–64. doi:10.1101/gr.094052.109
Ang KC, Ngu MS, Reid KP, Teh MS, Aida ZS, Koh DX, Berg A, Oppenheimer S, Salleh H, Clyde MM,
Md-Zain BM, Canfield VA, Cheng KC. 2012. Skin color variation in Orang Asli tribes of
Peninsular Malaysia. *PLoS One* **7**:e42752. doi:10.1371/journal.pone.0042752
Barsh GS. 2003. What controls variation in human skin color? *PLoS Biol* **1**:E27.
doi:10.1371/journal.pbio.0000027
Basu Mallick C, Iliescu FM, Möls M, Hill S, Tamang R, Chaubey G, Goto R, Ho SYW, Gallego Romero
I, Crivellaro F, Hudjashov G, Rai N, Metspalu M, Mascie-Taylor CGN, Pitchappan R, Singh L,
Mirazon-Lahr M, Thangaraj K, Villems R, Kivisild T. 2013. The Light Skin Allele of SLC24A5 in
South Asians and Europeans Shares Identity by Descent. *PLoS Genet* **9**:e1003912.
doi:10.1371/journal.pgen.1003912
Beleza S, Campos J, Lopes J, Araújo II, Almada AH, Silva AC e, Parra EJ, Rocha J. 2012. The
Admixture Structure and Genetic Variation of the Archipelago of Cape Verde and Its
Implications   for   Admixture   Mapping   Studies.   *PLOS   ONE*   **7**:e51103.
doi:10.1371/journal.pone.0051103
Beleza S, Santos AM, McEvoy B, Alves I, Martinho C, Cameron E, Shriver MD, Parra EJ, Rocha J.
2013. The Timing of Pigmentation Lightening in Europeans. *Mol Biol Evol* **30**:24–35.
doi:10.1093/molbev/mss207
Benn-Torres J, Bonilla C, Robbins CM, Waterman L, Moses TY, Hernandez W, Santos ER, Bennett
F, Aiken W, Tullock T, Coard K, Hennis A, Wu S, Nemesure B, Leske MC, Freeman V, Carpten
J, Kittles RA. 2008. Admixture and population stratification in African Caribbean populations.
Ann Hum Genet **72**:90–98. doi:10.1111/j.1469-1809.2007.00398.x
Brown LA, Sofer T, Stilp AM, Baier LJ, Kramer HJ, Masindova I, Levy D, Hanson RL, Moncrieft AE,
Redline S, Rosas SE, Lash JP, Cai J, Laurie CC, Browning S, Thornton T, Franceschini N. 2017.
Admixture Mapping Identifies an Amerindian Ancestry Locus Associated with Albuminuria in
Hispanics   in   the   United   States.   *J   Am   Soc   Nephrol*   **28**:2211–2220.
doi:10.1681/ASN.2016091010
Browning BL, Browning SR. 2013. Improving the Accuracy and Efficiency of Identity-by-Descent
Detection in Population Data. *Genetics* **194**:459–471. doi:10.1534/genetics.113.150029
Browning SR, Browning BL. 2007. Rapid and Accurate Haplotype Phasing and Missing-Data
Inference for Whole-Genome Association Studies By Use of Localized Haplotype Clustering.
The American Journal of Human Genetics **81**:1084–1097. doi:10.1086/521987

Carrasco A, Forbes EM, Jeambrun P, Brilliant MH. 2009. A splice site mutation is the cause of the
high prevalence of oculocutaneous albinism type 2 in the Kuna population. *Pigment Cell &*
Melanoma Research **22**:645–647. doi:10.1111/j.1755-148X.2009.00575.x

Chang CC, Chow CC, Tellier LC, Vattikuti S, Purcell SM, Lee JJ. 2015. Second-generation PLINK:
rising to the challenge of larger and richer datasets. *GigaSci* **4**:7. doi:10.1186/s13742-015-
0047-8

Cheung KH, Miller PL, Kidd JR, Kidd KK, Osier MV, Pakstis AJ. 2000. ALFRED: a Web-accessible
allele frequency database. *Pac Symp Biocomput* 639–50.

Choe YB, Jang SJ, Jo SJ, Ahn KJ, Youn JI. 2006. The difference between the constitutive and
facultative skin color does not reflect skin phototype in Asian skin. *Skin Res Technol* **12**:68–
72. doi:10.1111/j.0909-725X.2006.00167.x

Crawford NG, Kelly DE, Hansen MEB, Beltrame MH, Fan S, Bowman SL, Jewett E, Ranciaro A,
Thompson S, Lo Y, Pfeifer SP, Jensen JD, Campbell MC, Beggs W, Hormozdiari F, Mpoloka
SW, Mokone GG, Nyambo T, Meskel DW, Belay G, Haut J, Rothschild H, Zon L, Zhou Y, Kovacs
MA, Xu M, Zhang T, Bishop K, Sinclair J, Rivas C, Elliot E, Choi J, Li SA, Hicks B, Burgess S, Abnet
C, Watkins-Chow DE, Oceana E, Song YS, Eskin E, Brown KM, Marks MS, Loftus SK, Pavan WJ,
Yeager M, Chanock S, Tishkoff S. 2017. Loci associated with skin pigmentation identified in
African populations. *Science* eaan8433. doi:10.1126/science.aan8433

Derenko M, Malyarchuk B, Grzybowski T, Denisova G, Rogalla U, Perkova M, Dambueva I,
Zakharov I. 2010. Origin and Post-Glacial Dispersal of Mitochondrial DNA Haplogroups C and
D in Northern Asia. *PLoS One* **5**. doi:10.1371/journal.pone.0015214

Dessinioti C, Stratigos AJ, Rigopoulos D, Katsambas AD. 2009. A review of genetic disorders of
hypopigmentation: lessons learned from the biology of melanocytes. *Experimental*
Dermatology **18**:741–749. doi:10.1111/j.1600-0625.2009.00896.x

Diffey BL, Oliver RJ, Farr PM. 1984. A portable instrument for quantifying erythema induced by
ultraviolet radiation. *Br J Dermatol* **111**:663–72.

Edwards M, Bigham A, Tan J, Li S, Gozdzik A, Ross K, Jin L, Parra EJ. 2010. Association of the OCA2
Polymorphism His615Arg with Melanin Content in East Asian Populations: Further Evidence
of Convergent Evolution of Skin Pigmentation. *PLOS Genetics* **6**:1–8.

Engelsen O. 2010. The Relationship between Ultraviolet Radiation Exposure and Vitamin D Status.
Nutrients **2**:482–495. doi:10.3390/nu2050482

Fullerton A, Fischer T, Lahti A, Wilhelm KP, Takiwaki H, Serup J. 1996. Guidelines for measurement
of skin colour and erythema. A report from the Standardization Group of the European
Society of Contact Dermatitis. *Contact Dermatitis* **35**:1–10.

Gao J, D'Souza L, Wetherby K, Antolik C, Reeves M, Adams DR, Tumminia S, Wang X. 2017.
Retrospective analysis in oculocutaneous albinism patients for the 2.7 kb deletion in the
OCA2 gene revealed a co-segregation of the controversial variant, p.R305W. *Cell Biosci* **7**.
doi:10.1186/s13578-017-0149-3

Gargiulo A, Testa F, Rossi S, Di Iorio V, Fecarotta S, de Berardinis T, Iovine A, Magli A, Signorini S,
Fazzi E, Galantuomo MS, Fossarello M, Montefusco S, Ciccodicola A, Neri A, Macaluso C,
Simonelli F, Surace EM. 2011. Molecular and clinical characterization of albinism in a large
cohort of Italian patients. *Invest Ophthalmol Vis Sci* **52**:1281–9. doi:10.1167/iovs.10-6091

Garrison NA, Yi Z, Cohen-Barak O, Huizing M, Hartnell LM, Gahl WA, Brilliant MH. 2004. P gene
mutations in patients with oculocutaneous albinism and findings suggestive of Hermansky-
Pudlak syndrome. *J Med Genet* **41**:e86–e86. doi:10.1136/jmg.2003.014902
Gravel S, Zakharia F, Moreno-Estrada A, Byrnes JK, Muzzio M, Rodriguez-Flores JL, Kenny EE,
Gignoux CR, Maples BK, Guiblet W, Dutil J, Via M, Sandoval K, Bedoya G, Oleksyk TK, Ruiz-
Linares A, Burchard EG, Martinez-Cruzado JC, Bustamante CD, The 1000 Genomes Project.
2013. Reconstructing Native American Migrations from Whole-Genome and Whole-Exome
Data. *PLoS Genet* **9**:e1004023. doi:10.1371/journal.pgen.1004023
Grønskov K, Brøndum-Nielsen K, Lorenz B, Preising MN. 2014. Clinical utility gene card for:
Oculocutaneous albinism. *European Journal of Human Genetics* **22**.
doi:10.1038/ejhg.2013.307
Grønskov K, Dooley CM, Østergaard E, Kelsh RN, Hansen L, Levesque MP, Vilhelmsen K, Møllgård
K, Stemple DL, Rosenberg T. 2013. Mutations in C10orf11, a Melanocyte-Differentiation
Gene, Cause Autosomal-Recessive Albinism. *The American Journal of Human Genetics*
92:415–421. doi:10.1016/j.ajhg.2013.01.006
Gronskov K, Ek J, Brondum-Nielsen K. 2007. Oculocutaneous albinism. *Orphanet J Rare Dis* **2**:43.
doi:10.1186/1750-1172-2-43
Hanel A, Carlberg C. 2020. Skin colour and vitamin D: An update. *Experimental Dermatology* **n/a**.
doi:10.1111/exd.14142
Harvey, RG, Godber, MJ, Kopéc, AC, Mourant, AE, Tills, D. 1969 Frequency of Genetic Traits in the
Caribs of Dominica. *Human Biology* **41**:342–364.
Holick MF. 1981. The cutaneous photosynthesis of previtamin D3: a unique photoendocrine
system. *J Invest Dermatol* **77**:51–8.
Hong ES, Zeeb H, Repacholi MH. 2006. Albinism in Africa as a public health issue. *BMC Public
Health* **6**:212. doi:10.1186/1471-2458-6-212
Honychurch L. 2012. The Dominica Story: A History of the Island. Macmillan.
Honychurch L. 1998. Review of The Lesser Antilles in the Age of European Expansion. *NWIG: New
West Indian Guide / Nieuwe West-Indische Gids* **72**:305–307.
Integrative Genomics Viewer (IGV): high-performance genomics data visualization and
exploration | Briefings in Bioinformatics | Oxford Academic. n.d.
https://academic.oup.com/bib/article/14/2/178/208453
Island Caribs. 2016. . *Wikipedia, the free encyclopedia*.
Jablonski NG, Chaplin G. 2000. The evolution of human skin coloration. *J Hum Evol* **39**:57–106.
doi:10.1006/jhev.2000.0403
Kalinago Territory. 2021. . *Wikipedia*.
Kamaraj B, Purohit R. 2013. Computational Screening of Disease-Associated Mutations in OCA2
Gene. *Cell Biochem Biophys* **68**:97–109. doi:10.1007/s12013-013-9697-2
Kausar T, Bhatti M, Ali M, Shaikh R, Ahmed Z. 2013. OCA5, a novel locus for non-syndromic
oculocutaneous albinism, maps to chromosome 4q24. *Clin Genet* **84**:91–93.
doi:10.1111/cge.12019
King RA, Pietsch J, Fryer JP, Savage S, Brott MJ, Russell-Eggitt I, Summers CG, Oetting WS. 2003.
Tyrosinase gene mutations in oculocutaneous albinism 1 (OCA1): definition of the
phenotype. *Hum Genet* **113**:502–513. doi:10.1007/s00439-003-0998-1

Klimentidis YC, Miller GF, Shriver MD. 2009. Genetic admixture, self-reported ethnicity, self-
estimated admixture, and skin pigmentation among Hispanics and Native Americans. *Am J*
Phys Anthropol **138**:375–383. doi:10.1002/ajpa.20945
Lamason RL, Mohideen MA, Mest JR, Wong AC, Norton HL, Aros MC, Jurynec MJ, Mao X,
Humphreville VR, Humbert JE, Sinha S, Moore JL, Jagadeeswaran P, Zhao W, Ning G,
Makalowska I, McKeigue PM, O'Donnell D, Kittles R, Parra EJ, Mangini NJ, Grunwald DJ,
Shriver MD, Canfield VA, Cheng KC. 2005. SLC24A5, a putative cation exchanger, affects
pigmentation in zebrafish and humans. *Science* **310**:1782–6. doi:10.1126/science.1116238
Langmead B, Trapnell C, Pop M, Salzberg SL. 2009. Ultrafast and memory-efficient alignment of
short DNA sequences to the human genome. *Genome Biology* **10**:R25. doi:10.1186/gb-2009-
10-3-r25
Lee S-T, Nicholls RD, Bundey S, Laxova R, Musarella M, Spritz RA. 1994. Mutations of the P Gene
in Oculocutaneous Albinism, Ocular Albinism, and Prader-Willi Syndrome Plus Albinism. *New*
England Journal of Medicine **330**:529–534. doi:10.1056/NEJM199402243300803
Li H, Durbin R. 2009. Fast and accurate short read alignment with Burrows–Wheeler transform.
Bioinformatics **25**:1754–1760. doi:10.1093/bioinformatics/btp324
Li JZ, Absher DM, Tang H, Southwick AM, Casto AM, Ramachandran S, Cann HM, Barsh GS,
Feldman M, Cavalli-Sforza LL, Myers RM. 2008. Worldwide human relationships inferred
from genome-wide patterns of variation. *Science* **319**:1100–4. doi:10.1126/science.1153717
Loomis WF. 1967. Skin-pigment regulation of vitamin-D biosynthesis in man. *Science* **157**:501–
506.
Lucotte G, Yuasa I. 2013. Near Fixation of 374l Allele Frequencies of the Skin Pigmentation Gene
SLC45A2 in Africa. *Biochem Genet* **51**:655–665. doi:10.1007/s10528-013-9595-8
Maples BK, Gravel S, Kenny EE, Bustamante CD. 2013. RFMix: A Discriminative Modeling
Approach for Rapid and Robust Local-Ancestry Inference. *The American Journal of Human*
Genetics **93**:278–288. doi:10.1016/j.ajhg.2013.06.020
Marcheco-Teruel B, Parra EJ, Fuentes-Smith E, Salas A, Buttenschøn HN, Demontis D, Torres-
Español M, Marín-Padrón LC, Gómez-Cabezas EJ, Álvarez-Iglesias V, Mosquera-Miguel A,
Martínez-Fuentes A, Carracedo Á, Børglum AD, Mors O. 2014. Cuba: Exploring the History of
Admixture and the Genetic Basis of Pigmentation Using Autosomal and Uniparental Markers.
PLOS Genetics **10**:e1004488. doi:10.1371/journal.pgen.1004488
Martin AR, Gignoux CR, Walters RK, Wojcik GL, Neale BM, Gravel S, Daly MJ, Bustamante CD,
Kenny EE. 2017a. Human Demographic History Impacts Genetic Risk Prediction across
Diverse Populations. *The American Journal of Human Genetics* **100**:635–649.
doi:10.1016/j.ajhg.2017.03.004
Martin AR, Lin M, Granka JM, Myrick JW, Liu Xiaomin, Sockell A, Atkinson EG, Werely CJ, Möller
M, Sandhu MS, Kingsley DM, Hoal EG, Liu Xiao, Daly MJ, Feldman MW, Gignoux CR,
Bustamante CD, Henn BM. 2017b. An Unexpectedly Complex Architecture for Skin
Pigmentation in Africans. *Cell* **171**:1340-1353.e14. doi:10.1016/j.cell.2017.11.015
McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, Garimella K, Altshuler D,
Gabriel S, Daly M, DePristo MA. 2010. The Genome Analysis Toolkit: A MapReduce
framework for analyzing next-generation DNA sequencing data. *Genome Res* **20**:1297–1303.
doi:10.1101/gr.107524.110

Micheletti SJ, Bryc K, Esselmann SGA, Freyman WA, Moreno ME, Poznik GD, Shastri AJ, Agee M, Aslibekyan S, Auton A, Bell R, Clark S, Das S, Elson S, Fletez-Brant K, Fontanillas P, Gandhi P, Heilbron K, Hicks B, Hinds D, Huber K, Jewett E, Jiang Y, Kleinman A, Lin K, Litterman N, McCreight J, McIntyre M, McManus K, Mozaffari S, Nandakumar P, Noblin L, Northover C, O'Connell J, Petrakovitz A, Pitts S, Shelton J, Shringarpure S, Tian C, Tung J, Tunney R, Vacic V, Wang X, Zare A, Beleza S, Mountain JL. 2020. Genetic Consequences of the Transatlantic Slave Trade in the Americas. *The American Journal of Human Genetics* **0**. doi:10.1016/j.ajhg.2020.06.012

Moreno-Mayar JV, Potter BA, Vinner L, Steinrücken M, Rasmussen S, Terhorst J, Kamm JA, Albrechtsen A, Malaspinas A-S, Sikora M, Reuther JD, Irish JD, Malhi RS, Orlando L, Song YS, Nielsen R, Meltzer DJ, Willerslev E. 2018. Terminal Pleistocene Alaskan genome reveals first founding population of Native Americans. *Nature*. doi:10.1038/nature25173

Norton HL, Kittles RA, Parra E, McKeigue P, Mao X, Cheng K, Canfield VA, Bradley DG, McEvoy B, Shriver MD. 2007. Genetic evidence for the convergent evolution of light skin in Europeans and East Asians. *Mol Biol Evol* **24**:710–22. doi:10.1093/molbev/msl203

Park J-H, Lee M-H. 2005. A Study of Skin Color by Melanin Index According to Site, Gestational Age, Birth Weight and Season of Birth in Korean Neonates. *J Korean Med Sci* **20**:105–108. doi:10.3346/jkms.2005.20.1.105

Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. 2006. Principal components analysis corrects for stratification in genome-wide association studies. *Nature Genetics* **38**:904–909. doi:10.1038/ng1847

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de Bakker PIW, Daly MJ, Sham PC. 2007. PLINK: A Tool Set for Whole-Genome Association and Population-Based Linkage Analyses. *The American Journal of Human Genetics* **81**:559–575. doi:10.1086/519795

Rajeevan H, Soundararajan U, Kidd JR, Pakstis AJ, Kidd KK. 2012. ALFRED: an allele frequency resource for research and teaching. *Nucleic Acids Research* **40**:D1010–D1015. doi:10.1093/nar/gkr924

Reich D, Patterson N, Campbell D, Tandon A, Mazieres S, Ray N, Parra MV, Rojas W, Duque C, Mesa N, García LF, Triana O, Blair S, Maestre A, Dib JC, Bravi CM, Bailliet G, Corach D, Hünemeier T, Bortolini MC, Salzano FM, Petzl-Erler ML, Acuña-Alonzo V, Aguilar-Salinas C, Canizales-Quinteros S, Tusié-Luna T, Riba L, Rodríguez-Cruz M, Lopez-Alarcón M, Coral-Vazquez R, Canto-Cetina T, Silva-Zolezzi I, Fernandez-Lopez JC, Contreras AV, Jimenez-Sanchez G, Gómez-Vázquez MJ, Molina J, Carracedo Á, Salas A, Gallo C, Poletti G, Witonsky DB, Alkorta-Aranburu G, Sukernik RI, Osipova L, Fedorova SA, Vasquez R, Villena M, Moreau C, Barrantes R, Pauls D, Excoffier L, Bedoya G, Rothhammer F, Dugoujon J-M, Larrouy G, Klitz W, Labuda D, Kidd J, Kidd K, Di Rienzo A, Freimer NB, Price AL, Ruiz-Linares A. 2012. Reconstructing Native American population history. *Nature* **488**:370–374. doi:10.1038/nature11258

Rinchik EM, Bultman SJ, Horsthemke B, Lee S-T, Strunk KM, Spritz RA, Avidano KM, Jong MTC, Nicholls RD. 1993. A gene for the mouse pink-eyed dilution locus and for human type II oculocutaneous albinism. *Nature* **361**:72–76. doi:10.1038/361072a0

Rogoziński J. 2000. A Brief History of the Caribbean: From the Arawak and Carib to the Present:

Rosenberg NA. 2006. Standardized Subsets of the HGDP-CEPH Human Genome Diversity Cell Line
Panel, Accounting for Atypical and Duplicated Samples and Pairs of Close Relatives. *Annals*
of Human Genetics **70**:841–847. doi:10.1111/j.1469-1809.2006.00285.x
Schroeder H, Sikora M, Gopalakrishnan S, Cassidy LM, Delser PM, Velasco MS, Schraiber JG,
Rasmussen S, Homburger JR, Ávila-Arcos MC, Allentoft ME, Moreno-Mayar JV, Renaud G,
Gómez-Carballa A, Laffoon JE, Hopkins RJA, Higham TFG, Carr RS, Schaffer WC, Day JS,
Hoogland M, Salas A, Bustamante CD, Nielsen R, Bradley DG, Hofman CL, Willerslev E. 2018.
Origins and genetic legacies of the Caribbean Taino. *PNAS* 201716839.
doi:10.1073/pnas.1716839115
Shriver MD, Parra EJ, Dios S, Bonilla C, Norton H, Jovel C, Pfaff C, Jones C, Massac A, Cameron N,
Baron A, Jackson T, Argyropoulos G, Jin L, Hoggart CJ, McKeigue PM, Kittles RA. 2003. Skin
pigmentation, biogeographical ancestry and admixture mapping. *Hum Genet* **112**:387–99.
doi:10.1007/s00439-002-0896-y
Soejima M, Koda Y. 2007. Population differences of two coding SNPs in pigmentation-related
genes SLC24A5 and SLC45A2. *Int J Legal Med* **121**:36–9. doi:10.1007/s00414-006-0112-z
Spritz RA, Fukai K, Holmes SA, Luande J. 1995. Frequent intragenic deletion of the P gene in
Tanzanian patients with type II oculocutaneous albinism (OCA2). *Am J Hum Genet* **56**:1320–
1323.
Stevens G, Ramsay M, Jenkins T. 1997. Oculocutaneous albinism (OCA2) in sub-Saharan Africa:
distribution of the common 2.7-kb P gene deletion mutation. *Hum Genet* **99**:523–527.
Stevens G, van Beukering J, Jenkins T, Ramsay M. 1995. An intragenic deletion of the P gene is
the common mutation causing tyrosinase-positive oculocutaneous albinism in southern
African Negroids. *Am J Hum Genet* **56**:586–591.
Tamm E, Kivisild T, Reidla M, Metspalu M, Smith DG, Mulligan CJ, Bravi CM, Rickards O, Martinez-
Labarga C, Khusnutdinova EK, Fedorova SA, Golubenko MV, Stepanov VA, Gubina MA,
Zhadanov SI, Ossipova LP, Damba L, Voevoda MI, Dipierri JE, Villems R, Malhi RS. 2007.
Beringian Standstill and Spread of Native American Founders. *PLoS ONE* **2**.
doi:10.1371/journal.pone.0000829
The 1000 Genomes Project Consortium. 2015. A global reference for human genetic variation.
Nature **526**:68–74. doi:10.1038/nature15393
Thorvaldsdottir H, Robinson JT, Mesirov JP. 2013. Integrative Genomics Viewer (IGV): high-
performance genomics data visualization and exploration. *Briefings in Bioinformatics*
14:178–192. doi:10.1093/bib/bbs017
Torres JB, Stone AC, Kittles R. 2013. An anthropological genetic perspective on Creolization in the
Anglophone Caribbean. *Am J Phys Anthropol* **151**:135–143. doi:10.1002/ajpa.22261
Torres JB, Vilar MG, Torres GA, Gaieski JB, Hernandez RB, Browne ZE, Stevenson M, Walters W,
Schurr TG, Consortium TG. 2015. Genetic Diversity in the Lesser Antilles and Its Implications
for the Settlement of the Caribbean Basin. *PLOS ONE* **10**:e0139192.
doi:10.1371/journal.pone.0139192
Tsetskhladze ZR, Canfield VA, Ang KC, Wentzel SM, Reid KP, Berg AS, Johnson SL, Kawakami K,
Cheng KC. 2012. Functional assessment of human coding mutations affecting skin
pigmentation using zebrafish. *PLoS One* **7**:e47398. doi:10.1371/journal.pone.0047398
van Geel N, Speeckaert M, Chevolet I, De Schepper S, Lapeere H, Boone B, Speeckaert R. 2013.
Hypomelanoses in Children. *J Cutan Aesthet Surg* **6**:65–72. doi:10.4103/0974-2077.112665

Vogel P, Read RW, Vance RB, Platt KA, Troughton K, Rice DS. 2008. Ocular albinism and
hypopigmentation defects in Slc24a5-/- mice. *Vet Pathol* **45**:264–79. doi:10.1354/vp.45-2-
264

Willer CJ, Li Y, Abecasis GR. 2010. METAL: fast and efficient meta-analysis of genomewide
association scans. *Bioinformatics* **26**:2190–2191. doi:10.1093/bioinformatics/btq340

Woolf CM. 2005. Albinism (OCA2) in Amerindians. *Am J Phys Anthropol* **Suppl 41**:118–140.
doi:10.1002/ajpa.20357

Yang Z, Zhong H, Chen J, Zhang X, Zhang H, Luo X, Xu S, Chen H, Lu D, Han Y, Li J, Fu L, Qi X, Peng
Yi, Xiang K, Lin Q, Guo Y, Li M, Cao X, Zhang Y, Liao S, Peng Yingmei, Zhang L, Guo X, Dong S,
Liang F, Wang J, Willden A, Aun HS, Serey B, Sovannary T, Bunnath L, Samnom H, Mardon G,
Li Q, Meng A, Shi H, Su B. 2016. A genetic mechanism for convergent skin lightening during
recent human evolution. *Mol Biol Evol* msw003. doi:10.1093/molbev/msw003

Yi Z, Garrison N, Cohen-Barak O, Karafet TM, King RA, Erickson RP, Hammer MF, Brilliant MH.
2003. A 122.5-kilobase deletion of the P gene underlies the high prevalence of
oculocutaneous albinism type 2 in the Navajo population. *Am J Hum Genet* **72**:62–72.
doi:10.1086/345380

Yuasa I, Umetsu K, Harihara S, Kido A, Miyoshi A, Saitou N, Dashnyam B, Jin F, Lucotte G,
Chattopadhyay PK, Henke L, Henke J. 2006. Distribution of the F374 allele of the SLC45A2
(MATP) gene and founder-haplotype analysis. *Ann Hum Genet* **70**:802–11.
doi:10.1111/j.1469-1809.2006.00261.x

Zhou H, Alexander D, Lange K. 2011. A quasi-Newton acceleration for high-dimensional
optimization algorithms. *Stat Comput* **21**:261–273. doi:10.1007/s11222-009-9166-3

682