Title: Resting-state fMRI-based screening of deschloroclozapine in rhesus macaques predicts dosage-

dependent behavioral effects

Authors: Atsushi Fujimoto[1]*, Catherine Elorette[1]*, J. Megan Fredericks[1], Satoka H. Fujimoto[1], Lazar

Fleysher[4], Peter H. Rudebeck[1+], and Brian E. Russ[1,2,3+y]

Author affiliations:

[1] Nash Family Department of Neuroscience and Friedman Brain Institute, Icahn School of Medicine at

Mount Sinai, One Gustave L. Levy Place, New York, NY 10029

[2] Center for Biomedical Imaging and Neuromodulation, Nathan Kline Institute, 140 Old Orangeburg Road,

Orangeburg, NY 10962

[3] Department of Psychiatry, New York University at Langone, One, 8, Park Ave, New York, NY 10016

[4] BioMedical Engineering and Imaging Institute, Icahn School of Medicine at Mount Sinai, One Gustave L.

Levy Place, New York, NY 10029

* These authors contributed equally to this work

+ Joint last author

y Corresponding author

Correspondence:

Brian E. Russ

brian.russ@mssm.edu

ABSTRACT

Chemogenetic techniques such as Designer Receptors Exclusively Activated by Designer Drugs

(DREADDs) enable transient, reversible, and minimally invasive manipulation of neural activity in vivo.

Their development in non-human primates is essential for uncovering neural circuits contributing to

cognitive functions and their translation to humans. One key issue that has delayed the development of

chemogenetic techniques in primates is the lack of an accessible drug-screening method. Here, we utilize

resting-state functional MRI (rs-fMRI), a non-invasive neuroimaging tool, to assess the impact of

deschloroclozapine (DCZ) on brain-wide resting-state functional connectivity in seven rhesus macaques

without DREADDs. We found that systemic administration of 0.1 mg/kg DCZ did not alter the resting-

state functional connectivity. Conversely, 0.3 mg/kg of DCZ was associated with a prominent increase in

functional connectivity that was mainly confined to the connections of frontal regions. Additional

behavioral tests confirmed a negligible impact of 0.1 mg/kg DCZ on socio-emotional behaviors as well as

on reaction time in a probabilistic learning task. 0.3 mg/kg DCZ did, however, slow responses in the

probabilistic learning task, suggesting attentional or motivational deficits associated with hyperconnectivity

in fronto-temporo-parietal networks. Our study highlights both the excellent selectivity of DCZ as a

DREADD actuator, and the side-effects of its excess dosage. The results demonstrate the translational value

of rs-fMRI as a drug-screening tool to accelerate the development of chemogenetics in primates.

INTRODUCTION

The development of chemogenetics such as Designer Receptors Exclusively Activated by Designer Drugs

(DREADDs) has ushered in a new era of neuroscience research. DREADDs utilize a family of modified

G-protein coupled muscarinic cholinergic receptors to modulate neural activity *in vivo* [1]. In rodents,

DREADDs have been instrumental for identifying the function of specific cell populations and neural

circuits [2,3]. The development of DREADDs in primates has been slower for a number of reasons [4], but

holds the promise of enabling the translation of chemogenetic methods to treat psychiatric and neurological

disorders.

DREADD receptors alter the membrane potential of neurons only when bound by non-endogenous ligands

[2]. Recently, deschloroclozapine (DCZ) has been identified as a selective and high-affinity activator for

muscarinic-based DREADDs with few off-target effects, unlike the original actuator clozapine-N-oxide [5-

7]. At a dose of 0.1mg/kg, DCZ occupies up to 80% of DREADD receptors *in vivo* [5], producing rapid

and robust changes in neural activity and alterations in behavior. Despite this, there are open questions that

could limit the appropriateness of DCZ for use in both basic and clinical settings.

First, it is unclear how DCZ, in the absence of DREADDs, impacts brain-wide patterns of neural activity.

This is potentially vital for discerning the more subtle effects of systemic DCZ on the brain, such as changes

in functional connectivity between brain areas, information essential for assessing the clinical

appropriateness of DCZ. Second, while 0.1 mg/kg DCZ has little impact on tasks that assess executive

function or motivation [5,8], it is unknown if the same dose causes changes in translationally relevant

affective tasks. While DCZ has negligible affinity for most endogenous receptors, it has some affinity to

muscarinic acetylcholine receptors (mAchR) and serotonin type-2 receptors that are primarily located in

the cortex [9,10] and which are potentially important for affective behaviors [5]. It is possible therefore that

even at low, but functionally relevant doses DCZ may impact affective behavior by binding to these

endogenous receptors. Addressing these issues in macaque monkeys is critical as marked changes in brain-

wide activity patterns, behavior or both would limit the usefulness of DCZ as the actuator in basic

neuroscience and in future clinical settings.

Here we addressed these questions using resting-state functional magnetic resonance imaging (rs-fMRI)

and behavioral assessments in macaque monkeys. We measured the impact of either vehicle, low-dose DCZ

(0.1 mg/kg), or high-dose DCZ (0.3 mg/kg) on whole-brain functional connectivity as well as in conditioned

and unconditioned tasks that assess socio-emotional processing. We found no changes in overall functional

connectivity or affective behavior after administration of vehicle or 0.1 mg/kg of DCZ. Conversely, higher

doses of DCZ at 0.3 mg/kg affected brain-wide patterns of functional connectivity, in particular in the

connections of frontal regions to the temporal and parietal lobes. This dose also impacted decision-making

in a probabilistic learning task, increasing reaction times. Thus, our experiments reveal the dosage-

dependent impact of DCZ on brain-wide functional connectivity and demonstrate the potential value of rs-

fMRI as an additional drug-screening tool in basic and clinical neuroscience.

METHODS AND MATERIALS

Subjects

Seven adult rhesus macaques (*Macaca mulatta*) served as subjects (6 males and 1 female). All subjects

underwent anesthetized functional MRI scans, while subsets of 4 and 3 animals were assessed in 2 socio-

emotional tasks and a probabilistic learning task, respectively (see **Supplemental Table 1**). All procedures

were reviewed and approved by the Icahn School of Medicine Animal Care and Use Committee.

Resting-state fMRI experiments

Animals were first sedated with ketamine (5 mg/kg) and dexmedetomidine (0.0125 mg/kg), and then

intubated and maintained under low level (0.7-0.8 %) isoflurane anesthesia [11]. Vehicle (2% DMSO in

saline), low-dose DCZ (0.1 mg/kg), or high-dose DCZ (0.3 mg/kg) was administered intravenously halfway

through a session. In each session, three to four runs of echo planar image (EPI) functional scans (1.6 mm

isotropic, TR/TE 2120/16 ms, flip angle 45°, 300 volumes per run) were obtained for both before ('baseline'

scans) and after drug injection ('test' scans) (**Figure 1**).

Behavioral experiments

Four animals were assessed in the human intruder task [12] and social interaction tests [13]. Vehicle or low-

dose DCZ was administered intramuscularly 30-60 minutes prior to a test. Experiments were

counterbalanced for treatment order among animals.

Three monkeys were assessed in a probabilistic learning task on a touch-sensitive monitor. In this task,

subjects learned to discriminate which visual stimulus in two pairs of novel stimuli was associated with the

highest probability (80 or 20%) of receiving a food reward. Each session consisted of 200 trials.

Data analyses

Connectome analyses were computed using 3dNetCorr function in AFNI [14,15]. We computed the

correlation between anatomical ROIs at multiple cortical [16] and subcortical [17] levels in the macaque

hierarchical atlases. The correlation matrices, or connectomes, were Fisher's z-transformed and averaged

within condition (baseline and test for: vehicle, low-dose DCZ, and high-dose DCZ). Differences between

connectomes were calculated by subtracting the baseline from test data for each drug level. We also

calculated a modified group Independent Components Analysis (ICA) using FSL's melodic ICA function

[18,19], with 30 possible components. The extracted IC scores were then submitted to a 2 (baseline or test

scan) by 3 (vehicle, low-dose DCZ, or high-dose DCZ) ANOVA which was computed on a voxel-by-voxel

basis.

Data from unconditioned socio-emotional tests were scored in Observer XT 14.0 by a condition blind scorer

and analyzed using GraphPad Prism 9.0 (GraphPad, San Diego CA). Behavioral ethograms (**Supplemental**

Tables 2 and **3**) were adapted from previous studies [20,21]. Social interaction was analyzed via paired t-

test; human intruder was analyzed via a repeated measures two-way ANOVA with factors of drug and trial

type as main factors and monkey as a random effect. The probabilistic learning task was analyzed using

MATLAB 2021a. We assessed reaction time (RT, defined as the time from stimuli presentation to a choice

response) as a measure of general task-execution processes such as attention or motivation. The effect of

drug on RT was assessed using an ANOVA with factors of drug as a main effect and monkey and session

as random effects.

Across the rs-fMRI and behavioral testing, we timed injections so that the peak of DCZ concentration in

the CSF would occur as testing began [5]. For details of experimental procedures and analyses, see

Supplemental Information.

RESULTS

The impact of low- and high-dose DCZ on brain-wide fMRI functional connectivity

Seven subjects underwent anesthetized functional MRI scanning using isoflurane levels below those shown

to adversely impact functional connectivity in frontal cortex [22] (**Figure 1**). To assess the changes in brain-

wide functional connectivity following DCZ administration, we calculated functional connectivity using

predetermined regions of interest (ROIs) based on the cortical hierarchical atlas (CHARM) [16] and

subcortical hierarchical atlas (SARM) [17] for rhesus macaques (**Figure 2A** and **C**). The z-transformed

correlation coefficients between fMRI signal time courses in all ROIs were computed for each condition,

for both the baseline (before drug injection) and test (after drug injection) scans (**Figure 2B** and **D**, left and

center columns). Then, the difference in z-value (test minus baseline) was computed for each drug condition

(**Figure 2B** and **D**, right columns). This approach allowed us to control for session-to-session variation in

fMRI signals that could influence the results. **Figure 2** shows the full matrix of correlation coefficients

from this connectome-based analysis from a single subject (monkey Bu). As expected, we found clear

patterns of correlations between distinct brain areas that are consistent to previous reports in macaques [23]

in baseline scans (**Figure 2B**, left column).

Neither vehicle (2% DMSO in saline) nor low-dose DCZ (0.1 mg/kg) administration caused a detectable

change in the overall pattern of functional connectivity and consequently there were very minor differences

in functional connectivity between test and baseline scans for these conditions (**Figure 2B**, compare top-

right and middle-right panels). Conversely, high-dose DCZ administration (0.3 mg/kg) profoundly

increased functional connectivity across cortical regions (**Figure 2B**, bottom-right panel). This effect was

most apparent in frontal areas, where, compared to baseline, correlations increased, as well as between

frontal cortex and other parts of the brain. In contrast to cortex, correlations between subcortical areas were

not impacted by the different drug levels (**Figure 2D**).

To assess the group level impact of DCZ on brain-wide connectomes, we computed correlation matrices

for individual subjects for baseline and test scans (N = 5-6 subjects per condition, see **Supplemental Table**

1). Then the z-values differences were averaged across subjects (**Figure 3**). Similar to what we saw for a

single subject (**Figure 2**), vehicle or low-dose DCZ had little impact on overall functional connectivity. In

contrast, high-dose DCZ increased functional connectivity in cortical areas but not in subcortical areas

(**Figure 3A** and **D**). Collectively, average functional connectivity was stronger in the high-dose DCZ

condition compared to vehicle or low-dose DCZ in cortical areas but not in subcortical areas (**Figure 3B**

and **E**). Two-way ANOVA (Drug: vehicle, low-dose DCZ, or high-dose DCZ × Area category: cortical or

subcortical areas) revealed a significant main effect of drug ($F_{2,13450} = 31$, p = 0.0001) and an interaction

between drug and area category ($F_{2,13450} = 8.2$, p = 0.016). Post-hoc analyses confirmed that high-dose DCZ

modulated overall functional connectivity predominantly in cortical areas (vehicle or low-dose DCZ versus

high-dose DCZ, p < 0.0001).

Next, we assessed how high-dose DCZ differentially impacted functional connectivity across different

lobes. Frontal, temporal, parietal, but not occipital cortex, showed marked changes in functional

connectivity between the high-dose DCZ and vehicle conditions (**Figure 3C**, bottom, * denotes rank-sum

test p < 0.01 with Bonferroni correction). Furthermore, the proportion of brain areas that were impacted by

low-dose DCZ was far lower than those impacted by high-dose DCZ (**Figure 3C**, top, chi-square test, $\chi^2 =$

23, p = 1.6×10^{-6}). Neither DCZ dose level affected functional connectivity in subcortical areas (**Figure**

3F), suggesting the mechanism of functional-connectivity changes was exclusive to cortical regions.

Independent-components reflect brain-wide functional-connectivity changes caused by high-dose DCZ

While informative, the prior ROI-based approach is limited by predefined anatomical boundaries and may

overlook relevant changes in MRI signals not aligned to anatomical boundaries, and does not reveal

network-level changes that includes more than two areas. Thus, we performed a data-driven independent-

component analysis (ICA) to identify 30 independent components (ICs) that are spatially coherent time

courses in the whole-brain fMRI signal. We found a variety of ICs mainly located in cortical areas (**Figure**

4A-F, top). To identify ICs, and thus regions that are affected by high-dose DCZ, a two-way ANOVA was

performed (Drug: vehicle, low-dose or high-dose DCZ × Time: baseline or test). This analysis revealed

clusters (> 60 voxels with F > 3.5) in a subset of the ICs that showed a significant main effect of drug; these

included cortical areas such as the dorso-lateral prefrontal cortex (dlPFC), anterior cingulate cortex (ACC),

posterior cingulate cortex (PCC), ventromedial cortex (vmPFC), somatosensory and motor cortices (**Figure**

4A-F, bottom). Notably, IC 3 showed an effect of drug on the fronto-temporo-parietal network, implicated

in mediating attentional processes [24] (**Figure 4A**). Further, the ICs which showed drug effects overlapped

with regions that showed increased functional connectivity in our ROI-based analyses (**Figure 3**). Systemic

administration of high-dose DCZ influenced brain-wide spatially coherent patterns of neural activity,

especially in attentionally-linked fronto-temporo-parietal circuits.

Negligible effects of low-dose DCZ on socio-emotional behaviors

Our imaging results demonstrated a negligible impact of 0.1 mg/kg DCZ on brain-wide functional

connectivity with the least impact in subcortical structures (**Figure 3**). Although no adverse effects of low-

dose DCZ were reported in a working-memory task in macaque monkeys [5,8], no study has examined the

impact of DCZ on the socio-emotional responses of monkeys. To address this issue, a subset of 4 subjects

were assessed in the human intruder task, a well-validated socio-emotional task for non-human primates

[12,13]. Subjects received a I.M. injection of either low-dose DCZ or vehicle prior to testing. We analyzed

the time spent motionless on each trial type (alone, profile, or stare condition) as a measure of anxiety-like

behavior. We expected that the animals spent more time motionless in the 'profile' condition in response to

indirect threat compared to the 'alone' or directly threatening 'stare' condition [12]. Animals did spend more

time motionless in the 'profile' condition in both vehicle and low-dose DCZ conditions (**Figure 5A**). A

two-way ANOVA matched for subject (Trial type: alone, profile or stare × Drug: vehicle or low-dose DCZ)

revealed a significant main effect of trial type ($F_{1,3} = 16$, $p = 0.049$), but no main effect of drug ($F_{1,3} = 0.24$,

$p = 0.66$) nor an interaction between trial type and drug ($F_{2,6} = 1.4$, $p = 0.32$). We also analyzed the

frequency of anxious and hostile behaviors during the threatening stare condition. There were no differences

between low-dose DCZ and vehicle for either anxiety behaviors (Mean ± SD, vehicle: 10.5 ± 9.6 and low-

dose DCZ: 24.8 ± 35.7, respectively; $t = 0.68$, $p = 0.54$, paired t-test) or hostile behaviors (Mean ± SD,

vehicle: 28.3 ± 32.1 and low-dose DCZ: 37.5 ± 43.4, respectively; $t = 1.49$, $p = 0.23$, paired t-test).

We also examined whether low-dose DCZ alters social interaction. Subjects were paired with a cagemate,

and the total social contact, comprising grooming, play, and passive physical contact, was measured [13].

There were no statistical differences between vehicle and low-dose DCZ conditions on these measures

(Mean ± SD, vehicle: 435.4 ± 198.2 s and low-dose DCZ: 502.1 ± 342.3 s, respectively; $t = 0.26$, $p = 0.81$,

paired t-test). Thus, we found no evidence that macaque socio-emotional behaviors were affected by

systemic administration of low-dose DCZ.

Effect of high-dose DCZ on decision-making and its relation to resting-state functional connectivity

Our imaging results revealed an increase in overall functional connectivity in cortical areas following a

systemic administration of high-dose DCZ (0.3 mg/kg) (**Figures 3** and **4**). Because the changes occurred

mainly in frontal, temporal and parietal cortex, it likely influences behaviors dependent on functional

interaction between these areas (**Figure 4**). We trained a subset of 3 monkeys in a translationally-relevant

probabilistic learning task that relies on frontal and temporal cortices [25-27]. We theorized that high-dose

DCZ would specifically increase monkeys' reaction time (RT) as the drug slows decision-making. RTs did

increase after high-dose DCZ but not low-dose DCZ, compared to vehicle (**Figure 5B**). A one-way repeated

measures ANOVA (Drug: vehicle, low-dose DCZ or high-dose DCZ) revealed a significant main effect of

drug ($F_{2,49} = 17$, $p = 2.5 \times 10^{-6}$), suggesting that high-dose DCZ causes an impairment in general task-

execution processes, while low-dose DCZ did not. Based on these results, we decided not to include high-

dose DCZ in the socioemotional tasks, as an impairment in task execution or attention could compromise

the interpretation of the results.

To analyze the relationship between the neural and behavioral effects following systemic DCZ, we

correlated overall functional-connectivity changes in cortical (**Figure 5C**, left) or subcortical areas (**Figure**

5C, right) with RT in the 3 animals. There was a significant positive correlation between RT and cortical

functional connectivity (n = 58, p = 0.023) but no correlation with subcortical areas (n = 58, p = 0.49),

suggesting that cortical resting-state changes following DCZ administration were related to the impairment

in task-execution process observed after high-dose DCZ.

DISCUSSION

To assess the impact of DCZ on brain-wide function in primates, we conducted a resting-state fMRI (rs-

fMRI) experiments and behavioral assessments in macaque monkeys. We found that systemic

administration of low-dose DCZ, 0.1 mg/kg, did not affect overall functional connectivity or the socio-

emotional behaviors we measured. However, we did find that systemic administration of high-dose DCZ,

0.3 mg/kg, increased overall cortical functional connectivity, and was associated with slowed responses in

a probabilistic learning task. Together, our data indicate that at a functionally effective dose of 0.1 mg/kg

DCZ appears to be appropriate for use as a DREADD actuator in non-human primates, a key finding if

chemogenetics are to be translated further.

Negligible impact of low-dose DCZ on brain-wide functional connectivity or socio-emotional behaviors

Although DCZ was identified as a potent, selective, and metabolically stable DREADD actuator [5], prior

work only showed its local effects on the brain using electrophysiology and [18F] FDG-PET imaging [5].

What was not clear is if it impacts brain-wide neural activity in normal controls where DREADD receptors

are not present. Consequently, we conducted an rs-fMRI experiment with seven rhesus monkeys and

assessed the effect of different dose levels of DCZ. Animals were only lightly anesthetized throughout the

scan sessions so that the resting-state brain activity was preserved as closes to that of the awake state as

possible [11,22]. We hypothesized that low-dose DCZ, which is functionally effective in chemogenetic

experiments in non-human primates [5,28], would not affect resting-state functional connectivity in non-

transfected animals. As we expected, low-dose DCZ did not alter brain-wide functional connectivity either

in cortical or subcortical areas (**Figures 2** and **3**).

Although DCZ has a high-degree of selectivity to DREADD receptors, it shows some affinity to

endogenous muscarinic acetylcholine receptors (mAchR) and serotonin receptors [5]. Past studies

demonstrated a negligible effect of low-dose DCZ on performance in a working-memory task in control

monkeys [5,7,8], suggesting that this dose of DCZ does not affect working memory performance or

attention, processes linked to mAchR [29]. In the current study, we assessed socio-emotional functions of

monkeys in the human intruder and social interaction tasks [12,13], which are likely mediated by the

serotonergic system [30-33]. We asked whether even subtle agonism to endogenous serotonin receptors by

DCZ might impact unconditioned affective behaviors. Our results indicate that systemic low-dose DCZ did

not modify affective behaviors of the monkeys (**Figure 5**). In summary, there is a negligible effect of low-

dose DCZ on socio-emotional functions or brain-wide functional connectivity.

Functional-connectivity changes following high-dose DCZ administration

Systemic administration of high-dose, 0.3 mg/kg, DCZ occupies nearly 100% of DREADD receptors when

administered in monkeys expressing DREADDs in their brain [5]. The question remained as to whether

high-dose DCZ also causes significant non-specific binding of endogenous receptors leading to neural and

behavioral alterations. Indeed, DCZ at this dosage reduces performance in a working-memory task,

suggesting that the excess DCZ affects neural activity [8]. Our imaging results revealed that high-dose DCZ

caused a prominent increase in overall resting-state functional connectivity between frontal, temporal, and

parietal lobes that was not observed for vehicle or low-dose DCZ (**Figures 2** and **3**). Notably, the changes

were mainly observed in networks that include frontal regions, and no obvious effects were observed in

occipital cortex or subcortical structures. This coincides with the pattern of serotonin type-2 receptor (5-

HT$_{2A}$, 5-HT$_{2C}$) expression in the brain [34,35], suggesting that excess DCZ may alter brain function through

binding to endogenous serotonin receptors.

Behavioral changes following high-dose DCZ administration were associated with functional connectivity

changes

While low-dose DCZ has little impact on behavior, higher dose levels were associated with behavioral

changes. Specifically, high-dose DCZ significantly increased RT in the probabilistic learning task

compared to other drug conditions, indicating an impairment in general task-execution processes, such as

attention (**Figure 5**). Our ICA results revealed a prominent drug effect in a fronto-temporo-parietal network

(IC 3, **Figure 4**), heavily linked to attentional processes [24]. Alternatively, high-dose DCZ may increase

RT by affecting motivation; this seems less likely, as motivation is largely mediated by subcortical circuits

[36].

Past studies have demonstrated that patterns of rs-fMRI functional connectivity in humans are directly

related to subsequent behavioral measures recorded in separate sessions [37,38]. Consistently, we found

that the changes in cortical functional connectivity and RT were positively correlated (**Figure 5**). This

increase in cortical functional connectivity mirrors the slowing of responses may seem counterintuitive as

one could expect an improvement in task performance as a result of increased functional connectivity.

However, resting-state fMRI functional connectivity reflects low-frequency fluctuations of neural activity

between brain regions, and broad, multi-region, increases in functional connectivity do not necessarily

benefit coordinated activity across distributed areas [39]. Our data suggests that hyperconnectivity in frontal

regions could underlie the impairments in attention that are prominent in neurological and psychiatric

disorders [40-42].

Resting-state MRI as a tool for benchmarking in chemogenetic experiments and drug screening

Primate chemogenetic technique development has been relatively slow for a number of reasons, including

the limited number of animals available for development and physiological differences between rodents

and primates that hampers viral delivery [4,43]. Although PET imaging is useful for the quantification of

chemogenetic receptor expression *in vivo* [5,44-46], the more accessible rs-fMRI should play a

complementary role. Here we show that rs-fMRI provides insight into the correlated patterns of neural

activity across distinct brain areas and how this can be impacted by pharmacological agents, a level of

analysis unavailable with PET. Thus, rs-fMRI has advantages for screening chemogenetic actuators in naive

animals without specific task training. Moreover, rs-fMRI is potentially useful for "benchmarking" the

selectivity and potency of novel chemogenetic actuators [47], accelerating the development of

chemogenetics in primates.

Conclusion

The present study demonstrated that DCZ is inert in non-DREADD monkeys both in brain-wide activity

and in conditioned and unconditioned behaviors, if the dosage is carefully controlled. We also highlight a

link between resting-state hyperconnectivity and behavioral impairments, providing a neural mechanism

underlying the side-effects caused by excess DCZ. Our data underscore the value of fMRI to accelerate the

development of chemogenetics in primates, research that is essential for the translation of these approaches

to treat human psychiatric disorders.

ACKNOWLEDGEMENTS

AF, CE, BER and PHR are supported by grants from NIMH and the BRAIN initiative (R01MH110822 and

R01MH117040). BER is supported by grants from NIMH (R01MH111439) and NINDS (R01NS109498).

AF is supported by Overseas Research Fellowship from Takeda Science Foundation and a Brain &

Behavior Research Foundation Young Investigator grant (#28979). We would like to thank Dr Paula

Croxson for providing the foundation on which this work was built and Jairo Munoz and Niranjana

Bienkowska for assistance with data acquisition. For help with fMRI data pre-processing and analysis we

thank Drs Paul Taylor and Alex Franco, respectively.

FUNDING AND DISCLOSURE

314   The authors have nothing to disclose.

AUTHOR CONTRIBUTIONS

317   AF, CE, PHR and BER developed the experimental design and ideas for the research; AF, CE, JMF, SHF

318   and LF performed the research; AF, CE and BER analyzed the data; AF, CE, PHR and BER wrote the

319   manuscript; all authors edited the manuscript.

REFERENCES

1   Armbruster BN, Li X, Pausch MH, Herlitze S, Roth BL. Evolving the lock to fit the key to create

a family of G protein-coupled receptors potently activated by an inert ligand. Proceedings of the

National Academy of Sciences of the United States of America. 2007;104(12):5163-8.

2   Roth BL. DREADDs for Neuroscientists. Neuron. 2016;89(4):683-94.

3   Stachniak TJ, Ghosh A, Sternson SM. Chemogenetic synaptic silencing of neural circuits

localizes a hypothalamus→midbrain pathway for feeding behavior. Neuron. 2014;82(4):797-808.

4   Fredericks JM, Fujimoto A, Rudebeck PH. Trust, but verify: A cautionary tale of translating

chemogenetic methods (A commentray on Galvan et al). The European journal of neuroscience.

2019;50(5):2751-54.

5   Nagai Y, Miyakawa N, Takuwa H, Hori Y, Oyama K, Ji B, et al. Deschloroclozapine, a potent

and selective chemogenetic actuator enables rapid neuronal and behavioral modulations in mice

and monkeys. Nature neuroscience. 2020;23(9):1157-67.

6   Gomez JL, Bonaventura J, Lesniak W, Mathews WB, Sysa-Shah P, Rodriguez LA, et al.

Chemogenetics revealed: DREADD occupancy and activation via converted clozapine. Science

(New York, NY). 2017;357(6350):503-07.

7   Roseboom PH, Mueller SAL, Oler JA, Fox AS, Riedel MK, Elam VR, et al. Evidence in primates

supporting the use of chemogenetics for the treatment of human refractory neuropsychiatric

disorders. Molecular therapy : the journal of the American Society of Gene Therapy.

2021;29(12):3484-97.

8   Upright NA, Baxter MG. Effect of chemogenetic actuator drugs on prefrontal cortex-dependent

working memory in nonhuman primates. Neuropsychopharmacology : official publication of the

American College of Neuropsychopharmacology. 2020;45(11):1793-98.

9   Barnes NM, Sharp T. A review of central 5-HT receptors and their function. Neuropharmacology.

1999;38(8):1083-152.

10   Varastet M, Brouillet E, Chavoix C, Prenant C, Crouzel C, Stulzaft O, et al. In vivo visualization

of central muscarinic receptors using [11C]quinuclidinyl benzilate and positron emission

tomography in baboons. European journal of pharmacology. 1992;213(2):275-84.

11   Wu TL, Mishra A, Wang F, Yang PF, Gore JC, Chen LM. Effects of isoflurane anesthesia on

resting-state fMRI signals and functional connectivity within primary somatosensory cortex of

monkeys. Brain and behavior. 2016;6(12):e00591.

12   Kalin NH, Shelton SE. Defensive behaviors in infant rhesus monkeys: environmental cues and

neurochemical regulation. Science (New York, NY). 1989;243(4899):1718-21.

13   Kling A, Cornell R. Amygdalectomy and social behavior in the caged stump-tailed macaque

(Macaca speciosa). Folia primatologica; international journal of primatology. 1971;14(3):190-

208.

14   Cox RW. AFNI: software for analysis and visualization of functional magnetic resonance

neuroimages. Comput Biomed Res. 1996;29(3):162-73.

15   Taylor PA, Saad ZS. FATCAT: (an efficient) Functional and Tractographic Connectivity

Analysis Toolbox. Brain connectivity. 2013;3(5):523-35.

16   Jung B, Taylor PA, Seidlitz J, Sponheim C, Perkins P, Ungerleider LG, et al. A comprehensive

macaque fMRI pipeline and hierarchical atlas. NeuroImage. 2021;235:117997.

17   Hartig R, Glen D, Jung B, Logothetis NK, Paxinos G, Garza-Villarreal EA, et al. The Subcortical

Atlas of the Rhesus Macaque (SARM) for neuroimaging. NeuroImage. 2021;235:117996.

18   Beckmann CF, Mackay CE, Filippini N, Smith SM. Group comparison of resting-state FMRI

data using multi-subject ICA and dual regression. NeuroImage. 2009;47(Suppl 1):S148.

19   Beckmann CF, Smith SM. Probabilistic independent component analysis for functional magnetic

resonance imaging. IEEE transactions on medical imaging. 2004;23(2):137-52.

20   Raper J, Murphy L, Richardson R, Romm Z, Kovacs-Balint Z, Payne C, et al. Chemogenetic

Inhibition of the Amygdala Modulates Emotional Behavior Expression in Infant Rhesus

Monkeys. eNeuro. 2019;6(5).

21   Forcelli PA, Wellman LL, Malkova L. Blockade of glutamatergic transmission in the primate

basolateral amygdala suppresses active behavior without altering social interaction. Behavioral

neuroscience. 2017;131(2):192-200.

22   Giacometti C, Dureux A, Autran-Clavagnier D, Wilson CRE, Sallet J, Dirheimer M, et al. Frontal

Cortical Functional Connectivity Is Impacted by Anaesthesia in Macaques. Cerebral cortex (New

York, NY : 1991). 2021.

23   Grayson DS, Bliss-Moreau E, Machado CJ, Bennett J, Shen K, Grant KA, et al. The Rhesus

Monkey Connectome Predicts Disrupted Functional Networks Resulting from Pharmacogenetic

Inactivation of the Amygdala. Neuron. 2016;91(2):453-66.

24   Sani I, McPherson BC, Stemmann H, Pestilli F, Freiwald WA. Functionally defined white matter

of the macaque monkey brain reveals a dorso-ventral attention network. eLife. 2019;8.

25   Pizzagalli DA, Iosifescu D, Hallett LA, Ratner KG, Fava M. Reduced hedonic capacity in major

depressive disorder: evidence from a probabilistic reward task. Journal of psychiatric research.

2008;43(1):76-87.

26   Rudebeck PH, Saunders RC, Lundgren DA, Murray EA. Specialized Representations of Value in

the Orbital and Ventrolateral Prefrontal Cortex: Desirability versus Availability of Outcomes.

Neuron. 2017;95(5):1208-20.e5.

27   Chau BK, Sallet J, Papageorgiou GK, Noonan MP, Bell AH, Walton ME, et al. Contrasting Roles

for Orbitofrontal Cortex and Amygdala in Credit Assignment and Learning in Macaques. Neuron.

2015;87(5):1106-18.

28   Hirabayashi T, Nagai Y, Hori Y, Inoue KI, Aoki I, Takada M, et al. Chemogenetic sensory fMRI

reveals behaviorally relevant bidirectional changes in primate somatosensory network. Neuron.

2021;109(20):3312-22.e5.

29   Kuczewski N, Aztiria E, Gautam D, Wess J, Domenici L. Acetylcholine modulates cortical

synaptic transmission via different muscarinic receptors, as studied with receptor knockout mice.

The Journal of physiology. 2005;566(Pt 3):907-19.

30   Broekkamp CL, Berendsen HH, Jenck F, Van Delft AM. Animal models for anxiety and response

to serotonergic drugs. Psychopathology. 1989;22 Suppl 1:2-12.

31   Weinberg-Wolf H, Chang SWC. Differences in how macaques monitor others: Does serotonin

play a central role? Wiley interdisciplinary reviews Cognitive science. 2019;10(4):e1494.

32   Santangelo AM, Sawiak SJ, Fryer T, Hong Y, Shiba Y, Clarke HF, et al. Insula serotonin 2A

receptor binding and gene expression contribute to serotonin transporter polymorphism anxious

phenotype in primates. Proceedings of the National Academy of Sciences of the United States of

America. 2019;116(29):14761-68.

33   Santangelo AM, Ito M, Shiba Y, Clarke HF, Schut EH, Cockcroft G, et al. Novel Primate Model

of Serotonin Transporter Genetic Polymorphisms Associated with Gene Expression, Anxiety and

Sensitivity to Antidepressants. Neuropsychopharmacology : official publication of the American

College of Neuropsychopharmacology. 2016;41(9):2366-76.

34   Beliveau V, Ganz M, Feng L, Ozenne B, Højgaard L, Fisher PM, et al. A High-Resolution In

Vivo Atlas of the Human Brain's Serotonin System. The Journal of neuroscience : the official

journal of the Society for Neuroscience. 2017;37(1):120-28.

35   Mengod G, Vilaró MT, Cortés R, López-Giménez JF, Raurich A, Palacios JM. Chemical

neuroanatomy of 5-HT receptor subtypes in the mammalian brain. The serotonin receptors.

2006:319-64.

36   Robbins TW, Everitt BJ. Neurobehavioural mechanisms of reward and motivation. Current

opinion in neurobiology. 1996;6(2):228-36.

37   Kohno M, Morales AM, Ghahremani DG, Hellemann G, London ED. Risky decision making,

prefrontal cortex, and mesocorticolimbic functional connectivity in methamphetamine

dependence. JAMA psychiatry. 2014;71(7):812-20.

38   Li N, Ma N, Liu Y, He XS, Sun DL, Fu XM, et al. Resting-state functional connectivity predicts

impulsivity in economic decision-making. The Journal of neuroscience : the official journal of the

Society for Neuroscience. 2013;33(11):4886-95.

39   Hillary FG, Grafman JH. Injured Brains and Adaptive Networks: The Benefits and Costs of

Hyperconnectivity. Trends in cognitive sciences. 2017;21(5):385-401.

40   Di Martino A, Kelly C, Grzadzinski R, Zuo XN, Mennes M, Mairena MA, et al. Aberrant striatal

functional connectivity in children with autism. Biological psychiatry. 2011;69(9):847-56.

41   Siegel JS, Ramsey LE, Snyder AZ, Metcalf NV, Chacko RV, Weinberger K, et al. Disruptions of

network connectivity predict impairment in multiple behavioral domains after stroke. Proceedings

of the National Academy of Sciences of the United States of America. 2016;113(30):E4367-76.

42   Zhou J, Greicius MD, Gennatas ED, Growdon ME, Jang JY, Rabinovici GD, et al. Divergent

network connectivity changes in behavioural variant frontotemporal dementia and Alzheimer's

disease. Brain : a journal of neurology. 2010;133(Pt 5):1352-67.

43   Albaugh DL, Smith Y, Galvan A. Comparative analyses of transgene expression patterns after

intra-striatal injections of rAAV2-retro in rats and rhesus monkeys: A light and electron

microscopic study. The European journal of neuroscience. 2020;52(12):4824-39.

44   Bonaventura J, Eldridge MAG, Hu F, Gomez JL, Sanchez-Soto M, Abramyan AM, et al. High-

potency ligands for DREADD imaging and activation in rodents and monkeys. Nature

communications. 2019;10(1):4627.

45   Nagai Y, Kikuchi E, Lerchner W, Inoue KI, Ji B, Eldridge MA, et al. PET imaging-guided

chemogenetic silencing reveals a critical role of primate rostromedial caudate in reward

evaluation. Nat Commun. 2016;7:13605.

46   Roseboom PH, Mueller SAL, Oler JA, Fox AS, Riedel MK, Elam VR, et al. Evidence in primates

supporting the use of chemogenetics for the treatment of human refractory neuropsychiatric

disorders. Molecular therapy : the journal of the American Society of Gene Therapy. 2021.

47   Milham M, Petkov C, Belin P, Hamed SB, Evrard H, Fair D, et al. Toward next-generation

primate neuroscience: A collaboration-based strategic plan for integrative neuroimaging. Neuron.

2021.

FIGURE LEGENDS

Figure 1. Schematic of experiments. (**A**) An imaging session. Subjects were lightly anesthetized with

isoflurane gas throughout a session. A contrast agent (MION, 10 mg/kg) was administered intravenously

before functional scans. Drug (vehicle, 0.1 mg/kg DCZ or 0.3 mg/kg DCZ) was prepared and systemically

administered (i.v.) following baseline scans, and test scans were acquired after 15 min of drug injection.

(**B**) Data analysis pipeline. The scan data for each session initially went through preprocessing procedures.

Then seed-based analysis was performed using predetermined ROIs based on the CHARM and SARM

atlases. The temporally-concatenated scan data across all sessions was used for ICA.

Figure 2. fMRI based connectome changes after DCZ administration in an example subject. (**A**) CHARM

atlas level 3 visualized on an inflated brain (medial view). (**B**) Brain-wide functional connectivity in cortical

areas. Confusion matrices represent connectome in baseline scans (left column), test scans (center column),

and the difference between test and baseline (right column) for either in the vehicle (top), low-dose DCZ

(middle), and high-dose DCZ condition (bottom), respectively. Color indicates z-transformed correlation

coefficients. Labels on the right side show ROIs from CHARM atlas. (**C**) SARM atlas level 3 visualized

on an inflated brain. (**D**) Confusion matrices for subcortical areas with labels for SARM ROIs on the right

side. Conventions are as in *B*.

Figure 3. Group level changes in the fMRI-based connectome following DCZ administration. (**A-C**)

Functional connectivity changes in cortical areas. (**A**) Confusion matrices represent differences in z-value

(test - baseline) for vehicle (top), low-dose DCZ (middle), and high-dose DCZ (bottom) conditions. Labels

on the right side show ROIs from CHARM atlas (level 3). (**B**) Averaged functional connectivity. Bars

represent averaged z-value differences for vehicle, low-dose DCZ, and high-dose DCZ conditions. Error

bars are standard error. Symbols represent subjects. Asterisk indicates a significant interaction of drug by

area category (p = 0.016, two-way ANOVA). (**C**) Functional connectivity in each area. Bars indicate

averaged z-value differences averaged for each area of the CHARM atlas. Colors indicate frontal (red),

parietal (green), temporal (cyan), and occipital areas (yellow) for low-dose (top panel) and high-dose

(bottom panel) DCZ conditions. Gray bars indicate vehicle condition. Error bars are standard error.

Asterisks indicate significant difference between vehicle and DCZ conditions (p < 0.01 with Bonferroni

correction, rank-sum test). (**D-F**) Functional connectivity changes in subcortical areas. Conventions are as

in *A-C*. Colors in *F* indicate telencephalon (dark red), diencephalon (dark green), mesencephalon (dark

blue), metencephalon (dark yellow), and myelencephalon areas (magenta).

Figure 4. Independent component analysis of the effect of high-dose DCZ. (**A-E**) Top panels show

independent components (ICs) as correlation coefficients (*r*) on inflated brains. Bottom panels indicate

clusters that showed a significant main effect of drug condition ($p < 0.05$, two-way ANOVA).

Figure 5. The impact of systemic DCZ administration on socio-emotional behavior and probabilistic

learning. (**A**) The effect of low-dose DCZ in the human intruder task. Bars indicate the mean time spent

motionless with systemic administration of vehicle (gray) or low-dose DCZ (light blue) either in alone

condition (left panel), profile condition (center panel), or stare condition (right panel). Symbols indicate

individual animals. Error bars represent standard error. (**B**) Dosage-dependent changes in reaction time

(RT) in the probabilistic learning task. Bar plots represent the mean RT of three subjects in each drug

condition (vehicle, low-dose DCZ, high-dose DCZ). Error bars are standard error. Symbols represent

individual subjects. Asterisk indicates a significant main effect of drug condition ($p = 2.5 \times 10^{-6}$, one-way

repeated-measures ANOVA). (**C**) Relationship between RT and functional connectivity across conditions

in the probabilistic learning task. Scatter plots show the RT in each behavioral session (y axis) and the

change in overall functional connectivity in a corresponding subject (x axis) for cortical areas (left panel)

and for subcortical areas (right panel), respectively. Individual monkeys completed 6-8 behavioral sessions

per each condition. The colors represent drug conditions. The solid and dotted lines are linear fitting and

confidence intervals, and red lines indicate significant linear correlation between RT and z-value ($p =$

0.023).

A

Intubation
0.7-0.9 % isoflurane

Baseline EPIs
(3-4 runs)

Test EPIs
(3-4 runs)

MION (i.v.)

DCZ or Vehicle (i.v.)

B

Data

Run 1 - 4 (DCZ 0.1)
Run 1 - 4 (DCZ 0.3)
Run 1 - 4 (Vehicle)
Subject A

Run 1 - 3 (DCZ 0.3)
Run 1 - 3 (Vehicle)
Subject B

Run 1 - 4 (DCZ 0.3)
Run 1 - 4 (Vehicle)
Subject C

Preprocessing

Slice-timing correction
Registration and Normalization
Alignment and Motion correection
Smoothing
Masking and Scaling

Seed-based analysis
(Macaque atlas)

Independent component analysis (ICA)

Temporally-concatenated data = Estimated time courses * Independent components

Figure 1

Figure 2

Figure 3

A *IC 3: dlPFC and TE*

B *IC 20: dACC and SMA*

C *IC 6: vmPFC*

D *IC 10: PCC*

E *IC 1: Somatosensory*

F *IC 17: Sesorimotor*

Figure 4

Figure 5

SUPPLEMENTAL INFORMATION

Title: Resting-state fMRI-based screening of deschloroclozapine in rhesus macaques predicts dosage-

dependent behavioral effects

Authors: Atsushi Fujimoto[1]*, Catherine Elorette[1]*, J. Megan Fredericks[1], Satoka H. Fujimoto[1], Lazar

Fleysher[4], Peter H. Rudebeck[1+], and Brian E. Russ[1,2,3+y]

Author affiliations:

[1] Nash Family Department of Neuroscience and Friedman Brain Institute, Icahn School of Medicine at

Mount Sinai, One Gustave L. Levy Place, New York, NY 10029

[2] Center for Biomedical Imaging and Neuromodulation, Nathan Kline Institute, 140 Old Orangeburg

Road, Orangeburg, NY 10962

[3] Department of Psychiatry, New York University at Langone, One, 8, Park Ave, New York, NY 10016

[4] BioMedical Engineering and Imaging Institute, Icahn School of Medicine at Mount Sinai, One Gustave

L. Levy Place, New York, NY 10029

* These authors contributed equally to this work

+ Joint last author

y Corresponding author

Correspondence:

Brian E. Russ

brian.russ@mssm.edu

SUPPLEMENTAL METHODS

Drug preparation

To prepare a DCZ solution for fMRI or socio-emotional testing, DCZ powder (HY-42110, MedChemExpress) was dissolved in dimethyl sulfoxide (DMSO, 2% of total volume) and then diluted in saline to a concentration of 0.1 mg/kg or 0.3 mg/kg, with a total volume of 1ml. Vehicle solution was prepared as 2% DMSO diluted in 1ml saline. One vehicle scan was conducted without DMSO (monkey Wo). Both solutions underwent a sterilized filter (0.22 μm) before injection. DCZ or vehicle solution was prepared within 30 min of usage.

For the probabilistic learning testing, both vehicle and DCZ were prepared using sterile procedures and previously published drug preparation approaches for behavioral testing in macaques [1]. The vehicle consisted of 0.5% Acetic acid, 50% 1 M Sodium Acetate, and 49.5% 0.2 M NaOH. Each solution of DCZ or vehicle ultimately had a pH between 5.5 and 6.0. When DCZ was administered, the drug was first dissolved in acetic acid and sodium acetate and then diluted with NaOH. The vehicle remained consistent across both drug doses (0.1mg/kg, or 0.3mg/kg of DCZ). However, the concentration of each dose was altered so the total volume of each injection was the same no matter the dose, e.g. a 0.1mg/kg dose had a concentration of 1 mg/mL, while a 0.3 mg/kg dose had a concentration of 3 mg/mL. As with fMRI testing, DCZ or vehicle solution was prepared within 30 min of usage.

fMRI data acquisition

Animals were first sedated with ketamine (5 mg/kg) and dexmedetomidine (0.0125 mg/kg) at least 1.5 hours before the data collection to prevent detrimental effects of ketamine on neural activities. They were then intubated and maintained under low level (0.7-0.8 %) isoflurane throughout the session to ensure preservation of resting-state networks [2]. During all scanning sessions, anesthetized macaques were contained within a primate chair in the sphinx position with their heads supported by a sling. Because animals were not held in a stereotactic frame, this meant we were able to maintain sedation with lower levels of anesthesia than is often used. To minimize the effect of physiological changes in the neural activity, vital signs (end-tidal CO2, body temperature, blood pressure, capnograph) were continuously

monitored and maintained as steadily as possible throughout an experimental session. Structural and

functional scans were collected each session, as well as pre-injection (baseline) and post-injection (DCZ

or vehicle) functional scans. First, a set of setup scans were acquired which included shimming based off

the acquired fieldmap. Additionally, a 3D T1-weighted image (0.5 mm isotropic, TR/TE 2500/2.81 ms,

flip angle 8°) was acquired. Following MION injection (i.v.) [3,4], three to four runs of echo planar image

(EPI) functional scans (1.6 mm isotropic, TR/TE 2120/16 ms, flip angle 45°, 300 volumes per each run)

were obtained for baseline functional resting states. Then, approximately 30 minutes from the beginning

of the pre-injection rs-fMRI data collection, either DCZ (0.1 or 0.3 mg/kg) or vehicle was administered

(i.v.), and subsequently an additional three to four functional runs (with the same parameters) were

collected. Test EPI data were collected approximately 15 minutes after DCZ or vehicle injection to ensure

trafficking of the drug to the brain [5].

Unconditioned socio-emotional tasks

Four animals (Wo, He, La, Ha) were tested on two well validated behavioral tasks that assess socio-

emotional function: the human intruder [6] and social interaction tests [7]. Behavioral tests were

conducted following injections of either vehicle (2% DMSO in saline) or 0.1 mg/kg DCZ. Injections were

given intramuscularly (i.m.) 30-60 minutes before behavioral testing in keeping with the known

pharmacokinetics of DCZ (Nagai et al. 2020). Three of the animals received vehicle treatments before

low dose DCZ treatments, and a fourth animal received low dose DCZ treatment before vehicle. Each

animal was tested only once in each drug condition for each behavioral test. All behavioral data were

scored offline in Observer XT version 14.0 (Noldus, Wageningen, the Netherlands).

<u>Human Intruder:</u> Animals were tested on a human intruder paradigm, which has been shown to accurately

assess anxiety behaviors in macaques [6,8-10]. Testing was conducted in a Wisconsin General Testing

Apparatus (WGTA). The sliding doors of the WGTA were left open so the animal could observe the room

throughout the test session. We have adapted our testing procedure and behavioral ethogram from a

previous study [11].

The paradigm consisted of three conditions (alone, profile, stare) presented in the same order to all monkeys

for a total duration of 30 min. The experimenter wore a rubber mask depicting a different human male face

during each testing session, presented in pseudorandom order to subjects with no repetitions of masks per

subject. Different colored curtains hung behind the experimenter's seat were also used to increase novelty.

Within each testing session, monkeys were first left alone to observe the room and acclimatize to the

environment (alone condition). Next, the experimenter, wearing a novel mask, stepped into the animal's

field of view with only their profile visible to the animal, then sat in a chair at the eye level of the animal

for the duration of the condition. The experimenter remained still, with only the profile visible, for 9 minutes

(profile condition). Another alone condition elapsed for 3 minutes in order to allow the animal to return to

baseline behavior, before the experimenter stepped back into view, then sat in a chair at eye level facing

the animal and making eye contact. The experimenter remained still, attempting to make eye contact with

the animal for 9 minutes (stare condition) before stepping out of view and ending the session. A minimum

of one month elapsed between repetitions of this task with the same animal, as animals may habituate to

the human intruder (Kalin et al. 2018).

Social Interaction: Animals were removed from the home cage and placed in a play cage set up in an

unfamiliar room. A familiar cagemate was introduced to the play cage, and social interaction between the

subject and cagemate was observed. We used an ethogram previously defined by others to evaluate social

behaviors [12,13]. The cagemate chosen for this task was always the same across conditions. The cagemate

did not receive any injection prior to testing. A minimum of 48 hours elapsed between social interaction

tests.

Probabilistic stimulus-reward learning task

Three monkeys (He, Cy, and Wo) were trained to perform a two alternative forced choice probabilistic

learning task on a touch-sensitive monitor. All testing occurred while monkeys were seated in a wheeled

transport cage. In each 200-trial session, subjects had to learn to discriminate which visual stimulus in

two pairs of novel stimuli was associated with the highest probability of receiving a food reward pellet

(banana, grape, and raspberry flavored 20mg BIO-Serv Dustless Precision Pellets®). In each pair of

stimuli, one option was associated with an 80% probability of receiving a reward whereas the other was

associated with 20% probability of receiving a reward. Stimuli were novel at the beginning of each

session, the assignment of visual stimuli to high/low probability was random, and the probability of

receiving a reward was fixed throughout the session.

Each trial began with the presentation of a centrally located green square 'lever' stimulus. Once monkeys

pressed the lever stimulus, two stimuli were presented to the left and right of the screen, and monkeys

could choose between them. Pressing one stimulus caused the non-selected option to be removed from the

screen. The chosen stimulus remained on the screen for a further 0.5 seconds before being extinguished

and a 4 second inter-trial interval commenced. For one pair of stimuli, if a reward was to be delivered it

was dispensed when the chosen stimulus was removed from the screen. For the other pair of stimuli if a

reward was to be delivered, it was dispensed 1 second after the chosen stimulus was removed from the

screen. The presentation of stimuli pairs was randomized across trials and the left-right location of each

stimulus was also counterbalanced across trials.

Monkeys completed daily 200-trial sessions (100 trials of each pair), run 4-5 days per week. 0.1mg/kg,

0.3mg/kg DCZ, and vehicle were administered in sessions after stable performance was established each

week. In practice this meant that injections of 0.1mg/kg, 0.3mg/kg DCZ, or vehicle were administered on

either Wednesday, Thursday, or Friday.

fMRI analysis

Functional imaging data was preprocessed with standard AFNI/SUMA pipelines [14,15]. Raw images

were first converted into NIFTI data file format and ordered into BIDS format [16]. The T1 weighted

image from each session was first warped to the standard NMT atlas space using a newly developed

macaque skull stripping function followed by AFNI's @animal_warper command [15,17]. Specifically,

the T1w images were spatially normalized, then skull stripped using the U-Net model built from Primate

Data-Exchange open data sets [18], the mask was dilated out 6mm to ensure the whole brain was within

the mask, and finally the mask was applied to the original image, which was then renormalized. This

normalized skull-stripped T1w image was then aligned to the NMT atlas [17], along with subject specific

versions of the CHARM [15] and SARM atlases [19].

The EPI data were preprocessed using a customized version of the AFNI NHP preprocessing pipeline

(Jung et al 2020). For each session pre- and post-injection data were processed separately using the same

parameters. In addition to a set of dummy scans, the first 3 TRs of each EPI were removed to ensure that

any magnetization effects were removed from the data prior to functional connectivity analyses. The

images were first slice time corrected, then motion correction was applied and the EPIs were aligned to

the within session T1w image and warped to the standard space. Following alignment to the standard

space, the EPIs were blurred with a full width half max of 3mm, and then converted to percent signal

change. We then regressed the motion derivatives from each scan along with CSF and WM signal

regressors. The residuals of this analysis were then used to compute the functional connectivity analysis

described below.

The full connectome analyses were computed using 3dNetCorr function in AFNI [14,20]. The error time

series from the preprocessing steps described above were split into half runs and then we computed the

correlation between ROIs at various levels of the cortical [15] and subcortical [19] hierarchical atlas.

Following the creation of the individual correlation matrices, or connectomes, for each half run, the

connectomes were Fisher's z-transformed and then averaged within condition to create mean

connectomes for each condition (both pre-/post- injection for the vehicle, DCZ 0.1 mg/kg, and DCZ 0.3

mg/kg sessions) at all hierarchical levels. We then calculated the difference between connectomes by

subtracting the pre-injection data from the post- injection data for each session. Finally the average

difference connectome was computed across subjects for each drug level. To statistically determine the

effect of drug on the functional connectivity in CHARM and SARM atlases, the mean connectomes from

individual monkeys were then submitted to an ANOVA with main effects of drug and area category.

Monkey and brain area were modeled as random effects and brain area was nested under monkey.

In addition to the functional connectomes analyses, we used a modified group Independent Components

Analysis (ICA) to look for any changes in the component structure of our data following the drug

manipulations. The same split run error time series from the connectome analyses were submitted to

FSL's melodic ICA function [21,22] allowing for 30 possible components and a mixture model threshold

of 0.5. Following the computation of the group ICA, we submitted resultant data to FSL's dual regression

function, which was used to extract the spatial ICs for each subject and condition [21,23]. The extracted

condition IC scores were then submitted to a 2 (Pre-Injection vs Post-Injection) by 3 (Drug: Vehicle, DCZ

0.1 mg/kg, DCZ 0.3 mg/kg) ANOVA which was computed on a voxel by voxel basis.

Behavioral analysis

Data from socio-emotional tests were assessed by a scorer blinded to the condition. Data was scored in

Observer XT 14.0 and exported to GraphPad Prism 9.0 (GraphPad, San Diego CA). Behavioral ethograms

(**Supplemental Tables 2** and **3**) were adapted from previous studies [11,13]. Behavioral responses in the

human intruder and social interaction paradigms were assessed for normality using a Shapiro-Wilk

normality test. Human intruder data was analyzed using two-way repeated measures ANOVAs with

matching by subject. No assumption of sphericity was made. Sidak's multiple comparison test was used to

evaluate differences between categories (i.e. differences between DCZ and vehicle treatments within the

alone, profile, and stare conditions). Hostile and anxious behaviors during the stare condition of the human

intruder test were analyzed using a paired student's t-test, with matching by subject. Social interaction data

was also analyzed using a paired student's t-test, with matching by subject.

Data from the probabilistic learning task was analyzed using MATLAB 2021a. In all sessions analyzed,

each of the monkeys learned to discriminate the pairs of stimuli by the end of the session. This was based

on a running average of 10 trials and criterion was set at 8/10 trials or 80%. We assessed reaction time

(RT) as a measure of general task-execution process such as attention or motivation. RT was defined as

the time duration (in milliseconds) from the presentation of two stimuli to a choice response of the

monkeys. We analyzed RT using ANOVA with factors of drug as a main effect and monkey and session

as random effects.

SUPPLEMENTAL TABLES

Animal	rs-fMRI (vehicle)	rs-fMRI (DCZ 0.1 mg/kg)	rs-fMRI (DCZ 0.3 mg/kg)	Socio-emotional tasks	Probabilistic learning task
La	N	Y	N	Y	N
Ha	Y	Y	N	Y	N
Wo	Y	Y	Y	Y	N
He	N	Y	Y	Y	Y
Cy	Y	Y	Y	N	Y
Bu	Y	Y	Y	N	Y
Pi	Y	N	Y	N	N

179 **Supplemental Table 1:** Assignments of monkeys to fMRI and behavioral testing conditions. Y and N

180 indicate the condition that the data was collected and not collected, respectively.

Category and Specific Behaviors	Measurement	Examples
Affiliative	Cumulative Frequency	
Lipsmack	Frequency	Rapid movement of pursed lips, accompanied by a smacking sound
Present	Frequency	Rigid posture (knees locked) with tail elevated and rump oriented toward the stimulus object
Anxiety	Cumulative Frequency	
Scratch	Frequency	Rapid scratch of body with hands or feet

Body shake	Frequency	Shake of the whole body or just head and shoulders region
Tooth grind	Frequency	Repetitive rubbing or chewing of upper and lower teeth
Yawn	Frequency	Open mouth widely, exposing teeth
Fearful	Cumulative Frequency	
Withdrawal	Frequency	Quick, jerky motion away from the stimulus object (jump back)
Grimace	Frequency	Retracted lips, exposed clenched teeth
Hostile	Cumulative Frequency	
Cage aggression	Frequency	Slaps, shakes, bites, or slams body against cage
Threat	Frequency	Open mouth (no teeth exposed)
Bark	Frequency	Low pitch, high intensity, rasping, guttural
Lunge	Frequency	A quick, jerky movement toward the stimulus
Position	Duration	
Front of cage	Duration	No part of animal's body contacts back of the cage
Back of cage	Duration	Animal's body contacts back of the cage
Stereotypies	Cumulative Duration	
Pacing	Duration	Repetitive motor pattern around the test cage

Motor Stereotypy	Duration	Repetitive, abnormal voluntary or involuntary motor patterns (e.g. swinging, twirling, flipping)
Stillness	Duration	Motionless posture for at least 3 seconds (except small head movements)
Vocalizations	Cumulative Frequency	
Coo	Frequency	Clear soft pitch and intensity, sounds like "ooooh"
Scream	Frequency	High pitch, high intensity screech or loud chirp

182   **Supplemental Table 2.** Behavioral ethogram for human intruder task, adapted from Raper et al. 2019.

Category and Specific Behaviors	Measurement	Examples
Avoidance	Cumulative Frequency	
Attack	Frequency	Makes threatening gestures (i.e., mouth threat, head or body lunge, cage shake) towards or hits, grabs or bites the conspecific
Cower	Frequency	Withdraws from the conspecific in a crouched or recoiled position
Escape	Frequency	Sudden, explosive movement away from conspecific
Withdrawal	Frequency	Moves away at a normal pace from the conspecific when approached

Movement	Cumulative Duration	
Passive	Duration	Inactive, stays in one location
Manipulation	Duration	Handles, chews, licks, moves, or smells objects or cage parts
Locomotion	Duration	Walks, runs, climbs, or jumps
Self-directed	Duration	Engages in self-directed behaviors
Social Contact	Cumulative Duration	
Contact	Duration	Touches or holds the conspecific, not covered by any other behavior
Gives grooming	Duration	Subject grooms the conspecific
Receives grooming	Duration	The conspecific grooms the subject
Play	Duration	Chasing, wrestling, and "rough and tumble" behaviors

Supplemental Table 3. Behavioral ethogram for social interaction task, adapted from Forcelli et al. 2017.

SUPPLEMENTAL REFERENCES

1 Upright NA, Baxter MG. Effect of chemogenetic actuator drugs on prefrontal cortex-dependent

working memory in nonhuman primates. Neuropsychopharmacology : official publication of the

American College of Neuropsychopharmacology. 2020;45(11):1793-98.

2　Wu TL, Mishra A, Wang F, Yang PF, Gore JC, Chen LM. Effects of isoflurane anesthesia on

resting-state fMRI signals and functional connectivity within primary somatosensory cortex of

monkeys. Brain and behavior. 2016;6(12):e00591.

3　Russ BE, Petkov CI, Kwok SC, Zhu Q, Belin P, Vanduffel W, et al. Common functional

localizers to enhance NHP & cross-species neuroscience imaging research. NeuroImage.

2021;237:118203.

4　Leite FP, Tsao D, Vanduffel W, Fize D, Sasaki Y, Wald LL, et al. Repeated fMRI using iron

oxide contrast agent in awake, behaving macaques at 3 Tesla. NeuroImage. 2002;16(2):283-94.

5　Nagai Y, Miyakawa N, Takuwa H, Hori Y, Oyama K, Ji B, et al. Deschloroclozapine, a potent

and selective chemogenetic actuator enables rapid neuronal and behavioral modulations in mice

and monkeys. Nature neuroscience. 2020;23(9):1157-67.

6　Kalin NH, Shelton SE. Defensive behaviors in infant rhesus monkeys: environmental cues and

neurochemical regulation. Science (New York, NY). 1989;243(4899):1718-21.

7　Kling A, Cornell R. Amygdalectomy and social behavior in the caged stump-tailed macaque

(Macaca speciosa). Folia primatologica; international journal of primatology. 1971;14(3):190-

208.

8　Kalin NH, Shelton SE, Davidson RJ, Kelley AE. The primate amygdala mediates acute fear but

not the behavioral and physiological components of anxious temperament. The Journal of

neuroscience : the official journal of the Society for Neuroscience. 2001;21(6):2067-74.

9　Izquierdo A, Newman TK, Higley JD, Murray EA. Genetic modulation of cognitive flexibility

and socioemotional behavior in rhesus monkeys. Proceedings of the National Academy of

Sciences of the United States of America. 2007;104(35):14128-33.

10　Kalin NH, Shelton SE, Davidson RJ. Role of the primate orbitofrontal cortex in mediating

anxious temperament. Biological psychiatry. 2007;62(10):1134-9.

11   Raper J, Murphy L, Richardson R, Romm Z, Kovacs-Balint Z, Payne C, et al. Chemogenetic

Inhibition of the Amygdala Modulates Emotional Behavior Expression in Infant Rhesus

Monkeys. eNeuro. 2019;6(5).

12   Wellman LL, Forcelli PA, Aguilar BL, Malkova L. Bidirectional Control of Social Behavior by

Activity within Basolateral and Central Amygdala of Primates. The Journal of neuroscience : the

official journal of the Society for Neuroscience. 2016;36(33):8746-56.

13   Forcelli PA, Wellman LL, Malkova L. Blockade of glutamatergic transmission in the primate

basolateral amygdala suppresses active behavior without altering social interaction. Behavioral

neuroscience. 2017;131(2):192-200.

14   Cox RW. AFNI: software for analysis and visualization of functional magnetic resonance

neuroimages. Comput Biomed Res. 1996;29(3):162-73.

15   Jung B, Taylor PA, Seidlitz J, Sponheim C, Perkins P, Ungerleider LG, et al. A comprehensive

macaque fMRI pipeline and hierarchical atlas. NeuroImage. 2021;235:117997.

16   Gorgolewski KJ, Auer T, Calhoun VD, Craddock RC, Das S, Duff EP, et al. The brain imaging

data structure, a format for organizing and describing outputs of neuroimaging experiments.

Scientific data. 2016;3:160044.

17   Seidlitz J, Sponheim C, Glen D, Ye FQ, Saleem KS, Leopold DA, et al. A population MRI brain

template and analysis tools for the macaque. NeuroImage. 2018;170:121-31.

18   Wang X, Li XH, Cho JW, Russ BE, Rajamani N, Omelchenko A, et al. U-net model for brain

extraction: Trained on humans for transfer to non-human primates. NeuroImage.

2021;235:118001.

19   Hartig R, Glen D, Jung B, Logothetis NK, Paxinos G, Garza-Villarreal EA, et al. The Subcortical

Atlas of the Rhesus Macaque (SARM) for neuroimaging. NeuroImage. 2021;235:117996.

20   Taylor PA, Saad ZS. FATCAT: (an efficient) Functional and Tractographic Connectivity

Analysis Toolbox. Brain connectivity. 2013;3(5):523-35.

21   Beckmann CF, Mackay CE, Filippini N, Smith SM. Group comparison of resting-state FMRI

240   data using multi-subject ICA and dual regression. NeuroImage. 2009;47(Suppl 1):S148.

241   22   Beckmann CF, Smith SM. Probabilistic independent component analysis for functional magnetic

242   resonance imaging. IEEE transactions on medical imaging. 2004;23(2):137-52.

243   23   Nickerson LD, Smith SM, Öngür D, Beckmann CF. Using Dual Regression to Investigate

244   Network Shape and Amplitude in Functional Connectivity Analyses. Frontiers in neuroscience.

245   2017;11:115.