Single-domain antibodies efficiently neutralize SARS-CoV-2 variants of concern

Irina A. Favorskaya[1], Dmitry V. Shcheblyakov[2]*, Ilias B. Esmagambetov[2], Inna V. Dolzhikova[3], Irina A. Alekseeva[2], Anastasia I. Korobkova[2], Daria V. Voronina[2], Ekaterina I. Ryabova[2], Artem A. Derkaev[2], Anna V. Kovyrshina[3], Anna A. Iliukhina[3], Andrey G. Botikov[3], Olga L. Voronina[2], Daria A. Egorova[2], Olga V. Zubkova[2], Natalia N. Ryzhova[2], Ekaterina I. Aksenova[2], Marina S. Kunda[2], Denis Y. Logunov[1], Boris S. Naroditsky[2] and Alexandr L. Gintsburg[2]

[1]Medical Microbiology Department, Federal State Budget Institution «National Research Center for Epidemiology and Microbiology Named after Honorary Academician N.F. Gamaleya» of the Ministry of Health of the Russian Federation, Moscow, Russia

[2]Department of Genetics and Molecular Biology of Bacteria, Federal State Budget Institution «National Research Centre for Epidemiology and Microbiology Named after Honorary Academician N.F. Gamaleya» of the Ministry of Health of the Russian Federation, Moscow, Russia

[3]Department of the National Virus Collection, Federal State Budget Institution «National Research Centre for Epidemiology and Microbiology Named after Honorary Academician N.F. Gamaleya» of the Ministry of Health of the Russian Federation, Moscow, Russia

* Correspondence:
Dmitry V. Shcheblyakov
sdmitryv@yahoo.com

Keywords: SARS-CoV-2, COVID-19, single-domain antibodies, VHH, VOC, neutralizing antibodies, nanobodies.

ABSTRACT

Virus-neutralizing antibodies are one of the few treatment options for COVID-19. The evolution of SARS-CoV-2 virus has led to the emergence of virus variants with reduced sensitivity to some antibody-based therapies. The development of potent antibodies with a broad spectrum of neutralizing activity is urgently needed. Here we isolated a panel of single-domain antibodies that specifically bind to the receptor-binding domain of SARS-CoV-2 S glycoprotein. Three of the selected antibodies exhibiting most robust neutralization potency were used to generate dimeric molecules. We observed that these modifications resulted in up to a 200-fold increase in neutralizing activity. The most potent heterodimeric molecule efficiently neutralized each of SARS-CoV-2 variant of concern, including Alpha, Beta, Gamma and Delta variants. This heterodimeric molecule could be a promising drug candidate for a treatment for COVID-19 caused by virus variants of concern.

INTRODUCTION

The COVID-19 pandemic, caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2), has resulted in over 250 million of infections and over 5 million deaths worldwide (November 2021, WHO). These numbers are still rising and COVID-19 disease remains a great challenge to public health system. The development of safe and effective treatment together with vaccination is a highly important goal for scientists and health care professionals over the world.

One of the approaches to the development of an effective therapeutic agent is the isolation of monoclonal antibodies that efficiently neutralize SARS-CoV-2 virus. Currently, several monoclonal antibodies already received emergency use authorization for COVID-19 treatment and post-exposure prophylaxis. These antibodies are bamlanivimab plus etesevimab, casirivimab plus imdevimab and sotrovimab (1–3).

Along with conventional antibodies, camelid single-domain antibodies (also called nanobodies or VHHs) are promising candidates for the development of antibody-based therapies (4, 5). Camelids have unique heavy-chain antibodies that are devoid of light chains (6). Nanobodies are the minimal antigen-binding domains of these heavy-chain antibodies. They have several advantages, including the ability to recognize epitopes that are not accessible to conventional antibodies, increased stability of nanobodies, simplicity of generation of multivalent forms and low cost bacterial production (7). Currently, several nanobodies to SARS-COV-2 have been isolated showing neutralizing activity in *in vitro* and *in vivo* studies, which confirms their promising potential as therapeutic agents (8–14).

The evolution of SARS-CoV-2 virus has resulted in emergence of virus variants that have become more transmissible and less sensitive to neutralizing antibodies. The spread of these new virus variants has reduced the efficacy of vaccines and some therapeutic antibodies. The list of these variants of concern (VOCs) consist of B.1.1.7 (Alpha), B.1.351 (Beta), B.1.1.28/P.1 (Gamma) and B.1.617.2 (Delta) variants (November 2021, WHO). The reduction of neutralization by antibodies is caused by mutations in S glycoprotein, including K417N/T, L452R, T478K, E484K and N501Y substitutions (15, 16). In this regard, it became necessary to isolate antibodies, the neutralizing activity of which will not be affected due to observed mutations, therefore, these antibodies or their cocktails will retain activity against each of VOCs.

Here we identified and characterized a panel of single-domain antibodies isolated from immune VHH library that specifically bind RBD of S glycoprotein. We assessed the neutralizing activity of the isolated antibodies in a microneutralization assay with live SARS-CoV-2 and selected three most potent antibodies. To increase the therapeutic potential, these clones were modified to homodimeric and heterodimeric forms, and the neutralizing activity against SARS-CoV-2 VOCs was investigated. The most potent heterodimeric form, P2C5-P5F8, exhibited activity against all tested virus variants at low concentration. These results indicate that P2C5-P5F8 heterodimer is a promising candidate for further research to develop COVID-19 antibody-based therapy.

RESULTS

To identify SARS-CoV-2 neutralizing single-domain antibodies we immunized one Bactrian camel with SARS-CoV-2 RBD. The recombinant RBD protein was previously produced in CHO-S cells and purified. Immunization was performed using five sequential injections (Fig 1A). Blood was collected 5 days after final immunization and the serum RBD-specific antibodies titer was detected by ELISA. Post-immunization serum demonstrated potent binding to SARS-CoV-2 RBD with titer

more than 1/200 000 (Fig 1B). The neutralizing activity of antibodies in the immunized serum was measured by the microneutralization assay using live SARS-CoV-2 B.1.1.1, the neutralizing antibodies titer was 1/1280.

To isolate monoclonal RBD-specific nanobodies, we constructed a VHH phage display library by cloning VHH sequences from B cells of immunized camel into a phagmid vector. We performed one round of phage display followed by ELISA-based screening of individual clones to identify RBD-specific binders.

We used monoclonal phage-ELISA together with monoclonal ELISA to identify RBD binders with high expression levels and high solubility. A total of 212 clones showing positive phage-ELISA and ELISA signals were selected for sequencing (Fig 1C), which lead to the identification of 39 unique clones of nanobodies. After sequence alignment and phylogenetic analysis we selected 16 antibodies with highly diverse sequences for further research (Fig 1D). These nanobodies were expressed and purified, and the RBD binding activity of each antibody was analyzed by ELISA. We confirmed the binding of 14 clones with ELISA EC50 values ranged from 1.1 to 313.3 nM (Fig 2A). Most of the antibodies (13 out of 14) displayed strong positive signals with EC50 below 22 nM.

To examine the capacity of selected RBD binders to neutralize live SARS-CoV-2 virus we performed a microneutralization assay with inhibition of the cytopathic effect of virus as a marker of neutralization. We found that 10 of 14 nanobodies efficiently neutralized SARS-CoV-2 virus (B.1.1.1) at concentrations ranging from 12 nM to 1540 nM (Fig 2A). The three most potent antibodies were P2C5, P2G1 and P5F8, which completely inhibited the cytopathic effect of the virus at 24 nM, 12 nM and 48 nM, respectively.

We investigated the binding kinetics of P2C5, P2G1 and P5F8 neutralizing nanobodies via surface plasmon resonance (SPR). All antibodies tested bound RBD with high affinity, dissociation constants (K_D) were 3.97 nM, 5.36 nM and 1.94 nM for P2C5, P2G1 and P5F8, respectively (Fig 2C).

Since the capacity of an antibody to neutralize SARS-CoV-2 virus is frequently associated with blocking the interaction between S glycoprotein and ACE2 receptor, we analyzed the ability of isolated nanobodies to block the RBD binding to ACE2 by competitive ELISA (Fig 2B). We revealed a decrease in ELISA signal for most of neutralizing nanobodies indicating antibodies competition with ACE2 for RBD binding. We observed that one of the most potent neutralizing antibodies, P2C5, blocked the ACE2-RBD interaction with an inhibition rate greater than 90% at 0.5 µg/ml.

In order to improve the antiviral property of antibodies we produced homodimer and heterodimer forms of P2C5, P2G1 and P5F8 clones. Monomers were fused using a flexible GS-linker (Gly4Ser)$_4$ for dimerization. The binding of generated dimeric molecules to the recombinant RBD protein was confirmed by ELISA. Analysis of the neutralization capacity in the microneutralization assay revealed a pronounced improvement in the neutralizing activity of several dimeric forms (Table 1). Notable, P2C5 homodimer inhibited cytopathic effect of the virus at 89 pM, thus the homodimer activity increased more than 200-fold compared to the monomer activity. We also observed that P2C5-P5F8 heterodimer form was up to 100 times more active than the monomeric forms, this molecule exhibited neutralization activity at 178 pM.

With the emergence and spread of new variants of SARS-CoV-2 with increased transmissibility and reduced neutralization by antibodies it became necessary to isolate antibodies with a broad spectrum of neutralizing activity. We assessed the neutralizing activity of isolated antibodies against variants

of concern (VOCs), including B.1.1.7 (Alpha), B.1.351 (Beta), B.1.1.28/P.1 (Gamma) and B.1.617.2
(Delta) variants (Table 1). P5F8 and P2G1 monomeric forms remained active against all tested VOCs
(inhibited CPE at 12-96 nM). P5F8 monomer showed a 2-fold decrease in activity against B.1.351
and B.1.1.28/P.1. P2G1 showed a 2-4-fold decrease against VOCs. P2C5 monomer neutralized
B.1.1.7, B.1.351 and B.1.1.28/P.1 but was unable to neutralize B.1.617.2. The most potent
heterodimeric molecule P2C5-P5F8 exhibited activity against all VOCs neutralizing B.1.1.7,
B.1.351, B.1.1.28/P.1 and B.1.617.2 variants at 89 pM, 356 pM, 356 pM and 2.85 nM, respectively.
The most active P2C5 homodimer neutralized each of VOC except B.1.617.2 at a concentration 356
pM and lower, depending on the virus variant. Overall, P2C5-P5F8 heterodimer showed neutralizing
activity against each of the tested VOCs and was a promising candidate for further investigations as a
potential therapeutic agent.

DISCUSSION

The SARS-CoV-2 virus gave rise to ongoing pandemic, which continues to claim lives and poses a
threat to public health. Along with vaccination to prevent COVID-19, therapeutic antiviral agents are
urgently needed to treat the infection.

One of the promising approaches for the viral infection treatment is an antibody-based therapy.
Antibodies administration approved for immunoprophylaxis and treatment of viral infections caused
by respiratory syncytial virus and ebolavirus (17, 18). Today, several monoclonal antibodies are
already in use for COVID-19 treatment *(*bamlanivimab plus etesevimab, casirivimab plus imdevimab
and sotrovimab). It was shown that the administration of these antibodies to nonhospitalized patients
with symptomatic COVID-19 and risk factors for disease progression reduced the risk of
hospitalization and death (1–3)**.**

In addition to conventional antibodies, single-domain antibodies or nanobodies from camelid heavy-
chain antibodies are being actively investigated as a potential therapeutic agent with a number of
advantages. The small size and single-domain nature of nanobodies allow them to bind epitopes that
are not available for conventional antibodies, in particular concave epitopes such as grooves and
clefts (19–21). Nanobodies show strong binding affinities and also are highly stable across wide
range of temperatures and pH (22). The production of nanobodies in bacteria is less expensive than
classical antibody production. All this allows the use of single-domain antibodies for many of
applications, including research, diagnostics and therapy (5, 7).

The main mechanism of virus-neutralizing activity of antibodies is associated with the prevention of
viral entry into host cells. The entry of SARS-CoV-2 virus occurs after the interaction of S
glycoprotein with cell receptor angiotensin-converting enzyme 2 (ACE2). Therefore, S glycoprotein,
in particular its receptor-binding domain (RBD), is an attractive target for the development of
neutralizing antibodies.

In our work, we immunized a Bactrian camel with the recombinant RBD protein for following
isolation of effective neutralizing antibodies against SARS-CoV-2 virus. We generated a VHH phage
display library and selected a panel of RBD-specific nanobodies. Using an *in vitro*
microneutralization assay we identified three clones (P2C5, P5F8 and P2G1) of the most potent
antibodies that completely inhibited the cytopathic effect of SARS-CoV-2 virus at minimal
concentrations from 12 nM to 48 nM.

At the next stage, to enhance the neutralization potential of isolated nanobodies we generated homodimeric and heterodimeric forms of the most active antibodies. It was previously shown that multivalent forms of nanobodies are characterized by increased functionality by improving antigen-binding through avidity effect (23). For example, the dimeric form of anti-TNF VHH was up to 500 times more active than the monovalent form in rheumatoid arthritis model *in vivo* (24), homotrimeric molecule of anti-SARS-CoV-2 nanobody $Nb21_3$ were 30-fold more potent in pseudovirus neutralization (13), dimeric forms of nanobodies to botulinum neurotoxin showed a higher protective activity *in vivo* (25). In our study, we revealed a 200-fold improvement of neutralization potential after dimerization of P2C5 nanobody. The bivalent molecule effectively neutralized SARS-CoV-2 virus at 89 pM. We also observed a pronounced increase in neutralizing activity in the case of P2C5-P5F8 heterodimeric form, which inhibited SARS-CoV-2 virus at 178 pM. In the case of the bivalent form of P5F8 we observed only a slight increase in activity. We assume that this may be due to the monovalent binding of P5F8 dimer to the S glycoprotein, for example as a result of the insufficient linker length to span the distance between epitopes in the trimeric S glycoprotein.

The emergence of SARS-CoV-2 variants with mutations in their RBD epitopes leads to the viral escape from neutralizing antibodies. It was shown that the virus-neutralization activity of sera samples obtained from individuals vaccinated with Pfizer/BNT162b2, Moderna/mRNA-1273 or Sputnik V decreased in the case of the B.1.351, B.1.1.28/P.1, B.1.617.2 variants compared to wild-type virus (26–29). Escape mutations are also a challenge for antibody-based therapy development. Some therapeutic antibodies have become ineffective against VOCs, for example, the activity of bamlanivimab against Delta is reduced, the activity of etesevimab and casivimab against Beta variant is reduced (30). The activity of some previously isolated nanobodies also altered by escape mutations. The Beta strain drastically reduced the efficacy of the high-affinity nanobodies Nb20 and Nb21 (31). Thereby, it is urgently needed to identify nanobodies or their combinations that retain activity against SARS-CoV-2 variants of concern.

We investigated the activity of isolated potent nanobodies and their dimeric forms using a microneutralization assay against live SARS-CoV-2 VOCs. We found that P2C5-P5F8 heterodimer effectively neutralized all tested virus variants at low concentration. In particular it was observed that this dimeric molecule completely inhibited the cytopathic effect of the Alpha, Beta, Gamma and Delta variants at concentrations 89 pM, 356 pM, 356 pM and 2.85 nM, respectively.

For the improvement of antiviral efficacy of obtained P2C5-P5F8 molecule further modifications may include the fusion of the heterodimer to human IgG1 Fc domain. Fc modifications are often used to increase serum half-life of a nanobodies, as well as to involve Fc-mediated effector functions (antibody-dependent cell-mediated cytotoxicity, complement-dependent cytotoxicity, etc.) (25, 32).

Overall, we produced heterodimeric form of nanobodies that possess a broad spectrum of activity, overcoming escape mutations in VOCs. This heterodimeric molecule is a promising candidate for the further development of an effective antiviral agent for COVID-19 treatment especially for patients with risk factors for progression to severe COVID-19.

MATERIALS AND METHODS

Cell lines and viruses

CHO-S cells were obtained from Thermo Fisher Scientific (USA), cat. no. R80007. Vero E6 cells (ATCC CRL 1586) was obtained from Russian collection of vertebrate cell lines.

SARS-CoV-2 strains B.1.1.1 (hCoV-19/Russia/Moscow_PMVL-1/2020), B.1.351 (hCoV-
19/Russia/SPE-RII-27029S/2021) and B.1.617.2 (hCoV-19/Russia/SPE-RII-32758S/2021) were
isolated from nasopharyngeal swabs.

SARS-CoV-2 strains B.1.1.7 (hCoV-19/Netherlands/NoordHolland_20432/2020) and B.1.1.28/P.1
(hCoV-19/Netherlands/NoordHolland_10915/2021) were obtained from European Virus Archive
Global.

Cloning, expression and purification of recombinant SARS-CoV-2 RBD protein

The RBD nucleotide sequence of SARS-CoV-2 Wuhan-Hu-1 isolate (Genbank accession number
MN908947, from 319 to 545 aa) was synthesized (Evrogen, Russia) and cloned into the pCEP4
mammalian expression vector (Thermo Fisher Scientific, USA). The signal peptide of alkaline
phosphatase SEAP was inserted before RBD coding sequence, the His tag was fused to the end of the
sequence. The obtained recombinant vector was transfected into CHO-S cells using CHOgro High
Yield Expression System (Mirus Bio, USA). The culture supernatant containing the RBD protein was
harvested after 10 days, the recombinant RBD protein was purified by IMAC using an AKTA start
system (Cytiva, USA) and a HisTrap column (Cytiva, USA). The protein expression and purity was
confirmed with SDS-PAGE.

Bactrian camel immunization

Bactrian camel was immunized with recombinant SARS-CoV-2 RBD protein at a dose of 100 μg
with complete Freund`s adjuvant subcutaneously followed by four booster immunizations with 100
μg RBD with incomplete Freund`s adjuvant. The interval between the first and second injections was
14 days; the intervals between all subsequent injections were 10 days. Before immunization, a small
blood sample (5 ml) was collected as a control. 50 ml blood was collected 5 days after the last
immunization and was used for peripheral blood mononuclear cells (PBMC) isolation and VHH
phage display library construction.

Phage display library construction

228 The library construction was performed as described earlier (25). Briefly, total RNA was isolated
229 from PBMCs and then reverse transcribed to cDNA. VHHs coding sequences were PCR amplified
230 and cloned into a pHEN1 phagmid vector (33). Recombinant phagmids were electroporated into TG1
231 cells (Lucigen, USA). A VHH library of 7.3×10^6 individual clones was obtained.

Phage preparation and biopanning

Phage preparation was performed according to the previous description (25). To select RBD-specific
nanobodies one round of biopanning was performed. 5 μg of recombinant SARS-CoV-2 RBD protein
were immobilized in the well of microtiter plate. After rinsing three times with PBS with 0.1%
Tween 20 (TPBS), the well was blocked with blocking buffer (5% non-fat dried milk in TPBS) at 37
^{o}C for 1 h. A total of ~10^{11} phage particles in blocking buffer were added to the well and the plate
was incubated at 37 ^{o}C for 1 h. Unbound phages were removed by washing with TPBS. The bound
phages were eluted by trypsin (1 mg/ml). The eluted phage particles were used for infection of TG1
cells.

ELISA

Screening for RBD-specific binders was performed using monoclonal phage ELISA and monoclonal
ELISA. For monoclonal phage ELISA, individual clones were inoculated in 1 ml of 2xYT (Sigma,
USA) containing 100 μg/ml ampicillin in 96-deep-well plates. Bacterial cells were infected with
KM13 helper phage and incubated overnight at 30 ℃. The plates were centrifuged, 50 μl of the
supernatant were used for phage ELISA. For monoclonal ELISA, individual clones were cultured in
96-deep-well plates at 30 ℃ overnight. The next day, the cells were lysed by the freeze-thaw method,
and then the plates were centrifuged. 50 μl of the supernatant were used for monoclonal ELISA.
Immunoplates (Nunc MaxiSorp, Thermo Fisher Scientific, USA) were coated with recombinant RBD
protein (100 ng per well) at 4 ℃ overnight. The plates were rinsed three times with TPBS and the
wells were blocked with blocking buffer at 37 ℃ for 1 h. 50 μl of bacterial supernatants containing
monoclonal recombinant phages (monoclonal phage ELISA) or monoclonal nanobodies (monoclonal
ELISA) were added to the wells. After 1 h incubation at 37 ℃, the wells were rinsed five times with
TPBS. Bound phages were detected by HRP-conjugated anti-M13 antibody (Sino Biological, China),
bound VHH were detected by HRP-conjugated anti-Myc-tag antibody (Abcam, UK) followed by the
addition of 3,3`5,5`-tetramethylbenzidine (TMB) (Bio-Rad, USA) as a substrate. The reaction was
stopped by 1M H_2SO_4, the absorbance at 450 nm was read using a Varioskan LUX (Thermo Fisher
Scientific, USA).

To confirm RBD-specific binding of the selected nanobodies immunoplates (Nunc MaxiSorp,
Thermo Fisher Scientific, USA) were coated with RBD (100 ng per well). Serial two-fold dilutions
of antibodies were made, starting from 10 μg/ml to 0.6 ng/ml. 100 μl of each concentration was
added to the wells. Bound VHHs were detected by HRP-conjugated anti-Myc-tag antibody and TMB
substrate. EC50 values were calculated using four-parameter logistic regression using GraphPad
Prism 9 (GraphPad Software Inc, USA).

For competitive ELISA recombinant hACE2 protein were immobilized (400 ng per well) on
immunoplates (Nunc MaxiSorp, Thermo Fisher Scientific, USA). RBD-specific nanobodies were
mixed with HRP-conjugated recombinant RBD protein (0.2 μg/ml) to two final concentrations of
antibodies (0.5 μg/ml and 0.1 μg/ml) and incubated at 37 ℃ for 30 min. Nanobody to C. difficile
Toxin B was used as a negative control. After incubation samples (100 μl) were transferred to
hACE2 coated plates and incubated at 37 ℃ for 30 min. The TMB solution was added for 15 min,
the reaction was stopped by the addition of 1M H_2SO_4. The inhibition rate of ACE2-RBD binding
was calculated by comparing the signals in the sample and the negative control wells.

Expression and purification of nanobodies

Phagmids encoding the sequences of the selected nanobodies were transformed into BL21 cells
(NEB, USA) for expression and purification. The cells were cultured in 100 ml 2xYT (containing
100 μg/ml ampicillin) overnight at 30 ℃. The cells were harvested by centrifugation and lysed by a
BugBuster Protein Extraction Reagent (Novagen, USA). Nanobodies were purified by IMAC using
an AKTA start system and a HisTrap column, dialyzed against PBS and sterile filtered. Protein
expression and purity was confirmed with SDS-PAGE.

Microneutralization assay with live SARS-CoV-2

Nanobodies were serial two-fold diluted starting from 20 μg/ml to 1.2 ng/ml in complete Dulbecco`s
modified Eagle medium (DMEM) supplemented with 2% heat-inactivated fetal bovine serum (HI-
FBS). Triplicates of each dilution were mixed with 100 50% tissue culture infectious doses (TCID50)
of SARS-CoV-2 and incubated at 37 ℃ for 1 h. After incubation samples were added to a monolayer
of Vero E6 cells and incubated in a 5% CO_2 incubator at 37 ℃ for 96-120 h. The cytopathic effect

(CPE) of the virus was assessed visually. The minimal neutralizing concentration was defined as the
lowest antibody concentration (highest antibody dilution) that completely inhibited CPE of the virus
in two or three of the three replicable wells.

The following SARS-CoV-2 strains were used in the assay: B.1.1.1, B.1.1.7, B.1.351, B.1.1.28/P.1
and B.1.617.2.

All experiments were performed in a Biosafety Level 3 facility (BSL-3).

Binding kinetics measurements

The affinity of nanobodies was measured by surface plasmon resonance (Biacore 3000, GE
Healthcare Bio-Sciences AB, Sweden). Recombinant RBD protein at a concentration 20 μg/ml in 10
mM acetate buffer (pH 4.5) was immobilized on the surface of a CM5 sensor chip using a Amine
Coupling Kit (GE Healthcare Bio-Sciences AB, Sweden). VHHs were serial 4-fold diluted from 417
nM to 0 nM and flowed over the captured RBD surface at 15 μl/min for 3 minutes. Dissociation time
was 10 minutes. Working buffer was HBS-EP (0.01 M HEPES pH 7.4, 0.15 M NaCl, 2 mM EDTA,
0.005% v/v Surfactant P20). After each injection, the chip surface was regenerated with 10 mM Tris-
HCl, pH 1.5 for 40 s at a flow rate 30 μl/min. Calculations were performed using the BIAEvaluation
software (GE Healthcare Bio-Sciences AB, Sweden).

Generation of homodimeric and heterodimeric forms of nanobodies

Homodimeric and heterodimeric forms of nanobodies were generated using two rounds of PCR
followed by cloning PCR products into a pHEN1 vector. VHHs in dimeric forms were fused by GS-
linker (Gly4Ser)$_4$. In the first round of PCR, VHH monomers were amplified using Q5 High-fidelity
DNA polymerase (NEB, UK) and the primers listed in Table 2. In the second round PCR nanobodies
sequences were assembled together and amplified using pHEN1-F and pHEN1-R primers. PCR
products and the pHEN1 vector were digested with SfiI and NotI restriction enzymes (Thermo Fisher
Scientific, USA) and ligated using T4 ligase (Thermo Fisher Scientific, USA). Recombinant vectors
were used for TG1 cells transformation.

DNA isolation and sequencing

Phagemid DNA was isolated from bacterial cells using the Plasmid miniprep kit (Evrogen, Russia).
VHH-coding sequences were sequenced with Lac-prom (5'-CTTTATGCTTCCGGCTCGTATG-3')
and pIII-R (5` CTTTCCAGACGTTAGTAAATG 3`) primers according to the protocol of the
BigDyeTerminator 3.1 Cycle Sequencing kit for the Genetic Analyzer 3500 Applied Biosystems
(Waltham, MA, USA). The electrophoretic DNA separation was performed in 50-cm capillaries with
POP7 polymer. MEGA X was used for the generation of the single consensus sequences by
assembling of the forward and reverse sequences (34).

Phylogenetic tree analysis

A phylogenetic tree of isolated VHHs was constructed to display the sequence diversity of single-
domain antibodies. The analysis comprised the following traditional steps: alignment, phylogeny and
tree rendering. Sequences were aligned with MUSCLE (v3.8.31) (35). The phylogenetic tree was
reconstructed using the maximum likelihood method implemented in the PhyML program (v3.1/3.0
aLRT) (36). The WAG substitution model was selected assuming an estimated proportion of
invariant sites (of 0.000) and 4 gamma-distributed rate categories to account for rate heterogeneity

across sites. The gamma shape parameter was estimated directly from the data (gamma=0.448).
Reliability for internal branch was assessed using the aLRT test (SH-Like) (37). Graphical
representation and edition of the phylogenetic tree were performed with MEGA X (34).

AUTHOR CONTRIBUTIONS

Conception and design of the experiments: I.A.F., D.V.S., I.B.E., I.V.D., library construction, phage
display, cloning, production, purification and characterization of VHH and their dimeric forms:
I.A.F., I.A.A., A.I.K., I.B.E., E.I.R., A.A.D., O.V.Z., production and purification of recombinant
proteins: I.B.E., E.I.R., A.A.D., SARS-CoV-2 culture and microneutralization assay: I.V.D., A.V.K.,
A.A.I., A.G.B., SRP kinetic measurements: D.V,V., camel immunization and blood collection:
D.A.E., DNA sequencing, phylogenetic tree construction: O.L.V., N.N.R., E.I.A., M.S.K., data
analysis and interpretation, figures preparation: I.A.F., D.V.S., drafting the manuscript: I.A.F.,
revision the manuscript: D.V.S., I.B.E., I.V.D., O.L.V., approval of the final version for submission:
D.V.S., D.Y.L., B.S.N., A.L.G.

FUNDING

This work received funding from Sberbank Charitable Foundation "Investment to the Future",
donation agreement no. BM-03/2020.

ACKNOWLEDGMENTS

We would like to thank Y.S. Lebedin who kindly provided us with the recombinant hACE2 protein
and the HRP conjugated recombinant RBD protein.

REFERENCES

1. Dougan M, Nirula A, Azizad M, Mocherla B, Gottlieb RL, Chen P, et al. Bamlanivimab plus
    Etesevimab in Mild or Moderate Covid-19. *N Engl J Med* (2021) **385**:1382–92.
    doi:10.1056/NEJMoa2102685

2. Gupta A, Gonzalez-Rojas Y, Juarez E, Crespo Casal M, Moya J, Falci DR, et al. Early Treatment
    for Covid-19 with SARS-CoV-2 Neutralizing Antibody Sotrovimab. *N Engl J Med* (2021).
    doi:10.1056/NEJMoa2107934

3. Weinreich DM, Sivapalasingam S, Norton T, Ali S, Gao H, Bhore R, et al. REGEN-COV
    Antibody Combination and Outcomes in Outpatients with Covid-19. *N Engl J Med* (2021).
    doi:10.1056/NEJMoa2108163

4. Bessalah S, Jebahi S, Mejri N, Salhi I, Khorchani T, Hammadi M. Perspective on therapeutic and
    diagnostic potential of camel nanobodies for coronavirus disease-19 (COVID-19). *3 Biotech*
    (2021) **11**:89. doi:10.1007/s13205-021-02647-5

5. Jovčevska I, Muyldermans S. The Therapeutic Potential of Nanobodies. *BioDrugs* (2020) **34**:11–
    26. doi:10.1007/s40259-019-00392-z

6. Muyldermans S. Single domain camel antibodies: current status. *J Biotechnol* (2001) **74**:277–
    302. doi:10.1016/s1389-0352(01)00021-6

7. Muyldermans S. Nanobodies: natural single-domain antibodies. *Annu Rev Biochem* (2013)
    82:775–97. doi:10.1146/annurev-biochem-063011-092449

8.　Dong J, Huang B, Wang B, Titong A, Gallolu Kankanamalage S, Jia Z, et al. Development of
humanized tri-specific nanobodies with potent neutralization for SARS-CoV-2. *Sci Rep* (2020)
10:17806. doi:10.1038/s41598-020-74761-y

9.　Huo J, Mikolajek H, Le Bas A, Clark JJ, Sharma P, Kipar A, et al. A potent SARS-CoV-2
neutralising nanobody shows therapeutic efficacy in the Syrian golden hamster model of
COVID-19. *Nat Commun* (2021) **12**:5469. doi:10.1038/s41467-021-25480-z

10.　Lu Q, Zhang Z, Li H, Zhong K, Zhao Q, Wang Z, et al. Development of multivalent nanobodies
blocking SARS-CoV-2 infection by targeting RBD of spike protein. *J Nanobiotechnology* (2021)
19:33. doi:10.1186/s12951-021-00768-w

11.　Pymm P, Adair A, Chan L-J, Cooney JP, Mordant FL, Allison CC, et al. Nanobody cocktails
potently neutralize SARS-CoV-2 D614G N501Y variant and protect mice. *Proc Natl Acad Sci U
S A* (2021) **118**. doi:10.1073/pnas.2101918118

12.　Schoof M, Faust B, Saunders RA, Sangwan S, Rezelj V, Hoppe N, et al. An ultrapotent synthetic
nanobody neutralizes SARS-CoV-2 by stabilizing inactive Spike. *Science* (2020) **370**:1473–9.
doi:10.1126/science.abe3255

13.　Xiang Y, Nambulli S, Xiao Z, Liu H, Sang Z, Duprex WP, et al. Versatile and multivalent
nanobodies efficiently neutralize SARS-CoV-2. *Science* (2020) **370**:1479–84.
doi:10.1126/science.abe4747

14.　Xu J, Xu K, Jung S, Conte A, Lieberman J, Muecksch F, et al. Multimeric nanobodies from
camelid engineered mice and llamas potently neutralize SARS-CoV-2 variants. *bioRxiv* (2021).
doi:10.1101/2021.03.04.433768

15.　Harvey WT, Carabelli AM, Jackson B, Gupta RK, Thomson EC, Harrison EM, et al. SARS-
CoV-2 variants, spike mutations and immune escape. *Nat Rev Microbiol* (2021) **19**:409–24.
doi:10.1038/s41579-021-00573-0

16.　Hastie KM, Li H, Bedinger D, Schendel SL, Dennison SM, Li K, et al. Defining variant-resistant
epitopes targeted by SARS-CoV-2 antibodies: A global consortium study. *Science* (2021)
374:472–8. doi:10.1126/science.abh2315

17.　Markham A. REGN-EB3: First Approval. *Drugs* (2021) **81**:175–8. doi:10.1007/s40265-020-
01452-3

18.　Rocca A, Biagi C, Scarpini S, Dondi A, Vandini S, Pierantoni L, et al. Passive
Immunoprophylaxis against Respiratory Syncytial Virus in Children: Where Are We Now? *Int J
Mol Sci* (2021) **22**. doi:10.3390/ijms22073703

19.　Genst E de, Silence K, Decanniere K, Conrath K, Loris R, Kinne J, et al. Molecular basis for the
preferential cleft recognition by dromedary heavy-chain antibodies. *Proc Natl Acad Sci U S A*
(2006) **103**:4586–91. doi:10.1073/pnas.0505379103

20.　Muyldermans S. Applications of Nanobodies. *Annu Rev Anim Biosci* (2021) **9**:401–21.
doi:10.1146/annurev-animal-021419-083831

21.　Zavrtanik U, Lukan J, Loris R, Lah J, Hadži S. Structural Basis of Epitope Recognition by
Heavy-Chain Camelid Antibodies. *J Mol Biol* (2018) **430**:4369–86.
doi:10.1016/j.jmb.2018.09.002

22.　van der Linden RH, Frenken LG, Geus B de, Harmsen MM, Ruuls RC, Stok W, et al.
Comparison of physical chemical properties of llama VHH antibody fragments and mouse

monoclonal antibodies. *Biochim Biophys Acta* (1999) **1431**:37–46. doi:10.1016/s0167-
4838(99)00030-8

23. Saerens D, Ghassabeh GH, Muyldermans S. Single-domain antibodies as building blocks for
novel therapeutics. *Curr Opin Pharmacol* (2008) **8**:600–8. doi:10.1016/j.coph.2008.07.006

24. Coppieters K, Dreier T, Silence K, Haard H de, Lauwereys M, Casteels P, et al. Formatted anti-
tumor necrosis factor alpha VHH proteins derived from camelids show superior potency and
targeting to inflamed joints in a murine model of collagen-induced arthritis. *Arthritis Rheum*
(2006) **54**:1856–66. doi:10.1002/art.21827

25. Godakova SA, Noskov AN, Vinogradova ID, Ugriumova GA, Solovyev AI, Esmagambetov IB,
et al. Camelid VHHs Fused to Human Fc Fragments Provide Long Term Protection Against
Botulinum Neurotoxin A in Mice. *Toxins (Basel)* (2019) **11**. doi:10.3390/toxins11080464

26. Choi A, Koch M, Wu K, Dixon G, Oestreicher J, Legault H, et al. Serum Neutralizing Activity of
mRNA-1273 against SARS-CoV-2 Variants. *J Virol* (2021) **95**:e0131321.
doi:10.1128/JVI.01313-21

27. Gushchin VA, Dolzhikova IV, Shchetinin AM, Odintsova AS, Siniavin AE, Nikiforova MA, et
al. Neutralizing Activity of Sera from Sputnik V-Vaccinated People against Variants of Concern
(VOC: B.1.1.7, B.1.351, P.1, B.1.617.2, B.1.617.3) and Moscow Endemic SARS-CoV-2
Variants. *Vaccines (Basel)* (2021) **9**. doi:10.3390/vaccines9070779

28. Hoffmann M, Arora P, Groß R, Seidel A, Hörnich BF, Hahn AS, et al. SARS-CoV-2 variants
B.1.351 and P.1 escape from neutralizing antibodies. *Cell* (2021) **184**:2384-2393.e12.
doi:10.1016/j.cell.2021.03.036

29. Hoffmann M, Hofmann-Winkler H, Krüger N, Kempf A, Nehlmeier I, Graichen L, et al. SARS-
CoV-2 variant B.1.617 is resistant to bamlanivimab and evades antibodies induced by infection
and vaccination. *Cell Rep* (2021) **36**:109415. doi:10.1016/j.celrep.2021.109415

30. Planas D, Veyer D, Baidaliuk A, Staropoli I, Guivel-Benhassine F, Rajah MM, et al. Reduced
sensitivity of SARS-CoV-2 variant Delta to antibody neutralization. *Nature* (2021) **596**:276–80.
doi:10.1038/s41586-021-03777-9

31. Sun D, Sang Z, Kim YJ, Xiang Y, Cohen T, Belford AK, et al. Potent neutralizing nanobodies
resist convergent circulating variants of SARS-CoV-2 by targeting diverse and conserved
epitopes. *Nat Commun* (2021) **12**:4676. doi:10.1038/s41467-021-24963-3

32. Günaydın G, Yu S, Gräslund T, Hammarström L, Marcotte H. Fusion of the mouse IgG1 Fc
domain to the VHH fragment (ARP1) enhances protection in a mouse model of rotavirus. *Sci
Rep* (2016) **6**:30171. doi:10.1038/srep30171

33. Hoogenboom HR, Griffiths AD, Johnson KS, Chiswell DJ, Hudson P, Winter G. Multi-subunit
proteins on the surface of filamentous phage: methodologies for displaying antibody (Fab) heavy
and light chains. *Nucleic Acids Res* (1991) **19**:4133–7. doi:10.1093/nar/19.15.4133

34. Kumar S, Stecher G, Li M, Knyaz C, Tamura K. MEGA X: Molecular Evolutionary Genetics
Analysis across Computing Platforms. *Mol Biol Evol* (2018) **35**:1547–9.
doi:10.1093/molbev/msy096

35. Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput.
Nucleic Acids Res (2004) **32**:1792–7. doi:10.1093/nar/gkh340

36. Guindon S, Gascuel O. A simple, fast, and accurate algorithm to estimate large phylogenies by
maximum likelihood. *Syst Biol* (2003) **52**:696–704. doi:10.1080/10635150390235520
37. Anisimova M, Gascuel O. Approximate likelihood-ratio test for branches: A fast, accurate, and
powerful alternative. *Syst Biol* (2006) **55**:539–52. doi:10.1080/10635150600755453
38. Tillib SV, Vyatchanin AS, Muyldermans S. Molecular analysis of heavy chain-only antibodies of
Camelus bactrianus. *Biochemistry (Mosc)* (2014) **79**:1382–90. doi:10.1134/S000629791412013X
453

Figure legends

Figure 1. Isolation of RBD-specific nanobodies. (**A**) Immunization schedule. Bactrian camel was immunized with 100 mcg RBD subcutaneously (with complete Freund`s adjuvant), followed by four consecutive immunization with 100 mcg RBD subcutaneously (with incomplete Freund`s adjuvant). Blood samples were collected before immunization and five days after the last immunization. (**B**) RBD-specific antibodies in camel serum before and after immunization, detected by ELISA. The assay reveals a strong positive RBD-specific serological activity 5 days after the last immunization. (**C**) ELISA-based RBD-binders screening. A total of 212 individual clones with a strong positive ELISA signal were selected for sequencing. (**D**) Phylogenetic tree showing sequence diversity of 39 unique VHH clones from this study and four previously described naive single-domain antibodies of C. bactrianus (38), blue squares - naive single-domain antibodies of C. bactrianus, green squares – the clones selected for further analysis.

Figure 2. Characterization of the selected nanobodies. (**A**) RBD-binding activity of nanobodies by ELISA and neutralization activity of nanobodies by microneutralization assay using live SARS-CoV-2 virus. The minimal neutralizing concentration was defined as the lowest antibody concentration (highest antibody dilution) that completely inhibited the cytopathic effect of the virus in two or three from the three replicable wells. (**B**) Blocking of ACE2-RBD interaction measured by competitive ELISA. (**C**) Kinetic parameters of nanobodies interaction with RBD by SPR. Association (k_{on}), dissociation (k_{off}), maximal analyte binding capacity (R_{max}), equilibrium association constants (K_A), equilibrium dissociation constants (K_D) and Chi^2 for VHHs binding to RBD.

477 **Tables**

478 **Table 1**. Neutralizing activity of the most potent monomers of nanobodies, homodimeric and
479 heterodimeric forms of nanobodies in a microneutralization assay against live SARS-CoV-2 variants
480 of concern.

Antibody		Minimal neutralizing concentration of antibody, nM				
		B.1.1.1	Alpha (B.1.1.7)	Beta (B.1.351)	Gamma (B.1.1.28/P.1)	Delta (B.1.617.2)
Monomers	P2C5	24.04	48.08	96.15	48.08	>1500
	P2G1	12.02	48.08	96.15	48.08	24.04
	P5F8	48.08	48.08	96.15	96.15	48.08
Homodimers	(P2C5)2	0.089	0.178	0.356	0.356	>1500
	(P2G1)2	11.36	22.73	45.45	45.45	45.45
	(P5F8)2	5.68	5.68	22.72	22.72	11.36
Heterodimers	P2C5-P2G1	0.709	0.356	2.84	2.84	11.36
	P2C5-P5F8	0.178	0.089	0.356	0.356	2.85
	P2G1-P2C5	2.84	2.84	11.36	5.68	45.45
	P2G1-P5F8	45.45	45.45	181.82	45.45	90.91
	P5F8-P2C5	5.68	11.36	22.73	22.72	90.91
	P5F8-P2G1	11.36	22.72	90.91	22.72	22.73

482 **Table 2.** Primers used for generation of dimeric molecules

Primer	Sequence
PHEN1-F	CACACAGGAAACAGCTATGAC
VHH-GS-R	CCCAGGTCACTGTCTCCTCAGGTGGTGGCGGATCAGGTGGAGGTGGATCTGGCGGCGGCGGCTCAGGC
VH1-GS-F	TCTGGCGGCGGCGGCTCAGGCGGAGGAGGTTCCCAGGTGCAGCTGGTGCAGTCT
VH3-GS-F	TCTGGCGGCGGCGGCTCAGGCGGAGGAGGTTCCGAGGTGCAGCTGGTGGAGTCT
PHEN1-R	ACAACTTTCAACAGTCTAGCTCC

485 **Figure 1**

488   **Figure 2**

A

Nanobody	ELISA EC50, nM	Minimal neutralizing concentration, nM
P1G1	1.31	192
P1G9	1.32	>1540
P2C5	3.35	24
P2G1	21.39	12
P3D6	7.30	384
P4D5	2.10	769
P5C11	1.70	1540
P1B2	1.68	769
P1B4	2.26	>1540
P5A7	1.14	769
P5F8	11.73	48
P3B7	11.91	>1540
P3A9	313.3	>1540
P4H1	3.15	384

B

C

Nanobody	k_{on} (1/Ms)	k_{off} (1/s)	R_{max} (RU)	K_A (1/M)	K_D (M)	Chi^2
P2C5	1.2×10^5	4.74×10^{-4}	135	2.52×10^8	3.97×10^{-9}	8.5
P5F8	1.13×10^5	2.18×10^{-5}	90.7	5.16×10^8	1.94×10^{-9}	6.17
P2G1	1.06×10^5	5.94×10^{-4}	97.2	1.86×10^8	5.36×10^{-9}	1.74