High prevalence of *Klebsiella pneumoniae* in European food products: a multicentric study

comparing culture and molecular detection methods

Carla Rodrigues[1*], Kathrin Hauser [2], Niamh Cahill[3], Małgorzata Ligowska-Marzęta[4], Gabriella

Centorotola[5], Alessandra Cornacchia[5], Raquel Garcia Fierro[6], Marisa Haenni[6], Eva Møller

Nielsen[4], Pascal Piveteau[7], Elodie Barbier[8], Dearbháile Morris[3], Francesco Pomilio[5], Sylvain

Brisse[1*]

[1] Institut Pasteur, Université de Paris, Biodiversity and Epidemiology of Bacterial Pathogens,

Paris, France;

[2] Institute for Medical Microbiology and Hygiene, Austrian Agency for Health and Food

Safety, Vienna/Graz, Austria;

[3] Antimicrobial Resistance and Microbial Ecology Group, School of Medicine, National

University of Ireland Galway, Ireland;

[4] Statens Serum Institut, Copenhagen, Denmark;

[5] Istituto Zooprofilattico Sperimentale dell'Abruzzo e del Molise "G. Caporale", Teramo,

Italy;

[6] Unité Antibiorésistance et Virulence Bactériennes, Université Claude Bernard Lyon 1 -

ANSES, Lyon, France;

[7] INRAE, UR OPPALE, Rennes, France;

[8] Agroécologie, AgroSup Dijon, INRAE, Université Bourgogne Franche-Comté, Dijon, France

*Corresponding authors:

Sylvain Brisse

Institut Pasteur,

Biodiversity and Epidemiology of Bacterial Pathogens,

28 rue du Docteur Roux, F-75724 Paris, France.

Phone: +33 1 45 68 83 34 ; E-mail: sylvain.brisse@pasteur.fr

Carla Rodrigues

Institut Pasteur,

Biodiversity and Epidemiology of Bacterial Pathogens,

28 rue du Docteur Roux, F-75724 Paris, France.

E-mail: carla.parada-rodrigues@pasteur.fr

Keywords: *Klebsiella pneumoniae*, food sector, transmission, chicken meat, salads, antibiotic

resistance, One Health, genomics, surveillance, culture methods

Running Tittle: *K. pneumoniae* from chicken meat and salads

Abstract

Klebsiella pneumoniae species complex (KpSC) is a leading cause of multidrug-resistant

human infections. To better understand the potential contribution of food as a vehicle of

KpSC, we conducted a multicentric study to define an optimal culture method for its recovery

from food matrices, and to characterize food isolates phenotypically and genotypically.

Chicken meat (n=160) and salad (n=145) samples were collected in five European countries

and screened for KpSC presence using culture-based and ZKIR qPCR methods. Enrichment

using buffered peptone water followed by streaking on Simmons citrate agar with inositol

(44°C/48h) was defined as the most suitable selective culture method for KpSC recovery. High

prevalence of KpSC was found in chicken meat (60% and 52% by ZKIR qPCR and culture

approach, respectively) and salad (30% and 21%, respectively) samples. Genomic analyses

revealed high genetic diversity with the dominance of phylogroups Kp1 (91%) and Kp3 (6%).

82% of isolates presented a natural antimicrobial susceptibility phenotype and genotype, with

only four CTX-M-15-producing isolates detected. Notably, identical genotypes were found

across samples: same food type and same country (15 cases); different food types and same

country (1); same food type and two countries (1), suggesting high rates of transmission of

KpSC within the food sector. Our study provides a novel isolation strategy for KpSC from food

matrices and reinforces the view of food as a potential source of KpSC colonization in humans.

Importance

Bacteria of the *Klebsiella pneumoniae* species complex (KpSC) are ubiquitous and *K.*

pneumoniae (Kp) is a leading cause of antibiotic-resistant infections in humans and animals.

Despite the urgent public health threat represented by Kp, there is a lack of knowledge on

the contribution of food sources to colonization and subsequent infection in humans. This is

partly due to the absence of standardized methods for characterizing KpSC presence in food

matrices. Our multicentric study provides and implements a novel isolation strategy for KpSC

from food matrices and shows that KpSC members are highly prevalent in salads and chicken

meat, reinforcing the view of food as a potential source of KpSC colonization in humans.

Despite the large genetic diversity and the low-levels of resistance detected, the occurrence

of identical genotypes across samples suggests high rates of transmission of KpSC within the

food sector, which need to be further explored to define possible control strategies.

Introduction

Klebsiella pneumoniae (Kp), a common gut bacteria, is regarded as a critical priority pathogen (1, 2), given the depletion of therapeutic options to treat multidrug-resistant (MDR) Kp infections. Furthermore, hypervirulent community-acquired invasive infections represent a growing problem, especially in Asian countries (3). Concerning reports of the convergence of MDR and hypervirulent phenotypes are increasing (4). The clinical epidemiology of Kp is dominated by the clonal spread of important MDR and hypervirulent sublineages, such as clonal group (CG) 258, CG15, CG147 or CG23 (3).

The ecology and transmission of Kp outside the clinical setting is still underexplored; in particular, the reservoir and possible contribution of non-clinical sources in the transmission of prominent clonal groups to humans remain largely undefined. Given the broad ecological distribution of Kp in animals and the environment (5, 6), and its capacity to contaminate food (7–9), the contribution of food sources to colonization of humans, and to subsequent infection, is an important question to address. Currently, detection, isolation and identification of Kp is not well integrated in food microbiological surveillance programs. There is also a lack of standardized tools and procedures for the detection of Kp in food. To define the natural ecology and transmission of Kp, studies of the presence of Kp in food should be designed irrespective of antimicrobial resistance phenotypes.

In order to achieve precise detection and identification of Kp, it is important to define the target species. Recent taxonomic updates pointed out the existence of seven phylogroups (phylogroup 1 [Kp1] to Kp7), now defined as distinct taxa, within *Klebsiella pneumoniae sensu lato*. The resulting *K. pneumoniae* species complex *(*KpSC) consists of five different species: *K. pneumoniae sensu   stricto* (Kp1), *K. quasipneumoniae* subsp. *quasipneumoniae* (Kp2)   and

K. quasipneumoniae subsp. *similipneumoniae* (Kp4), *K. variicola* subsp. *variicola* (Kp3)   and

K. variicola   subsp. *tropica* (Kp5), '*K. quasivariicola*' (Kp6, which remains to be formally

defined), and *K. africana* (Kp7) (10–12). Recently, Barbier *et. al* developed the ZKIR qPCR (13)

for the detection of KpSC in soil and environmental samples, but its application to food

matrices was not evaluated. Regarding the isolation of *Klebsiella* spp., different selective

culture methods (7, 14, 15) have been described but comparisons of their efficiency for

Klebsiella isolation from food is lacking.

The aims of this study were (i) To compare the selectivity, productivity and specificity

of three agar media for the detection and isolation of *Klebsiella* spp., leading to propose a

standardized culture protocol for the recovery of *Klebsiella* spp. in food matrices and compare

its performance with the ZKIR qPCR; and (ii) To evaluate the presence and phenotypic and

genomic features of KpSC in two common globally consumed foodstuffs: chicken meat and

ready-to-eat salads, through a multicentric study in five European countries.

Results

Comparison of culture media

Productivity (P_R), selectivity (S_F) and specificity were calculated separately for each of

the media considered. The three media were compliant with ISO11133:2014 requirements

for productivity, as P_R was above 0.5 for almost all target strains **(Figure S3)**. However, two

strains had P_R < 0.50: *K. pneumoniae* SB132 in Chromatic Detection Agar and

K. quasipneumoniae subsp. *quasipneumoniae* SB1124, for which colony growth was only

observed on SCAI medium.

We observed that the media did not comply with the selectivity criteria ($S_F > 2$) for

most of the non-target strains considered: the S_F values ranged from -0.3 to 0.5 for *Klebsiella*

ChromoSelect Agar and from -0.5 to 0.9 for SCAI agar (Liofilchem was not tested as it is not

selective). We noted that *Klebsiella* ChromoSelect Agar was selective for *Cronobacter* spp.

and *C. freundii*, showing S_F values equal to 7.6 and 4, respectively. Hence, the media cannot

be considered as selective for *Klebsiella* according to the ISO11133:2014 requirements.

Regarding specificity, the selective media tested showed variable results regarding the

morphological features of observed colonies, based on the non-target strains considered. The

selective media, indeed, allowed the growth of some non-*Klebsiella* spp. strains, and in few

cases, colonies were morphologically similar to the target strains. Overall, because Kp

colonies were easily distinguishable on SCAI medium and because it was as performant as the

others regarding the other criteria, the SCAI medium was selected and used hereafter.

Definition of an optimized protocol for recovery of KpSC from food samples

Our initial comparisons of the four different culture protocols (**Figure S1**) showed that

the most effective method for recovery of KpSC from chicken meat was protocol 1B, which

consists of a one-step enrichment in BPW. KpSC was recovered from 7 out of 36 (19.4%)

samples using this protocol, while only 3 (8.3%) samples were positive using protocol 2

(enrichment in LB broth supplemented with ampicillin) and only 2 (5.6%) using protocol 1C

(double enrichment: enrichment in BPW followed by enrichment in LB broth supplemented

with ampicillin). No KpSC was isolated using direct plating (protocol 1A).

Further, using BPW enrichment of chicken meat (protocol 1B), a higher recovery of

KpSC was observed when streaking a 10 µl loop onto SCAI agar plates, *versus* spreading 10 µl

or 100 µl of bacterial enrichment. Of the 28 chicken meat samples examined using both

methods, KpSC was recovered from 24 (85.7%) samples using the 10 µl loop *versus* 19 (79.2%)

samples after spreading 10 µl, and 7 (29.2%) samples after spreading 100 µl, onto SCAI agar.

Thus, protocol 1B combined with streaking was used henceforth.

Evaluation of the SCAI agar plate incubation temperature

We compared two temperatures (37°C and 44°C) for the incubation of SCAI agar plates

(**Figure S2**). Collectively, 111 chicken samples were tested and 67.7% (64/111) of them were

positive for KpSC. KpSC was recovered from 59/111 (53%) samples when SCAI agar was

incubated at 37°C, as well as when incubated at 44°C, indicating no difference between

incubation temperatures (**Table S3**). However, when data from each individual institution was

analyzed, a slightly higher percentage of positive samples was recorded following incubation

of SCAI plates at 44°C at two institutions (AGES and NUIG), whereas contrasting results were

observed at the third institution (SSI) (**Table S3**). As a result, a decision was made to

implement incubation of SCAI agar plates at 44°C ± 1°C, which was incorporated into our

protocol (**Figure 2**) (dx.doi.org/10.17504/protocols.io.baxtifnn). All further food samples

were tested according to this protocol. Of note, no statistical differences were found in the

prevalence of KpSC in from free-range and not free-range chicken meat samples (**Table S3**).

Comparison of culture and ZKIR qPCR methods for KpSC detection

Chicken meat and salad leaf samples were tested for the presence of KpSC using both

the ZKIR qPCR assay (13) and the optimized culture protocol (**Figure 2**). Regarding the chicken

meat samples (n=160), the ZKIR qPCR method detected KpSC in 96 (60.0%) of the samples

(**Table 1**). In comparison, when the culture protocol was used, 83 (51.9%, *p*=0.177) of the

samples were positive for KpSC (82 *K. pneumoniae* and 1 *K. variicola* according to MALDI-TOF MS results). Regarding salad leaf samples (n=145), 29.7% (43/145) of the samples tested positive for KpSC using the ZKIR qPCR assay, versus 20.7% (30/145; *p*=0.104; 18 *K. pneumoniae* and 12 *K. variicola* according to MALDI-TOF MS results) using the culture method (**Table 1**). Considering the ZKIR qPCR assay as a reference, the optimized culture protocol showed a sensitivity (true positive rate) of 84.2%, a specificity (false positive rate) of 100%, a positive predictive value of 100%, and a negative predictive value of 86.6%.

Table 1. Comparison between the recovery rates of *K. pneumoniae* species complex using the ZKIR qPCR assay *versus* the optimized culture method.

Source	No. of samples	qPCR positive samples (%)	Culture positive samples (%)
Chicken meat	160	96 (60.0%)	83 (51.8%)
Salads	145	43 (29.7%)	30 (20.7%)
Total	**305**	**139 (45.6%)**	**113 (37.1%)**

Comparison of KpSC prevalence in chicken meat and salad samples

The overall prevalence of KpSC in chicken meat was twice as high as in salad samples (60.0% versus 29.7%; *p* < 0.00001; **Table 1**). However, prevalence differed among countries (**Figure 2**). The prevalence of KpSC in chicken meat samples ranged between 43.3% and 46.6% for three of the countries (Austria, Italy and Ireland), whereas in France and Denmark the prevalence was above 70% (72.5% and 90.0%, respectively). Regarding salad samples, the

prevalence of KpSC ranged from 3.3% in Ireland to 92% in France, with Austria, Italy and

France being the countries where the highest numbers of samples positive for KpSC were

detected.

In total, 131 KpSC isolates (92 from chicken meat and 39 from salads) were collected

from 113 positive samples using the optimized culture method; in some positive samples, up

to 5 different morphotypes were collected (**Figure 2, Table S2**).

Antimicrobial susceptibility

Antimicrobial susceptibility phenotypes were defined for the 131 KpSC isolates using

17 antimicrobial agents representing 11 antimicrobial classes (**Table S4**). 82% of the isolates

presented a wild-type phenotype, being susceptible to all the antibiotics tested except

ampicillin, for which constitutive resistance is a characteristic of all KpSC isolates. The highest

rates of resistance (detected in > 5% of the isolates) were observed for trimethoprim (12% in

chicken meat and 13% in salads), trimethoprim-sulfamethoxazole (8% for both), tetracycline

(9% in chicken meat and 5% in salads) and piperacillin-tazobactam (7% in chicken meat; **Figure**

S4; **Table S4**). Four isolates (2 from each source) were found to be resistant to extended-

spectrum cephalosporins, and ESBL gene presence in the genomic sequence was confirmed

(**Table S4**). No isolate was found to be resistant to cefoxitin, ertapenem or netilmicin. Similar

levels of resistance were detected among isolates from salads and from chicken meat.

Differences in antimicrobial resistance occurrence rates were observed among

countries (**Table S4**) but the low number of isolates implies precaution in the analysis. For

chicken meat, higher resistance rates were detected in Italy (8-50%), Austria (7-23%) and

Ireland (7-14%), whereas for salads it was in Italy (33-100%), Austria (20%) and France (4-12%)

(**Figure S4, Table S4**).

Nine (6.9%) multidrug resistant isolates were identified (**Table S5**), including the four

ESBL producers. MDR isolates were recovered in 7 chicken meat and 2 salad samples in Italy

(n=3), Austria (n=2), France (n=2) and Ireland (n=2) (**Table S5**).

Phylogenetic and genomic diversity

Based on genomic data we observed among the KpSC isolates the predominance of

Kp1 (90.8%, 119/131) followed by Kp3 (6.1%, 8/131), Kp2 (2.3%, 3/131) and Kp4 (0.8%,

1/131). The predominance of Kp1 was found in both sample types (98% in chicken meat; 74%

in salads); however, there was an association of Kp3 with salad samples (18% in salads versus

1% in chicken meat; p=0.001). Kp2 was only found in salads, whereas Kp4 was only detected

in chicken meat (**Table S6**).

Population structure analysis based on MLST and cgMLST genotyping unveiled a high

genetic diversity among the KpSC isolates, with 86 STs [19 of them defined in this study;

Simpson's index of diversity (SID) 0.989] and 107 cgSTs (SID 0.995; **Figure 3**). High-risk clonal

groups (CGs) common in the clinical setting represented 19.8% (26/131) of the isolates (**Table**

S6). Among them, CG45 (n=9), CG37 (n=5), CG17 (n=4), CG661 (n=4), CG147 (n=2), CG14 (n=1)

and CG15 (n=1) were found in chicken meat (n=18; 11.3%) and/or salad (n=6, 3.8%) samples.

A high diversity of capsular types was also found, with 71 *wzi*-alleles and 58 KL-types. In

contrast, O-type diversity was low, with O1 (25.1%; 33/131) and O2 (20.6%; 27/131)

representing almost 50% of the recovered KpSC, followed by O3 (28.2%; 37/131) (**Figure 3**).

Defining single strains (designed as genotypes) as isolates differing by no more than 5

cgMLST alleles (out of 629 loci) (16), we observed 15 strains that were isolated at least twice

across a total of 35 different samples. These were collected from the same food type (chicken

meat or salads) at different times in the same country (**Table S6, Figure S5**). To improve the

confidence inferring the close relationship between these cgMLST-based overlapped

genotypes, the SNP matrix based on 4,125 core genes obtained from Roary (representing

~75% of a typical *K. pneumoniae* genome length against 10% represented by the 629 cgMLST

genes) (16) was used, and confirmed the close genetic relatedness. For example, in France,

the same genotype (ST45/genotype 7 – 1 to 4 SNPs) was detected in 3 salad samples from

different brands (separated in time by 9 days), whereas in Ireland the same genotype

(ST4384/genotype 13 – 1 to 8 SNPs) was detected in six different chicken meat samples

(separated in time by 2 months) from different suppliers (**Table S6, Figure S5**). There was one

case of the same strain being detected in different countries, as two Kp1 ST290 isolates

(genotype 10; 15 SNPs) were detected in chicken meat samples from Denmark (April 2019)

and Austria (September 2019) (**Figure S5**). We also detected a single strain between chicken

meat and salads (ST1537/genotype 9 – 23 SNPs) between May and September 2019,

respectively, in France (**Table S6**).

In most isolates (82%; 108/131) only intrinsic $bla_{SHV/OKP/LEN}$ genes was detected,

consistent with a natural susceptibility genotype (**Table S6**). Few antibiotic resistance genes

were observed across the 131 isolates, with genes of non-prominent clinical significance *sul*

and *dfr*, (9.2%; 12/131 both), *tet* (8.4%; 11/131), *aadA* and *strAB* (6.1%; 8/131 both) being the

most frequent ones. However, four CTX-M-15 producing isolates were detected in two salad

samples from France and two chicken meat samples from Italy and Ireland. The genotype of

MDR isolates was compatible with their phenotype (**Table S5**). No carbapenemase or colistin

resistance genes or mutations were detected. In 45.8% (60/131) of the isolates, neither

plasmid replicons nor plasmid-encoded heavy metal tolerance genes were detected. In strains

where plasmid replicons were detected, the plasmids known to occur naturally in Kp, such as

IncFIB and IncFII$_K$, were the most prevalent (25.2% and 16.8%, respectively). Regarding

virulence genes, the yersiniabactin gene cluster was detected in 14.5% (19/131) of the isolates

and no hypervirulent genotypes (as defined by the presence of *rmpA* or aerobactin or

salmochelin) were detected.

Discussion

Despite the renewed interest in *Klebsiella pneumoniae* complex (KpSC) epidemiology, boosted by the increasing involvement of *K. pneumoniae sensu stricto* (Kp1) in human infections associated with high-levels of antibiotic resistance and/or virulence (1, 3), the contribution of non-clinical sources, such as food products, to the current emergence of KpSC is poorly understood. This is partly due to the lack of standardized protocols for the detection and isolation of Kp strains in environmental, food or animal samples. Different selective culture media have been previously developed for *Klebsiella* spp.. Among the most recognized are MacConkey-Inositol-Carbenicillin agar (MCIC) (17) and its variation containing adonitol (MCICA,) (18), SCAI (14, 19) and Brilliant green containing Inositol-Nitrate-Deoxycholate agar (BIND) (20). All of them rely on the ability of *Klebsiella* spp. to use inositol as carbon source, combined with the use of secondary selective compounds (e.g., citrate as carbon source in the case of SCAI; nitrate as nitrogen source in BIND; carbenicillin and adonitol in MCIC/MCICA, although only Kp1 and Kp2/Kp4 are able to metabolize adonitol) (11). In our study, we decided to evaluate the performance of three different culture media for the detection and isolation of *Klebsiella* spp. in accordance with ISO 11133:2014 and latest amendments. They included the non-commercial SCAI medium and two commercial culture media: the *Klebsiella* ChromoSelect Selective Agar produced by Sigma-Aldrich, that similarly to MCIC uses carbenicillin as selective compound (https://www.sigmaaldrich.com/deepweb/assets/sigmaaldrich/product/documents/240/58 0/90925dat.pdf); and Chromatic Detection agar developed by Liofilchem, which is a chromogenic non-selective medium allowing the identification of *Klebsiella* spp., *Enterobacter* spp. and *Serratia* spp. based on the green-blue color of colonies

(http://www.liofilchem.net/login/pd/ifu/11611_IFU.pdf). The productivity tests of the three

media did not show any critical issues, whereas none of the media tested fulfilled the

selectivity and specificity criteria defined by ISO 11133:2014. Nevertheless, the distinction of

Klebsiella spp. colonies from other species was achieved in general. This is a critical point, as

the routinely used media (e.g., MacConkey or CLED agar) do not provide easy differentiation

of *Klebsiella* spp. colonies from other species, such as *Citrobacter* spp., *Pantoea* spp. or

Serratia spp. Here, the performance of the tested culture media showed similar results and

SCAI was selected, due to its low cost and easy in-house preparation. The use of this media in

recent *Klebsiella* spp. epidemiological studies may also represent an advantage for global and

cross-sector comparisons (13, 21–23).

Once SCAI agar was selected, different protocols to isolate *Klebsiella spp*. from food

matrices were evaluated. Multiple conditions, involving pre-enrichment, enrichment,

incubation temperature and plating were assessed from chicken meat samples. The

optimized culture protocol was then compared with the reference ZKIR qPCR assay in a

multicentric design study involving chicken meat and salad samples (13). Overall, KpSC was

detected at a higher rate in both chicken meat and salad samples using the ZKIR qPCR method,

but even so the optimized culture protocol demonstrated good sensitivity (84%) and

specificity (100%) rates.

Approximately 50% of the food samples tested were KpSC positive, with significantly

higher occurrence in chicken meat compared to salads, although disparities in prevalence

were observed according to the country analyzed. It is difficult to contextualize our findings

in a global scenario since few studies have addressed KpSC prevalence in these type of

products without previous antibiotic selection: most of the studies addressing the prevalence

and characterization of KpSC from foodstuffs have focused on colistin resistant or ESBL- or

carbapenemase-producing *K. pneumoniae* (8, 24–26). As a consequence, knowledge on the

ecology and natural diversity of Kp populations has been very limited. Furthermore,

identification of the bacterial species was imprecise in most previous studies, in light of recent

taxonomic changes. Our broader approach also captures susceptible strains and led to several

important observations. First, the KpSC prevalence in chicken meat was higher than

previously described: 47% in USA (Arizona) in 2012 (7); 30% in Turkey in 2007-2008 (27); 14%

in China (Shijiazhuang) in 2013-2014 (28). Second, despite the high prevalence of KpSC in

chicken meat samples, ESBL-producing *K. pneumoniae* were only detected in 1.3% (2 CTX-M-

15) of the samples. These numbers are similar to those reported in other studies that targeted

ESBL strains directly by selective approaches (0-5%) (8, 26, 28–30). Noteworthy, the presence

of ESBL in chicken meat was scarce in *K. pneumoniae* when compared with *Escherichia coli*

(55%-92%) in recent European reports (30–33).

Third, the highest resistance rates detected in natural food Kp populations were to

tetracyclines, trimethoprim and trimethoprim-sulfamethoxazole, reflecting the recent

reports of sales of these antimicrobial classes in food-producing animals in different European

countries (34). Fourth, despite the predominance of Kp1 and the high genetic diversity found

among chicken meat isolates, some genotypes overlapped across samples. The relation of

these common genotypes (≤ 5 alleles mismatches) with the core-genome alignment obtained

from Roary was also explored, and in most of the cases, less than 21 SNPs were identified, a

threshold recently proposed for Kp ST258 to discriminate hospital clusters (35). Here, we

found a maximum of 42 SNPs detected within the same genotype. Some of these multi-

sample genotypes (ST290/genotype 10 recovered in Austria and Denmark and ST45/genotype

15 recovered in Denmark) were previously found circulating among healthy poultry in France

in 2015 (Rodrigues C and Haenni M., pers.comm.). The repeated isolation of single strains in

unrelated samples across time suggests transmission from the same source/reservoir or

transmission within chicken flocks during production. Importantly, MDR ST45/genotype 11

found in Italy has also been found in urinary tract infections among Italian patients in 2018

(22), suggesting a contribution of food KpSC to human infections (7, 24).

Salads are typically eaten raw, with a high risk of ingestion of live Kp bacteria if present.

The prevalence of KpSC among ready-to-eat salads was approximatively 30%, higher than

what was reported in the few comparable studies (6-15%) (36–39). ESBL-producers were

found in 1.4% of the samples (2 CTX-M-15), a lower prevalence than what is described in the

literature (15-25%) targeting ESBL strains (39–42). As observed for chicken meat samples, the

same genotype was observed within the same country, in salads from the same or different

chain supplier/supermarket, suggesting a common source of the vegetables (e.g., different

brands supplied by a common farm) or cross-contamination during their growth and

processing (e.g., use of untreated irrigation water, wild-life animals, contamination of fields

with manure, human contamination during packaging). Furthermore, one case of genotype

overlap was found between salads and chicken meat isolates from samples from France,

suggesting local transmission.

A previous study of risk factors of Kp carriage in community settings has linked the

consumption of raw vegetables and contact with chicken to MDR KpSC carriage in humans

(21). These findings together with our results, which are based on a small sampling survey,

highlight the possible role of food as a source of Kp and call for much broader studies to

address the ecology and transmission of this generalist pathogen in the food sector.

In our sampling, sublineages defined as high-risk clonal groups (3) represented 20% of

the isolates recovered. One of these, ST45 (in both type of samples) was highly prevalent, as

well as ST290 (in chicken meat), and the potential risk they pose should be addressed in future

studies. Both STs have already been linked to human infections (22, 43, 44) and human

carriage (21, 22) and also to broilers and pigs production (23, 45), raising the question of the

zoonotic potential of these STs.

Conclusions

In conclusion, our study addressed important knowledge gaps relating to Kp in food

sources. First, we addressed the lack of harmonized culture protocols for the detection and

isolation of *Klebsiella* spp. from food matrices, with the development of an optimized SCAI-

based culture protocol, which in the future may be more widely adopted to enable global

comparison regarding KpSC prevalence in these types of sources. Second, the high prevalence

of KpSC in chicken meat and ready-to-eat salads highlights the possible role of food as a

source of human colonization and infection by KpSC. Even though KpSC strains detected in

food products remain largely susceptible to antimicrobials, understanding the degree to

which food contamination by Kp contributes to human infections is an important topic for

future research in the One Health perspective.

Material and Methods

Comparison of productivity, selectivity and specificity of agar media

Three different selective solid media for *Klebsiella* spp. growth were tested in accordance with ISO11133:2014 (Microbiology of food, animal feed and water — Preparation, production, storage and performance testing of culture media; https://www.iso.org/standard/53610.html). Nutrient agar (Microbiol & C., Cagliari, Italy) was used as the non-selective reference medium. For productivity, selectivity and specificity assays a set of 58 strains [50 from Institut Pasteur and 8 from Istituto Zooprofilattico Sperimentale dell'Abruzzo e del Molise Giuseppe Caporale (IZSAM)] were used: 51 reference strains of *Klebsiella* spp. and closely related species (*Raoultella* spp.), which included representatives of the six main KpSC phylogroups, and 7 non-target strains (**Table S1**). For productivity (P_R), the three following media were tested: Simmons Citrate Agar with Inositol (SCAI) (14), *Klebsiella* ChromoSelect Selective agar produced by Sigma-Aldrich (Missouri, USA) and Chromatic Detection agar developed by Liofilchem (Roseto degli Abruzzi, Italy). The SCAI medium is not available commercially and was prepared in-house following previous protocols (14, 19) (https://www.protocols.io/view/isolation-of-klebsiella-strains-from-human-or-anim-662hhge/materials). P_R was calculated using 50 reference strains belonging to *Klebsiella* spp. and closely related species (*Raoultella* spp.) (**Table S1**). After incubation at 37°C for 24 h, broth cultures in Brain Heart Infusion (BHI) (Biolife, Milan, Italy) of each strain were diluted up to about 100 CFU/ml and spread on media. After incubation at 37°C for 48h, colonies compatible with the *Klebsiella* phenotype were enumerated. P_R is calculated as the ratio between the number of colonies grown on the selective agar medium and the number

390   of colonies grown on the non-selective medium, and according to ISO11133:2014, P_R values

391   should be ≥ 0.50 to consider the media productivity acceptable.

392   For selectivity (S_F), 7 non-target strains (*Cronobacter* spp., *Citrobacter koseri,*

Citrobacter freundii, *Serratia marcescens, Serratia liquefaciens, Serratia rubidaea* and *1*

Pantoea agglomerans) were tested on the SCAI and *Klebsiella* ChromoSelect Selective agar

(**Table S1**). After incubation at 37°C for 24h in BHI, broth cultures were ten-fold diluted serially

and each dilution was spread onto plates. After incubation at 37°C for 48 h, colony growth

was observed. S_F is defined as the difference between the highest dilution showing growth on

the non-selective reference medium ($\log_{10}$) and the highest dilution showing comparable

growth on the selective test medium ($\log_{10}$). In accordance with ISO11133:2014, S_F values of

non-target strains should be at least equal to 2 $\log_{10}$ growth differences, which reflects the

ability of the medium to partially or totally inhibit their growth.

402   For specificity, the 7 non-target strains and two control strains (1 *K. pneumoniae* and

1 *R. ornithinolytica*) were tested on the three media (**Table S1**). Broth cultures were prepared

in Buffered Peptone Water (BPW) (Biolife, Milan, Italy), incubated at 37°C for 24h, and spread

onto each medium considered. After incubation at 37°C for 48h of each medium, plates were

observed for the presence of typical and suspected colonies, recording if the characteristics

of the non-target colonies could be similar to the target ones.

Initial evaluation of protocols for recovery of *Klebsiella pneumoniae* from food

410   An initial assessment of four protocols for recovery of KpSC from chicken meat was

carried out. In total, 36 chicken meat samples [12 free-range (7 skin-on and 5 skin-off) and 24

non-free range (12 skin-on and 12 skin-off) samples] were collected together with the

following metadata: manufacturer, country of origin, and batch number. Each chicken meat

sample was cut into small slivers to a final weight of 50 g. This 50 g sample was divided into

two portions of 25 g and processed as follows and as illustrated in **Figure S1**:

<u>Initial suspension 1</u>: the first 25 g portion was diluted in 225 ml (1:10 dilution) of BPW and

the sample was homogenized using a stomacher/blender. Prior to incubation of initial

suspension 1, 100 µl of the suspension was cultured directly on SCAI medium using a sterile

spreader or sterile loop and incubated at 37°C ± 1°C for 48h ± 1 h (**Figure S1 - 1A** - detection

without enrichment). The remainder of initial suspension 1 was incubated at 37°C ± 1°C for

24h ± 1 h. Following 24h incubation of initial suspension 1, 100 µl of the suspension was

streaked on SCAI medium and incubated at 37°C ± 1°C for 48 h ± 1 h (**Figure S1 - 1B** - detection

with BPW enrichment only). Of the remaining incubated suspension 1, 1 ml was inoculated

into 9 ml of Lysogeny broth (LB) with ampicillin (final concentration 0.01 mg/mL) and

incubated at 37 °C ± 1 °C for 24h± 1 h. Following 24h incubation, 100 µl of the suspension was

streaked on SCAI medium and incubated at 37°C ± 1°C for 48h ± 1 h (**Figure S1 - 1C** - detection

with double enrichment).

<u>Initial suspension 2</u>: the second 25 g portion was diluted in 225 ml (1:10 dilution) of LB broth

supplemented with ampicillin (final concentration 0.01 mg/mL) and the sample was

homogenized using a stomacher/blender and incubated at 37 °C ± 1°C for 24 h ± 1 h. Following

24 h incubation, 100 µl of the suspension was cultured on SCAI medium and incubated at 37°C

± 1°C for 48 h ± 1 h (Figure **S1 - 2** detection with LB + Ampicillin enrichment only).

Typically, *Klebsiella* spp. appear yellow on SCAI medium (14). At least five suspected

Klebsiella colonies were collected from each SCAI plate and sub-cultured onto SCAI media or

a non-selective agar for identification using MALDI-TOF MS (Bruker Daltonics, Bremen,

Germany). Currently, databases only allow for the identification of *K. pneumoniae* or *K.*

variicola. Based on literature and in our test strain set, Kp1, Kp2 and Kp4 strains are identified

as *K. pneumoniae*, whereas Kp3, Kp5 and Kp6 are currently identified as *K. variicola* (46, 47).

For 28/36 chicken meat samples, the impact on KpSC recovery of culturing using a 10

µl loop versus the spreading of 10 µl and 100 µl of enrichments onto SCAI media was also

compared.

Evaluation of the optimized protocol and definition of an optimal temperature of

incubation

Following this initial assessment process, the optimized protocol as outlined below

and as illustrated in **Figure S2** was evaluated for recovery of KpSC from chicken meat samples,

using a total of 111 samples of chicken meat [51 free-range (25 skin-on and 26 skin-off) and

60 non-free range (30 skin-on and 30 skin-off)].

Each sample was cut into small slivers to a final weight of 25 g. This was added to 225

ml of BPW broth (1:10 dilution) and the sample was mixed using a stomacher or blender. The

suspension was then incubated at 37°C ± 1°C for 24h ± 1 h. Following incubation, the

suspension was cultured for single colonies on SCAI agar using a 10 µl loop and incubated at

two different temperatures, 37°C ± 1°C and at 44°C ± 1°C for 48 h ± 1 h (**Figure S2**), in order

to establish an optimal temperature of incubation of SCAI medium for recovery of Kp from

food matrices.

Comparison of the optimized SCAI culture protocol and ZKIR qPCR

A comparison of the optimized SCAI culture protocol and the ZKIR qPCR for the detection and isolation of KpSC in food matrices was carried out in 5 European institutions, including the Austrian Agency for Health and Food Safety (AGES), French Agency for Food, Environmental and Occupational Health & Safety (ANSES), National University of Ireland Galway (NUIG), Istituto Zooprofilattico Sperimentale dell'Abruzzo e del Molise Giuseppe Caporale (IZSAM) and the Statens Serum Institute (SSI, Denmark).

Each institution was asked to collect and test, where possible, an additional 30 samples of chicken meat, as well as 30 pre-packaged, pre-washed salad leaf samples (**Table S2**). In total, 160 chicken meat samples and 145 salad samples, collected between December 2018 and September 2019, were tested using the updated optimized protocol (dx.doi.org/10.17504/protocols.io.baxtifnn) for recovery of KpSC from food matrices (**Figure 1**).

The ZKIR qPCR was used to detect the presence of KpSC in food samples (13). After enrichment, DNA was prepared for ZKIR qPCR. 500 µl of the enrichment broth were centrifuged (5 min at 5800 × g), washed with sterile water, and resuspended in 500 µl sterile water before boiling for 10 min. qPCR conditions were as previously described (13) (dx.doi.org/10.17504/protocols.io.7n6hmhe).

Antimicrobial susceptibility testing

Antimicrobial susceptibility testing was performed on 131 KpSC isolates using either disk diffusion or microbroth dilution [GN2F panels (Sensititre, ThermoFischer Scientific)] methods. Clinical breakpoints were used for interpretation according to EUCAST guidelines

(https://www.eucast.org/fileadmin/src/media/PDFs/EUCAST_files/Breakpoint_tables/v_9.0

Breakpoint Tables.pdf), except for kanamycin and streptomycin (48). *Escherichia coli*

ATCC25922 was used as control strain. The panel of antimicrobials tested included:

amoxicillin/clavulanic acid (20/10 µg), piperacillin-tazobactam (both using disks of 30/6 µg, or

GN2F plates), cefpodoxime (10 µg, GN2F) or cefotaxime (5 µg), cefoxitin (30 µg, GN2F),

aztreonam (30 µg), ertapenem (10 µg), amikacin (30 µg, GN2F), gentamicin (10 µg, GN2F),

netilmicin (10 µg), tobramycin (10 µg, GN2F), kanamycin (10 µg), streptomycin (30 µg),

ciprofloxacin (5 µg, GN2F) or ofloxacin (5 µg), nalidixic acid (30 µg), trimethoprim (5 µg),

trimethoprim-sulfamethoxazole (23.75 + 1.25 µg, GN2F), and tetracycline (30 µg). The

presence of multidrug resistant isolates, which are defined as those resistant to at least one

agent in 3 different classes of antimicrobials (49), was analyzed.

Whole-genome sequencing and comparative genomic analyses

Whole-genome sequencing was performed on the 131 Kp isolates. Genomic DNA

libraries were prepared using the Nextera XT Library Prep Kit (Illumina, San Diego, USA)

following the manufacturer's protocol. Illumina sequencing was performed at the five

partners: AGES and SSI using MiSeq (2 x 250 bp paired-end sequencing; n=18 and n=31

isolates respectively); ANSES and NUIG using NovaSeq 6000 (2 x 150 bp paired-end

sequencing; n=52 and n=15, respectively); and IZSAM using NextSeq 500/550 (2 x 150 bp

paired-end sequencing; n=15). Genomic assemblies were obtained using SPAdes v3.9 (50) or

Velvet v1.2.10 (51) and were annotated using Prokka v1.12 (52).

Multilocus sequence typing (MLST) (53), core-genome MLST (cgMLST) and cgLIN codes

(16)   were   determined   using   the   BIGSdb-Kp   database

(https://bigsdb.pasteur.fr/klebsiella/klebsiella.html). This web tool and Kleborate (54)

(https://github.com/katholt/Kleborate) were used to look for antimicrobial resistance,

virulence, and heavy metal tolerance genes and to characterize the capsular synthesis gene

cluster. Plasmid replicons were searched using PlasmidFinder (55)

(https://cge.cbs.dtu.dk/services/PlasmidFinder/).

For phylogenetic analyses, a core-genome alignment based on the concatenation of

4,125 core genes was obtained with Roary v3.12 (56) using a blastP identity cut-off of 80%

and core genes defined as those being present in more than 90% of the isolates.

Recombination events were removed using Gubbins v2.2.0 (57), generating a recombination-

free alignment comprising 486,290 single-nucleotide variants (SNVs). This recombination-free

alignment was used to construct a maximum likelihood phylogenetic tree using IQ-TREE

v1.6.11 (model GTR+F+ASC+R5).

Chi square or Fisher exact test were used to compare the prevalence of KpSC in the

different sources and to check the association of the different categorical variables (P values

<0.05 were considered statistically significant).

Data availability

Detailed optimized SCAI culture protocol for the detection and isolation of KpSC in food

matrices was made publicly accessible to the scientific community in January 2020 through

the protocols.io platform (dx.doi.org/10.17504/protocols.io.baxtifnn). Genomic sequences

generated in this study were submitted to the European Nucleotide Archive and are

accessible under the BioProject number PRJEB34643, and are also publicly available in BIGSdb

through project ID 37 'MedVetKlebs_multicentric study'.

Authors Contribution

SB, CR, DM, KH, EMN and FP conceptualized the study. GC, AC and FP from IZSAM performed the culture media evaluation. KH, MLM, EMN, NC and DM performed the initial assessments for the establishment of an optimized culture protocol for recovery of KpSC from food matrices. KH, NC, MLM, GB, AC and RGF performed the comparison of culture media and ZKIR qPCR assay. EB and PP provided laboratory support and validated the analysis of the ZKIR qPCR assay. CR ad SB analyzed all the genomic data, coordinated the study and wrote the initial draft manuscript. SB acquired funding. All authors contributed to writing and editing the manuscript and reviewed the final version.

Acknowledgements

We thank to James Bray and Keith Jolley for technical assistance with genomic assemblies of KpSC isolates from Ireland. We thank the Institut Pasteur teams for the curation and maintenance of BIGSdb-Pasteur databases at http://bigsdb.pasteur.fr/.

Funding

This work was supported financially by the MedVetKlebs project, a component of European Joint Programme One Health EJP, which has received funding from the European Union's Horizon 2020 research and innovation programme under Grant Agreement No 773830. CR was also financed by a Roux-Cantarini grant from Institut Pasteur.

Authors license statement

Figure legends

Figure 1. Optimized protocol for the recovery and isolation of *Klebsiella* spp. from food

matrices.

Figure 2. Prevalence of *K. pneumoniae* species complex by type of sample and by country.

Figure 3. Maximum likelihood phylogeny based on 4,125 concatenated core genes generated

using IQ-Tree with the GTR+F+ASC+R5 model.

The scale bar indicates the number of nucleotide substitutions per site. Black dots on main

nodes indicate bootstrap values ≥95%. Tree tips are colored by sample type (see key). The

grey boxes delineate high-risk clonal groups (CG), as specified. European countries where the

samples were collected are colored as indicated in the legend. The presence of antimicrobial

resistance and virulence genes, and plasmid replicons is indicated.

Supplementary Material

Table S1. Bacterial reference strains used for agar media performance comparison.

Strains used for the different evaluation criteria are indicated. The correspondent strain code

for productivity results in Figure S3 is indicated.

Table S2. Chicken meat and salad samples analyzed in the different European countries for

the presence of *K. pneumoniae* species complex and its epidemiologic characteristics.

Table S3. Comparison between the recovery rates of *K. pneumoniae* species complex using

two different incubation temperatures of SCAI medium.

Table S4. Number of *K. pneumoniae* species complex resistant isolates for each of the

antimicrobials tested distributed by sampling source and by country.

Table S5. Multidrug resistant strains from chicken meat and salads, sequence-type (ST),

phenotypic resistance and identified AMR genes based on WGS.

Table S6. Characteristics of the 131 *K. pneumoniae* species complex isolates recovered:

population structure, surface polysaccharides, antimicrobial susceptibility, antimicrobial

resistance genes, virulence genes, metal tolerance gene clusters and plasmid replicons.

Figure S1. Evaluation of different protocols for recovery of *Klebsiella* spp. from food matrices.

Figure S2. Protocol to define the optimal temperature of incubation of SCAI medium plates

for the recovery of *Klebsiella* spp. from food matrices.

Figure S3. Productivity (P_R) results for the three media considered using a reference panel of

50 *Klebsiella* spp. and closely related species.

Figure S4. Antibiotic resistance of the *K. pneumoniae* species complex isolates recovered by

sample type and by country.

Upper panel: The x-axis represents the number of KpSC resistant isolates for each of the

antimicrobials tested and percentages (%) displayed on the graph were calculated based on

the total number of isolates.

Lower panel: The x-axis represents the number of KpSC resistant isolates for each of the

antimicrobials tested and percentages (%) displayed on the graph were calculated based on

the total number of isolates in each country.

Figure S5. Geographic distribution of the common genotypes found among *K. pneumoniae*

species complex recovered from food products.

References

1.   CDC. 2013. Antibiotic Resistance Threats in the United States, 2013.

2.   WHO. 2014. Antimicrobial resistance: global report on surveillance 2014.

3.   Wyres KL, Lam MMC, Holt KE. 2020. Population genomics of *Klebsiella pneumoniae*.

Nat Rev Microbiol. 18:344–359.

4.   Wyres KL, Wick RR, Judd LM, Froumine R, Tokolyi A, Gorrie CL, Lam MMC, Duchêne S,

Jenney A, Holt KE. 2019. Distinct evolutionary dynamics of horizontal gene transfer in

drug resistant and virulent clones of *Klebsiella pneumoniae*. PLoS Genet 15:e1008114.

5.   Brisse S, Grimont F, Grimont PAD. 2006. The Genus *Klebsiella*. J Bacteriol 51:159–196.

6.   Martin RM, Cao J, Brisse S, Passet V, Wu W, Zhao L, Malani PN, Rao K, Bachman MA.

2016. Molecular Epidemiology of Colonizing and Infecting Isolates of *Klebsiella*

pneumoniae. mSphere 1:1–12.

7.   Davis GS, Waits K, Nordstrom L, Weaver B, Aziz M, Gauld L, Grande H, Bigler R,

Horwinski J, Porter S, Stegger M, Johnson JR, Liu CM, Price LB. 2015. Intermingled

Klebsiella pneumoniae Populations Between Retail Meats and Human Urinary Tract

Infections. Clin Infect Dis 61:892–899.

8.   Ludden C, Moradigaravand D, Jamrozy D, Gouliouris T, Blane B, Naydenova P,

Hernandez-Garcia J, Wood P, Hadjirin N, Radakovic M, Crawley C, Brown NM, Holmes

M, Parkhill J, Peacock SJ. 2020. A One Health Study of the Genetic Relatedness of

Klebsiella pneumoniae and Their Mobile Elements in the East of England. Clin Infect Dis

70:219–226.

9.   Zadoks RN, Griffiths HM, Munoz MA, Ahlstrom C, Bennett GJ, Thomas E, Schukken YH.

2011. Sources of *Klebsiella* and *Raoultella* species on dairy farms: Be careful where you

walk. J Dairy Sci 94:1045–1051.

10.   Brisse S, Passet V, Grimont PAD. 2014. Description of *Klebsiella quasipneumoniae* sp.

nov., a novel species isolated from human infections, with two subspecies *Klebsiella*

quasipneumoniae subsp. *quasipneumoniae* subsp. nov. and *Klebsiella*

quasipneumoniae subsp. *similipneumoniae* subsp. nov., and a. Int J Syst Evol Microbiol

64:3146–3152.

11.   Rodrigues C, Passet V, Rakotondrasoa A, Diallo TA, Criscuolo A, Brisse S. 2019.

Description of *Klebsiella africanensis* sp. nov., *Klebsiella variicola* subsp. *tropicalensis*

subsp. nov. and *Klebsiella variicola* subsp. *variicola* subsp. nov. Res Microbiol 170:165–

170.

12.   Wesley Long S, Linson SE, Saavedra MO, Cantu C, Davis JJ, Brettin T, Olsen RJ. 2017.

Whole-Genome Sequencing of a Human Clinical Isolate of the Novel Species *Klebsiella*

quasivariicola sp. nov. Genome Announc 5.

13.   Barbier E, Rodrigues C, Depret G, Passet V, Gal L, Piveteau P, Brisse S. 2020. The ZKIR

assay, a real-time PCR method for the detection of *Klebsiella pneumoniae* and closely

related species in environmental samples. Appl Environ Microbiol 86.

14.   Van Kregten E, Westerdaal NAC, Willers JMN. 1984. New, simple medium for selective

recovery of *Klebsiella pneumoniae* and *Klebsiella oxytoca* from human feces. J Clin

Microbiol 20:936–941.

15.   Filius PMG, van Netten D, Roovers PJE, Vulto AG, Gyssens IC, Verbrugh HA, Endtz HP.

2003. Comparative evaluation of three chromogenic agars for detection and rapid

identification of aerobic Gram-negative bacteria in the normal intestinal microflora.

Clin Microbiol Infect 9:912–918.

16.   Hennart M, Guglielmini J, Maiden MCJ, Jolley KA, Criscuolo A, Brisse S. 2021. A dual

barcoding approach to bacterial strain nomenclature: Genomic taxonomy of *Klebsiella*

pneumoniae strains. bioRxiv 2021.07.26.453808.

17.   Bagley S, Seidler R. 1978. Primary *Klebsiella* identification with MacConkey-inositol-

carbenicillin agar. Appl Environ Microbiol 36:536–538.

18.   Gao H, Gao Q-L, Zhang X, Guan C, Luo M-H, Zhang H-B, Liu P, Zhang H-Y, Li J. 2010.

Improved Medium For Detection of *Klebsiella* In Powdered Milk. J Food Saf 30:12–23.

19.   Passet V, Brisse S. 2015. Association of tellurite resistance with hypervirulent clonal

groups of *Klebsiella pneumoniae*. J Clin Microbiol 53:1380–1382.

20.   Ohtomo R, Saito M. 2003. A New Selective Medium for Detection of *Klebsiella* from

Dairy Environments. Microbes Env 18:138–144.

21.   Huynh B-T, Passet V, Rakotondrasoa A, Diallo T, Kerleguer A, Hennart M, Lauzanne A

De, Herindrainy P, Seck A, Bercion R, Borand L, Pardos de la Gandara M, Delarocque-

Astagneau E, Guillemot D, Vray M, Garin B, Collard J-M, Rodrigues C, Brisse S. 2020.

Klebsiella pneumoniae carriage in low-income countries: antimicrobial resistance,

genomic diversity and risk factors. Gut Microbes 11:1287–1299.

22.   Thorpe H, Booton R, Kallonen T, Gibbon MJ, Couto N, Passet V, Fernandez JSL,

Rodrigues C, Matthews L, Mitchell S, Reeve R, David S, Merla C, Corbella M, Ferrari C,

Comandatore F, Marone P, Brisse S, Sassera D, Corander J, Feil EJ. 2021. One Health or

Three? Transmission modelling of *Klebsiella* isolates reveals ecological barriers to

transmission between humans, animals and the environment. bioRxiv

2021.08.05.455249.

23. Franklin-Alming F V., Kaspersen H, Hetland MAK, Bakksjø R-J, Nesse LL, Leangapichart

T, Löhr IH, Telke AA, Sunde M. 2021. Exploring *Klebsiella pneumoniae* in Healthy Poultry

Reveals High Genetic Diversity, Good Biofilm-Forming Abilities and Higher Prevalence

in Turkeys Than Broilers. Front Microbiol 0:2609.

24. Davis GS, Price LB. 2016. Recent Research Examining Links Among *Klebsiella*

pneumoniae from Food, Food Animals, and Human Extraintestinal Infections. Curr

Environ Heal reports 3:128–135.

25. Chaalal N, Touati A, Bakour S, Aissa MA, Sotto A, Lavigne J-P, Pantel A. 2021. Spread of

OXA-48 and NDM-1-Producing Klebsiella pneumoniae ST48 and ST101 in Chicken Meat

in Western Algeria. Microb Drug Resist 27:492–500.

26. Huizinga P, Bergh MK den, Rossen JW, Willemsen I, Verhulst C, Savelkoul PHM,

Friedrich AW, García-Cobos S, Kluytmans J. 2019. Decreasing prevalence of

contamination with extended-spectrum beta-lactamase-producing

Enterobacteriaceae (ESBL-E) in retail chicken meat in the Netherlands. PLoS One

14:e0226828.

27. Gundogan N, Citak S, Yalcin E. 2011. Virulence Properties of Extended Spectrum b-

Lactamase-Producing *Klebsiella* Species in Meat Samples. J Food Prot 74:559–564.

28. Guo Y, Zhou H, Qin L, Pang Z, Qin T, Ren H, Pan Z, Zhou J. 2016. Frequency, Antimicrobial

Resistance and Genetic Diversity of *Klebsiella pneumoniae* in Food Samples. PLoS One

11:e0153561.

29. Cardozo MV, Liakopoulos A, Brouwer M, Kant A, Pizauro LJL, Borzi MM, Mevius D, de

Ávila FA. 2021. Occurrence and Molecular Characteristics of Extended-Spectrum Beta-

Lactamase-Producing Enterobacterales Recovered From Chicken, Chicken Meat, and

Human Infections in Sao Paulo State, Brazil. Front Microbiol 0:901.

30.   Díaz-Jiménez D, García-Meniño I, Fernández J, García V, Mora A. 2020. Chicken and

turkey meat: Consumer exposure to multidrug-resistant Enterobacteriaceae including

mcr-carriers, uropathogenic *E. coli* and high-risk lineages such as ST131. Int J Food

Microbiol 331:108750.

31.   Day MJ, Hopkins KL, Wareham DW, Toleman MA, Elviss N, Randall L, Teale C, Cleary P,

Wiuff C, Doumith M, Ellington MJ, Woodford N, Livermore DM. 2019. Extended-

spectrum β-lactamase-producing *Escherichia coli* in human-derived and foodchain-

derived samples from England, Wales, and Scotland: an epidemiological surveillance

and typing study. Lancet Infect Dis 19:1325–1335.

32.   Casella T, Nogueira MCL, Saras E, Haenni M, Madec JY. 2017. High prevalence of ESBLs

in retail chicken meat despite reduced use of antimicrobials in chicken production,

France. Int J Food Microbiol 257:271–275.

33.   Huizinga P, Van Den Bergh MK, Rossen JW, Willemsen I, Verhulst C, Savelkoul PHM,

Friedrich AW, García-Cobos S, Kluytmans J. 2019. Decreasing prevalence of

contamination with extended-spectrum beta-lactamase-producing

Enterobacteriaceae (ESBL-E) in retail chicken meat in the Netherlands. PLoS One

14:e0226828.

34.   European Medicines Agency. 2020. Tenth ESVAC report: Sales of veterinary

antimicrobial agents in 31 European countries in 2018. Amsterdam.

35.   David S, Reuter S, Harris SR, Glasner C, Feltwell T, Argimon S, Abudahab K, Goater R,

Giani T, Errico G, Aspbury M, Sjunnebo S, Feil EJ, Rossolini GM, Aanensen DM, Grundmann H. 2019. Epidemic of carbapenem-resistant *Klebsiella pneumoniae* in Europe is driven by nosocomial spread. Nat Microbiol 2019 411 4:1919–1929.

36. Harada T, Taguchi M, Kawahara R, Kanki M, Kawatsu K. 2018. Prevalence and Antimicrobial Susceptibility of Bacterial Pathogens in Ready-to-Eat Foods Retailed in Osaka Prefecture, Japan. J Food Prot 81:1450–1458.

37. Falomir MP, Rico H, Gozalbo D. 2013. *Enterobacter* and *Klebsiella* Species Isolated from Fresh Vegetables Marketed in Valencia (Spain) and Their Clinically Relevant Resistances to Chemotherapeutic Agents. Foodborne Pathog Dis 10:1002–1007.

38. Boehme S, Werner G, Klare I, Reissbrodt R, Witte W. 2004. Occurrence of antibiotic-resistant enterobacteria in agricultural foodstuffs. Mol Nutr Food Res 48:522–531.

39. Giri S, Kudva V, Shetty K, Shetty V. 2021. Prevalence and Characterization of Extended-Spectrum β-Lactamase-Producing Antibiotic-Resistant *Escherichia coli* and *Klebsiella pneumoniae* in Ready-to-Eat Street Foods. Antibiotics 10:850.

40. Zurfluh K, Nüesch-Inderbinen M, Morach M, Berner AZ, Hächler H, Stephan R. 2015. Extended-spectrum-β-lactamase-producing Enterobacteriaceae isolated from vegetables imported from the Dominican Republic, India, Thailand, and Vietnam. Appl Environ Microbiol 81:3115–3120.

41. Bhutani N, Muraleedharan C, Talreja D, Rana SW, Walia S, Kumar A, Walia SK. 2015. Occurrence of multidrug resistant extended spectrum beta-lactamase-producing bacteria on iceberg lettuce retailed for human consumption. Biomed Res Int 2015.

42. Zekar FM, Granier SA, Touati A, Millemann Y. 2020. Occurrence of Third-Generation Cephalosporins-Resistant Klebsiella pneumoniae in Fresh Fruits and Vegetables

732   Purchased at Markets in Algeria. Microb Drug Resist 26:353–359.

733   43.   Mendes AC, Novais Â, Campos J, Rodrigues C, Santos C, Antunes P, Ramos H, Peixe L.

734   2018. mcr-1 in Carbapenemase-Producing *Klebsiella pneumoniae* in Hospitalized

735   Patients, Portugal, 2016–2017 24:2016–2017.

736   44.   Safavi M, Bostanshirin N, Hajikhani B, Yaslianifard S, van Belkum A, Goudarzi M,

737   Hashemi A, Darban-Sarokhalil D, Dadashi M. 2020. Global genotype distribution of

738   human clinical isolates of New Delhi metallo-β-lactamase-producing *Klebsiella*

739   *pneumoniae*: A systematic review. J Glob Antimicrob Resist 23:420–429.

740   45.   Leangapichart T, Lunha K, Jiwakanon J, Angkititrakul S, Järhult JD, Magnusson U, Sunde

741   M. 2021. Characterization of *Klebsiella pneumoniae* complex isolates from pigs and

742   humans in farms in Thailand: population genomic structure, antibiotic resistance and

743   virulence genes. J Antimicrob Chemother 76:2012–2016.

744   46.   Rodrigues C, Passet V, Rakotondrasoa A, Brisse S. 2018. Identification of *Klebsiella*

745   *pneumoniae*, *Klebsiella quasipneumoniae, Klebsiella variicola* and Related Phylogroups

746   by MALDI-TOF Mass Spectrometry. Front Microbiol 0:3000.

747   47.   Long SW, Linson SE, Ojeda Saavedra M, Cantu C, Davis JJ, Brettin T, Olsen RJ. 2017.

748   Whole-Genome Sequencing of Human Clinical *Klebsiella pneumoniae* Isolates Reveals

749   Misidentification and Misunderstandings of *Klebsiella pneumoniae* , *Klebsiella variicola*

750   , and *Klebsiella quasipneumoniae*. mSphere 2:e00290-17.

751   48.   Clinical and Laboratory Standards Institute (CLSI). 2017. Performance Standards for

752   Antimicrobial Susceptibility Testing *In* Wayne, P (ed.), Performance Standards for

753   Antimicrobial Susceptibility Testing. 27th ed. CLSI supplement M100.

49.   Magiorakos A-P, Srinivasan A, Carey RB, Carmeli Y, Falagas ME, Giske CG, Harbarth S,

Hindler JF, Kahlmeter G, Olsson-Liljequist B, Paterson DL, Rice LB, Stelling J, Struelens

MJ, Vatopoulos A, Weber JT, Monnet DL. 2012. Multidrug-resistant, extensively drug-

resistant and pandrug-resistant bacteria: an international expert proposal for interim

standard definitions for acquired resistance. Clin Microbiol Infect 18:268–81.

50.   Prjibelski A, Antipov D, Meleshko D, Lapidus A, Korobeynikov A. 2020. Using SPAdes De

Novo Assembler. Curr Protoc Bioinforma 70:e102.

51.   Zerbino DR, Birney E. 2008. Velvet: Algorithms for de novo short read assembly using

de Bruijn graphs. Genome Res 18:821.

52.   Seemann T. 2014. Prokka: Rapid prokaryotic genome annotation. Bioinformatics

30:2068–2069.

53.   Diancourt L, Passet V, Verhoef J, Grimont PAD, Brisse S. 2005. Multilocus Sequence

Typing of *Klebsiella pneumoniae* Nosocomial Isolates. J Clin Microbiol 43:4178–4182.

54.   Lam MMC, Wick RR, Watts SC, Cerdeira LT, Wyres KL, Holt KE. 2021. A genomic

surveillance framework and genotyping tool for *Klebsiella pneumoniae* and its related

species complex. Nat Commun 12:4188.

55.   Carattoli A, Zankari E, García-Fernández A, Voldby Larsen M, Lund O, Villa L, Møller

Aarestrup F, Hasman H. 2014. In silico detection and typing of plasmids using

PlasmidFinder and plasmid multilocus sequence typing. Antimicrob Agents Chemother

58:3895–903.

56.   Page AJ, Cummins CA, Hunt M, Wong VK, Reuter S, Holden MTG, Fookes M, Falush D,

Keane JA, Parkhill J. 2015. Roary: Rapid large-scale prokaryote pan genome analysis.

Bioinformatics 31:3691–3693.

57.   Croucher NJ, Page AJ, Connor TR, Delaney AJ, Keane JA, Bentley SD, Parkhill J, Harris

SR. 2015. Rapid phylogenetic analysis of large samples of recombinant bacterial whole

genome sequences using Gubbins. Nucleic Acids Res 43:e15.

Figure 1

Receive/Collect Sample

Dilute 25 g of sample in 225 ml Buffered Peptone Water (BPW) broth and mix

Incubate BPW suspension at 37°C for 24 h

Following incubation, using a 10 µl loop, plate onto SCAI media and incubate at 44 °C ± 1°C for 48 h ± 1 h

Remove plates from incubator following 48 h ± 1 h incubation. Record findings.

Purify and identify all suspected *Klebsiella* spp. colonies
(by MALDI-TOF or species-specific PCR)
Freeze strains confirmed as *Klebsiella* spp. at -80°C
(If several morphotypes are available, store one colony per morphotype)

Figure 2

Figure 3