1 **Young transposable elements rewired gene regulatory networks in human and chimpanzee**
2 **hippocampal intermediate progenitors**
3

4 Sruti Patoori[1], Samantha Barnada[1], and Marco Trizzino[1,*]
5
6 [1] Department of Biochemistry and Molecular Biology, Sidney Kimmel Medical College, Thomas Jefferson
7 University, Philadelphia, PA
8
9 * Corresponding author: Marco Trizzino, Department of Biochemistry and Molecular Biology, Sidney
10 Kimmel Medical College, Thomas Jefferson University, 233 S 10th Street, BLSB 826, Philadelphia, PA,
11 19104. E-mail: marco.trizzino@jefferson.edu

12
13
14
Abstract

The hippocampus is associated with essential brain functions such as learning and memory. Human
hippocampal volume is significantly greater than expected when compared to non-human apes,
suggesting a recent expansion. Intermediate progenitors, which are able to undergo multiple rounds of
proliferative division before a final neurogenic division, may have played a role in the evolutionary
hippocampal expansion. To investigate the evolution of gene regulatory networks underpinning
hippocampal neurogenesis in apes, we leveraged the differentiation of human and chimpanzee induced
Pluripotent Stem Cells into TBR2-positive hippocampal intermediate progenitors (hpIPCs). We find that
the gene networks active in hpIPCs are significantly different between humans and chimpanzees, with
~2,500 genes differentially expressed. We demonstrate that species-specific transposon-derived
enhancers contribute to these transcriptomic differences. Young transposons, predominantly
Endogenous Retroviruses (ERVs) and SINE-Vntr-Alus (SVAs), were co-opted as enhancers in a species-
specific manner. Human-specific SVAs provided substrates for thousands of novel TBR2 binding sites, and
CRISPR-mediated repression of these SVAs attenuates the expression of ~25% of the genes that are
upregulated in human intermediate progenitors relative to the same cell population in the chimpanzee.

30
31
32 **Summary statement:** Evolution of human and chimpanzee hippocampal development was mediated by
33 co-option of young retrotransposons into species-specific enhancers.
34
35 **Key words:** hippocampal development, transposable elements, iPSCs, intermediate progenitors, TBR2,
36 SVAs

Introduction

The hippocampus is associated with many traits relevant in the context of human evolution. These include traits such as tool use and language which require social cognition and learning, as well as spatial memory, navigation, and episodic memory (Burgess et al., 2002; Eichenbaum, 2017a; Eichenbaum, 2017b; Squire, 1992; Tomasello and Herrmann, 2010). This region of the brain is also greatly affected by Alzheimer's Disease (AD), a neurodegenerative disorder characterized by cell death, plaques and tangles of misfolded proteins, and cognitive decline (Duyckaerts et al., 2009). It has been hypothesized that the cognitive AD phenotype is uniquely human and that non-human primates, including chimpanzees, do not exhibit AD-related dementia (Edler et al., 2017; Finch and Austad, 2015; Walker and Jucker, 2017). If humans are uniquely susceptible to AD, it is crucial to understand how the human hippocampus differs from that of our closest biological relatives, the chimpanzees.

Human hippocampal volume is 50% greater than expected when compared to the hippocampal volumes of non-human apes, possibly indicating a recent hippocampal expansion specific to the human lineage (Barger et al., 2014). However, the evolution of the human hippocampus and the developmental mechanisms driving the human-specific volume increase have not yet been thoroughly studied.

Recent studies have suggested that evolutionary changes to neuronal progenitors may have had an impact on cortical volume in primates by increasing proliferative potential (Martínez-Cerdeño et al., 2006; Rétaux et al., 2013; Florio and Huttner, 2014). A specific class of neuronal progenitors known as intermediate progenitor cells (IPCs) or "transit amplifying cells" are able to undergo multiple rounds of proliferative division before a final neurogenic division (Englund, 2005; Arnold et al., 2008; Pontious et al., 2008; Hevner, 2019). These cells express the neurodevelopmental transcription factor TBR2 (EOMES) and are found in the sub-ventricular zone of the developing neocortex and hippocampus (Bulfone et al., 1999; Kimura et al., 1999; Englund, 2005; Cipriani et al., 2016). Genetic ablation of TBR2 in these progenitors results in reduced cortical thickness in mice (Sessa et al., 2008), abnormal cortical cell differentiation (Mihalas et al., 2016), and impaired neurogenesis in the hippocampal formation (Hodge et al., 2012). As these IPCs are hypothesized to play a role in neocortical expansion, they may also be involved in the lineage-specific hippocampal expansion seen in humans.

Many of the differences between humans and chimpanzees are due to diverging gene regulatory sequences (Agoglia et al., 2021; Enard et al., 2002; Gokhman et al., 2021; King and Wilson, 1975; Wray, 2007). A recent study comparing gene expression between adult human, chimpanzee and macaque brain regions identified several genes specifically upregulated in the human hippocampus (Sousa et al., 2017). However, transcriptomic differences between primate species during specific timepoints of hippocampal development have not been investigated.

As samples of developing human and chimpanzee brain tissue are extremely limited, Induced Pluripotent Stem Cells (iPSCs) are an ideal system to conduct comparative studies of human and chimpanzee hippocampal development. Previous studies have employed iPSC-derived cortical organoids and iPSC-derived neuronal progenitor cells (NPCs) from human and chimpanzee for comparative and developmental genomic purposes (Marchetto et al., 2019; Mora-Bermúdez et al., 2016). Here, we leverage human and chimpanzee iPSC-derived hippocampal progenitors as models for comparative developmental and genomic studies with the goal of identifying species-specific differences in gene regulation during hippocampal development.

Several recent papers have recently demonstrated that transposable elements (TEs) can alter existing regulatory elements or generate entirely novel ones, as well as expand in a species- or lineage-specific manner (reviewed in Sundaram and Wysocka, 2020). Species-specific TE expansion and co-option into gene regulatory networks have been demonstrated as a mechanism for evolutionary change (Chuong et al., 2016; Fuentes et al., 2018; Jacques et al., 2013; Lynch et al., 2011; Lynch et al., 2015; Miao et al., 2020; Mika et al., 2021; Pontis et al., 2019; Trizzino et al., 2017). Endogenous Retroviruses (ERVs)

and SINE-Vntr-Alus (SVAs) are among the TE families more frequently associated with gene regulatory
activity in the human genome (Chuong et al., 2016; Fuentes et al., 2018; Pontis et al., 2019; Trizzino et al.,
2017; Trizzino et al., 2018). SVAs encompass six subfamilies, denoted as SVA-A through -F. Of the ~3,000
SVA copies in the human genome, nearly half are human specific, including all the SVA-E and SVA-F (Quinn
and Bubb, 2014; Wang et al., 2005). The remaining half are also found in other great apes. SVAs are still
replication competent and thus able to transpose in the human genome. ERVs are retrotransposons
belonging to the Long Terminal Repeat (LTR) group. They are remnants of past retroviral infection events,
and make up ~8% of the human genome (Tokuyama et al., 2018). Both SVAs and ERVs were recently
found to be enriched within the sequences of active cis-regulatory elements (enhancers, promoters) in
hippocampal tissue as compared to other human brain regions where they are predominantly repressed
(Trizzino et al., 2018). Therefore, we hypothesize that species-specific ERV and SVA transposon activity
may influence the gene regulatory networks necessary for human and chimpanzee hippocampal
development.
Given the key function that intermediate progenitors played in the evolution of the primate brain
(Florio and Huttner, 2014; Martínez-Cerdeño et al., 2006), we sought to identify molecular differences
between iPSC-derived human and chimpanzee hippocampal intermediate progenitors (hpIPCs) in terms
of gene expression and the regulatory activity of non-coding regions. We specifically examined gene
expression (RNA-seq), gene regulation (ATAC-seq) and functional TE activity via CRISPR-interference.
After confirming that the hpIPC differentiated cells express the appropriate neurodevelopmental
markers, we conducted a transcriptomic comparison between human and chimpanzee hpIPCs. This
analysis revealed over 2,500 differentially expressed genes. We then used ATAC-seq to conduct extensive
analyses of differential chromatin accessibility between human and chimpanzee hpIPCs. In both species,
differentially accessible chromatin regions were more likely than expected to overlap a TE insertion.
Further, these regions were found as both enriched and depleted for specific TE families. Notably, species-
specific enrichment for ERV and SVA sequences within differentially accessible genomic sites correlated
with species-specific changes in nearby gene expression. This is likely driven by transcription factors
binding to the TE-derived regulatory sequences, as we demonstrate for TBR2 and SVA-derived enhancers.
Finally, we used CRISPR-interference to repress all the accessible SVAs in progenitor-like cells and
demonstrated that such repression results in global changes in gene expression and affects hundreds of
important neurodevelopmental genes.
This work demonstrates that two young TE families have contributed significantly to the gene
regulatory differences between human and chimpanzee hippocampal development, providing insight
into how the human hippocampus evolved both its unique cognitive capacity and its susceptibility to
neurodegenerative disease.
131
132
133 **Results**
134
An iPSC-derived model for human and chimpanzee hippocampal intermediate progenitors
We modeled hpIPCs in humans and chimpanzees using three human and three chimpanzee iPSC
lines. All six iPSC lines were validated as pluripotent in previous studies (Gallego Romero et al., 2015;
Pagliaroli et al., 2021; Pashos et al., 2017; Ward et al., 2018; Yang et al., 2015; Zhang et al., 2015). For both
human and chimpanzee iPSCs, we used two female cell lines and one male line.
We used a previously published method to generate hpIPCs from iPSCs (Yu et al., 2014). In this
protocol, the stem cells are treated with a media containing anticaudalizing factors and a Sonic Hedgehog
antagonist (DKK1, Nogging, SB431542) to generate forebrain progenitor cell types (Yu et al., 2014). It
should be noted that the hpIPCs are distinct from a more general neuronal progenitor cell (pan-NPC) as
the pan-NPC media is supplemented only with FGF2 and B27. Moreover, and the hpIPCs can be further

induced to generate mature hippocampal CA3 pyramidal or dentate gyrus granule neurons (Sarkar et al.,
2018; Yu et al., 2014).
As we were specifically interested in TBR2-positive intermediate hippocampal progenitors, we
first differentiated one of the human lines for five days and performed a TBR2 time-course western blot
on cells collected at day 0, day 5 and day 11. The blot revealed day-5 as a time-point in which TBR2 protein
is highly expressed (Fig. 1A).
To confirm that day 5 was the optimal time-point in both species, we differentiated one human
and one chimpanzee line for five days and used immunocytochemistry to confirm the presence of OTX2,
which labels neuronal progenitors, as well as TBR2 and PAX6 which specifically mark the intermediate
progenitor cell type (Florio and Huttner, 2014; Hevner, 2019; Fig. 1B). A previous study of human fetal
hippocampal development demonstrated that 47% and 57% of TBR2-positive cells express PAX6 at
gestational weeks 11 and 13, respectively (Cipriani et al., 2016). We found that nearly all of the
cells treated with hpIPC media for five days were positive for all three neurodevelopmental markers in
both species (Fig. 1B). Together, these data indicate that our iPSC-derived model is suitable to study TBR2-
positive hippocampal intermediate progenitors in both human and chimpanzee.
160
Important neurodevelopmental genes are differentially expressed between human and chimpanzee
hippocampal intermediate progenitors
To investigate the differences between human and chimpanzee hippocampal intermediate
progenitors, we first aimed to characterize differential gene expression between the hpIPCs of the two
species. After five days of treatment with the hpIPC differentiation media, we collected cells for RNA
extraction (Fig. 2A). To ensure statistical power, we conducted bulk RNA-seq on two replicates each of all
six cell lines (i.e. three biological replicates and six technical replicates per species). The differentiation,
the harvesting, and the RNA processing were performed in mixed batches with samples from both species
to prevent batch effects. The libraries were sequenced using Illumina NextSeq500, generating 100 bp
Paired-End reads. Non-orthologous genes were omitted from the analysis and a total of 2,588 genes were
identified as differentially expressed (FDR <0.05 and log2foldChange > 1.5 or < -1.5). The genes with
log2foldChange > 1.5 were more highly expressed in the human hpIPCs ("Human UP") while the genes
with log2foldChange < -1.5 were more highly expressed in the chimpanzee hpIPCs ("Chimp UP"). In total,
1,686 (65.1%) of the differentially expressed genes (DEG) were "Human UP" and 901 (34.9%) of the DEG
were "Chimp UP" genes (Fig. 2B, Supplementary Table 1.1).
Hippocampal neurodevelopmental markers *PAX6*, *OTX2*, *NEUROD1* and *FOXG1* were found as
highly expressed in both species. This is consistent with the immunostaining in demonstrating that
the hpIPCs differentiation was successful and that the progenitors from both species are comparable.
The "Human UP" genes include *FOXP2*, *MTRNR2L8*, *DHX40*, *VPS13B*, *WDFY2*, and *PURB*, all of
which are associated with neurodevelopment or neurodegenerative disease (Abrajano et al., 2009;
Hickey et al., 2019; Kamboh et al., 2019; Kolehmainen et al., 2003; MacDermot et al., 2005; Mathys et al.,
2019; Sin et al., 2015; Taher et al., 2014). The "Chimp UP" genes include *HIST1H3A*, *BLC2L2*, and
CLIC1 which have been reported as expressed in the hippocampus and associated with sleep, AD, and
neurite outgrowth (Averaimo et al., 2014; Datson et al., 2009; Wei, 2020).
To understand the transcriptional programs driving these differences in gene expression,
we conducted an Ingenuity Pathway Analysis. Three of the top five upstream regulators predicted by the
pathway analysis were the transcription factors CREB1, FOXA2 and TBR2 (EOMES) (Fig. 2C). CREB1 is
known to regulate genes involved in the nervous system and neurodevelopment (reviewed in Sakamoto
et al., 2011), FOXA2 controls dopaminergic neuronal development and disease (Kittappa et al., 2007), and
TBR2 plays a crucial role in cortical and hippocampal neurogenesis and is the signature marker of the
intermediate progenitor population (Cipriani et al., 2016; Englund, 2005). This pathway analysis was
consistent with the RNA-seq data as predicted targets of all three transcription factors were also found to

be among the 2,588 genes differentially expressed between human and chimpanzee hpIPCs (Fig. 2D,
Supplementary Table 1.2-1.4). The pathway analysis also determined that several of the differentially
expressed genes are involved in embryonic development (Fig. 2D, Supplementary Table 1.5). Overall,
these findings indicate that previously characterized neurodevelopmental gene regulatory networks are
utilized differently during human and chimpanzee hippocampal development.
198
Human-specific chromatin accessibility patterns in hippocampal intermediate progenitors
After identifying differentially expressed genes and the transcriptional networks that may be
involved, we sought to identify cis-regulatory differences between the human and chimpanzee hpIPCs. To
this end, we conducted ATAC-seq on the hpIPCs from both species. We used the same batches of
differentiated iPSCs for the ATAC-seq as we did for the RNA-seq (i.e. from the same batch of
differentiation), and generated 100 bp long Paired-End Illumina reads.
We first performed a "human-centric" analysis. We aligned the ATAC-seq reads from all six cell
lines to the respective reference genome assemblies (hg19 for the human cell lines, panTro5 for the
chimpanzee cell lines) and only retained uniquely mapped reads with high mapping quality (Samtools Q =
10 filtering). Next, we identified regions of accessible chromatin (peaks; FDR < 0.05) in all three human
cell lines. Only peaks replicated in all the three human lines were retained. To carry out a proper
comparison, we only retained replicated human ATAC-seq peaks with orthologs in the chimpanzee
genome (see methods; Fig. 3A). This filtering ultimately resulted in 82,235 human ATAC-seq peaks
replicated in all the human cell lines and with orthologs in the chimpanzee genome. These 82,235 regions
were used for downstream analysis. We found that the chromatin accessibility at these regions was highly
reproducible across all three human cell lines (Fig. 3B).
Next, we quantified the ATAC-seq read depth for each of the 82,235 regions and used DESeq2 to
identify sites exhibiting differential chromatin accessibility between the two species. In total, we identified
3,006 differentially accessible regions (FDR <0.05; log2foldChange > 1.5 or < -1.5; Supplementary Table
2.1-2.2). Of these regions, 92.3% were located at least 1 kb away from the closest transcription start site
(TSS), suggesting they could be putative enhancers, while the remaining were putative promoters (Fig.
3C). As expected, given that this analysis was performed with a human-centric approach, 90.1% of the
differentially accessible (DA) peaks were significantly more accessible in the human hpIPCs relative to
chimpanzee hpIPCs ("Human UP"; Fig. 3C).
As TE insertions can be a source of cis-regulatory evolution, we examined whether the DA regions
were more likely to overlap with a TE than those which were accessible to the same degree in both
species (non-DA, Fig. 3D and Supplementary Table 2.4-2.5). We observed that 1,335/3,006 (44.4%) DA
regions overlapped a TE (Supplementary Table 2.3). This is significantly higher than what was observed
for the non-DA peaks (33.2% overlapped a TE; Two-sided Fisher's Exact Test, p < 0.0001; Fig. 3D). This
indicates that chromatin regions with human-specific accessibility are significantly more likely to be TE-
derived than the regions with accessibility levels conserved between human and chimpanzee.
Next, we associated the nearest gene to each DA region and found that TE-derived DA regions
were significantly more likely to be near a differentially expressed gene relative to non-TE derived DA
regions (Two-sided Fisher's Exact Test, p = 0.016; Fig. 3E). Finally, we investigated whether specific TE
families were overrepresented among the TE-derived DA regions and found enrichment for LTRs. While
LTRs account for ~16% of all human annotated TEs, they represented 33.9% of the TEs overlapping DA
regions in our human-centric analysis (Two-Sided Fisher's Exact Test p < 0.0001; Fig. 3F and Supplementary
Table 2.6). Of these enriched LTRs, 97.1% were ERVs. Notably, 31.7% of those LTR-derived DA regions
were located near a differentially expressed gene (Two-Sided Fisher's Exact Test p < 0.0001).
Together, these data indicate that there are TE-derived cis-regulatory elements that have
significantly greater accessibility in humans than chimpanzees during hippocampal neurogenesis. These
TE insertions preceded the human-chimpanzee split, but the difference in accessibility is species-specific,

suggesting that the co-option into gene regulatory networks took place after the human-chimpanzee
divergence.
243
Chimpanzee-specific chromatin accessibility patterns in hippocampal intermediate progenitors
We repeated the ATAC-seq analysis as described above, but this time with a "chimpanzee-centric"
approach. We started from a set of 72,211 peaks found as replicated in all the three chimpanzee lines
and with orthologs in both species (Figs. 4A,B). With this approach, we identified 3,806 ATAC-seq peaks
as differentially accessible (DA) between human and chimpanzee (FDR <0.05; log2foldChange > 1.5 or < -
1.5; Supplementary Table 3.1-3.2), 82% of which displayed greater accessibility in chimpanzee as
compared to humans (i.e. "Chimp UP"; Fig. 4C). Similar to what we observed with the human-centric
analysis, 97.9% of the 3,806 DA regions were putative enhancers (distance > 1Kb from TSS; Fig. 4C).
As seen in the human-centric analysis, chimpanzee-specific DA peaks were more likely to be TE-
derived than those similarly accessible across species (Two-sided Fisher's Exact Test, p < 0.0001; Fig. 4D
and Supplementary Tables 3.3-3.5). Of the chimpanzee DA regions, 37.1% overlapped an annotated
chimpanzee TE, compared to only 28.1% of the non-DA regions (Two-sided Fisher's Exact Test, p < 0.0001;
Fig. 4D). However, the TE-derived enhancers in this chimpanzee-centric analysis were no more likely than
the non-TE derived ones to be located near differentially expressed genes (Fig. 4E).
We found enrichment for LTRs, which account for approximately 16% of the chimpanzee TEs but
represented 36.1% of the TE-derived DA regions (Two-sided Fisher's Exact Test, p-value < 0.0001; Fig. 4F
and Supplementary Table 3.6; 98.2% were ERVs), and SVAs, which account for just 0.25% of annotated
chimpanzee TEs but represented 1.1% of the TE-derived DA regions (Two-sided Fisher's Exact Test, p <
0.0001, Fig 4G and Supplementary Table 3.8). In particular, the SVA-B and C subfamilies were the most
enriched (Fig. 4H). Together, these data indicate that chimpanzee ERV and SVA transposons were co-
opted into regulatory elements important for the developing chimpanzee hippocampal intermediate
progenitors.
266
Genomic features underlying species-specific LTR enrichment at hpIPC enhancers
We aimed to further investigate genomic features potentially underlying the LTR enrichment
among the differentially accessible hippocampal progenitor regions. To this end, we conducted a motif
analysis using the MEME suite (Bailey et al., 2015). Binding motifs for CTCF and EGR2 were detected as
enriched in the LTR-derived differentially accessible regions identified from both the human-centric and
chimpanzee-centric analyses (Figs. 5A, 5C). EGR2 is an early response gene involved in learning and
memory, in the brain's response to stimuli, and in hippocampal synaptic plasticity (Cheval et al., 2012;
Mukherjee et al., 2021; Poirier et al., 2007). CTCF, a well-known regulator of chromatin structure, has
been implicated in various neurodevelopmental disorders (reviewed in Davis et al., 2018)
We identified 63 differentially expressed genes located near the human-enriched LTRs (Fig. 5B
and Supplementary Table 2.7). These included *GLUL*, a glutamine synthetase hypothesized to provide
neuroprotection in AD patients, (Kohane and Wood, 2021) and *DLK1*, a Notch ligand involved in SVZ
neurogenesis (Ferrón et al., 2011), both of which are upregulated in human hpIPCs compared to
chimpanzee hpIPCs.
We identified 34 differentially expressed genes located near the chimp-enriched LTRs (Fig. 5D and
Supplementary Table 3.7). For the chimp-centric analysis the differentially expressed genes located near
TE-derived enhancers with species-specific accessibility included *HIST1H3A* and *MTRNR2L8* (Fig. 5D).
These genes were highly upregulated in the chimpanzee and human hpIPCs, respectively (Fig. 5D).
Importantly, *HIST1H3A* has been associated with autism spectrum disorders and sleep deprivation
(Crawley et al., 2016; Wei, 2020) while *MTRN2L8* has been reported as upregulated in Alzheimer's patients
(Mathys et al., 2019). *BDNF*, which is involved in hippocampal neurogenesis and plays an important role

in the SVZ (Bath et al., 2012), was highly upregulated in the chimpanzee and also located near an LTR-
derived enhancer with chimpanzee-specific accessibility.
290
Human-Specific SVAs play a major role in hippocampal neurogenesis
As mentioned earlier, the chimpanzee-centric analysis also identified SVA transposons as enriched
within chimpanzee-specific enhancers (Fig. 6A and Supplementary Table 3.8). These SVAs were enriched
for the identified binding motifs of the neurodevelopmental factors ASCL1, ZIC1, and KLF8 (Andersen et
al., 2014; Aruga, 2004; Yi et al., 2014) as well as the JUN/FOS AP-1 dimer, which is a known enhancer
activator (Raivich, 2008; Raivich and Behrens, 2006) (Fig 6A).
It is important to remark that all the analyses shown so far were exclusively based on genomic
sites with characterized orthologs in both species, to ensure an "apple-to-apple" comparison. However,
nearly 2,000 SVA copies, including the entire SVA-E and F subfamilies, are exclusive to the human genome.
Given that previous studies found that SVAs are highly enriched in active enhancers and promoters of the
human hippocampus (Trizzino et al., 2018), we sought to investigate this further. We focused on the SVA
copies exclusively present in the human genome and not in any other primate genome (hereafter human-
specific SVAs). We performed sequence-based motif analysis for all the human-specific SVAs and
identified the binding motif for the hpIPCs signature factor TBR2 (EOMES) as the most enriched ($p = 10^{-2103}$; Fig. 6B). Motifs for other transcription factors associated to hippocampal neurogenesis and function
305
were also recovered (SMAD4, VDR, PLAG1; Fig. 6B).
Next, we annotated all of the genes found within 50 kb from each human-specific SVA. A total of
2,216 genes were recovered using this approach. We performed Pathway Analysis on this set of genes
and found that Melatonin Degradation and Nicotine Degradation were the two most significantly enriched
pathways ($p = 7.6 \times 10^{-6}$ and $p = 2.5 \times 10^{-5}$ respectively; Fig. 6C) and PHF8 was recovered as the top
upstream regulator for the gene network (Fig. 6C). Notably, both Melatonin and Nicotine Degradation
pathways are strongly active in the human hippocampus. PHF8 is a histone demethylase that contributes
to the regulation of mTOR. The mTOR pathway is hyperactive in the human hippocampus where it
regulates the protein synthesis-dependent plastic changes underlying learning and memory (Bekinschtein
et al., 2007; Fortress et al., 2013; Graber et al., 2013). Mutations in the *PHF8* gene cause cognitive
impairment and intellectual disability (Chen et al., 2018).
Therefore, we sought to determine the contribution of human-specific SVAs to the TBR2-
mediated gene regulatory network in human hpIPCs. Using our ATAC-seq data, we identified 1,816 human
SVAs as accessible in human hpIPCs (Fig. 6D). Of these, nearly a quarter (434) displayed high accessibility,
while 1,382 were moderately accessible (Fig. 6D). Next, we used chromatin immunoprecipitation
followed by sequencing (ChIP-seq) to profile TBR2 binding in two human lines at day 5 of hpIPC
differentiation. As with the previous sequencing experiments, we generated 100 bp long Paired-End reads
and only retained uniquely mapping high quality reads (Samtools q=10 filtering) in order to maximize our
chance to properly map reads on repetitive regions. This experiment revealed that 739 of the accessible
SVAs showed TBR2 signal in the two human lines (Fig. 6D and Supplementary Table 2.8). Notably, 257 of
the (48.3%) TBR2-bound SVAs were human-specific. (Fig. 6E and Supplementary Table 2.9). TBR2-bound
human-specific SVAs were located near 37 genes that our RNA-seq analysis identified as differentially
expressed between human and chimpanzee (Fig. 6F and Supplementary Table 2.10). These genes include
VPS13B (up in humans), which is responsible for a rare developmental disease known as Cohen Syndrome
(Kolehmainen et al., 2003), *NR4A2* (down in humans), which has been implicated in neurodevelopmental
language impairment (Reuter et al., 2017), *DHX40* (down in humans), which is implicated in Alzheimer's
(Taher et al., 2014), and *E2F1* (up in humans), which is a cell cycle regulator associated with several
neurodegenerative diseases such as Alzheimer's (Zhang et al., 2010).
In summary, these data support a model in which human-specific SVAs provided a substrate for
binding sites of TBR2 and other important hippocampal regulators. Therefore, it is likely they were co-

opted in the gene regulatory networks active during hippocampal neurogenesis, which led to human-
specific regulation of key genes.
338
CRISPR-mediated SVA repression has massive repercussions on global gene expression
To further assess the contribution of SVA transposons to the gene regulation of hippocampal
intermediate progenitors, we leveraged CRISPR-interference to simultaneously repress most of the active
SVAs. We utilized NCCIT cells treated with retinoic acid (RA) as the experimental system for this purpose.
The NCCIT cell line is derived from embryonal carcinoma and thus exhibits a gene expression signature
highly similar to human embryonic stem cells (Fuentes et al., 2018). Importantly, NCCITs treated for 7 days
with RA differentiate into intermediate neural progenitor-like cells expressing both PAX6 and TBR2
(Mandal et al., 2015; Fig 7A). We cloned a stable NCCIT line with a permanently incorporated doxycycline-
inducible, catalytically-dead Cas9 fused to a repressive KRAB domain (dCas9-KRAB). The KRAB domain
deposits repressive histone methylation (H3K9me3) to the regions targeted by the dCas9 via single guide-
RNAs (sgRNAs). Into this same line, we also permanently knocked-in two single guide-RNAs (sgRNAs) that
are able to target >80% of all SVAs (Pontis et al. 2019). Hereafter, we refer to the RA-treated CRISPR line
as RA-NCCITs. Remarkably, exposing the RA-NCCITs to doxycycline for 72 hours was sufficient to induce
dCas9 activation (Fig. 7B) and the deposition of repressive histone methylation (H3K9me3) on over 2,500
previously unmethylated SVAs (Fig. 7C). We next performed RNA-seq on the RA-NCCITs with or without
doxycycline treatment (three replicates per condition). First, we observed that the genome-wide
expression levels (TPMs) of the RA-NCCITs where highly correlated with those of the iPSC-derived human
hpIPCs (Pearson correlation = 0.93; $p < 2.2 \times 10^{-16}$). This indicates that RA-treated NCCITs are appropriate
to model hippocampal intermediate progenitors.
Then, we compared the expression levels of the RA-NCCITs with or without doxycycline treatment
and identified 5,795 differentially expressed genes (FDR < 0.05; Fig. 7D and Supplementary Table 4.1). Of
these genes, 677 were previously identified as differentially expressed when comparing human to
chimpanzee hpIPCs (Fig. 7E; Supplementary Table 4.2). In other words, the expression of over a quarter
(26.1%) of the genes that exhibited human-specific expression signature in hippocampal intermediate
progenitors seem to be under control of SVA transposons. Importantly, one of these genes was *FOXP2*,
which has been associated with the evolution of language and is implicated in several speech-disorders
(Enard, 2011; Liégeois et al., 2016; MacDermot et al., 2005).
Since the gRNAs used for this experiment were originally designed to target a DNA sequence
shared by the SVAs with the LTR5H family (Pontis et al. 2019), we restricted the analysis to the genes that
are associated to human SVAs (i.e. only considering the genes that represent the closest gene to an
annotated SVA: hereafter SVA-genes). By doing so, we found 611 SVA-genes as differentially expressed in
RA-NCCITs upon SVA repression (Supplementary Table 4.3). Of these, 90 were previously identified as
differentially expressed when comparing human to chimpanzee hpIPCs (Supplementary Table 4.4). Thus,
the expression of these 90 genes can be bona-fide considered as directly regulated by SVA-derived
enhancers in both primary hippocampal progenitors and in the NCCIT cell line. Remarkably, the large
majority of these genes (72.5%) had decreased expression upon SVA repression. These include *SOX2*,
FGF2, *PRDM1*, *NTRK2*, and *TFAP2B* (Supplementary Table 4.4).
Finally, we examined the genes previously identified as being near a TBR2-bound human-specific
SVA in iPSC-derived hpIPCs, and found that nearly a third of them lose expression in RA-NCCITs upon SVA
repression (Fig. 7F). In summary, our functional experiments indicate a widespread role for human-
specific SVA transposons as cis-regulatory elements during hippocampal neurogenesis.
380
Discussion
The hippocampus is susceptible to specific neurodegenerative disorders such as Alzheimer's
Disease but may have also played an important role in the evolution of human cognition. Spatial memory,

which is attributed to the hippocampus, may have contributed to the geographic expansion of ancient
humans. Characterizing the human-specific gene regulatory networks of hippocampal development
provides insight into its role in human evolution. Though there is no consensus on whether the cognitive
phenotypes seen in AD are uniquely human, understanding the unique properties of the human
hippocampus may lead the way to potential treatments. Thus, the work described here is relevant both
in terms of evolutionary developmental biology and evolutionary medicine.

To study the evolution of the human hippocampus from a developmental standpoint, we
investigated the extent to which TEs contributed to gene expression profiles of human and chimpanzee
hippocampal intermediate progenitors. TEs account for nearly ~50% of the human genome composition
and many elegant studies have established that at least a fraction of the TEs can regulate the host genes
in humans and other primates (Chuong et al., 2013; Chuong et al., 2016; Cosby et al., 2021; del Rosario et
al., 2014; Du et al., 2016; Fuentes et al., 2018; Jacques et al., 2013; Judd et al., 2021; Lynch et al., 2011;
Lynch et al., 2015; Mika et al., 2021; Modzelewski et al., 2021; Okhovat et al., 2020; Rayan et al., 2016;
Schmidt et al., 2012; Sundaram et al., 2014; Trizzino et al., 2017; Trizzino et al., 2018; Ward et al., 2018;
Xie et al., 2013).

The hpIPCs are a transient progenitor population during a critical developmental stage in the Sub-
ventricular zone (SVZ) of both the hippocampus and neocortex (Bulfone et al., 1999; Cipriani et al., 2016;
Englund, 2005; Kimura et al., 1999), and there is consensus that this progenitor population may have
played a role in the evolution of brain volume in mammals (Florio and Huttner, 2014; Martínez-Cerdeño
et al., 2006). To study the developmental evolution of human hpIPCs, we leveraged a comparative
approach centered on differentiating human and chimpanzee iPSCs into a neuronal population which
closely recapitulates hpIPCs, as demonstrated by the high expression of signature markers such as TBR2
and PAX6.

The transcriptomes of human and chimpanzee hpIPCs have not been previously compared, largely
due to the limited availability of primary tissue. Here, we carried out this comparison using our iPSC-
derived system and identified profound differences between the two species, with over 2,500 genes
differentially expressed at this stage. These genes include several which were previously associated with
cognitive function, language, neurodevelopment and neurodegeneration, many of which are upregulated
in humans relative to the chimpanzee (e.g. *MTRNR2L8*, *DHX40*, *VPS13B*, *WDFY2*, *PURB*).

We demonstrate that species-specific enhancers significantly contributed to the gene expression
differences we identified. This is consistent with recent studies that used primary brain tissues from
several species to profile species-specific cis-regulatory activity in mammals (Emera et al., 2016; Reilly et
al., 2015). Importantly, we find that these species-specific enhancers are enriched for young transposable
elements. Several studies have identified evolutionarily young L1 LINEs as active in the brain during
different developmental stages, suggesting that they could serve as alternative promoters for many genes
involved in neuronal functions (Coufal et al., 2009; Jönsson et al., 2019; Sur et al., 2017; Thomas et al.,
2012; Zhao et al., 2019). Here, we identified other young transposable elements, ERVs and SVAs, as
regulators of intermediate progenitor gene expression. The identification of ERVs as candidate enhancers
in human and chimpanzee intermediate progenitors is consistent with previous studies that
demonstrated that ERVs heavily impact the gene regulatory programs during immune response (Chuong
et al., 2013; Chuong et al., 2016), in pluripotency maintenance and development (Coluccio et al., 2018;
Fuentes et al., 2018; Miao et al., 2020), in the mammalian placenta (Lynch et al., 2011; Lynch et al., 2015;
Mika et al., 2021), in the primate liver (Trizzino et al., 2017), and in many cancer types (Ito et al.; Ivancevic
and Chuong, 2020; Shah et al., 2021). Similar to the L1s, the ERVs also possess a well-defined cis-regulatory
architecture (e.g. they have their own promoter), and this may have had a role in the co-option of these
TEs as functioning cis-regulatory elements.

The SVAs are particularly interesting from a human evolution standpoint, given that half of the
known copies are exclusively present in our species. Moreover, SVAs are among the few transposable

elements that still exhibit active transposition in the human genome. We and others have previously revealed that SVAs can be source of enhancers in primates (Playfoot et al., 2021; Pontis et al., 2019; Trizzino et al., 2017; Trizzino et al., 2018). The repression of some SVAs by specific zinc-finger proteins at specific stages of neuronal development is also a crucial mechanism for successful neurogenesis (Playfoot et al., 2021; Pontis et al., 2019; Turelli et al., 2020). Here, we demonstrate that SVAs are pervasive regulators of hippocampal neurogenesis and they act as enhancers in the hippocampal intermediate progenitor population. By using CRISPR-interference (CRISPR-i), we show that repressing hundreds of normally "de-repressed" SVAs alters the expression of thousands of genes. Intriguingly, our CRISPR-i experiments revealed that global SVA repression leads to the attenuation of the expression levels of over a quarter of the ~2,500 genes previously identified as with human-specific gene expression signal in hippocampal intermediate progenitors. These include critical neurodevelopmental regulators such as *FOXP2*, *HAND2*, *MEF2C*, *SOX2*, and *SOX4*. In normal conditions, these genes are more highly expressed in humans relative to the chimpanzee, but upon SVA repression these differences were diminished.

In conclusion, our findings indicate that the development of hippocampal neurons has been profoundly affected by the domestication of young transposable elements. These young TEs have been co-opted as functional enhancers and promoters and ultimately rewired the expression of hundreds of critically important neuro-regulators in the developing human hippocampus. The human-specific gene expression and the associated TE-derived enhancers we have identified here may play important roles both in human evolution and neurodegenerative disease.

Materials and Methods

Antibodies

Antibody	Catalog number	Use	Dilution
TBR2	Abcam ab23345	Immunofluorescence	1:250
TBR2	Abcam ab216870	ChIP	15 ug per ChIP
TBR2	Santa Cruz sc - 293481	Western Blot	1:500
PAX6	DHSB AB_528427	Immunofluorescence	1:20
OTX2	R&D Systems AF1979	Immunofluorescence	1:500
H3K9me3	Abcam ab8898	ChIP	3 ug per ChIP
GAPDH	CST #5174	Western Blot	1:1000

Human and Chimpanzee iPSC cultures
The human male iPSC line denoted as SV20 was obtained from the University of Pennsylvania, where it was generated, and validated by the expression of pluripotency markers and differentiation into various cell types in multiple studies (Pagliaroli et al., 2021; Pashos et al., 2017; Yang et al., 2015; Zhang et al., 2015). The human female iPSC line GM 23716 was obtained from the Coriell Institute for Medical Research (Camden, NJ) and validated by the expression of pluripotency markers and differentiation into cranial neural crest cells in a previous study (Pagliaroli et al., 2021). The human female iPSC line 21792 and all three of the chimpanzee iPSC lines were obtained from the laboratory of Yoav Gilad at the University of Chicago and validated in previous studies (Gallego Romero et al., 2015; Ward et al., 2018)
The iPSC lines were expanded in feeder-free, serum-free mTeSR™1 medium (STEMCELL Technologies). Cells were passaged ~1:10 at 80% confluency using ReLeSR (STEMCELL Technologies) and small cell clusters (50–200 cells) were subsequently plated on tissue culture dishes coated overnight with Geltrex™ LDEV-Free hESC-qualified Reduced Growth Factor Basement Membrane Matrix (Fisher-Scientific).

471
hpIPC Differentiation
472
The iPSC lines were differentiated into hpIPCs as previously described (Yu et al. 2014). 3 batches comprised
of one human and one chimpanzee iPSC line each were cultured until approximately 50-70% confluence
was reached and then treated with the hpIPC media for five days prior to collection for RNA-seq, ATAC-
seq, ChIP-seq or immunofluorescence. The hpIPC media consists of DMEM/F12, 0.5X N2, 0.5X B27, DKK-
1, cyclopamine, Noggin, and SB431542 as well as the antibiotic Penn-Strep.

Western Blot
For total lysate, cells were harvested and washed three times in 1X PBS and lysed in RIPA buffer (50mM
Tris-HCl pH7.5, 150mM NaCl, 1% Igepal, 0.5% sodium deoxycholate, 0.1% SDS, 500uM DTT) with proteases
inhibitors. Twenty µg of whole cell lysate were loaded in Novex WedgeWell 4-20% Tris-Glycine Gel
(Invitrogen) and separated through gel electrophoresis (SDS-PAGE) Tris-Glycine-SDS buffer (Invitrogen).
The proteins were then transferred to ImmunBlot PVDF membranes (ThermoFisher) for antibody probing.
Membranes were incubated with 10% BSA in TBST for 30 minutes at room temperature (RT), then
incubated for variable times with the suitable antibodies diluted in 5% BSA in 1X TBST, washed with TBST
and incubated with a dilution of 1:10000 of secondary antibody for one hour at RT. The antibody was then
visualized using Super Signal West Dura Extended Duration Substrat (ThermoFisher) and imaged with
Amersham Imager 680. Used antibodies are listed in the "Antibodies" section of the methods.

Immunofluorescence
Upon fixation (4% PFA for 10 minutes), cells were first treated with 10 mM sodium citrate (pH = 6) for 10
minutes at 95 C. The cells were then permeabilized in blocking solution (0.1% Triton X-100, re PBS, 5%
normal donkey serum) for one hour at room temperature and then incubated overnight at 4 C with the
primary antibody of interest. Subsequently, the cells were treated with blocking solution for 30 minutes
at room temperature and then fluorophore-conjugated secondary antibodies for 2 hours at room
temperature. The cell nuclei were stained with 4,6-diamidine-2-phenylindole dihydrochloride (DAPI;
Sigma-Aldrich; 50 mg/ml in PBS for 15 min at RT). Immunostained cells analyzed with confocal microscopy,
using a Nikon A1R+. Images were captured with a 40X objectives and a pinhole of 1.2 Airy unit. Analyses
were performed in sequential scanning mode to rule out cross-bleeding between channels. Used
antibodies are listed in the "Antibodies" section of the methods.

Real-time quantitative polymerase chain reaction (RT-qPCR)
Cells were lysed in Tri-reagent and RNA was extracted using the Direct-zol RNA MiniPrep kit (Zymo
research). 600ng of template RNA was retrotranscribed into cDNA using RevertAid first strand cDNA
synthesis kit (Thermo Scientific) according to manufacturer directions. 15ng of cDNA were used for each
real-time quantitative PCR reaction with 0.1 µM of each primer, 10 µL of PowerUp™ SYBR™ Green Master
Mix (Applied Biosystems) in a final volume of 20 µl, using QuantStudio 3 Real-Time PCR System (Applied
Biosystem). Thermal cycling parameters were set as following: 3 minutes at 95°C, followed by 40 cycles of
10 s at 95°C, 20 s at 63°C followed by 30 s at 72°C. Each sample was run in triplicate. 18S rRNA was used
as normalizer. Primer sequences are reported in Supplementary Data 1.

ChIP-Seq and ChiP-qPCR
Samples from different conditions were processed together to prevent batch effects. 15 million cells were
cross-linked with 1% formaldehyde for 5 min at room temperature, quenched with 125mM glycine,
harvested and washed twice with 1× PBS. The pellet was resuspended in ChIP lysis buffer (150 mM NaCl,
1% Triton X-100, 0,7% SDS, 500 µM DTT, 10 mM Tris-HCl, 5 mM EDTA) and chromatin was sheared to an
average length of 200–500 bp, using a Covaris S220 Ultrasonicator. The chromatin lysate was diluted with

SDS-free ChIP lysis buffer. For ChIP-seq, 10 μg of antibody (3 μg for H3K27ac) was added to 5 μg of
sonicated chromatin along with Dynabeads Protein A magnetic beads (Invitrogen) and incubated at 4 °C
overnight. On day 2, beads were washed twice with each of the following buffers: Mixed Micelle Buffer
(150 mM NaCl, 1% Triton X-100, 0.2% SDS, 20 mM Tris-HCl, 5 mM EDTA, 65% sucrose), Buffer 500 (500
mM NaCl, 1% Triton X-100, 0.1% Na deoxycholate, 25 mM HEPES, 10 mM Tris-HCl, 1 mM EDTA),
LiCl/detergent wash (250 mM LiCl, 0.5% Na deoxycholate, 0.5% NP-40, 10 mM Tris-HCl, 1 mM EDTA) and
a final wash was performed with 1× TE. Finally, beads were resuspended in 1× TE containing 1% SDS and
incubated at 65 °C for 10 min to elute immunocomplexes. Elution was repeated twice, and the samples
were further incubated overnight at 65 °C to reverse cross-linking, along with the untreated input (5% of
the starting material). On day 3, after treatment with 0.5 mg/ml Proteinase K for 1h at 65 °C, DNA was
purified with Zymo ChIP DNA Clear Concentrator kit and quantified with QUBIT.
For all ChIP-seq experiments, barcoded libraries were made with NEB ULTRA II DNA Library Prep Kit for
Illumina, and sequenced on Illumina NextSeq 500, producing 100 bp PE reads.
For ChIP-qPCR, on day 1 the sonicated lysate was aliquot into single immunoprecipitations of 2.5×10^6
cells each. A specific antibody or a total rabbit IgG control was added to the lysate along with Protein A
magnetic beads (Invitrogen) and incubated at 4 °C overnight. On day3, ChIP eluates and input were
assayed by real-time quantitative PCR in a 20 μl reaction with the following: 0.4 μM of each primer, 10 μl
of PowerUp SYBR Green (Applied Biosystems), and 5 μl of template DNA (corresponding to 1/40 of the
elution material) using the fast program on QuantStudio qPCR machine (Applied Biosystems). Thermal
cycling parameters were: 20sec at 95 °C, followed by 40 cycles of 1sec at 95°C, 20sec at 60°C. Used
antibodies are listed in the "Antibodies" section of the methods.
540
ChIP-seq Analyses
After removing the adapters with TrimGalore!, the sequences were aligned to the reference hg19, using
Burrows Wheeler Alignment tool (BWA), with the MEM algorithm[64] . Uniqueky mapping aligned reads
were filtered based on mapping quality (MAPQ > 10) to restrict our analysis to higher quality and likely
uniquely mapped reads, and PCR duplicates were removed. We called peaks for each individual using
MACS2[65] (H3K27ac) or Homer[66], at 5% FDR, with default parameters.
547
RNA-Seq
Cells were lysed in Tri-reagent (Zymo research) and total RNA was extracted using Quick-RNA Miniprep
kit (Zymo research) according to the manufacturer's instructions. RNA was further quantified using
DeNovix DS-11 Spectrophotometer while the RNA integrity was checked on Bioanalyzer 2100 (Agilent).
Only samples with RIN value above 8.0 were used for transcriptome analysis. RNA libraries were prepared
using 1 μg of total RNA input using NEBNext® Poly(A) mRNA Magnetic Isolation Module, NEBNext®
UltraTM II Directional RNA Library Prep Kit for Illumina® and NEBNext® UltraTM II DNA Library Prep Kit for
Illumina® according to the manufacturer's instructions (New England Biolabs).
556
557   **RNA-Seq Analyses**
558   After removing the adapters with TrimGalore!, Kallisto (Bray et al., 2016) was used to count reads mapping
559   to each gene. We analyzed differential gene expression levels with DESeq2 (Love et al., 2014) , with the
560   following model: design = ~condition, where condition indicates either Human or Chimpanzee.
561
562   **ATAC-Seq**
563   For ATAC-Seq experiments, 50,000 cells per condition were processed as described in the original ATAC-
564   seq protocol paper (Buenrostro et al. 2013). ATAC-seq data were processed with the same pipeline
565   described for ChIP-seq, with one modification: all mapped reads were offset by +4 bp for the forward-
566   strand and -5 bp for the reverse-strand. Peaks were called using MACS2 (Zhang et al., 2008).

567
Generation of the NCCIT-dCas9KRAB-SVAsgRNA Stable Cell Line
A dCas9-KRAB was cloned into a piggyBac transposon containing ampicillin and puromycin resistance (Addgene) which was obtained from the Wysocka Lab at Stanford University. The piggyBac dCas9-KRAB doxycycline-inducible plasmid, along with a piggyBac transposase (Cell Signaling), were transfected into NCCIT cells (ATCC) at ~70% confluency using a 6:1 ratio of Fugene HD (Promega) for 48 hours in ATCC-formulated RPMI media. Two days post-transfection, the media was changed and the transfected cells were selected for using 1µg puromycin per 1mL media. A piggyBac transposon plasmid containing two sgRNAs (SVAsgRNA1: 5'CTCCCTAATCTCAAGTACCC 3' and SVAsgRNA2: 5' TGTTTCAGAGAGCACGGGGT 3'; Integrated DNA Technologies) targeting ~80% of all annotated SVAs in humans (Pontis et al. 2019), along with a piggyBac transposase, were transfected into the NCCIT-dCas9KRAB cells using a 6:1 ratio of Fugene HD for 48 hours in ATCC-formulated RPMI media. Two days post-transfection, the media was changed and the transfected cells were selected for using 400µg geneticin per 1mL of media in addition to 1µg puromycin per 1mL media. The NCCIT-dCas9KRAB-SVAsgRNA cell line was maintained in ATCC-formulated RPMI media supplemented with 10% tet-free FBS, 1% L-glutamine, 1µg/mL puromycin, and 400µg/mL geneticin and incubated at 5% CO_2, 20% O_2 at 37°C.

Retinoic Acid-induced neuronal differentiation and CRISPR-interference of NCCIT-dCas9KRAB-SVAsgRNA cells
The NCCIT-dCas9KRAB-SVAsgRNA cells, at ~20% confluency, were treated with 10µM retinoic acid (RA) per 10mL media for 1 week to induce neuronal differentiation. At day 4, the media was refreshed and the cells were additionally treated with 2µg doxycycline per 1mL of media for 3 days. The cells were collected on day 7 of RA treatment (day 3 of doxycycline treatment) for qPCR and genomic experiments. Expression of the doxycycline-inducible dCas9 was verified via western blot.

Statistical and genomic analyses
All statistical analyses were performed using R v3.3.1 or Graphpad Prism version 9.2.0 for Mac OS X . BEDtools v2.27.1 (Quinlan and Hall, 2010) was used for genomic analyses. Pathway analysis was performed with Ingenuity Pathway Analysis Suite (QIAGEN Inc., https://www.qiagenbioinformatics.com/products/ingenuity-pathway-analysis). Motif analyses were performed using the MEME-Suite (Bailey et al., 2015), and specifically with the Meme-ChIP application. Fasta files of the regions of interest were produced using BEDTools v2.27.1. Shuffled input sequences were used as background. E-values < 0.001 were used as threshold for significance. Orthologous ATAC-seq regions were identified using the University of California Santa Cruz Genome Browser tool LiftOver.

Acknowledgements
The authors are grateful for Dr. Yoav Gilad (University of Chicago) for providing all the chimpanzee lines and one of the human lines, and to Dr. Joanna Wysocka's lab, and in particular Dr. Raquel Fueyo (Stanford University), for providing CRISPR reagents and critical advice. The authors thank the Thomas Jefferson Stem Cell and Regenerative Neuroscience Center for the support in the process of optimization of the iPSC differentiation. The authors are grateful to Trizzino lab members Luca Pagliaroli, PhD, Chiara Scopa, PhD and Connor Ott for help with specific analyses and experiments. The authors thank the Genomic Facility at The Wistar Institute (Philadelphia, PA) for the Next Generation Illumina Sequencing. The PAX6 antibody developed by Kawakami, A. was obtained from the Developmental Hybridoma Bank, created by the NICHD of the NIH and maintained at The University of Iowa, Department of Biology, Iowa City, IA 52242.

Competing interests
The authors declare no competing interests.

Funding
For this work, M.T. was funded by the National Institute of Health (NIH-NIGMS R35GM138344) and by the G. Harold and Leila Y. Mathers Foundation.

Data availability
The original genome-wide data generated in this study have been deposited in the GEO database under accession code GSE189347.

Author contributions
MT and SP designed the project. SP performed most of the experiments. SB carried out the CRISPR experiments in NCCIT cells. MT and SP analyzed the data and wrote the manuscript. All the authors read and approved the manuscript.

References
Abrajano, J. J., Qureshi, I. A., Gokhan, S., Zheng, D., Bergman, A. and Mehler, M. F. (2009). REST and CoREST Modulate Neuronal Subtype Specification, Maturation and Maintenance. *PLOS ONE* 4, e7936.

Agoglia, R. M., Sun, D., Birey, F., Yoon, S.-J., Miura, Y., Sabatini, K., Pașca, S. P. and Fraser, H. B. (2021). Primate cell fusion disentangles gene regulatory divergence in neurodevelopment. *Nature* 592, 421–427.

Andersen, J., Urbán, N., Achimastou, A., Ito, A., Simic, M., Ullom, K., Martynoga, B., Lebel, M., Göritz, C., Frisén, J., et al. (2014). A Transcriptional Mechanism Integrating Inputs from Extracellular Signals to Activate Hippocampal Stem Cells. *Neuron* 83, 1085–1097.

Arnold, S. J., Huang, G.-J., Cheung, A. F. P., Era, T., Nishikawa, S.-I., Bikoff, E. K., Molnar, Z., Robertson, E. J. and Groszer, M. (2008). The T-box transcription factor Eomes/Tbr2 regulates neurogenesis in the cortical subventricular zone. *Genes & Development* 22, 2479–2484.

Aruga, J. (2004). The role of Zic genes in neural development. *Mol Cell Neurosci* 26, 205–221.

Averaimo, S., Gritti, M., Barini, E., Gasparini, L. and Mazzanti, M. (2014). CLIC1 functional expression is required for cAMP-induced neurite elongation in post-natal mouse retinal ganglion cells. *Journal of Neurochemistry* 131, 444–456.

Bailey, T. L., Johnson, J., Grant, C. E. and Noble, W. S. (2015). The MEME Suite. *Nucleic Acids Research* 43, W39–W49.

Barger, N., Hanson, K. L., Teffer, K., Schenker-Ahmed, N. M. and Semendeferi, K. (2014). Evidence for evolutionary specialization in human limbic structures. *Front. Hum. Neurosci.* 8,.

Bath, K. G., Akins, M. R. and Lee, F. S. (2012). BDNF control of adult SVZ neurogenesis. *Developmental Psychobiology* 54, 578–589.

Bekinschtein, P., Katche, C., Slipczuk, L. N., Igaz, L. M., Cammarota, M., Izquierdo, I. and Medina, J. H. (2007). mTOR signaling in the hippocampus is necessary for memory formation. *Neurobiology of Learning and Memory* 87, 303–307.

Bray, N. L., Pimentel, H., Melsted, P. and Pachter, L. (2016). Near-optimal probabilistic RNA-seq
quantification. *Nat Biotechnol* 34, 525–527.

Bulfone, A., Martinez, S., Marigo, V., Campanella, M., Basile, A., Quaderi, N., Gattuso, C., Rubenstein, J.
L. R. and Ballabio, A. (1999). Expression pattern of the Tbr2 (Eomesodermin) gene during mouse
and chick brain development. *Mechanisms of Development* 84, 133–138.

Burgess, N., Maguire, E. A. and O'Keefe, J. (2002). The Human Hippocampus and Spatial and Episodic
Memory. *Neuron* 35, 625–641.

Chen, X., Wang, S., Zhou, Y., Han, Y., Li, S., Xu, Q., Xu, L., Zhu, Z., Deng, Y., Yu, L., et al. (2018). Phf8
histone demethylase deficiency causes cognitive impairments through the mTOR pathway. *Nat
Commun* 9, 114.

Cheval, H., Chagneau, C., Levasseur, G., Veyrac, A., Faucon-Biguet, N., Laroche, S. and Davis, S. (2012).
Distinctive features of Egr transcription factor regulation and DNA binding activity in CA1 of the
hippocampus in synaptic plasticity and consolidation and reconsolidation of fear memory.
Hippocampus 22, 631–642.

Chuong, E. B., Rumi, M. A. K., Soares, M. J. and Baker, J. C. (2013). Endogenous retroviruses function as
species-specific enhancer elements in the placenta. *Nat Genet* 45, 325–329.

Chuong, E. B., Elde, N. C. and Feschotte, C. (2016). Regulatory evolution of innate immunity through co-
option of endogenous retroviruses. *Science* 351, 1083–1087.

Cipriani, S., Nardelli, J., Verney, C., Delezoide, A.-L., Guimiot, F., Gressens, P. and Adle-Biassette, H.
(2016). Dynamic Expression Patterns of Progenitor and Pyramidal Neuron Layer Markers in the
Developing Human Hippocampus. *Cereb. Cortex* 26, 1255–1271.

Coluccio, A., Ecco, G., Duc, J., Offner, S., Turelli, P. and Trono, D. (2018). Individual retrotransposon
integrants are differentially controlled by KZFP/KAP1-dependent histone methylation, DNA
methylation and TET-mediated hydroxymethylation in naïve embryonic stem cells. *Epigenetics &
Chromatin* 11, 7.

Cosby, R. L., Judd, J., Zhang, R., Zhong, A., Garry, N., Pritham, E. J. and Feschotte, C. (2021). Recurrent
evolution of vertebrate transcription factors by transposase capture. *Science* 371, eabc6405.

Coufal, N. G., Garcia-Perez, J. L., Peng, G. E., Yeo, G. W., Mu, Y., Lovci, M. T., Morell, M., O'Shea, K. S.,
Moran, J. V. and Gage, F. H. (2009). L1 retrotransposition in human neural progenitor cells.
Nature 460, 1127–1131.

Crawley, J. N., Heyer, W.-D. and LaSalle, J. M. (2016). Autism and Cancer Share Risk Genes, Pathways,
and Drug Targets. *Trends in Genetics* 32, 139–146.

Datson, N. a., Morsink, M. c., Steenbergen, P. j., Aubert, Y., Schlumbohm, C., Fuchs, E. and de Kloet, E. r.
(2009). A molecular blueprint of gene expression in hippocampal subregions CA1, CA3, and DG is
conserved in the brain of the common marmoset. *Hippocampus* 19, 739–752.

Davis, L., Onn, I. and Elliott, E. (2018). The emerging roles for the chromatin structure regulators CTCF
and cohesin in neurodevelopment and behavior. *Cell. Mol. Life Sci.* 75, 1205–1214.

del Rosario, R. C. H., Rayan, N. A. and Prabhakar, S. (2014). Noncoding origins of anthropoid traits and a
new null model of transposon functionalization. *Genome Res* 24, 1469–1484.

Du, J., Leung, A., Trac, C., Lee, M., Parks, B. W., Lusis, A. J., Natarajan, R. and Schones, D. E. (2016).
Chromatin variation associated with liver metabolism is mediated by transposable elements.
Epigenetics & Chromatin 9, 28.

Duyckaerts, C., Delatour, B. and Potier, M.-C. (2009). Classification and basic pathology of Alzheimer
disease. *Acta Neuropathol* 118, 5–36.

Edler, M. K., Sherwood, C. C., Meindl, R. S., Hopkins, W. D., Ely, J. J., Erwin, J. M., Mufson, E. J., Hof, P. R.
and Raghanti, M. A. (2017). Aged chimpanzees exhibit pathologic hallmarks of Alzheimer's
disease. *Neurobiology of Aging* 59, 107–120.

Eichenbaum, H. (2017a). The role of the hippocampus in navigation is memory. *Journal of
Neurophysiology* 117, 1785–1796.

Eichenbaum, H. (2017b). Prefrontal–hippocampal interactions in episodic memory. *Nat Rev Neurosci* 18,
547–558.

Emera, D., Yin, J., Reilly, S. K., Gockley, J. and Noonan, J. P. (2016). Origin and evolution of
developmental enhancers in the mammalian neocortex. *PNAS* 113, E2617–E2626.

Enard, W. (2011). FOXP2 and the role of cortico-basal ganglia circuits in speech and language evolution.
Current Opinion in Neurobiology 21, 415–424.

Enard, W., Khaitovich, P., Klose, J., Zöllner, S., Heissig, F., Giavalisco, P., Nieselt-Struwe, K., Muchmore,
E., Varki, A., Ravid, R., et al. (2002). Intra- and Interspecific Variation in Primate Gene Expression
Patterns. *Science* 296, 340–343.

Englund, C. (2005). Pax6, Tbr2, and Tbr1 Are Expressed Sequentially by Radial Glia, Intermediate
Progenitor Cells, and Postmitotic Neurons in Developing Neocortex. *Journal of Neuroscience* 25,
247–251.

Ferrón, S. R., Charalambous, M., Radford, E., McEwen, K., Wildner, H., Hind, E., Morante-Redolat, J. M.,
Laborda, J., Guillemot, F., Bauer, S. R., et al. (2011). Postnatal loss of Dlk1 imprinting in stem
cells and niche astrocytes regulates neurogenesis. *Nature* 475, 381–385.

Finch, C. E. and Austad, S. N. (2015). Commentary: is Alzheimer's disease uniquely human? *Neurobiology
of Aging* 36, 553–555.

Florio, M. and Huttner, W. B. (2014). Neural progenitors, neurogenesis and the evolution of the
neocortex. *Development* 141, 2182–2194.

Fortress, A. M., Fan, L., Orr, P. T., Zhao, Z. and Frick, K. M. (2013). Estradiol-induced object recognition
memory consolidation is dependent on activation of mTOR signaling in the dorsal hippocampus.
Learn. Mem. 20, 147–155.

Fuentes, D. R., Swigut, T. and Wysocka, J. (2018). Systematic perturbation of retroviral LTRs reveals
widespread long-range effects on human gene regulation. *eLife* 7, e35989.

Gallego Romero, I., Pavlovic, B. J., Hernando-Herraez, I., Zhou, X., Ward, M. C., Banovich, N. E., Kagan, C.
L., Burnett, J. E., Huang, C. H., Mitrano, A., et al. (2015). A panel of induced pluripotent stem
cells from chimpanzees: a resource for comparative functional genomics. *eLife* 4, e07103.

Gokhman, D., Agoglia, R. M., Kinnebrew, M., Gordon, W., Sun, D., Bajpai, V. K., Naqvi, S., Chen, C., Chan,
A., Chen, C., et al. (2021). Human–chimpanzee fused cells reveal cis -regulatory divergence
underlying skeletal evolution. *Nature Genetics* 53, 467–476.

Graber, T. E., McCamphill, P. K. and Sossin, W. S. (2013). A recollection of mTOR signaling in learning and
memory. *Learn. Mem.* 20, 518–530.

Hevner, R. F. (2019). Intermediate progenitors and Tbr2 in cortical development. *J. Anat.* 235, 616–625.

Hickey, S. L., Berto, S. and Konopka, G. (2019). Chromatin Decondensation by FOXP2 Promotes Human
Neuron Maturation and Expression of Neurodevelopmental Disease Genes. *Cell Reports* 27,
1699-1711.e9.

Hodge, R. D., Nelson, B. R., Kahoud, R. J., Yang, R., Mussar, K. E., Reiner, S. L. and Hevner, R. F. (2012).
Tbr2 Is Essential for Hippocampal Lineage Progression from Neural Stem Cells to Intermediate
Progenitors and Neurons. *Journal of Neuroscience* 32, 6275–6287.

Ito, J., Kimura, I., Soper, A., Coudray, A., Koyanagi, Y., Nakaoka, H., Inoue, I., Turelli, P., Trono, D. and
Sato, K. Endogenous retroviruses drive KRAB zinc-finger protein family expression for tumor
suppression. *Science Advances* 6, eabc3020.

Ivancevic, A. and Chuong, E. B. (2020). Transposable elements teach T cells new tricks. *PNAS* 117, 9145–
9147.

Jacques, P.-É., Jeyakani, J. and Bourque, G. (2013). The Majority of Primate-Specific Regulatory
Sequences Are Derived from Transposable Elements. *PLOS Genetics* 9, e1003504.

Jönsson, M. E., Ludvik Brattås, P., Gustafsson, C., Petri, R., Yudovich, D., Pircs, K., Verschuere, S.,
Madsen, S., Hansson, J., Larsson, J., et al. (2019). Activation of neuronal genes via LINE-1
elements upon global DNA demethylation in human neural progenitors. *Nat Commun* 10, 3182.

Judd, J., Sanderson, H. and Feschotte, C. (2021). Evolution of mouse circadian enhancers from
transposable elements. *Genome Biol* 22, 193.

Kamboh, M. I., Fan, K.-H., Yan, Q., Beer, J. C., Snitz, B. E., Wang, X., Chang, C.-C. H., Demirci, F. Y.,
Feingold, E. and Ganguli, M. (2019). Population-based genome-wide association study of
cognitive decline in older adults free of dementia: identification of a novel locus for the
attention domain. *Neurobiol Aging* 84, 239.e15-239.e24.

Kimura, N., Nakashima, K., Ueno, M., Kiyama, H. and Taga, T. (1999). A novel mammalian T-box-
containing gene, Tbr2, expressed in mouse developing brain. *Developmental Brain Research*
115, 183–193.

King, M. and Wilson, A. (1975). Evolution at two levels in humans and chimpanzees. *Science* 188, 107–
116.

Kittappa, R., Chang, W. W., Awatramani, R. B. and McKay, R. D. G. (2007). The foxa2 Gene Controls the
Birth and Spontaneous Degeneration of Dopamine Neurons in Old Age. *PLOS Biology* 5, e325.

Kohane, A. A. and Wood, T. R. (2021). *Neurodevelopmental clustering of gene expression identifies lipid*
metabolism genes associated with neuroprotection and neurodegeneration.

Kolehmainen, J., Black, G. C. M., Saarinen, A., Chandler, K., Clayton-Smith, J., Träskelin, A.-L., Perveen, R.,
Kivitie-Kallio, S., Norio, R., Warburg, M., et al. (2003). Cohen syndrome is caused by mutations in
a novel gene, COH1, encoding a transmembrane protein with a presumed role in vesicle-
mediated sorting and intracellular protein transport. *Am J Hum Genet* 72, 1359–1369.

Liégeois, F. J., Hildebrand, M. S., Bonthrone, A., Turner, S. J., Scheffer, I. E., Bahlo, M., Connelly, A. and
Morgan, A. T. (2016). Early neuroimaging markers of FOXP2 intragenic deletion. *Sci Rep* 6,
35192.

Love, M. I., Huber, W. and Anders, S. (2014). Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biology* 15, 550.

Lynch, V. J., Leclerc, R. D., May, G. and Wagner, G. P. (2011). Transposon-mediated rewiring of gene
regulatory networks contributed to the evolution of pregnancy in mammals. *Nat Genet* 43,
1154–1159.

Lynch, V. J., Nnamani, M. C., Kapusta, A., Brayer, K., Plaza, S. L., Mazur, E. C., Emera, D., Sheikh, S. Z.,
Grützner, F., Bauersachs, S., et al. (2015). Ancient Transposable Elements Transformed the
Uterine Regulatory Landscape and Transcriptome during the Evolution of Mammalian
Pregnancy. *Cell Rep* 10, 551–561.

MacDermot, K. D., Bonora, E., Sykes, N., Coupe, A.-M., Lai, C. S. L., Vernes, S. C., Vargha-Khadem, F.,
McKenzie, F., Smith, R. L., Monaco, A. P., et al. (2005). Identification of FOXP2 Truncation as a
Novel Cause of Developmental Speech and Language Deficits. *The American Journal of Human*
Genetics 76, 1074–1080.

Mandal, C., Jung, K. H., Kang, S. C., Choi, M. R., Park, K. S., Chung, I. Y. and Chai, Y. G. (2015). Knocking
down of UTX in NCCIT cells enhance cell attachment and promote early neuronal cell
differentiation. *BioChip J* 9, 182–193.

Marchetto, M. C., Hrvoj-Mihic, B., Kerman, B. E., Yu, D. X., Vadodaria, K. C., Linker, S. B., Narvaiza, I.,
Santos, R., Denli, A. M., Mendes, A. P., et al. (2019). Species-specific maturation profiles of
human, chimpanzee and bonobo neural cells. *eLife* 8, e37527.

Martínez-Cerdeño, V., Noctor, S. C. and Kriegstein, A. R. (2006). The role of intermediate progenitor cells
in the evolutionary expansion of the cerebral cortex. *Cereb Cortex* 16 Suppl 1, i152-161.

Mathys, H., Davila-Velderrain, J., Peng, Z., Gao, F., Mohammadi, S., Young, J. Z., Menon, M., He, L.,
Abdurrob, F., Jiang, X., et al. (2019). Single-cell transcriptomic analysis of Alzheimer's disease.
Nature 570, 332–337.

Miao, B., Fu, S., Lyu, C., Gontarz, P., Wang, T. and Zhang, B. (2020). Tissue-specific usage of transposable
element-derived promoters in mouse development. *Genome Biol* 21, 255.

Mihalas, A. B., Elsen, G. E., Bedogni, F., Daza, R. A. M., Ramos-Laguna, K. A., Arnold, S. J. and Hevner, R.
F. (2016). Intermediate Progenitor Cohorts Differentially Generate Cortical Layers and Require
Tbr2 for Timely Acquisition of Neuronal Subtype Identity. *Cell Reports* 16, 92–105.

Mika, K., Marinić, M., Singh, M., Muter, J., Brosens, J. J. and Lynch, V. J. (2021). Evolutionary
transcriptomics implicates new genes and pathways in human pregnancy and adverse
pregnancy outcomes. *eLife* 10, e69584.

Modzelewski, A. J., Shao, W., Chen, J., Lee, A., Qi, X., Noon, M., Tjokro, K., Sales, G., Biton, A., Anand, A.,
et al. (2021). A mouse-specific retrotransposon drives a conserved Cdk2ap1 isoform essential for
development. *Cell* 184, 5541-5558.e22.

Mora-Bermúdez, F., Badsha, F., Kanton, S., Camp, J. G., Vernot, B., Köhler, K., Voigt, B., Okita, K., Maricic,
T., He, Z., et al. (2016). Differences and similarities between human and chimpanzee neural
progenitors during cerebral cortex development. *eLife* 5, e18683.

Mukherjee, D., Gonzales, B. J., Ashwal-Fluss, R., Turm, H., Groysman, M. and Citri, A. (2021). Egr2
induction in spiny projection neurons of the ventrolateral striatum contributes to cocaine place
preference in mice. *eLife* 10, e65228.

Okhovat, M., Nevonen, K. A., Davis, B. A., Michener, P., Ward, S., Milhaven, M., Harshman, L., Sohota, A.,
Fernandes, J. D., Salama, S. R., et al. (2020). Co-option of the lineage-specific LAVA
retrotransposon in the gibbon genome. *PNAS* 117, 19328–19338.

Pagliaroli, L., Porazzi, P., Curtis, A. T., Scopa, C., Mikkers, H. M. M., Freund, C., Daxinger, L., Deliard, S.,
Welsh, S. A., Offley, S., et al. (2021). Inability to switch from ARID1A-BAF to ARID1B-BAF impairs
exit from pluripotency and commitment towards neural crest formation in ARID1B-related
neurodevelopmental disorders. *Nat Commun* 12, 6469.

Pashos, E. E., Park, Y., Wang, X., Raghavan, A., Yang, W., Abbey, D., Peters, D. T., Arbelaez, J., Hernandez,
M., Kuperwasser, N., et al. (2017). Large, Diverse Population Cohorts of hiPSCs and Derived
Hepatocyte-like Cells Reveal Functional Genetic Variation at Blood Lipid-Associated Loci. *Cell*
Stem Cell 20, 558-570.e10.

Playfoot, C. J., Duc, J., Sheppard, S., Dind, S., Coudray, A., Planet, E. and Trono, D. (2021). Transposable
elements and their KZFP controllers are drivers of transcriptional innovation in the developing
human brain. *Genome Res.* 31, 1531–1545.

Poirier, R., Cheval, H., Mailhes, C., Charnay, P., Davis, S. and Laroche, S. (2007). Paradoxical Role of an
Egr Transcription Factor Family Member, Egr2/Krox20, in Learning and Memory. *Front Behav*
Neurosci 1, 6.

Pontious, A., Kowalczyk, T., Englund, C. and Hevner, R. F. (2008). Role of intermediate progenitor cells in
cerebral cortex development. *Dev Neurosci* 30, 24–32.

Pontis, J., Planet, E., Offner, S., Turelli, P., Duc, J., Coudray, A., Theunissen, T. W., Jaenisch, R. and Trono,
D. (2019). Hominoid-Specific Transposable Elements and KZFPs Facilitate Human Embryonic
Genome Activation and Control Transcription in Naive Human ESCs. *Cell Stem Cell* 24, 724-
735.e5.

Quinlan, A. R. and Hall, I. M. (2010). BEDTools: a flexible suite of utilities for comparing genomic
features. *Bioinformatics* 26, 841–842.

Quinn, J. P. and Bubb, V. J. (2014). SVA retrotransposons as modulators of gene expression. *Mobile
Genetic Elements* 4, e32102.

Raivich, G. (2008). c-Jun expression, activation and function in neural cell death, inflammation and
repair. *J Neurochem* 107, 898–906.

Raivich, G. and Behrens, A. (2006). Role of the AP-1 transcription factor c-Jun in developing, adult and
injured brain. *Prog Neurobiol* 78, 347–363.

Rayan, N. A., Del Rosario, R. C. H. and Prabhakar, S. (2016). Massive contribution of transposable
elements to mammalian regulatory sequences. *Semin Cell Dev Biol* 57, 51–56.

Reilly, S. K., Yin, J., Ayoub, A. E., Emera, D., Leng, J., Cotney, J., Sarro, R., Rakic, P. and Noonan, J. P.
(2015). Evolutionary genomics. Evolutionary changes in promoter and enhancer activity during
human corticogenesis. *Science* 347, 1155–1159.

Rétaux, S., Bourrat, F., Joly, J.-S. and Hinaux, H. (2013). Perspectives in Evo-Devo of the Vertebrate Brain.
In *Advances in Evolutionary Developmental Biology*, pp. 151–172. John Wiley & Sons, Ltd.

Reuter, M. S., Krumbiegel, M., Schlüter, G., Ekici, A. B., Reis, A. and Zweier, C. (2017). Haploinsufficiency
of NR4A2 is associated with a neurodevelopmental phenotype with prominent language
impairment. *Am J Med Genet A* 173, 2231–2234.

Sakamoto, K., Karelina, K. and Obrietan, K. (2011). CREB: a multifaceted regulator of neuronal plasticity
and protection. *Journal of Neurochemistry* 116, 1–9.

Sarkar, A., Mei, A., Paquola, A. C. M., Stern, S., Bardy, C., Klug, J. R., Kim, S., Neshat, N., Kim, H. J., Ku, M.,
et al. (2018). Efficient Generation of CA3 Neurons from Human Pluripotent Stem Cells Enables
Modeling of Hippocampal Connectivity In Vitro. *Cell Stem Cell* 22, 684-697.e9.

Schmidt, D., Schwalie, P. C., Wilson, M. D., Ballester, B., Gonçalves, A., Kutter, C., Brown, G. D., Marshall,
A., Flicek, P. and Odom, D. T. (2012). Waves of retrotransposon expansion remodel genome
organization and CTCF binding in multiple mammalian lineages. *Cell* 148, 335–348.

Sessa, A., Mao, C.-A., Hadjantonakis, A.-K., Klein, W. H. and Broccoli, V. (2008). Tbr2 directs conversion
of radial glia into basal precursors and guides neuronal amplification by indirect neurogenesis in
the developing neocortex. *Neuron* 60, 56–69.

Shah, A. H., Gilbert, M., Ivan, M. E., Komotar, R. J., Heiss, J. and Nath, A. (2021). The role of human
endogenous retroviruses in gliomas: from etiological perspectives and therapeutic implications.
Neuro-Oncology 23, 1647–1655.

Sin, C., Li, H. and Crawford, D. A. (2015). Transcriptional Regulation by FOXP1, FOXP2, and FOXP4
Dimerization. *Journal of Molecular Neuroscience* 55, 437–448.

Sousa, A. M. M., Zhu, Y., Raghanti, M. A., Kitchen, R. R., Onorati, M., Tebbenkamp, A. T. N., Stutz, B.,
Meyer, K. A., Li, M., Kawasawa, Y. I., et al. (2017). Molecular and cellular reorganization of neural
circuits in the human lineage. *Science* 358, 1027–1032.

Squire, L. R. (1992). Memory and the Hippocampus: A Synthesis From Findings With Rats, Monkeys, and
Humans.

Sundaram, V. and Wysocka, J. (2020). Transposable elements as a potent source of diverse cis-
regulatory sequences in mammalian genomes. *Philosophical Transactions of the Royal Society B:*
Biological Sciences 375, 20190347.

Sundaram, V., Cheng, Y., Ma, Z., Li, D., Xing, X., Edge, P., Snyder, M. P. and Wang, T. (2014). Widespread
contribution of transposable elements to the innovation of gene regulatory networks. *Genome*
Res 24, 1963–1976.

Sur, D., Kustwar, R. K., Budania, S., Mahadevan, A., Hancks, D. C., Yadav, V., Shankar, S. K. and Mandal, P.
K. (2017). Detection of the LINE-1 retrotransposon RNA-binding protein ORF1p in different
anatomical regions of the human brain. *Mobile DNA* 8, 17.

Taher, N., McKenzie, C., Garrett, R., Baker, M., Fox, N. and Isaacs, G. D. (2014). Amyloid-β Alters the DNA
Methylation Status of Cell-fate Genes in an Alzheimer's Disease Model. *Journal of Alzheimer's*
Disease 38, 831–844.

Thomas, C. A., Paquola, A. C. M. and Muotri, A. R. (2012). LINE-1 Retrotransposition in the Nervous
System. *Annual Review of Cell and Developmental Biology* 28, 555–573.

Tokuyama, M., Kong, Y., Song, E., Jayewickreme, T., Kang, I. and Iwasaki, A. (2018). ERVmap analysis
reveals genome-wide transcription of human endogenous retroviruses. *PNAS* 115, 12565–
12572.

Tomasello, M. and Herrmann, E. (2010). Ape and Human Cognition: What's the Difference? *Current*
Directions in Psychological Science 19, 3–8.

Trizzino, M., Park, Y., Holsbach-Beltrame, M., Aracena, K., Mika, K., Caliskan, M., Perry, G. H., Lynch, V. J.
and Brown, C. D. (2017). Transposable elements are the primary source of novelty in primate
gene regulation. *Genome Res.*

Trizzino, M., Kapusta, A. and Brown, C. D. (2018). Transposable elements generate regulatory novelty in
a tissue-specific fashion. *BMC Genomics* 19, 468.

Turelli, P., Playfoot, C., Grun, D., Raclot, C., Pontis, J., Coudray, A., Thorball, C., Duc, J., Pankevich, E. V.,
Deplancke, B., et al. (2020). Primate-restricted KRAB zinc finger proteins and target
retrotransposons control gene expression in human neurons. *Science Advances* 6, eaba3200.

Walker, L. C. and Jucker, M. (2017). The Exceptional Vulnerability of Humans to Alzheimer's Disease.
Trends in Molecular Medicine 23, 534–545.

Wang, H., Xing, J., Grover, D., Hedges, D. J., Han, K., Walker, J. A. and Batzer, M. A. (2005). SVA Elements:
A Hominid-specific Retroposon Family. *Journal of Molecular Biology* 354, 994–1007.

Ward, M. C., Zhao, S., Luo, K., Pavlovic, B. J., Karimi, M. M., Stephens, M. and Gilad, Y. (2018). Silencing
of transposable elements may not be a major driver of regulatory evolution in primate iPSCs.
eLife 7, e33084.

Wei, Y. (2020). Comparative transcriptome analysis of the hippocampus from sleep-deprived and
Alzheimer's disease mice. *Genet Mol Biol* 43, e20190052.

Wray, G. A. (2007). The evolutionary significance of cis-regulatory mutations. *Nat Rev Genet* 8, 206–216.

Xie, M., Hong, C., Zhang, B., Lowdon, R. F., Xing, X., Li, D., Zhou, X., Lee, H. J., Maire, C. L., Ligon, K. L., et
al. (2013). DNA hypomethylation within specific transposable element families associates with
tissue-specific enhancer landscape. *Nat Genet* 45, 836–841.

Yang, W., Liu, Y., Slovik, K. J., Wu, J. C., Duncan, S. A., Rader, D. J. and Morrisey, E. E. (2015). Generation
of iPSCs as a Pooled Culture Using Magnetic Activated Cell Sorting of Newly Reprogrammed
Cells. *PLoS One* 10, e0134995.

Yi, R., Chen, B., Zhao, J., Zhan, X., Zhang, L., Liu, X. and Dong, Q. (2014). Krüppel-like Factor 8 Ameliorates
Alzheimer's Disease by Activating β-Catenin. *J Mol Neurosci* 52, 231–241.

Yu, D. X., Di Giorgio, F. P., Yao, J., Marchetto, M. C., Brennand, K., Wright, R., Mei, A., Mchenry, L., Lisuk,
D., Grasmick, J. M., et al. (2014). Modeling Hippocampal Neurogenesis Using Human Pluripotent
Stem Cells. *Stem Cell Reports* 2, 295–310.

Zhang, Y., Liu, T., Meyer, C. A., Eeckhoute, J., Johnson, D. S., Bernstein, B. E., Nusbaum, C., Myers, R. M.,
Brown, M., Li, W., et al. (2008). Model-based Analysis of ChIP-Seq (MACS). *Genome Biology* 9,
R137.

Zhang, J., Li, H., Yabut, O., Fitzpatrick, H., D'Arcangelo, G. and Herrup, K. (2010). Cdk5 Suppresses the
Neuronal Cell Cycle by Disrupting the E2F1–DP1 Complex. *J. Neurosci.* 30, 5219–5228.

Zhang, H., Xue, C., Shah, R., Bermingham, K., Hinkle, C. C., Li, W., Rodrigues, A., Tabita-Martinez, J.,
Millar, J. S., Cuchel, M., et al. (2015). Functional Analysis and Transcriptomic Profiling of iPSC-
Derived Macrophages and Their Application in Modeling Mendelian Disease. *Circulation*
Research 117, 17–28.

Zhao, B., Wu, Q., Ye, A. Y., Guo, J., Zheng, X., Yang, X., Yan, L., Liu, Q.-R., Hyde, T. M., Wei, L., et al.
(2019). Somatic LINE-1 retrotransposition in cortical neurons and non-brain tissues of Rett
patients and healthy individuals. *PLOS Genetics* 15, e1008043.

937
938
Figure legends

940
941   **Figure 1** – Differentiation of human and chimpanzee iPSCs into hippocampal intermediate progenitors (A)
942 Western Blot of human differentiated iPSCs against TBR2 (73kDa) at day 0 (iPSCs), day 5, and day 11 of
943 treatment with hpIPC differentiation media. GAPDH is shown as the loading control. (B)
944 Immunofluorescent staining of human and chimpanzee hpIPCs after five days against
945 neurodevelopmental markers PAX6, TBR2 and OTX2. Nuclei are counterstained with DAPI.

946
Figure 2 – Differential Gene Expression In Human and Chimpanzee hpIPCs. (A) Schematic of iPSC
differentiation followed by RNA-seq library generation and analysis. (B) Volcano plot depicting "Human
UP" genes (right) and "Chimp UP" genes (left) with a log2FoldChange threshold of 1.5 and -1.5,
respectively, and a p-value threshold of 0.05. (C) Top upstream regulators of the differentially expressed
genes predicted by Ingenuity Pathway Analysis, ranked by -log10(p-value). (D) Heatmaps depicting the
expression of genes predicted to be under the control of the transcription factors CREB1, FOXA2 and TBR2
or predicted to play a role in embryonic development. The rows of the heatmaps indicate the transcript
names, columns indicate the species, sample number and replicate. "HS1 Rep 1" indicates Homo sapiens
sample 1 replicate 1 and "PT2 Rep 2" indicates Pan troglodytes sample 2 replicate 2

956
Figure 3 – Human Centric Chromatin Accessibility Analysis. (A) Schematic of iPSC differentiation followed
by ATAC-seq library generation and analysis. (B) UCSC Genome Browser visualization of human ATAC-seq
libraries (HS Rep 1, Rep 2, Rep 3) aligned to hg19 genome assembly. (C) Distribution of the differentially
accessible (DA) peaks into enhancers or promoters, and into Human UP (greater accessibility in humans)
or Chimp UP (greater accessibility in chimpanzees). (D) Differentially accessible chromatin regions (p <
0.05, n = 3006) are more likely to overlap with transposable elements (TE) than non-differentially
accessible regions (p > 0.9, n = 3,006). (E) The 1335 TE-derived DA regions are more likely to be near a
differentially expressed gene, as compared to DA regions that do not overlap a TE. (F) Distribution of the
five major TE classes in the human genome, compared with their distribution among the TE-derived DA
chromatin regions, and among the TE-derived DA chromatin regions proximal to a differentially expressed
gene

968
Figure 4 – Chimpanzee Centric Chromatin Accessibility Analysis. (A) Schematic of iPSC differentiation
followed by ATAC-seq library generation and analysis. (B) UCSC Genome Browser visualization of human
ATAC-seq libraries (PT Rep 1, Rep 2, Rep 3) aligned to hg19 genome assembly. (C) Distribution of the
differentially accessible (DA) peaks into enhancers or promoters, and into Human UP (greater accessibility
in humans) or Chimp UP (greater accessibility in chimpanzees). (D) Differentially accessible chromatin
regions (p < 0.05, n = 3,806) are more likely to overlap with transposable elements than other accessible
regions (p > 0.9, n = 3806. (E) Distribution of DA regions that overlap a TE and proximity to genes
differentially expressed between human and chimpanzee hpIPCs. (F) Distribution of the five major TE
classes in the chimpanzee genome, compared with their distribution among the 1410 TE-derived DA
chromatin regions (G) Breakdown of the 20 Miscellanous TEs represented in the 1,410 TE-derived DA
chromatin regions. (H) Distribution of SVA family of TEs in the human genome as compared to their
distribution in the TE-derived DA chromatin regions (n = 16).

981
Figure 5 – LTRs Are Enriched Among Human and Chimpanzee Differentially Accessible Transposons. (A)
Distribution of LTRs compared to non-LTRs in the human genome and in the 1,335 human TE-derived DA
regions, with predicted binding motifs. (B) Differentially expressed genes close to the human-enriched DA

LTRs. (C) Distribution of LTRs compared to non-LTRs in the chimpanzee genome and in the 1,410
chimpanzee TE-derived DA regions, with predicted binding motifs. (D) Differentially expressed genes close
to the chimpanzee-enriched DA LTRs.
988
Figure 6 – Human-specific SVAs Bind Tbr2 and Influence Neurodevelopment (A) The SVAs enriched within
the differentially accessible chimpanzee peaks are enriched for binding motifs of neurodevelopmental
transcription factors KLF8, ZIC1, and ASCL1 as well as the AP-1 dimer. (B) Human SVAs are enriched for
the binding motif of TBR2. (C) Genes proximal to human-specific SVAs are predicted to be regulated by
neurodevelopmental TFs and function in important neuronal pathways. (D) 1,816 human SVAs exhibit
ATAC-seq signal in hpIPCs (red = signal, white = noise) and of these, 739 exhibit TBR2 ChIP signal (black =
signal, yellow = background). (E) Breakdown of 739 accessible, TBR2 bound SVAs into "human-specific"
and "conserved in chimpanzees". (F) Differentially expressed genes close to the human-specific TBR2-
bound SVAs.
998
Figure 7 – CRISPR-interference on human SVAs. (A) Treating NCCITs with Retinoic Acid for 7 days leads to
induction of TBR2 and PAX6, as shown in qRT-PCR. (B) Treating the RA-NCCIT stable CRISPR line for 72
hours with doxycycline leads to dCas9 activation, as shown in the Cas9 immunoblot. (C) dCas9 activation
results in deposition of H3K9me3 at most human SVAs as shown in the H3K9me3 ChIP-seq heatmap. (D)
Volcano plot depicting genes differentially expressed upon doxycycline treatment. (E) Venn diagram
showing overlap between genes differentially expressed between human and chimpanzee's hippocampal
intermediate progenitors and between RA-NCCITs with and without doxycycline treatment (i.e. with and
without SVA repression). (F) Pie chart illustrating the fraction of TBR2-controlled genes that are
differentially expressed upon SVA repression in RA-NCCITs.

Figure 1 – Differentiation of human and chimpanzee iPSCs into hippocampal intermediate progenitors (A) Western Blot of human differentiated iPSCs against TBR2 (73kDa) at day 0 (iPSCs), day 5, and day 11 of treatment with hpIPC differentiation media. GAPDH is shown as the loading control. (B) Immunofluorescent staining of human and chimpanzee hpIPCs after five days against neurodevelopmental markers PAX6, TBR2 and OTX2. Nuclei are counterstained with DAPI.

Figure 2 – Differential Gene Expression In Human and Chimpanzee hpIPCs. (A) Schematic of iPSC differentiation followed by RNA-seq library generation and analysis. (B) Volcano plot depicting "Human UP" genes (right) and "Chimp UP" genes (left) with a log2FoldChange threshold of 1.5 and -1.5, respectively, and a p-value threshold of 0.05. (C) Top upstream regulators of the differentially expressed genes predicted by Ingenuity Pathway Analysis, ranked by -log10(p-value). (D) Heatmaps depicting the expression of genes predicted to be under the control of the transcription factors CREB1, FOXA2 and TBR2 or predicted to play a role in embryonic development.

Figure 3 – Human Centric Chromatin Accessibility Analysis. (A) Schematic of iPSC differentiation followed by ATAC-seq library generation and analysis. (B) UCSC Genome Browser visualization of human ATAC-seq libraries (HS Rep 1, Rep 2, Rep 3) aligned to hg19 genome assembly. (C) Distribution of the differentially accessible (DA) peaks into enhancers or promoters, and into Human UP (greater accessibility in humans) or Chimp UP (greater accessibility in chimpanzees). (D) Differentially accessible chromatin regions (p < 0.05, n = 3006) are more likely to overlap with transposable elements (TE) than non-differentially accessible regions (p > 0.9, n = 3,006). (E) The 1335 TE-derived DA regions are more likely to be near a differentially expressed gene, as compared to DA regions that do not overlap a TE. (F) Distribution of the five major TE classes in the human genome, compared with their distribution among the TE-derived DA chromatin regions, and among the TE-derived DA chromatin regions proximal to a differentially expressed gene

Figure 4 - Chimpanzee Centric Chromatin Accessibility Analysis. (A) Schematic of iPSC differentiation followed by ATAC-seq library generation and analysis. B) UCSC Genome Browser visualization of human ATAC-seq libraries (PT Rep 1, Rep 2, Rep 3) aligned to hg19 genome assembly. (C) Distribution of the differentially accessible (DA) peaks into enhancers or promoters, and into Human UP (greater accessibility in humans) or Chimp UP (greater accessibility in chimpanzees). (D) Differentially accessible chromatin regions (p < 0.05, n = 3,806) are more likely to overlap with transposable elements than other accessible regions (p > 0.9, n = 3806. (E) Distribution of DA regions that overlap a TE and proximity to genes differentially expressed between human and chimpanzee hpIPCs. (F) Distribution of the five major TE classes in the chimpanzee genome, compared with their distribution among the 1410 TE-derived DA chromatin regions (G) Breakdown of the 20 Miscellanous TEs represented in the 1,410 TE-derived DA chromatin regions. (H) Distribution of SVA family of TEs in the human genome as compared to their distribution in the TE-derived DA chromatin regions (n = 16).

Figure 5 – LTRs Are Enriched Among Human and Chimpanzee Differentially Accessible Transposons. (A) Distribution of LTRs compared to non-LTRs in the human genome and in the 1,335 human TE-derived DA regions, with predicted binding motifs. (B) Differentially expressed genes close to the human-enriched DA LTRs. (C) Distribution of LTRs compared to non-LTRs in the chimpanzee genome and in the 1,410 chimpanzee TE-derived DA regions, with predicted binding motifs. (D) Differentially expressed genes close to the chimpanzee-enriched DA LTRs.

Figure 6 – Human-specific SVAs Bind Tbr2 and Influence Neurodevelopment (A) The SVAs enriched within the differentially accessible chimpanzee peaks are enriched for binding motifs of neurodevelopmental transcription factors KLF8, ZIC1, and ASCL1 as well as the AP-1 dimer. (B) Human SVAs are enriched for the binding motif of TBR2 (C) Genes proximal to human-specific SVAs are predicted to be regulated by neurodevelopmental TFs and function in important neuronal pathways (D) 1,816 human SVAs exhibit ATAC-seq signal in hpIPCs (red = signal, white = noise) and of these, 739 exhibit TBR2 ChIP signal (black = signal, yellow = noise). (E) Breakdown of 739 accessible, TBR2 bound SVAs into "human-specific" and "conserved in chimpanzees". (F) Differentially expressed genes close to the human-specific TBR2-bound SVAs.

Figure 7 – CRISPR-interference on human SVAs. (A) Treating NCCITs with Retinoic Acid for 7 days leads to induction of TBR2 and PAX6, as shown in qRT-PCR. (B) Treating the RA-NCCIT stable CRISPR line for 72 hours with doxycycline leads to dCas9 activation, as shown in the Cas9 immunoblot. (C) dCas9 activation results in deposition of H3K9me3 at most human SVAs as shown in the H3K9me3 ChIP-seq heatmap (yellow = signal). (D) Volcano plot depicting genes differentially expressed upon doxycycline treatment. (E) Venn diagram showing overlap between genes differentially expressed between human and chimpanzee's hippocampal intermediate progenitors and between RA-NCCITs with and without doxycycline treatment (i.e. with and without SVA repression). (F) Pie chart illustrating the fraction of TBR2-controlled genes that are differentially expressed upon SVA repression in RA-NCCITs.