Population-level asymmetry of the cerebral cortex: reproducibility, lifespan changes, heritability, and individual differences

Authors and affiliations
James M. Roe[1]*, Didac Vidal-Piñeiro[1], Inge K. Amlien[1], Mengyu Pan[1], Markus H. Sneve[1], Michel Thiebaut de Schotten[2,3], Patrick Friedrich[4], Zhiqiang Sha[5], Clyde Francks[5,6], Yunpeng Wang[1], Kristine B. Walhovd[1,7], Anders M. Fjell[1,7] & René Westerhausen[8]

[1]Center for Lifespan Changes in Brain and Cognition (LCBC), Department of Psychology, University of Oslo, Norway.
[2]Groupe d'Imagerie Neurofonctionnelle, Institut des Maladies Neurodégénératives-UMR 5293, CNRS, CEA, University of Bordeaux, Bordeaux, France.
[3]Brain Connectivity and Behaviour Laboratory, Sorbonne Universities, Paris, France.
[4]Institute of Neuroscience and Medicine (INM-7: Brain and Behaviour), Research Centre Jülich, Jülich, Germany.
[5]Language and Genetics Department, Max Planck Institute for Psycholinguistics, Nijmegen, The Netherlands.
[6]Donders Institute Netherlands for Brain, Cognition and Behaviour, Radboud University, Nijmegen, The Netherlands.
[7]Department of Radiology and Nuclear Medicine, Oslo University Hospital, Oslo, Norway.
[8]Section for Cognitive and Clinical Neuroscience, Department of Psychology, University of Oslo, Norway.

*Address correspondence to James M. Roe, Department of Psychology, PO Box 1094 Blindern, 0317 Oslo, Norway.
Email: j.m.roe@psykologi.uio.no

Running title: Population-level cortical asymmetry
Keywords: Lateralization, population neuroscience, lifespan
Open access license: yes

The authors declare no competing interests

Abstract

Cortical asymmetry is a ubiquitous feature of brain organization that is altered in neurodevelopmental disorders, yet we lack knowledge of how its development proceeds across life in health. Achieving consensus on cortical asymmetries in humans is necessary to uncover the genetic-developmental mechanisms that shape them and factors moderating cortical lateralization. Here, we delineate population-level asymmetry in cortical thickness and surface area vertex-wise in 7 datasets and chart asymmetry trajectories across life (4-89 years; observations = 3937; 70% longitudinal). We reveal asymmetry interrelationships, heritability, and test associations in large-scale data (N=~37,500). Cortical asymmetry was robust across datasets. Whereas areal asymmetry is predominantly stable across life, thickness asymmetry grows in development and declines in aging. Areal asymmetry correlates in specific regions, whereas thickness asymmetry is globally interrelated across cortex in a pattern suggesting highly left-lateralized individuals tend towards left-lateralization also in population-level right-asymmetric regions (and vice versa). Areal asymmetry is moderately heritable (max h^2_{SNP} ~19%), and phenotypic correlations are reflected by high genetic correlations, whereas heritability of thickness asymmetry is low. Finally, we detected an asymmetry association with cognition and confirm recently-reported handedness links. Results suggest areal asymmetry is developmentally stable and arises in early life, whereas developmental changes in thickness asymmetry may lead to directional variability of global thickness lateralization. Our results bear enough reproducibility to serve as a standard for future brain asymmetry studies.

1. Introduction

The brain's hemispheres exhibit high contralateral symmetry [1,2] and homotopic regions are amongst the most genetically [3–5] and developmentally linked [3,6]. However, structural asymmetry is also a ubiquitous aspect of brain organization [7,8]. Cortical thickness (CT) and surface area (SA) are known to exhibit distinct asymmetry patterns [7,9], but these have been reported inconsistently [7,8,10–20]. Yet disrupted cortical asymmetry is a confirmed feature of neurodevelopmental disorders [21], aging [10], and Alzheimer's disease [10,22]. Hence, achieving consensus on cortical asymmetries and understanding the genetic-developmental and lifespan influences that shape and alter them is necessary to discover precise biomarkers for disease. To reach consensus, an atlas-free description of asymmetries that reliably replicate across international samples (i.e. population-level asymmetries) is needed. This would enable precision mapping of the genetic and individual-specific factors moderating cortical lateralization, and serve as a high-fidelity phenotype for future studies on brain asymmetry. Furthermore, it is unknown how cortical asymmetry development proceeds across life, as no previous study has mapped cortical asymmetry trajectories longitudinally across the lifespan.

Although several studies have mapped cortical asymmetry [7,8,18–20,23,10–17], conflicting results may be partly due to the use of brain atlases with varying spatial resolutions, especially if asymmetry conforms poorly to the predefined anatomical boundaries. Still, even amongst studies adopting an atlas-free approach, conflicting results abound [13–18]. For example, for CT asymmetry, medial prefrontal cortex (mPFC) has been reported to show both extensive rightward [14,16,17] and leftward [10–12,20] lateralization. Beyond regional inconsistencies, a recent meta-analysis confirmed the cortex is globally organized in a characteristic pattern of CT asymmetry, wherein anterior and posterior regions are thicker in the left and right hemisphere, respectively [7]. Notably, while this agrees with some reports [10–12,20,24], it is less compatible with many others [8,13–19,25]. Furthermore, that study applied a relatively course brain atlas, and there is currently no high-resolution complement to describe cortical asymmetries that reliably reproduce across international samples (but see [10,24]). For areal asymmetry, while results have been broadly more consistent [7,8,14,19,23,26], there nevertheless remain important discrepancies, such as reports of right- [7,27,28] and left- [14,26] lateralization of superior temporal sulcus (STS).

An accurate description of the lifespan trajectories of cortical asymmetry may shed light on mechanisms underlying diverse aspects of asymmetry across life. For CT, longitudinal increases in asymmetry have been shown during the first two years of life [11], with suggestions of rapid asymmetry growth from birth to 1 year [11], and continued growth until adolescence [29]. However, previous studies mapped CT asymmetry linearly across cross-sectional developmental and adult age-ranges [13,20], mostly concluding CT asymmetry is minimal in infancy and maximal age ~60. In contrast, recent work established CT asymmetry shows a non-linear decline from 20 to 90 years that is reproducible across aging cohorts [10]. Thus, although offering viable developmental insights [13,20], previous lifespan studies of CT asymmetry do not accurately capture the aging process, and likely conflate non-linear developmental and aging trajectories with linear models. A longitudinal exploration of the lifespan trajectories of CT asymmetry accounting for dynamic change is needed to further knowledge of normal brain development.

In addition, few studies have charted developmental [26,28] or aging effects [7] on SA asymmetry. However, indirect evidence suggests SA asymmetry may exhibit little change from birth to 2 years [26] despite rapid and concurrent developmental cortical expansion [30]. Determining the developmental timing of cortical asymmetry will provide a useful normative reference, as subtly altered asymmetry in neurodevelopmental disorders suggests early life perturbations in left-right brain organization contribute to lifelong detriment in brain health [21,31].

Correlations between cortical asymmetries in adults may provide a window on asymmetries formed under common genetic-developmental influence. Yet while there has been much research on whether asymmetries of various morphometric measures [8,14,19] or imaging modalities [27] relate to one another, few have focused on interrelationships between asymmetries derived from the same metric. Where reported, evidence suggests cortical asymmetries are mostly independent [32,33] – in line with a multifactorial view of asymmetry phenotypes [34–36] – and a recent study found asymmetry in anatomically connected regions of the cortical language network was no more related than in regions selected at random [27]. Currently, it is not known whether or how cortical asymmetries correlate within individuals, though this may suggest coordinated development of left-right brain asymmetries.

Altered lateralization has been hypothesized to relate to poorer cognitive outcomes [20,37,38]. In line with this, recent work suggests genetic overlap between cortical asymmetry, educational attainment, and neurodevelopmental disorders [31], and reduced brain torque [25,39,40] – a gross morphological asymmetry with a strong population-level bias – associates with lower cognition [41]. For CT and SA asymmetry, however, reported asymmetry-cognition associations have been conflicting [20,42,43] and remain untested in large-scale data. Furthermore, most large-scale studies of the factors moderating cortical asymmetry have adopted brain atlases offering limited spatial precision [7,31,44]. Accordingly, previous large-scale studies did not detect associations with handedness [7,45] that were not found until a recent study applied vertex-wise mapping in big data [24]. Similarly, it is unclear to what degree poor-fitting atlases drive down heritability estimates of cortical asymmetry [7,31], as estimates improve when brain measures better conform to the biology under genetic investigation [46,47]. However, no previous study has assessed heritability after precisely delineating regions of cortex that are asymmetric at the population level, and cortex-wide heritability maps have all used the same atlas we propose fits poorly to the asymmetry of cortex [48].

Here, we 1) delineate population-level cortical SA and CT asymmetries using vertex-wise analyses and their overlap in 7 international datasets, and 2) map their trajectories longitudinally across the lifespan. We 3) investigate interregional asymmetry correlations, asking whether and how asymmetries correlate within individuals. Next, we 4) tested heritability

117 of cortical asymmetry using both an extended twin design and genome-wide single nucleotide polymorphism (SNP) data.
118 Finally, we 5) screened our set of robust, population-level asymmetries for association with general cognitive ability,
119 handedness, sex, and brain size in UK Biobank (UKB) [49].
120
121
122
2. Results
2.1 Population-level asymmetry of the cerebral cortex
First, to delineate cortical regions exhibiting population-level SA and CT asymmetry, we assessed asymmetry vertex-
wise in 7 independent samples and quantified overlapping effects (Methods). SA asymmetries were markedly consistent
across all 7 datasets (Fig. 1A): the spatial overlap between AI maps ranged from *r* = .88 to .97 (Fig. 1C). Across all 7
datasets (Fig. 1D), strong leftward SA asymmetry was observed in a large cluster in supramarginal gyrus (SMG) that
spanned the length of postcentral gyrus, extended inferiorly into planum temporale and primary auditory regions and
conformed markedly to their anatomical boundaries (see Figure 1–figure supplement 1A for significance). We also
observed consistently strong leftward asymmetry in anterior insula, anterior temporal cortex, rostral anterior cingulate,
medially in superior frontal cortex, and precuneus, the latter extending the length of parahippocampal gyrus into
entorhinal cortex. Strong rightward SA asymmetry was consistently evident in cingulate cortex, inferior parietal cortex,
STS, lateral and medial occipital cortex, and in mPFC and rostral middle frontal cortex (Fig. 1A). The global pattern
agrees with previous reports [7,23,24], and effects showed markedly high overlap across datasets (Fig. 1D)
136
For CT, an anterior-posterior pattern of left-right asymmetry was evident in most datasets (Fig. 1B), consistent with recent
reports [7,10,24]. Though spatial correlations between AI maps were high, they were notably more variable (*r* = .33 - .93;
Fig. 1C); HCP showed lower correlation with all datasets (*r* = .33 - .46) whereas all other datasets correlated highly with
each other (min *r* = .78). Strong leftward CT asymmetry was evident in cingulate cortex, postcentral gyrus, and in superior
frontal cortex – with consistent effects across datasets (Fig. 1E) – and in medial and lateral prefrontal cortex, though the
latter two were less consistent among datasets. Strong rightward CT asymmetry was consistently evident in a large
cluster in and around STS and lateral temporal cortex (Fig 1E; Figure 1–figure supplement 1B), insula, lingual gyrus,
anterior parahippocampal and entorhinal cortex. Both SA and CT asymmetry extended beyond these described effects
(Figure 1–figure supplements 1-2).
146
147 Based on effect size criteria (Fig. 1D-E; Methods), we derived a set of robust clusters exhibiting population-level
148 asymmetry for SA (14 clusters) and CT (20 clusters) to be used in further analyses (see Supplementary file 1E-F for
149 anatomical descriptions, see Figure 1–figure supplement 3 for variances). We then formally compared our approach to
150 asymmetry estimates derived from a gyral-based atlas often used to assess asymmetry [7,19,31], finding fairly poor
151 correspondence with the vertex-wise structure of cortical asymmetry, particularly for CT (Figure 1–figure supplement 4).

Figure 1. A) Mean SA and *B)* CT asymmetry in each dataset. Warm and cold colours depict leftward and rightward asymmetry, respectively. *C)* Spatial overlap (Pearson's r) of the unthresholded maps between datasets for SA (lower matrix) and CT asymmetry (upper). *D)* Overlapping effects across datasets were used to delineate clusters exhibiting population-level SA (lower threshold = 5%) and *E)* CT asymmetry (lower threshold = 1%) based on a minimum 6-dataset overlap (black outlined clusters). Post=posterior; Lat=lateral; Med=medial; Ant=anterior; Sup=superior; Inf=inferior.

2.2 Lifespan trajectories of cortical asymmetry

We have recently shown that cortical regions exhibiting age-related reduction of CT asymmetry correspond with regions exhibiting strong asymmetry on average [10]. Thus, having delineated regions exhibiting population-level SA and CT asymmetry, we aimed to characterize the trajectories of SA and CT asymmetries longitudinally across the lifespan (4-89 years). To account for potentially non-linear trajectories, we used Generalized Additive Mixed Models (GAMMs) that enabled modelling the smooth left- (LH) and right hemisphere (RH) age-trajectories within our robust clusters (Methods).

In all clusters, SA asymmetry was strongly established already by age ~4 years, and the lifespan trajectories of both leftward (Fig. 2A) and rightward (Fig. 2B) SA asymmetries were largely parallel. Specifically, a large left-asymmetric region in and around SMG/perisylvian (#1; Fig. 2A) showed strong asymmetry by age ~4 that was maintained throughout life through steady aging-associated decline of both hemispheres, whereas leftward asymmetry of temporal cortex (#2,6) and anterior insular (#4) was maintained through developmental expansion and aging-associated decline of both hemispheres. Others (retrosplenial #5; mPFC #3,7) showed growth from pre-established asymmetry and more variable lifespan trajectories. On the other side, rightward asymmetries showed largely preserved asymmetry through aging-associated decline of both hemispheres (Fig 2B; medial occipital #1; lateral parietal #2; STS #5; orbitofrontal #7), through bilateral developmental expansion and aging-associated decline (mPFC #6), or steadily expanding bilateral SA until mid-life (cingulate; #3). Though asymmetry trajectories did show significant change at some point throughout life in most clusters (Supplementary file 1E), factor-smooth GAMM interaction analyses confirmed that asymmetry was significantly different from 0 across the entire lifespan in all SA clusters (Figure 3–figure supplements 1-2), and the average trajectories across all leftward and rightward clusters were clearly parallel (though still exhibited a significant difference; bordered plots in Fig. 2A-B; Supplementary file 1E).

In contrast, though homotopic trajectories of CT clusters were more variable, they were mostly characterized by developmental increase and aging-associated decrease in asymmetry (i.e. non-parallel lifespan trajectories), through unequal rates of continuous thinning between the hemispheres from age ~4 (Fig. 3A-B; see Figure 3–figure supplements 1-2). Specifically, leftward CT asymmetry developed through comparatively slower thinning trajectories of the LH, whereas rightward asymmetry developed through slower RH thinning. In general, asymmetry development was evident up to a peak around age ~25 for both leftward (Fig. 3A; superior frontal #2; precentral #4, frontal #8,9,10; calcarine #11) and rightward clusters (Fig 3B; #1-9) and declined thereafter (see also Figure 3–figure supplement 3). Factor-smooth GAMMs confirmed that the developmental foundation for CT asymmetry was already established by age ~4 (95% of CT clusters exhibited small but significant asymmetry at age ~4; Figure 3–figure supplement 2B), and again asymmetry trajectories showed significant change at some point throughout life (Supplementary file 1F). The average trajectories across all leftward and rightward clusters showed developmental asymmetry increase up to age ~25 and aging-associated asymmetry decrease from mid to old age (bordered plots; Fig 3). Results were robust to varying the number of knots used to estimate trajectories (Figure 2–figure supplement 1).

Figure 2: *Homotopic lifespan trajectories of SA in clusters exhibiting population-level **a)** leftward (yellow plots; yellow clusters) and **b)** rightward (pink plots; blue clusters) areal asymmetry (mm²). Larger plots on the left show the mean age trajectory across all clusters exhibiting leftward (top) and rightward (bottom) asymmetry. Note that the unit of measurement is the average surface area of a vertex within the cluster. Dark colours correspond to LH trajectories. All age trajectories were fitted using GAMMs. Data is residualized for sex, scanner and random subject intercepts. Clusters are numbered for reference.*

210
211

212
Figure 3: *Homotopic lifespan trajectories of CT in clusters exhibiting population-level **a)** leftward (yellow plots; yellow*
*clusters) and **b)** rightward (pink plots; blue clusters) thickness asymmetry (mm²). Larger plots on the left show the mean*
age trajectory across all clusters exhibiting leftward (top) and rightward (bottom) asymmetry. Note that the unit of
measurement is the average surface area of a vertex within the cluster. Dark colours correspond to LH trajectories. All
age trajectories were fitted using GAMMs. Data is residualized for sex, scanner and random subject intercepts. Clusters
are numbered for reference.
219
220
221
222
223 **2.3 Interregional asymmetry correlations**
224 We then investigated whether cortical asymmetry correlates within individuals (Methods).
225
For SA, a common covariance structure between asymmetries was detectable across datasets: LCBC, UKB and HCP
all correlated almost perfectly ($r >= 0.97$, all $p < 9.9^{-5}$; Fig. 4A; Figure 4–figure supplement 1A). The highest correlations
(or "hotspots") all reflected positive correlations between regions that are on average left-asymmetric and regions that
are on average right-asymmetric (i.e. higher leftward asymmetry in one region related to higher rightward asymmetry in
another; Fig 3A black outline); leftward asymmetry in SMG/perisylvian (#1L) was related to higher rightward asymmetry
in inferior parietal cortex (#2R; $r = .46$ [LCBC]), leftward anterior cingulate asymmetry (ACC; #3L) was related to higher

rightward asymmetry in mPFC (#6R, *r* = .46), and leftward asymmetry in a superior frontal cluster (#7L) was related to
rightward asymmetry in the cingulate (#3R, *r* = .67). None of the relationships could be explained by brain size, as
additionally removing the effect of intracranial volume (ICV) from cluster AI's had a negligible effect on their interrelations
(max correlation change = 0.003). Post-hoc tests confirmed that opposite-direction asymmetries were more correlated if
closer in cortex (Methods); geodesic distance was lower between cluster-pairs that were more correlated (*rho* = -.37, *p*
= .01 [LCBC]; -.38, *p* = .007 [UKB; Fig. 4C]; -.32, *p* = .02 [HCP]), though this was driven by the aforementioned "hotspots".
By contrast, same-direction SA asymmetries were not more correlated if closer in cortex (leftward [all *p* > .5]; rightward
[all *p* > .5]). This suggests specific SA asymmetries that are closer in cortex and opposite in direction may show
coordinated development.
241
For CT asymmetry, the correlation matrix exhibited a clear pattern in UKB that was less visible but still apparent in LCBC
and HCP (Fig. 4B; Figure 4–figure supplement 1B). Mantel tests confirmed that the covariance structure replicated
between all dataset-pairs (LCBC-UKB *r* = .43, *p* = .01; LCBC-HCP *r* = .45, *p* = .005; UKB-HCP *r* = .46, *p* = .01). The
observed pattern suggested higher leftward asymmetry in regions that are on average left-asymmetric was associated
with less rightward asymmetry in regions that are on average right-asymmetric. However, given that the AI measure is
bidirectional, closer inspection of the correlations revealed that higher leftward asymmetry in regions that are left-
asymmetric actually corresponded to more *leftward* asymmetry in right-asymmetric regions, and vice versa (and on
average; see Figure 4–figure supplement 2). In other words, individuals may tend towards either leftward lateralization
or rightward lateralization (or symmetry) on average, irrespective of the region-specific direction of mean asymmetry in
the cluster. Similarly, asymmetry in left-asymmetric regions was mostly positively correlated, and asymmetry in right-
asymmetric regions was mostly positively correlated. Again, additionally removing ICV-associated variance had
negligible effect (max correlation change = 0.007). Post-hoc principal components analysis (PCA) in UKB revealed PC1
explained 21.9% of the variance in CT asymmetry and strongly suggested a single global factor for CT asymmetry (Fig.
4D). Accordingly, we found a strong correlation between mean asymmetry across all leftward vs. mean asymmetry across
all rightward clusters in UKB (r = -.61, *p* < 2.2^{-16} [means weighted by cluster size]; see Fig. 4D; r = -.56, *p* < 2.2^{-16} [unweighted raw means];
r = .66, *p* < 2.2^{-16} [PC1 across all leftward vs. PC1 across all rightward]; r = -.56, *p* < 2.2^{-16} [spatial average across vertices]). Though less strong, all
relationships were significant in LCBC (r = -.10; *p* = 1.3^{-4} [weighted]; r = -.05; *p* = .03 [unweighted]; r = .17; *p* = 2.00^{-11} [PC1 vs. PC1];
r = -.05; *p* = .04 [spatial average]) and significant or trend-level in HCP (r = -.11; *p* = 1.6^{-4} [weighted]; r = -.04; *p* = .15 [unweighted]; r =
.13, *p* = 2.31^{-5} [PC1 vs. PC1]; r = -.05; *p* = .07 [spatial average]; see Figure 4–figure supplement 3). Opposite-direction CT
asymmetries that were closer in cortex were more negatively correlated in LCBC (*rho* = .29, *p* = .003) but not HCP (*p* =
.33) or UKB (*p* = .84), whereas CT asymmetry in left-asymmetric (*rho* = -.40, *p* = .003 [LCBC]; *rho* = -.44, *p* = 8.1^{-4} [UKB],
rho = -.28, *p* = .04 [HCP]) and right-asymmetric (*rho* = -.34, *p* = .04 [LCBC]; *rho* = -.48, *p* = .003 [UKB], *rho* = -.58, *p* =
2.0^{-4} [HCP]) regions was more positively correlated in cluster-pairs that were closer in cortex. These results suggest CT

asymmetry is globally interrelated across the cortex and shows high directional variability in the adult population.

Figure 4: *Interregional correlations between* **A)** *SA asymmetries and* **B)** *CT asymmetries for each replication dataset (AI's residualized for age, sex, scanner). AI's in rightward clusters are inversed, such that positive correlations denote positive asymmetry-asymmetry relationships regardless of direction. Yellow and blue brain clusters/colours denote leftward and rightward asymmetries, respectively (clusters numbered for reference). A consistent covariance structure was evident both for SA (r >= .97) and CT asymmetry (r >=.43; results above matrices). Black box in A highlights relationships between opposite-direction asymmetries (i.e. leftward vs rightward regions).* **C)** *For SA, opposite-direction cluster-pairs that were closer in cortex were more positively correlated (datapoints show cluster-pairs).* **D)** *A single component explained 21.9% variance in CT asymmetry in UKB (inset plot). Accordingly, we found a strong correlation (r = -.61; p < 10^{-16}) in UKB between mean asymmetry across leftward clusters (Y-axis) vs. mean asymmetry across rightward clusters (X-axis; AI's in rightward clusters inversed). Lines of symmetry (0) are in dotted grey (see also Figure 4–figure supplements 1-3).*

2.4 Heritability

Heritability of global AI measures was low and only significant for SA asymmetry in UKB (SA h^2_{SNP} = .07 [95% CI = .03 - .10]; *p* = 2.16e-4; CT h^2_{SNP} = .01 [-.02 - .05]; *p* = .22, Supplementary file 1G). For SA, only two clusters showed significant heritability in HCP and these did not survive multiple comparison correction. In contrast, SNP-based analyses revealed 71% (10/14) of SA asymmetry clusters exhibited significant heritability (Supplementary file 1H). Importantly, highest heritability was observed for leftward SA in the anterior insula cluster (h^2_{SNP} = 18.6%, *p* < 10^{-10}; see note in Supplementary file 1H), which was substantially higher than the next highest estimates in SMG/perisylvian (h^2_{SNP} = 10.7%, *p* = 3.01e-9), retrosplenial cortex, gyrus rectus and the cingulate (all h^2_{SNP} = 8-10%). For CT, no cluster survived in HCP, only 3/20 (15%) clusters exhibited significant SNP-heritability, and estimates were lower (h^2_{SNP} = 3-7%; Supplementary file 1I).

We then estimated cortex-wide heritability using a fine-grained parcellation [50] (Fig. 5). For SA, though no parcels survived correction in HCP, 53% (267/500) of parcels exhibited significant SNP-heritability (post-correction) in UKB (*p[FDR]*<.05; black outlines in Fig. 5A; parcels with suggestive significance in HCP [also in black outline] survived correction in UKB). Beyond significance, a consistent heritability pattern for SA asymmetry was clearly evident in both samples, notably in anterior insula, SMG, Sylvian fissure, STS, calcarine sulcus, cingulate, medial and orbitofrontal cortex, and fusiform (spatial correlation between maps; r = .38; *p* < 10^{-16}). Importantly, maximum SNP-heritability (yellow parcel in Fig. 5) was observed in anterior insula (parcel h^2_{SNP} = 16.4%; *p* < 10^{-10}), confirming this region constitutes the most heritable cortical asymmetry in humans (and not improving on the cluster-wise estimate). For CT, we observed little overlap in heritability estimates between datasets (spatial correlation was significant but low; r = .12; *p* = .01). Significant FDR-corrected SNP-heritability was observed in 11% (57/500) of parcels, including around superior temporal gyrus, planum temporale, the posterior insula/Sylvian fissure, anterior insula, and in orbitofrontal cortex (max h^2_{SNP} = 16.6%), along the cingulate and in medial visual cortex. However, SNP-heritability for CT asymmetry was substantially lower (β = -0.71, *p* < 2e-16), and higher estimates pertained to regions that were limited in extent but showed no clear global pattern.

For SA, large genetic correlations explained several phenotypic correlations in Fig. 4A (Fig. 5C): high genetic correlations were found between leftward asymmetry in SMG/perisylvian and higher rightward asymmetry in lateral parietal cortex (LPC; *rG* = .83; *p(FDR)* = 7.76e-05), between leftward superior frontal cortex asymmetry and rightward asymmetry along the cingulate (*rG* = .82; *p[FDR]* = 1.36e-02), and between leftward anterior temporal/parahippocampal asymmetry and rightward asymmetry in LPC (*rG* = .68; *p[FDR]* = 1.36e-02). Genetic correlations between anterior insula and two rightward superior frontal clusters were also observed (*rG* = .86; *p[FDR]* = 1.41e-06; rG = 0.42; *p[FDR]* = 7.74e-04) in the absence of phenotypic correlations (Fig. 4), and several same-direction asymmetries showed moderate genetic correlation. For CT, one cluster-pair survived FDR-correction (see Fig. 5C; *rG* = 0.68; *p[FDR]* = 3.03e-02).

2.5 Associations with Cognition, Handedness, Sex, and ICV

Several significant associations were observed between factors-of-interest and asymmetry in our clusters (Fig. 6; Supplementary file 1J-K). Notably, all effect sizes were small. For general cognitive ability, we found one association: higher SA asymmetry in the largest leftward cluster (SMG/perisylvian) was significantly associated with better cognition (β =.03 [CI = 0.02 – 0.04], p = 4.1e-7). This was checked in the substantially reduced non-imputed subset of data with no missing cognitive variables and retained the lowest p-value (N = 4696; β =0.04 [CI = 0.01 - 0.07]; p = 6.9e-3). For handedness, reduced leftward SA asymmetry in anterior insula and CT asymmetry along postcentral gyrus was found in left-handers, in line with our recent vertex-wise mapping in UKB [24]. For sex effects, which were also small, males typically exhibited slightly stronger SA asymmetry in large clusters (e.g. leftward SMG/perisylvian and temporal pole; rightward inferior parietal and superior frontal) but reduced leftward and rightward asymmetry in mPFC. For CT, males exhibited more rightward asymmetry in STS and posterior insula, more leftward CT asymmetry in superior frontal cortex, but reduced rightward CT asymmetry in entorhinal cortex and anterior insula, and reduced leftward asymmetry in caudal superior frontal cortex. As ICV effects were typically most nominal, these are shown in Figure 6–figure supplement 1.

Figure 6. *Associations with general cognitive ability (first principal component), Handedness, Sex, and estimated intracranial volume (ICV) in UKB, in clusters exhibiting population-level A) SA and B) CT asymmetry. Left plots denote significance (negative logarithm; corrected [p < 7.3e^{-5}] and uncorrected threshold [p = .01] shown by dotted and non-dotted line, respectively). X-axis displays the test for each cluster-association. As maximum sample size was used to test each association (Handedness, Sex and ICV: N=37,570), effects on cognition were tested in separate models with fewer observations (N = 35,199; separated association plots). Right plots denote effect sizes, 95% confidence intervals (error bars) and cortical location of associations surpassing Bonferroni-corrected significance. Right handers and females are coded 0, such that a negative effect for handedness / sex / ICV / cognition denotes less asymmetry in left handers / males / larger brains / higher cognition. Associations with ICV are shown in Figure 6–figure supplement 1. Yellow and blue clusters denote leftward and rightward asymmetries, respectively.*

3. Discussion

We provide a reference for population-level cortical asymmetries using 7 international datasets and offer the first description of the longitudinal lifespan trajectories of cortical asymmetry. Our results demonstrate the replicable interregional relationships between asymmetries within individuals, provide the most detailed heritability maps for cortical asymmetry to date, uncover novel and confirm previously-reported associations with factors reportedly related to asymmetry, and further knowledge on normal brain development. All maps are available at neurovault.org/XXXX.

Our vertex-wise description of cortical asymmetries that reproduce across cohorts replicates and completes a recent low-resolution meta-analysis [7], and can serve as a high-fidelity phenotype for future brain asymmetry studies. The marked consistency across samples here suggests consensus may now be reached regarding cortical asymmetry phenotypes in humans, as our results agree with most of the literature [7,8,14,19,23,26,28], including a recent large-scale mapping in mid-

old age [24]. This consensus, along with the genetic findings presented herein, suggests genetic-developmental programs
regulate mean brain lateralization with respect to SA, and the trajectories observed here suggest this form of cerebral
asymmetry is maintained throughout life and formed early on – likely *in utero* [26,31]. For CT asymmetry – for which findings
have been particularly mixed [7,8,18–20,10–17] – the left-right patterning observed here is compatible with recent reports [10,24],
studies examining CT asymmetry from birth [11], global meta-analyses [7], reports using alternative analysis streams [11,20],
anatomical asymmetries evident early in ontogeny [51,52], and leftward CT asymmetry overlapping language- [53] and motor-
related regions [54,55]. This consistency across adult samples may also indicate that mean CT asymmetry is genetically
regulated at the population-level in humans. However, our findings of development and decline of CT asymmetry across
life [10], higher directional variability in adult samples and lower heritability converge to suggest CT asymmetry may be
more prone to lifespan change, potentially more malleable to life experience, and susceptible to lifespan accumulation
of insult. Though it remains possible CT asymmetry change could be genetically regulated into old age, this interpretation
agrees with work suggesting SA may trace to prenatal factors [56,57] whereas CT relates more to postnatal lifespan
influences [57,58].
379
Conceivable sources of previously inconsistent results may be the age-distribution under study [10] and the existence of
varying directional asymmetries within the population (for CT asymmetry). This may partly explain why CT asymmetry
metrics are more variable across datasets compared with SA [7], though CT asymmetry effects are also smaller [8,14,19] and
likely contain more measurement error. Varying directional asymmetry within atlas-based parcels may also explain
inconsistent reports, such as in insular cortex where we observed consistent but discrepant asymmetry to that reported
in ENIGMA [7]. However, this does not account for the discrepancy that studies using the same atlas [48] typically report
areal asymmetry in STS to be left-lateralized [7,27,28], as the right-lateralization evidenced here and elsewhere [14,26] seems
unambiguous. And although we did not find strong SA asymmetry in inferior frontal regions as reported by Kong et al [7],
both the unthresholded significance maps and standard parcellation analyses were compatible with this (Figure 1–figure
supplement 2, Figure 1–figure supplement 5). The high overlap in effects between 7 datasets from 4 countries suggests
our results likely apply universally, though future studies will be needed to confirm this in non-American/North European
samples.
392
Our longitudinal description of cortical asymmetry lifespan trajectories gleaned novel insight into normal brain
development. For SA, adult-patterns of lateralization were strongly established already before ~4 years, indicating SA
asymmetry traces back further and does not primarily emerge through later cortical expansion [59]. Rather, the lifespan
trajectories predominantly show stability from childhood to old age, as SA asymmetry was generally maintained through
periods of developmental expansion and aging-associated change that were region-specific and bilateral. This agrees
with evidence indicating SA asymmetry is primarily determined *in utero* [26], and indirect evidence suggesting little change
in SA asymmetry from birth to 2 years despite rapid and concurrent cortical expansion [26,30,59]. It may also fit with the
principle that the primary microstructural basis of SA [60] – the number of and spacing between cortical minicolumns – is
determined in prenatal life [58,60], and agree with evidence suggesting asymmetry at this microstructural level may underly
hemispheric differences in SA [61]. The developmental trajectories agree with studies indicating SA asymmetry is
established and strongly directional early in life [26,28]. That anatomical change in later development specifically in SA
follows embryonic gene expression gradients may also agree with a prenatal account for SA asymmetry [58]. These results
may therefore constrain the extent to which SA asymmetry can be viewed as a plastic feature of brain organization, and
may even suggest SA asymmetry may sometimes be a marker for innate hemispheric specializations shared by most
humans. The high degree of precision with which leftward SA asymmetry follows the contours of auditory-related regions
in the Sylvian fissure (Figure 1–figure supplement 1) which show left functional lateralization in humans may be one
example [61–63].
410
In stark contrast, although weak CT asymmetry was evident by age 4, we observed considerable developmental growth
and lifespan change in CT asymmetry thereafter. Developmental trajectories showed non-linear asymmetry growth by
virtue of accelerated thinning of the non-dominant hemisphere, and led to maximally established asymmetry around ~25
years of age. These trajectories clearly suggest differentiation of the cortex is occurring with respect to CT asymmetry in
development, possibly, though not necessarily, suggesting CT asymmetry may be more amenable to experience-
dependent plastic change. Still, as cortical thinning in childhood is thought to partly reflect likely learning-dependent
processes such as intracortical myelination [64] and possibly pruning of initially overproduced synapses [65,66] and neuropil
reduction, CT asymmetry may reflect hemispheric differences in the developmental optimization of cortical networks at
least partly shaped by childhood experience. This raises the possibility CT asymmetry may be a marker of ontogenetic
hemispheric specialization within neurocognitive networks. Our findings in development agree with work finding a similar
left-right CT asymmetry pattern shows rapid asymmetry increase in the first years of life [11], with especially rapid increase
in leftward mPFC [11]. As we also observed rapid differentiation in mPFC that spanned across childhood and adolescence
(Fig. 3; Figure 3–figure supplement 3), we extend these earlier findings in neonates [11]. As prefrontal CT asymmetry
seems particularly vulnerable in neurodevelopmental disorders [21], aging, and Alzheimer's disease [10], these trajectories
may provide a useful normative reference. With regards to aging, most clusters exhibited the expected aging-associated
reduction of CT asymmetry we have previously shown is a feature of aging in heteromodal cortex [10]. The differentiation
and dedifferentiation of CT asymmetry at either end of life we show here underscores its proposed role in supporting
optimal brain organization and function.
429
430 For SA asymmetry, we uncovered a covariance structure that almost perfectly replicated across datasets. In general,
431 this fit with a multifaceted view [27,34,35], in which most asymmetries were either not or only weakly correlated, but reliably
432 so. However, we identified several regions wherein SA asymmetry reliably correlated within individuals, showing the

variance in cortical asymmetries is not always dissociable, as often thought [27,34,35]. The strongest relationships all pertained to asymmetries that were proximal in cortex but opposite in direction. Several of these were underpinned by high asymmetry-asymmetry genetic correlations, illustrating cerebral lateralizations in SA that are formed under common genetic-developmental influence, and in agreement with likely prenatal origins for SA asymmetry [26,60].

For CT asymmetry, we also uncovered a common covariance structure – particularly clear in UKB – that nevertheless replicated with moderate precision across datasets. Furthermore, a single global factor explained a relatively high proportion of variance in CT asymmetry in UKB, and a strong correlation across 38,172 individuals further suggested CT asymmetry is globally interrelated across the cortex (Fig. 4D). These data for CT indicate individuals tend towards either leftward asymmetry, rightward asymmetry, or symmetry, both globally across the cortex and irrespective of the region-specific average direction of asymmetry (Figure 4–figure supplements 2-3). This result seems in broad agreement with the notion that some lateralized genetic-developmental programs may trigger lateralization in either direction [35] or lose their directional bias through environmental interaction [35]. As CT asymmetry seems established at but minimal from birth [11], genetic effects may determine the average region-specific hemispheric bias in the population, but developmental change may subsequently confer major increases upon its directional variance [35]. Overall, the evidence converges to suggest a high degree of developmental change may shape CT asymmetry and lead to higher directional variability in the population. Thus, far from being independent phenotypes [27,34], CT asymmetries may be globally interrelated across the cortex and their direction coordinated through development.

For SA asymmetry, we found replicable patterns of moderate heritability across datasets and across twin and genomic methods. We also found SA asymmetry in the anterior insula is, to our knowledge, the most heritable brain or behavioural asymmetry yet reported with genomic methods [31,67,68], with common SNPs explaining ~19% variance. This is a substantial improvement on our recent report of < 5% [24], and illustrates a benefit of our data-driven population-mapping approach. As we reported recently [24], we confirm asymmetry in this region associates with handedness (see below). Furthermore, highest SNP-heritability for SA was found in all regions that constitute the earliest emerging cortical asymmetries *in utero* [51,69–71]: anterior insula, STS, PT, medial occipital cortex, and parahippocampal gyrus (Fig. 6A). Still, we found most SA asymmetries exhibited significant – albeit often lower – heritability, as did most parcels when estimated cortex-wide, and significant heritability was also evident in regions not found in the present analyses to show strong SA asymmetry, such as Broca's area (but see Figure 1–figure supplement 5). These effects agree with and elaborate on two genetic explorations using atlas-based methods [7,31] and reports of heritable SA asymmetry in handedness-associated clusters [24]. By contrast, CT asymmetry was generally not heritable, or showed low and localized heritability effects. We also observed divergent results using twin and genomic methods for CT, possibly due to low-power for twin-models, though we note the SNP-based effects we observed were somewhat in agreement with a previous twin study [7]. Overall, these reproducible results can guide phenotypic selection in future genomic and biological studies on cerebral laterality.

Considered together, lifespan stability possibly from birth [26], less interindividual directional variability, higher heritability, and phenotypic and genetic correlations all converge to suggest comparatively higher genetic influence upon SA asymmetry and possibly limited plasticity. This agrees with work showing genetic variants associated with (mostly SA) asymmetry are primarily expressed in prenatal life [31]. By contrast, developmental change, high interindividual directional variability and low heritability for CT asymmetry may fit a scenario whereby CT asymmetry may be more responsive to postnatal individual exposures [58], or driven by random developmental influences [72]. Whether region-specific CT asymmetry-change relates to the maturation of lateralized brain functions [72,73] will be an important question for future research. Regardless, our results support a relative prenatal-postnatal developmental dichotomy for SA and CT asymmetry.

Screening asymmetries for association with cognition revealed one region – SMG/perisylvian – wherein higher leftward asymmetry related to higher cognition. Across all samples tested, this cluster was consistently the most lateralized, with ~95% directional concordance (Figure 1–figure supplement 3), suggesting highly regulated genetic-developmental programs shape its laterality in humans. Asymmetry here is likely related to brain torque [25,40], a gross anatomical twist of the hemispheres leading to interhemispheric anatomical differences especially around the Sylvian fissure [25]. Given that brain torque also represents a population-level norm [74], this result suggests disruptions in prenatal cerebral lateralization may lead to cognitive deficits detectable in later life, and agrees with recent work suggesting brain torque may be related to cognitive outcomes [41,74]. That this was found specifically in the most lateralized SA cluster may agree with work suggesting general cognitive abilities that show inter-individual stability across life [75,76] relate primarily to SA phenotypes that depend mostly on prenatal factors [56,58].

Consistent with our recent vertex-wise analysis in UKB [24], we confirmed leftward SA asymmetry of the anterior insula, and leftward CT asymmetry of somatosensory cortex, is subtly reduced in left handers. Sha et al. [24] reported shared genetic influences upon handedness and cortical asymmetry in anterior insula and other more focal regions not identified with the approach used here. Anterior insula lies within a left-lateralized functional language network [77], and its structural asymmetry may relate to language lateralization [33,78,79] in which left-handers show increased incidence of atypicality [45,80,81]. Together with observations that its asymmetry emerges early *in utero* [69], we agree with others [33] that future research will find this ontogenetically foundational region of cortex [82,83] a fruitful line of inquiry for understanding genetic-developmental mechanisms influencing laterality phenotypes. Leftward CT asymmetry reduction in somatosensory cortex in left handers also echoes our recent report, where it was suggested to reflect plastic adaptation to an already-established hand preference [24]. We extend these results by showing CT asymmetry both postcentrally and in general shows developmental differentiation and lifespan change. Given this region overlaps with functional representations of

the hands [54,55,84] – as in Sha et al. [24] – and our approach also detected no significant heritability, these findings may also fit a scenario whereby CT asymmetries are amenable to alteration through use-dependent plasticity and possibly carry information regarding group-level hemispheric specializations of function. However, the small effects cast doubt on the utility of cortical asymmetry to predict individual hand preference.

Asymmetry-relationships with other factors were often compatible with those reported in the ENIGMA meta-analysis [7]. Concerning sex effects – which were small and differing in direction – we similarly observed leftward SA asymmetry in temporal and SMG/perisylvian regions to be larger in males [7], replicating earlier findings [32]. Previous genetic analyses imply steroid-hormone pathways underly this difference [32], and sex in general was found to be more predictive than ICV both here and elsewhere [32]. We also found lower SA asymmetry in medial prefrontal cortex in males that was compatible with this earlier report [7]. Inconsistencies evident between ours and the ENIGMA report include findings of increased (here) and decreased [7] lateral parietal SA asymmetry in males, and increased [7] and decreased (here) entorhinal CT asymmetry in males, and our approach detected other regions slightly more asymmetric in males (e.g. STS). Possibly, differences in sample median age (here UKB = ~64; Kong et al. = 26 [7]) and potential sex-differences in age decline trajectories [85] may underlie some inconsistencies, possibly moreso for CT measures in structures vulnerable to age-related degeneration [24].

Several limitations should be mentioned. First, our delineation of population-level asymmetry used a single analysis software. As with most current papers, we used FreeSurfer's default 'recon-all' function to delineate the cortex, which has been extensively validated against postmortem measurements [86] and is the software underlying most large-scale studies involving brain measures. It is currently unclear to what extent differences in pipelines account for previous mixed results [7,8,18–20,10–17]. Although we highlight there are clear commonalities between our results and studies using alternative pipelines [11,12,20], which may suggest our results generalize across analysis systems [11,12,20,23], we found one instance where the MRI pipeline leads to different results for CT asymmetry (Figure 1–figure supplement 6). It is not known what underlies this difference, though we find it is unrelated to the cross-hemispheric registration methods employed here, as our results reproduce using standard parcellation methods and thus are likely evident in most FreeSurfer-derived datasets (Figure 1–figure supplement 5). One possibility could be that thickness asymmetry may not reflect cortical thickness differences per se, but rather reflect biologically meaningful hemispheric differences in intracortical myelination levels that are consistently picked up on via FreeSurfer's standard delineation procedure. That the anterior-posterior CT asymmetry pattern shows a clear developmental trajectory also suggests it is a true biological effect (Figure 3; Figure 3–figure supplements 1-3). Relatedly, while the reported SA and CT asymmetry patterns and strengths using cross-hemispheric methods agree with standard analysis (Figure 1–figure supplement 5), it is possible the magnitude of some specific asymmetries near the boundary of the subcortex may be exaggerated via this approach [24]. Second, while GAMMs are considered an optimal modelling technique for longitudinal lifespan data and are robust to non-normal age distributions [87], relative underrepresentation of the mid-adulthood age-range may drive trajectory inflection points around this age [10], suggesting caution is warranted regarding interpreting mid-life inflection points as reflecting real change. Third, though the differing heritability methods applied enabled replication for SA, twin studies are prone to overestimating heritability due to unmet assumptions [88], whereas SNP-based methods may not capture all phenotype-relevant genetic variance and have their own assumptions [89]. Indeed, we found twin-based estimates were often substantially higher even where only nominally significant, agreeing with recent calls for caution when interpreting twin-based heritability estimates [88]. Fourth, we imposed a necessary cluster size limit for overlapping asymmetry effects across samples, and thus more focal asymmetries may also be informative in relation to the factors tested here [24]. Fifth, as only dichotomous handedness self-reports are available with UKB, future studies might benefit from incorporating more nuanced handedness assessments not currently available in data of this size. Relatedly, because UKB cognitive data is not exhaustive (e.g. fluid IQ ranges from 1-13), we extracted the common variance across core tests to index general cognitive ability. This approach did not permit testing associations with specific cognitive abilities, which may be highly informative in the context of asymmetry, particularly in the case of lateralized cognition [90].

Overall, we provide an openly-available comprehensive characterization of asymmetry in the cerebral cortex including longitudinal lifespan changes, heritability, and individual differences that bears enough reproducibility to be used as a standard in future research.

4. Methods
4.1 Samples
We used anatomical T$_1$-weighted (T$_{1w}$) scans from 7 independent international MRI datasets originating from 4 countries (see Supplementary file 1A for an overview of samples used for each analysis). Note that with the exception of vertex-wise analyses in UKB (see below), all analyses made use of all available observations from each sample meeting the stated criteria for each analysis (e.g. age-range).

4.1.1 Reproducibility across samples
To delineate average adult patterns of whole-cortical SA and CT asymmetry, we restricted the age-range of all samples used in the vertex-wise analyses to 18-55. **Dataset 1:** Here, the Center for Lifespan Changes in Brain and Cognition (*LCBC*) sample comprised 1572 mixed cross-sectional and longitudinal scans (N longitudinal = 812; timepoint range = 1-6) from 923 participants (mean age = 30.6 ± 9.6) collected across 2 scanners. Additionally, 125 individuals were double-scanned at the same timepoint on both scanners. **Dataset 2:** The Cambridge Centre for Ageing and Neuroscience (*Cam-CAN*) [91] sample comprised cross-sectional scans of 321 individuals (mean age = 38.7 ± 9.7) [92]. **Dataset 3:** The Dallas

Lifespan Brain Study (*DLBS*) [93] sample comprised cross-sectional scans of 160 individuals (mean age = 37.5 ± 10.7). **Dataset 4:** The Southwest University Adult Lifespan Dataset (*SALD*) [94] sample comprised cross-sectional scans of 301 individuals (mean age = 33.7 ± 11.5). **Dataset 5:** The *IXI* sample comprised cross-sectional scans of 313 healthy individuals collected across 3 scanners (mean age = 36.8 ± 9.6; http://brain-development.org/ixi-dataset). **Dataset 6:** Here, the Human Connectome Project (*HCP*) 1200 [95] sample comprised 1111 scans (mean age = 28.8 ± 3.7). **Dataset 7:** Here, the UKB sample consisted of 1000 randomly sampled cross-sectional scans (mean age = 52.1 ± 1.9), restricted to be comparable in size to the other datasets in this analysis.

4.1.2 Lifespan trajectories

Here, we used the full age-range of LCBC (4.1 - 89.4 years), with a sample comprising 3937 cross-sectional and longitudinal scans (N longitudinal = 2762) from 1886 individuals (females = 1139; mean age = 36.8) collected across 4 scanners (271 double-scans) [96,97].

4.1.3 Interregional correlations

Here, we used the three largest datasets; LCBC (N = 923; N obs = 1572), UKB (N = 38,172), and HCP (N = 1109; two outliers removed; see 4.3.3).

4.1.4 Heritability and individual differences

For twin heritability, we used HCP 1200 extended twin data (1037 scans from twins and non-twin siblings; age-range = 22-37; mean age = 28.9 ± 3.7). The various kinships are described in Supplementary file 1B. All included twin pairs were same-sex. For SNP-heritability, we used the UKB imaging sample with genome-wide data (N = 31,433; see 4.3.4). For individual differences analyses, we used the UKB imaging sample with the maximum number of available observations for each variable-of-interest (see 4.3.5).

4.2. MRI preprocessing

T1w anatomical images (see Supplementary file 1C for MRI acquisition parameters) were processed with FreeSurfer (v6.0.0) [98] and vertex-wise SA and CT morphometry estimates were obtained for each MRI observation. As the LCBC sample also contained longitudinal observations, initial cross-sectional reconstructions in LCBC were subsequently ran through FreeSurfer's longitudinal pipeline. As HCP data was acquired at a higher voxel resolution (0.7mm isotropic), the T1w scans were processed with the --hires flag to recon-all [99]. SA and CT maps of the LH and RH of each participant in each dataset were resampled from the native cortical geometry to a symmetrical surface template ("*LH_sym*") [14,100] based on cross-hemispheric registration [101]. This procedure achieves vertex-wise alignment of the data from each participant and homotopic hemisphere in a common analysis space. SA values were resampled with an additional Jacobian correction to ensure preservation of the areal quantities [102]. We then applied an 8mm FWHM Gaussian kernel to surface-smooth the LH and RH data.

4.3 Data analysis

All analyses were performed in FreeSurfer (v6.0) and R (v4.1.1).

4.3.1 Reproducibility across samples: population-level asymmetry

We assessed SA and CT asymmetry vertex-wise using FreeSurfer's Linear Mixed Effects (LME) tool [103]. Asymmetry was delineated via the main effect of Hemisphere (controlling for Age, Age × Hemisphere, Sex, Scanner [where applicable], with a random subject term). For each sample and metric, we computed mean Asymmetry Index maps (AI; defined as (LH-RH) / ((LH+RH)/2)). Spatial overlap of AI maps across datasets was quantified using Pearson's *r*. To delineate regions exhibiting robust SA and CT asymmetry across datasets, we thresholded and binarized the AI maps by a given absolute effect size (SA = 5%; CT = 1%; achieving *p*[FDR] < .001 in most datasets with FreeSurfer's 2-stage FDR-procedure [103]), and summed the binary maps. After removing the smallest clusters (<200 mm^2), a set of robust clusters was defined as those exhibiting overlapping effects in 6 out of 7 samples. We then extracted SA and CT data in symmetrical space for each cluster, subject, and hemisphere, spatially averaging across vertices.

4.3.2 Lifespan trajectories

Factor-smooth GAMMs ("gamm4" [104]) were used to fit a smooth Age trajectory per Hemisphere, and assess the smooth Age × Hemisphere interaction in our clusters. The linear predictor matrix of the GAMM was used to obtain asymmetry trajectories and their confidence intervals, computed as the difference between zero-centered (i.e. demeaned) hemispheric age-trajectories. We included Hemisphere as an additional fixed effect, sex and scanner as covariates-of-no-interest, and a random subject intercept. A low number of basis dimensions for each smoothing spline was chosen to guard against overfitting (knots = 6; see Figure 2–figure supplement 1). Here, outliers falling > 6SD from the trajectory of either hemisphere were removed on a region-wise basis (Supplementary file 1E-F).

4.3.3 Interregional correlations

We assessed covariance between asymmetries, separately for SA and CT. Here, we regressed out age, sex and scanner (where applicable) from each AI, and obtained a cluster-cluster correlation matrix. Individual AI's in clusters with rightward mean asymmetry were inversed, such that positive correlations denote asymmetry-asymmetry relationships regardless of the direction of mean asymmetry in the cluster. At this point, two outliers in HCP data were detected and discarded for this and all subsequent analyses (Figure 4–figure supplement 4). Replication was assessed using the Mantel test ("ade4" [105]) between each dataset-pair (LCBC, UKB, HCP) across 10,000 permutations. We then post-hoc tested whether covariance between asymmetries was related to proximity in cortex, obtaining the average geodesic distance between

all clusters along the ipsilateral surface ("SurfDist" Python package [106]), and correlating pair-wise distance with pair-wise correlation coefficient (Fisher's transformed coefficients; Spearman's correlation). To post-hoc test whether observed covariance patterns for CT reflected a global factor, we ran a PCA across z-transformed AI's for all CT clusters (pre-corrected for the same covariates). Based on the results, we computed the mean AIs across all leftward and across all rightward clusters and tested the partial correlation between mean leftward CT asymmetry in left-asymmetric clusters and mean rightward CT asymmetry in right-asymmetric clusters, in each of the three cohorts.

4.3.4 Heritability
Heritability of SA and CT asymmetry was assessed using both twin- and SNP-based methods, both for our set of robust clusters and cortex-wide across 500 parcels [50]. For cluster analyses, significance was considered at Bonferroni-corrected $p<.05$ applied separately across each metric. Cortex-wide significance was considered at $p(FDR) < .05$ (500 tests per map). Twin heritability was assessed using ACE models in "OpenMx" [107]. Using observed cross-twin and cross-sibling covariance, the ACE model decomposes the proportion of observed phenotypic variance into additive genetic effects [A], shared environmental effects [C], and unique environmental effects and/or error [E]. Data were reformatted such that rows represented family-wise observations. As is standard, we set A to be 1 for MZ twins assumed to share 100% of their segregating genes (but see [108]), 0.5 for DZ twins and siblings that share 50% on average, and shared environment was assumed equal for twins and non-twin siblings (but see [88,109]). For each phenotype we first regressed out age and sex and computed z-scores. Statistical significance was assessed by comparing ACE model fit to submodels with the parameter-of-interest removed.

For SNP-heritability, the final genetic sample consisted of 31,433 UKB participants (application #32048) with imaging and quality checked genetic data. We removed subjects that were outliers based on heterozygosity [field 22027] and missingness (> 0.05), mismatched genetic and reported sex [22001], sex chromosome aneuploidies [22019], and those not in the "white British ancestry" subset [22006] [110]. At variant level, after removing SNPs with minor allele frequency < 0.01, data from 654,584 autosomal SNPs were used to compute a genetic relationship matrix using GCTA (v1.93.2) [111]. For each phenotype, we first regressed out age and sex and computed z-scores. Genome-based restricted maximum likelihood (GREML) methods as implemented in GCTA were then used to compute SNP-heritability for each AI measure, applying a kinship coefficient cut-off of 0.025 (excluding one individual from each pair), and controlling for genetic population structure (first ten principal components). Bivariate GREML analysis was used to test genetic correlations between asymmetry clusters [111], with relationships tested only for cluster-pairs where both clusters exhibited significant SNP-heritability (p < .05; pre-corrected; 78 tests for SA, 48 for CT). Significance of genetic correlations was assessed at $p(FDR) <.05$.

4.3.5 Associations with Cognition, Sex, Handedness, & ICV
Finally, we assessed relationships between asymmetry in our robust clusters and cognitive ability, handedness, sex and ICV. For cognition, we used the first principal component across the following 11 core UK Biobank cognitive variables [112]: Mean reaction time (log transformed) [field 20023], Numeric memory [4282], Fluid reasoning [20016], Matrix completion [6373], Tower rearranging [21004], Symbol digit substitution [23324], Paired associate learning [20197], Prospective memory [20018], Pairs matching (log) [399], Trail making A (log) [6348], Trail making B (log) [6350]. Prior to the PCA, for participants with cognitive data, data was imputed for missing cognitive variables via the "imputePCA" R function (number of estimated components tentatively optimized using general cross validation; "missMDA" Package [113]). PC1 (explaining 39.2%; Supplementary file 1L) was inversed to correlate negatively with age (r = -.39), ensuring higher values reflected higher cognition. As fewer participants had cognitive data relative to the other variables, for each cluster we ran one set of linear models to assess the marginal effect of cognition (PC1 as predictor; age, sex, ICV controlled; N = 35,199), and one set of linear models to assess the marginal effects of Handedness, Sex, and ICV in a model including all three predictors (age controlled, N = 37,570 with available handedness data). For the cognitive analysis, effects identified in the imputed dataset were checked against the confidence intervals for the effect in the subset of the data with no missing cognitive variables (N = 4696). Participants who self-reported as mixed handed were not included as this can be unreliable over repeat time-points [24]. Significance was considered at Bonferroni-corrected $\alpha = p < 7.3^{-5}$ (.01/136 [34 clusters × 4]).

Data sharing/availability
All summary-level maps are available at neurovault.org/XXXX (upon acceptance). All code underlying the main analyses is available at https://github.com/jamesmroe/PopAsym and on the Open Science Framework (upon acceptance). All derived source data underlying all figures is also available here and in Supplementary files 2-3. All datasets used in this work are openly available, with the exception of LCBC, where participants, which include many children, have not consented to share their data publicly online. Other datasets used in this work are available without restrictions and are not subject to application approval (DLBS; https://fcon_1000.projects.nitrc.org/indi/retro/dlbs.html; CC BY-NC; SALD; http://fcon_1000.projects.nitrc.org/indi/retro/sald.html; CC BY-NC; IXI; https://brain-development.org/ixi-dataset; CC BY-SA 3.0). Accordingly, we have made the individual-level data for these samples available and our code can be used to reproduce vertex-wise analyses in these samples. Individual-level data for the remaining samples (LCBC; Cam-CAN, HCP; UKB) may be available upon reasonable request, given appropriate ethical, data protection, and data-sharing agreements where applicable. Requests must be submitted and approved via the relevant channel (details are provided in Supplementary File 2).

Acknowledgements

Scripts were ran on the Colossus processing cluster at the University of Oslo, and on resources provided by UNINETT Sigma2 (NN9769K). LCBC funding: European Research Council under grants 283634, 725025 (to A.M.F.), and 313440 (to K.B.W.); Norwegian Research Council (to A.M.F. and K.B.W.) under grants 249931 (TOPPFORSK) and 302854 (FRIPRO; to Y.W.), The National Association for Public Health's dementia research program, Norway (to A.M.F). Data used in the preparation of this work were obtained from the MGH-USC Human Connectome Project (https://ida.loni.usc.edu/login.jsp). Data used in this work was also provided by the Cambridge Centre for Ageing and Neuroscience (CamCAN). This research has been conducted using the UK Biobank Resource.

4.9 References

1. van Kesteren, E.-J. & Kievit, R. A. Exploratory factor analysis with structured residuals for brain network data. *Netw. Neurosci.* 1–45 (2020). doi:10.1162/netn_a_00162

2. Stark, D. E. *et al.* Regional Variation in Interhemispheric Coordination of Intrinsic Hemodynamic Fluctuations. *J. Neurosci.* **28**, 13754–13764 (2008).

3. Schmitt, J. E., Giedd, J. N., Raznahan, A. & Neale, M. C. The genetic contributions to maturational coupling in the human cerebrum: A longitudinal pediatric twin imaging study. *Cereb. Cortex* **28**, 3184–3191 (2018).

4. Chen, C. H. *et al.* Genetic topography of brain morphology. *Proc. Natl. Acad. Sci. U. S. A.* **110**, 17089–17094 (2013).

5. Eyler, L. T. *et al.* Conceptual and Data-based Investigation of Genetic Influences and Brain Asymmetry: A Twin Study of Multiple Structural Phenotypes. *J. Cogn. Neurosci.* **26**, 1100–1117 (2014).

6. Raznahan, A. *et al.* Patterns of coordinated anatomical change in human cortical development: A longitudinal neuroimaging study of maturational coupling. *Neuron* **72**, 873–884 (2011).

7. Kong, X.-Z. *et al.* Mapping cortical brain asymmetry in 17,141 healthy individuals worldwide via the ENIGMA Consortium. *Proc. Natl. Acad. Sci.* **115**, E5154–E5163 (2018).

8. Chiarello, C., Vazquez, D., Felton, A. & McDowell, A. Structural asymmetry of the human cerebral cortex: Regional and between-subject variability of surface area, cortical thickness, and local gyrification. *Neuropsychologia* **93**, 365–379 (2016).

9. Meyer, M., Liem, F., Hirsiger, S., Jäncke, L. & Hänggi, J. Cortical surface area and cortical thickness demonstrate differential structural asymmetry in auditory-related areas of the human cortex. *Cereb. Cortex* **24**, 2541–2552 (2014).

10. Roe, J. M. *et al.* Asymmetric thinning of the cerebral cortex across the adult lifespan is accelerated in Alzheimer's disease. *Nat. Commun.* **12**, 1–11 (2021).

11. Li, G., Lin, W., Gilmore, J. H. & Shen, D. Spatial patterns, longitudinal development, and hemispheric asymmetries of cortical thickness in infants from birth to 2 years of age. *J. Neurosci.* **35**, 9150–9162 (2015).

12. Luders, E. *et al.* Hemispheric asymmetries in cortical thickness. *Cereb. Cortex* **16**, 1232–1238 (2006).

13. Zhou, D., Lebel, C., Evans, A. & Beaulieu, C. Cortical thickness asymmetry from childhood to older adulthood. *Neuroimage* **83**, 66–74 (2013).

14. Maingault, S., Tzourio-Mazoyer, N., Mazoyer, B. & Crivello, F. Regional correlations between cortical thickness and surface area asymmetries: A surface-based morphometry study of 250 adults. *Neuropsychologia* **93**, 350–364 (2015).

15. Shaw, P., Lalonde, F. & Al, L. C. et. Development of Cortical Asymmetry in Typically Developing Children and Its Disruption in Attention-Deficit/Hyperactivity Disorder. **66**, 888–896 (2009).

16. Lou, Y. *et al.* Brain asymmetry differences between Chinese and Caucasian populations : a surface-based morphometric comparison study. (2019).

17. Hamilton, L. S. *et al.* Asymmetries of cortical thickness: Effects of handedness, sex, and schizophrenia. *Neuroreport* **18**, 1427–1431 (2007).

18. Zhou, D. *et al.* Preserved cortical asymmetry despite thinner cortex in children and adolescents with prenatal alcohol exposure and associated conditions. *Hum. Brain Mapp.* **39**, 72–88 (2018).

19. Koelkebeck, K. *et al.* The contribution of cortical thickness and surface area to gray matter asymmetries in the healthy human brain. *Hum. Brain Mapp.* **35**, 6011–6022 (2014).

20. Plessen, K. J., Hugdahl, K., Bansal, R., Hao, X. & Peterson, B. S. Sex, age, and cognitive correlates of asymmetries in thickness of the cortical mantle across the life span. *J Neurosci* **34**, 6294–6302 (2014).

21. Postema, M. C. *et al.* Altered structural brain asymmetry in autism spectrum disorder in a study of 54 datasets. *Nat. Commun.* **10**, 1–12 (2019).

22. Thompson, P. M. *et al.* Tracking Alzheimer's disease. *Ann. N. Y. Acad. Sci.* **1097**, 183–214 (2007).

23. Lyttelton, O. C. *et al.* Positional and surface area asymmetry of the human cerebral cortex. *Neuroimage* **46**, 895–903 (2009).

24. Sha, Z. *et al.* Handedness and its genetic influences are associated with structural asymmetries of the cerebral cortex in 31,864 individuals. *Proc. Natl. Acad. Sci.* **118**, 1–9 (2021).

25. Toga, a W. & Thompson, P. M. Mapping brain asymmetry. *Nat. Rev. Neurosci.* **4**, 37–48 (2003).

26. Li, G. *et al.* Mapping longitudinal hemispheric structural asymmetries of the human cerebral cortex from birth to 2 years of age. *Cereb. Cortex* **24**, 1289–1300 (2014).

27. Bain, J. S., Filo, S. & Mezer, A. A. The robust and independent nature of structural STS asymmetries. *Brain Struct. Funct.* **224**, 3171–3182 (2019).

28. Remer, J. *et al.* Quantifying cortical development in typically developing toddlers and young children, 1–6 years of age. *Neuroimage* **153**, 246–261 (2017).

29. Nie, J., Li, G. & Shen, D. Development of cortical anatomical properties from early childhood to early adulthood. *Neuroimage* **76**, 216–224 (2013).

30. Li, G. *et al.* Mapping region-specific longitudinal cortical surface expansion from birth to 2 years of age. *Cereb. Cortex* **23**, 2724–2733 (2013).

31. Sha, Z. *et al.* The genetic architecture of structural left–right asymmetry of the human brain. *Nat. Hum. Behav.* (2021). doi:10.1038/s41562-021-01069-w

32. Guadalupe, T. *et al.* Asymmetry within and around the human planum temporale is sexually dimorphic and influenced by genes involved in steroid hormone receptor activity. *Cortex* **62**, 41–55 (2015).

33. Chiarello, C., Vazquez, D., Felton, A. & Leonard, C. M. Structural asymmetry of anterior insula: Behavioral correlates and individual differences. *Brain Lang.* **126**, 109–122 (2013).

34. Rentería, M. E. Cerebral asymmetry: A quantitative, multifactorial, and plastic brain phenotype. *Twin Res. Hum. Genet.* **15**, 401–413 (2012).

35. Francks, C. Exploring human brain lateralization with molecular genetics and genomics. *Ann. N. Y. Acad. Sci.* **1359**, 1–13 (2015).

36. Liu, H., Stufflebeam, S. M., Sepulcre, J., Hedden, T. & Buckner, R. L. Evidence from intrinsic activity that asymmetry of the human brain is controlled by multiple factors. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 20499–20503 (2009).

37. Crow, T. J., Crow, L. R., Done, D. J. & Leask, S. Relative hand skill predicts academic ability: Global deficits at the point of hemispheric indecision. *Neuropsychologia* **36**, 1275–1282 (1998).

38. Ocklenburg, S., Güntürkün, O., Hugdahl, K. & Hirnstein, M. Laterality and mental disorders in the postgenomic age – A closer look at schizophrenia and language lateralization. *Neurosci. Biobehav. Rev.* **59**, 100–110 (2015).

39. Kong, X.-Z. *et al.* Large-Scale Phenomic and Genomic Analysis of Brain Asymmetrical Skew. *Cereb. Cortex* 1–18 (2021). doi:10.1093/cercor/bhab075

40. LeMay, M. Morphological Cerebral Asymmetries of Modern Man, Fossil Man, and Nonhuman Primate. *Ann. N. Y. Acad. Sci.* **280**, 349–366 (1976).

41. Zhao, L., Matloff, W., Shi, Y., Cabeen, R. P. & Toga, A. W. Mapping Complex Brain Torque Components and Their Genetic and Phenomic Architecture in 24 , 112 healthy individuals. (2021).

42. Moodie, J. E. *et al.* Fluctuating asymmetry in brain structure and general intelligence in 73-year-olds. *Intelligence* **78**, 101407 (2020).

43. Yeo, R. A., Ryman, S. G., Pommy, J., Thoma, R. J. & Jung, R. E. General cognitive ability and fluctuating asymmetry of brain surface area. *Intelligence* **56**, 93–98 (2016).

44. Kong, X. *et al.* Mapping brain asymmetry in health and disease through the ENIGMA consortium. *Hum. Brain Mapp.* 1–15 (2020).

45. Wiberg, A. *et al.* Handedness , language areas and neuropsychiatric diseases : insights from brain imaging and genetics. *Brain* 1–10 (2019). doi:10.1093/brain/awz257

46. van der Meer, D. *et al.* Quantifying the Polygenic Architecture of the Human Cerebral Cortex: Extensive Genetic Overlap between Cortical Thickness and Surface Area. *Cereb. Cortex* **30**, 5597–5603 (2020).

47. Eyler, L. T. *et al.* A comparison of heritability maps of cortical surface area and thickness and the influence of adjustment for whole brain measures: A magnetic resonance imaging twin study. *Twin Res. Hum. Genet.* **15**, 304–314 (2012).

48. Desikan, R. S. *et al.* An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *Neuroimage* **31**, 968–80 (2006).

49. Miller, K. L. *et al.* Multimodal population brain imaging in the UK Biobank prospective epidemiological study. *Nat. Neurosci.* **19**, 1523–1536 (2016).

50. Schaefer, A. *et al.* Local-Global Parcellation of the Human Cerebral Cortex from Intrinsic Functional Connectivity MRI. *Cereb. Cortex* 1–20 (2017). doi:10.1093/cercor/bhx179

51. Habas, P. A. *et al.* Early folding patterns and asymmetries of the normal human brain detected from in utero MRI. *Cereb. Cortex* **22**, 13–25 (2012).

52. Glasel, H. *et al.* A robust cerebral asymmetry in the infant brain: The rightward superior temporal sulcus. *Neuroimage* **58**, 716–723 (2011).

53. Foundas, A. L., Eure, K. F., Luevano, L. F. & Weinberger, D. R. MRI asymmetries of Broca's area: The pars triangularis and pars opercularis. *Brain Lang.* **64**, 282–296 (1998).

54. Penfield, W. & Boldrey, E. Somatic Motor and Sensory Representation in Man. *Brain* 389–443 (1937).

55. Roux, F. E., Djidjeli, I. & Durand, J. B. Functional architecture of the somatosensory homunculus detected by electrostimulation. *J. Physiol.* **596**, 941–956 (2018).

56. Walhovd, K. B. *et al.* Neurodevelopmental origins of lifespan changes in brain and cognition. *Proc. Natl. Acad. Sci. U. S. A.* **113**, 9357–9362 (2016).

57. Grasby, K. L. *et al.* The genetic architecture of the human cerebral cortex. *Science (80-.).* **367**, (2020).

58. Fjell, A. M. *et al.* Continuity and Discontinuity in Human Cortical Development and Change From Embryonic Stages to Old Age. *Cereb. Cortex* **29**, 3879–3890 (2019).

59. Wierenga, L. M., Langen, M., Oranje, B. & Durston, S. Unique developmental trajectories of cortical thickness and surface area. *Neuroimage* **87**, 120–126 (2014).

60. Rakic, P. A small step for the cell, a giant leap for mankind: a hypothesis of neocortical expansion during evolution. *Trends Neurosci.* **18**, 383–388 (1995).

61. Chance, S. A., Casanova, M. F., Switala, A. E. & Crow, T. J. Minicolumnar structure in Heschl's gyrus and planum temporale: Asymmetries in relation to sex and callosal fiber number. *Neuroscience* **143**, 1041–1050 (2006).

62. Tzourio-Mazoyer, N., Crivello, F. & Mazoyer, B. Is the planum temporale surface area a marker of hemispheric or regional language lateralization? *Brain Struct. Funct.* **223**, 1217–1228 (2018).

63. Ocklenburg, S. *et al.* Neurite architecture of the planum temporale predicts neurophysiological processing of auditory speech. *Sci. Adv.* **4**, 1–9 (2018).

64. Natu, V. S. *et al.* Apparent thinning of human visual cortex during childhood is associated with myelination. *Proc. Natl. Acad. Sci.* **116**, 20750–20759 (2019).

65. Petanjek, Z. *et al.* Extraordinary neoteny of synaptic spines in the human prefrontal cortex. *Proc. Natl. Acad. Sci. U. S. A.* **108**, 13281–13286 (2011).

66. Faust, T. E., Gunner, G. & Schafer, D. P. Mechanisms governing activity-dependent synaptic pruning in the developing mammalian CNS. *Nat. Rev. Neurosci.* **22**, 657–673 (2021).

67. Cuellar-Partida, G. *et al.* Genome-wide association study identifies 48 common genetic variants associated with handedness. *Nat. Hum. Behav.* (2020). doi:10.1038/s41562-020-00956-y

68. Carrion-Castillo, A. *et al.* Genetic effects on planum temporale asymmetry and their limited relevance to neurodevelopmental disorders, intelligence or educational attainment. *Cortex* **124**, 137–153 (2020).

69. Dubois, J. *et al.* Structural asymmetries of perisylvian regions in the preterm newborn. *Neuroimage* **52**, 32–42 (2010).

70. Kasprian, G. *et al.* The prenatal origin of hemispheric asymmetry: An in utero neuroimaging study. *Cereb. Cortex* **21**, 1076–1083 (2011).

71. Hill, J. *et al.* A Surface-Based Analysis of Hemispheric Asymmetries and Folding of Cerebral Cortex in Term-Born Human Infants. *J. Neurosci.* **30**, 2268–2276 (2010).

72. Bishop, D. V. M. & Bates, T. C. Heritability of language laterality assessed by functional transcranial Doppler ultrasound: A twin study. *Wellcome Open Res.* **4**, 1–52 (2020).

73. Somers, M. *et al.* Linkage analysis in a Dutch population isolate shows no major gene for left-handedness or atypical
language lateralization. *J. Neurosci.* **35**, 8730–8736 (2015).
74. Kong, X.-Z. *et al.* Handedness and Other Variables Associated with Human Brain Asymmetrical Skew. *bioRxiv* 756395
(2019). doi:10.1101/756395
75. Karama, S. *et al.* Childhood cognitive ability accounts for associations between cognitive ability and brain cortical
thickness in old age. *Mol. Psychiatry* **19**, 555–559 (2014).
76. Deary, I. J., Whiteman, M. C., Starr, J. M., Whalley, L. J. & Fox, H. C. The Impact of Childhood Intelligence on Later
Life: Following Up the Scottish Mental Surveys of 1932 and 1947. *J. Pers. Soc. Psychol.* **86**, 130–147 (2004).
77. Labache, L. *et al.* A SENtence Supramodal Areas AtlaS (SENSAAS) based on multiple task-induced activation
mapping and graph analysis of intrinsic connectivity in 144 healthy right-handers. *Brain Struct. Funct.* **224**, 859–882
(2019).
78. Biduła, S. P. & Króliczak, G. Structural asymmetry of the insula is linked to the lateralization of gesture and language.
Eur. J. Neurosci. **41**, 1438–1447 (2015).
79. Keller, S. S. *et al.* Can the language-dominant hemisphere be predicted by brain anatomy? *J. Cogn. Neurosci.* **23**,
2013–2029 (2011).
80. Mazoyer, B. *et al.* Gaussian mixture modeling of hemispheric lateralization for language in a large sample of healthy
individuals balanced for handedness. *PLoS One* **9**, 9–14 (2014).
81. Knecht, S. *et al.* Handedness and hemispheric language dominance in healthy humans. *Brain* **123**, 2512–2518 (2000).
82. Afif, A., Bouvier, R., Buenerd, A., Trouillas, J. & Mertens, P. Development of the human fetal insular cortex: Study of
the gyration from 13 to 28 gestational weeks. *Brain Struct. Funct.* **212**, 335–346 (2007).
83. Kalani, M. Y. S., Kalani, M. A., Gwinn, R., Keogh, B. & Tse, V. C. K. Embryological development of the human insula
and its implications for the spread and resection of insular gliomas. *Neurosurg. Focus* **27**, 1–4 (2009).
84. Saadon-Grosman, N., Loewenstein, Y. & Arzy, S. The 'creatures' of the human cortical somatosensory system. *Brain*
Commun. **2**, 1–10 (2020).
85. McCarrey, A. C., An, Y., Kitner-Triolo, M. H., Ferrucci, L. & Resnick, S. M. Sex differences in cognitive trajectories in
clinically normal older adults. *Psychol. Aging* **31**, 166–175 (2016).
86. Cardinale, F. *et al.* Validation of FreeSurfer-Estimated Brain Cortical Thickness: Comparison with Histologic
Measurements. *Neuroinformatics* **12**, 535–542 (2014).
87. Sørensen, Ø., Walhovd, K. B. & Fjell, A. M. A Recipe for Accurate Estimation of Lifespan Brain Trajectories,
Distinguishing Longitudinal and Cohort Effects. (2020).
88. Dalmaijer, E. S. Twin studies with unmet assumptions are biased towards genetic heritability. *bioRxiv*
2020.08.27.270801 (2020).
89. Hibar, D. P. *et al.* Novel genetic loci associated with hippocampal volume. *Nat. Commun.* **8**, (2017).
90. Ocklenburg, S., Hirnstein, M., Beste, C. & Güntürkün, O. Lateralization and cognitive systems. *Washington, DC APA,*
Guidel. Dev. Panel Treat. Posttraumatic Stress Disord. Adults. **5**, (2014).
91. Shafto, M. A. *et al.* The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) study protocol: A cross-sectional,
lifespan, multidisciplinary examination of healthy cognitive ageing. *BMC Neurol.* **14**, 1–25 (2014).
92. Taylor, J. R. *et al.* The Cambridge Centre for Ageing and Neuroscience (Cam-CAN) data repository: Structural and
functional MRI, MEG, and cognitive data from a cross-sectional adult lifespan sample. *Neuroimage* **144**, 262–269
(2017).
93. Kennedy, K. M. *et al.* Effects of beta-amyloid accumulation on neural function during encoding across the adult
lifespan. *Neuroimage* **62**, 1–8 (2012).
94. Wei, D. *et al.* Data Descriptor: Structural and functional brain scans from the cross-sectional Southwest University adult
lifespan dataset. *Sci. Data* **5**, 1–10 (2018).
95. Van Essen, D. C. *et al.* The WU-Minn Human Connectome Project: An overview. *Neuroimage* **80**, 62–79 (2013).
96. Fjell, A. M. *et al.* Self-reported sleep relates to hippocampal atrophy across the adult lifespan - results from the
Lifebrain consortium. *Sleep* 1–15 (2019). doi:10.1093/sleep/zsz280
97. Vidal-Pineiro, D. *et al.* Cellular correlates of cortical thinning throughout the lifespan. Preprint at
https://www.biorxiv.org/content/10.1101/585786v3. (2019).
98. Fischl, B. & Dale, A. M. Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proc.*
Natl. Acad. Sci. U. S. A. **97**, 11050–5 (2000).
99. Zaretskaya, N., Fischl, B., Reuter, M., Renvall, V. & Polimeni, J. R. Advantages of cortical surface reconstruction using
submillimeter 7 T MEMPRAGE. *Neuroimage* **165**, 11–26 (2018).
100. Marie, D., Maingault, S., Crivello, F., Mazoyer, B. & Tzourio-Mazoyer, N. Surface-Based Morphometry of Cortical
Thickness and Surface Area Associated with Heschl's Gyri Duplications in 430 Healthy Volunteers. *Front. Hum.*
Neurosci. **10**, (2016).
101. Greve, D. *et al.* A Surface-based Analysis of Language Lateralization and Cortical Asymmetry. *J. Cogn. Neurosci.* **25**,
1477–1492 (2013).
102. Winkler, A. M. *et al.* Measuring and comparing brain cortical surface area and other areal quantities. *Neuroimage* **61**,
1428–1443 (2012).
103. Bernal-Rusiel, J. L., Reuter, M., Greve, D. N., Fischl, B. & Sabuncu, M. R. Spatiotemporal linear mixed effects
modeling for the mass-univariate analysis of longitudinal neuroimage data. *Neuroimage* **81**, 358–370 (2013).
104. Wood, S. & Scheipl, F. gamm4: Generalized Additive Mixed Models using 'mgcv' and 'lme4'. R package version 0.2-5,
available at https://cran.r-project.org/web/packages/gamm4/gamm4.pdf. (2017).
105. Dray, S. & Dufour, A. B. The ade4 package: Implementing the duality diagram for ecologists. *J. Stat. Softw.* **22**, 1–20
(2007).
106. Margulies, D. S., Falkiewicz, M. & Huntenburg, J. M. A cortical surface-based geodesic distance package for Python.
Gigascience **5**, 1–26 (2016).
107. Neale, M. C. *et al.* OpenMx 2.0: Extended Structural Equation and Statistical Modeling. *Psychometrika* **81**, 535–549
(2016).
108. Jonsson, H. *et al.* Differences between germline genomes of monozygotic twins. *Nat. Genet.* **53**, 27–34 (2021).
109. Mayhew, A. J. & Meyre, D. Assessing the Heritability of Complex Traits in Humans: Methodological Challenges and
Opportunities. *Curr. Genomics* **18**, 332–340 (2017).
110. Bycroft, C. *et al.* The UK Biobank resource with deep phenotyping and genomic data. *Nature* **562**, 203–209 (2018).
111. Yang, J. *et al.* Common SNPs explain a large proportion of the heritability for human height. *Nat. Genet.* **42**, 565–569
(2010).

112.   Fawns-Ritchie, C. & Deary, I. J. Reliability and validity of the UK Biobank cognitive tests. *PLoS One* **15**, 1–24 (2020).
113.   Josse, J. & Husson, F. missMDA: A package for handling missing values in multivariate data analysis. *J. Stat. Softw.*
929   **70**, (2016).

930
931
932

Figure 1–figure supplement 1
Significance of asymmetry effects across samples
*A) Significance for **A)** SA and **B)** CT asymmetry across all samples. Note that because differences in sample size and*
test power affect the overall level of significance and consequently the FDR-correction, the visualization threshold is set
to match the FDR-corrected significance level for each cortical metric in the first sample (LCBC; p < .001). Warm and
cold colours depict significance of leftward and rightward asymmetry, respectively. To permit more fine grained
interpretation of anatomical correspondence, an outline of the Destrieux [1] cortical atlas is overlain. Compare with effect
sizes in Fig. 1 and 2 in main paper.
941

955
956
Figure 1–figure supplement 4
Comparison of vertex-wise and atlas-based asymmetry estimates
To assess to what extent vertex-wise SA and CT asymmetry estimates adhere to the anatomical boundaries of the Desikan-
Killiany (DK) atlas, we derived AI maps per subject and extracted a vertex × subject matrix. Then, within each of the 34 DK
parcels we correlated the AI at each vertex with the parcel mean (i.e. the mean asymmetry across parcel vertices) and
computed the mean correlation for each parcel. High correlations would be expected where atlas-derived parcels fit well to
the underlying vertex-wise structure of cortical asymmetry. We then repeated this analysis using our set of robust
asymmetry clusters. As a formal test, for each cortical metric the resulting coefficients were used as response variable in
linear regressions with Cluster Type (DK parcels vs. Robust clusters) as predictor, controlling for cluster size (nVertices)
and the Cluster Type × nVertices interaction. For SA asymmetry, the average vertex-mean correlations across all Desikan-
Killiany (DK) parcels were r = .53 ± .14 [LCBC], r = .51 ± .14 [UK Biobank], and r = .49 ± .15, which respectively increased
to r = .65 ± .18, r = .64 ± .18, and r = .62 ± .19 for SA asymmetry clusters. For CT, the average vertex-mean correlations
across all DK parcels were r = .36 ± .12 [LCBC], r = .39 ± .12 [UK Biobank] and r = .38 ± .12, which respectively increased
to r = .47 ± .11, r = .50 ± .11 and r = .48 ± .12. As would be expected, within parcel/cluster vertex-mean correlations were
significantly lower in larger parcels/clusters. Linear regressions (size controlled) revealed a significant main effect of Cluster
Type in all but one association test (see SI Table 5), confirming the visual impression that DK parcels conform poorly to the
underlying asymmetry of cortex. *A-C) Average correlation between vertex-wise estimates of SA asymmetry within DK*
parcels to the mean across each parcel (top rows) and between vertex-wise estimates of SA asymmetry within robust
clusters to the mean across each cluster (bottom rows) within each dataset. D) Vertex-mean correlation coefficients for SA
asymmetry in DK parcels and robust asymmetry clusters, shown as raw values and after correcting for number of vertices
*in the parcel/cluster. Comparable analyses for CT asymmetry are shown in **D-H)**. The complete set of raw vertex-mean*

correlation coefficients correlated highly between all dataset-pairs for both SA (min *r* = 0.97) and CT (min *r* = 0.95)
Post=posterior; Lat=lateral; Med=medial; Ant=anterior.
980
981

987　Mean SA and B) CT asymmetry in each dataset analysed using standard methods on the *fsaverage* template within
988　parcels from the HCP multimodal brain atlas [2]. We used this atlas here because it appeared best suited to assess
989　parcels that are homotopic. Warm and cold colours depict leftward and rightward asymmetry, respectively.
990　Post=posterior; Lat=lateral; Med=medial; Ant=anterior; Sup=superior; Inf=inferior.
991

CT asymmetry results in the HCP dataset vary depending on the preprocessing pipeline. A) Results from using the –
hires argument to recon-all (as in main paper). We used this method to best harmonize preprocessing across cohorts
whilst accounting for the higher resolution of HCP data. Shown again for comparison, the top row in A is akin to the
results in the main paper (analysed using cross-hemispheric registration methods) whereas the bottom row is the same
data analysed using standard parcellation methods (as in Figure 1–figure supplement 5). B) CT asymmetry results
using the HCP preprocessed data subject to extra preprocessing steps and inputs, analysed using cross-hemispheric
registration methods (top row) and standard parcellation methods (bottom). C) Results using the HCP preprocessed
data when calculating thickness asymmetry on the fs_LR template. Warm and cold colours depict leftward and
rightward asymmetry, respectively. Post=posterior; Lat=lateral; Med=medial; Ant=anterior; Sup=superior; Inf=inferior.
1004

Figure 3–figure supplement 1

Knot comparison

*Lifespan trajectories of population-level **a)** SA and **b)** CT asymmetries modelled with different smoothing parameters (number of knots 4-8, with the selected number of knots [6] shown in black dotted outline). Grey indicates overlap. All age trajectories were fitted using GAMMs. Inset plots show absolute asymmetry trajectories across life at the different smoothing parameters.*

1013

Figure 3–figure supplement 2

Smooth Age x Hemisphere interactions

*Generalized additive mixed Model (GAMM) results for population-level **a)** SA and **b)** CT asymmetries. A smooth Age ×*
Hemisphere interaction [s(LH)-s(RH-Age)] was fitted to model asymmetry changes across the lifespan. Specifically,
GAMMs were used to compute the zero-centered age-trajectories of the left [s(LH-Age)] and right [s(RH-Age)] hemisphere
for each cluster, and the asymmetry trajectory was computed as the difference between the two [s(LH-Age)-s(RH-Age)].
Gold and pink colours denote clusters defined by leftward or rightward asymmetry, respectively. For each cluster the main
effect of Hemisphere was added to visualize the lifespan trajectory of absolute asymmetry. Bands represent 95%
confidence intervals. Note that all SA clusters show asymmetry trajectories that are significantly different from 0 (symmetry;
dotted line) across the entire lifespan, and 19/20 CT clusters were significantly asymmetric already by ~age 4.

1024

Figure 3–

Figure 3–figure supplement 3

Lifespan CT trajectories in regions exhibiting age-related change in asymmetry

Homotopic lifespan CT trajectories in an alternative set of regions derived from a previous analysis (clusters from Roe et al., 2021 [3]; derived from vertex-wise analyses of age-related CT asymmetry change in LCBC adult data [20-89 years]). Dark colours correspond to LH trajectories (colours correspond to clustering solutions in Roe et al., 2021 [3]). To highlight development, datapoints are semitransparent after age 30. All GAMM models are equivalent to those in main paper.

1033

SI Fig. 10 – UK Biobank lower quadrant
Thickness asymmetry interrelationships between opposite-direction asymmetries (lower left quadrant of the UK Biobank
correlation matrix in SI. Fig. 3; N = 38,172) are visualized to describe whether negative asymmetry-asymmetry correlations
pertained to reduced or reversed thickness asymmetry. Lines of symmetry (0) are shown in grey. X-axis and Y-show raw
AI data after removing the fixed effects of age and sex (i.e. data is not scaled to have a mean of 0). AI's in rightward clusters
are inversed, such that positive correlations would denote positive asymmetry-asymmetry relationships regardless of
direction. Order of cortical locations is in SI. Fig. 3)
1054

1055
1056
Figure 4–figure supplement 3
SI Fig. 11 – Global thickness asymmetry relationships
Global thickness asymmetry relationships. Principal components analysis across AI's in all leftward and rightward CT
asymmetry clusters revealed a single component explained most of the variance in UK Biobank, and there was evidence
of a relatively stronger first component in LCBC and HCP (inset scree plots). Scatter plots show the partial correlation of
the mean asymmetry across all leftward vs. all rightward clusters (means weighted by cluster size) plotted for each cohort,
after AI's were corrected for age, sex and (where applicable) scanner. Lines of symmetry (0) are shown as dotted grey. AI's
in rightward clusters are inversed, such that positive correlations would denote positive asymmetry-asymmetry relationships
regardless of direction.
1066
1067
1068

1069   ***Figure 4–figure supplement 4***
1070   ***HCP outliers discarded***
1071   *Two outliers (black) were detected in the cortical thickness data of HCP and subsequently discarded for all subsequent*
1072   *analyses. Plot shows mean CT asymmetry across all leftward vs. mean CT asymmetry across all rightward clusters.*
1073

Areal asymmetry effects
UK Biobank

Thickness asymmetry effects
UK Biobank

Figure 6–figure supplement 1
ICV effects
Continuation of Fig. 5 in main paper. Cortical locations of the effects underlying the observed associations of robust SA
(left) and CT (right) asymmetry phenotypes with brain size (ICV) in UK Biobank. Plots denote the effect sizes
(standardized Betas) and cortical location of all associations surpassing Bonferroni-corrected significance (Fig. 5). Right
handers and females are coded 0, such that a negative effect size for handedness / sex / ICV / cognition denotes less
asymmetry in left handers / males / larger brains / higher cognition. Blue and yellow clusters denote leftward and rightward
asymmetry, respectively.
1083
1084
1085
1086