Cryo-EM structure of the
Smc5/6 holo-complex

Stephen T. Hallett[1], Isabella Campbell Harry[1],

Pascale Schellenberger[2], Lihong Zhou[1], Nora B. Cronin[3], Jonathan Baxter[1],

Thomas J. Etheridge[1*], Johanne M. Murray[1*] and Antony W. Oliver[1*]

1. Genome Damage and Stability Centre, School of Life Sciences,
   University of Sussex, Falmer, UK
2. Electron Microscopy Imaging Centre, School of Life Sciences,
   University of Sussex, Falmer, UK
3. London Consortium for CryoEM (LonCEM) Facility,
   The Francis Crick Institute, London, UK.

* To whom correspondence should be addressed:

TJE /// t.etheridge@sussex.ac.uk, ORCID: 0000-0001-8144-6917

JMM // j.m.murray@sussex.ac.uk, ORCID: 0000-0001-9225-6289

AWO // antony.oliver@sussex.ac.uk, ORCID: 0000-0002-2912-8273

1   ABSTRACT

The Smc5/6 complex plays an essential role in the resolution of recombination intermediates

formed during mitosis or meiosis, or as a result of the cellular response to replication stress.

It also functions as a restriction factor preventing viral integration. Here, we report the cryo-

EM structure of the six-subunit budding yeast Smc5/6 holo-complex, reconstituted from

recombinant proteins expressed in insect cells – providing a full overview of the complex in

its apo / non-liganded form, and revealing how the Nse1/3/4 subcomplex binds to the hetero-

dimeric SMC protein core. In addition, we demonstrate that a region within the head domain

of Smc5, equivalent to the 'W-loop' of Smc4 or 'F-loop' of Smc1, mediates an essential

interaction with Nse1. Taken together, these data confirm a degree of functional equivalence

between the structurally unrelated KITE and HAWK accessory subunits associated with SMC

complexes.

INTRODUCTION

The eukaryotic Structural Maintenance of Chromosomes (SMC) family includes the protein complexes cohesin, condensin and Smc5/6. At their respective 'hearts' sits an obligate heterodimer of two SMC proteins, either Smc1/Smc3, Smc2/Smc4 or Smc5/Smc6. Globular entities found at both the N- and C-termini of each SMC protein are brought together in space, to form a so-called 'head' domain that is capable of binding to, and turning over, ATP. The two halves of the ATPase are connected by a structural excursion known as the 'arm'; formed from sequential alpha-helical regions that coalesce to form a highly extended anti-parallel coiled-coil. The arm is interrupted at its apex (most distant point from the head) by the 'hinge' domain; a region of SMC proteins responsible for hetero-dimerisation with their obligate binding partner. The binding, hydrolysis, and release of ATP by the two head domains (one from each SMC protein) provides a secondary, more transient, and regulated dimerisation interface. The 'core' of each SMC complex is then elaborated through binding of additional 'non-SMC' protein subunits or 'elements', to provide the distinct functionalities required for their respective cellular functions.

For more expansive reviews of the SMC-family, including their respective functions and subunit compositions, please see: (Cutts and Vannini 2020; Datta, Lecomte, and Haering 2020; Hassler, Shaltiel, and Haering 2018; Matityahu and Onn 2021; Palecek 2018; Sole-Soler and Torres-Rosell 2020; Uhlmann 2016; Yatskevich, Rhodes, and Nasmyth 2019).

All three complexes are required for the organisation and management of chromosome architecture and structure throughout the cell cycle. Cohesin has well described roles in sister chromatid cohesion and the organisation of the interphase chromosomes into topologically associated domains or TADs, whereas condensin is required to compact chromosomes at mitosis. The Smc5/6 complex has roles in the processes of DNA replication and DNA damage repair; the complex acting to suppress / prevent formation of inappropriate structures that can form during homologous recombination-mediated rescue of replication forks that have stalled (or collapsed) on encountering replication 'road-blocks' or obstacles.

Alterations to the coding sequence of human Nse2 (Non-SMC-element; generally written as either Nse or NSMCE) have been linked to primordial dwarfism and insulin resistance (Payne et al. 2014), with changes in Nse3 linked to severe lung disease immunodeficiency and chromosome breakage syndrome (LICS, van der Crabben et al. 2016). Interestingly, Smc5/6 is also specifically targeted for ubiquitylation and degradation by the regulatory protein X of hepatitis B virus (HbX) to alleviate restriction of viral replication by the complex (Murphy et al. 2016). The complex has also been shown to act as a restriction factor working against other viruses (Bentley et al. 2018; Gibson and Androphy 2020; Xu et al. 2018).

As well as containing subunits that provide both ubiquitin and SUMO E3-ligase activity (through Nse1 and Nse2 respectively), Smc5/6 is further differentiated from the cohesin and condensin complexes by the fact that the two non-SMC proteins (Nse1 and Nse3) that bind to its kleisin subunit (Nse4) belong to the 'KITE' family (kleisin-interacting tandem winged-helix element; Palecek and Gruber 2015) rather than the distinct and structurally unrelated 'HAWK' family (HEAT proteins associated with kleisins; Wells et al. 2017). As proteins of the KITE-family are also found in prokaryotic SMC complexes, it has led to the hypothesis that Smc5/6 best represents the eukaryotic 'cousin' of *Bacillus subtilis* Smc/ScpAB and *Escherichia coli* MukBEF (Palecek and Gruber 2015) — however, it is still unclear as to when in the evolutionary timescale KITEs were replaced by HAWK subunits, to create cohesin and condensin complexes.

Here, we present the cryo-EM structure of the budding yeast Smc5/6 complex, in its apo or non-liganded form. Our study serves to confirm the overall architecture of the complex, plus provide details of how and where the Nse1/Nse3/Nse4 KITE-kleisin subcomplex interacts with the Smc5/Smc6/Nse2 core. Additional experiments reveal the presence of a crucial interface formed between the head domain of Smc5 and Nse1, which utilises the equivalent of the 'W-loop' or 'F-loop' found in Smc4 and Smc1 respectively (Hassler et al. 2019). Taken together, our data uncover an unanticipated degree of functional equivalence between KITE and HAWK accessory subunits.

RESULTS

In this manuscript, for simplicity and brevity, we refer to the protein components of the *Saccharomyces cerevisiae* SMC complexes, unless otherwise indicated.

We have previously described reconstitution and characterisation of the *S. cerevisiae* Smc5/6 complex, using recombinant proteins expressed in insect cells (Hallett et al. 2021). Here, cryo-EM data were collected for the six component 'holo-complex', where Smc5 and Smc6 contain Walker B mutations E1015Q and E1048Q, respectively (**Fig. 1a, Materials and Methods**).

A neural network was trained with a set of manually picked particles (Bepler et al. 2019). The resultant 'picking model' was then used to identify a total of 380,714 particles (see **Sup. Fig. 1** for data-processing summary). Two sets of two-dimensional (2D) class averages emerged from processed data, with either the 'hinge' (**Fig. 1b, top**) or the 'head-end' of the complex more clearly in focus (**Fig. 1b, bottom**); the blurred density at either end consistent with a degree of conformational flexibility within the coiled coil 'arms' (Hallett et al. 2021). From this, a sub-set of 17,162 particles were used to generate a medium-resolution 3-dimensional (3D) map that covered the entire length of the holo-complex, reconstructed at a resolution of 10.8 Å as judged by the 0.143 Fourier shell correlation (FCS) criterion (Rosenthal and Henderson 2003) (**Fig. 1c**). Focussed refinement allowed a slightly higher resolution map to be calculated for the upper (106,660 particles, 8.5 Å) segment of the complex (**Fig. 1d**). Pleasingly, resultant maps were consistent with the envelope we previously obtained by uranyl acetate negative stain transmission electron microscopy (Hallett et al. 2021), **Fig. 1d, inset**).

In parallel, we trained a second neural network with manually picked particles that encompassed just the head-end of the complex (824,644 particles). The resultant 2D class averages and 3D map from processed data (84,180 particles, 6.5 Å) are shown in **Fig. 1e** and **1f** respectively. Segmentation analysis (Pettersen et al. 2021; Pintilie and Chiu 2012; Pintilie et al. 2010) allowed portions of the map to be readily assigned to the head domains, as well as the Nse1/Nse3/Nse4 (Nse1/3/4) subcomplex (coloured red, blue and grey respectively in **Fig. 1f**). Additional segments of density associated with either the 'head' or 'arm' (dark or light

green, **Fig. 1f**) allowed the identity of each head-domain to be determined, due the expected

binding positions of the N- and C-terminal domains of the kleisin Nse4 (please see additional

text below).

A model for the Smc5/6 holo-complex

A sharpened composite map, generated from the highest resolution maps for each segment

of the holo-complex, was of sufficient quality to allow positioning of the coiled-coil regions

for both Smc5 and Smc6, as well as placement of the structure of *S. cerevisiae* Nse2 (Mms21)

bound to the arm of Smc5 (PDB: 3HTX; Duan et al. 2009). This, combined with homology

models for the 'hinge', 'heads' and Nse1/3/4, allowed an initial pseudo-atomic model to be

constructed; this model was subsequently updated using AlphaFold predictions made

available through the EMBL-EBI repository (https://alphafold.ebi.ac.uk; Jumper et al. 2021)

(see **Materials and Methods**).

Working from the 'top' of the complex downwards, we see that the hinge domain is tilted

with respect to an axis defined by the coiled-coil arms; consistent with our previous analysis

of this region in the fission yeast complex and with data published for condensin (Alt et al.

2017; Soh et al. 2015). Below this, there is an apparent discontinuity / break in the helices of

the arms, located between the hinge and Nse2 subunit — a feature likely to represent the

'elbow' found in the other complexes of the SMC-family (Burmann et al. 2019; Kong et al.

2020; Lee et al. 2020) — notably, at this point the two arms of Smc5/6 also cross over each

other (**Fig. 2a**, inset *ii*).

In our previous lower-resolution study, we observed that the arms of Smc5/6 remain in close

proximity ('arms-together', 'rod-like', 'I'-conformation) for the majority of their length

(Hallett et al. 2021), in agreement with parallel studies published by other laboratories

(Gutierrez-Escribano et al. 2020; Taschner et al. 2021; Yu et al. 2021). With the additional

resolution now afforded by cryo-EM, we see that the arms separate slightly from each other

after the junction with Nse2 (**Fig. 2a**, inset *iii*) but then remain approximately parallel until the

head-end of the complex. Our model positions the first alpha-helix (α1) of Nse2 such that it

can 'talk' across to the descending helix (hinge to head) of Smc6, as well as make its previously

documented set of interactions with Smc5 (Duan et al. 2009); the helix appearing to 'glue'

the two arms together, in agreement with crosslinking data for the holo-complex recently

published by Taschner et al. (2021) (**Sup. Fig. 2a**). The two arms then briefly re-contact each

other, through the two inward facing helices (αC3) of the 'joint' (**Fig. 2a,** inset *iv*) — a

molecular feature common to SMC proteins (Diebold-Durand et al. 2017) formed from three

helix-loop repeats that encircle the more continuous ascending alpha-helix, thus generating

both an interruption and point of flexure within the coiled-coiled structure of the arm; again

supported by crosslinking data (**Sup. Fig. 2b**; Taschner et al. 2021).

In our model, the head domains of Smc5 and Smc6 are not in direct contact. Comparison of

head domain positions indicates that our structure most resembles that of the apo / ATP-free

/ 'non-engaged' / juxtaposed / or 'J'-state of budding yeast condensin deposited under PDB

accession code 6YVU (Lee et al. 2020) rather than the ATP-bound / 'engaged' / 'E'-state seen

for budding yeast cohesin when in complex with Scc2 and DNA (PDB: 6ZZ6; Collier et al. 2020)

(**Fig. 2b**). Sections of the map corresponding to the RecA lobe of both the Smc5 and Smc6

head domain become less apparent at higher contour levels, suggesting a degree of

conformational flexibility in this part of the complex, as supported by estimates of local

resolution (**Sup. Fig. 2c**).

Revealing the binding location of the Nse1/3/4 sub-complex

The Nse1/3/4 subcomplex is located to one side of the central axis defined by the coiled-coil

arms of Smc5 and Smc6. The winged-helix 2 (WH/2) domain of Nse1 and the head domain of

Smc5 are in direct contact, with a short loop protruding from the equivalent winged-helix

domain of Nse3 (WH/B) positioned to interact with the arm of Smc5 (amino acids 244-254,

Fig. 2a inset *v*). Notably, neither Nse1 nor Nse3 directly interact with Smc6.

The recent publication of an X-ray crystal structure for *Xenopus laevis* Nse1/3/4, has provided

molecular details for how the kleisin subunit interacts with the Nse1/Nse3 KITE heterodimer

(kleisin-interacting tandem winged-helix element). The central section of Nse4 follows a path

through the centre of both KITE proteins, interacting with the linker regions that serve to

connect their component winged-helix domains together (Jo et al. 2021). Pleasingly, our cryo-

EM data allows integration of this kleisin path with the set of interactions made by the

globular domains found at each of its termini: the helical N-terminal domain binding to the

'neck' of Smc6 (**Fig. 1f and 2c**) and the C-terminal domain to the 'cap' of the Smc5 head

domain (**Fig. 1f and 2d**, **left**) thus confirming at the structural level the set of kleisin-facilitated

interactions conserved across the SMC-family of complexes, including prokaryotic ScpAB and

MukBEF as well as eukaryotic condensin and cohesin (Burmann et al. 2013; Diebold-Durand

et al. 2017; Fennell-Fezzie et al. 2005; Gligoris et al. 2014; Haering et al. 2004; Kamada et al.

2017; Woo et al. 2009; Zawadzka et al. 2018). Saying this, the resolution of our composite

map is not sufficient to allow an unambiguous tracing of the parts of Nse4 that serve to

connect the N-terminal domain to the central section (amino acids 125-183) or the central

section to the C-terminal domain (aa 46-283), indicating either intrinsic disorder or a high

degree of conformational flexibility; an observation consistent with the kleisin moieties in

cryo-EM structures of other eukaryotic SMC complexes (Collier et al. 2020; Higashi et al. 2020;

Lee et al. 2020; Shi et al. 2020).

Additional structural features are predicted by AlphaFold

AlphaFold predicts (albeit with a lower level of confidence) the presence of a hereto unknown

alpha-helical element in the C-terminal domain of Nse4 (amino acids Ser360 through to

Ala372). Nicely, this provides a facile explanation for a region of additional density visible in

our composite map, not accounted for by the initial homology model (**Fig. 2d left**). Similarly,

AlphaFold predicts the presence of a budding yeast-specific extended loop within the NH-

RING domain of Nse1 (amino acids Glu287 to Gln303), which again provides a better overall

fit to the experimental map (**Fig. 2d right, Supp. Fig 2d**).

A loop in Smc5 has functional equivalence to the W-loop of Smc4 and F-loop of Smc1

The location of Nse1/3/4 binding is particularly striking, as it echoes that of the structurally

unrelated HAWK accessory proteins (HEAT proteins associated with kleisins) found in

condensin and cohesin complexes; in each case, the head domain of the κ-SMC (Smc5, Smc4,

Smc1) serving to provide a major interaction surface (**Fig. 3a**).

Closer inspection reveals that a region of Smc5 (amino acids Gly947—Gly978), which 'talks'

across to the Nse1 subunit, is structurally equivalent to the 'W-loop' of the Smc4 head domain

(Hassler et al. 2019) (**Fig. 3b, 3c inset**). Notably, defined mutations (S1316D or W1317A)

introduced into the W-loop render budding yeast cells non-viable. Furthermore, in the

context of a fully recombinant 'head complex', mutation of the equivalent tryptophan residue in *Chaetomium thermophilum* (*Ct*) Smc4 resulted in a dramatic reduction of its ability to turnover ATP (Hassler et al. 2019). In Smc1, the equivalent region has been given the alternative title of 'F-loop', due to the presence of a conserved phenylalanine residue (F1123 in budding yeast Smc1 (Hassler et al. 2019; Petela et al. 2021; Shi et al. 2020).

In Smc5, Leu978 appears to be the structurally equivalent 'F' or 'W-loop' residue (**Fig. 3b**) — a relationship supported by a strong preference for a leucine in this position, as revealed by a cross-species multiple amino acid sequence alignment (**Fig. 3c**). However, two additional amino acids within the same loop, Phe972 and Leu981, are also strictly conserved (label F in **Fig. 3b** and **c**). A second region of conservation was also evident in the preceding loop (labelled Y in **Fig. 3b and c**). Here, structural comparison suggested that one or more aromatic residues might potentially act to anchor or connect the helical lobe of the head domain back to its coiled-coil arm. Interestingly, in *E. coli* MukB, this region is highly elaborated to form the so-called 'larynx', residues of which contribute to its DNA-binding interface (Bürmann et al, 2021; https://doi.org/10.1101/2021.06.29.450292).

When working with homology models and structural data of limited resolution, there is a degree of uncertainty as to the precise amino acids involved in a molecular interface. With this in mind, we introduced sets of mutations designed to either disrupt localised structure or alter key features within a loop (**Materials and Methods**). Yeast harbouring the (Y) Y961A, W964A double mutation were viable and surprisingly displayed no sensitivity to a range of genotoxic agents. In sharp contrast, introduction of the (F) F972A, L978D, L981N triple mutant rendered budding yeast cell inviable, as determined by dissecting sporulated diploids (**Fig. 3c, Sup. Fig. 3a**); confirming the importance of this loop to cellular function. We also introduced a set of complementary mutations into Nse1, designed to perturb the interaction with the Smc5 loop (N: F217A, E228R, R242A). Here, yeast were viable, but had a slow growth phenotype and mild sensitivity to treatment with MMS or camptothecin (**Fig. 3c and Sup. Figs. 3c and 3d**).

DISCUSSION

Using recombinant proteins expressed in insect cells we have reconstituted, then visualised by cryo-EM, the six-subunit budding yeast Smc5/6 'holo-complex' — in its apo or unliganded form — to provide a structural overview of the complex's architecture and reveal the position of the bound Nse1/3/4 sub-complex. Whilst structurally unrelated to the HAWK-family of proteins (HEAT proteins associated with kleisins; Wells et al. 2017) the set and type of interactions made by the KITE (kleisin-interacting tandem winged-helix element; Palecek and Gruber 2015) hetero-dimer of Nse1/3 and the kleisin Nse4 indicate several unifying features that span the SMC-family of complexes:

i) KITE and HAWK subunits can bind to double-stranded DNA

Smc5/6: The ability of the Smc5/6 complex to bind dsDNA is provided by the Nse1/3 heterodimer (see also text below, Zabrady et al. 2016). Furthermore, reconstituted complexes lacking both Nse1/3 and the kleisin Nse4 do not bind dsDNA (Hallett et al. 2021). In condensin and cohesin, a similar dsDNA-binding activity is provided by the HAWK subunits, with the notable exception of Pds5 (Li et al. 2018). **Condensin:** Both the Brn1-Ycg1-Ycs4 hetero-trimer (Piazza et al. 2014) as well as Ycg1 in complex with the C-terminal portion of Brn1 (Kschonsak et al. 2017) can interact with nucleic acid. Whilst DNA-binding by Ycs4 has not been formally demonstrated, an examination of its structure and surface charge suggests the presence of a compatible region located between HEAT repeats 7 and 8 (Lee et al. 2020); a view supported by experiments using *Ct*Ycs4 and *Ct*Ycg1 which, as isolated proteins, have affinity for dsDNA (Piazza et al. 2014). **Cohesin:** Scc3, when bound to Scc1, is capable of binding to dsDNA (Li et al. 2018). DNA-binding has been reported for Mis4, the *S. pombe* equivalent of Scc2, both in the context of its hetero-dimeric complex with Ssl3 (equivalent to Scc4) or as an isolated protein (Murayama and Uhlmann 2014; Kurokawa and Murayama 2020). Recent data indicates a similar pattern of behaviour for both the Scc2/Scc4 hetero-dimer and for Scc2 alone [Collier *et al.*, 2020; https://doi.org/10.1101/2020.06.03.132233].

ii) Addition of dsDNA stimulates ATPase activity

Smc5/6: Addition of dsDNA to Smc5/6 strongly stimulates its ATPase activity (Fousteri and Lehmann 2000) with enhancement being strictly dependent on the presence of the Nse1/3/4

sub-complex (Hallett et al. 2021). **Condensin:** Similarly, the Brn1-Ycg1-Ycs4 or 'non-SMC'

complex is essential for dsDNA-dependent simulation of ATPase activity (Piazza et al. 2014).

Cohesin: Here, the Scc2 subunit is required, as determined by experiments that added the

protein to pre-assembled complexes containing either the core hetero-trimer

(Smc1/Smc3/Scc1) or hetero-tetramer (Smc1/Smc3/Scc1/Scc3) (Petela et al. 2018). A similar

requirement has been demonstrated for the fission yeast cohesin complex (Murayama and

Uhlmann 2014). In each case, the presence of the kleisin subunit is required to connect the

dsDNA binding activity of the KITE/HAWK protein to the core SMC complex.

iii) dsDNA-binding is intrinsically linked to chromatin-binding, loading or retention

Smc5/6: Hypomorphic (non-lethal) mutations that reduce the ability of the Smc5/6 complex

to bind to dsDNA, result in a loss of chromatin association, as shown by chromatin immuno-

precipitation (ChIP) and live-cell single-molecule tracking experiments (Etheridge et al. 2021;

Zabrady et al. 2016). **Condensin:** Mutations that reduce the ability of Brn1 to interact with

dsDNA, as part of its composite interaction surface with Ycg1, reduce the amount of

condensin associated with chromatin at defined loci (Kschonsak et al. 2017). **Cohesin:**

Mutations that affect the ability of Scc2 or Mis4 to bind DNA result in reduced chromatin

binding/loading, at several known cohesin-enriched loci (Collier et al. 2020; Kurokawa and

Murayama 2020).

iv) mutations within the W-loop (or equivalent) of the κ-SMC perturb function

The 'W-loop' is a conserved structural feature, present in each of the κ-SMC subunits (Smc1,

Smc4, Smc5) albeit with different patterns of amino acid conservation. Mutation of highly

conserved amino acids within the loop are incompatible with cell viability of yeast when

introduced into Smc4 (Hassler et al. 2019) or Smc5 (this manuscript).

By taking each of the above points in hand, we can now add the Nse1/3 hetero-dimer into

the statement made by Petela et al. (2018) "A role in contacting DNA might therefore be a

feature conserved between Scc2 and Ycg1" and also amend the statement made by Hassler

et al. (2019) that '…interaction of a HEAT-repeat subunit with the κ-SMC$_{hd}$ (head domain) is a

central feature of all condensin and cohesin complexes' — to also include Smc5/6 — to state

that 'interaction of a kleisin-associated subunit, whether HAWK or KITE, with the κ-SMC head

domain is a central feature of the hetero-dimeric SMC complexes, condensin, cohesin and

Smc5/6.'

A speculative model for DNA-binding and engagement

Zabrady et al. (2016) previously reported a docking pose for binding of a double-stranded

DNA duplex to a positively charged cleft formed between human Nse1 and Nse3

(NSMCE1/NSMCE3) supported by a range of biochemical and biophysical data. A more recent

study using *Xenopus laevis* Nse1/3/4 has reported a similar mode of interaction (Jo et al.

2021). By simple superposition, one end of the docked DNA duplex would physically clash

with the arm of Smc5 (**Fig. 4a, left**). However, the DNA duplex could be accommodated by a

relatively small movement of the Nse1/3/4 subcomplex away from the Smc5 head domain

(or vice versa). Saying this, the resulting pose would still be at odds with the expected dsDNA

binding location, as illustrated here by comparison to the DNA-bound *S. cerevisiae* cohesin

complex (**Fig. 4a, right**. SMC1-SMC3-SCC1-SCC2 complex; PDB: 6ZZ6), but also observed for

other proteins of the SMC-family, including the more distantly related Rad50 (Kashammer et

al. 2019).

In their 2019 paper, Hassler et al. state that 'It is likely that tilting and repositioning of the W-

loop residues dissociates Ycs4, which binds this part of Smc4$_{hd}$ (head-domain) and sterically

blocks access of Smc2$_{hd}$'. The follow-on study builds on this initial hypothesis to uncover a

'flip-flop' mechanism, which involves a physical switch of HEAT subunit from Ycs4 (binding to

the head domain of Smc4) in the apo/unliganded state, to Ycs1 (binding to the head domain

of Smc2) in the engaged/ATP-bound state (Lee et al. 2020). It is therefore tempting to

speculate that a similar mechanism may well be at play in Smc5/6, as in order for the head-

domain of Smc5 to become engaged with that of Smc6, a conformational change is dictated

— potentially linked to breaking of the Smc5 'W-loop' / Nse1 interface, to release and allow

repositioning of the KITE sub-complex, and thus facilitate engagement of a bound dsDNA

substrate with the top section of each head domain (see speculative model presented in **Fig.**

4b). Here, however, a 'flip-flop' mechanism might invoke the Nse3 subunit and a potential

direct interaction with the head domain of Smc6. This will, however, require experimental

validation, especially given that in PDB entry 6ZZ6 (cohesin) the F-loop of Smc1 remains

engaged with the Scc2 subunit, indicating that the precise molecular choreography
underpinning the initial 'engagement' complex is still unclear, and may well be (subtly?)
different for each complex of the SMC-family in accordance with their respective cellular
functions.

Does Nse2 prevent or control bending at the elbow?
We and others have observed some degree of flexure at the 'elbow' of Smc5/6 (Gutierrez-
Escribano et al. 2020; Hallett et al. 2021; Serrano et al. 2020; Taschner et al. 2021; Yu et al.
2021) but without the acute bend observed in condensin, cohesin and MukBEF (Anderson et
al. 2002; Burmann et al. 2019; Collier et al. 2020; Higashi et al. 2020; Kong et al. 2020; Lee et
al. 2020; Shi et al. 2020). However, the set of interactions made by Nse2 with Smc5, and now
Smc6, may provide insight, as these are likely to prevent separation and rotation of the arms,
and therefore the concerted set of motions thought to produce the bend (Burmann et al.,
2019).

Interestingly, a major hotspot for auto-SUMOylation has been mapped to a region within the
coiled-coil arm of Smc5, sitting between amino acids Lys310 and Lys327. Whilst the precise
site(s) of modification appears to be redundant, Lys311 has been identified with high
confidence as a major site of auto-SUMOylation (Zapatka et al. 2019). Saying this, it is not
known what effect SUMOylation has on the overall structure / conformation of the Smc5/6
holo-complex, but the proximity of the primary modification site to the α1-helix of Nse2 hints
at a regulatory role; potentially one that acts to control arm 'architecture' (**Fig. 2**, inset *iii*) as
well as performing the documented roles in promoting interactions with, and / or modulating
activity of, replication fork associated-DNA helicases such as Sgs1[BLM] (Bermudez-Lopez et al.
2016) and Mph1[FANCM] (Xue et al. 2014).

Interestingly, SUMOylation catalysed by the Smc5/6 complex has been shown to be
stimulated by the addition of single-stranded DNA (ssDNA) to reaction mixes (Varejao et al.
2018) with circular dichroism experiments showing localised changes in the secondary
structure of Smc5 upon DNA-engagement. We speculate that this is likely to occur as a direct
consequence of an initial binding event at the hinge-region of the complex, that leads to
stimulation and auto-modification of Smc5/6, which then allows the complex to

control/modulate homologous recombination activity at a stalled replication fork,

presumably through regulation of specific helicases and/or other modified substrates;

supported by our experimental observations in fission yeast, where mutations affecting the

ability of the hinge to bind ssDNA lead to gross chromosomal rearrangements, but do not

impact chromatin-binding or retention of the complex {Etheridge, 2021 #164}.

Finally, our cryo-EM structure of the apo/non-liganded state provides a highly valuable

'stepping stone' along the way to a fuller understanding of the set of molecular interactions

and dynamics than underpin the cellular function(s) of the Smc5/6 complex. Clearly, a

desirable goal is now the determination of structures for the ATP-bound and DNA-engaged

states.

MATERIAL AND METHODS

Expression and purification

Detailed experimental procedures for both expression and purification of the *S. cerevisiae* Smc5/6 holo-complex are available in (Hallett et al. 2021). For convenience, the composition of the two buffers that have been used in this study are listed below:

BUFFER C: 20 mM HEPES.NaOH pH 7.5, 100 mM NaCl, 0.5 mM TCEP

BUFFER F: 20 mM HEPES.NaOH pH 7.5, 0.5 mM TCEP

Cryo-EM

Sample preparation

Fractions eluting from a Superose 6 size exclusion chromatography column (equilibrated in BUFFER C; Cytiva Life Sciences, Little Chalfont, UK), corresponding to BS3-crosslinked holo-complex, were immediately used for grid preparation. The sample was diluted by a factor of two (with addition of BUFFER F) to reduce the overall NaCl concentration to 50 mM and yield a final concentration of 0.1 mg/ml. From this, 3 μl was applied to a freshly glow-discharged grid (Quantifoil R0.6/1 Cu 300 mesh grid; 60 seconds, 15 mA, PELCO easiGlow — Agar Scientific, Stansted, UK). Using an EM GP2 automatic plunge freezer (Leica Microsystems, Wetzlar, Germany) the grid was held in a chamber at 10 °C and 90% relative humidity, for a period of 10 seconds, before blotting for 2.5 - 4.5 seconds using the auto-sensor. The grid was immediately plunged into liquid ethane at -182 ° and then stored under liquid nitrogen until data collection.

Data collection

Data were collected at LonCEM (The Frances Crick Institute, London, UK) at 300 kV on a Titan Krios electron microscope (Thermo Fisher Scientific, Waltham, MA USA), equipped with a Gatan K3 detector operating in counted super-resolution mode. Movies were acquired using EPU (Thermo Fisher Scientific). Data were collected from four grids across separate sessions, without hardware binning at a calibrated pixel size of 0.55 Å. Target defocus was -1 to -3.5 μm, with a total dose of 50 electrons per $Å^2$, during an exposure time of 3.9 to 5 seconds, fractionated into 38-50 frames.

Data processing

An overview of the workflow used to process data is provided in **Fig. S1**. In summary, all movie frames were aligned using 5 x 5 patches in Motioncor2 with dose-weighting (Zheng et al. 2017). All data were binned by a factor of 2. Contrast transfer function (CTF) parameters were estimated using CTFFIND4 (Rohou and Grigorieff 2015). Particles were picked using Topaz (Bepler et al. 2019) integrated into CryoSPARC. Initial training used a set of particles manually picked from a few micrographs, with the resultant 'trained' model being used to auto-pick particles across all micrographs. Apart from the 'Head-only' approach, all processing was carried out in parallel, using both CryoSPARC (v3.1.0, Punjani et al. 2017) and RELION (v3.1, (Zivanov et al. 2018)) software suites to yield similar set of maps. For the sake of simplicity and brevity, only the maps with the highest resolution estimates are reported here. All refinements were performed using independent data half-sets (gold-standard refinement) and resolutions determined based on the Fourier shell correlation (FSC = 0.143) criterion (Rosenthal and Henderson 2003).

Holo-complex

After several rounds of 2D classification in cryoSPARC, particles from 'good' classes were used for an initial *ab initio* reconstruction. The resultant particles were exported to Relion and then re-classified in 2D. Subsequent rounds of 3D classification in Relion yielded 6 classes, of which one (1: HOLO-COMPLEXl; 17,152 particles, representing 13% of the input) provided the highest resolution after 3D refinement with a soft-edged mask and solvent flattening. Post-processing produced a map at 10.8 Å resolution.

Hinge/Arm and Head

Working in parallel, using cryoSPARC, particle subtraction was used to remove the 'head-end' of the complex, before additional rounds of 3D classification, refinement and post-processing yielded a map at 8.53 Å (2: HINGE/ARM; 106,660 particles, 62.4% of input). A similar strategy was used to remove the 'hinge-end' of the complex, however further rounds of processing was halted due to the superior results obtained via the 'head-only' approach described below.

Head-only

A second trained 'picking model' was used to identify particles corresponding to just the

'head-end' of the complex. After several rounds of 2D classification, particles from 'good'

classes were used for an initial *ab initio* reconstruction. After iterative rounds of 3D

classification and refinement, one class was selected to take forward into non-uniform

refinement, after removal of density corresponding to the HALO-tag attached to the Nse4

subunit by particle subtraction (3: HEAD-ONLY; 84,810 particles, 51.7% of input). Post-

processing yielded a map at 6.5 Å resolution.

Composite Map and Model Building

An initial pseudo-atomic model for the Smc5/6-complex was generated by fitting of Phyre2-

generated homology models (Kelley et al. 2015) into map segments using programs of either

the PHENIX software suite (Liebschner et al. 2019) or ChimeraX (Pettersen et al. 2021; Afonine

et al. 2018; Liebschner et al. 2019), with additional manual positioning in either Coot (Emsley

et al. 2010) or PyMOL (The PyMOL Molecular Graphics System, Version 2.3.2, Schrödinger

LLC). Phenix.combine_focused_maps was used to generate a composite map, using each of

the reported maps with the initial model as an alignment reference. The initial homology

models were subsequently replaced with those generated by AlphaFold upon their public

release. The fit of the overall model was optimised by real-space refinement in PHENIX

against the composite map.

Data availability

The two maps used to generate the composite cryo-EM volume have been deposited in the

Electron Microscopy Data Bank (EMDB) with accession codes EMD-13893 (head-end of

complex) and EMD-13894 (hinge and arm-region). Real-space refined coordinates for the

Smc5/6 model have been deposited in the Protein Data Bank (PDB) with accession code PDB-

7QCD. The accompanying composite cryo-EM volume has been deposited with accession

code: EMD-13895.

Experiments in yeast

Yeast strains were generated using the following procedure. Synthetic DNA encoding the

genomic sequence for both WT and mutant versions of each gene were purchased as 'Strings

DNA Fragments' from GeneArt (Thermo Fisher Scientfic, UK). These were cloned into the

vector pAW8-natMX6 (Watson et al. 2008), at the PaeI / SalI restriction sites, through Gibson

assembly via short regions of homology included during synthesis. PCR was then used to

amplify the gene and associated nourseothricin resistance module for introduction into the

endogenous locus of diploid yeast cells, using lithium acetate transformation.

Yeast Strains

Strains were generated from 3 individual haploid isolates, generated from tetrad dissections.

Strain #	Genotype	Reference
2655	*MATalpha [leu2-3 trp1-1 ura3-1 can1-100 ADE2 his3-11,15]; haploid isolate, derived from W303*	(Thomas and Rothstein 1989)
2747 diploid	*MATa/alpha [leu2-3 trp1-1 ura3-1 can1-100 ade2-1/ADE2 his3-11,15 SMC5/smc5-Y961A, W964A]*	This study
2753 isolate 1	*MATa [leu2-3 trp1-1 ura3-1 can1-100 ADE2 his3-11,15 smc5-Y961A, W964A]*	This study
2755 isolate 2	*MATalpha [leu2-3 trp1-1 ura3-1 can1-100 ADE2 his3-11,15 smc5-Y961A, W964A]*	This study
2756 isolate 3	*MATalpha [leu2-3 trp1-1 ura3-1 can1-100 ADE2 his3-11,15 smc5-Y961A, W964A]*	This study
2745 diploid	*MATa/alpha [leu2-3 trp1-1 ura3-1 can1-100 ade2-1/ADE2 his3-11,15 SMC5/smc5-F972A, L978D, L981N]*	This study
2793 diploid	*MATa/alpha [leu2-3 trp1-1 ura3-1 can1-100 ade2-1/ADE2 his3-111,15 NSE1/nse1-F217A, E228R, R242A]*	This study

2794 isolate 1	*MATa [leu2-3 trp1-1 ura3-1 can1-100 ADE2 his3-111,15 nse1-F217A, E228R, R242A]*	This study
2795 isolate 2	*MATa [leu2-3 trp1-1 ura3-1 can1-100 ADE2 his3-111,15 nse1-F217A, E228R, R242A]*	This study
2796 isolate 3	*MATalpha [leu2-3 trp1-1 ura3-1 can1-100 ADE2 his3-111,15 nse1-F217A, E228R, R242A]*	This study

Figures

Molecular images were generated using either PyMOL (v. 2.3.2) or ChimeraX (v. 1.1.1).

ACKNOWLEDGEMENTS

The authors would like to thank Dr. Adam Watson, Dr. Alessandro Bianchi, Prof. Uli Rass,

Prof. Antony Carr [AMC] and Prof. Laurence Pearl [LHP] (University of Sussex) for advice,

discussion, and access to reagents/equipment. We are also grateful to the cryo-EM facility at

the University of Sussex (funded by a Wellcome Trust award enhancement grant

095605/Z/11/A to LHP and the RM Phillips Trust). The work was supported by funding from:

University of Sussex Strategic Development Fund [JB], Wellcome [110047/Z/15/Z; AMC],

Cancer Research UK [C302/A24386; AWO and LHP] and MRC [MR/PO18955/1; JMM and

AWO]. Funding for open access charge: Medical Research Council, University of Sussex Open

Access Team (Library).

Conflict of interest statement: None declared.

Fig. 1. Cryo-EM of the budding yeast Smc5/6 complex

(a) Representative SDS-PAGE gel for the purified Smc5/6 holo-complex. (b) Representative 2D

class averages (side views). Conformational flexibility leads to blurring of density at either the

head (top) or hinge-end (bottom) of the complex. (c) Initial 3D map from cryo-EM at a

resolution of 10.8 Å. (d) Maps from focussed refinement at 8.5Å and 6.5 Å for the indicated

segments of the Smc5/6 holo-complex (d, inset) model obtained by uranyl acetate negative

stain electron microscopy for comparison (Hallett et al. 2021). (e) Representative 2D class

averages (side views) and (f) resultant 3D map at 6.5 Å for the 'head'-end of the complex. The

cryo-EM map has been segmented and coloured with respect to its assigned component (see

associated key for additional detail).

FIGURE 2

Fig. 2. A pseudo-atomic model for the Smc5/6 holo-complex

(a) Overview of the pseudo-atomic model. (a, inset *i*) Schematic showing the overall

architecture of the Smc5/6 complex and selected molecular features. (a, inset *ii*) Expanded

view of the 'Elbow', highlighting the crossover of the coiled-coil 'arms' of Smc5 and Smc6 at

this point. (a, inset *iii*) The first alpha-helix of Nse2 (α1) is situated between the two arms of

the complex. The position of Lys311, a known site of auto-SUMOylation is also indicated (a,

inset *iv*) Expanded view of the interface between the 'Joint' features of Smc5 and Smc5,

involving the two αC3 helices. (a, inset *v*) A short beta-hairpin (amino acids Glu244-Ile254)

protruding from Nse3 is in close proximity to the arm of Smc5. For each inset, the

directionality of the ascending helix (head to hinge) is indicated by a blue or red arrow, for

Smc5 and Smc6 respectively. (b) Comparison of the relative head domain positions in the

cryo-EM structures of budding yeast condensin (PDB: 6YVU), Smc5/6 (this manuscript) and

cohesin (PDB: 6ZZ6); in each, using the head of the κ-SMC as a fixed reference point.

(c) Expanded view for the N-terminal helical domain of Nse4 (aa Lys40-Asp124) bound to the

'arm' of Smc6. (d, left) AlphaFold predicts the presence of an additional helical element (aa

Ser360-Ala372) in the C-terminal domain of Nse4. (d, right). AlphaFold predicts a budding

yeast-specific loop insertion in the NH-RING of Nse1 (aa Glu287-Gln303). Where shown,

sections of density from the composite cryo-EM map are represented by a semi-transparent

molecular surface, shaded in grey. Please also see associated key for additional detail.

Fig. 3. KITES and HAWKS share a common interaction interface that involves the κ-SMC 'W-loop'

(a) Side-by-side visualisation of the κ-SMC head domain from Smc5/6 (left), condensin (middle) and cohesin (right) in complex with their respective kleisin C-terminal domain; Nse4, Brn1 and Scc1. In each case, the interacting partner, whether KITE or HAWK, makes a similar set of interactions with the head domain of the κ-SMC.   (b) Expanded view, showing secondary structure molecular cartoons for each κ-SMC head domain, highlighting the position of conserved amino acids within the 'W-loop' or equivalent (stick representation, carbon atoms coloured cyan) plus aromatic residues within the preceding loop (stick representation, carbon atoms coloured magenta). The ABC-signature motif is additionally highlighted in orange. (c, left) Tetrad dissections. Spores derived from diploid *S. cerevisiae* strains carrying both wild-type allele and indicated mutant allele plus associated NAT-selectable marker (natMX6). Genotypes were confirmed by replica plating of spores on selective media (not shown). (c, right) Multiple sequence alignment, across selected species, showing conservation and consensus of amino acids within the W-loop and preceding region of Smc5 (produced using Jalview 2 with Clustal X colour scheme; (Waterhouse et al. 2009)). (c, top left) Relative positions of the Y and F mutation sets, with respect to the head domain of Smc5 and Nse1 subunit (c, inset) Molecular secondary structure cartoon / surface representation, showing the relative locations of the introduced mutation sets within the Nse1/Smc5 interface. Sets of compound mutations introduced into budding yeast: Smc5-Y = Y961A, W964A; Smc5-F = F972A, L978D, L981N, Nse1-N = F217A, E228R, R242A. Please also see associated key for additional detail.

Fig. 4. A speculative model for dsDNA binding by the Smc5/6 complex

(a) Secondary structure molecular cartoons comparing a docked pose for dsDNA-binding to

the Smc5/6 complex with cohesin in complex with ATP and dsDNA (PDB: 6ZZ6). The trajectory

of bound DNA was obtained by superposition of the docked pose reported in (Zabrady et al.

2016) for the human NSE1/3 hetero-dimer. (b) Model for conformational changes upon DNA

binding. We propose that transition from the apo, non-engaged state (J-state) to ATP-bound,

engaged (E-state) includes an intermediary 'encounter' complex — where dsDNA is first

bound to a positively charged surface/groove (created by the interface between Nse1 and

Nse3) — before a secondary reconfiguration step, triggered by either DNA or ATP-binding,

acts to release the Smc5 'W-loop' from its interaction with Nse1.   This allows stimulation of

ATP turnover by the head domains and repositions the dsDNA such that it is bound with the

expected configuration (i.e., as in cohesin). It is not clear how, or indeed if, ubiquitylation,

SUMOylation or other post-translational modification affects either conformation or ATPase

activity. It is also not known if the presence of Nse2, acts to block binding or transition of

bound dsDNA into the S-K ring (SMC-kleisin) compartment, or if it serves to restrict it to the

J-K / E-K (juxtaposed-heads/kleisin or engaged head-kleisin) spaces. Binding of the Nse5/6

heterodimer blocks the ability of Smc5/6 to turn over ATP (Hallett et al. 2021; Taschner et al.

2021), but it is not known what effect this has on the overall conformation at the head-end

of the complex.

a

DATASET 1   DATASET 2   DATASET 3   DATASET 4
6678 movies   10,413 movies   3212 movies   10,669 movies

CRYOSPARC

Topaz picking

| HOLO-COMPLEX | | HEAD-ONLY |

380,714 particles   824,644 particles

Extract 4.4 Å/px   Extract 2.2 Å/px

RELION

2D classification   2D classification   2D classification
with alignment   without re-centering   with re-centering
134,533 particles   132,800 particles   163,960 particles

Recentre and   2.2 Å/px
re-extract

Ab initio model

| HINGE/ARM | | HEAD |

3D classification   3D classification   3D classification   3D classification
+ refinement   + refinement   + refinement   + refinement

Particle subtraction   Particle subtraction
(remove head density)   (remove hinge/arm density)

6 classes   2 classes   4 classes

23.8% 3.5% 11.4% 28.5% 19.4% 13%   37.6% 62.4%   51.7% 19.5% 12.7% 16.1%

Re-extract 2.2 Å/px   Particle subtraction
                                                                        (remove HALO-tag* density)   *Nse4

3D refinement   3D homologous refinement   3D non-uniform refinement
17,152 particles   106,660 particles   84,810 particles

1   **2**   **3**

b

HOLO-COMPLEX   Final resolution = 10.8 Å

1

Legend:
— No Mask
— Corrected
— Masked
— Phase Randomised

Fourier Shell Correlation vs Resolution, Å
(20.0, 10.0, 6.7, 5.0)

HINGE/ARM   GSFSC Resolution: 8.53Å

2

Legend:
— No Mask (16Å)
— Spherical (13Å)
— Loose (8.8Å)
— Tight (8.6Å)
— Corrected (8.5Å)

Fourier Shell Correlation vs Resolution
(DC, 35Å, 18Å, 12Å, 8.8Å, 7Å, 5.9Å, 5Å)

HEAD-ONLY   GSFSC Resolution: 6.50Å

3

Legend:
— No Mask (7.9Å)
— Spherical (7.4Å)
— Loose (6.9Å)
— Tight (6.4Å)
— Corrected (6.5Å)

Fourier Shell Correlation vs Resolution
(DC, 35Å, 18Å, 12Å, 8.8Å, 7Å, 5.9Å, 5Å)

553   **Supplementary Figure 1**

554   (a) Summary of image-processing workflow, used to determine the structure of the budding

555   yeast Smc5/6 holo-complex. (b) Fourier Shell Correlation (FSC) curves estimating the average

556   resolution of the indicated maps.

SUP. FIGURE 2

a

Nse2
Smc6   Smc5

α1

— crosslink

Author:	Taschner	Gutierrez-Escribano	Yu
Crosslinker:	BS402	BS3	DSSO
Complex:	Holo-complex	Holo-complex + Nse5/6	Holo-complex + Nse5/6
+ ATP:	No	No	Yes

b

Smc5   Smc6

Taschner

BS402

Holo-complex

No

c

90°

	Smc5		Nse1
	Smc6		Nse3
	Nse2		Nse4

6.5   7.5   8.5   9.5 Å

increasing map contour

d

| NSE1_YEAST | NSE1_SCHPO | NSE1_HUMAN | NSE1_MOUSE | NSE1_RAT | NSE1_DROME |
| Glu287-Gln303 | Cys199, Gly200 | Cys207, Gly208 | Cys207, Gly208 | Cys207, Gly208 | Cys200, Ser201 |

Supplementary Figure 2

(a) Selected cross-linking mass spectrometry (XL-MS) data taken from 3 separate studies, supporting interaction of the first helix of Nse2 (α2) with the arm of Smc5. Study, type of crosslinker used, complex composition and nucleotide status are summarised below the molecular graphics. Cross-links were visualised using the PyXLinkViewer plugin (Serrano et al. 2020) for PyMOL (The PyMOL Molecular Graphics System, Version 2.3.2, Schrödinger, LLC). XL-MS data reported in (Taschner et al. 2021), (Gutierrez-Escribano et al. 2020), (Yu et al. 2021). (b) Selected XL-MS data taken from (Taschner et al. 2021) supporting interaction of the two SMC 'joints' in the Smc5/6 holo-complex. (c) Different representations for the 'head-end' of the Smc5/6 complex. Sequentially from left to right: secondary structure molecular cartoon; estimate of local resolution as calculated by ResMap-1.1.4 (Kucukelbir, Sigworth, and Tagare 2014) mapped onto the surface of the composite cryo-EM map; composite map shown at increasing contour levels. See associated keys for additional details. (d) AlphaFold predicts the presence of a budding yeast-specific loop insertion in Nse1 (amino acids 287-303, coloured dark blue).

SUP. FIGURE 3

a

Smc5 Y Y961A W964A

Nse1 N F217A E228R R242A

(WT) W303
#1
#2
#3

24 hours / YP

(WT) W303
#1
#2
#3

48 hrs / 100 mM HU

(WT) W303
#1
#2
#3

48 hrs / 50 J UV

(WT) W303
#1
#2
#3

48 hrs / 40 μM CPT

(WT) W303
#1
#2
#3

48 hrs / 0.005% MMS

b

N F217A E228R R242A

Glu228
Leu978
Phe217
Arg242

Nse1
Smc5
Zn
NH-RING

c

(WT) W303

k = 0.40
dt = 1.7 hours
r² = 0.99

nse1-N isolate 1

k = 0.33
dt = 2.1 hours
r² = 0.99

nse1-N isolate 2

k = 0.32
dt = 2.1 hours
r² = 0.99

nse1-N isolate 3

k = 0.30
dt = 2.3 hours
r² = 0.99

Supplementary Figure 3

(a) Yeast strains carrying the Y-set of Smc5 mutations are not sensitive to a range of genotoxic

agents and have no obvious growth defect. Those carrying the N-set of Nse1 mutations are

mildly sensitive to treatment with camptothecin (CPT) or methyl methanesulphonate (MMS).

Dose, type of treatment and duration of growth are as indicated; where HU = hydroxyurea,

UV = ultraviolet light. The W303 strain is included as a wild-type (WT) control. (b) Molecular

secondary structure cartoon highlighting the position of the mutated amino acid residues in

Nse1 (carbon atoms coloured in orange and shown in 'stick' representation) relative to that

of the Smc5 'W-loop' (coloured in dark grey, with the side chain of Leu978 shown in 'stick'

representation). Please also see associated key for additional details. (c) Representative

growth curves for the 3 isolates containing the N-set of mutations introduced into Nse1. The

W303 strain is included as a wild-type (WT) control. Data are fitted to an Exponential

(Malthusian) grown curve, with k defining a rate constant (slope), dt indicating the calculated

doubling time, and r^2 a goodness of fit parameter.

REFERENCES

Afonine, P. V., B. K. Poon, R. J. Read, O. V. Sobolev, T. C. Terwilliger, A. Urzhumtsev, and P. D. Adams. 2018. 'Real-space refinement in PHENIX for cryo-EM and crystallography', *Acta Crystallogr D Struct Biol*, 74: 531-44.

Alt, A., H. Q. Dang, O. S. Wells, L. M. Polo, M. A. Smith, G. A. McGregor, T. Welte, A. R. Lehmann, L. H. Pearl, J. M. Murray, and A. W. Oliver. 2017. 'Specialized interfaces of Smc5/6 control hinge stability and DNA association', *Nat Commun*, 8: 14011.

Anderson, D. E., A. Losada, H. P. Erickson, and T. Hirano. 2002. 'Condensin and cohesin display different arm conformations with characteristic hinge angles', *J Cell Biol*, 156: 419-24.

Bentley, P., M. J. A. Tan, A. A. McBride, E. A. White, and P. M. Howley. 2018. 'The SMC5/6 Complex Interacts with the Papillomavirus E2 Protein and Influences Maintenance of Viral Episomal DNA', *J Virol*, 92.

Bepler, T., A. Morin, M. Rapp, J. Brasch, L. Shapiro, A. J. Noble, and B. Berger. 2019. 'Positive-unlabeled convolutional neural networks for particle picking in cryo-electron micrographs', *Nat Methods*, 16: 1153-60.

Bermudez-Lopez, M., M. T. Villoria, M. Esteras, A. Jarmuz, J. Torres-Rosell, A. Clemente-Blanco, and L. Aragon. 2016. 'Sgs1's roles in DNA end resection, HJ dissolution, and crossover suppression require a two-step SUMO regulation dependent on Smc5/6', *Genes Dev*, 30: 1339-56.

Burmann, F., B. G. Lee, T. Than, L. Sinn, F. J. O'Reilly, S. Yatskevich, J. Rappsilber, B. Hu, K. Nasmyth, and J. Lowe. 2019. 'A folded conformation of MukBEF and cohesin', *Nat Struct Mol Biol*, 26: 227-36.

Burmann, F., H. C. Shin, J. Basquin, Y. M. Soh, V. Gimenez-Oya, Y. G. Kim, B. H. Oh, and S. Gruber. 2013. 'An asymmetric SMC-kleisin bridge in prokaryotic condensin', *Nat Struct Mol Biol*, 20: 371-9.

Collier, J. E., B. G. Lee, M. B. Roig, S. Yatskevich, N. J. Petela, J. Metson, M. Voulgaris, A. Gonzalez Llamazares, J. Lowe, and K. A. Nasmyth. 2020. 'Transport of DNA within cohesin involves clamping on top of engaged heads by Scc2 and entrapment within the ring by Scc3', *Elife*, 9.

Cutts, E. E., and A. Vannini. 2020. 'Condensin complexes: understanding loop extrusion one conformational change at a time', *Biochem Soc Trans*, 48: 2089-100.

Datta, S., L. Lecomte, and C. H. Haering. 2020. 'Structural insights into DNA loop extrusion by SMC protein complexes', *Curr Opin Struct Biol*, 65: 102-09.

Diebold-Durand, M. L., H. Lee, L. B. Ruiz Avila, H. Noh, H. C. Shin, H. Im, F. P. Bock, F. Burmann, A. Durand, A. Basfeld, S. Ham, J. Basquin, B. H. Oh, and S. Gruber. 2017. 'Structure of Full-Length SMC and Rearrangements Required for Chromosome Organization', *Mol Cell*, 67: 334-47 e5.

Duan, X., P. Sarangi, X. Liu, G. K. Rangi, X. Zhao, and H. Ye. 2009. 'Structural and functional insights into the roles of the Mms21 subunit of the Smc5/6 complex', *Mol Cell*, 35: 657-68.

Emsley, P., B. Lohkamp, W. G. Scott, and K. Cowtan. 2010. 'Features and development of Coot', *Acta Crystallogr D Biol Crystallogr*, 66: 486-501.

Etheridge, T. J., D. Villahermosa, E. Campillo-Funollet, A. D. Herbert, A. Irmisch, A. T. Watson, H. Q. Dang, M. A. Osborne, A. W. Oliver, A. M. Carr, and J. M. Murray. 2021. 'Live-cell single-molecule tracking highlights requirements for stable Smc5/6 chromatin association in vivo', *Elife*, 10.

Fennell-Fezzie, R., S. D. Gradia, D. Akey, and J. M. Berger. 2005. 'The MukF subunit of Escherichia coli condensin: architecture and functional relationship to kleisins', *EMBO J*, 24: 1921-30.

Fousteri, M. I., and A. R. Lehmann. 2000. 'A novel SMC protein complex in Schizosaccharomyces pombe contains the Rad18 DNA repair protein', *EMBO J*, 19: 1691-702.

Gibson, R. T., and E. J. Androphy. 2020. 'The SMC5/6 Complex Represses the Replicative Program of High-Risk Human Papillomavirus Type 31', *Pathogens*, 9.

Gligoris, T. G., J. C. Scheinost, F. Burmann, N. Petela, K. L. Chan, P. Uluocak, F. Beckouet, S. Gruber, K. Nasmyth, and J. Lowe. 2014. 'Closing the cohesin ring: structure and function of its Smc3-kleisin interface', *Science*, 346: 963-7.

Gutierrez-Escribano, P., S. Hormeno, J. Madariaga-Marcos, R. Sole-Soler, F. J. O'Reilly, K. Morris, C. Aicart-Ramos, R. Aramayo, A. Montoya, H. Kramer, J. Rappsilber, J. Torres-Rosell, F. Moreno-Herrero, and L. Aragon. 2020. 'Purified Smc5/6 Complex Exhibits DNA Substrate Recognition and Compaction', *Mol Cell*, 80: 1039-54 e6.

Haering, C. H., D. Schoffnegger, T. Nishino, W. Helmhart, K. Nasmyth, and J. Lowe. 2004. 'Structure and stability of cohesin's Smc1-kleisin interaction', *Mol Cell*, 15: 951-64.

Hallett, S. T., P. Schellenberger, L. Zhou, F. Beuron, E. Morris, J. M. Murray, and A. W. Oliver. 2021. 'Nse5/6 is a negative regulator of the ATPase activity of the Smc5/6 complex', *Nucleic Acids Res*, 49: 4534-49.

Hassler, M., I. A. Shaltiel, and C. H. Haering. 2018. 'Towards a Unified Model of SMC Complex Function', *Curr Biol*, 28: R1266-R81.

Hassler, M., I. A. Shaltiel, M. Kschonsak, B. Simon, F. Merkel, L. Tharichen, H. J. Bailey, J. Macosek, S. Bravo, J. Metz, J. Hennig, and C. H. Haering. 2019. 'Structural Basis of an Asymmetric Condensin ATPase Cycle', *Mol Cell*, 74: 1175-88 e9.

Higashi, T. L., P. Eickhoff, J. S. Sousa, J. Locke, A. Nans, H. R. Flynn, A. P. Snijders, G. Papageorgiou, N. O'Reilly, Z. A. Chen, F. J. O'Reilly, J. Rappsilber, A. Costa, and F. Uhlmann. 2020. 'A Structure-Based Mechanism for DNA Entry into the Cohesin Ring', *Mol Cell*, 79: 917-33 e9.

Jo, A., S. Li, J. W. Shin, X. Zhao, and Y. Cho. 2021. 'Structure Basis for Shaping the Nse4 Protein by the Nse1 and Nse3 Dimer within the Smc5/6 Complex', *J Mol Biol*, 433: 166910.

Jumper, J., R. Evans, A. Pritzel, T. Green, M. Figurnov, O. Ronneberger, K. Tunyasuvunakool, R. Bates, A. Zidek, A. Potapenko, A. Bridgland, C. Meyer, S. A. A. Kohl, A. J. Ballard, A. Cowie, B. Romera-Paredes, S. Nikolov, R. Jain, J. Adler, T. Back, S. Petersen, D. Reiman, E. Clancy, M. Zielinski, M. Steinegger, M. Pacholska, T. Berghammer, S. Bodenstein, D. Silver, O. Vinyals, A. W. Senior, K. Kavukcuoglu, P. Kohli, and D. Hassabis. 2021. 'Highly accurate protein structure prediction with AlphaFold', *Nature*.

Kamada, K., M. Su'etsugu, H. Takada, M. Miyata, and T. Hirano. 2017. 'Overall Shapes of the SMC-ScpAB Complex Are Determined by Balance between Constraint and Relaxation of Its Structural Parts', *Structure*, 25: 603-16 e4.

Kashammer, L., J. H. Saathoff, K. Lammens, F. Gut, J. Bartho, A. Alt, B. Kessler, and K. P. Hopfner. 2019. 'Mechanism of DNA End Sensing and Processing by the Mre11-Rad50 Complex', *Mol Cell*, 76: 382-94 e6.

Kelley, L. A., S. Mezulis, C. M. Yates, M. N. Wass, and M. J. Sternberg. 2015. 'The Phyre2 web portal for protein modeling, prediction and analysis', *Nat Protoc*, 10: 845-58.

Kong, M., E. E. Cutts, D. Pan, F. Beuron, T. Kaliyappan, C. Xue, E. P. Morris, A. Musacchio, A.
Vannini, and E. C. Greene. 2020. 'Human Condensin I and II Drive Extensive ATP-
Dependent Compaction of Nucleosome-Bound DNA', *Mol Cell*, 79: 99-114 e9.
Kschonsak, M., F. Merkel, S. Bisht, J. Metz, V. Rybin, M. Hassler, and C. H. Haering. 2017.
'Structural Basis for a Safety-Belt Mechanism That Anchors Condensin to
Chromosomes', *Cell*, 171: 588-600 e24.
Kucukelbir, A., F. J. Sigworth, and H. D. Tagare. 2014. 'Quantifying the local resolution of cryo-
EM density maps', *Nat Methods*, 11: 63-5.
Kurokawa, Y., and Y. Murayama. 2020. 'DNA Binding by the Mis4(Scc2) Loader Promotes
Topological DNA Entrapment by the Cohesin Ring', *Cell Rep*, 33: 108357.
Lee, B. G., F. Merkel, M. Allegretti, M. Hassler, C. Cawood, L. Lecomte, F. J. O'Reilly, L. R. Sinn,
P. Gutierrez-Escribano, M. Kschonsak, S. Bravo, T. Nakane, J. Rappsilber, L. Aragon, M.
Beck, J. Lowe, and C. H. Haering. 2020. 'Cryo-EM structures of holo condensin reveal
a subunit flip-flop mechanism', *Nat Struct Mol Biol*, 27: 743-51.
Li, Y., K. W. Muir, M. W. Bowler, J. Metz, C. H. Haering, and D. Panne. 2018. 'Structural basis
for Scc3-dependent cohesin recruitment to chromatin', *Elife*, 7.
Liebschner, D., P. V. Afonine, M. L. Baker, G. Bunkoczi, V. B. Chen, T. I. Croll, B. Hintze, L. W.
Hung, S. Jain, A. J. McCoy, N. W. Moriarty, R. D. Oeffner, B. K. Poon, M. G. Prisant, R.
J. Read, J. S. Richardson, D. C. Richardson, M. D. Sammito, O. V. Sobolev, D. H.
Stockwell, T. C. Terwilliger, A. G. Urzhumtsev, L. L. Videau, C. J. Williams, and P. D.
Adams. 2019. 'Macromolecular structure determination using X-rays, neutrons and
electrons: recent developments in Phenix', *Acta Crystallogr D Struct Biol*, 75: 861-77.
Matityahu, A., and I. Onn. 2021. 'Hit the brakes - a new perspective on the loop extrusion
mechanism of cohesin and other SMC complexes', *J Cell Sci*, 134.
Murayama, Y., and F. Uhlmann. 2014. 'Biochemical reconstition of topological DNA binding
by the cohesin ring', *Nature*, 505: 367-71.
Murphy, C. M., Y. Xu, F. Li, K. Nio, N. Reszka-Blanco, X. Li, Y. Wu, Y. Yu, Y. Xiong, and L. Su.
2016. 'Hepatitis B Virus X Protein Promotes Degradation of SMC5/6 to Enhance HBV
Replication', *Cell Rep*, 16: 2846-54.
Palecek, J. J. 2018. 'SMC5/6: Multifunctional Player in Replication', *Genes (Basel)*, 10.
Palecek, J. J., and S. Gruber. 2015. 'Kite Proteins: a Superfamily of SMC/Kleisin Partners
Conserved Across Bacteria, Archaea, and Eukaryotes', *Structure*, 23: 2183-90.
Payne, F., R. Colnaghi, N. Rocha, A. Seth, J. Harris, G. Carpenter, W. E. Bottomley, E. Wheeler,
S. Wong, V. Saudek, D. Savage, S. O'Rahilly, J. C. Carel, I. Barroso, M. O'Driscoll, and R.
Semple. 2014. 'Hypomorphism in human NSMCE2 linked to primordial dwarfism and
insulin resistance', *J Clin Invest*, 124: 4028-38.
Petela, N. J., T. G. Gligoris, J. Metson, B. G. Lee, M. Voulgaris, B. Hu, S. Kikuchi, C. Chapard, W.
Chen, E. Rajendra, M. Srinivisan, H. Yu, J. Lowe, and K. A. Nasmyth. 2018. 'Scc2 Is a
Potent Activator of Cohesin's ATPase that Promotes Loading by Binding Scc1 without
Pds5', *Mol Cell*, 70: 1134-48 e7.
Petela, N. J., A. Gonzalez Llamazares, S. Dixon, B. Hu, B. G. Lee, J. Metson, H. Seo, A. Ferrer-
Harding, M. Voulgaris, T. Gligoris, J. Collier, B. H. Oh, J. Lowe, and K. A. Nasmyth. 2021.
'Folding of cohesin's coiled coil is important for Scc2/4-induced association with
chromosomes', *Elife*, 10.
Pettersen, E. F., T. D. Goddard, C. C. Huang, E. C. Meng, G. S. Couch, T. I. Croll, J. H. Morris,
and T. E. Ferrin. 2021. 'UCSF ChimeraX: Structure visualization for researchers,
educators, and developers', *Protein Sci*, 30: 70-82.

Piazza, I., A. Rutkowska, A. Ori, M. Walczak, J. Metz, V. Pelechano, M. Beck, and C. H. Haering. 2014. 'Association of condensin with chromosomes depends on DNA binding by its HEAT-repeat subunits', *Nat Struct Mol Biol*, 21: 560-8.

Pintilie, G., and W. Chiu. 2012. 'Comparison of Segger and other methods for segmentation and rigid-body docking of molecular components in cryo-EM density maps', *Biopolymers*, 97: 742-60.

Pintilie, G. D., J. Zhang, T. D. Goddard, W. Chiu, and D. C. Gossard. 2010. 'Quantitative analysis of cryo-EM density map segmentation by watershed and scale-space filtering, and fitting of structures by alignment to regions', *J Struct Biol*, 170: 427-38.

Rohou, A., and N. Grigorieff. 2015. 'CTFFIND4: Fast and accurate defocus estimation from electron micrographs', *J Struct Biol*, 192: 216-21.

Rosenthal, P. B., and R. Henderson. 2003. 'Optimal determination of particle orientation, absolute hand, and contrast loss in single-particle electron cryomicroscopy', *J Mol Biol*, 333: 721-45.

Serrano, D., G. Cordero, R. Kawamura, A. Sverzhinsky, M. Sarker, S. Roy, C. Malo, J. M. Pascal, J. F. Marko, and D. D'Amours. 2020. 'The Smc5/6 Core Complex Is a Structure-Specific DNA Binding and Compacting Machine', *Mol Cell*, 80: 1025-38 e5.

Shi, Z., H. Gao, X. C. Bai, and H. Yu. 2020. 'Cryo-EM structure of the human cohesin-NIPBL-DNA complex', *Science*, 368: 1454-59.

Soh, Y. M., F. Burmann, H. C. Shin, T. Oda, K. S. Jin, C. P. Toseland, C. Kim, H. Lee, S. J. Kim, M. S. Kong, M. L. Durand-Diebold, Y. G. Kim, H. M. Kim, N. K. Lee, M. Sato, B. H. Oh, and S. Gruber. 2015. 'Molecular basis for SMC rod formation and its dissolution upon DNA binding', *Mol Cell*, 57: 290-303.

Sole-Soler, R., and J. Torres-Rosell. 2020. 'Smc5/6, an atypical SMC complex with two RING-type subunits', *Biochem Soc Trans*, 48: 2159-71.

Taschner, M., J. Basquin, B. Steigenberger, I. B. Schafer, Y. M. Soh, C. Basquin, E. Lorentzen, M. Raschle, R. A. Scheltema, and S. Gruber. 2021. 'Nse5/6 inhibits the Smc5/6 ATPase and modulates DNA substrate binding', *EMBO J*: e107807.

Thomas, B. J., and R. Rothstein. 1989. 'The genetic control of direct-repeat recombination in Saccharomyces: the effect of rad52 and rad1 on mitotic recombination at GAL10, a transcriptionally regulated gene', *Genetics*, 123: 725-38.

Uhlmann, F. 2016. 'SMC complexes: from DNA to chromosomes', *Nat Rev Mol Cell Biol*, 17: 399-412.

van der Crabben, S. N., M. P. Hennus, G. A. McGregor, D. I. Ritter, S. C. Nagamani, O. S. Wells, M. Harakalova, I. K. Chinn, A. Alt, L. Vondrova, R. Hochstenbach, J. M. van Montfrans, S. W. Terheggen-Lagro, S. van Lieshout, M. J. van Roosmalen, I. Renkens, K. Duran, I. J. Nijman, W. P. Kloosterman, E. Hennekam, J. S. Orange, P. M. van Hasselt, D. A. Wheeler, J. J. Palecek, A. R. Lehmann, A. W. Oliver, L. H. Pearl, S. E. Plon, J. M. Murray, and G. van Haaften. 2016. 'Destabilized SMC5/6 complex leads to chromosome breakage syndrome with severe lung disease', *J Clin Invest*, 126: 2881-92.

Varejao, N., E. Ibars, J. Lascorz, N. Colomina, J. Torres-Rosell, and D. Reverter. 2018. 'DNA activates the Nse2/Mms21 SUMO E3 ligase in the Smc5/6 complex', *EMBO J*, 37.

Waterhouse, A. M., J. B. Procter, D. M. Martin, M. Clamp, and G. J. Barton. 2009. 'Jalview Version 2--a multiple sequence alignment editor and analysis workbench', *Bioinformatics*, 25: 1189-91.

Watson, A. T., V. Garcia, N. Bone, A. M. Carr, and J. Armstrong. 2008. 'Gene tagging and gene
replacement using recombinase-mediated cassette exchange in Schizosaccharomyces
pombe', *Gene*, 407: 63-74.

Wells, J. N., T. G. Gligoris, K. A. Nasmyth, and J. A. Marsh. 2017. 'Evolution of condensin and
cohesin complexes driven by replacement of Kite by Hawk proteins', *Curr Biol*, 27: R17-
R18.

Woo, J. S., J. H. Lim, H. C. Shin, M. K. Suh, B. Ku, K. H. Lee, K. Joo, H. Robinson, J. Lee, S. Y. Park,
N. C. Ha, and B. H. Oh. 2009. 'Structural studies of a bacterial condensin complex
reveal ATP-dependent disruption of intersubunit interactions', *Cell*, 136: 85-96.

Xu, W., C. Ma, Q. Zhang, R. Zhao, D. Hu, X. Zhang, J. Chen, F. Liu, K. Wu, Y. Liu, and J. Wu. 2018.
'PJA1 Coordinates with the SMC5/6 Complex To Restrict DNA Viruses and Episomal
Genes in an Interferon-Independent Manner', *J Virol*, 92.

Xue, X., K. Choi, J. N. Bonner, T. Chiba, Y. Kwon, Y. Xu, H. Sanchez, C. Wyman, H. Niu, X. Zhao,
and P. Sung. 2014. 'Restriction of replication fork regression activities by a conserved
SMC complex', *Mol Cell*, 56: 436-45.

Yatskevich, S., J. Rhodes, and K. Nasmyth. 2019. 'Organization of Chromosomal DNA by SMC
Complexes', *Annu Rev Genet*, 53: 445-82.

Yu, Y., S. Li, Z. Ser, T. Sanyal, K. Choi, B. Wan, H. Kuang, A. Sali, A. Kentsis, D. J. Patel, and X.
Zhao. 2021. 'Integrative analysis reveals unique structural and functional features of
the Smc5/6 complex', *Proc Natl Acad Sci U S A*, 118.

Zabrady, K., M. Adamus, L. Vondrova, C. Liao, H. Skoupilova, M. Novakova, L. Jurcisinova, A.
Alt, A. W. Oliver, A. R. Lehmann, and J. J. Palecek. 2016. 'Chromatin association of the
SMC5/6 complex is dependent on binding of its NSE3 subunit to DNA', *Nucleic Acids
Res*, 44: 1064-79.

Zapatka, M., I. Pocino-Merino, H. Heluani-Gahete, M. Bermudez-Lopez, M. Tarres, E. Ibars, R.
Sole-Soler, P. Gutierrez-Escribano, S. Apostolova, C. Casas, L. Aragon, R. Wellinger, N.
Colomina, and J. Torres-Rosell. 2019. 'Sumoylation of Smc5 Promotes Error-free
Bypass at Damaged Replication Forks', *Cell Rep*, 29: 3160-72 e4.

Zawadzka, K., P. Zawadzki, R. Baker, K. V. Rajasekar, F. Wagner, D. J. Sherratt, and L. K.
Arciszewska. 2018. 'MukB ATPases are regulated independently by the N- and C-
terminal domains of MukF kleisin', *Elife*, 7.

Zheng, S. Q., E. Palovcak, J. P. Armache, K. A. Verba, Y. Cheng, and D. A. Agard. 2017.
'MotionCor2: anisotropic correction of beam-induced motion for improved cryo-
electron microscopy', *Nat Methods*, 14: 331-32.

Zivanov, J., T. Nakane, B. O. Forsberg, D. Kimanius, W. J. Hagen, E. Lindahl, and S. H. Scheres.
2018. 'New tools for automated high-resolution cryo-EM structure determination in
RELION-3', *Elife*, 7.