The reproduction of Gram-negative protoplasts and the influence of

environmental conditions on this process

Authors

Dheeraj Kanaparthi[1,2,3*], Marko Lampe[4], Jan-Hagen Krohn[1,3], Baoli Zhu[2&5], Andreas Klingl[6],

& Tillmann Lueders[2*]

Affiliations

[1]Max-Planck Institute for Biochemistry, Munich, Germany.

[2]Chair of Ecological Microbiology, BayCeer, University of Bayreuth, Germany.

[3]Excellence Cluster ORIGINS, Garching, Germany.

[4]Advanced Light Microscopy Facility, European Molecular Biology Laboratory, Heidelberg,

Germany.

[5]Key Laboratory of Agro-ecological Processes in Subtropical Regions, CAS, Changsha,

China.

[6]Department of Biology, LMU, Planegg-Martinsried, Germany.

*Corresponding author

Keywords: Protoplasts, Protoplast reproduction, Gram-negative Protoplasts, & Minimal

Cells.

Summary

Bacterial protoplasts are known to reproduce independently of canonical molecular biological processes. Their reproduction is shown to be mediated entirely by the physicochemical properties of cell constituents. However, the physiochemical properties of the cell constituents are influenced by the environmental conditions like salinity, salt composition, and mechanical stresses experienced by a cell in natural environments. The influence of such environmental conditions on protoplast reproduction is seldom investigated. Here, we studied protoplast reproduction in their native environmental conditions. Contrary to the previous perceptions of protoplasts reproducing in an erratic manner, cells in our study reproduced in a defined sequence of steps. The process of their reproduction can be explained by an interplay between intracellular metabolism, the physicochemical properties of cell constituents, and the nature of cations in the growth media. We observed a minimal leakage of intracellular constituents during protoplast reproduction, suggesting an efficient reproduction. However, the efficiency of reproduction is determined by the environmental conditions. Under favorable environmental conditions, protoplasts reproduce with nearly similar efficiency to cells that possess a cell wall. In short, here we demonstrate the simplest method of cellular reproduction and the influence of environmental conditions on this process.

Introduction

Bacterial species typically possess a peptidoglycan cell wall, which imparts structural rigidity and forms the foundation of their morphology[1]. In addition to its structural functions, the cell wall plays a vital role in maintaining proper turgor pressure, transporting nutrients into the

cell, and facilitating reproduction [2]. However, bacteria are known to lose their cell wall and

transform into their protoplast state when exposed to cell wall-degrading enzymes and

bactericidal antibiotics [3,4]. Although this transformation is thought to make cells susceptible

to changes in environmental conditions like osmolarity [5], it is also known to confer some

advantages to the bacteria [3,6,7]. Most of the antibody-binding sites are located on the surface

of the cell wall [8,9]. When these cells transition to their protoplast state, they avoid being

detected by the immune system of the host cell, facilitating the unchecked proliferation of

pathogens within the host [10]. Given these clinical implications, most research on protoplasts

primarily focuses on understanding either the clinical implications of this transformation [11–13]

or the intracellular processes within the protoplasts [14,15].

In contrast to our conventional understanding, some recent studies showed that the

transformation of bacteria into their protoplast state could happen in natural environments

without antibiotic stress [6,16]. This transformation is observed to have been induced by

environmental conditions like osmolarity and exposure to lytic phases [6,16]. This

transformation of cells into the protoplast state is shown to confer cells with phase resistance,

increased survival rates in natural environments, and receptivity to incorporating DNA from

the surrounding environment [6,7,17]. Apart from the well-documented clinical implications of

bacterial transformation into protoplasts, these studies imply the ecological benefits of such

transformation [6,16,17]. However, we currently lack a good understanding of the ecological

aspects of this cell state, especially how the environmental conditions influence the protoplast

life cycle.

Protoplasts are known to reproduce independent of canonical molecular biological processes

[14]. Reproduction in such cells is shown to have been primarily mediated by the

physiochemical properties of the cell constituents, like the membrane fluidity and ratio of cell

volume to surface area [15,18–20]. The physicochemical properties of cell membranes, like the

fluidity and spontaneous curvature, are influenced by external conditions like osmolarity, the

nature of the cations, and the mechanical strain experienced by a cell in natural environments

[21–24]. This suggests that conditions surrounding the cell exert a considerable influence on the

morphology and reproduction of protoplasts. Despite this possibility, much of our current

understanding of the protoplasts is a result of studying them under well-controlled

environmental conditions rather than in their native environment [4,14]. In this study, we

examined protoplast reproduction in their native environment and how various environmental

conditions influence this process.

The cytoplasmic osmolarity of a non-halophilic bacteria ranges between 300-500 mosM [25,26].

This was a result of the densely packed intracellular environment with proteins and nucleic

acids. Maintaining this high intracellular osmolarity is necessary for the cells to absorb water

and nutrients from the surroundings [27]. One essential function of the rigid peptidoglycan cell

wall is to counteract the uncontrolled expansion of cytoplasm due to its high osmolarity [28].

Given that the protoplasts lack a cell wall, they are prone to osmotic lysis [5]. Hence it is

necessary to grow them in an isotonic media [14,18], and their distribution in natural

environments seems restricted to mildly saline environments [11–13,17]. To emulate these

conditions, we grew the bacterial protoplasts in media containing 7%Dead Sea salt (DSS) at

30°C. DSS was used in our incubations rather than pure NaCl to replicate the complex salt

composition of the natural environments. In the below sections of the manuscript, we present

the lifecycle of protoplasts under environmental conditions close to their native habitat.

Results

Generation and physiological characterization of protoplasts

We transformed *R. sphaeroides (RS)* into its protoplast (*R. sphaeroides-protoplast, RS-P*)

state by the procedure described in the methods section. This transformation was evident

from the change in the morphology of cells from coccoid to relatively bigger spherical cells

(Figure 1). The absence of the cell wall was also confirmed by transmission electron

microscopy (TEM) (presented below). *RS-P* did not transform back into its native state when

the use of lysozyme and penicillin was discontinued. But a step-by-step reduction of salt

content in the media or gradually replacing salt with an equivalent concentration of sucrose

led to the transformation of cells into their native state with a cell wall. No reversal of *RS-P*

to native-*RS* state was observed in control incubations where the salt concentration is

consistently maintained at 7% or in sugar-rich media with lysozyme and penicillin.

Substantial differences in the physiological behavior were observed between *RS-P* and

"wildtype" *R. sphaeroides*. *RS-P* exhibited similar superoxide dismutase (SOD) activity

under aerobic (14.8 ± 1.3 U/ml) and anaerobic (15.2 ± 2.1 U/ml) conditions. In contrast, the

SOD activity of WT *R. sphaeroides* depended on environmental conditions. Considerably

lower cytoplasmic SOD activity was observed under aerobic growth conditions (8 ± 0.04

U/ml) compared to *RS-L*. Negligible to no SOD activity was observed when *R. sphaeroides*

was grown in anaerobic media (0.16 ± 0.05 U/ml).

Fermentation products like lactate, butyrate, and acetate were observed when *RS-P* was

grown in well-aerated media containing glucose (Figure S1). Such volatile fatty acid

accumulation was not observed in *R. sphaeroides* incubations under aerobic conditions

(Figure S1). All the experiments were replicated (at least n=5), as indicated in the individual

methods sections. The lifecycle and other data presented below were repeated numerous

times over the course of this work.

The life cycle of *RS-P* in 7%DSS-TSB.

Once we obtained stable protoplasts, we transferred them into fresh growth media containing

7% Dead Sea salt (DSS). We then monitored these cells at regular intervals using various

microscopic techniques. The lifecycle of these cells is shown in Figure 1.

During the lag phase, growth was restricted to the outer membrane of the cell, which

expanded to form a hollow filamentous extension (Figure 1A-C, 2 & S2). The length of these

extensions varied from cell to cell, ranging between 0.5 to 5 µm (Figures 1A-C, 2 & S2).

Most cells in this growth stage were observed to have developed hollow membrane vesicles

within their cytoplasm and in the periplasmic space between the outer and inner membranes

(Figure 2). These vesicles initially appeared as tiny buds attached to the inner cell membrane

(Figure 2D, arrow), followed by their expansion into or out of the cytoplasm (Figure 2D-F).

Figure 1. The lifecycle of *RS-P*

All panels show STED microscopy images of cells stained with FM^{TM}5-95 (membrane,

white) and PicoGreen^{TM} (DNA, red). White arrows guide the eye in following the life cycle

progressing from A to J. Arrows (yellow) in A highlight a cell with outer membrane

extensions (are also seen in B & C). The steps involved in forming outer membrane

extensions are shown in Figure 2. Boxed regions in G – I highlight cytoplasmic

compartments within the filamentous cell undergoing binary fission (in sequence from G-I).

J: Individual daughter cells with membrane overhangs formed from the fragmentation of

filamentous cells. Also, see Figure S2 for phase-contrast images. Scale bars: 2μm (A-G & J)

and 10μm (H & I).

Figure 2. TEM images of early growth stage *RS-P* cells.

Images A – C show cells with folded excess membrane within the cell (highlighted regions).

Image D shows *RS-P* cells with intracellular vesicles. The arrow in D points to the inner

membrane undergoing budding into the periplasmic space. A cell with hollow vesicles in the

periplasmic space was shown in E. Image F shows the magnified periplasmic region of the

cell in E. Images G – I show cells with the elongated outer membrane. Image J shows the

cytoplasmic compartment's growth, filling in the void left by the outer membrane (arrow).

Scale bars: 0.5μm (A-J).

During the early-log phase, with an increase in the growth rate (Figure 3), the volume of the

cytoplasm increased, gradually filling the hollow template created by the outer membrane

(Figure 1C&D, 2J, and S1A-C). This led to the transformation of cells in sequence from

spherical to tubular (Figures 1, 3, and S2), then to dumbbell-shaped cells with cytoplasm

mostly restricted to either end of the cell (Figure 3, early log-phase). In the subsequent

growth stages, cells developed surface depressions along their length resembling tubular

vesicles subjected to curvature-induced destabilization at their surface (resembling Rayleigh-

Plateau instability) (Figures 1D-G, 3A, & S3). These surface depressions were formed due to

the cytoplasmic compartment surrounded by the inner membrane undergoing binary fission

(Figure 1D-H & G-I).

Over time, these surface depressions grew deeper, and two adjacent cytoplasmic

compartments were linked to each other only by a hollow strand of the outer membrane

(boxed regions in Figures 1H&I and 3 – late log phase). The rupture of the outer membrane

connections led to the formation of individual daughter cells (Figure 1J). This sequential

transformation of cells was not observed when we repeatedly transferred log-phase cells into

fresh media every few hours to keep them in a prolonged log phase. Cells in these

experiments grew into longer and thicker filamentous cells without developing surface

deformations (Figure S5) or transforming into a "string-of-beads" morphology. The

cytoplasm of these cells was observed to have multiple intracellular vesicles (Figure S5C-E).

In the stationary growth phase (5-8 days from the start of the incubation), cell morphologies

were a mix of all the above-described morphologies and a considerable amount of membrane

debris (Figure S5).

Figure 3. *RS-P* cells in different growth stages and comparison of their morphology with

LVs. Image A shows the growth curve and morphologies of *RS-P* in different growth stages

(individual plots represent multiple repetitions, n=6). Cells were stained, imaged, and color-

coded, as in Figure 1. Closed and open arrows point to cytoplasmic compartments and hollow

outer membrane connecting the cytoplasmic compartments, respectively (Figure 4). Images B

and C show theoretically predicted and experimentally observed morphologies of vesicles

subjected to elongational stress over time (t) (images B & C were originally published by

Narasimhan et al., 2015 [29], reproduced here with permission from Cambridge University

Press). A similar comparison of *RS-P* cells with LVs undergoing expansion is shown in

Figure S3. Scale bar: 1μm (lag & early log-phase), 10μm (late log-phase).

Figure 4. Membrane phase separation in *RS-P*. Cells in images A, C, D, F, H, I, & J are

stained with FM^{TM}5-95 (all membrane, red) and *FAST*^{TM} DiI (L$_d$-membrane, green). A & B:

confocal and TEM images of early growth stage cells. Arrows in image A point to regions of

the cell with L$_d$-membrane. Arrows in image B point to similar intracellular vesicles. C - G:

confocal and TEM images of later growth stage cells. Arrows in C & B point to the

cytoplasmic compartment, exclusively enclosed in the L$_d$-membrane. The Arrow (red) in E

point to the vesicles in periplasmic space. The arrow (red) in image F shows similar

periplasmic vesicles composed of L$_o$-membrane. H: Late log-phase filamentous *RS-P* cells

(see Figure S7 for multiple optical sections of this stack). I & J: Magnified regions of the

boxed region in H in individual L$_o$ & L$_d$-membrane channels. K: *RS-P* cell similar in

morphology to the cell shown in I. The cell in this image is stained with membrane and DNA

stain, as in Figure 1. A comparison of cells in H & K shows the inner cell membrane enriched

in L$_d$-membrane. Scale bars: 1µm (A, C, D, & F), 500nm (B, E & G), and 10µm (H & K).

Membrane phase separation in *RS-P*

Staining *RS-P* cells with both universal membrane dyes (FM^{TM}5-95) and a dye specific for

the liquid-disordered membrane (L$_d$) (*FAST*^{TM}DiI) showed the existence of two distinct

phases within the cell membrane: liquid-ordered (L$_o$, red) and liquid-disordered (L$_d$, green)

phases (Figures 4, and S7). During early growth stage, the cell membrane underwent phase

separation, and both L$_d$ and L$_o$-membrane underwent invagination to form hollow buds. All

the buds extending out of the cytoplasm into the periplasmic space were exclusively

composed of the L$_o$-membrane (Figure 4F, arrow), and buds extending into the cytoplasm

were composed of the L$_d$-membrane (Figure 4A-E). Over time we observed a gradual

enrichment of a more fluid L$_d$-phase within the inner cell membrane and a more rigid L$_o$-

phase in the outer cell membrane (Figures 4 and S7).

Influence of environmental conditions on *RS-P*'s morphology

To understand the influence of individual salts on the morphology of *RS-P*, we repeated the

above experiments replacing DSS in the growth media with an equivalent concentration of

either only divalent or only monovalent salts like MgCl$_2$ or KCl/NaCl. Cells grown under

these conditions exhibited morphologies similar to those grown in DSS (Figure 5). However,

considerable differences were observed in their size – considerably thicker and longer

filamentous cells (Figure 5A-D) and considerably smaller individual cells (Figure 5E-H)

were observed when cells were grown in media containing 7%KCl and 7%MgCl$_2$,

respectively. Thick tubular or long filamentous cells with diameter >1-2μm were rarely

observed in incubations with MgCl$_2$, often only transiently during the log phase (Figure

5F&G). When grown on an orbital shaker, hardly any filamentous cells were observed in our

incubations (Figure 5I-L). Instead, most cells during all growth stages are unicellular or cells

undergoing binary fission. Filamentous cells were rarely observed (Figure 5I-L). Treatment

of *RS-P* with 5mM of 3-methoxybenzamide [30] did not result in any apparent morphological

changes compared to the control incubation (Figure S8).

Figure 5. Influence of salt composition on *RS-P*'s morphology. Images A – D shows *RS-P*

cells grown in 7% KCl-NB from early to late growth stages. Images E – H show *RS-P* cells

grown in 7% MgCl₂-NB from early to late growth stages. Images I – J show *RS-P* cells

grown in 7% DSS-NB on an orbital shaker. Arrows in image L point to cells undergoing

binary fission. Cells were stained, imaged, and color-coded, as in Figure 1. Scale bars: 10

μm.

Reproductive efficiency and viability of *RS-P* daughter cells

The concentration of extracellular DNA during the stationary growth phase was used as a proxy to determine the leakage of intracellular constituents during reproduction. The results of this quantification are presented in Figure 6. The concentration of extracellular DNA in all our incubations was an order of magnitude lower than the intracellular DNA. Quantifiable but no significant differences in the concentration of extracellular DNA was observed between *RS-P* and the native *R. spheroides* cells, which possess a well-functioning molecular biological regulatory process (Figure 6A). Neither were significant differences observed between cells grown in media containing different salts. The reproductive efficiency was marginally higher when *RS-P* was incubated on an orbital shaker rather than under static conditions (Figure 6B). Staining and imaging of the stationary growth phase incubations also suggest the absence of extracellular DNA (Figures 5, 6C, and S6).

The metabolic viability of *RS-P* daughter cells was determined by staining daughter cells from the stationary growth phase with dyes specific for cell membrane, DNA, and cytoplasmic activity (Figure 6D). Quantification of live cells (cells with intact membrane, intracellular DNA, and cytoplasmic activity) was done by flow cytometry. Our results suggest that most of the daughter cells ($< 70\%$) received DNA from the parent cell and are physiologically active ($< 80\%$) with intracellular enzyme activity (Figure 6E). Subsequent transfer of daughter cells of different ages – 10, 20, or 100-day-old cultures either directly or after passing through a 0.45µm filter (to separate daughter cells from larger parent cells and membrane debris) into a new media always resulted in growth. Transfer of daughter cells into minimal salt media[31] containing glucose as a sole carbon and energy source resulted in growth (Figure S1).

Figure 6. Reproductive efficiency and viability of *RS-P* daughter cells. Graphs A & B show intracellular (blue bars) and extracellular (orange bars) DNA concentrations (n=5 for each condition). A: *RS-P* grown under static conditions. B: *RS-P* grew on an orbital shaker. Annotations: WT (wild type, *R. sphaeroides* with a cell wall grown in NB); DSS, KCl & MgCl$_2$ – *RS-P* grown in NB with respective salts. Image C shows *RS-P* cells grown in 7%DSS-NB in the late growth stage (Scale bars: 10 µm). Cells in this image are stained, as described in Figure 1. D: Schematic workflow of the experiment to quantify live daughter cells. Cells were stained with different combinations of FM^{TM}5-95 (cell membrane), PicoGreen^{TM} (PG, DNA), and CellTrace^{TM} Violet (CTV, cell integrity) (n=5, in the figure but multiple repetitions over the course of the work). E: The percentage of *RS-P* daughter cells

identified positive for different viability markers in flow cytometry. The original flow

cytometry plots are shown in Figure S9.

Discussion

The above-described life cycle of *RS-P* could be explained by the lack of molecular

biological coordination between physiological processes within the cell, the biophysical

properties of its cell constituents, growth rate, and environmental conditions. This loss of

intracellular coordination between cellular processes is evident from the excessive lipid

synthesis, production of volatile fatty acids from under aerobic conditions, and high

intracellular SOD while fermenting glucose. In a normal functioning cell, the rate of lipid

synthesis is tightly regulated by its growth rate [32]. In contrast, *RS-P* was in a perpetual state of

excess lipids. This is evident by the excessive membrane within *RS-P* during all growth

stages (Figures 1, 2B, 2C, S2A, and S4).

The flow of electrons in the electron transport chain is known to produce reactive oxygen

species (ROS) [33]. Cells were known to synthesize catalase and SOD enzymes to prevent

ROS-related oxidative damage. Such enzymes are selectively expressed only during the

availability of terminal electron acceptors and were repressed during growth by the

fermentation [33]. High and low cytoplasmic SOD activity of *RS,* when grown under aerobic

and fermentation conditions, aligns with our current understanding of SOD-related gene

expression. In contrast, *RS-P* exhibited high cytoplasmic SOD activity irrespective of its

physiology or the availability of terminal electron acceptors. Moreover, we observed the

accumulation of volatile fatty acids like lactate, butyrate, and acetate in growth media when

RS-P was grown under well-aerated conditions (Figure S1). Approximately 30-35% of the

glucose was converted to fermentation products. This transformation of glucose to volatile

fatty acids happens only under anaerobic conditions when the growth media lacks terminal

electron acceptors, and the formation of such compounds was not expected during aerobic

growth. No fermentation products were observed when *RS* was cultured under aerobic

conditions (Figure S1). These differences in the physiological behavior between *RS-P* and *RS*

also suggest the loss of mechanisms to sense the environmental conditions or the loss of

molecular biological coordination required to adapt cellular physiology to the ecological

conditions. A loss of molecular biological coordination leading to chaotic behavior of cells

[4,14,20], like the excessive lipids synthesis [34], and lack of ability to assemble a peptidoglycan

cell wall, despite upregulation of the genes for synthesis of a cell wall components [35], was

previously reported in protoplasts. Although our observations suggest a lack of coordination

within the cytoplasmic processes and are in tune with the previous studies [4,34,35], further

substantiating this claim requires an in-depth molecular biological investigation, which is

beyond the scope of understanding the protoplast reproduction in natural environments.

Morphological features observed in *RS-P,* like the outer membrane extensions, hollow

vesicles within the cells, and transformation of tubular cells to a string-of-beads morphology

(Figure 3), can be explained as a mechanism of accommodating this excess membrane.

Transfer of lipids synthesized within the cytoplasm to the outer membrane and its subsequent

expansion into a hollow filamentous structure (Figures 1&2) happened by a two-step process.

In the first step, disproportionately high lipid synthesis compared to cytoplasmic volume

increase (low growth rate during lag-phase) facilitated the transformation of the excess

membrane into a bud (Figure 2D-F). Subsequently, these hollow buds detached from the

inner membrane and were released into the periplasmic space (Figure 2E-F). In step two,

these hollow vesicles fused and incorporated into the outer membrane, expanding into a

hollow filamentous structure (Figures 1B & 2D-J).

The formation of these buds and associated membrane phase separation (Figure 4) can be

explained by the model proposed by Lipowsky [36]. It was hypothesized that when membranes

phase-separate, more fluid L_d-membrane spontaneously undergoes out-of-plane invagination

and transforms into a bud to reduce line tension at the phase-boundary [37]. The transformation

of the L_d-membrane into intracellular vesicles (Figure 4) observed in our study is in tune with

this hypothesis. Nevertheless, apart from a more fluid L_d-membrane, we also observed L_o-

membrane with high bending rigidity and low spontaneous curvature undergoing budding

(Figure 4F). The formation of buds by the L_o-membrane can be explained by the chemical

composition of the cell membrane [38]. In contrast to thermodynamic considerations, recent

studies attempting to understand the behavior of cell membranes demonstrated the possibility

of a more rigid L_o-membrane undergoing budding. This selective invagination of L_o-

membranes was shown to have been induced by specific sterols within the membrane [38]. In

accordance with these studies, Gram-negative bacteria like *RS-P* and, in particular, anoxic

phototrophs were known to have a diverse sterol composition in their plasma membrane [39],

which could have played a role in the budding of L_o-membranes.

Apart from the chemical composition, the nature of the membrane (L_o or L_d) and the direction

of its budding (into or out of the cytoplasm) could also have been determined by the

differences in the macromolecular crowding between the cytoplasm and the periplasmic

space [22,40]. Previous studies attempting to understand budding and asymmetric cell division in

aqueous two-phase LVs observed that phase separation of the aqueous core induced

membrane phase separation and preferential association of L_d & L_o-membrane with the

aqueous phases of higher and lower densities, respectively. This preferential association

between the membrane and aqueous phases was explained by the interaction between the

membrane and the intracellular constituents [41,42]. Such interactions were shown to alter the

spontaneous curvature of the membrane and the direction of its budding [41,42]. Our observation

of membrane phase separation and subsequent budding in *RS-P* is in tune with these studies.

L_d-membrane in *RS-P* was always observed to have been in association with densely packed

cytoplasm (~17-35 wt% molecules), always budding into the cell and whereas L_o-membrane

preferentially invaginating out of the cell into less densely packed periplasmic space (~6-7

wt% molecules) (Figure 4). These differences in the packing densities between the cytoplasm

and the periplasmic space can also be inferred from the TEM images (Figure 2) [43].

Biophysical properties of the cell membrane and its phase separation into L_d & L_o phases was

proposed to have played an essential role in the protocell reproduction [40,44]. Nevertheless,

such studies were primarily restricted to LVs (Figure 4). To our knowledge, ours is the first

study to test these theories on self-replicating live cells. Unlike the studies that reported

membrane phase separation in live cells[45], we propose that the phase separation in *RS-P* was

due to a passive reorganization of membrane components to reach a thermodynamically

stable state rather than a functional membrane phase separation induced by an intracellular

process or linked to a specific cellular function [45,46].

Morphologies of *RS-P* observed during log and stationary growth stages can also be

explained from a physicochemical perspective. Growth (or expansion) of the cytoplasmic

compartment within the cell, gradually filling the void left by the outer membrane, resembles

LVs undergoing stretching when placed in an elongational flow path (Figures 2 & S3) [29].

Spherical LVs experiencing a low elongational strain were shown to transform into prolate or

tubular shapes[47] (Figures 3B, 3C & S3). However, at a certain strain level (threshold strain),

vesicles transformed into a dumbbell shape. Upon further strain increase, vesicles formed

elongated dumbbells with beaded structures appearing along their length. *RS-P* exhibited a

similar pattern of morphological transformation (Figure 3 & S3). Like in LVs, morphologies

of *RS-P* can be explained by the physical properties of the cell membrane, like its elasticity

and expansion strain exerted on the membrane [29,47] due to cytoplasmic volume increase. In

the case of *RS-P*, high and low elongational strains on the membrane translate into high and

low growth rates (i.e., rate of cytoplasmic volume increase). In accordance with this

hypothesis, short prolate or tubular cells were most frequently observed during the early lag

phase, when the cell's growth rate was low compared to other growth stages (Figure 3). With

an increase in their growth rate during the mid/late-log phase, *RS-P* transformed into

elongated dumbbells and, subsequently, dumbbell-shaped cells with beaded structures along

their length (Figure 3).

The morphologies of *RS-P* observed in our study were similar to the LVs undergoing an

osmotic deflation [48,49]. When placed in a hyperosmotic solution, LVs undergo an osmotic

deflation[50] and transform into a plethora of morphologies, to attain a thermodynamically

stable state. According to Area Differential Elasticity (ADE) theory [48,49], the end morphology

of such vesicles is determined by the ratio of their surface area to volume. Although not

undergoing osmotic deflation, *RS-P* is in a similar physical state due to excess membrane

(Figure 2), i.e., high surface area: volume. However, in the case of *RS-P*, the surface area-to-

volume ratio is everchanging and is determined by the growth stage (Figure 3). The

morphologies of *RS-P* within a given growth stage and the change in its morphologies from

lag, log, to stationary growth phases can be explained by the changing surface area-to-volume

ratio. During the lag phase, the slow growth rate (rate of cytoplasmic volume) resulted in the

formation of outer membrane extensions to accommodate the excess membrane.

During the early and mid-log phases, the cytoplasmic volume increase kept pace with the

surface area increase of the cytoplasmic compartment, i.e., low surface area-to-volume ratio.

This resulted in tubular cells with smooth surfaces or slight surface depressions (Figure 3A

and S4). Cells with outer membrane extensions were rarely observed in the mid-log phase.

Nevertheless, we do not presume this low ratio was due to the reduced rate of lipid synthesis.

Closer observation of cells showed the presence of lipid droplets or hollow intracellular

vesicles within the cytoplasm of the cells (Figure S4). Hence, we propose that the low surface

area-to-volume ratio was due to the accommodation of excess lipids within the cytoplasm and

the slow transfer of these lipids to the outer cell membrane. The extensive formation of

intracellular vesicles also contributed to the rapid cytoplasmic volume increase in log-phase

cells. A gradual decrease in the growth rate by the end of the log phase reverted cells to a

state of excess outer membrane (Figure 1H&I). This transfer of lipids to the outer membrane

transformed the cells from filamentous into a "string-of-beads" morphology.

A similar accumulation of lipids within the cytoplasm was observed when cells were kept in

a constant state of logarithmic growth (Figure S5). Cells kept in a prolonged log phase

gradually increased in length and diameter (Figure S5) but did not transform into "string-of-

beads" morphologies. Such a transformation in these cells was only observed when their

growth rate was reduced by stopping the transfers of these cells into fresh media. These

results suggest that the growth rate had an influence on the morphology of the cells, and a

faster rate of cytoplasmic volume increase reduced the rate of lipid transfer from the

cytoplasm to the outer membrane. In spite of these observations, precise measurement of cell

volume or the amount of excess membrane is technically challenging due to the lack of

uniformity in cell morphology (Figure S10- S12). Moreover, from the mid-log phase, the

cytoplasm of most cells had numerous intracellular vesicles (Figure S12). No uniformity was

observed in the size, number, and morphology of these vesicles (Figures S4, S5, and S10-12).

Cell morphologies within any given growth stage were seldom uniform and, in general, were

a mix of morphologies (Figure S10- S12). The length and width of the cells differed

considerably from one incubation to another, even when cultivated under the exact same

media composition. These morphological variations could be explained by the physiological

differences among the cells within a population. Bacterial growth curves and their partition

into different stages (lag, log & stationary), in general, represent the growth dynamics of the

bacterial population. Nevertheless, averaging the data obscure considerable variation in the

behavior of individual cells [51,52]. It is known that genetically identical cells within a single

bacterial population could exhibit considerable cell-to-cell variation in gene expression [53,54]

and growth rates [55]. The reason for such stochastic behavior of monoclonal cells has to date,

not been well understood. Manifestation of these variations in the morphology of normal

bacteria was not observed due to the presence of a rigid cell wall. Given the absence of a cell

wall in *RS-P*, we assume such cell-to-cell variations in growth rate were manifested in its

morphology (Figure S10- S12).

The morphological unpredictability of *RS-P* could also be explained from a bio-physical

perspective. Recent studies attempting to understand similar unpredictability in LVs

suggested that the initial aspect ratio (width: height) of the vesicle determines its end

morphology[56,57]. An ever-changing aspect ratio of the vesicles resulted in unpredictable end

morphologies. During the initial growth stages, we observed *RS-P* in a constant state of

oscillation (ever-changing aspect ratio) (Video S1). We presume these oscillations could have

originated within the cell due to asymmetric conditions on either side of the membrane [21] or

random movement of cytoplasmic constituents (Video S2), or due to the presence of an

internal metabolism [58,59]. It was also hypothesized that internal metabolism could lead to

shape instabilities on the cell surface ultimately leading to cell division [58].

A gradual reduction in the salt concentration of the media while maintaining a similar

osmolarity with sucrose led to the transformation of *RS-P* back to its native form. This

suggests that DSS played an essential role in stabilizing the protoplast state and preventing

the formation of peptidoglycan cell walls [60]. This also suggests that cells can reversibly

transform from a stable protoplast state to their native state or *vice versa*, depending on the

environmental conditions. Morphological differences between cells grown under different

salt conditions (Figure 5) can be explained by the interaction between phospholipid

membranes and cations [24]. It was reported that divalent cations like Mg^{+2} organize lipids into

clusters of 10-15 molecules each [61], dehydrating and stiffening the membrane and hindering

its area expansion. On the other hand, monovalent cations like K^+ reduce the bending rigidity

of the membrane, making the membrane more flexible [21], conducive to the formation of

larger cells. The absence of filamentous cells when *RS-P* was grown on a shaker (Figure 5I-

L) was likely due to the constant shearing of long filamentous cells. As changes in cell

morphology were reversible, when we cells were transferred between growth media

containing different salt compositions, suggests that cell morphology was determined by the

environmental conditions rather than the information encoded in its genome. In tune with the

earlier studies that canonical structural proteins like FtsZ plays no role in regulating

protoplast morphology [14], we did not observe an influence of FtsZ inhibitors on the lifecycles

of *RS-P* (Figure S8).

During the late log phase, most *RS-P* cells are individual daughter cells formed by the fission

of long filamentous cells. In the case of minimal cells or LVs, the proxy minimal cells, some

earlier studies have hypothesized that the physiochemical forces of the environment could

mediate such fission. Hence to achieve division, these studies employed methods like

extrusion of LVs through a narrow pore size filter or application of an external force [62–64], to

simulate the physical forces faced by such cells in the natural environments. Given that all

our experiments are conducted under static conditions, we do not presume such forces could

have contributed to the reproduction in *RS-P*. We instead propose that the force required to

mediate fission was intrinsic to the cell. Recent studies showed that an internal metabolism or

Brownian movement of intracellular constituents [65,66] could result in surface instabilities

leading to cell division [58,66]. In support of this presumption, cells in our incubations were

intrinsically unstable (Video S1, early growth phase cells with expanded outer membrane).

They were in a constant state of oscillation, which should have resulted from the Brownian

movement of intracellular constituents [65] (Video S1 & movement of DNA in spherical *RS-P*

cells in the top half of Video S2). This constant movement of intracellular constituents could

have led to the generation of force required for the fission of daughter cells in *RS-P* [66]. In

support of this presumption, we observed that cells within a filament were in a constant pull-

and-push motion, which could have resulted in cell fission (Video S3 and Figure 1H-J).

The absence of intracellular constituents like DNA in the growth media suggests that loss of

molecular biological coordination within the cells did not considerably affect the

reproductive efficiency (Figure 6). Nevertheless, we do not presume *RS-P's* reproduction was

efficient. A considerable amount of membrane debris (Figures S6) and large clumps of cells

in the stationary growth phase cultures (Figure 6C) suggest a lower reproductive efficiency

than the cells in their original cell state. These clumps could have formed by the

entanglement of filamentous cells (Figure S13). Observation of these clumps at regular

intervals after their formation showed no evidence of them forming individual daughter cells

and, over time, transforming into cell debris (Figure S13). Despite the wastefulness of *RS-P*'s

reproduction in our static incubations, most daughter cells received DNA from the parent

cell. They grew into filamentous cells upon transfer to fresh media, suggesting that this

method of reproduction always resulted in metabolically viable daughter cells (Figure 6).

Given the absence of long filamentous cells and large cell clumps when grown on an orbital

shaker (Figure 5I-L), we presume *RS-P* reproduced more efficiently under more turbulent

environmental conditions (Figure 6E). Cells in natural environments are more likely to

experience a similar turbulent force than being in a static state. The growth of *RS-P* daughter

cells in media with glucose as the only carbon and energy source suggests that daughter cells

received a full complement of genes from the parent cell to maintain an autonomous growth,

capable of synthesizing all the required cell constituents from glucose alone (Figure S1).

Processes like growth and reproduction are punctiliously regulated by the timely expression

of specific genes and precise localization of proteins within the cell [67]. This thigh

coordination between anabolic processes, growth, and reproduction is considered essential for

the proper functioning of the cell [68]. In contrast to this current understanding, we demonstrate

that cells can still reproduce without such intracellular coordination. Excessive lipid

synthesis, the multilamellar nature of the cell membrane, and favorable environmental

conditions facilitated the formation of long filamentous cells by the sequential expansion of

the outer membrane followed by the inner cell membrane. This two-step formation of long

filamentous cells rather than large spherical cells by the random expansion of cytoplasm

could have played an essential role in mediating leak-proof cell fission [64].

Conclusions

In the present study, we demonstrated the influence of environmental conditions on

protoplast reproduction. Our study shows that cells can reproduce independently of canonical

molecular biological processes without significantly impacting reproductive efficiency under

favorable environmental conditions. Given the simplicity and efficiency with which these

cells reproduced, we propose primitive Gram-negative bacteria likely reproduced by a similar

process before the evolution of complex intracellular processes.

Acknowledgments

We want to thank Petra Schwille for her support and scientific input throughout the work and

in preparing the manuscript. Gabriella Berthal and Markus Oster for excellent technical

support. We thank the Advanced Light Microscopy Facility at EMBL, Heidelberg, Ulf

Schwartz from Leica Microsystems, and colleagues at the departments of Ecological

Microbiology (Bayreuth University) and of Cellular and Molecular Biophysics (Max Planck

Institute for Biochemistry) for their support throughout the work.

Author contributions

Conceptualization: DK, ML, BZ & TL. Culture experiments: DK. Electron microscopy: DK

& AK. Light microscopy: DK & ML. Analysis: DK & JHK. Manuscript Writing: DK &

JHK.

Funding

This research was funded by the European Research Council (ERC) grant agreement 616644

(POLLOX) and by the Deutsche Forschungsgemeinschaft (DFG) grant agreements DFG-

TRR174 and Seed funding from Excellence Cluster ORIGINS EXC2094 – 390783311.

Conflicts of Interest

The authors declare that there are no conflicts of interest regarding the publication of this

article.

Data Availability

Dr. Dheeraj Kanaparthi will share all data, materials, and methods upon reasonable request.

References

1. Vollmer, W., Blanot, D., and De Pedro, M.A. (2008). Peptidoglycan structure and architecture. FEMS Microbiol. Rev. *32*. 149-167. 10.1111/j.1574-6976.2007.00094.x.

2. Kern, T., Giffard, M., Hediger, S., Amoroso, A., Giustini, C., Bui, N.K., Joris, B., Bougault, C., Vollmer, W., and Simorre, J.P. (2010). Dynamics characterization of fully hydrated bacterial cell walls by solid-state NMR: Evidence for cooperative binding of metal ions. J. Am. Chem. Soc. *132*. 10911-10919. 10.1021/ja104533w.

3. Westblade, L.F., Errington, J., and Dörr, T. (2020). Antibiotic tolerance. PLOS Pathog. *16*, e1008892. 10.1371/journal.ppat.1008892.

4. Kawai, Y., Mercier, R., Wu, L.J., Domínguez-Cuevas, P., Oshima, T., and Errington, J. (2015). Cell growth of wall-free L-form bacteria is limited by oxidative damage. Curr. Biol. *25*, 1613–1618.

5. Eisenberg, A.D., and Corner, T.R. (1973). Osmotic behavior of bacterial protoplasts: temperature effects. J. Bacteriol. *114*. 10.1128/jb.114.3.1177-1183.1973.

6. Wohlfarth, J.C., Feldmüller, M., Schneller, A., Kilcher, S., Burkolter, M., Meile, S., Pilhofer, M., Schuppler, M., and Loessner, M.J. (2023). L-form conversion in Gram-positive bacteria enables escape from phage infection. Nat. Microbiol. *8*. 387-399. 10.1038/s41564-022-01317-3.

7. Kapteijn, R., Shitut, S., Aschmann, D., Zhang, L., de Beer, M., Daviran, D., Roverts, R., Akiva, A., van Wezel, G.P., Kros, A., et al. (2022). Endocytosis-like DNA uptake by cell wall-deficient bacteria. Nat. Commun. *13*. 10. 5524. 1038/s41467-022-33054-w.

8. Sukhithasri, V., Nisha, N., Biswas, L., Anil Kumar, V., and Biswas, R. (2013). Innate immune recognition of microbial cell wall components and microbial strategies to evade such recognitions. Microbiol. Res. *168*. 396-406. 10.1016/j.micres.2013.02.005.

9. Thirumalapura, N.R., Ramachandran, A., Morton, R.J., and Malayer, J.R. (2006). Bacterial cell microarrays for the detection and characterization of antibodies against surface antigens. J. Immunol. Methods *309*. 48-54. 10.1016/j.jim.2005.11.016.

10. Clasener, H. (1972). Pathogenicity of the L-phase of bacteria. Annu. Rev. Microbiol. *26*. 10.1146/annurev.mi.26.100172.000415.

11. Errington, J., Mickiewicz, K., Kawai, Y., and Wu, L.J. (2016). L-form bacteria, chronic diseases and the origins of life. Philos. Trans. R. Soc. B Biol. Sci. *371*. 10.1098/rstb.2015.0494.

12. Gutman, L.T., Schaller, J., and Wedgwood, R.J. (1967). Bacterial L-forms in relapsing urinary-tract infection. Lancet *1*. 10.1016/s0140-6736(67)91090-2.

13. Mickiewicz, K.M., Kawai, Y., Drage, L., Gomes, M.C., Davison, F., Pickard, R., Hall, J., Mostowy, S., Aldridge, P.D., and Errington, J. (2019). Possible role of L-form switching in recurrent urinary tract infection. Nat. Commun. *10*. 4379. 10.1038/s41467-019-12359-3.

14. Leaver, M., Dominguez-Cuevas, P., Coxhead, J.M., Daniel, R.A., and Errington, J. (2009). Life without a wall or division machine in Bacillus subtilis. Nature *457*, 849-853. https://doi.org/10.1038/nature07742.

15. Mercier, R., Kawai, Y., and Errington, J. (2013). Excess membrane synthesis drives a primitive mode of cell proliferation. Cell *152*, 997–1007.

16. Ramijan, K., Ultee, E., Willemse, J., Zhang, Z., Wondergem, J.A.J., van der Meij, A., Heinrich, D., Briegel, A., van Wezel, G.P., and Claessen, D. (2018). Stress-induced formation of cell wall-deficient cells in filamentous actinomycetes. Nat. Commun. *9*. 5164. 10.1038/s41467-018-07560-9.

17. Malek, M.A., Bitam, I., Levasseur, A., Terras, J., Gaudart, J., Azza, S., Flaudrops, C., Robert, C., Raoult, D., and Drancourt, M. (2017). Yersinia pestis halotolerance illuminates plague reservoirs. Sci. Rep. *7*. 40022. 10.1038/srep40022.

18. Studer, P., Staubli, T., Wieser, N., Wolf, P., Schuppler, M., and Loessner, M.J. (2016). Proliferation of Listeria monocytogenesL-form cells by formation of internal and external vesicles. Nat. Commun. *7*, 13631. https://doi.doi: 10.1038/ncomms13631

19. Briers, Y., Staubli, T., Schmid, M.C., Wagner, M., Schuppler, M., and Loessner, M.J. (2012). Intracellular vesicles as reproduction elements in cell wall-deficient L-form bacteria. PLoS ONE *7*, e38514.

20. Wu, L.J., Lee, S., Park, S., Eland, L.E., Wipat, A., Holden, S., and Errington, J. (2020). Geometric principles underlying the proliferation of a model cell system. Nat. Commun. *11*. Article no. 4149. https://doi:10.1038/s41467-020-17988-7.

21. Karimi, M., Steinkühler, J., Roy, D., Dasgupta, R., Lipowsky, R., and Dimova, R. (2018). Asymmetric Ionic Conditions Generate Large Membrane Curvatures. Nano Lett. 12, 7816-7821. https://doi:10.1021/acs.nanolett.8b03584.

22. Liu, Y., Agudo-Canalejo, J., Grafmüller, A., Dimova, R., and Lipowsky, R. (2016). Patterns of flexible nanotubes formed by liquid-ordered and liquid-disordered membranes. ACS Nano. 10, 463-474. https://doi:10.1021/acsnano.5b05377.

23. Liu, Y., Lipowsky, R., and Dimova, R. (2012). Concentration dependence of the interfacial tension for aqueous two-phase polymer solutions of dextran and polyethylene glycol. Langmuir *28*. 3831-3839. https://doi:10.1021/la204757z.

24. Tsai, H.H.G., Lai, W.X., Lin, H.D., Lee, J.B., Juang, W.F., and Tseng, W.H. (2012). Molecular dynamics simulation of cation-phospholipid clustering in phospholipid bilayers: Possible role in stalk formation during membrane fusion. Biochim. Biophys. Acta - Biomembr. *1818*. 2742-2755. https://doi:10.1016/j.bbamem.2012.05.029.

25. Rojas, E., Theriot, J.A., and Huang, K.C. (2014). Response of Escherichia coli growth rate to osmotic shock. Proc. Natl. Acad. Sci. U. S. A. *111*. 7807-7812 10.1073/pnas.1402591111.

26. Cesar, S., Anjur-Dietrich, M., Yu, B., Li, E., Rojas, E., Neff, N., Cooper, T.F., and Huang, K.C. (2020). Bacterial evolution in high-osmolarity environments. mBio *11*. 1191-1120. 10.1128/mBio.01191-20.

27. Sleator, R.D., and Hill, C. (2002). Bacterial osmoadaptation: the role of osmolytes in bacterial stress and virulence. FEMS Microbiol. Rev. *26*. 49-71. 10.1111/j.1574-6976.2002.tb00598.x.

28. Kysela, D.T., Randich, A.M., Caccamo, P.D., and Brun, Y.V. (2016). Diversity Takes Shape: Understanding the Mechanistic and Adaptive Basis of Bacterial Morphology. PLOS Biol. *14*, 1–15. 10.1371/journal.pbio.1002565.

29. Narsimhan, V., Spann, A.P., and Shaqfeh, E.S.G. (2015). Pearling, wrinkling, and buckling of vesicles in elongational flows. J. Fluid Mech. 777, 1-26. 10.1017/jfm.2015.345.

30. Stokes, N.R., Baker, N., Bennett, J.M., Berry, J., Collins, I., Czaplewski, L.G., Logan, A., Macdonald, R., MacLeod, L., Peasley, H., et al. (2013). An improved small-molecule inhibitor of FtsZ with superior in vitro potency, drug-likeproperties, and in vivo efficacy. Antimicrob. Agents Chemother. *57*. 317-325, https://doi:10.1128/AAC.01580-12.

31. Kanaparthi, D., and Conrad, R. (2015). Role of humic substances in promoting autotrophic growth in nitrate-dependent iron-oxidizing bacteria. Syst. Appl. Microbiol. *38*, 184–188. http://dx.doi.org/10.1016/j.syapm.2015.02.009.

32. Zhang, Z., Zhang, Q., Guan, S., and Shi, H. (2020). Quantitative Connection between Cell Size and Growth Rate by Phospholipid Metabolism. Cells *9*. 1-28, https://doi:10.3390/cells9020391.

33. Hertel, C., Schmidt, G., Fischer, M., Oellers, K., and Hammes, W.P. (1998). Oxygen-dependent regulation of the expression of the catalase gene katA of Lactobacillus sakei LTH677. Appl. Environ. Microbiol. *64*. 1359-1363, https://doi:10.1128/aem.64.4.1359-1365.1998.

34. Mercier, R., Kawai, Y., and Errington, J. (2013). Excess membrane synthesis drives a primitive mode of cell proliferation. Cell *152*, 997–1007.

35. Studer, P., Borisova, M., Schneider, A., Ayala, J.A., Mayer, C., Schuppler, M., Loessner, M.J., and Briers, Y. (2016). The Absence of a Mature Cell Wall Sacculus in Stable Listeria monocytogenes L-Form Cells Is Independent of Peptidoglycan Synthesis. PLoS ONE *11*. e0154925. https://doi.org/10.1371/journal.pone.0154925.

36. Lipowsky, R. (1992). Budding of membranes induced by intramembrane domains. J. Phys. II. *2*, 1825-1840. https://doi:10.1051/jp2:1992238.

37. Baumgart, T., Hess, S.T., and Webb, W.W. (2003). Imaging coexisting fluid domains in biomembrane models coupling curvature and line tension. Nature. *425*, 821-824. https://doi:10.1038/nature02013.

38. Bacia, K., Schwille, P., and Kurzchalia, T. (2005). Sterol structure determines the separation of phases and the curvature of the liquid-ordered phase in model membranes. Proc. Natl. Acad. Sci. U. S. A. *102*, 3272-3277. https://doi:10.1073/pnas.0408215102.

39. Rashby, S.E., Sessions, A.L., Summons, R.E., and Newman, D.K. (2007). Biosynthesis of 2-methylbacteriohopanepolyols by an anoxygenic phototroph. Proc. Natl. Acad. Sci. *104*, 15099–15104.

40. Andes-Koback, M., and Keating, C.D. (2011). Complete budding and asymmetric division of primitive model cells to produce daughter vesicles with different interior and membrane compositions. J. Am. Chem. Soc. *133*. 9545-9555. https://doi:10.1021/ja202406v.

41. Liu, Y., Lipowsky, R., and Dimova, R. (2012). Concentration dependence of the interfacial tension for aqueous two-phase polymer solutions of dextran and polyethylene glycol. Langmuir *28*. 3831-3839. https://doi:10.1021/la204757z.

42. Breidenich, M., Netz, R.R., and Lipowsky, R. (2005). The influence of non-anchored polymers on the curvature of vesicles. Mol. Phys. *103*. 3169-3183. https://doi:10.1080/00268970500270484.

43. Sandoz, P.A., Tremblay, C., Gisou van der Goot, F., and Frechin, M. (2019). Image-based analysis of living mammalian cells using label-free 3D refractive index maps

reveals new organelle dynamics and dry mass flux. PLoS Biol. *17*. e3000553. https://doi:10.1371/journal.pbio.3000553.

44. Dreher, Y., Jahnke, K., Bobkova, E., Spatz, J.P., and Göpfrich, K. (2021). Division and Regrowth of Phase-Separated Giant Unilamellar Vesicles**. Angew. Chem. - Int. Ed. *60*. 10661-10669. https://doi:10.1002/anie.202014174.

45. Strahl, H., Bürmann, F., and Hamoen, L.W. (2014). The actin homologue MreB organizes the bacterial cell membrane. Nat. Commun. *5*. Article no. 3442. https://doi:10.1038/ncomms4442.

46. Simons, K., and Ikonen, E. (1997). Functional rafts in cell membranes. Nature. *387*, 569-572. https://doi:10.1038/42408.

47. Kantsler, V., Segre, E., and Steinberg, V. (2008). Critical dynamics of vesicle stretching transition in elongational flow. Phys. Rev. Lett. *101*, 048101-048104. https://doi:10.1103/PhysRevLett.101.048101.

48. Yanagisawa, M., Imai, M., and Taniguchi, T. (2010). Periodic modulation of tubular vesicles induced by phase separation. Phys. Rev. E - Stat. Nonlinear Soft Matter Phys. *82*, 1-9. https://doi:10.1103/PhysRevE.82.051928.

49. Yanagisawa, M., Imai, M., and Taniguchi, T. (2008). Shape deformation of ternary vesicles coupled with phase separation. Phys. Rev. Lett. *148*, 102-104. https://doi:10.1103/PhysRevLett.100.148102.

50. Sanborn, J., Ogl□cka, K., Kraut, R.S., and Parikh, A.N. (2012). Transient pearling and vesiculation of membrane tubes under osmotic gradients. Faraday Discuss. 161, 161-176, 2012. https://doi:10.1039/c2fd20116j.

51. Cooper, S. (2006). Distinguishing between linear and exponential cell growth during the division cycle: Single-cell studies, cell-culture studies, and the object of cell-cycle research. Theor. Biol. Med. Model. 3, 10, https://doi:10.1186/1742-4682-3-10.

52. Mitchison, J.M. (2005). Single cell studies of the cell cycle and some models. Theor. Biol. Med. Model. 2, 4, https://doi:10.1186/1742-4682-2-4.

53. Kærn, M., Elston, T.C., Blake, W.J., and Collins, J.J. (2005). Stochasticity in gene expression: From theories to phenotypes. Nat. Rev. Genet. 6, 451-464. https://doi:10.1038/nrg1615.

54. Elowitz, M.B., Levine, A.J., Siggia, E.D., and Swain, P.S. (2002). Stochastic gene expression in a single cell. Science. 297, 1183-1186, https://doi:10.1126/science.1070919.

55. Strovas, T.J., Sauter, L.M., Guo, X., and Lidstrom, M.E. (2007). Cell-to-cell heterogeneity in growth rate and gene expression in Methylobacterium extorquens AM1. J. Bacteriol. 189, 7132-7133, https://doi:10.1128/JB.00746-07.

56. Narsimhan, V., Spann, A.P., and Shaqfeh, E.S.G. (2014). The mechanism of shape instability for a vesicle in extensional flow. J. Fluid Mech. 750, 144-190, https://doi:10.1017/jfm.2014.248.

57. Zhao, H., and Shaqfeh, E.S.G. (2013). The shape stability of a lipid vesicle in a uniaxial extensional flow. J. Fluid Mech. 719, 345-361, https://doi:10.1017/jfm.2013.10.

58. Zwicker, D., Seyboldt, R., Weber, C.A., Hyman, A.A., and Jülicher, F. (2017). Growth and division of active droplets provides a model for protocells. Nat. Phys. *13*. 408-413. https://doi:10.1038/nphys3984.

59. Oglęcka, K., Rangamani, P., Liedberg, B., Kraut, R.S., and Parikh, A.N. (2014). Oscillatory phase separation in giant lipid vesicles induced by transmembrane osmotic differentials. eLife. 3, e03695, https://doi:10.7554/elife.03695.

60. Ramijan, K., Ultee, E., Willemse, J., Zhang, Z., Wondergem, J.A.J., van der Meij, A., Heinrich, D., Briegel, A., van Wezel, G.P., and Claessen, D. (2018). Stress-induced formation of cell wall-deficient cells in filamentous actinomycetes. Nat. Commun. *9*. Article number: 5164, https://doi:10.1038/s41467-018-07560-9.

61. Schultz, Z.D., Pazos, I.M., McNeil-Watson, F.K., Lewis, E.N., and Levin, I.W. (2009). Magnesium-induced lipid bilayer microdomain reorganizations: Implications for membrane fusion. J. Phys. Chem. B. *113*, 9932-9941. https://doi:10.1021/jp9011944.

62. Deshpande, S., Spoelstra, W.K., Van Doorn, M., Kerssemakers, J., and Dekker, C. (2018). Mechanical Division of Cell-Sized Liposomes. ACS Nano *12*. 2560-2568. https://doi:10.1021/acsnano.7b08411.

63. Hanczyc, M.M., Fujikawa, S.M., and Szostak, J.W. (2003). Experimental models of primitive cellular compartments: encapsulation, growth, and division. Science *302*, 618–622.

64. Zhu, T.F., and Szostak, J.W. (2009). Coupled growth and division of model protocell membranes. J. Am. Chem. Soc. *131*, 5705. https://doi.org/10.1021/ja900919c.

65. Di Rienzo, C., Piazza, V., Gratton, E., Beltram, F., and Cardarelli, F. (2014). Probing short-range protein Brownian motion in the cytoplasm of living cells. Nat. Commun. *5*. Article no. 5891. https://doi:10.1038/ncomms6891.

66. Fisher, J.K., Bourniquel, A., Witz, G., Weiner, B., Prentiss, M., and Kleckner, N. (2013). Four-dimensional imaging of E. coli nucleoid organization and dynamics in living cells. Cell *153*. 822-895. https://doi:10.1016/j.cell.2013.04.006.

67. Khanna, K., Lopez-Garrido, J., Sugie, J., Pogliano, K., and Villa, E. (2021). Asymmetric localization of the cell division machinery during bacillus subtilis sporulation. eLife *10*. e62204. https://doi:10.7554/eLife.62204.

68. Lozada-Chávez, I., Janga, S.C., and Collado-Vides, J. (2006). Bacterial regulatory networks are extremely flexible in evolution. Nucleic Acids Res. *34*. 3434-3445. https://doi:10.1093/nar/gkl423.

Methods and Materials

Cultivation of bacteria and generation of bacterial protoplasts

Gram-negative *Rhodobacter sphaeroides* (*DSM159*) was obtained from DMSZ

(Braunschweig, Germany). Lyophilized cells were revived in nutrient broth (Sigma Aldrich,

Germany) according to the procedure described by DSMZ. Protoplasts were generated from

R. sphaeroides (*RS-P*) by inoculating overnight cultures into 3X nutrient broth (NB)

containing 2% w/v sucrose, 80 µg/ml lysozyme, and 200µg/ml penicillin G (Sigma Aldrich,

Germany). Cells were observed under a microscope to confirm the transformation of cells to

their protoplast state. Over the next 3-4 weeks, the cells were repeatedly subcultured into

fresh media with penicillin G and lysozyme, as described above. Subsequently, *RS-P* was

transferred into NB with 7% Dead Sea Salt (DSS) (Amazon.de). The stability of these

protoplasts (*RS-P*) was determined after ten sub-cultures by transferring cells into media

without penicillin G or lysozyme. As *RS-P* did not revert to its native state with the cell wall

under these conditions, lysozyme, and penicillin G were not added to the growth media for

subsequent experiments. Unless otherwise stated, cells in the rest of the experiments were

grown in 7%DSS-NB. In some experiments, we replaced DSS in the growth media with

equivalent concentrations of other salts like $MgCl_2$ and KCl (7%w/v). Cell growth was

quantified at intervals by measuring their optical density at regular intervals (as mentioned in

Figure 3) using a Biophotometer[®] D30 (Eppendorf, Germany). The morphology of cells at

different growth stages was determined by a variety of microscopic techniques (as described

in the below sections).

Influence of salt on the stability of protoplasts

The influence of salt on the stability of protoplasts was tested by two different methods. In the first method, the osmolarity of the media was reduced by dilution with NB. This was done in a step-by-step manner of replacing 10ml of the 7%DSS-NB (from the original 100ml) with 10ml of NB with no additional salt, day over seven days. All test and control experiments were replicated (n=6 each). To the control bottles, we added 10ml of the fresh 7%DSS-NB media. The morphology of the cells was observed using a phase-contrast microscope. No lysozyme and penicillin G were added to both of these media.

In the second method, we transferred 2ml of mid-log phase cells grown in 7%DSS-NB into 6xNB with 2% sucrose, 6xNB with 2% sucrose containing lysozyme and penicillin G and into 7%DSS-NB. 6xNB with 2% sucrose was used in these experiments to maintain the osmolarity similar to the media similar to 7%DSS-NB. All experiments were replicated (n=6 each).

To understand if the intracellular processes played a role in regulating the protoplast morphology, we inoculated *RS-P* into 7%DSS-NB with and without FtsZ functional inhibitor, 5mM 3-methoxybenzamide[30]. Cells were observed under a phase-contrast microscope at regular intervals for differences in the morphologies between the control and test incubations.

Quantification of Superoxide dismutase activity and metabolic end products under aerobic growth

Minimal salt media (MSM) with 2% w/v glucose was used for these experiments. The composition of the minimal salt media was previously described by Kanaparthi et al., 2015 [31],

with the exception of using 0.1g/l of yeast extract. 100ml of media was dispensed into 500ml

Schott bottles (Schott AG, Germany). One set of growth media was made anaerobic using a

vacuum manifold by repeatedly flushing the headspace with N_2 gas and sealed with butyl

rubber stoppers before autoclaving. After autoclaving, glucose and vitamin solutions were

added to both aerobic and anaerobic media from the sterile anoxic stock solution. Aerobic

and anaerobic media were inoculated (500µl) with overnight cultures of *RS*, grown in aerobic

and anaerobic half-strength NB. A similar media composition was used to cultivate *RS-P*

under aerobic and anaerobic conditions, except for adding 7%DSS. All the bottles were

incubated on an orbital shaker at 30^0C. To ensure uniform oxygen solubility, cultures grown

under aerobic conditions were subject to shaking speeds of 150 rpm. All culture bottles were

incubated in the dark to ensure non-phototrophic growth. All experiments were done in

replicates (n=5). OD was determined every 3h interval for 15h. 1ml of the culture was

centrifuged, and the supernatant was filter sterilized and stored at -20^0C, for volatile fatty acid

analysis. Glucose was quantified using a Glucose colorimetric assay kit (Invitrogen,

Germany), according to manufacturers' instructions.

SOD activity was determined using the 19160 SOD detection kit (Sigma Aldrich, Germany).

Anaerobic culture bottles were transferred to an anaerobic chamber (Braun, Germany), and 1

ml of the culture was transferred into a 2 ml screw cap tube. The optical density of the cell

was adjusted to a similar value in all the samples. Cells were homogenized using silica beads

(Sigma Aldrich, Germany) and vertexing for 3min. All tubes were briefly centrifuged, and

250µl of the cell lysate was transferred into the microtiter plate, and other reagents were

added according to the manufacturer's instructions. The plates were then transferred out of

the anaerobic chamber and incubated at 37^0C for 1h. The colorimetric measurements were

done using a 96-well plate reader (Thermo Fischer Scientific, Germany). A similar

experimental procedure was used for determining the SOD activity of *RS* and *RS-P* cells grown under aerobic conditions. All the experiments were done in replicates n=10 for the data presented in the manuscript, but the experiment was repeated several times over the course of our work.

Phase-contrast and Super-resolution STED microscopy

Morphology *RS-P* was determined at regular intervals using an Axioskop 2plus microscope (Carl Zeiss, Germany) using a Plan-NEOFLUAR 100X/1.3 objective. Images were acquired with Leica DSF9000 (Leica, Germany) cameras. STED microscopy was performed with an inverted TCS SP8 STED 3X microscope (Leica Microsystems, Mannheim Germany) using an 86x/1.2 NA water immersion objective (Leica HC PL APO CS2 - STED White). Fluorophores were excited with either 488, 561nm, 594nm, or 633m laser light derived from an 80 MHz pulsed White Light Laser (Leica Microsystems, Mannheim Germany). For stimulated emission, either a pulsed 775 nm laser or a 592nm CW laser (Leica Microsystems, Mannheim, Germany) was used depending on the fluorophore. The emitted fluorescence light was collected with Hybrid Detectors (HyD, Leica Microsystems, Mannheim Germany) using appropriate emission band pass-filter setting and typically a gate of 0.3ns-6.0ns for depletion with the 775nm laser or 1.0-6.0ns with the 592nm laser. Images were recorded in photon counting mode and line accumulation. Image deconvolution was performed on selected images and video with Huygens Professional (version 16.10.1p2, Scientific Volume Imaging, Hilversum, The Netherlands).

Transmission electron microscopy

RS-P cells at different growth stages were harvested from the growth media by centrifugation

at 500rpm for 5min. Cells were then preserved in 2.5% v/v glutaraldehyde (Carl Roth,

Karlsruhe, Germany). Cells were subsequently stained with 0.25% w/v uranyl acetate. Cells

were subsequently dehydrated in increasing concentrations of acetone. Cells were embedded

in epoxy resin and left for polymerization for 72h. Epoxy blocks were sectioned, and cells

were counter-stained with 1% lead citrate. TEM was carried out using a Zeiss EM 912 (Zeiss,

Oberkochen) with an integrated OMEGA filter at 80kV. Images were acquired with 2k x 2k

pixel slow scan CCD camera (TRS, TrÖndle Restlichtverstrkersysteme, Moorenweis,

Germany) and an ImageSP software (version 1.2.9.77) (SysProg, Minsk, Belarus).

Quantifying reproductive efficiency and metabolic viability of daughter cells

Reproductive efficiency was determining the leakage of cell constituents during the

formation of daughter cells. The viability of the daughter cell was determined by staining and

quantifying cells by flow cytometry.

RS-P was inoculated into nutrient broth containing 7% w/v of DSS, NaCl, or $MgCl_2$. The

culture bottles were incubated under static conditions or on an orbital shaker at 160 rpm. 2ml

of cells were harvested every 24h intervals from all the incubations. In the case of static

cultures, incubations were briefly placed on an orbital shaker for 10 min at 100 rpm to

disperse the biofilm. Intact cells were precipitated by centrifugation at 6000 rpm for 10 min.

DNA was extracted from both the cell and supernatant. In the final step, DNA was washed

with 70% ethanol three times to remove any residual salts from DNA. The resultant DNA

was dissolved in 2ml of TE buffer. DNA was quantified using the Qubit dsDNA

quantification kit (Invitrogen). All the experiments were done in replicates (biological

replication, n=5).

The viability of daughter cells was determined by the growth of daughter cells either by

directly sub-culturing 10-20 day old cultures into fresh media or passing 10-20 day old

cultures through 0.45μm cellulose acetate filter (Millipore) and then transferring the filtrate

into new media. The presence of intracellular DNA and cytoplasmic activity of daughter cells

was also tested by staining them with DNA stain, PicoGreen^{TM} (Invitrogen), and live cell

stain CellTrace^{TM} violet stain (Invitrogen). Cells with intracellular DNA and cytoplasmic

activity were quantified using flow cytometry (Attune NxT, Thermo Fisher Scientific).