1

The reproduction process of Gram-positive protocells

Dheeraj Kanaparthi[1,2&3*], Marko Lampe[4], Jan-Hagen Krohn[1&3], Baoli Zhu[2&5], Falk

Hildebrand[6], Thomas Boesen[7], Andreas Klingl[8], Prasad Phapale[9], & Tillmann Lueders[2*]

Affiliations:

[1]Department of Cellular and Molecular Biophysics, Max-Planck Institute for Biochemistry,

 Munich, Germany.

[2]Chair of Ecological Microbiology, BayCeer, University of Bayreuth, Germany.

[3]Excellenzcluster ORIGINS, Garching, Germany.

[4]Advanced Light Microscopy Facility, European Molecular Biology Laboratory, Heidelberg,

 Germany.

[5]Key Laboratory of Agro-ecological Processes in Subtropical Regions, CAS, Changsha, China.

[6]Quadrum institute, Norwich, UK.

[7]Department of Biosciences, Center for Electromicrobiology, Aarhus, Denmark.

[8]Department of Biology, LMU, Planegg-Martinsried, Germany.

[9]European Molecular Biology Laboratory, Heidelberg, Germany.

* Corresponding author

Abstract

Prokaryotes are hypothesized to have evolved from more primitive protocells. Unlike present-day cells, protocells are thought to have been devoid of complex molecular biological processes. They are believed to have mediated reproduction entirely by biophysical forces under favorable environmental conditions. Despite this proposition, little is known about the actual mechanism of their reproduction. To understand the reproduction process of protocells in their native habitat, here we used a top-down approach to transform bacterial cells into a primitive lipid vesicle-like state. Given that environmental conditions are thought to have played an essential role in mediating protocell reproduction, we then studied these cells under the presumed environmental conditions of Archaean Eon Earth. Even in the absence of functioning biological processes, cells in our study reproduced in a defined sequence of steps, always leading to the formation of viable daughter cells. Their reproduction mechanism can be explained by the interaction between intracellular metabolism, physicochemical properties of cell constituents, and, most importantly, environmental conditions. Given the simplicity of this reproduction mechanism and its suitability to environmental conditions of early Earth, we propose that protocells reproduced by this process. Moreover, this method of reproduction is also in tune with the earlier theoretical propositions on protocells, the results of the top-down approach of building a minimal cell, and the paleontological record of the Achaean Eon. Our study is the first to bridge the gap between non-living systems like lipid vesicles, living cells, and the paleontology of the Archaean Eon.

1 **Introduction**

2

3 The theory of chemical evolution proposes that prokaryotes evolved from more primitive

4 protocells (1). Molecular biological processes in these primitive cells are hypothesized to

5 have been simpler and far less complex than the present-day cells (2,3). Because protocells

6 lack complex molecular biological processes, they were believed to reproduce through simple

7 biophysical processes under favorable environmental conditions (3–5). However, much of

8 what we know about protocells has been derived from studying lipid vesicles (used as model

9 protocells) under well-controlled laboratory settings (4,5). Despite achieving significant

10 progress, we believe lipid vesicles (LVs) do not represent the true physiological complexity

11 of protocells. Moreover, studying an organism outside its natural habitat also fails to provide

12 a complete picture of its ecological function and behavior (6). This is especially true for

13 protocells, given their greater dependence on environmental conditions to mediate essential

14 cellular functions like reproduction. To understand the reproduction of protocells in their

15 natural habitat, here we worked with protoplasts of Gram-positive bacteria as model

16 protocells (*Exiguobacterium strain M* protoplasts –*EM-P*) under the environmental

17 conditions of early Earth.

18

19 Our choice of using bacterial protoplasts is based on three key considerations. First,

20 protocells like bacterial protoplasts were likely devoid of a peptidoglycan cell wall. Synthesis

21 of individual cell wall components, transporting them across the cell membrane, and

22 assembling them into a cell wall is a physiologically complex process requiring many

23 enzymes working in a coordinated manner (7,8). Evolutionarily primitive protocells most

24 certainly are considered to have been devoid of such complex intracellular processes (4).

25 Moreover, high-resolution FIB-SEM imaging of the earliest known microfossils was shown

to be surrounded by a 5nm carbon layer (9) resembling the thickness of a cell membrane

rather than a much thicker cell wall(10). Second, like the theoretical propositions on

protocells, protoplasts lack an innate ability to regulate their morphology or mediate

reproduction (11–13). They can be described as a sack of cytoplasm reproducing in a

haphazard manner (14). Third, even the most primitive cell must possess three essential

characteristics – biochemical processes for harvesting free energy, a mechanism of using this

energy to sustain minimal biosynthetic processes, and a mechanism of reproduction involving

the transfer of its genetic material to the daughter cells. Even these basic processes

incorporates considerable cytoplasmic complexity. Therefore, the biophysical properties of

protocells may be more similar to bacterial cytoplasm than compositionally simpler LVs.

Given these similarities between protocells and protoplasts, we argue that bacterial

protoplasts could be a better proxy than LVs for studying protocell reproduction.

Self-assembly of lipids into LVs happens within a narrow pH range and is influenced by

environmental conditions like temperature and salt content (15). Hence, most protocell

studies are conducted under narrow range of environmental conditions. Despite the

theoretical propositions on protocells suggesting that their reproduction is greatly dependent

on the environmental conditions, currently used experimental conditions rarely resemble the

natural habitat of protocells. To understand the protocell reproduction in its native habitat, we

grew *EM-P* under environmental conditions mimicking those that could have been prevalent

on early Earth. The exact environmental conditions of Archaean Earth are currently unclear.

Nevertheless, an emerging consensus among the scientific community suggests that at least

by 3.8 billion years ago, surface temperatures likely ranged between 26° and 35°C (16,17).

Early Earth's oceans are also hypothesized to be 1.5 to 2 times saltier than the present-day

oceans (18). Moreover, all known paleontological evidence of life is restricted to Archaean

Eon coastal marine environments (19). This suggests primitive life (if not the very first cells)

thrived under saline conditions. To replicate these conditions, we grew *EM-P* in half-strength

tryptic soy broth with 7% (w/v) Dead Sea Salt (TSB-DSS) at 30°C. We used DSS, rather than

pure NaCl, in our experiments to better simulate the complex salt compositions of natural

environments. We then monitored these cells at regular intervals using a variety of super-

resolution microscopic techniques to understand if and how these cells grow and reproduce.

In the following sections of the manuscript, we present the lifecycle of *EM-P* under Archaean

Eon environmental conditions, followed by a mechanistic explanation.

Methods

Exiguobacterium strain-Molly (*EM*) was isolated from submerged biofilms in the Dead Sea

(20). Protoplasts of *Exiguobacterium* Strain-Molly (*EM-P*) were generated by growing the

cells in 7%DSS-TSB with Lysozyme and Penicillin G (Sigma Aldrich, Germany).

Transformation to protoplast was confirmed by a change in cell morphology (Carl Zeiss,

Germany). Given the stability of protoplasts, the use of both lysozyme and penicillin G was

discontinued for subsequent experiments. For some experiments, *EM-P* was also grown in

TSB with 4%DSS, 15%DSS, 6%MgCl$_2$, and 7%KCl (Sigma Aldrich, Germany). We also

grew the cells in 3X TSB containing 0.3M sucrose, 0.5% glycerol, 300mg of BSA, and 20mg

of extracellular DNA.

Physiological characterization of *EM* was done in aerobic and anaerobic minimal salt media

(21) with 2% w/v glucose as a sole carbon source. Aerobic and anaerobic media were

inoculated (500ml) with overnight cultures of *EM*. A similar media composition was used to

cultivate *EM-P* under aerobic and anaerobic conditions, except for adding 7%DSS. All the

bottles were incubated on an orbital shaker at 30^0C. All experiments were done in replicates

(n=5). OD was determined every 3h interval for 15h. 1ml of the culture was centrifuged, and

the supernatant was filter sterilized and stored at -20C for volatile fatty acid analysis (22).

Glucose was quantified using a Glucose colorimetric assay kit (Invitrogen, Germany)

according to manufacturers' instructions. Catalase activity was determined using the 19160

SOD detection kit (Sigma Aldrich, Germany). For anaerobic cultures, OD of all the test

cultures was adjusted to similar values by dilution in original media, and cells were lysed by

bead beating within the anaerobic chamber (Braun, Germany). Cell lysates were then taken

out to measure catalase activity. All the experiments were done in replicates n=10 for the data

presented in the manuscript, but the experiment was repeated several times throughout our

work.

TEM and SEM were carried out using a Zeiss EM 912 (Carl Zeiss, Germany) with an

integrated OMEGA filter as described before (23). Fluorescence microscopy was performed

using an Olympus SpinSR10 spinning disk confocal microscope (Olympus, Tokyo, Japan).

STED super-resolution fluorescence imaging was performed with an inverted TCS SP8

STED 3X microscope (Leica Microsystems, Germany) using 86x/1.2 NA water immersion

objective (Leica HC PL APO CS2 - STED White).

To understand the relative contribution of intracellular processes and environmental

conditions on the *EM-P*'s surface instability, we treated cells either with 2% methanol-free

formaldehyde of 0.5% glutaraldehyde (ThermoFisher Scientific, Germany) to polymerize and

immobilize the movement of proteins inside the cell. To equilibrate the conditions in and out

of the cells, cells were permeabilized using 0.3% Triton X-100 (Sigma-Aldrich, Germany)

for 10 min.

To quantify extracellular and intracellular DNA, we inoculated *EM-P* into the respective media. We incubated them either under static conditions or on an orbital shaker at 160 rpm. 2 ml of cells were harvested every day (24h) from all the incubations. Intact cells were precipitated by centrifugation at 6000 rpm for 10 min. DNA was extracted from both cells and supernatant using phenol-chloroform extraction. The resultant DNA was dissolved in 0.2 ml of TE buffer. Quantification of DNA was done using the Qubit dsDNA quantification kit (Invitrogen, Germany).

The viability of daughter cells was determined by inoculating 10-20 day-old cultures into fresh media either directly or after passing them through a 0.45μm cellulose acetate filter (Schematic shown in Figure 8) to get rid of large cells and membrane debris. Cytoplasmic activity in daughter cells was tested by staining them with DNA stain, PicoGreen[TM,] and live cell labeling CellTrace[TM] violet stain (Invitrogen, Germany) and imaging the cells using a confocal microscope. The relative percentage of cells with and without cytoplasmic activity and intracellular DNA was quantified by Attune CytPix flow cytometer (Thermo Fischer Scientific, Germany). The metabolic viability of *EM-P* daughter cells was also tested using the minimal salt media described above, with glucose as the sole carbon source.

Results and Discussion

Given the lack of a cell wall, protoplasts were known to exhibit random morphologies and reproduce in a chaotic manner (11,14). This reproduction method is shown to have been independent of canonical molecular biological processes that govern bacterial reproduction (11). In addition to losing control over its morphology and reproduction, more recent studies

demonstrated that the loss of cell wall has more profound implications on cell physiology. These studies reported several discrepancies in intracellular functions, like an uncontrolled synthesis of cell wall precursors, even among stable protoplasts (24), the inability of protoplasts to maintain a proper intracellular oxidative state (25), uncontrolled replication of its genome (26) and excessive lipid (12). These studies suggest that cells in their protoplast state lack proper coordination between different cellular processes.

EM-P exhibited similar behavior. Cells in our incubations were in a perpetual state of lipid (membrane) excess (Fig. S1). The presence of excess membrane, together with the limited cytoplasmic volume of the cell, resulted in the formation of hollow extracellular and intracellular membrane structures as a mechanism of accommodating this excess membrane. We often observed log-phase *EM-P* cells with excess lipids within the cytoplasm in the form of lipid globules (Fig. S1a&b). Together with excessive lipid synthesis, *EM-P* also exhibited reduced efficiency in using terminal electron acceptors (Fig. S2) and a chaotic catalase expression (Supplementary results and discussion) compared to the wild-type cells with a cell wall. In tune with previous studies, these results suggest a lack of coordination among the processes in *EM-P*. However, in sharp contrast to the earlier studies (14), this loss of coordination did not result in an entirely random reproduction method. Under the environmental conditions of early Earth, *EM-P* reproduced in a defined sequence of steps, always leading to the formation of viable daughter cells. This consistency in its behavior was not due to the influence of molecular biological processes within the cell. Instead, it could be explained by the biophysical properties of its cell constituents, like the growth rate, excess lipid synthesis, and, most importantly, the environmental conditions.

Over the course of its life cycle, we observed *EM-P* reproducing by different processes. During the lag and early-log phase, cells remained spherical and gradually grew by an order of magnitude in diameter (Fig. 1a, 1b & Video 1). Subsequently, these spherical cells transformed into oval cells (Fig. 1c & Video 2). They then developed surface invaginations (Fig. 1c, 1d & Video 2), which gradually got deeper, transforming an oval-shaped cell into two visibly distinct daughter cells (Fig. 1d & Video 3). These cells then underwent fission, resulting in individual daughter cells. We occasionally saw cells that underwent fission but are still connected by a thin membrane tether (Video 3).

One key aspect of the cell cycle is the fission of the parent cell into metabolically viable daughter cells (3). Archiving cell division without the aid of molecular biological processes has been the primary focus of multiple studies and is of crucial importance in enhancing our understanding of most primitive cell replication processes (27) and building autonomously replicating synthetic cells (3,28). Previous studies archived fission either by mechanical means (29), by encapsulating protein involved in cell division (30), or by an uncontrolled splitting of LVs by gradually increasing the size of the LVs to the point of instability. Given *EM-P* was grown under static conditions as a thin film attached to the base of the culture flask, we do not presume that the physical stimulus required for splitting the parent cell into two daughter cells (binary fission) originated out of the cell.

In accordance with our proposition, most cells in the log-phase were observed to be in a constant state of surface oscillations/deformations (Video 1-3, constantly shivering cell surface). A similar surface instability was theoretically predicted in liquid droplets (31) and experimentally demonstrated in LVs packed with vibrating nanoparticles (32). Such surface deformations in droplets and LVs are shown to have been caused by the Brownian movement

of the intracellular molecules and their interactions with the cell membrane. This constant

movement of intracellular constituents is also thought to have provided the energy required to

split the cell into two daughter cells (32). In tune with these studies, we observed DNA in a

constant state of movement (video 4) in *EM-P*. In a normal bacterial cell, DNA forms highly

organized and condensed into a structure like a nucleoid or attached to the cell membrane via

a cation-mediated salt-bridge (33,34). The formation of DNA-DNA and DNA-membrane

complexes is known to have been mediated by divalent cations, which neutralize negative

charges of the phosphate groups (35). In the case of *EM-P*, no such organizational structure

was observed. DNA is distributed evenly throughout the cytoplasm as loose strands (video 4).

This suggests little interaction between individual DNA strands or DNA and cell membranes.

The absence of such structures and high concentration of intracellular DNA due to

uncontrolled genome replication (26), could have resulted in repulsion between DNA and

membrane. This constant movement of DNA within the cell could have led to surface

instability and ultimately to the fission of a parent cell. In addition to the movement of

intracellular constituents, similar surface deformations were also known to have been caused

by the action of proteins involved in cell division (30).

To understand if the surface deformation were a result of the Brownian movement of

intracellular constituents or due to the specific action of some intracellular protein, we treated

the cells with bactericidal chemicals like glutaraldehyde and formaldehyde (36). These

compounds were known to denature and polymerize proteins with little influence on the

biophysical properties of DNA (no influence of on DNA under the experimental conditions

of our study)(37). Treatment of cells either with glutaraldehyde and formaldehyde or with 3-

methoxybenzamide (38) did not resulted in a static cell. 3-methoxybenzamide inhibits the

activity of FtsZ, the tubulin homologue in prokaryotes known to play an important role in

regulating bacterial cell morphology and reproduction (39). Moreover, culturing of cells in

media containing 3-methoxybenzamide, did not affect the growth characteristics or the

morphology of the cells (Fig. S3). These results suggests that innate physiochemical forces

rather than the canonical molecular biological processes mediated the cell fission. Similar

electrostatic repulsion between old and newly synthesized copies of the genome is recently

shown to play an important role in genome segregation and fission of bacterial cells (40).

Together with the Brownian motion of intracellular constituents, surface deformations could

also have been caused by the transient binding of cations onto the membrane surface (41) or

due to the differences in the osmolarities and composition of molecules between the

cytoplasm and the growth media (42–44). In the case of *EM-P*, the osmolarity of the

cytoplasm was a result of high packing densities of intracellular constituents (organic in

nature)(42). While the osmolarity of the surrounding media was primarily a result of

inorganic salts (7%DSS) added to the growth media. To understand the influence of this

process on surface instability and cell fission, we transferred cells from 7%DSS-TSB into an

isotonic growth media. Unlike 7%DSS-TSB, this media's high osmolarity resulted from high

concentrations of organic compounds like sucrose, glycerol, DNA, and proteins. The

presence of these organic compounds could have resulted in similar conditions on either side

of the cell membrane. Apart from not observing surface deformations, cells in organic

carbon-rich media grew into random morphologies (Fig. 1 f-k). Segregation of the parent cell

into two daughter cells did not happen by binary fission. We instead observed random growth

of membrane (Fig. 1 f-k), ultimately compartmentalizing single parent cells into multiple

daughter cells. Transfer of log-phase cells between these two different media compositions

resulted in a reversible change in the cell morphology. Cell surface deformations in 7%DSS-

TSB can also be halted by osmotically equilibrating cytoplasm with the surrounding media

by treating the cells with membrane pore-forming compounds like Triton X100. These results suggest that cell division in *EM-P* was influenced by the environmental conditions (media composition) and the biophysical properties of cell constituents rather than the action of specific proteins within the cell (45).

Depending on the size of the daughter cells, cells in this growth stage appear to be reproducing either by budding (Fig. 1e-g) or by binary fission (Fig. 1c&d, & 2). Despite numerous repetitions, no uniformity was observed either in the sizes or number of daughter cells formed from a single *EM-P* cell (Fig. 1, 2, S4 & S5d). This asymmetric cell division and variability in the sizes of the daughter cells was theoretically predicted to happen in LVs made from compositionally diverse lipid species (46). Most previous attempts at understanding the biophysical properties of protocells were conducted on LVs of simple chemical composition. Usually made from one to three lipid species(5,42). In such LVs, the physical properties of vesicle membranes, like elasticity and spontaneous curvature, remain homogeneous along their surface (46). In contrast, live cell membranes comprise numerous lipids species, pigments, quinones, and membrane proteins (47). This diversity results in the formation of lipid-lipid (48) and lipid-protein (49) nanodomains. The physical properties of these nanodomains are dependent on the nature of the interactions between these components (50). Hence unlike the LVs, live cell membranes are considered to be an assembly of nanodomains that differ in their physical properties, i.e., their physical properties like elasticity and curvature remain non-homogenous throughout their surface (46) (Fig. 2a & 2b). This lateral heterogenicity in membrane physical properties, together with the amount of excess membrane, is theoretically predicted to result in cell division resembling budding (46).

Asymmetric cell division and the lack of uniformity in the sizes of the *EM-P* daughter cells

can be explained by these theoretical predictions (46). The excess lipid synthesis and

diversity of lipid species in *EM-P*'s cell membrane (47) could have created a state of low $\mathcal{V}$

(below 1) and H^B_m-values (a measure of heterogeneity in membrane physical properties) (Fig.

2). The $\mathcal{V}$-value of the cell is defined as the ratio of the actual volume of the cell (V) and its

volume in its spherical form (V_s), expresses as reduced volume ($\mathcal{V}$; $\mathcal{V}$=V/V_s). A spherical cell

with no excess membrane has $\mathcal{V}$-values of 1. Among all the cell morphologies, spherical cells

have the lowest volume compared to the surface area. The presence of excess membrane, as

observed in *EM-P* (Fig. S1), increases the cell's surface area. This, in turn, results in an

increased cell volume. This disproportional increase in the cell volume in relation to the

cytoplasmic volume results in the transformation of spherical *EM-P* cells into a plethora of

non-spherical morphologies. A large difference in the volume of the cell (~ membrane

surface area) and the volume of the cytoplasm results in low $\mathcal{V}$-values (Fig. 2). On the other

hand, a smaller difference between these two parameters results in high $\mathcal{V}$-values (closer to

1). Despite the excess lipid synthesis in *EM-P*, most lipids within the log-phase cell were not

incorporated into the cell membrane and remained within the cytoplasm as lipid globules

(Fig. S1a and S5). This resulted only in a slight elongation of the cells ($\mathcal{V}$-values below but

close to 1) and facilitated reproduction by budding (Fig. 2). The diversity in the sizes of the

daughter cells can be explained by the regional differences that are to be expected for a

bacterial cell, i.e., regional differences in the H^B_m-values. To our knowledge, ours is the first

study to link the theoretically predicted behavior of LVs with division in live cells (Fig. 2).

During the late log-phase, most cells developed filamentous extensions (video 5) or intracellular vesicles (Fig. 3-6). These structures could have formed due to the gradual transfer of lipids from the lipid globules to the cell membrane (Fig. S5-S7). This transfer of lipids is evident from the absence of lipid globules in stationary-phase cells. This influx of lipids, together with a reduced growth rate (~slow rate of cytoplasmic volume increase) (Fig. 1) of stationary phase cells, could have resulted in a state of the excess cell membrane (Fig. 3 & S1c-f). The formation of hollow filamentous or intracellular vesicles can be explained as a mechanism for accommodating this excess membrane. In accordance with this inference, cells kept in a continuous log-phase by transferring them to fresh media every few hours did not develop either intracellular or extracellular membrane extensions (Fig. S6 & S7). Like the log-phase cells, these cells continued to reproduce by binary fission (Fig. 1, S5 & S6g). Moreover, we also observed the presence of lipid globules within these cells (Fig. S7). The reason for this slow transfer of lipids from the cytoplasm to the cell membrane in actively growing cells is currently unclear.

The formation of filamentous extensions (Fig. 3a-d) and intracellular vesicles (Fig. 3e-i) did not happen in a random manner. These structures were observed to have formed in a sequence of steps previously reported from LVs (Fig. 3 n&o)(51). When LVs are placed in a hypertonic solution, they undergo a step-by-step morphological transformation to reach a thermodynamically stable state (52). The morphologies of such LVs are determined by the osmolarity of the media and by the rate of osmotic shrinkage. The gradual step-by-step increase in the osmolarity of the media resulted in a slow transfer of lipids from the LV membrane to the filamentous structures. From a thermodynamic perspective, this slow transfer of lipids will lead to the formation of hollow bud-like structures from multiple sites on the cell surface (51). Whereas a sudden increase in osmolarity resulted in a rapid

shrinkage of the LVs (i.e., a relatively faster rate of lipid transfer), which was shown to

thermodynamically favor the formation of fewer but longer filamentous extensions (Fig.

3n&o).

Like in LVs, the sequential steps involved in the formation of membrane extensions can be

explained by the differences in the growth rates of log and stationary phase cells and by the

rate of lipid transfer from the LV into the membrane extensions. We observed lipids

accumulated within the cytoplasm during the log-phase as lipid droplets (Fig. S1). We

noticed a slow rate of lipid transfer between these droplets and the cell membrane during the

log-phase cells (Fig. S7). This slow transfer of lipids to the cell membrane and the high

growth rate (~ rate of cytoplasmic volume increase) and nearly spherical morphologies

suggest that cells maintained a close match between the cell volume and surface area of the

cell membrane, i.e., less membrane was available for forming hollow filamentous extensions.

This led to the slow transfer of lipids from the cytoplasm to the membrane resulting in the

formation of tiny hollow membrane buds from multiple sites on the cell surface (Fig. 3a, b,

and e-h). Complete transfer of lipids from lipid droplets to the cell membrane during the late-

log and early stationary-phase, resulted in a sudden influx of lipids from the cytoplasm into

the cell membrane. Together with the reduced growth rate during the stationary phase, could

have energetically favored an increase in the length of the existing membrane extensions

(both internal and external Fig. 3d & 3l-m) rather than forming new membrane invaginations

on the cell surface. Transfer of late log-phase cells from media with 7%DSS to 15%DSS to

create a sudden state of lipid excess led to osmotic shrinkage of cells and abrupt increase in

the membrane area. In accordance with previous observations on LVs(51), these cells were

observed to have formed longer and fewer filamentous extensions (video 6) without the

initial appearance of multiple membrane buds on the cell membrane. These results suggest

that growth rate and environmental conditions played an important role in determining the

formation and morphology of filamentous extensions (Fig. 3).

All the above discussed morphological changes like the budding and formation of external

and internal membrane structures is associated with phase-separation of membrane into more

fluid, liquid-disordered (L_d) and more rigid, liquid-ordered (L_o) phases. This appearance of

distinct phases in the cell membrane and associated formation of external and internal

membrane extensions can be explained by the model proposed by Julicher & Lipowsky

(53,54). It was hypothesized that phase-separated membranes minimize interfacial energy at

the phase boundary to attain a thermodynamically stable state. They do so by forming an out-

of-plane invagination (bud formation), gradually transforming the softer L_d-membrane into a

bud. In accordance with this theory (53,54), the formation of buds in *EM-P* is associated with

the appearance of a distinct membrane phase – more fluid, L_d and more rigid L_o membrane

(Fig. 4). L_d membrane initially appeared as small patches on the cell surface, less than 180nm

in diameter (Fig. 4a-c). As the area of the L_d membrane increased, it underwent budding due

to an increase in the line tension at the phase boundary (55)(Fig. 4a, 4d, 4g & S8). However,

we observed some deviations from this expected behavior. First was the difference in the

direction of budding. Contrary to the predictions (53), we observed L_d-membrane underwent

a negative (into the cell) rather than positive (out of the cell) curvature (Fig. 4g). Second,

despite its higher bending rigidity and unfavorable thermodynamics, we also observed a more

rigid L_o-membrane undergoing budding. All the membrane structures extending out of the

cell are exclusively composed of L_o, and all the membrane structures extending into the cell

were composed of L_d membrane (Fig. 4d-f, & arrows 4g, cell within the highlighted region).

The direction and physical nature of a membrane that underwent budding in *EM-P* could be explained by its lipid composition (56) and high molecular crowding of the cytoplasm (57). Bacia *et al.* studied the influence of membrane composition on budding (56) and reported that the membrane sterols composition influences the direction and nature of the membrane that undergoes budding. Despite higher bending rigidities, this study showed that some sterols could preferentially induce the budding of a more rigid L_o membrane instead of L_d membrane. Prokaryotes lack sterols like cholesterol and their derivatives but possess functional cholesterol analogs, like hopanoids (58). The properties of these compounds in inducing membrane phase separation and budding were rarely investigated. Apart from sterols, invagination of a more rigid L_o membrane was also reported to happen when LVs (52) experience a state of an excess membrane (during the osmotic shrinkage in LVs or slow growth rate in protoplasts), possibly due to an increase in the spontaneous curvature of the L_o under such conditions (59).

The nature and direction of the membrane invagination could also have been determined by the differences in the molecular crowding between the cytoplasm and the growth media (57). Studies showed that when the aqueous core (cytoplasm) undergoes phase-separation into regions of higher and lower densities, results in a spontaneous membrane phase-separation (57). In such two-phase LVs, the L_d and L_o phases of the membrane were observed to have been associated with cytoplasmic regions of higher and lower densities, respectively. The mechanism behind this preferential association is currently not well understood. Nevertheless, this could explain the selective invagination of the L_d into the cell given the higher densities of cytoplasm (~17-35 wt% molecules) and L_o membranes into and out of the cell due to the comparatively less dense growth media (~6-8 wt% molecules) (Fig. 4). TEM

images of cells shown in Fig. 6, show these differences in the packing densities between the

cytoplasm and surroundings (60).

The live cell membranes are thought to undergo phase-separation, a process proposed to

profoundly influence cellular functions (48). Despite these propositions, the existence of

distinct phases within the cell membrane has been a point of contention among researchers

(61). To the best of our knowledge, ours is among the very few studies that provide visual

evidence of membrane phase-separation in live cells (Fig. 4)(62).

The formation of filamentous extensions and intracellular vesicles (Fig. 3) led to two

different reproduction methods. In short, *EM-P* reproduction either by forming external (Fig.

5) or intracellular (Fig. 6) daughter cells. Sequential steps involved in forming external and

internal daughter cells are shown in videos 7-11 and 12-18, respectively.

During the stationary growth phase, we observed cytoplasm transfer from the parent cell into

filamentous extensions (Fig. 5c & Video 7-9). Consequently, the hollow tubular filaments

transformed into a "string-of-individual daughter cells" (Fig. 5c-g & Videos 8&9). The

transformation of hollow filamentous extensions into string-of-individual daughter cells

could be explained by two different mechanisms. One could have been the spontaneous

transformation of tubular vesicles into a "string-of-beads" morphology to reduce the

interfacial area to reach a thermodynamically stable lower energy state(63). A similar

transformation of tubular vesicles during osmotic shrinkage when placed in a hyperosmotic

solution (identical to *EM-P,* video 6) was previously reported by multiple studies (63,64).

Alternately, the formation of daughter cells could also be explained simply by the lower

volume of cytoplasm that was transferred into the filaments compared to the total volume of

the filament (51). The excess membrane within these tubular structures should have collapsed

and transformed into tethers (Fig. 5 e&f, arrows) connecting adjacent daughter cells.

Lowering the salt content (osmolarity) of the media from 7% to 4 %DSS during the late log-

phase prevented the formation of filamentous extensions and transfer of cytoplasm to leading

to the formation of daughter cells (Fig. S9). Possibly due to the influx of water these cells we

observed these cells to be larger than the cells in the control experiment (Fig. S9). This

absence of daughter cells could be explained by the osmotic expansion of cells which could

have prevented the cytoplasm transfer from the cell into the filamentous extensions. In

comparison, the transfer of cells into media of higher osmolarity (15%DSS) resulted in the

osmotic deflation of cells and the formation of longer filamentous extensions with individual

daughter cells (Fig. S10). These strings of daughter cells are then detached from the parent

cell and fragmented into individual daughter cells in a sequence of steps, as shown in Figure

3F-I (video 9-11).

We also observed the formation of daughter cells within the intracellular vesicles. These

daughter cells were formed by a process similar to budding (Fig. 6d, arrow). These buds

either detached from the vesicle surface and released into the vesicle as individual daughter

cells (Fig. 6f&h & video 14) or grew into a string of daughter cells (Fig. 6e & video 14 – cell

on the left). These "string-of-daughter cells" subsequently detached from the vesicle and

fragmented (Fig. 6h) into individual daughter cells (video 17). Due to the gradual cytoplasm

transfer from parent to daughter cells (Fig. 6d, & 6e), the cytoplasmic content of the *EM-P*

cells gradually depleted (Fig. S11). By the late stationary-phase, many *EM-P* cells were

hollow with intracellular vesicles (Fig. 6g, S4 & video 14) containing highly motile daughter

cells. Daughter cells were released into the surrounding media by rupture of the cell or

vesicle membrane (Fig. 6h & Video 12-18). The Reproduction by formation of intra vesicular

daughter cells is observed in previous studies (65). Unlike in our study which shows that daughter cells were released into the intracellular vesicles, these studies presumed that hollow vesicles to be daughter cells. To the best of our knowledge, ours is the first study to show the sequential steps involved in the formation of internal daughter cells. The final growth stage of *EM-P* was characterized by the presence of cells with all the above-described morphologies, along with a considerable amount of membrane debris (Fig. S12).

Much of the focus in the above sections was dedicated to understanding *EM-P*'s life cycle from a biophysical perspective. Apart from the overall salinity, the biophysical properties of the cell constituents were known to have been influenced by the nature of the salts in the media. Different cations interact differently with the cell membrane and alter their physical properties (42,66,67). To understand how individual components of DSS influenced the morphology and life cycle of *EM-P*, we replaced DSS in the growth media with an equivalent concentration of pure salts like KCl and $MgCl_2$ (6%w/v). When grown in TSB-7%KCl, cells reproduced predominantly via the formation of intracellular daughter cells (Fig. S13). Cells in these incubations are larger than cells grown in DSS. In contrast, cells grown in TSB-6%$MgCl_2$ were significantly smaller than those grown in equivalent concentrations of DSS or KCl (Fig. S14, video 19 & 20). The diameter of these cells was restricted to 0.5-2μm and reproduced by forming extracellular daughter cells (Fig. S14). The nature of interactions between cations and phospholipid head groups(42) can explain these morphological differences. Earlier studies reported that monovalent cations like potassium increase the membrane's fluidity, whereas divalent cations like magnesium stiffen the membrane by reducing its fluidity(66). This increase in membrane fluidity could have thermodynamically facilitated the easier expansion of cell membranes in incubations containing DSS and KCl (42), resulting in relatively larger cells. Increased stiffness of the membrane in the presence

of MgCl$_2$ could have prevented the easier expansion of the cell membrane resulting in the formation of smaller cells (Fig. S14). The reasons for the reproduction predominantly via extracellular daughter in the presence of MgCl$_2$ and via intracellular daughter cells in the presence of KCl are currently unclear. Nevertheless, a similar phenomenon of intracellular vesicle formation in the presence of KCl was observed previously in LVs (42). *EM-P* reversibly changes its morphology when transferred between media of different salt compositions.

Apart from the salinity, one other characteristic feature of the coastal marine environments is tidal activity. Given the moon's proximity during the Archaean Eon, it is thought that coastal marine environments of early Earth could have experienced high-intensity wave action (68). To understand the influence of such physical forces on *EM-P's* reproduction, we grew cells on an orbital shaker. Under these growth conditions, *EM-P* reproduced exclusively by a process resembling binary fission, like the cells in the early log-phase (Fig. 1c-e). None of the above-described morphologies, like large cells with intracellular vesicles or strings of daughter cells (Fig. 5&6), were observed in these incubations. However, we observed the re-appearance of filamentous extensions and intracellular vesicles when we transferred late log-phase cells from an orbital shaker to a static incubator. These observations suggest that reproduction by binary fission is a result of culturing the cells under non-static conditions. Transfer of *EM-P* between media containing different salts or from an orbital shaker to static conditions led to a reversible change in cell morphology.

A considerable amount of membrane debris (Fig. S12) and extracellular DNA (eDNA) was observed during all growth stages (Fig. 7 & S15). These observations imply that *EM-P's* reproduction is leaky, resulting in a significant loss of intracellular constituents. DNA could

have been released into the surroundings when the cell underwent lysis to release the

daughter cell (Fig. 6h, S5 & video 14-18) or was a result of cells undergoing explosive lysis

during all growth stages (Fig. S16 & video 21). To our knowledge, such an explosion of

bacterial cells was not reported previously. However, a similar rupture of LVs by two

different mechanisms is currently known (69,70). This lysis of *EM-P* could have been caused

either due to osmotic pressure created by Reactive Oxygen Species (ROS)(70) or by

interactions between DNA and lipid head groups (69). To understand the role of ROS in *EM-P* lysis, we grew *EM-P* under anaerobic conditions in media devoid of terminal electron

acceptors (fermentation conditions). ROS are generally produced by the activity of ETC (71).

Given the absence of terminal electron acceptors and the high catalase activity of *EM-P*, we

presume low ROS production and their effective neutralization under anaerobic conditions.

Given the possibility of photoinduced ROS production (72), all bottles were incubated in the

dark and imaged them using a phase-contrast microscope at fewer intervals (48h – 72h).

There were no significant differences in the amount of eDNA between the aerobic control

and anaerobic incubations (Fig. S16). Apart from the eDNA, we also observed all the end

products of cell lysis, like membrane debris, in these incubations. These results suggest that

cell lysis is a natural part of *EM-P*'s cell cycle rather than an artifact of the imaging process.

The efficiency of *EM-P* reproduction also depended on the environmental conditions. We

observed less leakage of cell constituents when cells were grown in the presence of $MgCl_2$,

compared to other salts (Fig. 7 & S15). Significantly less leakage of cell constituents was

observed when *EM-P* was grown on an orbital shaker than under static conditions (Fig. 7 &

S15). These observations suggest that the reproductive efficiency of *EM-P*-like cells could be

higher in natural environments than in our static incubations. Despite the low reproductive

efficiency, *EM-P* batch cultures always resulted in viable daughter cells (Fig. 8, S17 & S18).

Staining and quantifying cells by flow cytometry showed that most daughter cells received

DNA from the parent cell and exhibited cytoplasmic activity (Fig. 8). Moreover, inoculating

smaller daughter cells after passing through a 0.44μm filter to separate them from membrane

debris and other larger cells always resulted in growth, suggesting the viability of daughter

cells. The transfer of daughter cells from 7%DSS-TSB to media containing glucose as the

sole carbon and energy source always resulted in growth. This suggests that daughter cells

received a full complement of genes from the parent cells, which is necessary for

synthesizing all the cell components.

Throughout our study, we observed the morphology and method of reproduction was

determined by the environmental conditions. Transferring cells from salt rich to sugar rich

media (Fig. 1), changing the osmolarities (Fig. S9&S10) or salt composition within the media

(Fig. S13&S14), or the incubation conditions (static or stirring) had a significant influence on

the cell morphology, reproduction, and reproductive efficiency (Fig. 7&S15). Transfer of

EM-P between media containing different salts or from an orbital shaker to static conditions

led to a reversible change in cell morphology. This suggests that environmental conditions

rather than the information encoded in its genome determined all aspects of *EM-P*'s life

cycle. Such profound influence of environmental conditions on bacterial cell reproduction has

never been reported before. Our study also demonstrates the working with protoplasts rather

than lipid vesicles as proxy-protocells and working under natural environmental conditions

rather than well-controlled laboratory conditions provides better understanding of the

behavior of protocells.

Our current understanding of protocells results from either a bottom-up approach of building

a synthetic cell from scratch (2,3) or a less frequently used top-down approach of

transforming extant bacteria into a primitive state by reducing its genome size to a bare

minimum (73). The method of reproduction we observed in *EM-P* resembles the reproductive

process of minimal cells resulting from both these approaches (73) together with the existing

theoretical predictions on protocells (74)(Fig. S19 – S21). Despite these similarities,

suggesting protocells reproduced by this process first requires us to prove the existence of

such cells on early Earth. To date, protocells remain a theoretical possibility with no

definitive proof of their existence. To understand if protocells ever existed on early Earth, we

compared the distinctive *EM-P* morphologies with the morphologies of the earliest known

microfossils from Paleo-Archaean Eon. In this follow-up study, we demonstrated the

remarkable similarities between *EM-P* and microfossils in terms of their morphology,

elemental, and isotopic composition (Additional materials). The results of this study provide

an independent paleontological conformation of our proposition that protocells inhabiting

early Earth reproduced by a process similar to *EM-P* (10).

Conclusion

Our results suggest that reproduction is an intrinsic biophysical property of cells with an

internal metabolism. Even in the absence of canonical molecular biological mechanisms that

regulate reproduction, cells can still reproduce efficiently. In such cells, environmental

conditions play an essential role in determining their method of reproduction and

reproductive efficiency. Despite the long-held notion that protocells reproduced independent

of canonical molecular biological processes and environmental conditions influenced

protocell reproduction, ours is the first study to demonstrate this process experimentally.

Given the suitability of this process to the environmental conditions of early Earth, its

simplicity, and its efficiency, we propose that uni-lamellar or Gram-positive protocells inhabiting early Earth likely reproduced by a similar process.

Acknowledgements

We want to thank Petra Schwille for her support and scientific input throughout the work and in preparing the manuscript. Gabriella Berthal, and Markus Oster for excellent technical support. We thank the Advanced Light Microscopy Facility at EMBL, Heidelberg, Ulf Schwartz from Leica Microsystems, and colleagues at the departments of Ecological Microbiology (Bayreuth University) and of Cellular and Molecular Biophysics (Max Planck Institute for Biochemistry) for their support throughout the work.

Competing Interests

The authors declare that there are no conflicts of interest regarding the publication of this article.

Data Availability Statement

All data, materials, and methods will be shared by Dr. Dheeraj Kanaparthi upon reasonable request.

References

1. Oparin AI. Chemistry and the origin of life. R Inst Chem Rev. 1969;2:1-12.

2. Szostak JW, Bartel DP, Luisi PL. Synthesizing life. Nature. 2001;409:387-390.

3. Schwille P. Division in synthetic cells. Emerg Top Life Sci. 2019;3:551-558.

4. Kudella PW, Preißinger K, Morasch M, Dirscherl CF, Braun D, Wixforth A, et al. Fission of Lipid-Vesicles by Membrane Phase Transitions in Thermal Convection. Sci Rep. 2019;9:Article no. 18808.

5. Zhu TF, Szostak JW. Coupled growth and division of model protocell membranes. J Am Chem Soc. 2009;131:5705-5713.

6. Heard E. Molecular biologists: let's reconnect with nature. Nature. 2022;601:9.

7. van Heijenoort J. Lipid Intermediates in the Biosynthesis of Bacterial Peptidoglycan. Microbiol Mol Biol Rev. 2007;71:620-635.

8. Bouhss A, Trunkfield AE, Bugg TDH, Mengin-Lecreulx D. The biosynthesis of peptidoglycan lipid-linked intermediates. FEMS Microbiol Rev. 2008;32:208-233.

9. Wacey D, Saunders M, Kong C, Brasier A, Brasier M. 3.46 Ga Apex chert 'microfossils' reinterpreted as mineral artefacts produced during phyllosilicate exfoliation. Gondwana Res. 2016;36:296–313.

10. Kanaparthi D, Lampe M, Zhu B, Boesen T, Klingl A, Schwille P, et al. On the nature of the earliest known lifeforms. bioRxiv. 2021 Jan 1;2021.08.16.456462.

11. Leaver M, Dominguez-Cuevas P, Coxhead JM, Daniel RA, Errington J. Life without a

wall or division machine in Bacillus subtilis. Nature. 2009;457:849-853.

12. Mercier R, Kawai Y, Errington J. Excess membrane synthesis drives a primitive mode of

cell proliferation. Cell. 2013;152:997–1007.

13. Errington J. L-form bacteria, cell walls and the origins of life. Open Biol. 2013;

14. Wu LJ, Lee S, Park S, Eland LE, Wipat A, Holden S, et al. Geometric principles

underlying the proliferation of a model cell system. Nat Commun. 2020;11: Article no.

4149..

15. Monnard PA, Apel CL, Kanavarioti A, Deamer DW. Influence of ionic inorganic solutes

on self-assembly and polymerization processes related to early forms of life: Implications

for a prebiotic aqueous medium. Astrobiology. 2002;2:139–52.

16. Blake RE, Chang SJ, Lepland A. Phosphate oxygen isotopic evidence for a temperate and

biologically active Archaean ocean. Nature. 2010;464: 11029-11032.

17. Catling DC, Zahnle KJ. The Archean atmosphere. Sci Adv. 2020;6:eaax 1420.

18. Knauth LP. Salinity history of the Earth's early ocean. Nature. 1998;395:554.

19. Westall F. Barberton Greenstone Belt, Traces of Early Life BT - Encyclopedia of

Astrobiology. In: Gargaud M, Amils R, Quintanilla JC, Cleaves HJ (Jim), Irvine WM,

Pinti DL, et al., editors. Berlin, Heidelberg: Springer Berlin Heidelberg; 2011. p. 147–50.

Available from: https://doi.org/10.1007/978-3-642-11274-4_150.

20. Häusler S, Noriega-Ortega BE, Polerecky L, Meyer V, de Beer D, Ionescu D.

Microenvironments of reduced salinity harbour biofilms in Dead Sea underwater springs.

Environ Microbiol Rep. 2014;6:152-158.

21. Kanaparthi D, Conrad R. Role of humic substances in promoting autotrophic growth in

nitrate-dependent iron-oxidizing bacteria. Syst Appl Microbiol. 2015;38:184–188.

22. Kanaparthi D, Reim A, Martinson GO, Pommerenke B, Conrad R. Methane emission

from feather moss stands. Glob Change Biol. 2017;23:4884-4895.

23. Cerri MR, Wang Q, Stolz P, Folgmann J, Frances L, Katzer K, et al. The ERN1

transcription factor gene is a target of the CCaMK/CYCLOPS complex and controls

rhizobial infection in Lotus japonicus. New Phytol. 2017;215:323-337.

24. Studer P, Borisova M, Schneider A, Ayala JA, Mayer C, Schuppler M, et al. The

Absence of a Mature Cell Wall Sacculus in Stable Listeria monocytogenes L-Form Cells

Is Independent of Peptidoglycan Synthesis. PLoS ONE. 2016;11:e0154925.

25. Kawai Y, Mercier R, Wu LJ, Domínguez-Cuevas P, Oshima T, Errington J. Cell growth

of wall-free L-form bacteria is limited by oxidative damage. Curr Biol. 2015;25:1613–

1618.

26. Studer P, Staubli T, Wieser N, Wolf P, Schuppler M, Loessner MJ. Proliferation of

Listeria monocytogenesL-form cells by formation of internal and external vesicles. Nat

Commun. 2016;7:13631.

27. Pelletier JF, Sun L, Wise KS, Assad-Garcia N, Karas BJ, Deerinck TJ, et al. Genetic

requirements for cell division in a genomically minimal cell. Cell. 2021;184:2430-2440.

28. Olivi L, Berger M, Creyghton RNP, De Franceschi N, Dekker C, Mulder BM, et al. Towards a synthetic cell cycle. Nat Commun. 2021;12:4531.

29. Deshpande S, Spoelstra WK, Van Doorn M, Kerssemakers J, Dekker C. Mechanical Division of Cell-Sized Liposomes. ACS Nano. 2018;12:2560-2568.

30. Litschel T, Ramm B, Maas R, Heymann M, Schwille P. Beating Vesicles: Encapsulated Protein Oscillations Cause Dynamic Membrane Deformations. Angew Chem - Int Ed. 2018;57:16286-16290.

31. Zwicker D, Seyboldt R, Weber CA, Hyman AA, Jülicher F. Growth and division of active droplets provides a model for protocells. Nat Phys. 2017; 13:408-413.

32. Vutukuri HR, Hoore M, Abaurrea-Velasco C, van Buren L, Dutto A, Auth T, et al. Active particles induce large shape deformations in giant lipid vesicles. Nature. 2020;586:52-56.

33. Morzy D, Rubio-Sánchez R, Joshi H, Aksimentiev A, Di Michele L, Keyser UF. Cations Regulate Membrane Attachment and Functionality of DNA Nanostructures. J Am Chem Soc. 2021;143:7358-7367.

34. Spahn C, Endesfelder U, Heilemann M. Super-resolution imaging of Escherichia coli nucleoids reveals highly structured and asymmetric segregation during fast growth. J Struct Biol. 2014;185:243-249.

35. Bloomfield VA. DNA condensation by multivalent cations. Biopolymers. 1997;44:269-282.

1   36. Spahn C, Grimm JB, Lavis LD, Lampe M, Heilemann M. Whole-Cell, 3D, and

2   Multicolor STED Imaging with Exchangeable Fluorophores. Nano Lett. 2019;19:500-

3   505.

4   37. McKenzie AT. Glutaraldehyde: A review of its fixative effects on nucleic acids, proteins,

5   lipids, and carbohydrates. OSF Prepr. 2019; https://doi.org/10.31219/osf.io/8zd4e.

6   38. Stokes NR, Baker N, Bennett JM, Berry J, Collins I, Czaplewski LG, et al. An improved

7   small-molecule inhibitor of FtsZ with superior in vitro potency, drug-likeproperties, and

8   in vivo efficacy. Antimicrob Agents Chemother. 2013;57:317-325.

39. Szwedziak P, Wang Q, Bharat TAM, Tsim M, Löwe J. Architecture of the ring formed

by the tubulin homologue FtsZ in bacterial cell division. eLife. 2014;3:e04601.

40. Fisher JK, Bourniquel A, Witz G, Weiner B, Prentiss M, Kleckner N. Four-dimensional

imaging of E. coli nucleoid organization and dynamics in living cells. Cell.

2013;153:882-895.

41. Tarun OB, Okur HI, Rangamani P, Roke S. Transient domains of ordered water induced

by divalent ions lead to lipid membrane curvature fluctuations. Commun Chem.

2020;3:Article number 17.

42. Karimi M, Steinkühler J, Roy D, Dasgupta R, Lipowsky R, Dimova R. Asymmetric Ionic

Conditions Generate Large Membrane Curvatures. Nano Lett. 2018; 12:7816-7821.

43. Sanborn J, Oglęcka K, Kraut RS, Parikh AN. Transient pearling and vesiculation of

membrane tubes under osmotic gradients. Faraday Discuss. 2012;161:167-176.

44. Oglecka K, Rangamani P, Liedberg B, Kraut RS, Parikh AN. Oscillatory phase separation in giant lipid vesicles induced by transmembrane osmotic differentials. eLife. 2014;3:e03695.

45. Litschel T, Ramm B, Maas R, Heymann M, Schwille P. Beating Vesicles: Encapsulated Protein Oscillations Cause Dynamic Membrane Deformations. Angew Chem - Int Ed. 2018;57:16286-16290.

46. Bobrovska N, Góźdź W, Kralj-Iglič V, Iglič A. On the Role of Anisotropy of Membrane Components in Formation and Stabilization of Tubular Structures in Multicomponent Membranes. PLoS ONE. 2013;8:e73941.

47. Jinendiran S, Dahms HU, Dileep Kumar BS, Kumar Ponnusamy V, Sivakumar N. Diapolycopenedioic-acid-diglucosyl ester and keto-myxocoxanthin glucoside ester: Novel carotenoids derived from Exiguobacterium acetylicum S01 and evaluation of their anticancer and anti-inflammatory activities. Bioorganic Chem. 2020;103:104149.

48. Simons K, Ikonen E. Functional rafts in cell membranes. Nature. 1997; 387:569-572.

49. Corradi V, Sejdiu BI, Mesa-Galloso H, Abdizadeh H, Noskov SY, Marrink SJ, et al. Emerging Diversity in Lipid-Protein Interactions. Chem Rev. 2019;119:5775-5848.

50. Stachowiak JC, Brodsky FM, Miller EA. A cost–benefit analysis of the physical mechanisms of membrane curvature. Nat Cell Biol. 2013;15:1019-1027.

51. Liu Y, Agudo-Canalejo J, Grafmüller A, Dimova R, Lipowsky R. Patterns of flexible nanotubes formed by liquid-ordered and liquid-disordered membranes. ACS Nano. 2016; 10:463-474.

52. Yanagisawa M, Imai M, Taniguchi T. Shape deformation of ternary vesicles coupled with phase separation. Phys Rev Lett. 2008; 148:102-104.

53. Lipowsky R. Budding of membranes induced by intramembrane domains. J Phys II. 1992; 2:1820-1840.

54. Jülicher F, Lipowsky R. Domain-induced budding of vesicles. Phys Rev Lett. 1993;

55. Baumgart T, Hess ST, Webb WW. Imaging coexisting fluid domains in biomembrane models coupling curvature and line tension. Nature. 2003; 425:821-824.

56. Bacia K, Schwille P, Kurzchalia T. Sterol structure determines the separation of phases and the curvature of the liquid-ordered phase in model membranes. Proc Natl Acad Sci U S A. 2005; 102:3272-3277.

57. Andes-Koback M, Keating CD. Complete budding and asymmetric division of primitive model cells to produce daughter vesicles with different interior and membrane compositions. J Am Chem Soc. 2011; 133:9545-9555.

58. Sáenz JP, Grosser D, Bradley AS, Lagny TJ, Lavrynenko O, Broda M, et al. Hopanoids as functional analogues of cholesterol in bacterial membranes. Proc Natl Acad Sci U S A. 2015; 112:11971-11976.

59. Liu Y, Agudo-Canalejo J, Grafmüller A, Dimova R, Lipowsky R. Patterns of flexible nanotubes formed by liquid-ordered and liquid-disordered membranes. ACS Nano. 2016;10:463-474.

60. Sandoz PA, Tremblay C, Gisou van der Goot F, Frechin M. Image-based analysis of living mammalian cells using label-free 3D refractive index maps reveals new organelle dynamics and dry mass flux. PLoS Biol. 2019;17: e3000553.

61. Sezgin E, Levental I, Mayor S, Eggeling C. The mystery of membrane organization: Composition, regulation and roles of lipid rafts. Nat Rev Mol Cell Biol. 2017;10:361-374.

62. Strahl H, Bürmann F, Hamoen LW. The actin homologue MreB organizes the bacterial cell membrane. Nat Commun. 2014;5: Article no. 3442.

63. Li Y, Lipowsky R, Dimova R. Membrane nanotubes induced by aqueous phase separation and stabilized by spontaneous curvature. Proc Natl Acad Sci U S A. 2011;108:4731-4736.

64. Sanborn J, Oglęcka K, Kraut RS, Parikh AN. Transient pearling and vesiculation of membrane tubes under osmotic gradients. Faraday Discuss. 2012;161:161-176.

65. Dell'Era S, Buchrieser C, Couvé E, Schnell B, Briers Y, Schuppler M, et al. Listeria monocytogenes l-forms respond to cell wall deficiency by modifying gene expression and the mode of division. Mol Microbiol. 2009;73:306-322.

66. Tsai HHG, Lai WX, Lin HD, Lee JB, Juang WF, Tseng WH. Molecular dynamics simulation of cation-phospholipid clustering in phospholipid bilayers: Possible role in stalk formation during membrane fusion. Biochim Biophys Acta - Biomembr. 2012;1818: 2742-2755.

67. Blöchliger E, Blocher M, Walde P, Luisi PL. Matrix effect in the size distribution of fatty acid vesicles. J Phys Chem B. 1998;102:10383–10390.

68. Crawford EB, Arbic BK, Sheldon ND, Ansong JK, Timko PG. Investigating the behavior

of mid-Archean tides and potential implications for biogeochemical cycling. Precambrian

Res. 2022;380:106799.

69. Angelova MI, Hristova N, Tsoneva I. DNA-induced endocytosis upon local

microinjection to giant unilamellar cationic vesicles. Eur Biophys J. 1999;28(2):142–50.

70. Zhu TF, Szostak JW. Exploding vesicles. J Syst Chem. 2011;2: Article no.4.

71. Muller FL, Liu Y, Van Remmen H. Complex III releases superoxide to both sides of the

inner mitochondrial membrane. J Biol Chem. 2004;279:49064-49073.

72. Dixit R, Cyr R. Cell damage and reactive oxygen species production induced by

fluorescence microscopy: Effect on mitosis and guidelines for non-invasive fluorescence

microscopy. Plant J. 2003;36:280-290.

73. Hutchison CA, Chuang RY, Noskov VN, Assad-Garcia N, Deerinck TJ, Ellisman MH, et

al. Design and synthesis of a minimal bacterial genome. Science. 2016;351:1414.

74. Ruiz-Herrero T, Fai TG, Mahadevan L. Dynamics of growth and form in prebiotic

vesicles. 2019;123:038102.

Figure abstracts

Fig. 1 Growth characteristics of *EM-P* in batch culture. *EM-P* exhibited a typical sigmoid

growth curve with an initial phase of slow growth rate (lag-phase), a log-phase with

exponentially increasing cell numbers and a gradual decline in the growth rate during the

stationary-phase. The X and Y-axis in the plot indicate the time in hours (h) and optical

density. Images a & b shows typical morphologies of cells in early and late lag-phase. Images

c-i show early log-phase cells that seems to be reproducing either by binary fission (c, d and

Video 1-3) or budding (f-e, arrows). Images f – k show cells in media isotonic to 7%DSS but

devoid of salt (but contain sucrose, BSA and DNA, to maintain appropriate osmolarity).

Morphologies of cells under these growth conditions is entirely random. Arrows in the

images point to membrane septs formed at random places. Scale bars: 5µm.

Fig. 2 Asymmetric division of *EM-P* cells. Images a & b show theoretically predicted

morphologies of vesicles with lateral membrane heterogeneity (originally published by

Bobrovska *et al*., 2013)(46). ν is the ratio between the actual volume of a vesicle to the

volume of a vesicle in its spherical form, H^B_m is the bending rigidity of the anisotropic

component (lipid-lipid or lipid-protein nanodomains within the membrane), and Φ_{tot} is the

concentration of the membrane component that differed in its intrinsic membrane curvature.

Images c-i show, phase-contrast (c-e) and SEM (h & i) images of *EM-P* cells undergoing a

similar method of reproduction. Similarities between *EM-P* cells and predicted morphologies

is indicated in image a (letters in red). Scale bars: 10μm (c-e) & 2μm (h & i).

Fig 3 Comparison between theoretically predicted steps in filament formation to

observation in *EM-P*.

Images a-m show images of *EM-P* cells from different growth stages with external (c-f) or

internal membrane (e-m) structures that were formed in a pattern similar to what was

theoretically predicted (n & o). Arrows in images a & b point to late-log phase *EM-P* cells

with tiny membrane extensions. Image-c shows the cell surface covered with short

filamentous extensions. Image - d shows cells transferred to 15%DSS with longer

filamentous extensions. Cells in these images were stained with universal membrane stain,

FM™5-95 (white), and DNA stain, PicoGreen™ (red). Images e-i show sequential stages observed in the development of intracellular vesicles. Images e-I show vesicles forming at the periphery of the cells before their detachment and migration into the cell. Images j-m show optical sections of a Nile red-stained *EM-P* cell at different planes. Arrows in these images point to the interconnected nature of these vesicles, similar to $L^{(3)}$ and $L^{(4)}$ in image-o. Images n & o were originally published by Liu *et al.,* 2015(51). Image-n shows the energy landscape of different equilibrium shapes as a function of reduced volume (v). ΔE indicates deflation-induced (or excess membrane-induced) reduction in the membrane bending energy. Image-o shows the shapes of filamentous extensions corresponding to eight vertical lines in image-n. (please refer to Figure 5 in Liu *et al.,* 2015(51) for further information). Scale bars: 10μm.

Fig. 4 Membrane phase-separation in *EM-P*. Membrane phase-separation in *EM-P*

Cells in these images were stained with FM™5-95 (all membrane, red) and *FAST*™ DiI (L_d-

membrane, green) and imaged either using a super-resolution STED microscope (a-j) or a

spinning disk confocal (k). Images a-c, d-f, and g-i are images of the same cells in different

confocal channels. Open arrow in a-f, point to small patches of L_d-membrane on the cell

surface ($\approx$ 180nm, beyond the diffraction limit of the conventional microscopes) (a-c). Closed

arrows in d-i: L_o-membrane extensions out of the cell. Image j is the 3D-rendered image of

intracellular vesicles composed of L_d-membrane within the cell. A series of optical sections

of the stacks is shown in Fig. S8. Image k shows *EM-P* cells with filamentous extensions, as

shown in Figure 3c, entirely composed of L_o membrane. Scale bars: 10µm (a-i) & 5µm (j &

k).

Fig. 5 Life cycle of *EM-P* reproducing via formation of extracellular daughter cells.

Image a & b show a late log phase cell with bud like structures on *EM-P* cell surface. Image-

c show a late log phase cell with open arrow pointing to surface buds and a closed arrow

pointing towards filamentous extensions. Image-d shows a cell from a subsequent growth

stage cells with multiple filamentous extensions (also see video 5). Images e-f show

formation of daughter cells as "string-of-beads" attached to a parent *EM-P* cell (boxed

region) (also see video 6-8). Images f-i show the detachment of these strings of daughter

cells and their subsequent fragmentation into individual daughter cells (also see video 9-11).

Scale bars: 10µm (a-i).

Fig. 6 The life cycle of *EM-P* reproducing via the formation of intracellular daughter

cells. Images a-h is TEM images of the life cycle of *EM-P* reproducing by the formation of

intracellular daughter cells. Image-a show the process by which intracellular (white arrows)

vesicles and bud-like structures are formed (black arrows). Closed red arrows in images b & c

point to the narrow cytoplasmic region of the cell. Arrows in images d-f points to the

formation of individual and string of daughter cells or string of multiple daughter cells.

Image-g shows internal daughter cells with intracellular vesicles. Image-h shows the daughter

cell release process by rupturing the vesicle membrane (open red arrows). Scale bars: 1µm.

Fig. 7 Leakage of cell constituents during *EM-P*'s reproduction. Image a-c shows lysis of

EM-P cells and release of intracellular DNA (red). Cells in these images were stained with

FM™5-95 (membrane, white) and PicoGreen™ (DNA, red) and imaged using a STED

microscope. Image-d shows a similar process in the biofilm from the static incubations.

Highlighted regions in this image show hollow intracellular vesicles released by the lysis of

cells. Image-e show late log-phase *EM-P* cells with considerable amount of extracellular

DNA, hollow intracellular vesicles (boxed regions), and membrane debris (arrows). Scale

bars: 10μm. Image-f & g show the results of the quantification of DNA leakage during *EM-*

P's late log growth stage (Figure S15, for all growth stages) grown under static conditions

and on an orbital shaker respectively. wt – wild type (*Exiguobacterium strain M* with a cell

wall); DSS, KCl & MgCl₂: *EM-P* grown in TSB amended with 7% of the respective salts.

Data in images f & g represent mean values of biological repetitions from different batches of

EM-P's inoculum's (n=5).

Fig. 8 Metabolic viability of daughter cells. Panel-a shows the workflow of the experiment. Image-b shows the results of this quantification of these cells by flow cytometry. The green bar shows the percentage of cells containing both an intact membrane and intracellular DNA (FM^{TM}5-95 & PG). The subsequent blue, yellow, and red bars show the percentages of cells that were either deficient in intracellular DNA or an intact membrane or both respectively. Image-c shows the results of this quantification of different cell population flow cytometry. The green bar shows the percentage of cells containing both an intact membrane and intracellular enzyme activity (FM^{TM}5-95 & CTV). The subsequent blue, yellow and red bars show the percentages of cells that were either deficient in intracellular enzyme activity or an intact membrane or both respectively. The original flow cytometry plots together with the confocal microscope images of daughter cells were shown in Figures S17 & S18.