1

Meisosomes, folded membrane platforms, link the epidermis to the cuticle in *C. elegans*

Dina Aggad[1#], Nicolas Brouilly[2#], Shizue Omi[1#], Clara L. Essmann[3,$], Benoit Dehapiot[2], Cathy Savage-Dunn[4],
Fabrice Richard[2], Chantal Cazevieille[5], Kristin A. Politi[6], David H. Hall[6], Remy Pujol[5] and Nathalie Pujol[1]*

[1]Aix Marseille Univ, INSERM, CNRS, CIML, Turing Centre for Living Systems, Marseille, France
[2]Aix Marseille Univ, CNRS, IBDM, Turing Centre for Living Systems, Marseille, France
[3]UCL, University College London, Department of Computer Science, Engineering Building, Malet Place,
London, WC1E 7JG UK
[4]Department of Biology, Queens College and the Graduate Center, CUNY, 65-30 Kissena Blvd, Flushing, NY
11367, USA.
[5]INM, Institut des Neurosciences de Montpellier, Plateau de microscopie électronique, INSERM, Université
de Montpellier, Montpellier, France
[6]Department of Neuroscience, Albert Einstein College of Medicine, Bronx, NY 10461, USA.

# equal contribution

[$] present address: Bio3/Bioinformatics and Molecular Genetics, Albert-Ludwigs-University, Freiburg,
Germany

* Author for correspondence

Abstract

Apical extracellular matrices (aECMs) form a physical barrier to the environment. In *C. elegans*, the
epidermal aECM, the cuticle, is composed mainly of different types of collagen, associated in circumferential
ridges separated by furrows. Here, we show that in mutants lacking furrows, the normal intimate connection
between the epidermis and the cuticle is lost, specifically at the lateral epidermis, where, in contrast to the
dorsal and ventral epidermis, there are no hemidesmosomes. At the ultrastructural level, there is a profound
alteration of structures that we term "meisosomes", in reference to eisosomes in yeast. We show that
meisosomes are composed of stacked parallel folds of the epidermal plasma membrane, alternately filled with
cuticle. We propose that just as hemidesmosomes connect the dorsal and ventral epidermis, above the
muscles, to the cuticle, meisosomes connect the lateral epidermis to it. Moreover, furrow mutants present
marked modifications of the biomechanical properties of their skin and exhibit a constitutive damage response
in the epidermis. As meisosomes co-localise to macrodomains enriched in phosphatidylinositol (4,5)
bisphosphate, they might act, like eisosomes, as signalling platforms, to relay tensile information from the
aECM to the underlying epidermis, as part of an integrated stress response to damage.

33 Introduction

34 Apical extracellular matrices (aECMs) are associated with all epithelia and are essential for animal life. In 35 *C. elegans*, a collagen-rich aECM covers the entire surface of the worm, and is called the cuticle. It is a 36 complex multilayer structure that acts as an exoskeleton, to which body-wall muscles are connected via 37 structures called hemidesmosomes that traverse the intervening epidermis (Davies & Curtis, 2011; Johnstone, 38 2000; Page & Johnstone, 2007). Specific subsets of the more than 170 collagens are enriched in the different 39 layers of the cuticle. Some are involved in the formation of distinct structures, including the circumferential 40 parallel furrows that cover the entire animal (Cox & Hirsh, 1985; Cox et al., 1980; McMahon et al., 2003; 41 Page & Johnstone, 2007; Thein et al., 2003), and the longitudinal lateral alae. The latter have been proposed 42 to be involved in facilitating the traction of *C. elegans* to its substrate during locomotion, although, notably 43 they are not present from the L2 through the L4 larval stages (Cox et al., 1981; Katz et al., 2022).

44 The cuticle also constitutes a physical barrier, protecting the underlying epidermal syncytium from biotic and 45 abiotic stresses. When the cuticle is damaged, mechanically or through infection, the epidermis reacts, 46 activating an immune response, reflected by the increased expression of antimicrobial peptide (AMP) genes, 47 including those of the *nlp-29* cluster (Belougne et al., 2020; Pujol, Cypowyj, et al., 2008; Taffoni et al., 2020). 48 An elevated level of AMP gene expression is also observed in a subset of mutants that affect the cuticle, 49 specifically those that lack furrows (Dodd et al., 2018; Pujol, Zugasti, et al., 2008; Zugasti et al., 2016). These 50 6 known furrowless mutants (*dpy-2*, *3*, *7*, *8*, *9*, and *10*) exhibit other characteristic physiological alterations, 51 including an activation of detoxification genes, dependent on the Nrf transcription factor SKN-1, and the 52 induction of genes required for osmolyte accumulation (Dodd et al., 2018).

53 If the pathway leading to AMP induction in the epidermis is well described (reviewed in (Martineau et al., 54 2021)), exactly how the epidermis senses cuticular damage remains obscure. Part of the induction seen in 55 *dpy-10* mutants is the consequence of an increase in the levels of hydroxyphenyllactic acid (HPLA). This 56 metabolite, derived from tyrosine by transamination and reduction, activates the G-protein coupled receptor 57 (GPCR) DCAR-1 (Zugasti et al., 2014), switching on a signalling cascade that leads to AMP gene expression 58 (Polanowska et al., 2018; Zugasti et al., 2016). What provokes elevated HPLA levels in *dpy-10* mutants is, 59 however, currently unknown. Further, the HPLA/DCAR-1 signalling pathway only accounts for part of the 60 elevated *nlp-29* expression seen in furrow collagen mutants (Zugasti et al., 2014). We have therefore proposed 61 that a hypothetical, cuticle-associated, damage sensor exists that would control, in an as yet undefined 62 manner, AMP gene expression. This damage sensor would also function to coordinate antimicrobial responses 63 with the distinct detoxification and hyperosmotic responses that are simultaneously activated in furrow 64 collagen mutants (Dodd et al., 2018; Rohlfing et al., 2010; Wheeler & Thomas, 2006).

In yeast, eisosomes, single invaginations of the plasma membrane underneath the aECM, the cell wall, are 66 responsible for detecting changes in nutrient availability, but also cell wall integrity and membrane tension.

They relay information primarily via BAR domain proteins to induce the appropriate responses to
environmental changes (Appadurai et al., 2020; Lanze et al., 2020; Moseley, 2018). While nematodes lack
eisosomes, the apical plasma membrane of the epidermis, which is in direct contact with the aECM (the
cuticle), is characterised by localised regions of folds that can be observed by electron microscopy (Wood,
1988). Given their superficial similarity, we refer to these structures as meisosomes, for multifold-eisosomes.
In this study, we undertook a detailed ultrastructural analysis of meisosomes in adults, as well as
characterising them during development, and in furrowless mutants. This mutant analysis proposed a role for
meisosomes in maintaining the structural integrity of the cuticle and the epidermis, and has opened the way to
future, more detailed, characterisation of their function.

Results

Meisosomes: epidermal plasma membrane folds interfacing the apical ECM

The stacked organelles that we refer to as meisosomes were mentioned during early electron microscopy
characterisation of *C. elegans* (Wood, 1988). A survey of the long transverse transmission electron
microscopy (TEM) series "N2U" from the MRC archive (White et al., 1986), which is of a 4-5 day old adult
hermaphrodite, found hundreds of meisosomes across the 400 odd available transverse prints in the midbody.
As a first step in the detailed investigation of meisosomes, we undertook a focused TEM study to determine
their 2D organization and their 3D structure. Meisosomes are repeated folded structures at the interface of the
aECM (the cuticle) and the epidermis (Figure 1). They can be found in similar locales at all larval stages,
predominantly in the epidermal syncytium hyp7 at the lateral, dorsal and ventral ridges, and in the tail tip
epidermal cells. They are not present on the basal side of the epidermis, nor in the seam cells, the rectal
epithelia, nor in the pharynx (Figure 1 & Figure 1-figure supplement 1). In adults, meisosomes typically
comprised 4 to 10 closely apposed parallel folds of the plasma membrane, although we observed some with
up to 30 folds (Figure 1C). Most folds formed an indentation 200-400 nm deep (Figure 1C-G). The folds were
regularly spaced. The gap between each cytoplasmic-facing plasma membrane fold was 35 nm, 75 % wider
than that between the folds made from cuticle-facing plasma membrane (20 nm) (Figure 1G). The
cytoplasmic faces of the folds were free of ribosomes but contained dense material close to the plasma
membrane, separated by a thin less electron-dense area (Figures 1G). Meisosomes were frequently found in
close proximity to mitochondria (85 %, n=355) (Figures 1C-E). On their apical side, some folds were found
close to a furrow (Figure 1C&E). Although very variable in a single worm, meisosomes of similar dimensions
were observed in both transverse and longitudinal sections (Table 1), consistent with a random orientation
relative to the animal's antero-posterior axis. This random orientation was clearly visible in electron
micrographs of freeze-fractured samples (Figure 1-figure supplement 2A-B). It contrasted with a much more
regular pattern in moulting larvae, in which meisosomes were in-between the position of circumferential
furrows, (Figure 1-figure supplement 2C-D). As described below, a similar organisation could be observed

through moulting using *in vivo* markers. Some much smaller meisosomes, typically with only 2-4 shallow
folds were seen in the thin epidermal tissue that lies between body-wall muscles and the cuticle (dorsal and
ventral epidermis, see Figure 1A) and that is largely devoid of cytoplasmic content (Figure 1-figure
supplement 3).

To understand meisosomes' 3D structure, we undertook a tomographic analysis on serial 350 nm-thick
sections. This approach confirmed the existence of groups of parallel folds, all found in continuity with the
plasma membrane (Figure 2). The tomographic analysis also revealed variability in the geometry of the folds.
Although most groups of folds were perpendicular to the apical surface, some were tilted. The folds had a
relatively uniform depth, but were deeper at the centre of each stack. No break in the plasma membrane was
observed on neither the apical nor the basal side of the meisosomes. Despite their close apposition with
mitochondria, no membrane continuity was observed between meisosomes and mitochondria (Figure 2A-D,
Figure 2 - Video 1).

In order to evaluate not just the topology, but also the distribution of these organelles, we developed a fixation
protocol for serial block-face scanning electron microscopy (SBF-SEM) of *C. elegans* samples. Starting with
protocols previously described (Deerinck et al., 2010; D.H. Hall et al., 2012; D. H. Hall et al., 2012), we
adapted the solvents and temperatures for each contrasting agent, including lead and uranyl acetate, to
maximize sample contrast (see Materials and Methods). We acquired series of electron micrographs of the
lateral epidermis as transversal views along 12 µm. We produced a voxel-based classification within the
Waikato Environment for Knowledge Analysis (WEKA in Fiji) and then used its machine learning algorithms
to perform semi-automated organelle recognition. This revealed that the meisosomes were irregularly spaced
at the apical surface of the lateral epidermis, with various sizes and orientations and confirmed their frequent
apposition to mitochondria (Figure 2-figure supplement 1).

VHA-5 is a marker of meisosomes

VHA-5, one of four α-subunits of the transmembrane V0 complex of the vacuolar ATPase (V-ATPase) (Oka
et al., 2001; Pujol et al., 2001), and RAL-1, ortholog of human RALA (RAS like proto-oncogene A) (Frische
et al., 2007), are the only known markers of meisosomes, with both proteins being also associated with
multivesicular bodies (MVB) that play a role in exosome secretion (Hyenne et al., 2015; Liegeois et al.,
2006). The expression pattern of VHA-5 is the better characterised of the two proteins. A TEM/immunogold
staining study showed that more than 85 % of the VHA-5 signal can be attributed to meisosomes (Liegeois et
al., 2006).

We have used several VHA-5 reporter strains, including one expressing a GFP-tagged version of VHA-5 from
a CRISPR/Cas9 engineered genomic locus [KI], a single copy insertion under an epidermis promoter [Si], or
classic multi-copy integrated [Is] or extrachromosomal [Ex] transgenic arrays. In all the strains, and in line

with previous reports (Liegeois et al., 2006), we observed the same punctate fluorescence at the apical surface
of the lateral epidermis, and in the ventral and dorsal ridges, from the head to the tail. As expected, it was
almost completely absent from the dorsal and ventral epidermis above the body-wall muscles, and underneath
the seam cells (Figure 3A-C). By combining a MUP-4::GFP (Suman et al., 2019) and a VHA-5::RFP reporter
(Liegeois et al., 2006), we confirmed the complementary pattern in the epidermis of hemidesmosomes above
the muscles, and meisosomes in the lateral epidermis and dorso/ventral ridges (Figure 3D). Higher resolution
analysis in the lateral epidermis revealed the VHA-5-marked structures to have an irregular shape in no
preferred direction in the lateral epidermis, consistent with the TEM and SBF analyses (Figure 3E).

When we examined worms co-expressing VHA-5::GFP and HGRS-1::mScarlet, a known marker of MVBs
and the endosome degradation pathway (Liegeois et al., 2006; Norris et al., 2017; Roudier et al., 2005;
Serrano-Saiz et al., 2020), we observed essentially no co-localisation. Further, the patterns of the two markers
were distinct, with HGRS-1::mScarlet labelling structures that were less apical, larger and more scattered that
those labelled by VHA-5::GFP (Figure 3F & Figure 3-figure supplement 1A). We also did not observe any
co-localisation between VHA-5::GFP and SNX-1::mScarlet, a marker for the recycling endosomes, nor with
LGG-1::mScarlet, an autophagolysosome marker (Serrano-Saiz et al., 2020) (Figure 3-figure supplement
1B&C). We have previously shown that the plasma membrane of the epidermis in young adult worms
contains heterogeneous macrodomains that can be revealed with a prenylated, or a pleckstrin homology
domain-tagged form of GFP (GFP::CAAX and GFP::PH-PLC1δ, respectively) (Taffoni et al., 2020).
Interestingly, in worms co-expressing VHA-5::RFP and either one of these membrane probes, we observed a
high degree of co-localisation (Figure 3G&H & Figure 3-figure supplement 1D-F). This further reinforces the
notion that VHA-5 is primarily a marker of subdomains of the plasma membrane.

The structures labelled with both VHA-5::GFP and CAAX::GFP or PH-PLCδ::GFP in the adult epidermis
were similar in size and spatial distribution to the meisosomes reconstituted from the SBF data (Figure 2-
figure supplement 1D). To confirm that the observable fluorescence signal from VHA-5::GFP indeed
primarily originated from meisosomes, we performed Correlation Light and Electron Microscopy (CLEM)
using the VHA-5::GFP [Si] strain in which the strong and potentially confounding excretory canal GFP signal
is absent, due to the use of an heterologous epidermis-specific promoter. As we used a different fixation
technique to preserve the GFP signal (Johnson et al., 2015) and we worked on semi-thin section, meisosomes
were revealed by electron tomography. After alignment of the confocal and TEM images, we could show that
the fluorescence foci matched meisosomes (Figure 3I-J & Figure 3-figure supplement 2). Together with
previous CLEM observations in the excretory duct (Kolotuev et al., 2009), these results indicate that the
VHA-5::GFP signal that we observe at the apical membrane in the epidermis corresponds to meisosomes and
that VHA-5::GFP can be used *in vivo* as a *bona fide* meisosome marker for this study.

Meisosomes align in between furrows before moulting

VHA-5 has been shown to have an essential role in alae formation and secretion of Hedgehog-related peptides
through exocytosis via MVBs, but not to be involved in secretion of the collagen DPY-7, nor in meiososome
morphology (Liegeois et al., 2006). Indeed, to date, no gene has been assigned a role in determining
meiososome structure. As a path to understanding their function, we first observed their morphology during
development. Consistent with previous reports (Liegeois et al., 2007), we observed that VHA-5::GFP aligns
parallel to the actin fibres in animals entering the L4/adult moult, a stage we refer to here as "late L4". We
refined this observation by precisely staging the worms throughout the L4 stage on the basis of vulval
morphology and the shape of the lumen as previously described in (Cohen et al., 2020; Mok et al., 2015)
(Figure 4A & 5). The parallel circumferential alignment of VHA-5::GFP could be observed at the beginning
of the L4 stage, in L4.1 worms, but was then lost at the L4.2 stage. It reappeared progressively starting in the
L4.3 stage, culminating between the L4.7 and L4.9 stages, just before the moult. This is consistent with the
EM observations of meiososome alignment before moulting (Figure 1-figure supplement 2C-D).

As meiososomes, like the rest of the apical plasma membrane, are in direct contact with the aECM, the cuticle,
we investigated the relation of meiososomes to different cuticle components. Different classes of cuticular
collagen exist that form either the circumferential constricted furrows, or the cuticle in the regions between
the furrows, called the annuli (Cox & Hirsh, 1985; Cox et al., 1980; McMahon et al., 2003; Page & Johnstone,
2007; Thein et al., 2003). As revealed with one marker of a furrow collagen, DPY-7::GFP (Miao et al., 2020),
in combination with VHA-5::RFP, meiososomes align in between the furrows at the late L4 stage (Figures 4B).
Notably, during the L4.7 stage, some DPY-7::GFP can be observed in small vesicles on the apical side of the
epidermis, that could represent the ongoing secretion of furrow collagen at that stage. Interestingly, these
vesicles do not co-localise with VHA-5::RFP (Figure 4B, right panel). We further show that the CAAX and
PH-PLC1δ markers that co-localise with VHA-5 in young adult animals (Figure 3G-H) also align during the
L4 stage, but other vesicular components, like the one marked by HGRS-1, SNX-1 or LGG-1 do not (Figure
4C-D). Thus, meiososomes together with specific membrane subdomains align in between the furrow before
moulting.

Furrow collagens determine the organisation of the cytoskeleton and meiososomes in L4 larvae

Before moulting, there is profound reorganisation of the cytoskeleton in the lateral epidermis. Microtubules
and actin fibres align in a series of circumferential bands that are not present in adults (Castiglioni et al., 2020;
Costa et al., 1997; Taffoni et al., 2020). Interestingly, meiososomes exhibited the same sequence of dynamic
changes in alignment as microtubules and actin. After the moult, there was a concomitant loss of alignment of
VHA-5, actin and microtubules, so that in wild-type adult animals, as described above, there was no clear
overall pattern to the organisation of meiososomes, actin or microtubules (Figures 5).

We then examined the consequence of knocking down the expression of one of the furrow collagen genes,
dpy-7, on the organised VHA-5::GFP pattern of late L4s worms. Compared to the control, RNAi of *dpy-7*

provoked a loss of meisosome alignment (Figures 5 and Figure 5-figure supplement 1). A similar phenotype
was observed in *dpy-3* mutant worms; DPY-3 is another furrow collagen (Figure 5-figure supplement 1B).
Strikingly, this loss of expression of furrow collagens was also associated with a disruption of the normal
organized pattern of both actin fibres and microtubules from the L4.3 stage (Figure 5). It was previously
proposed that the formation of actin fibres and microtubules in apposed circumferential bands plays an
instructive role in positioning the furrows (Costa et al., 1997). Our results suggest, on the contrary, that furrow
collagens in the cuticle govern the alignment of the underlying cytoskeleton as development progresses into
the final moult. Thus, furrow collagens appear to be required to align both meisosomes and the actin-
microtubule cytoskeleton.

Abnormal meisosomes in adult furrow mutant worms

As previously mentioned, different classes of cuticular collagen exist that are expressed and form either
furrows or annuli (Figure 6A) (Cox & Hirsh, 1985; Cox et al., 1980; McMahon et al., 2003; Page &
Johnstone, 2007; Thein et al., 2003). While mutant in these collagens all have a Dumpy (i.e. short and fat;
Dpy) phenotype, only the furrow-less mutants, in contrast to mutants of annuli collagens, exhibit an increased
expression of the AMP reporter *nlp-29*p::GFP (Dodd et al., 2018; Pujol, Zugasti, et al., 2008; Zugasti et al.,
2014; Zugasti et al., 2016). This is one reason that furrow collagens have been proposed to be part of a
damage sensor that relays information about cuticle integrity to the epidermis (Dodd et al., 2018).
Interestingly, this reporter is also induced at the late L4 stage in the wild type before the last moult when the
cuticle is reshaping (Figure 6-figure supplement 1A; (Miao et al., 2020)).

We examined the consequences of knocking down the expression of all these different collagen genes on the
pattern of meisosomes in the adult. Collagen inactivation was always confirmed by observing the effect on
body size, as well as the change in the expression of *nlp-29*p::GFP in parallel experiments (Figure 6-figure
supplement 1B-C). Compared to control RNAi or to annuli collagen inactivation (*dpy-4, dpy-5* and *dpy-13*),
inactivation of all 6 furrow collagen genes (*dpy-2, dpy-3, dpy-7, dpy-8, dpy-9* and *dpy-10)* provoked a marked
and specific alteration in the pattern of VHA-5::GFP. The meisosomes' normal reticulated pattern was
fragmented, as reflected by a decrease in their average size and Feret's diameter, and a >25 % increase in their
density (Figures 6B-C). A similar fragmentation was observed with the different VHA-5 reporter strains, and
either following inactivation of the furrow collagen gene's expression with RNAi or in null mutants (Figure 6-
Source data file 1).

To test if the fragmentation was associated with a change in VHA-5's relation to other organelles, we
inactivated furrow collagen genes in strains combining VHA-5::GFP and different mScarlet-tagged vesicular
membrane markers, LGG-1 for autophagosomes, HGRS-1 for multivesicular bodies and SNX-1 for recycling
endosomes. In contrast to VHA-5::GFP, we observed no marked alteration in their patterns, and there was still
no overlap between the signal from VHA-5::GFP (Figure 6-figure supplement 2A). Further, neither *dpy-3* nor

dpy-7 inactivation had any effect on the size and density of the vesicular pattern of EEA-1::GFP (Figure 6-
figure supplement 2B), a marker of early endosomes (Shi et al., 2009). Thus, loss of furrow collagen gene
expression leads to a substantial fragmentation of meisosomes, without affecting vesicular organelles in the
epidermis. This suggests that furrow collagens play an important and specific role in maintaining meisosome
integrity.

Furrow mutants, with small meisosomes, display a detached cuticle

The cuticle is connected, through the epidermis, to the underlying body-wall muscles via hemidesmosomes.
The maintenance of hemidesmosome integrity is vital; their complete loss causes a fully penetrant lethality.
On the other hand, partial loss of the hemidesmosome component MUA-3 causes the cuticle to detach above
the muscles (Bercher et al., 2001). We have shown above that hemidesmosomes and meisosomes are present
in complementary and non-overlapping regions of the epidermis (Figure 3B-D). This suggest that
hemidesmosomes cannot play a role in the attachment of the lateral epidermis and the dorso/ventral ridge to
the cuticle. As the meisosomes, containing numerous folds of the plasma membrane, increase the surface
contact between the epidermis and the cuticle, we asked whether meisosome fragmentation could impact the
attachment of the epidermis to the cuticle. Examination by TEM first confirmed that the meisosomes are
significantly smaller in furrow collagen mutants compared with another Dpy mutant (*dpy-13*) or the wild-
type, irrespective of the direction of the section, as both longitudinal and sagittal sections show the same
phenotype (Figure 7A-D, Table 1 and Figure 1 for the wild type). In all furrow collagen mutants examined,
there was a frequent disruption of the contact between the epidermal plasma membrane and the cuticle, either
in the lateral epidermis or the dorso or ventral ridge, but not above the muscle quadrants (Figures 7B&G and
Figure 7-figure supplement 1). This detachment is clearly distinct from what is observed in so-called Blister
mutants where, due to the absence of the connective struts, the detachment happens between the 2 main layers
of the cuticle (Page & Johnstone, 2007). To confirm the phenotype, we analysed by SBF entire transversal
worm sections over a length of 21.5 and 34.4 µm, for a wild-type for a *dpy-2* young adult mutant worm,
respectively. This confirmed that the detachment between the cuticle and the epidermis was always found in
the furrow collagen mutant outside the region of the body-wall muscles (Figures 7H). In furrow collagen
mutants, the space between the cuticle and the underlying epidermal cell was often filled with a diverse range
of cytoplasmic content, including membrane-bound vesicles with the appearance of endosomes, lysosomes,
mitochondria, as well as electron-dense particles the size of ribosomes (Figures 8A).

To exclude the remote possibility that this detachment was an artefact linked to the different fixation protocols
used for electron microscopy, we carried out live imaging on two independent strains in which the cuticle was
labelled with a collagen tagged with mScarlet (ROL-6::mScarlet [KI]) and the epidermal plasma membrane
was labelled with GFP::CAAX or GFP::PH-PLC1δ. Compared to the wild-type, where the GFP signal is
restricted to heterogeneous macrodomains in the plasma membrane (Taffoni et al., 2020), in a *dpy-3* furrow

collagen mutant, the GFP was seen in numerous brightly-stained vesicular structures that accumulated outside
the epidermis at the level of the mScarlet cuticular signal (Figure 8B). Together, these phenotypes suggest that
the meisosomes may play an important role in attaching the cuticle to the underlying epidermal cell and that
loss of this intimate connection causes a profound alteration of epidermal integrity.

Furrow mutants have abnormal biomechanical properties

We predicted that the changes in cuticle attachment seen in the furrow mutants would impact the
biomechanical properties of worms. It was previously shown that furrows are stiffer than the rest of the cuticle
in wild-type worms (Essmann et al., 2016). We therefore used atomic force microscopy to measure the
resistance to force in wild-type and mutant worms, as previously described (Essmann et al., 2016; Essmann et
al., 2020). While topographic AFM imaging (Figures 9A) provided further corroboration of the fact that in the
absence of furrow collagens the cuticle has a disorganised aspect with irregular folds, lacking the usual
repeated linear pattern of annuli and furrows, force spectroscopy AFM revealed differences in stiffness. In
contrast to the non-furrow *dpy-13* mutant that had a rigidity similar to wild-type, the different *dpy* furrow
mutants (*dpy-2*, *dpy-3*, *dpy-7* and *dpy-8*) exhibited markedly less steep force-indentation curves (Figure 9B),
and hence lower stiffness or Young's Moduli (Figure 9C). This suggests that furrow collagens are required for
normal stiffness. While lack of certain collagens in the cuticle could directly affect cuticle stiffness, we
hypothesise that the effect on stiffness is a consequence of the fact that furrow collagens are essential for the
presence of normal meisosomes.

Discussion

In this study, we undertook the characterisation of structures that link the nematode epidermis to the cuticle.
Across species, interfaces exist between flexible and dynamic cell membranes and more rigid extracellular
matrices. Because of requirements for growth, signal transduction, and repair, the microstructures of the ECM
need to be tightly linked to the plasma membrane and cytoskeleton of the underlying cell (Chebli et al., 2021).
In yeast, eisosomes are single membrane invaginations located under the cell wall that bridge this boundary
and fulfil this function. They can disassemble in minutes to buffer changes in membrane tension, protecting
cells from osmotic shock (Lemiere et al., 2021). Eisosomes are specific to yeast; there are no orthologues for
core components, such as LSP-1, in animals. Conversely, the meisosomes that we describe here in *C. elegans*,
with their multiple membrane invaginations that individually are similar in appearance to eisosomes, are, to
the best of our knowledge, distinct from interfacial structures in non-nematode species. Interestingly, we show
here that they are enriched in a PH-PLCδ marker, which is known to bind phosphatidylinositol 4,5-
bisphosphate (PIP$_2$) (Lemmon et al., 1995). PIP$_2$ has a major role in signal transduction and in regulating
cellular processes including actin cytoskeleton and membrane dynamics (Katan & Cockcroft, 2020).
Moreover, we have previously shown that the same PH-PLCδ marker rapidly reorganises upon wounding of

the lateral epidermis (Taffoni et al., 2020). So, it is tempting to propose that analogous to eisosomes,
meisosomes could have a role as a signalling platform in response to stress.

While the presence of meisosomes had been noted in earlier studies (Hyenne et al., 2015; Liegeois et al.,
2006), we have been able to go beyond their previous characterisation, in part because of improvements in
electron microscopy techniques. Specifically, we adapted the fixation protocol after high pressure freezing to
have a better membrane contrast in serial block scanning electron microscopy, allowing semi-automated *in*
silico segmentation of meisosomes. Moreover, adapting a CLEM protocol, we were able to match the
VHA-5::GFP observed by fluorescence microscopy to meisosomes revealed by tomography. VHA-5, together
with RAL-1, are currently the only known meisosome components. In contrast to the well-defined roles of
these two proteins in alae formation and exosomes biogenesis (Hyenne et al., 2015; Liegeois et al., 2006),
their function in meisosomes remains to be characterised. Notably, inactivation of *ral-1* did not eliminate
VHA-5::GFP fluorescence in the epidermis (Hyenne et al., 2015), and knocking down the expression of *vha-5*
did not affect the secretion of DPY-7 (Liegeois et al., 2006). This suggests that the V-ATPase on meisosomes
is not involved in cuticle synthesis. Further study will be required to determine the catalogue of proteins that
are needed for meisosome formation and maintenance.

Notably, a recent study reported the isolation of mutants with an abnormal pattern of VHA-5::RFP in the
epidermis but attributed this to a change in MVBs (Shi et al., 2022), despite a lack of substantial co-
localisation with HGRS-1, a well characterised MVBs marker, part of the ESCRT-0 complex that sorts
endosomes to MVBs (Babst, 2011). Since previous studies (Liegeois et al., 2006), and the results presented
here, show that VHA-5 is predominantly a marker of meisosomes, more so than of MVBs, we hypothesise
that the one gene that Shi *et al.* characterised in detail, *fln-2*, which encodes the F-actin cross-linking protein
filamin (Zhao et al., 2019), could actually be involved in the formation and/or maintenance of meisosomes.
Interestingly, a *fln-2* loss of function mutation has been serendipitously found in several *C. elegans* strains
originating from a different wild-type stocks (Zhao et al., 2019), so careful attention to genotypes will be
needed in future work. Regardless, *fln-2* may represent an important tool to investigate meisosome function.

Setting this issue aside, by taking an ultrastructural approach, we were able to build up a detailed picture of
the organisation of meisosomes. One of their defining features is the constant 25 nm spacing of their
constituent plasma membrane folds. This raises the question of how the membrane folds with such precision.
One possibility is that the striking electron-dense material that is apposed to each side of the membrane on the
cytoplasm-facing folds, spaced less than 10 nm apart, will contain specific structural protein that maintain the
uniform width of each meisosome fold, and influence their mechanical properties. These structures will
require more precise characterisation. We equally have yet to establish whether the frequent proximity of
meisosomes to mitochondria, with a close apposition of membranes, has a functional significance.

Contrary to the cuticle of many adult insects, the nematode cuticle is flexible enough to allow bending during
locomotion. It is also thought to stretch to accommodate growth between moults. When the old cuticle is shed,
it leaves in its place the new cuticle that had been moulded underneath it. The circumferential furrows of the
new cuticle thus appear exactly in register with the position of old furrows. Before moulting, the cytoskeleton
aligns in the apical epidermis, underneath and parallel to each furrow. Although this had been proposed to be
important for positioning the furrows of the new cuticle (Costa et al., 1997; McMahon et al., 2003; Page &
Johnstone, 2007), a recent study found unexpectedly, that actin is dispensable for the alignment of furrows
(Katz et al., 2018). On the other hand, we found that the furrows are required for the alignment of actin fibres
before the last moult. We propose therefore that only the old furrows are required to pattern the new furrows.
Consistent with such a model, the LPR-3 protein that is part of the transient pre-cuticle that is formed between
the old and the new cuticles before each moult is absent from the region of the furrows (Forman-Rubinsky et
al., 2017). We have shown that furrow determines the regular parallel and circumferential positioning of
meiososomes. It is not yet clear whether this alignment of meiososomes is functionally important. It could result
from steric constraints during moulting, in the limited space between nascent furrows of the new cuticle and
the closely apposed circumferential actin fibres. It should, however, be noted that this alignment is not seen
for vesicular organelles like MVBs, endosome or autophagosomes.

As adults, furrowless collagen mutants have fragmented meiososomes and a detached cuticle. We hypothesise
that this fragmentation causes the detachment, and that the multiple folds of plasma membrane normally
increase its contact surface with the cuticle thus ensuring a more robust connection of the aECM to the lateral
epidermis. While the lateral epidermis is rich in meiososomes, it is devoid of hemidesmosomes. Conversely, in
the dorsal and ventral quadrants, there are essentially no meiososomes, but abundant hemidesmosomes. These
latter structures secure the muscles to the cuticle through epidermis and are indispensable for worm
development and viability. Above the muscles, the epidermis is extremely thin, with the apical and basal
plasma membranes juxtaposed, linked via intermediate filaments that bridge apical and basal hemidesmosome
protein complexes (Zhang & Labouesse, 2010). MUA-3 is an hemidesmosome transmembrane protein in
direct contact with the cuticle. In hypomorphic *mua-3* mutants, large gaps form between the apical epidermal
surface and the cuticle in the dorso-ventral quadrants, reflecting a loss of attachment of apical
hemidesmosomes to the cuticle. Unlike the cytoplasm-filled gaps we observed in furrowless mutants, in *mua-*
3(rh195) worms, these spaces appear devoid of contents, and the apical epidermal membrane is intact
(Bercher et al., 2001). So, in contrast to the loss of hemidesmosomes, fragmentation of meiososomes in
furrowless mutants affects the integrity of the apical epidermal membrane in the lateral epidermis, potentially
explaining the permeability phenotype of furrowless mutants (Sandhu et al., 2021). Despite these differences,
both meiososomes in the lateral epidermis, and hemidesmosomes in the dorso-ventral quadrants, do appear to
have an analogous function, ensuring the attachment of the apical plasma membrane to the cuticle.

In animals, ECMs provide mechanical support for tissue assembly and organ shape. During embryogenesis in
C. elegans, the aECM is essential during elongation as it not only maintains embryonic integrity, but also
relays the mechanical stress produced by the actomyosin cytoskeleton and the muscles (T. T. Vuong-Brender
et al., 2017; T. T. K. Vuong-Brender et al., 2017). In the adult, the mechanical properties of the aECM have
only recently started to be explored. Atomic force microscopy revealed that the furrows have a higher
stiffness than the annuli (Essmann et al., 2016). Here, we show that loss of specific furrow collagens, but not
of non-furrow collagens, decreases stiffness. Part or all of this could be a direct consequence of the altered
cuticle morphology, an analogy being the increased stiffness that corrugation provides. Furrow Dpy mutants
are known to have a higher internal concentration of glycerol (Wheeler & Thomas, 2006), which will decrease
their internal hydrostatic pressure. We propose that this decreased hydrostatic pressure is a consequence of the
decrease in the stiffness of the cuticle. It would ensure the necessary balance of inward and outward pressures
required for body integrity. Since we used a 10 micrometer diameter AFM probe to indent the worm, and the
indentation depth was greater than the thickness of the cuticle (ca. 800 nm compared to 500 nm for the
cuticle), our measurements did not directly assess the cuticle stiffness, so further investigations will be needed
to confirm our hypothesis. It is interesting to note, however, that a decrease in stiffness and an increase in the
activity of innate immune signalling pathways in the epidermis are signatures of ageing in *C. elegans* (E et al.,
2018; Essmann et al., 2020). How physiological and pathological modifications of the biomechanical
properties of the aECM are surveyed by the epidermis remains an open question for future studies.

Materials and Methods

Nematode strains

All *C. elegans* strains were maintained on nematode growth medium (NGM) and fed with *E. coli* OP50, as
described (Stiernagle, 2006), the wild-type N2, IG274 *frIs7[col-12p::DsRed, nlp-29p::GFP]* IV (Pujol,
Cypowyj, et al., 2008), IG1697 *dpy-2(e8) II; frIs7[nlp-29p::GFP, col-12p::DsRed] IV*, IG1685 *dpy-3(e27) X;*
frIs7[nlp-29p::GFP, col-12p::DsRed] IV, IG1689 dpy-*7(e88) X; frIs7[nlp-29p::GFP, col-12p::DsRed] IV*
(Dodd et al., 2018), IG1699 *dpy-8(e130) X; frIs7[nlp-29p::GFP, col-12p::DsRed] IV*, IG344 *dpy-13(e184)*
frIs7[nlp-29p::GFP, col-12p::DsRed] IV, RT424 *pwIs126[eea-1p::GFP::EEA-1* (Shi et al., 2009), RT3657
pwSi46[hyp7p::mScarlet::HGRS-1+G418-R], RT3640 *pwSi65[hyp7p::mScarlet::SNX-1+G481-R]*, RT3635
pwSi62[hyp7p::mScarlet::LGG-1+G418-R] (Serrano-Saiz et al., 2020), ML2113
mcIs67[dpy-7p::LifeAct::GFP; unc-119(+)] V; stIs10088[hlh-1::his-24::mCherry, unc-119(+)] (Lardennois
et al., 2019) to visualise actin in larval stages, IG1813 *frSi9[pNP151(col-62p::Lifeact::mKate_3'c-nmy), unc-*
119(+) ttTi5605]/+ II; tbb-2(tj26[GFP::TBB-2]) III (Taffoni et al., 2020) to visualise actin in the adult,
IG1935 *frSi9[pNP151(col-62p::Lifeact::mKate_3'c-nmy), unc-119(+) ttTi5605] II; Is[wrt-2p::GFP::PH-*
PLC1δ, wrt-2p::GFP::H2B, lin-48p::mCherry], XW18042 *qxSi722[dpy-7p::DPY-7::sfGFP; ttTi5605] II*

(Miao et al., 2020) and MBA365 *Ex[dpy-7p::GFP::CAAX, myo-2p::GFP]* kindly provided by M. Barkoulas
(UCL).

Extrachromosomal transgenic strain [Ex] containing GFP tagged version of VHA-5 were generated by
microinjection in N2 worms of the *vha-5*pVHA-5::GFP construct pML670 (Liegeois et al., 2006) kindly
provided by M. Labouesse (LBD/IBPS) at 3 ng/µl together with *unc-122*p::GFP at 100 ng/µl to generate
IG1930. The transgene *frSi26* is a single copy insertion on chromosome II (ttTi5605 location) of pNP165
(*dpy-7*p::VHA-5::GFP) by CRISPR using a self-excising cassette (SEC) (Dickinson et al., 2015). pNP165 was
obtained by insertion of the *dpy-7* promoter, which leads to an epidermal specific expression, in front of
VHA-5::GFP into the pNP154 vector. pNP154 was made from a vector containing the SEC cassette for single
insertion on Chromosome II at the position of ttTi5605 (pAP087, kindly provided by Ari Pani) (Watts et al.,
2020). Constructs were made using Gibson Assembly (NEB Inc., MA) and confirmed by sequencing. pNP165
was injected in N2 at 20 ng/µl together with pDD122 (*eft-3*p::Cas9) at 50 ng/µl, pCFJ90 (*myo-2*p::mCherry)
at 2 ng/µl, and #46168 (*eef-1A.1*p::CAS9-SV40_NLS::3'*tbb-2*) at 30 ng/ml. pCFJ90 was a gift from Erik
Jorgensen (Addgene plasmid # 19327; http://n2t.net/addgene:19327; RRID:Addgene_19327) (Frokjaer-
Jensen et al., 2008). Non-fluorescent roller worms were selected then heat shocked to remove the SEC by
FloxP as described in (Dickinson et al., 2015) to generate IG2118 *frSi26[pNP165(dpy-7p::VHA-5::GFP)*
ttTi5605] II. Fluorescent knock-in [KI] reporter strains were generated through CRISPR editing
(SunyBiotech) at the C-terminus of the gene to generate PHX5715 *vha-5(syb5715[VHA-5::sfGFP]) IV*, that
was further outcrossed 2 times to generate IG2144, and PHX2235 *rol-6(syb2235[ROL-6::mScarlet]) II*. All
the multiple reporter strains generated in this study were obtained by conventional crosses (see Supplementary
File 1 for a list of all strains).

424 Transmission Electron Microscopy (TEM)

Day 1 adult worms were frozen in NaCl 50 mM medium containing 5 % of BSA and *E. coli* bacteria using
Leica EM Pact 2 high pressure freezer. After freezing, samples were freeze-substituted at -90 °C in acetone
containing 2 % OsO_4 for 96 hours. The temperature was gradually increased to -60 °C and maintained for 8
hours. The temperature was then raised to -30 °C and maintained for 8 hours, before to be raised again to RT.
Samples were finally washed in acetone and embedded in Epoxy resin. Resin was polymerised at 60 °C for 48
hours. 70 nm ultrathin and 350 nm semithin sections were performed using a Leica UC7 ultramicrotome and
post-stained with 2 % uranyl acetate and Reynolds' lead citrate. Images were taken with a Tecnai G2
microscope (FEI) at 200 kV. For tomography acquisitions, tilted images (+60°/-60° according to a Saxton
scheme) were acquired using Xplorer 3D (FEI) with a Veleta camera (Olympus, Japan). Tilted series
alignment and tomography reconstruction was performed using IMOD (Mastronarde, 1997).

435 Freeze fracture

Wild-type adults were fixed in buffered 2.5 % glutaraldehyde, then cryoprotected in 30 % glycerol overnight
prior to freezing. Fixed animals were positioned between two gold discs, and plunge frozen in liquid nitrogen-
chilled isopentane. Frozen worms were placed into a double replica holder for a Balzer's 301 freeze etch
device. Samples were cleaved within the freeze etch device by free breaks, then shadowed with Pt/C to form a
metal replica. Replicas were washed in bleach to remove all tissue prior to mounting on slot grids for
examination by TEM.

442 Scanning electron microscopy by Serial Block Face (SBF)

After freezing in the aforementioned conditions, samples were incubated at -90 °C in acetone containing 2 %
OsO_4 for 106 hours. The temperature was gradually increased to 0 °C and samples were washed over 1h in
acetone at RT. Samples were then incubated in acetone containing 0.1 % TCH for 60 min, washed over 1h in
acetone, and incubated in acetone containing 2 % OsO_4 for 1 hour at RT. After rehydration in ethanol
decreasing graded series, samples were incubated ON in 1 % aqueous uranyl acetate at 4 °C and in 30 nM
lead aspartate for 30 min at 60 °C. Samples were finally dehydrated in graded series of ethanol baths and pure
acetone and embedded in Durcupan resin. Resin was polymerised at 60 °C for 48 hours. For the segmentation
of meisosomes, regions of the lateral epidermis were acquired over a length of 12 µm, with a resolution of 10
nm par pixels. For scanning the cuticle detachment, entire transversal sections were acquired over a length of
21.5 and 34.4 µm, for a wild-type and a *dpy-2* mutant worm, respectively, with a resolution of 10 nm par
pixels.

454 Correlative Light Electron Microscopy (CLEM)

Sample for CLEM experiments were treated as in (Johnson et al., 2015). Briefly, the worms were high
pressure frozen (EMPACT2, Leica) and then freeze-substituted (AFS2, Leica) for 20 hours from -130 °C to -
45 °C in an acetone-based cocktail containing 0.2 % uranyl acetate, 0.1 % tannic acid and 5 % H_2O. After 2 h
of acetone washes at -45 °C, the samples were infiltrated with gradients of HM20 resin over 9 h, with pure
resin for 18 h at -45 °C and the resin was polymerized under UV for 24 h at -45 °C and for 12 h at 0 °C. 350
nm semithin sections were processed as described for TEM tomography above. TEM grids were first analysed
at by confocal imaging where a bright field image is overlaid with the fluorescent image, then analysed in
TEM at low magnification. Brightfield, GFP confocal and TEM images were aligned using AMIRA. Several
positions with 2 or 3 GFP spots were chosen to do a high magnification tomography, as described above, to
reveal the meisosomes.

465 Segmentations and 3D Image analysis

For electron tomography datasets, a binned version of the reconstructed tomogram was segmented using the
Weka 3D segmentation plugin in Fiji/ImageJ to visualize the mitochondria and the meisosomes. The cuticle

was visualized by the Amira-embedded Volume Rendering plugin from a manually segmented mask. A
cropped area of interest of the full resolution electron tomogram was segmented in iLastik to visualize a
representative portion of the organelle. For Serial Block-face datasets, the segmentation of the meisosome
mitochondria was generated using the Weka 3D segmentation plugin in Fiji/ImageJ. Animations and
snapshots were generated in Amira.

RNA interference

RNAi bacterial clones were obtained from the Ahringer or Vidal libraries and verified by sequencing (Kamath
et al., 2003; Rual et al., 2004). RNAi bacteria were seeded on NGM plates supplemented with 100 μg/ml
ampicillin and 1 mM Isopropyl-β-D-thiogalactopyranoside (IPTG). Worms were transferred onto RNAi plates
as L1 larvae and cultured at 20 °C or 25 °C until L4 or young adult stage. In all our experiments, we are using
sta-1 as our control, as we have shown over the last decade that it does not affect the development nor any
stress or innate response in the epidermis (Dierking et al., 2011; Lee et al., 2018; Taffoni et al., 2020; Zhang
et al., 2021; Zugasti et al., 2014; Zugasti et al., 2016).

Fluorescent Image acquisition

Live young adult worms were placed on a 2 % agarose pad containing 0.25 mM levamisole in NaCl to
immobilize the worms. Images were acquired using a confocal laser scanning microscopy: Zeiss LSM780 and
its acquisition software Zen with a Plan-Apochromat 40 X/1.4 Oil DIC M27 objective with a zoom 2 to 4, a
Plan-Apochromat 63 X/1.40 Oil DIC M27 with a zoom 1. Spectral imaging combined with linear unmixing
was used in most confocal images to separate the autofluorescence of the cuticle.

Airyscan super-resolution microscopy

Airyscan imaging were performed using a commercial Zeiss confocal microscope LSM 880 equipped with an
Airyscan module (Carl Zeiss AG, Jena, Germany) and images were taken with a 63x/1.40NA M27 Plan
Apochromat oil objective. In this mode, emission light was projected onto an array of 32 sensitive GaAsP
detectors, arranged in a compound eye fashion. The Airyscan processing was done with Zen Black 2.3
software by performing filtering, deconvolution and pixel reassignment to improve SNR. The Airyscan
filtering (Wiener filter associated with deconvolution) was set to the default filter setting of 6.1 in 2D.

Fluorescent Image analysis

To extract the morphological properties of meisosomes, we devised an automatic Fiji segmentation procedure
(GitHub https://github.com/centuri-engineering/BD_BlobsSeg). We first restricted the analysis to manually
drawn ROIs and isolated organelles (foreground image) from the background by using the "remove outliers"
function of Fiji (radius = 30 pixels and threshold = 30). We next applied a Gaussian blur (sigma = 1 pixel) on

the foreground image and automatically defined a threshold value to binarize the newly blurred image. This

threshold was determined automatically by multiplying the background value (retrieved by averaging the

fluorescent levels of the background image) by a constant coefficient. This allowed us to normalize the

segmentation, since the expression levels of fluorescent proteins may vary from one animal to another.

Finally, after filtering out smaller objects (less than ~ 0.15 µm²), we measured the averaged organelles area,

Feret's diameter (longest axis) and density in the different conditions. Unpaired t test was used to compare the

samples which passed the normality test (Shapiro-Wilk normality test) and with homogeneity variances

(Fisher test) and unpaired nonparametric Mann-Whitney test for the others. For co-localisation analysis, we

counted the percentage of segmented objects in a given channel, green (G) or red (R), whose centroid is

located in an object of the other channel. We then averaged these percentages across images, each

representing a different worm (n=10 for each strain analysed).

Atomic Force Microscopy (AFM)

Worms were prepared as described before (Essmann et al., 2016). Briefly, staged 1-day-old young adult

worms were paralysed in 15 mg/ml 2, 3-butanedione monoxime (Sigma) for 2 h at room temperature and

transferred to a ~2 mm thick 4 % agarose bed in a petri dish (30 mm). Heads and tails were fixed with tissue

glue (Dermabond, Ethicon) and the dish filled with a 2.5 ml M9 buffer. AFM data of worms were obtained

using a NanoWizard3 (JPK) under aqueous conditions. Type qp-CONT-10 (0.1 N/m; nanosensors) cantilevers

were used for imaging in contact-mode at setpoint 0.3 V and 0.5 Hz scanning speed, and NSC12 tipless

cantilevers (7.5 N/m; MikroMash) with a 10 µm borosilicate bead attached (produced by sQUBE

www.sQUBE.de) were used in force spectroscopy mode to obtain force-indentation measures at 450 nN force

setpoint and 0.5 µm/s indentation speed. Cantilever sensitivity and stiffness (k) were calibrated using the JPK

system calibration tool before each experiment. AFM raw data were analysed using the JPK analysis

software. All force curves were processed to zero the baseline, to determine the tip-sample contact point and

to subtract cantilever bending. The Young's Modulus was calculated within the software by fitting the

Hertz/Sneddon model respecting the indenter shape (10 µm bead) to each curve. All topographical images are

flattened using the plane fitting option of the JPK software at 1-2 degree to correct for sample tilt and natural

curvature of the worm.

Acknowledgements

We thank Jonathan Ewbank for support and input throughout the project, Ken C Q Nguyen for some of the

TEM imbedding, Michel Labouesse, Barth Grant, Michalis Barkoulas and Ari Pani for sharing strains and

reagents, Chris Crocker at Wormatlas for diagrams, Roxane Fabre for the Airyscan images, Meera Sundaram,

Thomas Sontag and Michel Labouesse for critical reading of the MS and Barth Grant and Erik Jorgensen for

discussions. Worm sorting was performed by Jerome Belougne using the facilities of the French National

Functional Genomics platform, supported by the GIS IBiSA and Labex INFORM. Electron tomography in
Figure 2S1 were performed at the New York Structural Biology Center, with help from KD Derr and William
Rice. We thank John White and Jonathan Hodgkin for sharing the MRC/LMB archive of nematode
micrographs. Some *C. elegans* strains were provided by the CGC, which is funded by NIH Office of Research
Infrastructure Programs (P40 OD010440). We acknowledge the PICsL-FBI photonic microscopy facility of
the CIML (ImagImm) and the PICsL-FBI electron microscopy facility of the IBDM, members of the national
infrastructure France-BioImaging supported by the French National Research Agency (ANR-10-INBS-04).
The project leading to this publication has received funding from France 2030, the French Government
program managed by the French National Research Agency (ANR-16-CONV-0001) and from Excellence
Initiative of Aix-Marseille University - A*MIDEX.

542 Funding

543 Work is funded by the French National Research Agency ANR-22-CE13-0037-01, ANR-16-CE15-0001-01
544 and ANR-10-INBS-04-01 (France Bio Imaging), by the "Investissements d'Avenir" French Government
545 program (ANR-16-CONV-0001) and from Excellence Initiative of Aix-Marseille University - A*MIDEX and
546 institutional grants from CNRS, Aix Marseille University, National Institute of Health and Medical Research
547 (Inserm) to the CIML; and by a NIH OD 010943 to DHH.

549 Figure legends

550 Figure 1: Meisosomes are membrane folded structures on the apical side of the lateral epidermis.

(A) Schematic view of the outer tissues of an adult *C. elegans* worm near the mid-body. The different regions
of the epidermis containing meisosomes are boxed in blue (lateral, and dorsal and ventral ridges), the regions
above the muscles that are extremely thin are boxed in red. (B) Schematic view of the connection between the
cuticle and the plasma membrane (pm) of the lateral epidermis highlighting the position of meisosomes
relative to multivesicular bodies (mvb) and mitochondria. (C-G) TEM images of longitudinal sections of
young adult wild-type hermaphrodite worms reveal epidermal structures, meisosomes (m), contacting the
cuticle (cut), composed of parallel plasma membrane (pm) folds. (C-F) Meisosomes typically comprise less
than 10 folds, but can have more than 30, as shown in (C) where the meisosome is 4 µm wide, and extends on
both sides of a furrow (f). Some folds are apposed to mitochondria (mit) (D), can vary in size and orientation
(C-F) and can appear not to be in direct contact with the cuticle in some EM sections (F); scale bar, 200 nm.
(G) High magnification view of plasma membrane (pm) folds. The "internal" folds (i) are 35 nm wide,
contain a ribosome-free cytoplasm but ladder-like banding (indicated by the black arrowheads), and are less

electron dense in their middle (*). The "outer" folds (o) on the cuticle side are 20 nm wide. Epidermis (epi);
scale bar, 50 nm.

Figure 1-figure supplement 1: Meisosomes are present in epidermis at all development stages.

(A) Schematic view of the tissues of an adult *C. elegans* worm. MRC archival transversal TEM images
showing that meisosomes (red arrow) are present in different body regions in the epidermis (hyp7, B) and in
the tail (hyp9/10, C-D), as well as in different larval stages L1 (E), L2 (F), L3 (G), L4 (H), adult 5 days (I),
adult 15 days (J), dauer stage (K).
570

Figure 1-figure supplement 2: Meisosomes are randomly orientated except before moulting.

(A) Schematic view of the outer tissues of an adult *C. elegans* worm near the mid-body. The plane of
longitudinal EM sections and the *en face* view obtained by freeze fracture are shown in blue and red dotted
lines, respectively. (B) Freeze fracture view of adult epidermal plasma membrane (anterior to the top right)
shows meisosome with random orientation vs the body axis, highlighted by red arrows. MRC archival image.
(C-D) Lengthwise TEM view of the epidermis at pre-moult stages reveals an alignment of meisosomes in
between the position of the furrows (f) facing the annuli (a) of the newly synthesised cuticle (nc), while the
old cuticle (oc) lies to the outside, on its way to be shed. (C) Late L4 stage, scale bar, 500 nm. (D) Late L1
stage, MRC archival image, scale bar, 1 µm.

Figure 1-figure supplement 3: Smaller meisosomes can be found over the muscle quadrants.

(A) Schematic view of the outer tissues of an adult *C. elegans* worm near the mid-body. The different regions
of the epidermis analysed are boxed: either the lateral side as in Figure 1 (blue), or the ventral (or dorsal) side
overlaying the muscle quadrants (red). (B) TEM image of longitudinal section of young adult wild-type worm
reveals small meisosomes in the epidermis above the body-wall muscle; the thickness of the epidermal cell is
less than 500 nm at these positions. Scale bar, 500 nm. (C) A tomogram of a transversal section reveals
several smaller meisosomes (green) with typically not more than 4 folds; other small organelles lie in this
region, including multivesicular bodies (pink), ribosomes (red), and a possible small mitochondrion (dark
blue). (D) One selected TEM image of the same tomogram, red arrows point to small meisosomes. Scale bars
in C&D, 200 nm.
590

Figure 2: Meisosomes are formed by epidermal plasma membrane facing the apical ECM.

Adjacent meisosomes in two serial thick (300 nm) sections were analysed with by electron tomography. (A)
Selected virtual image from the serial reconstruction. (B-E) Segmentation of membranes and mitochondria
reveal their 3D topology. Meisosomes (in yellow) are in close apposition to, but not in continuity with,

mitochondria (orange) and are formed by epidermal plasma membrane folds, as observed in an *en face* view
after removing the cuticle *in silico* (E). (F) Two folds were extracted and manually filled for a schematic
view. Cuticle (cut), epidermis (epi), furrow (f), mitochondria (mit), meisosomes (m); scale bar, 200 nm.

Figure 2-figure supplement 1: Meisosome distribution.

(A) Schematic view of the outer tissues of an adult *C. elegans* worm near the mid-body. The plane of
transverse EM sections and the orthogonal *en face* view for EM and fluorescence microscopy are indicated
with the blue and red dashed rectangles, respectively. (B-C) Representative image from an SBF acquisition,
the boxed area in (B) is shown at a higher magnification in (C). Scale bar, 500 nm. (D) SBF segmentation and
3D reconstruction revealed the *en face* spatial distribution of meisosomes (yellow) compared to mitochondria
(orange). Scale bar, 1 µm.

Figure 3: VHA-5 is a marker of plasma membrane containing meisosomes

(A) Schematic view of the outer tissues of an adult *C. elegans* worm near the mid-body, with the boxed *en*
face view (red) or z projection (blue) for fluorescence microscopy. (B-H) Confocal images of young adult
worms using different strains expressing either a single copy insertion of VHA-5::GFP under an epidermis
promoter [SI], or a classic multi-copy integrated VHA-5::RFP [Is] or extrachromosomal VHA-5::GFP [Ex]
transgenic array. (B) In a young adult worm, VHA-5::GFP fluorescence from a multicopy transgene [Ex] is
observed in the lateral epidermis, as well as in the excretory canal (exc), but largely excluded from the ventral
(v) and dorsal (d) regions above the muscle quadrants, known to contain hemidesmosomes, as well as the
region above the seam cell (sc), as observed in *en face* (left panel), and orthogonal (right) projections of
reconstructed confocal stacks. (C) VHA-5::GFP is also expressed in the ventral ridge, as observed in a ventral
view. (D) High resolution imaging on Airyscan mode revealed VHA-5::GFP from a single copy insertion [Si]
to be associated with long and convoluted apical membranous structures. (D) Meisosomes (m), visualised
using a VHA-5::RFP integrated transgene [Is] (magenta), are complementary with hemidesmosomes (hd)
visualised in a MUP-4::GFP KI (green), autofluorescence, in white, is from intestinal granules. Tiled image
acquired with the spectral mode z projection of 6 section of 1 µm; exc, excretory canal. Attachment of
different structures to the epidermis with hd are highlighted: vulval muscles (vm), ALM and PLM neurones,
anal muscles (a). (F-H) Confocal images of young adult worms expressing VHA-5::GFP [Ex] (green) together
with an HGRS-1::mScarlet marker (magenta) (F), a VHA-5::RFP [Is] (magenta) together with a CAAX::GFP
(G) or PH-PLC1δ::GFP marker (green) (H); see associated Figure 3-figure supplement 1 for the individual
channel as well as the co-localisation quantifications. Scale bar, 10 µm. (I-J) Correlative Light Electron
Microscopy (CLEM) aligns the position of 3 meisosomes revealed by tomography (I) with 3 spots of
VHA-5::GFP signal imaged by confocal imaging (J), see associated Figure 3-figure supplement 2 for the
detailed procedure and another example. Epidermis is pseudo-coloured in beige. Scale bar, 1 µm.

Figure 3-figure supplement 1: VHA-5::FP colocalise with membrane markers but not with vesicular markers in the epidermis.

(A-C) Confocal images of young adult worms expressing VHA-5::GFP [Ex] (green, left panel) together with an mScarlet marker (magenta, middle panel) in the epidermis; right panel overlay; in A, HGRS-1, in B, SNX-1 and in C, LGG-1. (D-E) Confocal images of young adult worms expressing VHA-5::RFP [Is] (magenta, middle panel) together with a GFP marker (green, left panel) in the epidermis; right panel overlay; in D, CAAX, and in E, PH-PLC1δ. Scale bar, 10 μm. (F) Quantification of VHA-5::RFP co-localisation with either the CAAX::GFP or PH-PLC1δ::GFP marker. Segmentation of objects in each channel define green (G) and red (R) objects (left panels), the G objects whose centroid is located in a R object are represented in white and counted as 1, the G objects whose centroid is located outside a R object are represented in green and counted as 0, the R objects whose centroid is located outside a G object are represented in magenta and counted as 0. The average % of all objects is represented, for each reverse situation, with each dot being an analysed ROI in one worm, n=10 for each strain.

Figure 3-figure supplement 2: Correlative Light Electron Microscopy (CLEM).

Two different VHA-5::GFP [Si] worms (A-D and E-H) fixed and sliced for TEM were first analysed by confocal imaging where a bright field view is overlaid with the green GFP signal (A&E). The same preparation was then analysed in TEM at low magnification and both the GFP confocal and the TEM images were aligned using AMIRA (B&F). Then a position with 2 or 3 GFP spots was chosen to do a high magnification tomography TEM acquisition to reveal the meisosomes, 2 meisosomes (indicated by the thin yellow lines) in C correlate with 2 GFP spots in D, 3 meisosomes in G correlate with 3 GFP spots in H. Scale bar, 1 μm.

Figure 4: Meisosome aligned in between furrow before moulting.

(A) Confocal images of worms expressing VHA-5::GFP [Si] from early L4 to young adult (YA) stage. To define the precise L4 stage of all observed worms, the vulva was observed and worms classified according to (Cohen et al., 2020; Mok et al., 2015). (B) Confocal images of worms expressing both DPY-7::GFP and VHA-5::RFP, red channel on the left, merge channel on the right. The inserts show selected regions at a 1.6 X higher magnification. (C) Merged red and green channel confocal images of lateral epidermis of late L4 worms expressing: upper panel, a VHA-5::RFP [Is] with a CAAX::GFP (left panel) or a PH-PLC1δ::GFP (right panel); lower panel, a HGRS-1::mScarlet (left panel), SNX-1::mScarlet (middle panel) or LGG-1::mScarlet (right panel) with VHA-5::GFP [Ex]. Scale bar, 10 μm.

Figure 5: Furrow collagen inactivation provokes a loss of meisosome and cytoskeleton alignment during the L4 stage.

Confocal images of worms expressing VHA-5::GFP [Ex] (upper paired panels), TBB-2::GFP (middle paired
panels), LIFEACT::GFP (lower paired panels) from early L4 to young adult (YA) stage, treated with the
control (*sta-1*) or furrow Dpy (*dpy-7*) RNAi clones, n>4. Scale bar, 5 μm. To define the precise L4 stage, the
vulva was observed and worms classified according to (Cohen et al., 2020). A representative example of the
vulva at each stage is shown on the top row in worms expressing the marker TBB-2::GFP.

Figure 5-figure supplement 1: Furrow collagen inactivation provokes a loss of meisosomes alignment
during the L4 stage.

(A) Confocal images of worms expressing VHA-5::GFP [KI] at the L4.6 (upper panels) and L4.8 stages
(lower panels) treated with a control (*sta-1*; left panels) or a *dpy-7* RNAi (right panels), n>5. (B) A selection
of confocal images of selected regions on the lateral epidermis of late L4 wild-type worms expressing
VHA-5::GFP [Ex] and treated with a control RNAi (*sta-1*; left panels), a *dpy-7* RNAi (middle panels) or in a
dpy-3(e27) mutant background (right panels). Scale bar, 5 μm.

Figure 6: Furrow collagen inactivation provokes meisosome fragmentation.

(A) Schematic view of the outer tissues of an adult *C. elegans* worm near the mid-body, highlighting the
furrows that separate the annuli, the different collagen genes in the annuli (green) or furrow (blue) and
indicating the region analysed with the red dashed rectangle. (B, C) L1 larvae expressing VHA-5::GFP [Si]
were treated with the indicated RNAi clones and analysed at the young adult stage: control (*sta-1*), annuli
collagens *dpy-4*, *dpy-5* and *dpy-13* (green) and the furrow collagens: *dpy-2, dpy-3*, *dpy-7, dpy-8*, *dpy-9* or
dpy-10 (blue). (B) Confocal images of selected regions on the lateral epidermis, at a constant position relative
to the vulva, scale bar, 5 μm. (C) Quantification of VHA-5 signal average size, Feret's diameter and density.
All statistics are in Figure 6-Source data file 1, ****p<0.0001 and ***p<0.001. Control: *sta-1* (n= 17), *dpy-4*
(n= 11), *dpy-5* (n= 10), *dpy-13* (n= 11); furrow Dpy: *dpy-2* (n= 11) *dpy-3* (n= 10), *dpy-7* (n= 14), *dpy-8*
(n=13), *dpy-9* (n=13) or *dpy-10* (n= 11).

Figure 6-figure supplement 1: Worms increase AMP gene expression at the late L4 stage and in furrow
collagen mutants.

IG274 worms carrying the *frIs7* reporter transgene constitutively express the *col-12*p::DsRed reporter and
express *nlp-29*p::GFP only under certain conditions, red and green fluorescence were visualized
simultaneously in all images. (A) Compared to young adults, in late L4 animals entering the moult, the
expression of the *nlp-29*p::GFP reporter is markedly increased. Scale bar, 200 μm (left). Quantification of
relative green fluorescence, L4 stage (n=260), young adult stage (n=149) (right). (B-C) RNA inactivation of
selected collagen genes leads to the induction of *nlp-29*p::GFP in young adult worms. Compared to wild-type
or annuli Dpy mutants, in furrow Dpy mutants, the expression of the *nlp-29*p::GFP reporter is markedly
increased (left panels). Quantification of relative green fluorescence in worms carrying *frIs7* (middle panel)

and the length of the worm (TOF, right panel). Note that these 2 phenotypes (induction of the *nlp-29*p::GFP
reporter and short size) serve as controls for the correct inactivation of the selected genes, and were always
run in parallel to RNAi inactivation on other reporter strains, as the one used in all Figure 5,6 & 8. (B)
Control: *sta-1* (n= 539), *dpy-13* (n= 492); furrow Dpy: *dpy-3* (n= 680), *dpy-7* (n= 688). (C) Control: *sta-1* (n=
195), *dpy-4* (n= 68), *dpy-5* (n=81); furrow Dpy: *dpy-3* (n= 72) or *dpy-7* (n= 95).

Figure 6-figure supplement 2: Inactivation of furrow Dpy do not change the VHA-5 relationship to
endosomes and MVB, nor affect early endosomes.

(A) Confocal images of selected regions on the lateral epidermis of young adult epidermis expressing
LGG-1::mScarlet (left panel), SNX-1::mScarlet (middle panel) or HGRS-1::mScarlet (right panel) with
VHA-5::GFP [Ex] and treated with the indicated RNAi clones. Control: *sta-1*, *dpy-4* and *dpy-5;* furrow Dpy:
dpy-3 or *dpy-7*, n>7. Scale bar, 5 µm. (B) Confocal images and segmentation views (left panels), with the
corresponding quantification (right panels) of selected regions on the lateral epidermis of young adults
expressing EEA-1::GFP and treated with the indicated RNAi clones. Scale bar, 5 µm. Control: *sta-1* (n= 9),
dpy-13 (n=10)*;* furrow Dpy: *dpy-3* (n=10) or *dpy-7* (n=9).

Figure 7: Furrow collagen inactivation leads to smaller and abnormal meisosomes and detachment of
the cuticle.

TEM images of young adult worms. Compared to wild-type (Figure 1) or *dpy-13* (A), *dpy-2* (B) *dpy-7* (C-D)
have abnormally small meisosomes with irregular spacing between the membrane folds (D); number of
worms analysed are given in Table 1. Compared to wild-type (E) or *dpy-13* (F), lower magnification reveals
detachment of the cuticle (cut) from the plasma membrane (pm) in *dpy-3* mutant worms over the whole lateral
surface of the epidermal cell, on both sides of the alae (al) (G). (H) Compared to wild-type (left), SBF
analysis of the entire transversal worm also reveals the detachment (red arrow) in a *dpy-2* mutant of the
cuticle from the lateral epidermis, contrary to the regions above the muscles, delineated in beige and green,
respectively (one representative slice per animal, entire transversal sections were acquired over a length of
21.5 and 34.4 µm, for a wild-type and a *dpy-2* mutant worm, respectively.). Scale bar 500 nm in A&B, 250
nm in insert in B, 200 nm in C&D, 5 µm in E-G and 10 µm in H.

Figure 7-figure supplement 1: Furrow collagen inactivation leads to detachment of the cuticle in lateral
and ventral/dorsal ridges.

TEM images of sagittal sections of young adult worms. Compared to *dpy-13* (A), *dpy-7* (B-C) and *dpy-8* (D)
present detachment of the cuticle (cut) from the plasma membrane, and have abnormally small meisosomes
(C, and inset), in the lateral epidermis (C, D) or the dorsal cord in between muscles (mus) (B). Scale bar, 500
nm in all including C inset, except in C, 1 µm.

Figure 8: Furrow collagen inactivation provokes extrusion of membrane and cytoplasmic contents into the cuticle.

(A) TEM images of *dpy-2* (left) and *dpy-7* (right) young adult mutant worms revealing the presence of membranous organelles (*) and cytoplasmic content, including ribosome-like particles (#), between the cuticle and the plasma membrane, epidermis is pseudo-coloured in beige. Scale bar, 500 nm. (B) Confocal images of wild-type (left) and *dpy-3* mutant (right) young adult worms expressing ROL-6::mScarlet and VHA-5::GFP, CAAX::GFP or PH-PLC1δ::GFP, one confocal plane was selected at the level of the cuticle using the ROL-6::mScarlet. Scale bar, 10 µm. (C) Quantification of the percentage of worm presenting the abnormal presence of GFP extrusion at the level of the cuticle for the different markers; number of worms observed is noted in parenthesis.

Figure 9: Furrow collagen inactivation provokes a reduction in stiffness of the cuticle.

(A) AFM topography of the cuticle in wild-type, *dpy-13*, *dpy-2*, *dpy-3*, *dpy-7* and *dpy-8* mutant adult worms. Scale bar, 1 µm. (B) Mean force-indentation curves of wild-type and collagen mutants acquired by AFM. (C) Young's Modulus estimation from force curves by applying the Hertz model for contact mechanics. Data are from 3 independent experiments with a total of n= 30, 25, 16, 40, 34 and 32, for wild-type, *dpy-13(e184), dpy-2(e8), dpy-3(e27), dpy-7(e88)* and *dpy-8(e130)* mutant worms, respectively.

genotype	cut	n (worms)	n (meisosome)	mean length (µm)	smallest length (µm)	longest length (µm)
wild-type	Transv.	5	22	0,71	0,2	2,6
wild-type	Longit.	4	41	0,77	0,1	4
dpy-13(e184)	Transv.	3	31	0,77	0,3	4
dpy-13(e184)	Longit.	1	11	1,07	0,2	4
dpy-2(e8)	Transv.	1	45	0,39	0,05	1
dpy-2(e8)	Longit.	4	6	0,28	0,1	0,6
dpy-3(e8)	Longit.	1	16	0,49	0,1	1
dpy-7(e88)	Transv.	2	24	0,28	0,05	0,7
dpy-7(e88)	Longit.	4	37	0,37	0,1	1,5

Table 1: Quantification of the length of the meisosomes on TEM images in young adult wild type and
different collagen mutants. The orientation of the section, transversal or longitudinal, the number of
different worms observed and the number of meisosomes analysed are reported.

Supplementary File 1: *C. elegans* **strains used in this study.**

Figure 2 - Video 1: Visualisation of the electron tomography and the 3D segmentation of meisosomes.
Two serial tomograms were stitched and segmentations of the plasma membrane, mitochondria and cuticle
were made (see material and methods for details). The continuity of the plasma membrane could be revealed
by semi-automated segmentation, except under certain orientation due the high membrane curvature of the
meisosomes and the missing wedge artefact, which is inherent to electron tomography. At the end of the
video, the segmentation of two folds was completed manually to present a schematic view.

Figure 6 - Source data file 1: Quantification of the fragmentation of the meisosomes. Using different
VHA-5 reporter strains in wild type and different collagen RNA inactivation or mutants, VHA-5 positive
objects were segmented and three parameters were quantified, average size, Feret's diameter and density; the
number of worms, total surface analysed per condition are presented, together with statistical analysis (see
material and methods for details).

References

Appadurai, D., Gay, L., Moharir, A., Lang, M. J., Duncan, M. C., Schmidt, O., Teis, D., Vu, T.
N., Silva, M., Jorgensen, E. M., & Babst, M. (2020). Plasma membrane tension
regulates eisosome structure and function. *Molecular biology of the cell*, *31*(4), 287-
303. https://doi.org/10.1091/mbc.E19-04-0218
Babst, M. (2011). MVB vesicle formation: ESCRT-dependent, ESCRT-independent and
everything in between. *Current opinion in cell biology*, *23*(4), 452-457.
https://doi.org/10.1016/j.ceb.2011.04.008
Belougne, J., Ozerov, I., Caillard, C., Bedu, F., & Ewbank, J. J. (2020). Fabrication of sharp
silicon arrays to wound *Caenorhabditis elegans*. *Scientific reports*, *10*(1), 3581.
https://doi.org/10.1038/s41598-020-60333-7
Bercher, M., Wahl, J., Vogel, B. E., Lu, C., Hedgecock, E. M., Hall, D. H., & Plenefisch, J. D.
(2001). mua-3, a gene required for mechanical tissue integrity in *Caenorhabditis*
elegans, encodes a novel transmembrane protein of epithelial attachment complexes.
The Journal of cell biology, *154*(2), 415-426. https://doi.org/10.1083/jcb.200103035
Castiglioni, V. G., Pires, H. R., Rosas Bertolini, R., Riga, A., Kerver, J., & Boxem, M. (2020).
Epidermal PAR-6 and PKC-3 are essential for larval development of *C. elegans* and
organize non-centrosomal microtubules. *eLife*, *9*. https://doi.org/10.7554/eLife.62067
Chebli, Y., Bidhendi, A. J., Kapoor, K., & Geitmann, A. (2021). Cytoskeletal regulation of
primary plant cell wall assembly. *Curr Biol*, *31*(10), R681-R695.
https://doi.org/10.1016/j.cub.2021.03.092
Cohen, J. D., Sparacio, A. P., Belfi, A. C., Forman-Rubinsky, R., Hall, D. H., Maul-Newby, H.,
Frand, A. R., & Sundaram, M. V. (2020). A multi-layered and dynamic apical
extracellular matrix shapes the vulva lumen in *Caenorhabditis elegans*. *eLife*, *9*.
https://doi.org/10.7554/eLife.57874
Costa, M., Draper, B. W., & Priess, J. R. (1997). The role of actin filaments in patterning the
Caenorhabditis elegans cuticle. *Developmental biology*, *184*(2), 373-384.
https://doi.org/10.1006/dbio.1997.8530
Cox, G. N., & Hirsh, D. (1985). Stage-specific patterns of collagen gene expression during
development of *Caenorhabditis elegans*. *Molecular and cellular biology*, *5*(2), 363-
372. https://doi.org/10.1128/mcb.5.2.363
Cox, G. N., Laufer, J. S., Kusch, M., & Edgar, R. S. (1980). Genetic and Phenotypic
Characterization of Roller Mutants of *Caenorhabditis elegans*. *Genetics*, *95*(2), 317-
339. https://doi.org/10.1093/genetics/95.2.317
Cox, G. N., Staprans, S., & Edgar, R. S. (1981). The cuticle of *Caenorhabditis elegans.* II.
Stage-specific changes in ultrastructure and protein composition during
postembryonic development. *Developmental biology*, *86*(2), 456-470.
https://doi.org/10.1016/0012-1606(81)90204-9
Davies, K. G., & Curtis, R. H. (2011). Cuticle surface coat of plant-parasitic nematodes. *Annu*
Rev Phytopathol, *49*, 135-156. https://doi.org/10.1146/annurev-phyto-121310-111406
Deerinck, T. J., Bushong, E., Thor, A., & Ellisman, M. (2010). NCMIR methods for 3D EM: A
new protocol for preparation of biological specimens for serial block face scanning
electron microscopy. *Nat Center Microsc Imag Res*, 6-8. https://ncmir.ucsd.edu/sbem-
protocol
Dickinson, D. J., Pani, A. M., Heppert, J. K., Higgins, C. D., & Goldstein, B. (2015).
Streamlined Genome Engineering with a Self-Excising Drug Selection Cassette.
Genetics, *200*(4), 1035-1049. https://doi.org/10.1534/genetics.115.178335
Dierking, K., Polanowska, J., Omi, S., Engelmann, I., Gut, M., Lembo, F., Ewbank, J. J., &
Pujol, N. (2011). Unusual regulation of a STAT protein by an SLC6 family transporter
in *C. elegans* epidermal innate immunity. *Cell Host Microbe*, *9*(5), 425-435.
https://doi.org/10.1016/j.chom.2011.04.011.
Dodd, W., Tang, L., Lone, J. C., Wimberly, K., Wu, C. W., Consalvo, C., Wright, J. E., Pujol,
N., & Choe, K. P. (2018). A Damage Sensor Associated with the Cuticle Coordinates

Three Core Environmental Stress Responses in *C. elegans*. *Genetics*, *208*(4), 1467-1482. https://doi.org/10.1534/genetics.118.300827

E, L., Zhou, T., Koh, S., Chuang, M., Sharma, R., Pujol, N., Chisholm, A. D., Eroglu, C., Matsunami, H., & Yan, D. (2018). An Antimicrobial Peptide and Its Neuronal Receptor Regulate Dendrite Degeneration in Aging and Infection. *Neuron*, *97*(1), 125-138. https://doi.org/10.1016/j.neuron.2017.12.001

Essmann, C. L., Elmi, M., Shaw, M., Anand, G. M., Pawar, V. M., & Srinivasan, M. A. (2016). In-vivo high resolution AFM topographic imaging of *Caenorhabditis elegans* reveals previously unreported surface structures of cuticle mutants. *Nanomedicine*. https://doi.org/10.1016/j.nano.2016.09.006

Essmann, C. L., Martinez-Martinez, D., Pryor, R., Leung, K. Y., Krishnan, K. B., Lui, P. P., Greene, N. D. E., Brown, A. E. X., Pawar, V. M., Srinivasan, M. A., & Cabreiro, F. (2020). Mechanical properties measured by atomic force microscopy define health biomarkers in ageing *C. elegans*. *Nature communications*, *11*(1), 1043. https://doi.org/10.1038/s41467-020-14785-0

Forman-Rubinsky, R., Cohen, J. D., & Sundaram, M. V. (2017). Lipocalins Are Required for Apical Extracellular Matrix Organization and Remodeling in *Caenorhabditis elegans*. *Genetics*, *207*(2), 625-642. https://doi.org/10.1534/genetics.117.300207

Frische, E. W., Pellis-van Berkel, W., van Haaften, G., Cuppen, E., Plasterk, R. H., Tijsterman, M., Bos, J. L., & Zwartkruis, F. J. (2007). RAP-1 and the RAL-1/exocyst pathway coordinate hypodermal cell organization in *Caenorhabditis elegans*. *The EMBO journal*, *26*(24), 5083-5092. https://doi.org/10.1038/sj.emboj.7601922

Frokjaer-Jensen, C., Davis, M. W., Hopkins, C. E., Newman, B. J., Thummel, J. M., Olesen, S. P., Grunnet, M., & Jorgensen, E. M. (2008). Single-copy insertion of transgenes in *Caenorhabditis elegans* [Research Support, Non-U.S. Gov't]. *Nature genetics*, *40*(11), 1375-1383. https://doi.org/10.1038/ng.248

Hall, D. H., Hartweig, E., & Nguyen, K. C. (2012). OTO Fixation for SEM and Blockface Imaging. https://www.wormatlas.org/EMmethods/OTOFix.htm

Hall, D. H., Hartwieg, E., & Nguyen, K. C. (2012). Modern electron microscopy methods for *C. elegans*. *Methods Cell Biol*, *107*, 93-149. https://doi.org/10.1016/B978-0-12-394620-1.00004-7

Hyenne, V., Apaydin, A., Rodriguez, D., Spiegelhalter, C., Hoff-Yoessle, S., Diem, M., Tak, S., Lefebvre, O., Schwab, Y., Goetz, J. G., & Labouesse, M. (2015). RAL-1 controls multivesicular body biogenesis and exosome secretion. *The Journal of cell biology*, *211*(1), 27-37. https://doi.org/10.1083/jcb.201504136

Johnson, E., Seiradake, E., Jones, E. Y., Davis, I., Grunewald, K., & Kaufmann, R. (2015). Correlative in-resin super-resolution and electron microscopy using standard fluorescent proteins. *Scientific reports*, *5*, 9583. https://doi.org/10.1038/srep09583

Johnstone, I. L. (2000). Cuticle collagen genes. Expression in *Caenorhabditis elegans*. *Trends in genetics : TIG*, *16*(1), 21-27. https://doi.org/10.1016/s0168-9525(99)01857-0

Kamath, R. S., Fraser, A. G., Dong, Y., Poulin, G., Durbin, R., Gotta, M., Kanapin, A., Le Bot, N., Moreno, S., Sohrmann, M., Welchman, D. P., Zipperlen, P., & Ahringer, J. (2003). Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature*, *421*(6920), 231-237. https://doi.org/10.1038/nature01278

Katan, M., & Cockcroft, S. (2020). Phosphatidylinositol(4,5)bisphosphate: diverse functions at the plasma membrane. *Essays Biochem*, *64*(3), 513-531. https://doi.org/10.1042/EBC20200041

Katz, S. S., Barker, T. J., Maul-Newby, H. M., Sparacio, A. P., Nguyen, K. C. Q., Maybrun, C. L., Belfi, A., Cohen, J. D., Hall, D. H., Sundaram, M. V., & Frand, A. R. (2022). A transient apical extracellular matrix relays cytoskeletal patterns to shape permanent acellular ridges on the surface of adult *C. elegans*. *PLoS genetics*, *18*(8), e1010348. https://doi.org/10.1371/journal.pgen.1010348

Katz, S. S., Maybrun, C., Maul-Newby, H. M., & Frand, A. R. (2018). Non-canonical apical constriction shapes emergent matrices in *C. elegans*. *bioRxiv*, 189951. https://doi.org/10.1101/189951

Kolotuev, I., Schwab, Y., & Labouesse, M. (2009). A precise and rapid mapping protocol for correlative light and electron microscopy of small invertebrate organisms. *Biol Cell*, *102*(2), 121-132. https://doi.org/10.1042/BC20090096

Lanze, C. E., Gandra, R. M., Foderaro, J. E., Swenson, K. A., Douglas, L. M., & Konopka, J. B. (2020). Plasma Membrane MCC/Eisosome Domains Promote Stress Resistance in Fungi. *Microbiol Mol Biol Rev, 84*(4). https://doi.org/10.1128/MMBR.00063-19

Lardennois, A., Pasti, G., Ferraro, T., Llense, F., Mahou, P., Pontabry, J., Rodriguez, D., Kim, S., Ono, S., Beaurepaire, E., Gally, C., & Labouesse, M. (2019). An actin-based viscoplastic lock ensures progressive body-axis elongation. *Nature*, *573*(7773), 266-270. https://doi.org/10.1038/s41586-019-1509-4

Lee, S. H., Omi, S., Thakur, N., Taffoni, C., Belougne, J., Engelmann, I., Ewbank, J. J., & Pujol, N. (2018). Modulatory upregulation of an insulin peptide gene by different pathogens in *C. elegans*. *Virulence*, *9*(1), 648-658. https://doi.org/10.1080/21505594.2018.1433969

Lemiere, J., Ren, Y., & Berro, J. (2021). Rapid adaptation of endocytosis, exocytosis and eisosomes after an acute increase in membrane tension in yeast cells. *eLife*, *10*. https://doi.org/10.7554/eLife.62084

Lemmon, M. A., Ferguson, K. M., O'Brien, R., Sigler, P. B., & Schlessinger, J. (1995). Specific and high-affinity binding of inositol phosphates to an isolated pleckstrin homology domain. *Proc Natl Acad Sci U S A*, *92*(23), 10472-10476. https://doi.org/10.1073/pnas.92.23.10472

Liegeois, S., Benedetto, A., Garnier, J. M., Schwab, Y., & Labouesse, M. (2006). The V0-ATPase mediates apical secretion of exosomes containing Hedgehog-related proteins in *C. elegans*. *The Journal of cell biology*, *173*(6), 949-961. https://doi.org/10.1083/jcb.200511072

Liegeois, S., Benedetto, A., Michaux, G., Belliard, G., & Labouesse, M. (2007). Genes required for osmoregulation and apical secretion in *C. elegans*. *Genetics*, *175*(2), 709-724. https://doi.org/10.1534/genetics.106.066035

Martineau, C. N., Kirienko, N. V., & Pujol, N. (2021). Innate immunity in *C. elegans*. *Current topics in developmental biology*, *144*, 309-351. https://doi.org/10.1016/bs.ctdb.2020.1

McMahon, L., Muriel, J. M., Roberts, B., Quinn, M., & Johnstone, I. L. (2003). Two sets of interacting collagens form functionally distinct substructures within a *Caenorhabditis elegans* extracellular matrix. *Molecular biology of the cell*, *14*(4), 1366-1378. https://doi.org/10.1091/mbc.E02-08-0479

Miao, R., Li, M., Zhang, Q., Yang, C., & Wang, X. (2020). An ECM-to-Nucleus Signaling Pathway Activates Lysosomes for *C. elegans* Larval Development. *Dev Cell*, *52*(1), 21-37 e25. https://doi.org/10.1016/j.devcel.2019.10.020

Mok, D. Z., Sternberg, P. W., & Inoue, T. (2015). Morphologically defined sub-stages of C. elegans vulval development in the fourth larval stage. *BMC Dev Biol*, *15*, 26. https://doi.org/10.1186/s12861-015-0076-7

Moseley, J. B. (2018). Eisosomes. *Curr Biol*, *28*(8), R376-R378. https://doi.org/10.1016/j.cub.2017.11.073

Norris, A., Tammineni, P., Wang, S., Gerdes, J., Murr, A., Kwan, K. Y., Cai, Q., & Grant, B. D. (2017). SNX-1 and RME-8 oppose the assembly of HGRS-1/ESCRT-0 degradative microdomains on endosomes. *Proc Natl Acad Sci U S A*, *114*(3), E307-E316. https://doi.org/10.1073/pnas.1612730114

Oka, T., Toyomura, T., Honjo, K., Wada, Y., & Futai, M. (2001). Four subunit a isoforms of Caenorhabditis elegans vacuolar H+-ATPase. Cell-specific expression during development. *The Journal of biological chemistry*, *276*(35), 33079-33085. https://doi.org/10.1074/jbc.M101652200

Page, A. P., & Johnstone, I. L. (2007). The cuticle. *WormBook : the online review of C elegans biology*, 1-15. https://doi.org/10.1895/wormbook.1.138.1

Polanowska, J., Chen, J. X., Soule, J., Omi, S., Belougne, J., Taffoni, C., Pujol, N., Selbach, M., Zugasti, O., & Ewbank, J. J. (2018). Evolutionary plasticity in the innate immune function of Akirin. *PLoS genetics*, *14*(7), e1007494. https://doi.org/10.1371/journal.pgen.1007494

Pujol, N., Bonnerot, C., Ewbank, J. J., Kohara, Y., & Thierry-Mieg, D. (2001). The *Caenorhabditis elegans unc-32* gene encodes alternative forms of a vacuolar ATPase a subunit. *The Journal of biological chemistry*, *276*(15), 11913-11921. https://doi.org/10.1074/jbc.M009451200

Pujol, N., Cypowyj, S., Ziegler, K., Millet, A., Astrain, A., Goncharov, A., Jin, Y., Chisholm, A. D., & Ewbank, J. J. (2008). Distinct innate immune responses to infection and wounding in the *C. elegans* epidermis. *Curr Biol*, *18*(7), 481-489. https://doi.org/10.1016/j.cub.2008.02.079.

Pujol, N., Zugasti, O., Wong, D., Couillault, C., Kurz, C. L., Schulenburg, H., & Ewbank, J. J. (2008). Anti-fungal innate immunity in *C. elegans* is enhanced by evolutionary diversification of antimicrobial peptides. *PLoS pathogens*, *4*(7), e1000105. https://doi.org/10.1371/journal.ppat.1000105

Rohlfing, A. K., Miteva, Y., Hannenhalli, S., & Lamitina, T. (2010). Genetic and physiological activation of osmosensitive gene expression mimics transcriptional signatures of pathogen infection in *C. elegans*. *PloS one*, *5*(2), e9010. https://doi.org/10.1371/journal.pone.0009010

Roudier, N., Lefebvre, C., & Legouis, R. (2005). CeVPS-27 is an endosomal protein required for the molting and the endocytic trafficking of the low-density lipoprotein receptor-related protein 1 in *Caenorhabditis elegans*. *Traffic*, *6*(8), 695-705. https://doi.org/10.1111/j.1600-0854.2005.00309.x

Rual, J. F., Ceron, J., Koreth, J., Hao, T., Nicot, A. S., Hirozane-Kishikawa, T., Vandenhaute, J., Orkin, S. H., Hill, D. E., van den Heuvel, S., & Vidal, M. (2004). Toward improving *Caenorhabditis elegans* phenome mapping with an ORFeome-based RNAi library. *Genome research*, *14*(10B), 2162-2168. https://doi.org/10.1101/gr.2505604

Sandhu, A., Badal, D., Sheokand, R., Tyagi, S., & Singh, V. (2021). Specific collagens maintain the cuticle permeability barrier in *Caenorhabditis elegans*. *Genetics*, *217*(3). https://doi.org/10.1093/genetics/iyaa047

Serrano-Saiz, E., Vogt, M. C., Levy, S., Wang, Y., Kaczmarczyk, K. K., Mei, X., Bai, G., Singson, A., Grant, B. D., & Hobert, O. (2020). SLC17A6/7/8 Vesicular Glutamate Transporter Homologs in Nematodes. *Genetics*, *214*(1), 163-178. https://doi.org/10.1534/genetics.119.302855

Shi, A., Sun, L., Banerjee, R., Tobin, M., Zhang, Y., & Grant, B. D. (2009). Regulation of endosomal clathrin and retromer-mediated endosome to Golgi retrograde transport by the J-domain protein RME-8. *The EMBO journal*, *28*(21), 3290-3302. https://doi.org/10.1038/emboj.2009.272

Stiernagle, T. (Ed.). (2006). *Maintenance of C. elegans* (Vol. doi/10.1895/wormbook.1.101.1). The *C. elegans* Research Community ed. http://www.wormbook.org.

Suman, S. K., Daday, C., Ferraro, T., Vuong-Brender, T., Tak, S., Quintin, S., Robin, F., Grater, F., & Labouesse, M. (2019). The plakin domain of *C. elegans* VAB-10/plectin acts as a hub in a mechanotransduction pathway to promote morphogenesis. *Development*, *146*(24). https://doi.org/10.1242/dev.183780

Taffoni, C., Omi, S., Huber, C., Mailfert, S., Fallet, M., Rupprecht, J. F., Ewbank, J. J., & Pujol, N. (2020). Microtubule plus-end dynamics link wound repair to the innate immune response. *eLife*, *9*, e45047, Article e45047. https://doi.org/10.7554/eLife.45047

Thein, M. C., McCormack, G., Winter, A. D., Johnstone, I. L., Shoemaker, C. B., & Page, A. P. (2003). *Caenorhabditis elegans* exoskeleton collagen COL-19: an adult-specific

marker for collagen modification and assembly, and the analysis of organismal
morphology [Research Support, Non-U.S. Gov't]. *Developmental dynamics : an*
official publication of the American Association of Anatomists, *226*(3), 523-539.
https://doi.org/10.1002/dvdy.10259

Vuong-Brender, T. T., Ben Amar, M., Pontabry, J., & Labouesse, M. (2017). The interplay of
stiffness   and   force   anisotropies   drives   embryo   elongation.   *eLife*,   *6*.
https://doi.org/10.7554/eLife.23866

Vuong-Brender, T. T. K., Suman, S. K., & Labouesse, M. (2017). The apical ECM preserves
embryonic integrity and distributes mechanical stress during morphogenesis.
Development, *144*(23), 4336-4349. https://doi.org/10.1242/dev.150383

Watts, J. S., Harrison, H. F., Omi, S., Guenthers, Q., Dalelio, J., Pujol, N., & Watts, J. L.
(2020). New Strains for Tissue-Specific RNAi Studies in *Caenorhabditis elegans*. *G3*
(Bethesda), *10*(11), 4167-4176. https://doi.org/10.1534/g3.120.401749

Wheeler, J. M., & Thomas, J. H. (2006). Identification of a novel gene family involved in
osmotic stress response in *Caenorhabditis elegans*. *Genetics*, *174*(3), 1327-1336.
http://www.ncbi.nlm.nih.gov/entrez/query.fcgi?cmd=Retrieve&db=PubMed&dopt=Citat
ion&list_uids=16980399

White, J. G., Southgate, E., Thomson, J. N., & Brenner, S. (1986). The structure of the
nervous system of the nematode *C. elegans*. *Philosophical Transactions of the Royal*
Society of London, *314B*, 1-340.

Wood, W. B. (Ed.). (1988). *The nematode Caenorhabditis elegans*. Cold Spring Harbor
Laboratory Press.

Zhang, H., & Labouesse, M. (2010). The making of hemidesmosome structures in vivo.
Developmental dynamics : an official publication of the American Association of
Anatomists, *239*(5), 1465-1476. https://doi.org/10.1002/dvdy.22255

Zhang, X., Harding, B. W., Aggad, D., Courtine, D., Chen, J. X., Pujol, N., & Ewbank, J. J.
(2021). Antagonistic fungal enterotoxins intersect at multiple levels with host innate
immune   defences.   *PLoS   genetics*,   *17*(6),   e1009600.
https://doi.org/10.1371/journal.pgen.1009600

Zugasti, O., Bose, N., Squiban, B., Belougne, J., Kurz, C. L., Schroeder, F. C., Pujol, N., &
Ewbank, J. J. (2014). Activation of a G protein-coupled receptor by its endogenous
ligand triggers the innate immune response of *C. elegans. Nature immunology*, *15*(9),
833-838. https://doi.org/10.1038/ni.2957

Zugasti, O., Thakur, N., Belougne, J., Squiban, B., Kurz, C. L., Soule, J., Omi, S., Tichit, L.,
Pujol, N., & Ewbank, J. J. (2016). A quantitative genome-wide RNAi screen in *C.*
elegans for antifungal innate immunity genes. *BMC biology*, *14*(1), 35.
https://doi.org/10.1186/s12915-016-0256-3

1012

FIGURE 1

FIGURE 1-Supplement 1

FIGURE 1-Supplement 2

A

surface view
freeze crack (B)

longitudinal
view (C,D)

B

C L4 pre-moulting

f   a   f   a   f   a

oc

nc

f

D L1 pre-moulting   f

oc

nc   f

FIGURE 1-Supplement 3

A

lateral
(Figure 1)

epidermis

muscle

ventral
(B-D)

B

cut

f

muscle

C

cut

muscle

D

FIGURE 2

FIGURE 2-Supplement 1

FIGURE 3

FIGURE 3-Supplement 1

FIGURE 4

FIGURE 5

FIGURE 5-Supplement 1

A

control (RNAi)   *dpy-7* (RNAi)

VHA-5::GFP (KI)

L4.6   L4.6

L4.8   L4.8

B

control (RNAi)   *dpy-7* (RNAi)   *dpy-3(e27)*

VHA-5::GFP (Ex)

FIGURE 6

FIGURE 6-Supplement 1

FIGURE 6-Supplement 2

FIGURE 7

FIGURE 7-Supplement 1

FIGURE 8

% of worms with GFP at the level of the cuticle

	wt	dpy-3
VHA-5::GFP	0 % (n>50)	0 % (n>50)
GFP::CAAX	0 % (n=20)	100 % (n=21)
GFP::PH-PLC1δ	0 % (n=27)	89,6 % (n=29)

FIGURE 9