Heterocomplexes between the Atypical Chemokine MIF and the CXC-Motif Chemokine CXCL4L1 Regulate Inflammation and Thrombus Formation

Markus Brandhofer[1,#], Adrian Hoffmann[1,2,#], Xavier Blanchet[3], Elena Siminkovitch[1], Anne-Katrin Rohlfing[4], Omar El Bounkari[1], Jeremy A. Nestele[4], Alexander Bild[4], Christos Kontos[5], Kathleen Hille[5], Vanessa Rohde[1], Adrian Fröhlich[1], Jona Golemi[6], Ozgun Gokce[6,7], Patrick Scheiermann[2], Nikolaos Tsilimparis[8], Wolfgang E. Kempf[9], Lars Maegdefessel[9], Remco T.A. Megens[3], Hans Ippel[10], Rory R. Koenen[10], Kevin H. Mayo[10,11], Meinrad Gawaz[4], Aphrodite Kapurniotu[5], Christian Weber[3,7,10,12], Philipp von Hundelshausen[3,*], Jürgen Bernhagen[1,7,12*]

[1]Division of Vascular Biology, Institute for Stroke and Dementia Research (ISD), LMU University Hospital, Ludwig-Maximilians-Universität (LMU) München, 81377 Munich, Germany; [2]Department of Anesthesiology, LMU University Hospital, Ludwig-Maximilians-Universität (LMU) München, 81377 Munich, Germany; [3]Institute for Cardiovascular Prevention, LMU University Hospital, Ludwig-Maximilians-Universität (LMU) München, 80336 Munich, Germany; [4]Department of Cardiology and Angiology, University Hospital Tübingen, Eberhard Karls University Tübingen, Tübingen, Germany; [5]Division of Peptide Biochemistry, TUM School of Life Sciences, Technische Universität München (TUM); [6]Systems Neuroscience Group, Institute for Stroke and Dementia Research (ISD), LMU University Hospital, Ludwig-Maximilians-Universität (LMU) München, 81377 Munich, Germany; [7]Munich Cluster for Systems Neurology (SyNergy), 81377 Munich, Germany; [8]Department of Vascular Surgery, LMU University Hospital, Ludwig-Maximilians-Universität (LMU) München, 81377 Munich, Germany; [9]Department for Vascular and Endovascular Surgery, Klinikum rechts der Isar, Technische Universität München (TUM), 81675 Munich, Germany; [10]Cardiovascular Research Institute Maastricht (CARIM), Maastricht University, 6229 ER Maastricht, The Netherlands; [11] Department of Biochemistry, Molecular Biology and Biophysics, Health Sciences Center, University of Minnesota, Minneapolis, MN, USA; [12]Munich Heart Alliance, 80802 Munich, Germany.

Correspondence:

Jürgen Bernhagen, PhD

Professor and Chair of Vascular Biology

Institute for Stroke and Dementia Research (ISD)

LMU University Hospital (LMU Klinikum)

Ludwig-Maximilians-Universität (LMU) München

Feodor-Lynen-Straße 17, 81377 Munich, Germany

Tel.:   0049-89 4400 - 46151

E-Mail: juergen.bernhagen@med.uni-muenchen.de

Philipp von Hundelshausen, MD

Institute for Cardiovascular Prevention (IPEK)

LMU University Hospital (LMU Klinikum)

Ludwig-Maximilians-Universität (LMU) München

Pettenkofer Straße 8a/9, 80336 Munich, Germany

Tel.:   0049-89 4400 - 54359

E-Mail: Philipp.von_Hundelshausen@med.uni-muenchen.de

Abstract

To fulfil its orchestrating function in immune cell trafficking, a network of chemokines and receptors capitalizes on specificity, redundancy, and functional selectivity. The discovery of heteromeric interactions in the chemokine interactome has expanded the complexity within the network. Moreover, some inflammatory mediators, not structurally linked to classical chemokines, bind to chemokine receptors and behave as atypical chemokines (ACKs). We identified macrophage migration inhibitory factor (MIF) as an ACK that binds to the chemo-kine receptors CXCR2 and CXCR4 to promote atherogenic leukocyte recruitment. Here, we hypothesized that chemokine-chemokine interactions extend to ACKs and that MIF may form heterocomplexes with classical chemokines. We tested this hypothesis by an unbiased chemokine protein array. The platelet chemokine CXCL4L1, but not its variant CXCL4 or the CXCR2/CXCR4 ligands CXCL8 or CXCL12, was identified as a candidate interactor. MIF/CXCL4L1 complexation was verified by co-immunoprecipitation, surface plasmon-resonance analysis, and microscale thermophoresis, also establishing high-affinity binding. We next determined whether heterocomplex formation modulates inflammatory/atherogenic activities of MIF. Complex formation abrogated MIF-elicited T-cell chemotaxis as assessed in a 3D-matrix-based live cell-imaging set-up. Heterocomplexation also blocked MIF-triggered migration of microglia in cortical cultures *in situ*. Of note, CXCL4L1 blocked the binding of Alexa-MIF to a soluble surrogate of CXCR4 and co-incubation with CXCL4L1 attenuated MIF responses in HEK293-CXCR4 transfectants, indicating that complex formation interferes with MIF/CXCR4 pathways. As MIF and CXCL4L1 are platelet products, we finally tested their role in platelet activation. Multi-photon microscopy, FLIM-FRET, and proximity-ligation assay visualized heterocomplexes in platelet aggregates and clinical human thrombus sections. Moreover, heterocomplexes inhibited MIF-stimulated thrombus formation under flow and skewed the lamellipodia phenotype of adhering platelets. Our study establishes a novel molecular interaction, adding to the complexity of the chemokine interactome and chemokine/receptor-network. MIF/CXCL4L1, or more generally, ACK/CXC-motif chemokine heterocomplexes may be target structures to modulate inflammation and thrombosis.

Introduction

Chemokines orchestrate immune cell trafficking in health and disease (Charo & Ransohoff, 2006; Noels *et al*, 2019; Weber & Noels, 2011). Chemokine-directed targeting strategies are pursued in acute and chronic inflammatory conditions, autoimmunity, cancer, and athero-sclerosis (Hutchings *et al*, 2017; Noels *et al.*, 2019; Zlotnik *et al*, 2011). The chemokine network encompasses 49 classical chemokines (CKs) and 18 classical chemokine receptors (CKRs), which belong to the class of Giα protein-coupled receptors (GPCRs) (Bachelerie *et al*, 2014a; Bachelerie *et al*, 2014b; Murphy *et al*, 2000). Depending on the particular chemokine ligand/receptor pair and various disease and microenvironmental factors, chemokine signaling through CKRs overall capitalizes on the principles of specificity, promiscuity, and biased agonism. Accordingly, multiple chemokines can bind to a certain chemokine receptor and *vice versa*, while 'biased agonism' can occur on a ligand, receptor, or tissue basis (Eiger *et al*, 2021; Kleist *et al*, 2016; Steen *et al*, 2014). Fine-tuning of chemokine responses within this network is further expanded by five atypical chemokine receptors (ACKRs) that serve as decoy receptors and promiscuously bind many chemokines to shape their gradients, but also elicit specific signaling responses (Nibbs & Graham, 2013).

Chemokines are well-known to form homodimers, but the discovery of the chemokine interactome additionally suggested a multitude of heteromeric chemokine-chemokine inter-actions even across CC- and CXC-chemokine class borders (Koenen *et al*, 2009; von Hundelshausen *et al*, 2017). CC-type heterodimers between CCL5 and CCL17 or CCL5 and CXCL4 (also termed platelet factor 4, PF4) were found to lead to functional synergism by receptor retention or auxiliary proteoglycan binding and enhancement of chemotactic responses, respectively, while CXC-type heterodimers between CXCL12 and CCL5 or CXCL12 and CXCL4 led to signaling inhibition. This has demonstrated yet another level of complexity within the chemokine network and offers novel intervention strategies in inflammatory and cardiovascular diseases (von Hundelshausen *et al.*, 2017).

Moreover, some alarmin-like inflammatory mediators such as human β-defensins (HBDs) and secreted fragments of amino acyl tRNA-synthetases (AARSs), which do not

belong to one of the four structural classes of CC-, CXC-, CX_3C-, or C-chemokines, can bind

to chemokine receptors by molecular mimicry and exhibit chemokine-like activities (Rohrl *et*

al, 2010; Wakasugi & Schimmel, 1999). These proteins are also referred to as atypical

chemokines (ACKs) (Degryse & de Virgilio, 2003; Kapurniotu *et al*, 2019; Oppenheim &

Yang, 2005).

Macrophage migration inhibitory factor (MIF) is an evolutionarily conserved pleiotropic

inflammatory cytokine (David, 1966; Michelet *et al*, 2019). MIF is an upstream regulator of

the host innate immune response and, when dysregulated, is a pivotal mediator of

inflammatory diseases, autoimmunity, cancer, and cardiovascular diseases (Calandra &

Roger, 2003; Tilstam *et al*, 2017). MIF is a structurally unique cytokine (Sun *et al*, 1996) and,

contrary to its eponymous name, has chemokine-like activities and functions as a

prototypical ACK (Bernhagen *et al*, 2007; Kapurniotu *et al.*, 2019). Accordingly, MIF not only

signals through its cognate receptor CD74/invariant chain, but engages in high-affinity

interactions with the CXC chemokine receptors CXCR2 and CXCR4 to promote atherogenic

monocyte and T-/B-cell recruitment, cancer metastasis, and inflammation (Bernhagen *et al.*,

2007; Kapurniotu *et al.*, 2019; Klasen *et al*, 2014; Leng *et al*, 2003; Pawig *et al*, 2015; Sinitski

et al, 2019; Tillmann *et al*, 2013). We elucidated the structural determinants of the binding

interface between MIF and its CXC-motif chemokine receptors and found that MIF mimics

chemokine receptor binding regions such as the ELR motif and the N-loop (Kraemer *et al*,

2011b; Krammer *et al*, 2021; Lacy *et al*, 2018; Rajasekaran *et al*, 2016; Weber *et al*, 2008).

Interestingly, CXCL12/SDF-1α (stromal-derived factor-1α), the cognate ligand of CXCR4,

was recently found to bind to the non-chemokine proteins galectin-3 (Eckardt *et al*, 2020) and

high-mobility group box-1 (HMGB1) (De Leo *et al*, 2019; Schiraldi *et al*, 2012), but potential

interactions between MIF and CXCL12 or CXCL8, the cognate ligand of its chemokine

receptors CXCR2, have remained unclear.

Here, we hypothesized that chemokine-chemokine interactions are not only possible

between different types of classical chemokines, as demonstrated by chemokine interactome

mapping (von Hundelshausen *et al.*, 2017), but might extend to ACKs. Choosing MIF as a

prototypical ACK, we thus asked whether this mediator would form heterocomplexes with

classical chemokines. We tested this hypothesis applying an unbiased chemokine protein

array and validated candidate interactors by a battery of biochemical and biophysical

methods. We identified the platelet chemokine CXCL4L1 (also termed PF4var1), but not its

variant CXCL4, nor the CXCR2 ligand CXCL8 or the CXCR4 ligand CXCL12, as a high

affinity interactor of MIF and tested the potential functional role of CXCL4L1/MIF

heterocomplex formation for MIF binding to its receptor CXCR4, and in cell systems that are

relevant for the inflammatory, atherogenic, and thrombogenic activities of MIF. Finally, we

also asked whether such heterocomplexes can be detected in clinical thrombus specimens.

Our study extends the chemokine interactome to ACK/CK interactions and demonstrates a

functional role for the MIF/CXCL4L1 heterocomplex in disease-relevant activities.

Materials and Methods

Proteins and reagents

Biologically active and endotoxin-free recombinant human MIF was prepared as previously described and was obtained at a purity of ~98% as confirmed by SDS-PAGE analysis in combination with silver staining (Bernhagen *et al*, 1994; Kontos *et al*, 2020). For the preparation of Alexa Fluor-488- and MST-Red-labeled MIF, a 90-95% pure MIF fraction was used. Alexa Fluor-488-labeled MIF was generated using the Microscale Protein Labeling Kit from Invitrogen-Molecular Probes (Karlsruhe, Germany) and MST-Red-MIF was prepared using the Monolith Protein Labeling Kit RED-NHS 2nd Generation from NanoTemper (Munich, Germany), following the manufacturers' instructions. Biotinylated human MIF was produced using D-biotinoyl-ε-aminocaproic acid-N-hydroxy-succinimide ester (Biotin-7-NHS) with the Biotin Protein Labeling Kit from Roche (Mannheim, Germany). Alternatively, biotin-amidohexanoic acid N-hydroxysuccinimide ester from Sigma-Aldrich (Taufkirchen, Germany) was used.

For the fluorescence polarization assay, a hexahistidine-tagged variant of CXCL4L1 was used. Briefly, the coding sequence of human CXCL4L1 with a methionine-flanked N-terminal His$_6$-tag was cloned into the pET21a vector for recombinant bacterial expression using *Xho*I and *Nde*I restriction sites. This construct was then used to transform Rosetta-gami™ 2 (DE3) competent *E. coli* (Novagen, Merck KGaA, Darmstadt, Germany) for subsequent recombinant protein production following induction with 1 mM IPTG (Carl Roth, Karlsruhe, Germany) according to a standard protocol. For purification, bacteria pellets were resuspended in lysis buffer (20 mM Tris-HCl, pH 8.0, 0.5 M NaCl, 5 mM EDTA, 0.1% Triton X-100, with added protease inhibitor tablets according to manufacturer's instructions) and cells disrupted in a EmlsiFlex-C5 high pressure homogenizer (Avestin Europe GmbH, Mannheim, Germany), the raw extract cleared via centrifugation at 20.000 × g and the resulting pellet, containing recombinant His-CXCL4L1 in inclusion bodies, was washed in lysis buffer with and without detergent. Inclusion bodies were solubilized by gentle shaking

overnight in 50 mM Tris-HCl, pH 8.0, 6 M guanidine-HCl, 0.5 M NaCl, 10 mM DTT and His-CXCL4L1 purified from the solubilized pellet via IMAC on a HisTrap HP column on an ÄKTA Pure 25 M FPLC system (Cytiva Europe GmbH, Freiburg, Germany). The obtained protein was subjected to two dialysis steps in refolding buffer (50 mM Tris-HCl, pH 8.0, 0.5 M NaCl, 5 mM methionine, 5 mM cysteine) with and subsequently without 0.9 M guanidine-HCl, followed by a final purification step by size exclusion chromatography in 20 mM sodium phosphate buffer, pH 7.4, using a Superdex 75 10/300 GL column (Cytiva Europe GmbH) on an ÄKTA Pure 25 M FPLC system. A purity degree of 90-95% was verified by SDS-PAGE followed by Coomassie staining and Western Blot according to standard protocols.

Recombinant human peroxiredoxins 1 and 6 (PRX1, PRX6) were purchased from Abcam (Abcam PLC, Cambridge, UK), while recombinant human β-defensin-1 and 2 (HBD-1, HBD-2) were obtained from ProSpec (ProSpec-Tany TechnoGene Ltd., Ness Ziona, Israel). Recombinant human HMGB1 was purchased from Novus (Novus Biologicals Europe, Abingdon, UK). Recombinant human CXCL4L1 (PF4var1) as well as the CXCL4 (PF4) were purchased from ChromaTec (Greifswald, Germany). The other recombinant human chemokines were obtained from Peprotech (Hamburg, Germany). All other reagents and chemicals were purchased from Merck KGaA (Darmstadt, Germany), Carl Roth GmbH (Karslruhe, Germany), or Sigma-Aldrich and were of the highest purity degree available.

Cell culture and cultivation of mammalian cell lines

Jurkat T cells were cultured in RPMI1640 medium (Gibco) supplemented with 10% fetal calf serum (FCS), 1% penicillin/streptomycin, and 1x non-essential amino acids (NEAAs, Gibco). The human monocytic cell line MonoMac6 (Ziegler-Heitbrock *et al*, 1988) was cultured in RPMI1640 medium + GlutaMAX (1x), supplemented with 1x NEAAs, 10% FCS, and 1% penicillin/streptomycin. HEK293 cells stably transfected with human CXCR4 (HEK293-CXCR4) were used at passage 5 and were cultivated in DMEM medium (Gibco), supplemented with 10% FCS and 1% penicillin/streptomycin (Gibco), and used for the experiment between passage 6 and 8.

Unless stated otherwise, cells were cultivated in a temperature- and humidity-

controlled incubator at a temperature of 37°C and 5% CO_2. FCS from an EU-approved origin

was obtained from Invitrogen-Thermo Fisher Scientific and heat-inactivated prior to usage.

Other cell culture reagents, media and supplements were bought from Invitrogen-Thermo

Fisher Scientific, unless stated otherwise. Cell lines were originally obtained from the

German Society for Microorganisms and Cell Cultures (DSMZ, Braunschweig, Germany) or

from the American Type Culture Collections (ATCC).

Isolation of primary human CD4+ T cells

Primary human CD4-positive T cells were isolated from enriched peripheral blood

mononuclear cell (PBMC) fractions using the human CD4+ T cell isolation kit from Miltenyi

Biotec (Bergisch Gladbach, Germany) according to the manufacturer's instructions. Cells

were cultivated in RPMI1640 medium, supplemented with 10% FCS, 1% penicillin-

/streptomycin, and 1x NEAAs in a cell culture incubator at 37°C and 5% CO_2 and used for

functional assays on the next day. PBMC fractions were obtained by apheresis from conical

chambers of a Leucoreduction System Chamber sourced from anonymous platelet donations

at the Department of Transfusion Medicine, Cell Therapeutics and Hemostaseology of LMU

University Hospital. Studies abide by the Declaration of Helsinki principles and were

approved by ethics approval 18-104 of the Ethics Committee of LMU Munich, which

encompasses the use of anonymized tissue and blood specimens for research purposes.

Isolation of human platelets

For immunofluorescent stainings

Human platelets were isolated from blood, freshly drawn from healthy donors, using a

syringe containing 1/10 volume of CTAD–buffer (0.105 M tri-sodium citrate, 10 mM

theophylline, 3.7 mM adenosine, 0.198 mM dipyridamole) (Polack *et al*, 2001). To prevent

platelet activation, the blood was supplemented with prostaglandine E1 (Merck KGaA),

Apyrase (New England Biolabs GmbH, Frankfurt am Main, Germany), and EGTA (Sigma-

Aldrich). Briefly, platelets were isolated by sequential centrifugation steps, performed at room

temperature (RT) with reduced brake settings. Platelet-rich plasma (PRP) was separated

from whole blood by centrifugation for 5 min at 300 × g, diluted with an equal volume of

phosphate-buffered saline (PBS), pH 7.4, and centrifuged again for 10 min at 200 × g to

remove remaining leukocytes. Finally, platelets were sedimented by centrifugation for 10 min

at 400 × g.

For functional studies

Washed human platelets were isolated as previously described (Borst *et al*, 2012) and

subsequently used for functional flow chamber or platelet spreading assays.

Mice and preparation and cultivation of primary mixed cortical cultures for the

microglia motility assay

CX3CR1$^{GFP/+}$ mice, which were originally obtained from the Jackson Laboratories (strain

005582; (Niess *et al*, 2005)), were established on a pure C57BL/6 background and housed

under standardized light-dark cycles in a temperature-controlled air-conditioned environment

under specific pathogen-free conditions at the Center for Stroke and Dementia Research

(CSD), Munich, Germany, with free access to food and water. Animals were sacrificed under

anaesthesia with a mixture of midazolam (5 mg/mL), medetomidine and fentanyl (MMF).

Mouse maintenance and experiments were reviewed and overseen by the institutional

animal use and care committee of the local authorities (Regierung von Oberbayern, ROB,

Germany) and performed in accordance with the procedures provided by the animal

protection representative of CSD.

Primary mixed cortical cultures containing CX3CR1$^{GFP/+}$ microglia were prepared in

96-well imaging plates based on a previously established protocol (Gokce & Sudhof, 2013)

from the cortices of 5 newborn pups of the CX3CR1$^{GFP/+}$ mouse line (postnatal day 0) in

plating medium, consisting of modified Minimum Essential Medium (MEM without glutamine

and phenol red) (Gibco), supplemented with 0.5% glucose, 0.02% sodium bicarbonate, 1x

ITS-supplement (Sigma-Aldrich), 2 mM L-glutamine (Gibco), 1% penicillin/streptomycin, and

10% FCS. Cultures were incubated in a humidified atmosphere at 37 °C and 5% CO_2 for 10

d. One day after plating, 80% of the plating medium was replaced with growth medium,

prepared from MEM (without glutamine and phenol red) supplemented with 0.5% glucose,

0.02% sodium bicarbonate, 5% FCS, 0.5 mM L-glutamine, and serum-free B-27™

supplement (Gibco). On the fourth day after dissection, 50% of the medium was replaced

with growth medium additionally supplemented with 4 µM cytosine-1-β-D-arabinofuranoside

(Sigma-Aldrich).

Chemokine protein array

Human chemokines and selected atypical chemokines were spotted on a nitrocellulose

membrane at 100 ng per spot and left to dry at RT. Membranes were blocked with 1x

ROTI®Block (Carl Roth) for 2 h at RT and then probed overnight with biotinylated human

MIF (biotin-MIF) at a concentration of 1 µg/mL in either 10 mM Tris-HCl pH, 8.0 or 10 mM

MES, pH 6.0. Subsequently, membranes were washed three times with 0.01% Tween®20 in

water and developed with horseradish-peroxidase (HRP)-conjugated streptavidin (Bio-

Techne GmbH, Wiesbaden-Nordenstadt, Germany), diluted 1:200 in 1x ROTI®Block, for 2 h.

After another washing step, bound biotin-MIF was revealed via chemiluminescence using

SuperSignal™ West Pico Chemiluminescent Substrate (Thermo Fisher Scientific) on a LAS-

3000 Imaging System (Fuji Photo Film Co., LTD., Japan).

Pull-down of CXCL4L1 from cell lysates

MonoMac-6 cells were first washed with PBS and then lysed on ice for 30 min with

immunoprecipitation (IP) lysis buffer (1x cell lysis buffer, Cell Signaling, cat#9803), 100 mM

PMSF, and 1x protease and phosphatase inhibitors (ThermoFisher). The purified cell lysates

were then incubated with pre-washed streptavidin-conjugated paramagnetic beads

(DYNAL™ Dynabeads™ M-280 Streptavidin; Invitrogen, cat#11205D) for 2 h at 4 °C

(preclearing step). After centrifugation, the supernatant was incubated with biotinylated

human MIF by gentle, constant shaking on a rotary shaker overnight at 4 °C. To capture

MIF/CXCL4L1 complexes, prewashed streptavidin-conjugated beads were added to the

precleared lysates, and the mixture was incubated for 2 h at 4 °C on a rotary shaker. Beads

were separated from the lysate using a magnetic stand (Dynal™ MCP-S) and washed three

times with lysis buffer. The supernatant was removed and the beads were resuspended in

LDS sample buffer (Invitrogen) and boiled at 95 °C for 15 min. Samples were subjected to

SDS-PAGE and analyzed by Western blotting. For this purpose, equal amounts of protein

were loaded onto 11% SDS-polyacrylamide gels (NuPAGE, Thermofisher) and transferred to

polyvinylidene difluoride (PVDF) membranes (Carl Roth, Karlsruhe, Germany). Membranes

were blocked in PBS-Tween-20 containing 5% BSA for 1 h and incubated overnight at 4 °C

with rabbit polyclonal anti-MIF antibody Ka565 (Bernhagen *et al.*, 2007) or rabbit polyclonal

anti-PF4V1 IgG PA5-21944 (Invitrogen) diluted in blocking buffer. Proteins were revealed

using anti-rabbit HRP as a secondary antibody. Signals were detected by chemilumi-

nescence on an Odyssey® Fc Imager (LI-COR Biosciences GmbH, Bad Homburg,

Germany) using SuperSignal™ West Dura ECL substrate from ThermoFisher Scientific and

specific primary antibodies as indicated.

CelluSpot peptide array

The CelluSpot peptide array method has been described previously (Lacy *et al.*, 2018).

Briefly, 15-meric peptides, positionally frame-shifted by three residues and spanning the

entire sequence of CXCL4 and CXCL4L1, were synthesized on modified cellulose disks

(Intavis MultiPep RSi/CelluSpot Array, Cologne, Germany). Peptides were then further

processed by dissolving the cellulose, and spotted on coated glass slides using a slide

spotting robot from Intavis. Slides were incubated in blocking buffer (50 mM Tris-buffered

saline, pH 7.4, 1% BSA, 0.1% Tween® 20), washed (50 mM Tris-buffered saline, pH 7.4,

0.1% Tween® 20) and probed with biotinylated human MIF (3 µM in blocking buffer). After

washing, slides were developed with a dilution of streptavidin-conjugated horseradish

peroxidase (Roche) in blocking buffer. Bound MIF was revealed by chemiluminescence on

an Odyssey® Fc imager using the SuperSignal™ West Dura ECL substrate

Microscale thermophoresis (MST)

Protein-protein interactions were analyzed via microscale thermophoresis on a Monolith NT.115 instrument equipped with green/red filters (NanoTemper Technologies, Munich, Germany). Measurements were performed at 25°C at both 40% and 80% MST power. LED excitation power was adjusted to 90 or 95% in order to obtain an initial fluorescence count of 700 to 800. MST traces were recorded for 40 s (-5 s to +35 s), according to default settings with the sample being heated from 0 to 30 s. All measurements were performed in assay buffer (10 mM Tris-HCl, pH 8.0, 0.01% BSA). MST-Red-MIF was used at a fixed concentration, mixed 1:1 with serial dilutions of either CXCL4 (Peprotech, Hamburg, Germany) or CXCL4L1 (ChromaTec, Greifswald, Germany) (final MIF concentrations: 456 nM or 312 nM, respectively). Prior to measurement, the prepared samples were incubated for at least 30 min on ice. MST traces of multiple experiments were analyzed according to the K_D model using the default T-jump settings, focusing on the temperature related intensity change (TRIC) of the fluorescent label ("cold region" from -1 to 0 s, "hot region" from 0.5 to 1.5 s) using the MO.AffinityAnalysis V2.3 software (NanoTemper Technologies). Curve fitting for data presentation was performed by GraphPad Prism Version 6.07 ('one site – total binding').

Analysis of protein-protein interactions by surface plasmon resonance (SPR)

Surface plasmon resonance measurements were performed using a Biacore X100 instrument (GE Healthcare Europe GmbH) and neutravidin-modified C1 sensor chips. Biotin-MIF was immobilized on flow cells to 1064.8 RU. CXCL4 (Peprotech, Hamburg, Germany) and CXCL4L1 (ChromaTec, Greifswald, Germany), used at concentrations in the range of 0.125 to 20 µg/mL in running buffer (HBS-EP+ Buffer: 0.01 M HEPES, 0.15 M NaCl, 0.003 M EDTA and 0.05% v/v surfactant P20) were injected at a flow rate of 60 µL/min. The complex was allowed to associate and dissociate for 90 s and 240 s, respectively. Surfaces were regenerated with 2 pulses (60 s) of 30 mM NaOH and 2 M NaCl. Responses from analyte

325   injections were fitted to a 1:1 Langmuir interaction profile using Biacore X100 evaluation

326   2.0.1 Plus package software.

Transwell migration assay

Transwell migration experiments to study the influence of CXCL4L1 on MIF-mediated

chemotaxis responses were performed with Jurkat T cells. Briefly, Jurkat cells were diluted in

RPMI1640 medium at a density of 1×10^7 cells/mL. Cells were placed in the upper chamber

of a 24-well Transwell insert with 5 µm pore size (Corning, Kaiserslautern, Germany). 16 nM

MIF, either alone or pre-incubated (30 min on ice to allow for complex formation) with 32 nM

of CXCL4L1, as well as 32 nM CXCL4L1 alone were added to the lower chamber as a

chemoattractant. After a 12 h migration interval at 37 °C and 5% CO_2, migrated cells were

recovered from the lower chamber and counted via flow cytometry by using CountBright™

absolute counting beads (Molecular Probes-Invitrogen). In a similar experimental setup, the

influence of CXCL4 on MIF-mediated chemotaxis was tested as well. MIF was used at a

concentration of 16 nM and CXCL4 (ChromaTec, Greifswald, Germany) at 32 nM.

3D migration of human CD4[+] T cells

The migratory behavior of primary human T cells was assessed by three-dimensional (3D)

migration methodology using time-lapse microscopy and single cell tracking using the 3D

chemotaxis µ-Slide system from Ibidi GmbH (Munich, Germany). The method was performed

following a slight modification of the established Ibidi dendritic cell protocol for human

monocytes, as described previously (Kontos *et al.*, 2020). Briefly, isolated CD4[+] human T

cells (3.5×10^6) were seeded in a rat tail collagen type-I gel (Ibidi, Munich, Germany) in

DMEM and subjected to a gradient of human MIF, CXCL4L1, or a pre-incubated combination

of both. Cell motility was monitored performing time-lapse imaging every 0.5 or 2 min at 37

°C for a period of 120 min to cover either a short or extended migration period, using a Leica

DMi8 inverted microscope (Leica Microsystems, Wetzlar, Germany) and Leica live cell-

imaging software (LAS X version 3.7.4). Images were imported as stacks to ImageJ version

1.51n and analyzed with the manual tracking and Chemotaxis and Migration tool (Ibidi

GmbH) plugin for ImageJ.

Motility measurement of primary murine microglia

The motility of mouse microglia was determined using mixed cortical cultures, established

and cultivated as stated above. A day prior to imaging, the medium was changed to

Hibernate A medium (Gibco) in order to maintain a physiological pH value during imaging.

Prior to imaging, different wells of cells from each individual pup were treated with either 8

nM MIF, 1.6 nM CXCL4L1, or both (pre-incubated for 30 min on ice to allow for complex

formation). A control group was treated with 20 mM sodium phosphate buffer, pH 7.4. Cell

motility was monitored by time-lapse imaging for 15 h at 37 °C with recordings every 5 min,

using a Leica DMi8 inverted Life Cell Imaging System using the FITC channel for visualizing

the GFP-positive cells. Images were imported as stacks to ImageJ software version 1.51n

and analyzed with the manual tracking and Chemotaxis and Migration tool Plugin for ImageJ

from Ibidi. In order to quantify microglial motility from the time-lapse videos, 20-25 GFP-

positive microglia per treatment group were randomly selected and manually tracked

throughout all frames. Cells that died or moved out of the frame were excluded from the

analysis. Accumulated distance of each tracked microglia was calculated with Chemotaxis

and Migration Tool (Ibidi).

Fluorescence polarization spectroscopy

Fluorescence polarization was measured using a JASCO FP-6500 fluorescence

spectrophotometer equipped with FDP-223 and FDP-243 manual polarizers (JASCO

Deutschland GmbH, Pfungstadt, Germany). Preparation of stock solutions, measurements

and analysis were performed essentially following a previously published protocol (Kontos *et*

al., 2020). For binding/inhibition experiments, mixtures of Alexa 488-labeled MIF (10 nM) in

the absence/presence of CXCL4L1 (1.6 µM) (or 20 mM sodium phosphate buffer, pH 7.2),

and non-labeled msR4M-L1 peptide (concentration between 1 nM to 10 µM) were prepared

in 10 mM sodium phosphate, pH 7.2, containing 2% hexafluoro-isopropanol (HFIP). Where

CXCL4L1 was added as a putative inhibitor, Alexa488-MIF and CXCL4L1 were mixed and

incubated for 30 min prior to measurements. Bandwidth for excitation and emission was set

at 5 nm and time response at 0.5 s. The excitation wavelength was 492 nm and emission

was recorded at 519 nm. Measurements were taken at RT within 2 to 3 min upon preparation

of the solutions. Polarization P was calculated according to the equation $P = (I_\parallel - G \cdot I_\perp) / (I_\parallel +$

$G \cdot I_\perp)$, with $I_\parallel$ as the intensity of emitted light polarized parallel to the excitation light and $I_\perp$ as

the intensity of emitted light polarized perpendicular to the excitation light. The G factor was

calculated based on the instrumental documentation (Moerke, 2009). Apparent K_D values

were calculated assuming a 1:1 binding model (Yan *et al*, 2006), using sigmoidal curve fitting

with OriginPro 2016 (OriginLab Corporation, Northampton, MA, USA).

Label-free dynamic mass redistribution (DMR) assay

Analysis of dynamic mass redistribution of adherent cells was performed on an EnSpire

Multimode plate reader equipped with an Epic® label-free measurement module

(PerkinElmer Inc., Waltham MA, USA) according to the manufacturer's instructions for cell-

based label-free DMR measurements (Schroder *et al*, 2010). The assay protocol was

adapted according to a previous publication to be performed in EnSpire label-free 96-well

fibronectin-coated cell assay microplates (Corning GmbH, Amsterdam, The Netherlands)

with HEK293 cells stably expressing human CXCR4 (Krammer *et al*., 2021). Briefly, 40.000

cells were seeded into each well and cultivated overnight (37°C, 5% CO_2) to achieve a

confluency of >70%. Prior to the assay, the medium was exchanged with DMR assay buffer

(20 mM HEPES and 1% DMSO in HBSS, pH 7.4) and the assay plate left for 6 h to

equilibrate to ambient temperature. Baseline measurements for each well were recorded for

10 min every 30 s prior to treatment of the cells with chemokines, inhibitors, or the

corresponding buffers as control. Treatments were applied to each well as a 5x concentrated

stock, prepared in assay buffer. For treatment with MIF/CXCL4L1 complexes, both proteins

were mixed in assay buffer and incubated for 5 h at RT. Directly after addition of the stimuli,

the DMR response was recorded for the indicated duration. The DMR response resembles

the wavelength shift of the light reflected from the sensor integrated in the assay microplates

and serves as a cumulative cellular response signal. Measurements were performed on two

replicates per treatment and results are presented as their mean value.

Staining of human thrombus specimens

Human thrombus tissue specimens, obtained as disposable material from vascular surgery

procedures (ethics allowance LMU Munich 18-104 and TUM-MRI project # 2799/10), were

embedded in Tissue Tek O.C.T. Compound (Sakura Finetek Germany GmbH, Staufen,

Germany), frozen, and cut into 5 µm sections using a CM 1950 Cryostat (Leica Biosystems).

The cryosections were transferred to microscopy slides and stored at -80°C until use.

Hematoxylin & eosin (HE) staining of thrombus sections

Cryosections of human thrombus tissue were stained with Mayer's hematoxylin and eosin

(HE) according to standard protocols. Briefly, after thawing and brief air drying and

rehydration, sections were incubated in Mayer's hematoxylin solution (Sigma-Aldrich) for 15

min. After thorough rinsing of the samples, 0.5% eosin Y solution (Sigma-Aldrich) was

applied as a counterstain for 30 s. After dehydration of the tissue in 95 % ethanol, 100 %

ethanol (Merck KGaA) and xylene (VWR International GmbH), the sample was covered with

resinous mounting medium (Eukitt, Sigma-Aldrich), covered with a glass coverslip and

examined by light microscopy (Leica Dmi8 inverted microscope using a DMC2900 digital

camera (Leica Microsystems; 10x objective).

Staining of platelets

Freshly isolated platelets were fixed with 4% paraformaldehyde (PFA) in PBS (Morphisto

GmbH, Frankfurt a. M., Germany) for 10 min and subsequently permeabilized using 1x Perm

buffer (Invitrogen) for 15 min. After washing, platelets were blocked in ROTI®Block (Carl

Roth) for 1 h. Immunofluorescent staining was performed as described above for the human

thrombus specimens, except that the anti-human MIF antibody was used at a dilution of 1:20

and the anti-human CXCL4L1 antibody at a dilution of 1:50. Stained platelets were then

washed in blocking buffer and mounted on poly-L-ornithine-coated glass slides using

ProLong^{TM} Glass Antifade mountant (Invitrogen), covered with coverslips and stored at 4 °C

until imaging by multiphoton microscopy.

Multiphoton laser-scanning microscopy (MPM) and FLIM-FRET

Imaging was conducted using a multispectral TCS SP8 DIVE FALCON LIGHTNING

microscope (Leica, Germany) equipped with filter-free 4TUNE NDD detection module, an

extended IR spectrum tunable laser (New InSight® X3™, Spectra-Physics) (680-1300 nm)

and fixed IR laser (1045 nm), advanced Vario Beam Expander (VBE), Ultra-high-speed

resonance scanner (8kHz), HC PL IRAPO 25x/1.0 WATER objective, and FLIM-FRET

modality. Images were collected in a sequential scanning mode using hybrid diode detectors

Reflected Light Hybrid Detectors (HyD-RLD) (Alexa Fluor-488: excitation 965 nm / emission

479-568 nm; Cy3: excitation 1095 nm / emission 538-650 nm) and were handled using the

LAS-X software package. Deconvolution microscopy was performed using the Leica

LIGHTNING (adaptive deconvolution) application.

For fluorescence lifetime imaging (FLIM) and FLIM-FRET measurements, up to 1000

photons per pixel were captured in a time-correlated single photon counting (TCSPC) mode.

Fluorescence lifetime decay data were fitted using Leica FALCON (FAstLifetime CONtrast)

software. The fitting was assessed by randomly distributed residuals and by low Chi-square

(χ^2) values. The number of components used for the fittings was manually fixed to a value

(n=2-3) to minimize χ^2 values. The fluorescence lifetime of the donor was acquired similarly

in the absence of the acceptor.

Proximity ligation assay (PLA)

For detection of protein complexes by proximity ligation assay (PLA), the Duolink™ InSitu Orange Starter Kit Mouse/Rabbit (DUO92102) from Sigma Aldrich was used. Following scouting experiments to establish the PLA methodology in thrombus material, cryosections of the thrombi were prepared by treatment with cold acetone for 6 min at 4 °C and for 30 min at RT. Samples were rehydrated in PBS for 10 min and hydrophobic barriers were applied to the microscopy slide using an ImmoEdgeTM Pen (Vector Laboratories Inc. Burlingame, USA).

For PLA detection, the Duolink® PLA Fluorescence protocol provided by the manufacturer was essentially followed, using primary antibodies against human MIF (mouse anti-MIF D2, sc-271631, Santa Cruz Biotechnology Inc., Dallas, USA; 1:20) and against human CXCL4L1 (rabbit anti-CXCL4L1, PA5-21944, Invitrogen; 1:50). Samples were then prepared for microscopy using Duolink® mounting medium with DAPI, and coverslips sealed with commercially available nail polish and stored at 4 °C until imaging by confocal microscopy using a Zeiss LSM880 AiryScan microscope was performed.

Flow chamber assay with platelets

Chemokines were diluted in calcium-free PBS, pH 7.4, at their final concentrations (MIF: 16 nM; CXCL4L1: 32 nM) and allocated into separate reaction tubes. 200 µL of each solution were distributed onto separate collagen-coated cover slips (100 µg/mL) and incubated for 2 h. Cover slips were blocked with PBS, pH 7.4, containing 1% BSA for 1 h. Next, human whole-blood was diluted at a 5:1 ratio with PBS, pH 7.4, containing calcium. Before perfusion, the blood was incubated with fluorochrome 3,3′-dihexyloxacarbocyanine iodide (DiOC$_6$, 1 mM; Sigma Aldrich) for 10 min at RT. Thereafter, the blood was allocated into 1 mL syringes and perfused over the different cover slips, through a transparent flow chamber with high shear rate (1000 s^{-1}) for 5 min. Per run, one 2-minute video clip was recorded (200 ms/frame, Nikon Eclipse Ti2-A, 20x objective). Afterwards, the chamber was rinsed and pictures were taken of five representative areas using the same objective. The covered area

was analyzed using the NIS-Elements AR software (Nikon) and the mean percentage of the

covered area, the mean thrombus area as well as the mean thrombus count were

determined.

Platelet spreading analysis

Fibrinogen-coated (100 µg/mL, Sigma Aldrich) coverslips were preincubated with MIF (16

nM), CXCL4L1 (32 nM), or MIF (16 nM) and CXCL4L1 (32 nM) together, for 2 h. Afterwards,

isolated human platelets were diluted in Tyrodes buffer (pH 7.4) to match a concentration of

15.000 cells/µL. Platelets were supplemented with 1 mM $CaCl_2$, activated with 1 µg/mL CRP-

XL (CambCol, Cambridge, UK), and incubated on the previously prepared fibrinogen-coated

coverslips for 30 or 60 min at RT. Thereafter, platelets were fixed with 4% formaldehyde

(Sigma Aldrich) for 10 min, and washed three times with PBS, pH 7.4. The coverslips were

mounted onto slides and five images from randomly selected areas were taken using a Nikon

Eclipse Ti2-A microscope with a 100x DIC objective. Subsequently a quarter of each image

with at least 20 cells was analyzed.

Protein structure visualization

Three-dimensional structures as well as the surface charge distribution of human MIF,

CXCL4 and CXCL4L1 were visualized using the PyMOL Molecular Graphics System

software, version 1.8.2.2 (Schrödinger, LLC). The structures represent the Protein Data Bank

(PDB) files for MIF (PDB ID: 3DJH), CXCL4 (PDB ID: 1F9Q), and CXCL4L1 (PDB ID: 4HSV),

orour molecular docking results.

Protein-protein docking

To simulate the interaction of monomeric MIF with CXCL4 and CXCL4L1 in their monomeric

forms, rigid protein-protein docking, followed by clustering of the 1000 lowest energy

structures and removal of steric clashes was performed using the ClusPro 2.0 webserver,

with single chains of MIF and CXCL4L1 defined as 'receptor' and 'ligand', respectively

(Kozakov *et al*, 2017; Vajda *et al*, 2017).

Statistical analysis

Statistical analysis was performed using GraphPad Prism Version 6.07 software. Unless

stated otherwise, data are represented as means ± standard deviation (SD). After testing for

normality, data were analyzed either by two-tailed Student's T-test, Mann-Whitney U test, or

Kruskal-Wallis test as appropriate. Differences with $P<0.05$ were considered to be

statistically significant.

Results

High affinity binding between the atypical chemokine MIF and the platelet CXC

chemokine CXCL4L1

To begin to test the hypothesis that chemokine-chemokine interactions may extend to ACKs

and that MIF may form heterocomplexes with classical chemokines, we applied unbiased

chemokine protein array technology (*Figure 1A-B*), as previously successfully used to map

formation of heterocomplexes between different classical chemokines (von Hundelshausen

et al., 2017). In addition to 47 human chemokines covering all four sub-classes (CXC-, CC-,

CX3C- and C-type CKs) we also included structurally related and positively charged protein

mediators including ACKs/DAMPs such as HMGB1, HBDs, and peroxiredoxins (Prxs)

(Shichita *et al*, 2012; He *et al*, 2019), as well as MIF itself and the MIF homolog D-

dopachrome tautomerase (D-DT)/MIF-2 as spotted proteins in the protein array. Probing of

the array with biotin-conjugated MIF and streptavidin-POD (StrAv-POD) revealed high-

intensity spots indicative of a tight interaction of MIF with CXCL4L1 and Prx1 (*Figure 1B-C*).

Weaker spots were detected for CCL28, CXCL9, Prx6, and MIF itself. No spot intensity

whatsoever was observed for any of the other immobilized proteins, indicating that none of

the other 44 chemokines interacts with MIF. This also included CXCL8 and CXCL12, which

share their receptors CXCR2 and CXCR4, respectively, with MIF (Bernhagen *et al.*, 2007).

Similarly, no binding signal of biotin-MIF was detected with HMGB1, a DAMP which

has been demonstrated to form heterodimers with CXCL12 and for which a functional

interaction with MIF has been suggested (Ma *et al*, 2017; Schiraldi *et al.*, 2012), nor for the

human β-defensins HBD1 or HBD2 (*Figure 1B-C*). Importantly, when testing a control

chemokine array developed with StrAv-POD without a biotin MIF incubation step, only one

signal was not fully specific. This was the signal for Prx1 so that its interpretation was not

possible (*Supplementary Figure 1A*). Biotin-MIF also bound to MIF itself, but not to MIF-2

(*Figure 1B-C*). As MIF is known to form homo-oligomers (Sun *et al.*, 1996) and has been

reported to form higher-order hexameric complexes (Bai *et al*, 2012), this result further

verified the validity of the chemokine array approach for MIF.

A striking observation was that biotin-MIF specifically interacted with the immobilized

platelet chemokine CXCL4L1, but not with CXCL4 (*Figure 1C and Supplementary Figure*

1B). CXCL4 and CXCL4L1 are highly homologous chemokines, their sequences only differ

by three amino acids, and CXCL4L1 has also been suggested to be a decoy chemokine

paralog of CXCL4. Given this remarkable specificity of the interaction with MIF and that the

spot corresponding to biotin-MIF and CXCL4L1 was the strongest interaction detected on the

array, we focused on CXCL4L1 as a novel candidate interactor of MIF.

We first verified the interaction by co-immunoprecipitation using whole cell lysates of

MonoMac6 cells, which we found to express substantial amounts of CXCL4L1. Semi-

endogenous pulldown of proteins from MonoMac6 lysates by biotin-MIF and StrAv magnetic

beads and Western blot using an anti-human CXCL4L1 antibody revealed a specific band for

CXCL4L1, which was absent when the pulldown was performed without biotin-MIF

preincubation (*Figure 2A*). Pulldown specificity was further confirmed by Western blot against

MIF. We next applied surface plasmon resonance ('Biacore') methodology, which was

previously successfully used to characterize interactions within the classical chemokine

interactome (von Hundelshausen *et al.*, 2017). To study the MIF/CXCL4L1 interaction, MIF

chips were exposed to increasing concentrations of CXCL4L1 in the soluble phase. The

obtained surface plasmon resonance response curves indicated that MIF specifically binds to

CXCL4L1 (*Figure 2B-C*). Quantitative analysis determined a K_D value of 116 ± 16 nM (mean

± SD) indicating high-affinity binding between MIF and CXCL4L1. By contrast, no

appreciable signal was detectable for the incubation with increasing concentrations of

CXCL4 and no K_D could be derived, verifying the specificity of the MIF/CXCL4L1 interaction

in this set-up. To further confirm the MIF/CXCL4L1 interaction, we next applied microscale

thermophoresis (MST), which relied on the interaction between MST-Red-labeled MIF and its

binding partner, with both partners in the soluble phase. This methodology was recently

established for MIF (Kontos *et al.*, 2020). MST titrations of MST-Red-MIF with increasing

concentrations of CXCL4L1 revealed a typical sigmoidal binding curve with a derived binding constant (K_D = 159.8 ± 16.8 nM) that was similar to that obtained by surface plasmon resonance (*Figure 2D-E*). In contrast, binding was much weaker when CXCL4 was titrated and accordingly a low affinity K_D in the micromolar range was determined (K_D = 2.0 ± 0.8 µM).

Heterodimer formation between classical chemokines relies on CC-type or CXC-type interactions. To determine which residues in CXCL4L1 are critical for the interaction with MIF, we employed peptide array technology. A set of 15-meric peptides derived from the CXCL4L1 sequence, positionally frame-shifted by three amino acids to cover the entire sequence of the processed chemokine, were synthesized and immobilized on glass slides and arrays, and probed with biotin-MIF. The most pronounced binding signal was observed for peptides representing the sequence region, which corresponds to the β2-strand motif IKAGPHCPTAQLIAT of CXCL4L1 (*Supplementary Figure 2A-B*). A second peak encompasses the N-terminal sequence QCLCVKTTSQVRPRH. The difference in the 3D structures of CXCL4 and CXCL4L1 is characterized by a significant conformational rearrangement of the α-helix (Kuo *et al*, 2013), although the sequence of CXCL4 differs from that of CXCL4L1 in only three α-helical residues (L58P, K66E, L67H with the conformational difference being mainly governed by the L67H exchange). In this respect, CXCL4 showed an essentially identical peptide binding profile as that of CXCL4L1 at the N-terminus as expected, but a slightly different pattern at the β2 strand region GPHCPTAQLIATLKN, that is packed onto the C-terminal α-helix (*Supplementary Figure 2A-B*). Peptide array-based mapping of the CXCL4L1 residues involved in MIF binding was confirmed by molecular docking simulations. Docking applying the ClusPro software predicted that the β-sheet region including the IKAGPHCPTAQLIAT motif is located near the MIF contact site, facing the 4-stranded β-sheet of a single MIF monomer chain. This interaction could be promoted by an energetically favorable complementary electrostatic interaction between the two surfaces (*Supplementary Figure 2C*).

Together, the co-immunoprecipitation, Biacore, and MST studies confirmed specific

binding between MIF and CXCL4L1 and determined a high-affinity binding constant in the

100-150 nM range for the interaction. Analysis of the binding interface by peptide array-

based mapping and molecular docking provides an initial prediction of the residues involved

in the CXCL4L1/MIF binding site.

MIF/CXCL4L1 heterocomplex formation attenuates MIF-mediated inflammatory/athero-

genic activities

We next wished to determine a potential functional role of MIF/CXCL4L1 heterocomplex

formation. CXCL4L1 is a potent angiostatic chemokine acting through CXCR3 (Struyf *et al*,

2011), but its role in inflammatory responses and atherogenesis is not well understood. Pro-

atherogenic activities of MIF have been extensively characterized and are mainly mediated

through non-cognate interaction of MIF with CXCR2 and CXCR4 (Bernhagen *et al.*, 2007;

Sinitski *et al.*, 2019). Here, we hypothesized that MIF/CXCL4L1 complex formation could

predominantly influence CXCR4-mediated pathways of MIF.

We first asked whether MIF-elicited T-cell chemotaxis, a well-characterized

atherogenic MIF effect mediated via T-cell-expressed CXCR4 (Bernhagen *et al.*, 2007), is

affected by CXCL4L1. Scouting experiments using Jurkat T-cells confirmed that, when added

to the lower chamber of a Transwell migration device as a chemotattractant, MIF elicited

chemotaxis with a chemotactic index (CTX) of approximately 2. Moreover, when CXCL4L1

was preincubated with MIF to allow for complex formation, no upregulation of Jurkat T-cell

chemotaxis was observed, while CXCL4L1 alone exhibited neither a chemotactic nor

inhibitory effect (*Supplementary Figure 3A*). In line with the observed lack of binding between

MIF and CXCL4, MIF-mediated Jurkat T-cell chemotaxis was not attenuated by co-

incubation with CXCL4, which by itself did not significantly enhance Jurkat T-cell chemotaxis

(*Supplementary Figure 3B*). To test the physiological relevance of this finding, we next

studied primary CD4[+] T-cell chemotaxis and also applied a three-dimensional migration set-

up, following individual cell migration trajectories by live cell imaging. MIF potently triggered

T-cell migration as evidenced by a significant increase in forward migration index (FMI) (*Figure 3A-B*), confirming previous data showing CXCR4-dependent stimulation of monocyte migration by MIF (Kontos *et al.*, 2020). This effect was abrogated when MIF was coincubated with CXCL4L1, while CXCL4L1 alone had no effect on 3D T-cell motility. This suggested that MIF/CXCL4L1 heterocomplex formation interferes with MIF/CXCR4-stimulated chemotaxis of T cells.

To study the potential relevance of these findings for other inflammatory/immune cell types, we next evaluated the effect of MIF/CXCL4L1 complex formation on microglial motility in the physiological setting of cortical brain cultures. MIF promotes the motility of Egfp[+] microglia in murine cortical brain cultures *ex vivo* in a Cxcr4-dependent manner, as read out by live microscopy and as indicated by blockade of the MIF effect by the soluble CXCR4 mimicking peptide msR4M-L1(*Supplementary Figure 3C*). Importantly, MIF-triggered microglia migration in this setting was fully ablated when CXCL4L1 was added together with MIF following preincubation, while CXCL4L1 alone had no effect on microglia motility (*Figure 3C-D*). This indicated that CXCL4L1/MIF heterocomplex formation attenuates MIF's CXCR4-dependent effect on microglia migration.

MIF/CXCL4L1 heterocomplex formation inhibits MIF binding to CXCR4

The cell migration experiments implied, but did not directly test, the notion that MIF/CXCL4L1 complex formation affects MIF signaling through the CXCR4 pathway. To test the involvement of CXCR4 directly, we performed a binding competition experiment that capitalized on our recent identification of a MIF-binding CXCR4 ectodomain-mimicking peptide msR4M-L1 (Kontos *et al.*, 2020). Employing fluorescence polarization spectroscopy (FP), titration of increasing concentrations of msR4M-L1 with Alexa 488-MIF led to a pronounced sigmoidal change in the FP signal (*Figure 4A*), in line with previous data showing high affinity binding between MIF and msR4M-L1 (Kontos *et al.*, 2020). By contrast, when Alexa 488-MIF was preincubated with CXCL4L1 before the titration, the FP signal was

ablated (*Figure 4A*), suggesting that MIF/CXCL4L1 heterocomplex formation interfered with

MIF binding to the CXCR4 mimic.

To further confirm an interference of heterocomplex formation with the MIF/CXCR4

pathway, we next studied dynamic mass redistribution (DMR) responses in HEK293 cells

stably transfected with human CXCR4. Incubation of HEK293-CXCR4 transfectants with MIF

but not control buffer led to a pronounced time-dependent increase in the DMR signal as a

real-time readout of an integrated cellular response of living HEK293 cell activation through

the MIF/CXCR4 receptor signaling pathway (*Figure 4C*). This signal was markedly

attenuated by the small molecule CXCR4 inhibitor AMD3100, whereas the DMR curve of

AMD3100 alone was similar to the control buffer curve, confirming CXCR4-dependency of

the MIF-induced signal. Of note, preincubation of MIF with CXCL4L1 led to an appreciable

reduction in the DMR response curve as well, when compared to cell stimulation with MIF

alone, while CXCL4L1 alone and buffer control showed no effect (*Figure 4C*).

Together, the competition binding study and the DMR experiment confirmed the

notion that complexation by CXCL4L1 interferes with binding of MIF to CXCR4 and its ability

to activate CXCR4-mediated cell responses.

MIF and CXCL4L1 colocalize and form complexes in human platelet aggregates and

clinical thrombus specimens

CXCL4L1 is an abundant platelet chemokine (Karshovska *et al*, 2013; von Hundelshausen *et*

al, 2007) and we previously found that platelets also are a rich source of MIF (Strüßmann *et*

al, 2013). The colocalization of MIF and CXCL4L1 in sub-cellular platelet compartments has

not yet been studied, but a cell biological characterization of CXCL4 suggested that this

paralog may be localized in a different intracellular platelet compartment than MIF

(Strüßmann *et al.*, 2013). Notwithstanding, we surmised that colocalization and complex

formation between MIF and CXCL4L1 may occur extracellularly after secretion from

activated platelets.

Initial evidence for a colocalization of CXCL4L1 and MIF following co-secretion from

activated platelets came from human platelet preparations that aggregated due to handling

stress. Examination of these aggregates by multi-photon microscopy (MPM) using an Alexa

488 signal to label MIF and Cy3 immunofluorescence for CXCL4L1 revealed several areas

with an apparent colocalization of MIF and CXCL4L1 (*Figure 5A*). Colocalization was also

detectable in areas with more isolated non-aggregated platelets (*Figure 5B*). These areas

were then subjected to an in-depth analysis by fluorescence lifetime imaging-Förster

resonance energy transfer (FLIM-FRET) capitalizing on the Alexa 488/Cy3 FRET donor-

/acceptor pair. For molecule-molecule interactions within a distance range of 1-10 nm, FLIM-

FRET monitors the change in fluorescence lifetime of the donor via FRET and directly

visualizes the proximity of the donor (*here*: Alexa 488-labeled anti-mouse IgG secondary

antibody in combination with mouse anti-MIF) and the acceptor molecule (*here*: Cy3-labelled

anti-rabbit secondary antibody in combination with rabbit anti-CXCL4L1). We detected

significant donor lifetime shortening (from 2.019 ± 0.069 ns to 1.496 ± 0.033 ns) and FRET

events (FRET efficiency peak at 20-25%), when Alexa 488/Cy3 FLIM-FRET was recorded in

appropriate regions-of-interest (ROIs) (*Figure 5C-D*), an observation that is consistent with

the notion that MIF and CXCL4L1 not only colocalize in activated platelet preparations but

form true heterocomplexes.

To further investigate the physiological relevance of these findings, we next examined

clinical thrombus specimens derived from vascular surgery procedures. To determine

whether colocalized MIF and CXCL4L1 formed heterocomplexes in thrombus tissue, a

proximity ligation assay (PLA) was performed which detects inter-molecular interactions

within a distance of <10 nm. Specific PLA signals were detected in an atherosclerotic

thrombus specimen (*Figure 6A-B*), suggesting the abundant occurrence of MIF/CXCL4L1

heterocomplexes in the context of clinical thrombus tissue and confirming the FLIM-FRET

data obtained in platelet preparations from healthy blood samples. Thus, both FLIM-FRET

and PLA demonstrated that MIF and CXCL4L1 form heteromeric complexes upon release

from activated platelets.

Heterocomplex formation inhibits MIF-stimulated thrombus formation and alters the effect of MIF on platelet morphology

Thrombus formation and clot retraction are relevant processes upon vessel injury and in advanced atherosclerotic vessels. MIF was found to modulate these processes (Wirtz *et al*, 2015). As our data showed that MIF/CXCL4L1 heterocomplexes form in the micro-environment of a thrombus, we next determined whether heterocomplex formation affects thrombus characteristics. Thrombus formation under flow perfusing diluted human blood over a collagen-coated surface harboring combinations of MIF and CXCL4L1 was studied as established (Chatterjee *et al*, 2014) and was found to double following exposure to MIF when applying a shear rate of 1000 s^{-1} (*Figure 7*). CXCL4L1 alone did not affect thrombus characteristics, but when added together with MIF following preincubation, MIF-elicited thrombus formation was blocked. These effects were mainly related to thrombus size/coverage (*Figure 7B*, *Supplementary Figure 5*) rather than thrombus numbers (*Figure 7C*). These data indicated that heterocomplex formation inhibited MIF-stimulated thrombus formation.

The role of platelet morphology and lamellipodia in stable thrombus formation has been controversial, but platelet lamellipodia formation is critical for thrombus formation under flow (Fotinos *et al*, 2015; Kraemer *et al*, 2011a; Schurr *et al*, 2019). To further study the above observed effect of heterocomplex formation on thrombus behavior, we examined the morphology of flow-stressed adhered platelets exposed to MIF or heterocomplexes in detail. Platelet flow stress responses were recorded after 30 and 60 min, with significant changes observed for the 30 min time point. Morphological changes encompassed increased platelet numbers with filopodia, small lamellipodia, large lamellipodia, as well as fully spread platelets. Interestingly, the strong increase in large lamellipodia under control buffer conditions was significantly reduced by MIF and a further significant reduction was observed for platelets coincubated with MIF and CXCL4L1. Inversely, the incubation with the heterocomplex resulted in a significant increase in platelets with small lamellipodia compared to stimulation with MIF alone (*Figure 7D*). *Figure 7E* further illustrates the inverse effect of

1   MIF/CXCL4L1 on large *versus* small lamellipodia formation. Together, these experiments

2   indicated MIF/CXCL4L1 heterocomplex formation skewed the morphology of adhering flow-

3   stressed platelets from a large to a small lamellipodia phenotype compared to treatment with

4   MIF alone.

Discussion

Chemokines control numerous pathogenic pathways contributing to inflammation and atherogenesis. The recent systematic characterization of the chemokine interactome revealed that heteromeric interactions between classical CC- and/or CXC-type chemokines represent an important molecular adjustment screw that serves to amplify, inhibit, or modulate chemokine activity (von Hundelshausen *et al.*, 2017). Here, we have identified a heteromeric interaction between MIF, a pleiotropic inflammatory cytokine and ACK, and the classical platelet chemokine CXCL4L1. We also show that CXCL4L1/MIF complex formation affects inflammatory/atherogenic and thrombogenic activities of MIF. The scheme in *Figure 8* summarizes the main findings of this study. This suggests that disease-relevant activities of MIF may be fine-tuned by heterocomplexation with CXCL4L1 and that the chemokine interactome extends to heteromeric interactions between classical and atypical chemokines.

In fact, binding of classical chemokines to non-CC- or CXC-chemokine mediators is not unprecedented. Three examples have been documented: i) the CXC-chemokine CXCL12 binds to the alarmin HMGB1 and HMGB1/CXCL12 complex formation promotes chemotactic activity through CXCR4 (De Leo *et al.*, 2019; Schiraldi *et al.*, 2012); ii) the anti-microbial peptide and α-defensin HNP1 binds to CCL5 and enhances monocyte adhesion through CCR5 (Alard *et al*, 2015); iii) macrophage-expressed galectins such as galectin-3 (Gal-3) bind to CXCL12 and attenuate CXCL12-stimulated signaling via CXCR4 (Eckardt *et al.*, 2020). However, while these studies underscore that classical chemokine activity may be modulated by interaction with various soluble mediators, HMGB1 and Gal-3 have no chemotactic activity on their own; HNP1 has been reported to exhibit chemoattractive properties, but the mediating chemoattractant receptor has remained elusive. In contrast, despite lacking the signature structural elements of classical chemokines such as the chemokine-fold and the N-terminal cysteine motif, MIF is a chemoattractant and depending on the microenvironmental context, can signal through the CXC chemokine receptors CXCR2, CXCR4, and/or ACKR3 to promote atherogenic and inflammatory leukocyte recruitment. Its CXC receptor binding capacity is based on the presence of a pseudo-ELR

motif and an extended N-like loop, structurally mimicking the site 1 and 2 receptor binding

motifs of the corresponding cognate ligands CXCL1/8 and CXCL12, respectively. Together

with the β-defensins HDB1/2 and HBD3, which bind to CCR6 and CXCR4, respectively, and

secreted fragments of certain AARSs, which bind to CXCR1 and CXCR2, MIF has therefore

been designated an ACK (Degryse & de Virgilio, 2003; Kapurniotu *et al.*, 2019; Oppenheim &

Yang, 2005; Rohrl *et al.*, 2010; Sinitski *et al.*, 2019; Wakasugi & Schimmel, 1999). Our

current identification of MIF/CXCL4L1 heterocomplexes thus also shows that the chemokine

interactome is not strictly limited to interactions between classical CC- and/or CXC-type

chemokines, but also encompasses heteromeric interactions between classical and atypical

chemokines, with potential functional modulation of the chemokine receptor pathway of both

the classical or atypical chemokine. Although not further validated and pursued in our current

study, the detection of additional candidate interactors of MIF in our performed unbiased

chemokine array, i.e. CCL28, CXCL9, as well as Prx6 leads us to hypothesize that

interactions between classical and atypical chemokines could represent a broader principal

of an "expanded ACK/CK interactome".

The validity of the solid phase chemokine array as an unbiased screening approach

for candidate chemokine interactors has been previously established (von Hundelshausen *et*

al., 2017). The general utility and specificity of this methodology was further confirmed in the

current study. Out of 47 immobilized classical chemokines, in addition to CXCL4L1, only two

other classical chemokines, i.e. CCL28 and CXCL9, were revealed to have positivity. While a

functional link between MIF and CCL28 has yet to be unveiled, it is interesting to note that

the other detected CXC chemokine was CXCL9, a CXCR3 agonist like CXCL4L1.

Intriguingly, biotin-MIF neither bound to CXCL12 nor to CXCL8, indicating that implicated

functional interactions between MIF and the cognate CXCR4 and CXCR2 ligands,

respectively, are independent of heterocomplex formation.

Futhermore, the specificity of the performed array is underscored by the notion that

CXCL4, the highly homologous sister variant of CXCL4L1, did not bind to MIF, both at pH 8

and also when we tested for this interaction at pH 6 (data not shown) to account for pH-

dependent charge differences. We hypothesize that the striking difference between CXCL4L1 and CXCL4 in binding to MIF might be due to the suggested different conformation of these two chemokines, e.g. the more exposed and flexible α-helix of monomeric CXCL4L1 (Kuo *et al.*, 2013). While CXCL4 has been amply characterized by us and others as a pro-atherogenic platelet chemokine, in part also via its intriguing capacity to hetero-oligomerize with CCL5 (Koenen *et al.*, 2009; von Hundelshausen *et al.*, 2007), very little is known about the role of CXCL4L1 in chronic inflammatory diseases and atherosclerosis. Like its sister molecule, CXCL4L1 is also abundantly expressed in platelets; however, it apparently is not localized in α-granules but resides in a different sub-cellular compartment, from where it is constitutively secreted (Lasagni *et al*, 2007). It is also found in other cell types including mononuclear cells and smooth muscle cells (Lasagni *et al.*, 2007). CXCL4L1 serves as an inhibitor of angiogenesis and has pro-inflammatory effects by inducing the release of CCL2 and CXCL8 from monocytes, while – contrary to CXCL4 – it does not promote monocyte survival (Domschke & Gleissner, 2019; Gouwy *et al*, 2016; Sarabi *et al*, 2011). There is only one *in vivo* study, in which CXCL4L1 was investigated as prognostic marker in cardiovascular disease. Interestingly, below-median levels of CXCL4L1 were found to correlate with a worse outcome in stable coronary artery disease patients, as indicated by a higher rate of cardiac death, stroke, or myocardial infarction (De Sutter *et al*, 2012). This finding might argue for a beneficial role of this chemokine in cardiovascular disease, even though the mechanisms behind this remain unclear, but certainly more studies are required. Of note, there is no equivalent of CXCL4L1 in mice (Eisman *et al*, 1990), limiting functional *in vivo* studies of this chemokine and its complex with MIF, as predicted from our study.

Importantly, we validated the binding between MIF and CXCL4L1 by semi-endo-genous pulldown from monocytes, as well as two different biophysical *in vitro* methods, i.e. SPR and MST. The combination of both methods also addresses potential disadvantages of having one interaction partner immobilized (Zhou *et al*, 2016). The binding affinity constants derived from the SPR and MST experiments (116 and 160 nM, respectively) are in reasonable agreement with each other. The observed (small) difference could be due to a

number of factors, including surface immobilization effects, fluorescence *versus* biotin labeling, or buffers employed. Together, the results are suggestive of a relatively high binding affinity between MIF and CXCL4L1. Moreover, the obtained nanomolar K_D is consistent with the reported concentrations of both proteins in inflammatory disease settings (Sinitski *et al.*, 2019). Flanking evidence for MIF/CXCL4L1 complex formation was obtained by our peptide array mapping and molecular docking results. As expected given their high sequence identity, the peptide array predicted identical binding sites for CXCL4 and CXCL4L1. Also, the peptide array methodology interrogates linear binding epitopes but cannot delineate conformational differences. In fact, Kuo *et al.* suggested that the three-amino acid difference between CXCL4 and CXCL4L1, although marginal, leads to a slight tilting of the C-terminal α-helix (Kuo *et al.*, 2013). We hypothesize that this moderate conformational change could be the basis for the observed preferred binding of MIF to CXCL4L1 compared to CXCL4. Differences in their binding affinity to CCL5 have already been reported for CXCL4 and CXCL4L1 and also the availability of their monomers, regulated by the stability of their tetrameric complexes, differs between these two chemokines (Sarabi *et al.*, 2011). Future structural studies, e.g. by nuclear magnetic resonance (NMR) spectroscopy, may help to further address these and other conformational questions.

To investigate the functional consequences of MIF/CXCL4L1 heterocomplex formation, we focused on inflammatory and atherosclerosis-relevant activities of MIF. T-cell migration is one such activity that is regulated by the MIF/CXCR4 pathway (Bernhagen *et al.*, 2007). In line with previous results, MIF promoted T-cell migration in a physiologically relevant 3D migration setting. Although T cells generally express the CXCL4L1 receptor CXCR3, CXCL4L1 alone had no effect on the chemotaxis of human PBMC-derived T cells. Lack of CXCL4L1 activity in this assay is likely due to the fact that CXCL4L1 is not a *bona fide* T-cell chemoattractant (Gouwy *et al.*, 2016) and that the preferential CXCL4L1 receptor variant CXCR3B is poorly expressed on T cells (Korniejewska *et al*, 2011). The 3D T-cell migration data are supported by the result that MIF, but not the combination of MIF and CXCL4L1, promoted Jurkat T-cell migration in a 2D Transwell assay. Confirming the

remarkable specificity of MIF binding to CXCL4L1 *versus* CXCL4, coincubation of MIF with

CXCL4 did not result in reduced Jurkat T-cell migration. Of note, heterocomplex formation

with MIF led to a complete blockade of MIF's pro-migratory effect on primary T cells in the 3D

migration setting. While *in vivo* T-cell recruitment studies were beyond the scope of our

study, inhibition of MIF-mediated T-cell migration by CXCL4L1 complexation could potentially

be relevant in atherosclerosis, where it might represent a feedback mechanism that could

serve to dampen the atherogenic response. In fact, abundant CXCL4L1 levels may be

released by activated platelets in an atherogenic microenvironment, where they could

colocalize with endothelial-immobilized or monocyte-secreted MIF and infiltrating T cells.

That complexation of MIF by CXCL4L1 can interfere with MIF's chemoattractant activities

was confirmed in a microglia assay, in which the motility of Egfp$^+$ microglia in murine cortical

brain cultures *ex vivo* was studied. In addition to representing an independent cell migration

system, the data obtained from the microglia-containing cortical cultures further confirmed

that complex formation interferes with MIF signaling through the CXCR4 pathway and

underscored that the mechanism could be relevant in *in vivo*-like physiological tissue

settings. That MIF/CXCL4L1 heterocomplex formation interferes with MIF signaling through

CXCR4 was independently validated by biochemical experiments using FP spectroscopy and

DMR analysis of HEK293-CXCR4 transfectants.

This identified interaction of MIF with CXCL4L1, supposedly resulting in local

inhibition of MIF's pro-inflammatory effects was especially interesting to us in the context of

previous studies, in which we identified human and mouse platelets as an abundant source

of MIF (Strüßmann *et al.*, 2013; Wirtz *et al.*, 2015). Here, we verified expression and

localization of MIF in human platelets as well as in platelet-rich clinical thrombus tissue by

confocal (CLSM) and multiphoton microscopy (MPM). As expected, these experiments also

showed the abundant presence of CXCL4L1 in platelets and thrombi, and suggested the

colocalization and/or complex formation of MIF and CXCL4L1 in the vicinity of platelets. Due

to the optical resolution limits of the CLSM and MPM methods, true colocalization and the

specific subcellular compartment could not be determined. Evidence for the presence of

MIF/CXCL4L1 heteromers is suggested by PLA performed on cryosections of a human

thrombus. In fact, PLA is an established method to detect CK heteromers as shown

previously for HNP1/CCL5 complexes (Alard *et al.*, 2015).

Having confirmed the occurrence of this novel complex in platelet preparations and

thrombus tissue, lastly the effect of MIF, CXCL4L1 and their complex on platelet function and

thrombus formation was assessed. MIF promoted thrombus formation leading to a larger

thrombus-covered area in an *in vitro* setting under flow conditions. Confirming our previous

results, this effect was abrogated upon co-incubation with CXCL4L1. It is interesting to note

that in the settings used in our experiment applying a shear rate of 1000 s^{-1} for 5 min, MIF

acted to enhance thrombus formation. Instead, in a previous study employing a shear rate of

1700 s^{-1} MIF was found to reduce thrombus size, confirming that MIF is a modulator of

thrombus formation, but also indicating that the directionality of the effect may depend on the

specific microenvironmental context.

Moreover, studying the morphology change of isolated platelets during adhesion and

activation on a fibrinogen-coated surface revealed that both MIF and CXCL4L1 favored a

switch from large to small lamellipodia at an early time point. Interestingly, in this setting no

inhibition by the complex on MIF-mediated effects was observed, but a synergistic behavior

of MIF and CXCL4L1 was observed, suggesting that this effect may occur independently of

CXCR4.

In addition to their classical role in wound closure and haemostasis, thrombus formation

and platelet activation are processes that are closely linked to inflammatory processes

driving atherosclerosis (Gawaz, 2006; Lippi *et al*, 2011; Nording *et al*, 2020; von

Hundelshausen & Weber, 2007). MIF has been amply linked to atherosclerotic pathogenesis

both clinically and experimentally, with evidence for a number of contributing mechanisms

including leukocyte recruitment and platelet activation (Bernhagen *et al.*, 2007; Chatterjee *et*

al., 2014; Muller *et al*, 2013; Sinitski *et al.*, 2019; Zernecke *et al*, 2008) The identified

heteromerization of MIF and CXCL4L1 in our current study and the observed effect of

MIF/CXCL4L1 complex formation on immune cell migration as well as thrombus size and

platelet morphology might imply that CXCL4L1 could have a protective role in atherosclerosis by mitigating the pro-atherosclerotic effects of MIF via complex formation. This hypothesis would warrant future studies in corresponding experimental *in vivo* models, albeit the lack of CXCL4L1 expression in rodents will impose a particular challenge here.

In summary, we provide evidence that MIF does not only behave as a chemokine-like mediator by way of engaging classical chemokine receptors but also by direct binding to classical chemokines. Interestingly, the identified chemokine interactor of MIF is not one of the cognate ligands of the MIF receptors CXCR2 or CXCR4, but CXCL4L1, a prominent platelet chemokine not previously implicated in MIF biology or MIF-mediated pathologies. While evidence from experimental *in vivo* disease models will have to be obtained in future studies, our data suggest that MIF/CXCL4L1 complex formation could serve to attenuate inflammatory/atherogenic activities of MIF through the CXCR4 receptor axis. Our study also gives insight into the growing "chemokine interactome" with a particular focus on ACKs. While modulatory effects on the interactome by mediators not belonging to the class of chemokines have already been exemplified by intriguing studies involving HMGB1, HNP1, and the galectins (Alard *et al.*, 2015; Eckardt *et al.*, 2020; Schiraldi *et al.*, 2012), the current study is first in demonstrating a role for MIF family proteins in particular, and *bona fide* ACKs in general, as defined by their chemotactic activity mediated through engagement of classical chemokine receptors. While not yet validated by follow up analyses, the identification of additional potential interactors in our array indicates that this could represent a broader principle of an ACK/CK interactome.

Acknowledgements

This work was supported by Deutsche Forschungsgemeinschaft (DFG) grant SFB1123-A3 to J.B. and A.K., DFG INST 409/209-1 FUGG to J.B., SFB1123-A2 to P.v.H., SFB1123-A1 to C.W., SFB1123-B5 to L.M., SFB240-B01 (374031971 – TRR 240) to M.G., and by DFG under Germany's Excellence Strategy within the framework of the Munich Cluster for Systems Neurology (EXC 2145 SyNergy—ID 390857198) to J.B., C.W., and O.G. A.H. was supported by a Metiphys scholarship of LMU Munich and by the Friedrich-Baur-Foundation e.V. at LMU University Hospital. E.S. and J.B. received funding from the LMU-FöFoLe program under project 52114016. C.W. is Van de Laar Professor of Atherosclerosis. We thank Simona Gerra, Priscila Bourilhon, and Lusine Saroyan for technical support, and Christine Krammer for help with the DMR assay. We are grateful to the mouse core facility of the Center for Stroke and Dementia Research (CSD) for their support. We also thank the Biophysics Core Facility at the School of Biology of LMU Munich and Dr. Sophie Brameyer for usage of the MST instrument.

 Figure legends

Figure 1: Unbiased chemokine protein array identifies CXCL4L1, but not CXCL4, as a novel

interaction candidate of MIF. (**A**) Schematic illustrating binding of biotinylated MIF to the

chemokine protein array. (**B**) Layout of the immobilized chemokines, atypical chemokines

and alarmins (*top*) and membrane of chemokine solid phase assay performed at pH 8.0,

developed against bound biotin-MIF (*bottom*). (**C**) Close-up of the membrane with a focus on

CXCL4 and CXCL4L1 with the corresponding negative control membrane, incubated without

biotin-MIF.

Figure 2: Validation of MIF/CXCL4L1 complex formation by a variety of protein-protein

interaction assays and verification of the specificity of MIF complexation with CXCL4L1 over

CXCL4*.* (**A**) Semi-endogenous pull-down assay, in which endogenous CXCL4L1 from

MonoMac6 lysates was captured with recombinant biotinylated MIF and pulled down by

streptavidin-coated paramagnetic beads. Blots, developed against MIF (*left*) and CXCL4L1

(*right*), show representative results of three independent experiments. Input corresponds to

5% cell lysate without pull-down and control (Ctrl) refers to pull-downs performed in the

absence of biotin-MIF. Molecular weight markers were lelectrophoresed in the same gel and

relevant marker sizes are indicated. (**B**) Interrogation of MIF/CXCL4L1 complex formation by

surface plasmon resonance (SPR) spectroscopy using chip-immobilized biotin-MIF titred

against increasing concentrations of CXCL4L1. Measurements indicate an interaction

between MIF and CXCL4L1 with an estimated K_D of 116 ± 16 nM. The SPR response signal

is given in relative units (RU). (**C**) Same as (**B**), except that titration was performed with

CXCL4. Corresponding SPR spectroscopy data for MIF and CXCL4. No detectable binding

signal was obtained and no K_D could be derived. (**D**) Interrogation of MIF/CXCL4L1 complex

formation by microscale thermophoresis (MST) utilizing fluorescently labeled MIF and

CXCL4L1 in solution. MST analysis revealed a K_D of 159.8 ± 16.8 nM for the interaction of

MIF and CXCL4L1. (**E**) Same as (**D**), except that CXCL4 was tested. The derived apparent

K_D of 2.0 ± 0.8 µM was ten-fold higher compared to MIF/CXCL4L1.

Figure 3. Co-incubation with CXCL4L1 inhibits MIF-mediated immune cell chemotaxis. (**A**)

Migration of human CD4$^+$ T-cells embedded in a gel matrix, subjected to gradients of MIF,

CXCL4L1 or both. Movement of cells was followed by live cell imaging and individual tracks

reconstructed from acquired images. Tracks of cells migrating towards the indicated stimuli

are marked in the corresponding color. Starting point was centered to x = y = 0. The black

crosshair indicates the cell population's center of mass after migration. (**B**) Quantification of

the 3D chemotaxis experiment in (**A**), indicating that complexation of MIF by CXCL4L1

attenuates MIF-mediated directed migration of human CD4$^+$ T-cells. Plotted is the calculated

forward migration index (FMI), based on manual tracking of at least 30 individual cells per

treatment. (**C**) Migration trajectories of murine microglia, obtained by live cell imaging for 15

h, treated with MIF, CXCL4L1, or both. Used concentrations: MIF: 8 nM, CXCL4L1: 1.6 nM;

n=5 independent experiments; horizontal bar: 100 µm. (**D**) Analysis of microglia motility,

based on each tracked cell accumulated distance, shown in (**C**). Data is presented as mean

± SD. Statistical significance is indicated as described: *, P < 0.05; **, P < 0.01; ***, P <

0.001.

Figure 4. MIF/CXCL4L1 complex formation inhibits binding of MIF to CXCR4 and signaling

of MIF through the CXCR4 signaling axis. (**A**) Fluorescence polarization (FP) spectroscopy

shows the interaction of Alexa488-labeled MIF with the soluble CXCR4 receptor mimic

msR4M-L1 with an apparent K_D of 237.2 ± 24.2 nM. Data is presented as mean of 3

independent experiments; error bars represent the SD. (**B**) Pre-incubation of MIF with

CXCL4L1 (160-fold molar excess) prevents the interaction of MIF with msR4M-L1 (app. K_D >

10 µM). Mean of 3 experiments ± SD. (**C**) Dynamic mass redistribution (DMR)

measurements with HEK293 cells stably expressing CXCR4 indicate that the cellular

response to MIF is reduced, when MIF is pre-incubated with CXCL4L1. The DMR response

of CXCR4-expressing HEK293 cells to MIF in the presence or absence of the CXCR4-

antagonist AMD3100 is also shown, confirming the CXCR4-dependency of the cellular

response to MIF.

Figure 5. Co-localization and interaction of MIF and CXCL4L1 in human platelet

preparations, detected in multiphoton microscopy (MPM). (**A**): MPM images of isolated

platelets, forming small aggregates, stained for MIF and CXCL4L1. White arrowheads

indicate areas of colocalization. Size bar: 5 µm. (**B**) MPM images of isolated, more separated

platelets, stained as in (**A**), showing colocalization of MIF and CXCL4L1. Size bar: 5 µm. (**C**)

Fluorescence lifetime imaging (FLIM) of platelets isolation as shown in (**B**). Color-code

corresponds to lifetime of the donor, Alexa 488, the dye used for the antibody-based staining

of MIF. (**D**) Histogram of the Förster Resonance Energy Transfer (FRET) efficiency in (**C**).

(**E**) Donor lifetime shortening, presented as the mean lifetime (τ), average weighted, of the

donor (Alexa 488, MIF staining) alone, and in combination with the acceptor fluorophore

(Cy3, CXCL4L1 staining), where FRET occured.

Figure 6. Proximity ligation assay (PLA) indicates that MIF/CXCL4L1 heterocomplexes are

present in human thrombus tissue. (**A**) MIF/CXCL4L1 complex formation in thrombus

specimen revealed by PLA. PLA-positive signals are depicted in yellow; tissue was

counterstained with fluorescent-labeled phalloidin (cyan). Stained tissue samples were

imaged by CLSM; size bar: 50 µm. (**B**) HE staining of thrombus tissue specimen; size bar: 75

µm.

Figure 7. (**A**) Thrombus formation in human blood under flow stress is enhanced by MIF,

and this effect is diminished by pre-incubation of MIF with CXCL4L1. Fluorescent staining

with $DiOC_6$. Shown are representative images of one experiment, performed at a shear rate

of 1000 s^{-1}; size bar: 100 µm. (**B**) Quantification of thrombi sizes from flow chamber

experiments, as depicted exemplarily in (**A**). MIF-mediated increase in thrombus-covered

area is diminished, when MIF is pre-incubated with CXCL4L1. n = 6 experiments and

platelets coming from 4 donors. (**C**) Quantification of total thrombi numbers per treatment

group. As thrombus numbers remain unchanged, effects on thrombus-covered area originate

from the size of the formed thrombi (see also Supplementary Figure 5); n = 6 experiments.

(**D**) Analysis and quantification of platelet morphology upon adhesion on fibrinogen-coated

coverslips. Activated platelets were allowed to adhere on fibrinogen-coated coverslips that

were pre-treated with MIF, CXCL4L1 or a mixture of both for the indicated times. After fixing

with PFA, images of randomly selected areas were taken and platelet morphology analyzed.

Treatment with a combination of MIF and CXCL4L1 led to a reduction in the large

lamellopodia phenotype, favoring small lamellopodia, with the MIF/CXCL4L1 complex

showing a stronger effect then the individual proteins; n = 6 experiments. (**E**) Platelet

morphology distribution after 30 min for each treatment group according to panel (**D**).

Figure 8: Summary scheme and suggested model of CXCL4L1/MIF complex formation and

functions. The atypical chemokine MIF and the classical chemokine CXCL4L1, e.g. present

in an inflammatory or atherogenic microenvironment after release from platelets, form

heteromeric complexes. Complexes inhibit inflammatory effects of MIF on leukocyte

recruitment as well as its pro-thrombotic effects through impairing MIF interactions with its

non-cognate receptor CXCR4.

Supplementary figure legends

Supplementary Figure 1. Additional data for chemokine protein array. (**A**) Negative control

membrane related to the experiment in **Figure 1**, incubated in buffer at pH 8.0 without biotin-

MIF. (**B**) Close-up of membrane from a chemokine protein array experiment with a focus on

CXCL4 and CXCL4L1. The membrane was incubated with biotin-MIF and the incubation was

performed at pH 6.0.

Supplementary Figure 2. Investigation of the MIF/CXCL4L1 interaction interface and *in*

silico studies. (**A**) CelluSpot peptide array experiments, where overlapping peptides of

CXCL4 (*left*) and CXCL4L1 (*right*) were spotted on an array and probed with biotin-MIF.

Chemiluminescence signal intensity indicates binding of biotin-MIF to the respective peptide.

Arrows indicate peptides of interest that are most likely to be involved in the interaction with

MIF. (**B**) Sequences of peptides identified in **A** are highlighted in the 3D structure of

monomeric CXCL4 and CXCL4L1, showing their localization in the folded proteins. For both

chemokines, these peptides of interest represent almost identical amino acid sequences,

corresponding to highly similar regions of the protein. This indicates that not only the

sequence but also the three-dimensional conformation of the chemokines might play a role in

the interaction with MIF. Amino acid residues, in which CXCL4L1 differs from CXCL4 are in

italics. PyMOL was used to visualize a CXCL4 (PDB ID: 1F9Q Chain A) and CXCL4L1

monomer (PDB ID: 4HSV Chain A). (**C**) To visualize the proposed MIF/CXCL4L1 complex,

an unbiased *in silico* protein-protein docking approach was taken. The ClusPro 2.0

webserver was used to simulate a complex consisting of both a MIF and CXCL4L1

monomer. Depicted here is the highest-ranking docking result, with peptides identified in **A** to

be potentially part of the interaction interface highlighted in CXCL4L1. According to this *in*

silico prediction, they are partially directed towards MIF, allowing parts of their sequences

being involved in complex formation. PyMOL was used to calculate the surface charge

distribution of these proteins (red: negatively charged; blue: positively charged), revealing an

area of opposite charges in the proposed contact region of MIF and CXCL4L1 that partially

matches the peptide array results.

Supplementary Figure 3. Effects on cell migration in Jurkat T cells and microglia. (**A**) Effect

of CXCL4L1 on MIF-mediated chemotaxis of Jurkat T cells as analyzed in a Transwell

migration assay. Used concentrations: MIF: 16 nM, CXCL4L1: 32 nM; Data is presented as

mean ± SD. n = 2-4 independent experiments. (**B**) Same as (**A**), except that co-incubation

with CXCL4 was analyzed. Data is presented as mean ± SD. n = 4 independent experiments

with duplicates each. (**C**) Quantification of murine microglia motility, based on the

accumulated distance of GFP-positive microglia tracked during live cell imaging (n = 5). MIF

was used at a concentration of 8 nM, the soluble CXCR4-mimicking peptide msR4M-L1 at 40

nM and the cognate ligand of CXCR4, CXCL12, at 16 nM. Data presented as mean ± SD.

Statistical significance: *, $P < 0.05$; **, $P < 0.01$; ***, $P < 0.005$; ****, $P < 0.0001$.

Supplementary Figure 4. Quantification of mean thrombus sizes from **Figure 5A**, showing a

trend for CXCL4L1 inhibiting the MIF-mediated increase in thrombus size in samples, in

which MIF and CXCL4L1 were pre-incubated together.

References

Alard JE, Ortega-Gomez A, Wichapong K, Bongiovanni D, Horckmans M, Megens RT, Leoni G, Ferraro B, Rossaint J, Paulin N *et al* (2015) Recruitment of classical monocytes can be inhibited by disturbing heteromers of neutrophil HNP1 and platelet CCL5. *Sci Transl Med* 7: 317ra196

Bachelerie F, Ben-Baruch A, Burkhardt AM, Combadiere C, Farber JM, Graham GJ, Horuk R, Sparre-Ulrich AH, Locati M, Luster AD *et al* (2014a) International Union of Basic and Clinical Pharmacology. LXXXIX. Update on the extended family of chemokine receptors and introducing a new nomenclature for atypical chemokine receptors. *Pharmacol Rev* 66: 1-79

Bachelerie F, Graham GJ, Locati M, Mantovani A, Murphy PM, Nibbs R, Rot A, Sozzani S, Thelen M (2014b) New nomenclature for atypical chemokine receptors. *Nat Immunol* 15: 207-208

Bai F, Asojo OA, Cirillo P, Ciustea M, Ledizet M, Aristoff PA, Leng L, Koski RA, Powell TJ, Bucala R *et al* (2012) A novel allosteric inhibitor of macrophage migration inhibitory factor (MIF). *J Biol Chem* 287: 30653-30663

Bernhagen J, Krohn R, Lue H, Gregory JL, Zernecke A, Koenen RR, Dewor M, Georgiev I, Schober A, Leng L *et al* (2007) MIF is a noncognate ligand of CXC chemokine receptors in inflammatory and atherogenic cell recruitment. *Nat Med* 13: 587-596

Bernhagen J, Mitchell RA, Calandra T, Voelter W, Cerami A, Bucala R (1994) Purification, bioactivity, and secondary structure analysis of mouse and human macrophage migration Inhibitory factor (MIF). *Biochemistry* 33: 14144-14155

Borst O, Schmidt EM, Munzer P, Schonberger T, Towhid ST, Elvers M, Leibrock C, Schmid E, Eylenstein A, Kuhl D *et al* (2012) The serum- and glucocorticoid-inducible kinase 1 (SGK1) influences platelet calcium signaling and function by regulation of Orai1 expression in megakaryocytes. *Blood* 119: 251-261

Calandra T, Roger T (2003) Macrophage migration inhibitory factor: a regulator of innate immunity. *Nat Rev Immunol* 3: 791-800

Charo IF, Ransohoff RM (2006) The many roles of chemokines and chemokine receptors in inflammation. *N Engl J Med* 354: 610-621

Chatterjee M, Borst O, Walker B, Fotinos A, Vogel S, Seizer P, Mack A, Alampour-Rajabi S, Rath D, Geisler T *et al* (2014) Macrophage migration inhibitory factor limits activation-induced apoptosis of platelets via CXCR7-dependent Akt signaling. *Circ Res* 115: 939-949

David JR (1966) Delayed hypersensitivity in vitro: its mediation by cell-free substances formed by lymphoid cell-antigen interaction. *Proc Natl Acad Sci U S A* 56: 72-77

De Leo F, Quilici G, Tirone M, De Marchis F, Mannella V, Zucchelli C, Preti A, Gori A, Casalgrandi M, Mezzapelle R *et al* (2019) Diflunisal targets the HMGB1/CXCL12 heterocomplex and blocks immune cell recruitment. *EMBO Rep* 20: e47788

De Sutter J, Van de Veire NR, Struyf S, Philippe J, De Buyzere M, Van Damme J (2012) PF-4var/CXCL4L1 predicts outcome in stable coronary artery disease patients with preserved left ventricular function. *PLoS One* 7: e31343

Degryse B, de Virgilio M (2003) The nuclear protein HMGB1, a new kind of chemokine? *FEBS Lett* 553: 11-17

Domschke G, Gleissner CA (2019) CXCL4-induced macrophages in human atherosclerosis. *Cytokine* 122: 154141

Eckardt V, Miller MC, Blanchet X, Duan R, Leberzammer J, Duchene J, Soehnlein O, Megens RT, Ludwig AK, Dregni A *et al* (2020) Chemokines and galectins form heterodimers to modulate inflammation. *EMBO Rep* 21: e47852

Eiger DS, Boldizsar N, Honeycutt CC, Gardner J, Rajagopal S (2021) Biased agonism at chemokine receptors. *Cell Signal* 78: 109862

Eisman R, Surrey S, Ramachandran B, Schwartz E, Poncz M (1990) Structural and

functional comparison of the genes for human platelet factor 4 and PF4alt. *Blood* 76: 336-

344

Fotinos A, Klier M, Gowert NS, Munzer P, Klatt C, Beck S, Borst O, Billuart P, Schaller M,

Lang F *et al* (2015) Loss of oligophrenin1 leads to uncontrolled Rho activation and increased

thrombus formation in mice. *J Thromb Haemost* 13: 619-630

Gawaz M (2006) Platelets in the onset of atherosclerosis. *Blood Cells Mol Dis* 36: 206-210

Gokce O, Sudhof TC (2013) Membrane-tethered monomeric neurexin LNS-domain triggers

synapse formation. *J Neurosci* 33: 14617-14628

Gouwy M, Ruytinx P, Radice E, Claudi F, Van Raemdonck K, Bonecchi R, Locati M, Struyf S

(2016) CXCL4 and CXCL4L1 Differentially affect monocyte survival and dendritic cell

differentiation and phagocytosis. *PLoS One* 11: e0166006

Hutchings CJ, Koglin M, Olson WC, Marshall FH (2017) Opportunities for therapeutic

antibodies directed at G-protein-coupled receptors. *Nat Rev Drug Discov* 16: 787-810

Kapurniotu A, Gokce O, Bernhagen J (2019) The multitasking potential of alarmins and

atypical chemokines. *Front Med (Lausanne)* 6: 3

Karshovska E, Weber C, von Hundelshausen P (2013) Platelet chemokines in health and

disease. *Thromb Haemost* 110: 894-902

Klasen C, Ohl K, Sternkopf M, Shachar I, Schmitz C, Heussen N, Hobeika E, Levit-Zerdoun

E, Tenbrock K, Reth M *et al* (2014) MIF promotes B cell chemotaxis through the receptors

CXCR4 and CD74 and ZAP-70 signaling. *J Immunol* 192: 5273-5284

Kleist AB, Getschman AE, Ziarek JJ, Nevins AM, Gauthier PA, Chevigne A, Szpakowska M,

Volkman BF (2016) New paradigms in chemokine receptor signal transduction: Moving

beyond the two-site model. *Biochem Pharmacol* 114: 53-68

Koenen RR, von Hundelshausen P, Nesmelova IV, Zernecke A, Liehn EA, Sarabi A, Kramp BK, Piccinini AM, Paludan SR, Kowalska MA *et al* (2009) Disrupting functional interactions between platelet chemokines inhibits atherosclerosis in hyperlipidemic mice. *Nat Med* 15: 97-103

Kontos C, El Bounkari O, Krammer C, Sinitski D, Hille K, Zan C, Yan G, Wang S, Gao Y, Brandhofer M *et al* (2020) Designed CXCR4 mimic acts as a soluble chemokine receptor that blocks atherogenic inflammation by agonist-specific targeting. *Nat Commun* 11: 5981

Korniejewska A, McKnight AJ, Johnson Z, Watson ML, Ward SG (2011) Expression and agonist responsiveness of CXCR3 variants in human T lymphocytes. *Immunology* 132: 503-515

Kozakov D, Hall DR, Xia B, Porter KA, Padhorny D, Yueh C, Beglov D, Vajda S (2017) The ClusPro web server for protein-protein docking. *Nat Protoc* 12: 255-278

Kraemer BF, Schmidt C, Urban B, Bigalke B, Schwanitz L, Koch M, Seizer P, Schaller M, Gawaz M, Lindemann S (2011a) High shear flow induces migration of adherent human platelets. *Platelets* 22: 415-421

Kraemer S, Lue H, Zernecke A, Kapurniotu A, Andreetto E, Frank R, Lennartz B, Weber C, Bernhagen J (2011b) MIF-chemokine receptor interactions in atherogenesis are dependent on an N-loop-based 2-site binding mechanism. *FASEB J* 25: 894-906

Krammer C, Kontos C, Dewor M, Hille K, Dalla Volta B, El Bounkari O, Tas K, Sinitski D, Brandhofer M, Megens RTA *et al* (2021) A MIF-Derived Cyclopeptide that Inhibits MIF Binding and Atherogenic Signaling via the Chemokine Receptor CXCR2. *Chembiochem* 22: 1012-1019

Kuo JH, Chen YP, Liu JS, Dubrac A, Quemener C, Prats H, Bikfalvi A, Wu WG, Sue SC (2013) Alternative C-terminal helix orientation alters chemokine function: structure of the anti-angiogenic chemokine, CXCL4L1. *J Biol Chem* 288: 13522-13533

Lacy M, Kontos C, Brandhofer M, Hille K, Groning S, Sinitski D, Bourilhon P, Rosenberg E, Krammer C, Thavayogarajah T *et al* (2018) Identification of an Arg-Leu-Arg tripeptide that contributes to the binding interface between the cytokine MIF and the chemokine receptor CXCR4. *Sci Rep* 8: 5171

Lasagni L, Grepin R, Mazzinghi B, Lazzeri E, Meini C, Sagrinati C, Liotta F, Frosali F, Ronconi E, Alain-Courtois N *et al* (2007) PF-4/CXCL4 and CXCL4L1 exhibit distinct subcellular localization and a differentially regulated mechanism of secretion. *Blood* 109: 4127-4134

Leng L, Metz CN, Fang Y, Xu J, Donnelly S, Baugh J, Delohery T, Chen Y, Mitchell RA, Bucala R (2003) MIF signal transduction initiated by binding to CD74. *J Exp Med* 197: 1467-1476

Lippi G, Franchini M, Targher G (2011) Arterial thrombus formation in cardiovascular disease. *Nat Rev Cardiol* 8: 502-512

Ma F, Kouzoukas DE, Meyer-Siegler KL, Hunt DE, Leng L, Bucala R, Vera PL (2017) Macrophage migration inhibitory factor mediates protease-activated receptor 4-induced bladder pain through urothelial high mobility group box 1. *Physiol Rep* 5: e13549.

Michelet C, Danchin EGJ, Jaouannet M, Bernhagen J, Panstruga R, Kogel KH, Keller H, Coustau C (2019) Cross-Kingdom Analysis of Diversity, Evolutionary History, and Site Selection within the Eukaryotic Macrophage Migration Inhibitory Factor Superfamily. *Genes (Basel)* 10: 740

Moerke NJ (2009) Fluorescence Polarization (FP) Assays for Monitoring Peptide-Protein or Nucleic Acid-Protein Binding. *Curr Protoc Chem Biol* 1: 1-15

Muller I, Schonberger T, Schneider M, Borst O, Ziegler M, Seizer P, Leder C, Muller K, Lang M, Appenzeller F *et al* (2013) Gremlin-1 Is an inhibitor of macrophage migration inhibitory

factor and attenuates atherosclerotic plaque growth in ApoE-/- Mice. *J Biol Chem* 288: 31635-31645

Murphy PM, Baggiolini M, Charo IF, Hebert CA, Horuk R, Matsushima K, Miller LH, Oppenheim JJ, Power CA (2000) International union of pharmacology. XXII. Nomenclature for chemokine receptors. *Pharmacol Rev* 52: 145-176

Nibbs RJ, Graham GJ (2013) Immune regulation by atypical chemokine receptors. *Nat Rev Immunol* 13: 815-829

Niess JH, Brand S, Gu X, Landsman L, Jung S, McCormick BA, Vyas JM, Boes M, Ploegh HL, Fox JG *et al* (2005) CX3CR1-mediated dendritic cell access to the intestinal lumen and bacterial clearance. *Science* 307: 254-258

Noels H, Weber C, Koenen RR (2019) Chemokines as therapeutic targets in cardiovascular disease. *Arterioscler Thromb Vasc Biol* 39: 583-592

Nording H, Baron L, Langer HF (2020) Platelets as therapeutic targets to prevent atherosclerosis. *Atherosclerosis* 307: 97-108

Oppenheim JJ, Yang D (2005) Alarmins: chemotactic activators of immune responses. *Curr Opin Immunol* 17: 359-365

Pawig L, Klasen C, Weber C, Bernhagen J, Noels H (2015) Diversity and Inter-Connections in the CXCR4 Chemokine Receptor/Ligand Family: Molecular Perspectives. *Front Immunol* 6: 429

Polack B, Schved JF, Boneu B, Groupe d'Etude sur l'Hemostase et la T (2001) Preanalytical recommendations of the 'Groupe d'Etude sur l'Hemostase et la Thrombose' (GEHT) for venous blood testing in hemostasis laboratories. *Haemostasis* 31: 61-68

Rajasekaran D, Groning S, Schmitz C, Zierow S, Drucker N, Bakou M, Kohl K, Mertens A, Lue H, Weber C *et al* (2016) Macrophage Migration Inhibitory Factor-CXCR4 Receptor

Interactions: evidence for partial allosteric agonism in comparison with CXCL12 chemokine. *J Biol Chem* 291: 15881-15895

Rohrl J, Yang D, Oppenheim JJ, Hehlgans T (2010) Specific binding and chemotactic activity of mBD4 and its functional orthologue hBD2 to CCR6-expressing cells. *J Biol Chem* 285: 7028-7034

Sarabi A, Kramp BK, Drechsler M, Hackeng TM, Soehnlein O, Weber C, Koenen RR, Von Hundelshausen P (2011) CXCL4L1 inhibits angiogenesis and induces undirected endothelial cell migration without affecting endothelial cell proliferation and monocyte recruitment. *J Thromb Haemost* 9: 209-219

Schiraldi M, Raucci A, Munoz LM, Livoti E, Celona B, Venereau E, Apuzzo T, De Marchis F, Pedotti M, Bachi A *et al* (2012) HMGB1 promotes recruitment of inflammatory cells to damaged tissues by forming a complex with CXCL12 and signaling via CXCR4. *J Exp Med* 209: 551-563

Schroder R, Janssen N, Schmidt J, Kebig A, Merten N, Hennen S, Muller A, Blattermann S, Mohr-Andra M, Zahn S *et al* (2010) Deconvolution of complex G protein-coupled receptor signaling in live cells using dynamic mass redistribution measurements. *Nat Biotechnol* 28: 943-949

Schurr Y, Sperr A, Volz J, Beck S, Reil L, Kusch C, Eiring P, Bryson S, Sauer M, Nieswandt B *et al* (2019) Platelet lamellipodium formation is not required for thrombus formation and stability. *Blood* 134: 2318-2329

Sinitski D, Kontos C, Krammer C, Asare Y, Kapurniotu A, Bernhagen J (2019) Macrophage Migration Inhibitory Factor (MIF)-Based Therapeutic Concepts in Atherosclerosis and Inflammation. *Thromb Haemost* 119: 553-566

Steen A, Larsen O, Thiele S, Rosenkilde MM (2014) Biased and g protein-independent signaling of chemokine receptors. *Front Immunol* 5: 277

Strüßmann T, Tillmann S, Wirtz T, Bucala R, von Hundelshausen P, Bernhagen J (2013) Platelets are a previously unrecognised source of MIF. *Thromb Haemost* 110: 1004-1013

Struyf S, Salogni L, Burdick MD, Vandercappellen J, Gouwy M, Noppen S, Proost P, Opdenakker G, Parmentier M, Gerard C *et al* (2011) Angiostatic and chemotactic activities of the CXC chemokine CXCL4L1 (platelet factor-4 variant) are mediated by CXCR3. *Blood* 117: 480-488

Sun HW, Bernhagen J, Bucala R, Lolis E (1996) Crystal structure at 2.6-A resolution of human macrophage migration inhibitory factor. *Proc Natl Acad Sci U S A* 93: 5191-5196

Tillmann S, Bernhagen J, Noels H (2013) Arrest functions of the MIF ligand/receptor axes in atherogenesis. *Front Immunol* 4: 115

Tilstam PV, Qi D, Leng L, Young L, Bucala R (2017) MIF family cytokines in cardiovascular diseases and prospects for precision-based therapeutics. *Exp Opin Ther Targ* 21: 671-683

Vajda S, Yueh C, Beglov D, Bohnuud T, Mottarella SE, Xia B, Hall DR, Kozakov D (2017) New additions to the ClusPro server motivated by CAPRI. *Proteins* 85: 435-444

von Hundelshausen P, Agten SM, Eckardt V, Blanchet X, Schmitt MM, Ippel H, Neideck C, Bidzhekov K, Leberzammer J, Wichapong K *et al* (2017) Chemokine interactome mapping enables tailored intervention in acute and chronic inflammation. *Sci Transl Med* 9

von Hundelshausen P, Petersen F, Brandt E (2007) Platelet-derived chemokines in vascular biology. *Thromb Haemost* 97: 704-713

von Hundelshausen P, Weber C (2007) Platelets as immune cells: bridging inflammation and cardiovascular disease. *Circ Res* 100: 27-40

Wakasugi K, Schimmel P (1999) Two distinct cytokines released from a human aminoacyl-tRNA synthetase. *Science* 284: 147-151

Weber C, Kraemer S, Drechsler M, Lue H, Koenen RR, Kapurniotu A, Zernecke A,

Bernhagen J (2008) Structural determinants of MIF functions in CXCR2-mediated

inflammatory and atherogenic leukocyte recruitment. *Proc Natl Acad Sci U S A* 105: 16278-

16283

Weber C, Noels H (2011) Atherosclerosis: current pathogenesis and therapeutic options. *Nat*

Med 17: 1410-1422

Wirtz TH, Tillmann S, Strussmann T, Kraemer S, Heemskerk JW, Grottke O, Gawaz M, von

Hundelshausen P, Bernhagen J (2015) Platelet-derived MIF: a novel platelet chemokine with

distinct recruitment properties. *Atherosclerosis* 239: 1-10

Yan LM, Tatarek-Nossol M, Velkova A, Kazantzis A, Kapurniotu A (2006) Design of a mimic

of nonamyloidogenic and bioactive human islet amyloid polypeptide (IAPP) as nanomolar

affinity inhibitor of IAPP cytotoxic fibrillogenesis. *Proc Natl Acad Sci U S A* 103: 2046-2051

Zernecke A, Bernhagen J, Weber C (2008) Macrophage migration inhibitory factor in

cardiovascular disease. *Circulation* 117: 1594-1602

Zhou M, Li Q, Wang R (2016) Current Experimental Methods for Characterizing Protein-

Protein Interactions. *ChemMedChem* 11: 738-756

Ziegler-Heitbrock HW, Thiel E, Futterer A, Herzog V, Wirtz A, Riethmuller G (1988)

Establishment of a human cell line (Mono Mac 6) with characteristics of mature monocytes.

Int J Cancer 41: 456-461

Zlotnik A, Burkhardt AM, Homey B (2011) Homeostatic chemokine receptors and organ-

specific metastasis. *Nat Rev Immunol* 11: 597-606

A

Streptavidin-POD

Substrate

Biotinylated MIF

Protein array

B

Immobilized chemokines

CCL1	CCL2	CCL3	CCL3L1	CCL4	CCL4L1	CCL5	CCL7	CCL8	CCL11	CCL13	CCL14
CCL15	CCL16	CCL17	CCL18	CCL19	CCL20	CCL21	CCL22	CCL23	CCL24	CCL25	CCL26
CCL27	CCL28	XCL1	XCL2	CXCL1	CXCL2	CXCL3	CXCL4	CXCL4L1	CXCL5	CXCL6	CXCL7
CXCL8	CXCL9	CXCL10	CXCL11	CXCL12a	CXCL12b	CXCL13	CXCL14	CXCL16	CXCL17	CX$_3$CL1	
HMGB1	BD1	BD2	Prx1		Prx6	MIF	MIF-2				

Detection of bound Biotin-MIF

C

Immobilized chemokine		Probe
CXCL4	CXCL4L1	

Biotin-MIF

Control

Brandhofer*, Hoffmann* *et al.* Fig. 4

A

MPM - platelet aggregates

MIF | CXCL4L1 | Merge

B

MPM - isolated platelets

MIF | CXCL4L1 | Merge

C

Fluorescence lifetime imaging (FLIM)

τ [ns]

D

FRET Efficiency Histogram

Occurence [Events]

FRET efficiency [%]

E

Donor/Alexa488 Lifetime Shortening

FLIM:	τ = 2.019 ± 0.069 ns
FLIM FRET:	τ = 1.496 ± 0.033 ns

A

Proximity Ligation Assay

B

HE staining

CXCL4L1 + MIF

CXCL4L1

MIF

No primary AB

A

Negative control membrane

B

A

CXCL4 array probed with biotin-MIF

CXCL4L1 array probed with biotin-MIF

B

CXCL4 monomer (PDB ID: 1F9Q)

CXCL4L1 monomer (PDB ID: 4HSV)

Peptide G16

Peptide G24

CXCL4:   EAEEDGDL**QCLCVKTTSQVRPRH**ITSLEVIKA**GPHCPTAQLIATLKN**GRKICLDLQA*P*LYKKIIK*K*LLES

CXCL4L1: *A*EAEEDGDL**QCLCVKTTSQVRPRH**ITSLEVI**KAGPHCPTAQLIAT**LKNGRKICLDLQA*L*LYKKIIK*EH*LES

Peptide H23

Peptide I06

C

Protein-Protein Docking via ClusPro

CXCL4L1
(PDB ID: 4HSV:A)

MIF
(PDB ID: 3DJH:A)

Surface charge distribution

MIF
(PDB ID: 3DJH:A)

CXCL4L1
(PDB ID: 4HSV:A)

−100.000   100.000

Thrombus size