Timing is everything: event-related transcranial direct current stimulation improves motor adaptation

Authors: Matthew Weightman[1,3,4*], John-Stuart Brittain[2,4], Alison Hall[1,4], R. Chris Miall[2,3,4], Ned Jenkinson[1,3,4]

Affiliations:

[1] School of Sport, Exercise and Rehabilitation Sciences, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

[2] School of Psychology, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

[3] MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

[4] Centre for Human Brain Health, University of Birmingham, Edgbaston, Birmingham, B15 2TT, UK

*Corresponding Author:

Matthew Weightman
School of Sport, Exercise and Rehabilitation Sciences,
The University of Birmingham
Edgbaston
Birmingham
B15 2TT
Email: mcw423@adf.bham.ac.uk

Abstract

Background: There is a current discord between the foundational theories underpinning motor learning and how we currently apply transcranial direct current stimulation (TDCS): the former is dependent on tight coupling of events while the latter is conducted with very low temporal resolution.

Objective: Here we aimed to investigate the temporal specificity of stimulation by applying TDCS in short epochs, and coincidentally with movement, during a motor adaptation task.

Methods: Participants simultaneously adapted a reaching movement to two opposing velocity-dependent force-fields (clockwise and counter-clockwise), distinguished by a contextual leftward or rightward shift in the task display and cursor location respectively. Brief bouts (< 3 second) of event-related TDCS (er-TDCS) were applied over M1 or the cerebellum during movements for only one of these learning contexts.

Results: We show that when short duration stimulation is applied to the cerebellum and yoked to movement, only those reaching movements performed simultaneously with stimulation are selectively enhanced, whilst similar and interleaved movements are left unaffected. We found no evidence of improved adaptation following M1 er-TDCS, as participants displayed equivalent levels of error during both stimulated and unstimulated movements. Similarly, participants in the sham stimulation group adapted comparably during left and right-shift trials.

Conclusions: It is proposed that the coupling of cerebellar stimulation and movement influences timing-dependent (i.e., Hebbian-like) mechanisms of plasticity to facilitate enhanced learning in the stimulated context.

Keywords: TDCS; Motor Adaptation; Plasticity

Introduction

Coincident, time-dependent mechanisms of synaptic plasticity are the canonical basis of theories of motor learning [1, 2]. These 'Hebbian' mechanisms are ubiquitous throughout the mammalian brain, having been described in the hippocampus, cerebellum, and sensory-motor cortices [3–5], and are believed to underpin all forms of learning and memory. Yet, protocols for non-invasive brain stimulation intended to promote motor learning and rehabilitation - particularly transcranial direct current stimulation (TDCS) - largely ignore timing-dependent mechanisms.

TDCS is a non-invasive form of brain stimulation often used to induce plasticity in the motor system by modulating neural excitability. Studies measuring the physiological effects of TDCS using transcranial magnetic stimulation (TMS), electroencephalogram (EEG), functional MRI (fMRI), magnetic resonance spectroscopy (MRS) etc. have reported alterations in neural excitability that are dependent on the polarity of stimulation [6–11]. Changes in neuronal excitation have been shown to be almost instantaneous in terms of increases in firing rates [12–14] and motor evoked potentials [6, 15] following a brief application of TDCS and other forms of polarising currents. Despite this, most conventional studies apply TDCS for 15-20 minutes in a continuous stimulation period, prior to and/or during a motor task and these changes in neural excitability do not always reliably translate into modulation behaviour, including motor learning. One potential reason for these apparently inconsistent results may be the low temporal resolution of stimulation. If TDCS can instantaneously modulate neural activity, applying short duration epochs of TDCS temporally aligned with movement has the potential to specifically and

selectively enhance learning, by driving coincident mechanisms of plasticity in the

circuits of the brain that are active during the movement.

Here we demonstrate (and replicate) that brief epochs of stimulation, applied in

synchrony with movement, selectively enhanced motor adaptation whilst, importantly

leaving adaptation of the non-stimulated (yet interleaved) movements unaffected.

We believe this first proof-of-principle study using a novel stimulation protocol

harnesses mechanisms of Hebbian plasticity, resulting in the selective, transient,

potentiation of those neurobehavioural circuits that are active concomitant with

stimulation [16–18].

96   Materials & Methods

97   Participant Details

A total of seventy-eight participants (aged 18-32 years, mean = 20.6 ± 3.0 years; 39

male) gave written informed consent to take part in the study (approved by the

Science, Technology, Engineering and Mathematics Ethical Review Committee at

the University of Birmingham). All participants were right-handed, had normal or

corrected to normal vision and completed a safety screening questionnaire for TMS

and TDCS prior to beginning the session. For the main experiment sixty participants

were pseudo-randomised into one of three experimental groups: M1 er-TDCS group

(n = 20; mean age = 19.7 ± 0.8 years, 10 males), Cerebellar er-TDCS group (n = 20;

mean age = 19.8 ± 1.9 years, 11 males) or a Sham TDCS group (n = 20; mean age

= 19.5 ± 0.8 years, 9 males). A further eighteen naive participants (n = 18, mean age

= 23.9 ± 4.7 years, 9 males) were recruited for a secondary experimental group.

Participants in this secondary group also received cerebellar er-TDCS. Importantly,

data from the secondary group were collected after the conclusion of the main

experiment in order to replicate the main experimental finding. No a priori statistical

methods were used to determine sample size. Instead, our sample size was chosen

to be consistent with, or greater than similar existing literature [19-21].

Experimental Design

Participants were seated in an armless chair so they could comfortably reach and

manipulate the handle of a custom-built robotic manipulandum (vBOT; [22]) with their

right arm. The vBOT measured and stored the position and velocity of the handle at

1000Hz, only allowing movements in the horizontal plane. The visual display screen

(Mac Cinema HD Display) was reflected in a horizontal mirror (60 x 76 cm) to appear

as a virtual image co-planar with the manipulandum. The screen displayed two grey

circular markers (2 cm diameter) which represented the home position and the

target, located 20 cm and 10 cm away from the edge of the screen respectively. The

screen also displayed a white cursor (1 cm diameter) which showed the position of

the vBOT handle. The home and target markers were displaced 10cm left or right of

the screen midline, depending on the context of the trial, and the cursor was

displaced 10cm left or right of the vBOT handle (Figure 1a).

Participants were told that the aim of the task was to make fast movements from

the home marker to the target marker, so that the cursor moved in a straight line

between the two (10 cm movement). At the beginning of each trial the participant

entered the home position and were held there for 3 seconds by stiff spring forces on

the vBOT handle, before being allowed to move. The home marker then changed

from grey to blue indicating that participants were allowed to make their movement

and the holding forces were released. During the hold period participants were asked

to keep relaxed and not to 'pull' or 'lean' on the handle. To encourage fully

completed movements, participants were told that it was acceptable to overshoot the

target slightly, but were discouraged from making excessively large movements. If

the movement was accurate and hit the target marker it would flash yellow for 1

second indicating a successful trial. If the movement failed to hit the target or

deviated ± 2 cm from the midline at any point during the movement path the target

would flash red for 1 second, indicating an unsuccessful trial. Once the movement

was completed, the vBOT would actively guide the handle back to the home position

with a spring force, ready for the next trial to start. Each trial took roughly 5 seconds

from start to finish (3 second hold, 1 second movement, 1 second return). Vision of

the upper arm was blocked during the task using a curtain and all lights extinguished

prior to starting the task.

(Figure 1)

Behavioural Protocol

The behavioural task consisted of 600 trials and was split into three phases:

Baseline (100 trials), Adaptation (400 trials) and Washout (100 trials). On each trial,

in all three phases, the task display would either be presented with a 10 cm leftward

or rightward shift from the midline position. The order of this shift was pseudo-

randomised, so that there was an equal number of leftward and rightward shift trials

in each phase and no more than 3 consecutive trials with the same contextual shift.

The trial/shift order was the same for all participants. During baseline trials no forces

were imposed on the handle so participants could move between the home marker

and the target unperturbed (Figure 1b). For adaptation trials the leftward or rightward

contextual shift in the task display was consistently associated with either a

161   clockwise (CW, for left-shift) or counter-clockwise (CCW, for right-shift) velocity

162   sensitive curl force-field (Figure 1c) - thus creating two distinct trial contexts [23]. The

163   strength of both imposed force-fields was 12 N/m/s (see equation 1).

165
$$\begin{bmatrix} Fx \\ Fy \end{bmatrix} = \begin{bmatrix} 0 & -12 \\ 12 & 0 \end{bmatrix} \begin{bmatrix} Vx \\ vy \end{bmatrix} \qquad (1)$$

Washout trials immediately followed the adaptation phase, which were once again

performed without any forces i.e. identical to baseline. Participants were not given

any explicit information regarding the link between the visual shift and associated

direction of force-field.

In order to measure compensatory forces applied against the curl field, 60 error-

clamp trials were pseudorandomly interleaved throughout the task (10% of all trials).

These trials were distributed proportionally throughout the three task phases. During

error- clamp trials, movements were constrained to a 'virtual channel' between the

home position and target marker, so that forces produced against the channel walls

could be measured (see Supplementary Figures 2 & 3 for data and analysis). Error-

clamp trials took place twice every 20 trials and the contextual shift was

pseudorandomly ordered so that no more than two successive error-clamp trials

were of the same shift. Additionally, error-clamp trials only occurred on trials where

there was a switch in context and not after one or two trials of the same shift. To

avoid directional feedback, the vBOT position was indicated via an expanding semi-

circle representing only the distance from start location [24]. At the end of the task

each participant was asked if they had noticed the relationship between the

visuospatial context and direction of the force in order to understand their explicit

knowledge during the task.

Transcranial Direct Current Stimulation

Anodal TDCS was delivered via two sponge electrodes (5 x 7 cm) soaked in saline

solution, using a nurostym tES device (Neuro Device Group S.A., Poland). For

cerebellar stimulation the anodal electrode was placed over the right cerebellar

cortex (3 cm lateral to the inion; [25]) and the cathode was positioned on the superior

aspect of the right trapezius muscle [19, 21, 26]. For M1 stimulation the anodal

electrode was positioned over the hand area of the left motor cortex, identified by

single pulse TMS (Magstim Rapid2 stimulator; Magstim Ltd, UK) delivered at

suprathreshold stimulus intensity so as to elicit a visible twitch of the first dorsal

interosseous muscle. The cathode electrode was placed on the skin over the

contralateral supraorbital ridge [6]. TDCS was only delivered during the adaptation

phase, the parameters of which depended on the stimulation group. Once the task

had ended, participants rated their perceived comfort and confidence in their belief

that they received real stimulation on a 10-point visual analogue scale (VAS). They

were also asked if they noticed anything specific regarding the timing of the

stimulation, with respect to the task.

In the main experiment, participants in the M1 and cerebellar stimulation groups

received brief epochs of TDCS during the adaptation phase, with each epoch,

temporally overlapping with movements made through the CCW force-field, when

the task display was shifted to the right. Stimulation was ramped up over 1 second

during the hold period (1 second prior to the movement cue). It was then held at 2

mA for 1 second during the movement and then ramped down over 1 second as the

vBOT returned the participant's hand to the home position. TDCS was applied on

both force-field and error-clamp trials. In the secondary experimental group, the er-

TDCS protocol was conducted as per the main experiment, however, stimulation

was only applied over the cerebellum and concurrently with movements made during

the CW force-field and associated left-shift in visual display. Data from this group

served as an attempt to replicate the effect found in the main experiment, with

stimulation applied during movements through the opposing force-field direction. For

the sham group, stimulation was ramped up over 10 seconds at the start of the

adaptation phase, held at 2 mA for 10 seconds and then ramped down over a further

10 seconds. The electrode montage was randomly assigned to either M1 or

cerebellar prior to starting the session.

Data & Statistical Analysis

Data collected from the vBOT were analysed offline in MATLAB (The Mathworks,

version R2018b). The lateral deviation (LD) of movements at peak velocity was

calculated for null and force-field trials and subsequently averaged into bins. For

error-clamp trials, a force compensation ratio (FC) was calculated, see equation 2.

$$\frac{\int_{t0}^{t\ end} actual\ force}{\int_{t0}^{t\ end} ideal\ force} \times 100 \qquad (2)$$

Actual force was the forces generated against the channel walls, integrated across

the movement; ideal force was the force required to fully compensate for the

perturbation in force-field trials (ideal force: velocity x field constant). Any LD or FC

values that fell outside ± 2 SD of the mean across the group were excluded prior to

averaging and thus removed from further analysis. LD and FC values were analysed

separately for the two contexts. Area under the learning curve for LD and FC during

each phase of null, force-field and error-clamp trials was calculated to be used for

statistical analysis, providing a measure of total error/adaptation during the task.

Statistical analyses were conducted in MATLAB, R (R Core Team, version 3.6.3)

and SPSS (IBM, version 26). All ANOVAs were run in general linear model format

and followed up with Bonferroni-corrected multiple comparisons, when a significant

main effect or interaction was found. The threshold for statistical significance was set

at $p < 0.05$ and we report partial eta squared (ηp^2) effect sizes for ANOVAs.

Estimation statistics were conducted as per [27], reporting paired Cohen's d effect

sizes and bias-corrected and accelerated confidence intervals (following 5000

bootstrap samples).

Data from the secondary experimental group were processed and analysed

separately as they were collected after the main experiment and this group was not

included in the original study design.

Results and Discussion

We first sought to determine any differences in adaptation between trial contexts for

each stimulation group during the task. We measured the area under the learning

curve (calculated from lateral deviation of movements from the target midline; see

methods for further details) in order to compare adaptation performance on trials

performed with simultaneous stimulation and those without, using a 3x3x2 mixed-

design ANOVA. The ANOVA contained within-subject factors of adaptation phase

(baseline, adaptation, washout) and trial context (CW (left-shift), CCW (right-shift))

and the between-subject factor of stimulation group (M1 er-TDCS, cerebellar er-

TDCS, sham). The ANOVA revealed a significant main effect of adaptation phase (F

$(2, 342) = 1312.45$, $p < 0.001$, $\eta^2 p = 0.89$), stimulation group ($F (2, 342) = 18.93$, $p <$

0.001, $\eta^2 p = 0.1$), and a significant three-way interaction between group, phase and

trial context (F (4,342) = 3.44, p = 0.009, $\eta^2 p = 0.04$). Bonferroni corrected multiple

comparisons revealed that, during adaptation, participants who received cerebellar

er-TDCS made significantly less error, and thus adapted better, on stimulated CCW

trials compared to unstimulated CW trials, p < 0.001 (Figure 2). In contrast, we found

no evidence of enhanced adaptation following M1 er-TDCS, as participants made

similar levels of error on both stimulated and unstimulated trials (p = 0.97). Similarly,

participants in the sham group adapted comparably during left and right-shift trials (p

= 0.72), suggesting there was no in-built task bias (Figure 2).

270   (Figure 2)

Testing the robustness of these effects using estimation statistics [27], confirmed

that er-TDCS applied to the cerebellum selectively improved the adaptation on CCW

trials, while CW trials were unaffected. The effect size for this comparison was

substantial (paired Cohen's d = -1.3) with 95.0% confidence intervals (CI) that did

not overlap zero ([-2.05, -0.53]; Figure 3). Conversely, the paired Cohen's d between

CW and CCW trials for the M1 er-TDCS group was close to zero, which fell well

within the CI (d = -0.0059, 95.0% CI = [-0.66, 0.64]); for the sham group d = -0.061

and 95.0% CI = [-0.69, 0.57].

Further multiple comparisons from the three-way interaction suggest that there no

differences in adaptive performance between trial contexts during baseline for any of

the stimulation groups (all p > 0.504), indicating that any differences during

adaptation were not a result of uneven performance during baseline (Figure 2).

There were also no differences in de-adaptation between trial contexts during

washout trials (all p > 0.23), which is likely a result of similar levels of learning

reached towards the end of the adaptation phase. Additionally, when considering

overall adaptation (adaptation to both CW and CCW trials, sign transformed and

combined), results from the ANOVA revealed that participants in the cerebellar er-

TDCS group made significantly less error compared to the M1 er-TDCS and sham

group (both p < 0.001), with no significant differences between the latter two groups,

p = 0.078.

These results show that event-related stimulation of the cerebellum selectively

enhanced adaptation of stimulated CCW trials compared to CW trials, with no effect

of M1 er-TDCS on context-dependent adaptation or task bias (indicated by sham

data). The reduction of error on CCW (right-shift) trials following cerebellar er-TDCS

also contributed to (at least in part) an overall improvement in adaptation during the

task (Figure 2 & Supplementary Figure 1).

299   (Figure 3)

Results from our additional experimental group helped to confirm the findings from

our main experiment. We found that er-TDCS over the cerebellum enhanced

adaptation of movements through the stimulated context compared to the

unstimulated control – a replication of the effect initially found (see Figure 4a). Area

under the learning curve was compared in a 3x2 way ANOVA (Task Phase:

baseline, adaptation, washout; Trial Context: CW (left-shift), CCW (right-shift)), and

revealed significant main effects of task phase ($F_{(2,102)} = 248.23$, $p < 0.001$, $\eta^2 p = $

0.83), trial context ($F_{(1,102)} = 12.49$, $p < 0.001$, $\eta^2 p = 0.11$) and a significant

interaction ($F_{(2,102)} = 6.05$, $p = 0.003$, $\eta^2 p = 0.11$). Bonferroni corrected multiple

comparisons showed that during the adaptation phase participants made

significantly less error on stimulated CW trials, compared to CCW trials (p < 0.001).

We found no evidence to suggest performance during the two trial contexts were

different during baseline or washout phases (both p > 0.53). Estimation statistics

affirmed this result and revealed a large effect size (paired cohen's d = 0.82, 95% CI

= [0.095, 1.43]), with zero falling outside the 95% CI range (see Figure 4b).

317   (Figure 4)

One concern was that short epochs of stimulation, with rapid onset/offset, might be

perceived by the participant and act as a form of attentional cue. However, we found

no significant differences between their self-reported Confidence in Stimulation (F

$(3,74) = 0.078$, $p = 0.97$, $\eta^2 p = 0.003$), suggesting equivalent blinding to the

stimulation condition was achieved. Given that this er-TDCS protocol is relatively

novel, it is also pleasing that the short bouts of TDCS did not cause excessive

discomfort (Perceived Comfort: F $(3,74) = 2.31$, $p = 0.084$, $\eta^2 p = 0.085$), with all

groups reporting low levels of discomfort. Additionally, comfort levels were similar to

those reported after 17 minutes of continuous TDCS [21]). Crucially, only one

participant (in the M1 er-TDCS group) noticed that the stimulation only occurred on

CCW force-field trials with a corresponding right-shift in task display. The lack of

awareness of stimulation timing in the cerebellar group suggests enhanced

adaptation cannot be due to explicit cueing or other explicit mechanism such as

increased attention towards CCW (right-shift) trials.

These initial results suggest that brief periods of TDCS applied over the

cerebellum and in synchrony with movement can selectively and specifically improve

motor adaptation of that movement, while leaving adaptation of interleaved

movements unaffected. Although many studies have reported positive effects of

TDCS on motor learning and rehabilitation, when applied continuously for 10-20

minutes [21, 28–34], there are a growing number of studies reporting null or mixed

effects [35–38], leading to uncertainty around the effectiveness of TDCS [39]. During

continuous stimulation, for 10-20 minutes, any number of different behaviours may

be performed alongside the specific task that is the 'target' of stimulation. The

concatenation of all these behaviours under the same stimulation conditions may

lead to changes in excitation levels in multiple cortical circuits that confound the

results, contributing to some of the conflicting findings [40]. In contrast, event-related

TDCS may selectively modulate only those circuits and task-related synapses that

are contemporaneously active and undergoing concurrent plasticity [16, 41]. As

such, er-TDCS could prove beneficial when long experimental or rehabilitative

protocols are required, as recent research suggests that long continuous bouts of

TDCS may cause the resultant modulatory effects to diminish and even reverse over

time, i.e., from excitation to inhibition [42]. Neural recording studies have also found

that the effects of direct current stimulation may attenuate during long stimulation

blocks. These changes have been seen in spontaneous neural activity and evoked

potentials, due to short-term habituation-like adaptation processes [43–45].

The plastic mechanisms responsible for the improvement in adaptation of specific

movements associated with er-TDCS in this study are still unclear. We propose that

(1) applying anodal TDCS concurrently during movement enhances activity within

the neural circuit associated with and activated by this specific behaviour [16]. This

increased activity in turn could potentiate the synapses within this circuit that are

'eligible' for Hebbian change during the stimulated behaviour. (2) There are

homeostatic mechanisms that act to down-regulate the increase in activity [43-45],

that might be of slow onset, most prominent during prolonged TDCS. (3) By

providing brief er-TDCS, it is more likely that only those circuits involved in the

particular (concurrent) behavioural context are the ones that are potentiated, without

homeostatic reduction. In essence, er-TDCS provides transient heightened

potentiation that 'focuses' Hebbian learning onto specific neurobehavioural circuits

[41, 46].

However, there are other plausible mechanisms that are less reliant on synaptic

plasticity mechanisms. For instance, by boosting the excitability of circuits of

cerebellar neurons during movement, er-TDCS may act to increase the number of

active neurons within the circuit associated with the specific behaviour. This would

result in a greater network of cerebellar neurons selectively activated during adaptive

movements, which in turn, could plausibly drive better, more accurate movements

and result in improved adaptation trial-by-trial.

It is also important to consider which learning processes which may be responsible

for this enhanced adaptation. Given that stimulation was ramped up for 1 second

prior to movement initiation, and outlasted the movement, there is scope to suggest

that activity related to motor planning and preparation may have been facilitated

before each stimulated trial, or that error processing after each trial was facilitated.

Additionally, as sensorimotor adaptation is highly dependent on the cerebellum [47–

49], it is possible that er-TDCS modulates forward model processing, leading to

enhanced prediction error and thus more rapid learning during the stimulated

context. Experiments testing some of these suggestions will be reported in due

course.

We should also note that there is some non-uniformity in the apparent mass of the

vBOT when moving in different directions due to the direction of the 'elbow' joint

(Figure 1a). This may have resulted in a small disparity in the errors between the two

movement conditions when the CW and CCW forces were first introduced, as seen

for example in the sham group data (Figure 2). However, this difference was quickly

overcome in the sham and M1 group, and we found significantly improved

adaptation in both CW and CCW cerebellar conditions, so it is unlikely to have had

major impact on the overall results.

Hebbian-like plasticity can also be induced in the motor cortex, for example using

paired associative stimulation (PAS) and use-dependent plasticity (UDP) protocols

[50–53]. However, we found no specific effect of M1 er-TDCS. Adaptation of CW and

CCW trials were not significantly different, and participants performed similarly to the

sham stimulation group. The null effect of M1 er-TDCS may be explained by the

nature of the task. Reach adaptation to force-fields is believed to be predominantly

cerebellar-dependant [48, 53–56], and is dominated by the proximal muscles of the

arm [21, 26]. Thus, targeting M1 may have had relatively little effect on the

processes governing motor adaptation/performance during the task.

The results presented here suggest ways in which TDCS could be utilised in

research and rehabilitation, with a focus on increased temporal resolution. For

example, er-TDCS could be used in conjunction with therapy protocols comprising of

precise temporal epochs, with further scope beyond the motor modality e.g.,

targeting memory encoding or recall [57, 58]. While TMS remains a more temporally

precise non-invasive brain stimulation intervention, er-TDCS may now be

considered, especially given that direct current stimulation has the potential to

modulate neural excitability bi-directionally. It is yet to be determined how this novel

stimulation protocol will transfer to other settings, and future research in diverse

learning contexts, targeting different brain regions, will be required to determine its

robustness.

In conclusion, we have shown that brief epochs of TDCS over the cerebellum can

selectively modulate motor adaptation when delivered coincidentally with reaching

movements. We suggest that the coupling of stimulation and movement influences

mechanisms of Hebbian-like plasticity and thus facilitates learning in the stimulated

context. These initial results potentially open up new possibilities for the use of

TDCS in both research and clinical settings, to improve its effectiveness and

specificity. They also demonstrate that stimulation can be applied in very short bouts

without inducing severe discomfort or side-effects, and without the participants'

explicit knowledge of the specific stimulation protocol.

Acknowledgements

This work was supported by the MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research (CMAR). We thank an anonymous reviewer for their helpful suggestions for the discussion of our results.

Author Contributions

Matthew Weightman: Conceptualization, Methodology, Investigation, Formal analysis, Visualization, Writing - original draft. **John-Stuart Brittain:** Conceptualization, Methodology, Formal analysis, Writing - review & editing. **Alison Hall:** Methodology, Writing - review & editing. **R. Chris Miall:** Conceptualization, Formal analysis, Writing - review & editing.

Ned Jenkinson: Conceptualization, Methodology, Formal analysis, Writing - review & editing.

Competing Interests

The authors declare no competing interests exist

Data and Code Availability

Data used in this manuscript to create figures have been deposited to Mendeley Data and are available at:

DOI: 10.17632/2k33svf244.2

For further information and requests for resources please communicate with corresponding author, Matthew Weightman (mcw423@adf.bham.ac.uk).

References

References

[1] Hebb, D. O. (1949). *The organization of behavior: A neuropsychological theory.*
Wiley, New York.

[2] Kandel, E. R., & Hawkins, R. D. (1992). The biological basis of learning and
individuality. *Scientific American*, *267*(3), 78–87.

[3] Shatz, C. J. (1992). The developing brain. *Scientific American*, *267*(3), 60–67.

[4] Ito, M. (2001). Cerebellar long-term depression: characterization, signal
transduction, and functional roles. *Physiological Reviews*, *81*(3), 1143–1195. doi:
https://doi.org/10.1152/physrev.2001.81.3.1143

[5] Malenka, R. C., & Bear, M. F. (2004). Ltp and ltd: an embarrassment of riches.
Neuron, *44*(1), 5–21. doi: https://doi.org/10.1016/j.neuron.2004.09.012

[6] Nitsche, M. A., & Paulus, W. (2000). Excitability changes induced in the human
motor cortex by weak transcranial direct current stimulation. *The Journal of*
Physiology, *527*(3), 633–639. doi: https://doi.org/10.1111/j.1469-7793.2000.t01-1-
00633.x

[7] Nitsche, M. A., & Paulus, W. (2001). Sustained excitability elevations induced by
transcranial dc motor cortex stimulation in humans. *Neurology*, *57*(10), 1899–1901.
doi: https://doi.org/10.1212/WNL.57.10.1899

[8] Kwon, Y. H., Ko, M.-H., Ahn, S. H., Kim, Y.-H., Song, J. C., Lee, C.-H., ... Jang,
S. H. (2008). Primary motor cortex activation by transcranial direct current
stimulation in the human brain. *Neuroscience Letters*, *435*(1), 56–59. doi:
https://doi.org/10.1016/j.neulet.2008.02.012

[9] Stagg, C. J., Best, J. G., Stephenson, M. C., O'Shea, J., Wylezinska, M., Kincses,
Z. T., . . . Johansen-Berg, H. (2009). Polarity-sensitive modulation of cortical
neurotransmitters by transcranial stimulation. *Journal of Neuroscience*, *29*(16),
5202–5206. doi: https://doi.org/10.1523/JNEUROSCI.4432-08.2009

[10] Kwon, Y. H., & Jang, S. H. (2011). The enhanced cortical activation induced by
transcranial direct current stimulation during hand movements. *Neuroscience*
Letters, *492*(2), 105–108. doi: https://doi.org/10.1016/j.neulet.2011.01.066

[11] Pellicciari, M. C., Brignani, D., & Miniussi, C. (2013). Excitability modulation of
the motor system induced by transcranial direct current stimulation: a multimodal
approach. *Neuroimage*, *83*, 569–580. doi:
https://doi.org/10.1016/j.neuroimage.2013.06.076

[12] Bindman, L. J., Lippold, O., & Redfearn, J. (1964). The action of brief polarizing
currents on the cerebral cortex of the rat (1) during current flow and (2) in the
production of long-lasting after-effects. *The Journal of Physiology*, *172*(3), 369–382.
doi: https://doi.org/10.1113/jphysiol.1964.sp007425

[13] Landau, W. M., Bishop, G. H., & Clare, M. (1964). Analysis of the form and
distribution of evoked cortical potentials under the influence of polarizing currents.
Journal of Neurophysiology, 27(5), 788–813. doi:
https://doi.org/10.1152/jn.1964.27.5.788

[14] Purpura, D. P., & McMurtry, J. G. (1965). Intracellular activities and evoked
potential changes during polarization of motor cortex. *Journal of Neurophysiology,*
28(1), 166-185. doi: https://doi.org/10.1152/jn.1965.28.1.166

[15] Priori, A., Berardelli, A., Rona, S., Accornero, N., & Manfredi, M. (1998).
Polarization of the human motor cortex through the scalp. *Neuroreport, 9*(10), 2257–
2260.

[16] Bikson, M., & Rahman, A. (2013). Origins of specificity during tdcs: anatomical,
activity- selective, and input-bias mechanisms. *Frontiers in Human Neuroscience, 7,*
688. doi: https://doi.org/10.3389/fnhum.2013.00688

[17] Fertonani, A., & Miniussi, C. (2017). Transcranial electrical stimulation: what we
know and do not know about mechanisms. *The Neuroscientist, 23*(2), 109–123. doi:
10.1177/1073858416631966

[18] Bikson, M., Paulus, W., Esmaeilpour, Z., Kronberg, G., & Nitsche, M. A. (2019).
Mechanisms of acute and after effects of transcranial direct current stimulation. In
Practical guide to transcranial direct current stimulation (pp. 81–113). Springer. doi:
https://doi.org/10.1007/978-3-319-95948-1-3

[19] Panouillères, M. T., Joundi, R. A., Brittain, J.-S., & Jenkinson, N. (2015).
Reversing motor adaptation deficits in the ageing brain using non-invasive
stimulation. *The Journal of Physiology, 593*(16), 3645–3655. doi:
https://doi.org/10.1113/JP270484

[20] Spampinato, D. A., Satar, Z., & Rothwell, J. C. (2019). Combining reward and
m1 transcranial direct current stimulation enhances the retention of newly learnt
sensorimotor mappings. *Brain stimulation, 12*(5), 1205–1212. doi:
https://doi.org/10.1016/j.brs.2019.05.015

[21] Weightman, M., Brittain, J.-S., Punt, D., Miall, R. C., & Jenkinson, N. (2020).
Targeted tdcs selectively improves motor adaptation with the proximal and distal
upper limb. *Brain Stimulation, 13*(3), 707–716. doi:
https://doi.org/10.1016/j.brs.2020.02.013

[22] Howard, I. S., Ingram, J. N., & Wolpert, D. M. (2009). A modular planar robotic
manipulandum with end-point torque control. *Journal of Neuroscience Methods,*
181(2), 199–211. doi: https://doi.org/10.1016/j.jneumeth.2009.05.005

[23] Howard, I. S., Wolpert, D. M., & Franklin, D. W. (2013). The effect of contextual
cues on the encoding of motor memories. *Journal of Neurophysiology, 109*(10),
2632–2644. doi: https://doi.org/10.1152/jn.00773.2012

[24] Lago-Rodriguez, A., & Miall, R. C. (2016). Online visual feedback during error-
free channel trials leads to active unlearning of movement dynamics: evidence for
adaptation to trajectory prediction errors. *Frontiers in Human Neuroscience*, *10*, 472.
doi: https://doi.org/10.3389/fnhum.2016.00472

[25] Galea, J. M., Jayaram, G., Ajagbe, L., & Celnik, P. (2009). Modulation of
cerebellar excitability by polarity-specific noninvasive direct current stimulation.
Journal of Neuroscience, *29*(28), 9115–9122. doi:
https://doi.org/10.1523/JNEUROSCI.2184- 09.2009

[26] Weightman, M., Brittain, J.-S., Miall, R. C., & Jenkinson, N. (2021). Direct and
indirect effects of cathodal cerebellar tdcs on visuomotor adaptation of hand and arm
movements. *Scientific Reports*, *11*(1), 1–10. doi: https://doi.org/10.1038/s41598-
021-83656-5

[27] Ho, J., Tumkaya, T., Aryal, S., Choi, H., & Claridge-Chang, A. (2019). Moving
beyond p values: data analysis with estimation graphics. *Nature Methods*, *16*(7),
565–566. doi: 10.1038/s41592-019-0470-3

[28] Hummel, F. C., Celnik, P., Giraux, P., Floel, A., Wu, W.-H., Gerloff, C., & Cohen,
L. G. (2005). Effects of non-invasive cortical stimulation on skilled motor function in
chronic stroke. *Brain*, *128*(3), 490–499. doi: https://doi.org/10.1093/brain/awh369

[29] Fregni, F., Boggio, P. S., Santos, M. C., Lima, M., Vieira, A. L., Rigonatti, S. P., .
. . Pascual-Leone, A. (2006). Noninvasive cortical stimulation with transcranial direct
current stimulation in parkinson's disease. *Movement Disorders*, *21*(10), 1693–1702.
doi: https://doi.org/10.1002/mds.21012

[30] Galea, J. M., Vazquez, A., Pasricha, N., Orban de Xivry, J.-J., & Celnik, P.
(2010). Dissociating the roles of the cerebellum and motor cortex during adaptive
learning: the motor cortex retains what the cerebellum learns. *Cerebral Cortex*,
21(8), 1761– 1770. doi: https://doi.org/10.1093/cercor/bhq246

[31] Hardwick, R. M., & Celnik, P. A. (2014). Cerebellar direct current stimulation
enhances motor learning in older adults. *Neurobiology of Aging*, *35*(10), 2217–2221.
doi: https://doi.org/10.1016/j.neurobiolaging.2014.03.030

[32] Allman, C., Amadi, U., Winkler, A. M., Wilkins, L., Filippini, N., Kischka, U., ...
Johansen-Berg, H. (2016). Ipsilesional anodal tdcs enhances the functional benefits
of rehabilitation in patients after stroke. *Science Translational Medicine*, *8*(330),
330re1–330re1. doi: https://doi.org//10.1126/scitranslmed.aad5651

[33] Benussi, A., Dell'Era, V., Cotelli, M. S., Turla, M., Casali, C., Padovani, A., &
Borroni, B. (2017). Long term clinical and neurophysiological effects of cerebellar
transcranial direct current stimulation in patients with neurodegenerative ataxia.
Brain Stimulation, *10*(2), 242–250. doi: https://doi.org/10.1016/j.brs.2016.11.001

[34] Chiou, S.-Y., Morris, L., Gou, W., Alexander, E., & Gay, E. (2020). Motor cortical
circuits contribute to crossed facilitation of trunk muscles induced by rhythmic arm

movement. *Scientific Reports*, *10*(1), 1–12. doi: https://doi.org/10.1038/s41598- 020-
74005-z

[35] Jalali, R., Miall, R. C., & Galea, J. M. (2017). No consistent effect of cerebellar
transcranial direct current stimulation on visuomotor adaptation. *Journal of*
Neurophysiology, *118*(2), 655–665. doi: https://doi.org/10.1152/jn.00896.2016

[36] Hulst, T., John, L., Ku☐per, M., van der Geest, J. N., Go☐ricke, S. L., Donchin,
O., & Timmann, D. (2017). Cerebellar patients do not benefit from cerebellar or m1
transcranial direct current stimulation during force-field reaching adaptation. *Journal*
of Neurophysiology, *118*(2), 732–748. doi: https://doi.org/10.1152/jn.00808.2016

[37] Mamlins, A., Hulst, T., Donchin, O., Timmann, D., & Claassen, J. (2019). No
effects of cerebellar transcranial direct current stimulation (tdcs) on force field and
visuomotor reach adaptation in young and healthy subjects. *Journal of*
Neurophysiology. doi: https://doi.org/10.1152/jn.00352.2018

[38] Wiltshire, C. E., & Watkins, K. E. (2020). Failure of tdcs to modulate motor
excitability and speech motor learning. *Neuropsychologia*, *146*, 107568. doi:
https://doi.org/10.1016/j.neuropsychologia.2020.107568

[39] Horvath, J. C., Forte, J. D., & Carter, O. (2015). Evidence that transcranial direct
current stimulation (tdcs) generates little-to-no reliable neurophysiologic effect
beyond mep amplitude modulation in healthy human subjects: a systematic review.
Neuropsychologia, 66, 213–236. doi:
https://doi.org/10.1016/j.neuropsychologia.2014.11.021

[40] Antal, A., Terney, D., Poreisz, C., & Paulus, W. (2007). Towards unravelling
task-related modulations of neuroplastic changes induced in the human motor
cortex. *European Journal of Neuroscience*, *26*(9), 2687–2691. doi:
https://doi.org/10.1111/j.1460- 9568.2007.05896.x

[41] Kronberg, G., Bridi, M., Abel, T., Bikson, M., & Parra, L. C. (2017). Direct current
stimulation modulates ltp and ltd: activity dependence and dendritic effects. *Brain*
Stimulation, *10*(1), 51–58. doi: 10.1016/j.brs.2016.10.001

[42] Hassanzahraee, M., Nitsche, M. A., Zoghi, M., & Jaberzadeh, S. (2020).
Determination of anodal tdcs duration threshold for reversal of corticospinal
excitability: An investigation for induction of counter-regulatory mechanisms. *Brain*
Stimulation, *13*(3), 832–839. doi: https://doi.org/10.1016/j.brs.2020.02.027

[43] Creutzfeldt, O. D., Fromm, G. H., & Kapp, H. (1962). Influence of transcortical dc
currents on cortical neuronal activity. *Experimental Neurology*, *5*(6), 436–452. doi:
https://doi.org/10.1016/0014-4886(62)90056-0

[44] Kunori, N., & Takashima, I. (2019). Evaluation of acute anodal direct current
stimulation- induced effects on somatosensory-evoked responses in the rat. *Brain*
Research, *1720*, 146318. doi: https://doi.org/10.1016/j.brainres.2019.146318

[45] Asan, A. S., Lang, E. J., & Sahin, M. (2020). Entrainment of cerebellar purkinje cells with directional ac electric fields in anesthetized rats. *Brain Stimulation*. doi: https://doi.org/10.1016/j.brs.2020.08.017

[46] Kronberg, G., Rahman, A., Sharma, M., Bikson, M., & Parra, L. C. (2020). Direct current stimulation boosts hebbian plasticity in vitro. *Brain Stimulation*, *13*(2), 287–301. doi: https://doi.org/10.1016/j.brs.2019.10.014

[47] Bastian, A. J. (2006). Learning to predict the future: the cerebellum adapts feedforward movement control. Current opinion in neurobiology, 16(6), 645–649. doi: https://doi.org/10.1016/j.conb.2006.08.016

[48] Tseng, Y.-w., Diedrichsen, J., Krakauer, J. W., Shadmehr, R., & Bastian, A. J. (2007). Sensory prediction errors drive cerebellum-dependent adaptation of reaching. Journal of Neurophysiology, 98(1), 54–62. doi: https://doi.org/10.1152/jn.00545.2007

[49] Wong, A. L., Marvel, C. L., Taylor, J. A., & Krakauer, J. W. (2019). Can patients with cerebellar disease switch learning mechanisms to reduce their adaptation deficits? Brain, 142(3), 662–673. doi: https://doi.org/10.1093/brain/awy334

[50] Bütefisch, C. M., Khurana, V., Kopylev, L., & Cohen, L. G. (2004). Enhancing encoding of a motor memory in the primary motor cortex by cortical stimulation. Journal of Neurophysiology, 91(5), 2110–2116. doi: https://doi.org/10.1152/jn.01038.2003

[51] Ziemann, U., Ilia ́c, T. V., Pauli, C., Meintzschel, F., & Ruge, D. (2004). Learning modifies subsequent induction of long-term potentiation-like and long-term depression-like plasticity in human motor cortex. Journal of Neuroscience, 24(7), 1666–1672. doi: https://doi.org/10.1523/JNEUROSCI.5016-03.2004

[52] Rosenkranz, K., Kacar, A., & Rothwell, J. C. (2007). Differential modulation of motor cortical plasticity and excitability in early and late phases of human motor learning. Journal of Neuroscience, 27(44), 12058–12066. doi: https://doi.org/10.1523/JNEUROSCI.2663-07.2007

[53] Koch, G., Ponzo, V., Di Lorenzo, F., Caltagirone, C., & Veniero, D. (2013). Hebbian and anti-hebbian spike-timing-dependent plasticity of human cortico-cortical connections. Journal of Neuroscience, 33(23), 9725–9733. doi: 10.1523/JNEUROSCI.4988-12.2013

[54] Maschke, M., Gomez, C. M., Ebner, T. J., & Konczak, J. (2004). Hereditary cerebellar ataxia progressively impairs force adaptation during goal-directed arm movements. Journal of Neurophysiology, 91(1), 230–238. doi: https://doi.org/10.1152/jn.00557.2003

[55] Rabe, K., Livne, O., Gizewski, E. R., Aurich, V., Beck, A., Timmann, D., & Donchin, O. (2009). Adaptation to visuomotor rotation and force field perturbation is correlated to different brain areas in patients with cerebellar degeneration. Journal of Neurophysiology, 101(4), 1961–1971. doi: https://doi.org/10.1152/jn.91069.2008

640   [56] Criscimagna-Hemminger, S. E., Bastian, A. J., & Shadmehr, R. (2010). Size of
641   error affects cerebellar contributions to motor learning. Journal of Neurophysiology,
642   103(4), 2275–2284. doi: https://doi.org/10.1152/jn.00822.2009
643
[57] Donchin, O., Rabe, K., Diedrichsen, J., Lally, N., Schoch, B., Gizewski, E. R., &
Timmann, D. (2012). Cerebellar regions involved in adaptation to force field
and visuomotor perturbation. Journal of Neurophysiology, 107(1), 134–147. doi:
https://doi.org/10.1152/jn.00007.2011

[58] Javadi, A. H., Cheng, P., & Walsh, V. (2012). Short duration transcranial direct
current stimulation (tdcs) modulates verbal memory. Brain Stimulation, 5(4), 468–
474. doi: https://doi.org/10.1016/j.brs.2011.08.003
651
652   [59] MacDonald, H. J., Brittain, J.-S., Spitzer, B., Hanslmayr, S., & Jenkinson,
653   N. (2019). Memory deficits in parkinson's disease are associated with
654   reduced beta power modulation. Brain Communications, 1(1), fcz040. doi:
655   https://doi.org/10.1093/braincomms/fcz040
656

Figure Legends

Figure 1: Experimental design and set-up.
*a. A schematic of the task set-up, with an example screen display. Movements were always made in the midline position, but the cursor, home and target markers would be shifted either 10cm to the right or left. The order of the contextual shift in the task display was pseudorandomised, with an equal number in each task phase and runs of no more than three trials of the same contextual shift. Examples of the task display during null field **(b)** and context-dependent force-field **(c)** trials for both trial types.*

Figure 2: Cerebellar er-TDCS selectively improves context-dependent force-field adaptation.
***Top panel:** Mean lateral deviation for CW (left-shift) and CCW (right-shift) trials (± standard error, shaded regions), averaged into bins of two trials for the M1 er-TDCS, cerebellar er-TDCS and sham stimulation groups. The M1 and cerebellar groups received er-TDCS on CCW trials during the adaptation phase. **Lower panel:** mean LD (bins of two trials) for all three stimulation groups during either CW or CCW contextual trials.*

Figure 3: Direct comparisons of adaptation during CW and CCW force-fields.
***Top panel:** area under the curve (cm x bins) for CW vs CCW trials for each participant during the adaptation phase, with each participant's data connected by a line. **Lower panel:** paired Cohen's d plotted as a bootstrap sampling distribution. Mean differences are depicted as dots, with 95.0% CIs indicated by black vertical bars.*

Figure. 4: Adaptation performance and context selectivity for the secondary experimental group receiving er-TDCS during CW trials.
*a: Mean lateral deviation for CW (left-shift) and CCW (right-shift) trials (± standard error, shaded regions), averaged into bins of two trials for the secondary experimental group. Participants in this group received er-TDCS over the cerebellum during CW adaptation trials. Panel **(b.)** depicts the area under the curve (cm x bins) for CW vs CCW trials for each participant during the adaptation phase, with each participant's data connected by a line in a Paired Gardner-Altman plot, with the paired Cohen's d plotted as a bootstrap sampling distribution alongside. The mean difference is shown as a dot, with 95.0% CIs indicated by black vertical bars.*

a.

b.